

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

Evidence of the impact of deep convection on reactive volatile organic compounds in the upper tropical troposphere during the AMMA experiment in West Africa

J. Bechara¹, A. Borbon¹, C. Jambert¹, A. Colomb^{1,*}, and P. E. Perros¹

¹Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), Universités Paris 12 et Paris 7, CNRS, 61, avenue du Général de Gaulle, Créteil, France

* now at: Laboratoire d'Aérodynamique (LA), Université Paul Sabatier, CNRS, Toulouse, France

Received: 18 July 2009 – Accepted: 12 August 2009 – Published: 29 September 2009

Correspondence to: J. Bechara (joelle.bechara@lisa.univ-paris12.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

A large dataset of reactive trace gases was collected for the first time over West Africa during the African Monsoon Multidisciplinary Analysis (AMMA) field experiment in August 2006. Volatile Organic Compounds (VOC from C5–C9) were measured onboard the two French aircrafts the ATR-42 and the Falcon-20 by a new instrument AMOVOC. The goal of this study is (i) to characterize VOC distribution in the tropical region of West Africa (ii) to determine the impact of deep convection on VOC distribution and chemistry in the tropical upper troposphere (UT) and (iii) to characterize its spatial and temporal extensions. Experimental strategy consisted in sampling at altitudes between 0 and 12 km downwind of Mesoscale Convective Systems (MCS) and at cloud base. Biogenic and anthropogenic VOC distribution in West Africa is clearly affected by North to South emission gradient. Isoprene, the most abundant VOC, is at maximum level over the forest (1.26 ppb) while benzene reaches its maximum over the urban areas (0.11 ppb). First, a multiple physical and chemical tracers approach using CO, O₃ and relative humidity was implemented to distinguish between convective and non-convective air masses. Then, additional tools based on VOC observations (tracer ratios, proxy of emissions and photochemical clocks) were adapted to characterize deep convection on a chemical, spatial and temporal basis. VOC vertical profiles show a “C-shaped” trend indicating that VOC-rich air masses are transported from the surface to the UT by deep convective systems. VOC mixing ratios in convective outflow are up to two times higher than background levels even for reactive and short-lived VOC (e.g. isoprene up to 0.19 ppb at 12 km-altitude) and are dependent on surface emission type. As a consequence, UT air mass reactivity increases from 0.52 s⁻¹ in non-convective conditions to 0.95 s⁻¹ in convective conditions. Fractions of boundary layer air contained in convective outflow are estimated to be 40±15%. Vertical transport timescale is calculated to be 25±10 min. These results characterize deep convection occurring over West Africa and provide relevant information for tropical convection parameterization in regional/global models.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Tropospheric ozone (O₃) plays an essential role in determining the oxidizing capacity of the atmosphere and contributes largely to the global greenhouse effect. It is a major tropospheric oxidant and controls global atmospheric chemistry. Tropospheric ozone production is considered to be mainly driven by carbon monoxide (CO) and volatile organic compounds (VOC) oxidation in presence of nitrogen oxides (NO_x) and sunlight. Trace gases originate from both biogenic and anthropogenic emissions. Tropical regions are major sources of these ozone precursors and undergo intensive solar radiation. This induces highly active photochemistry leading to high ozone formation (Kesselmeier et al., 2000; Guenther et al., 2006; Saxton et al., 2007). Nevertheless, the ozone budget is not well established yet over these regions due to complex coupling between chemical processes and dynamics. Indeed, significant deviations exist between observed and modeled global distribution of ozone in tropical regions (Martin et al., 2002; Stevenson et al., 2006). Moreover, modeling studies show discrepancies in ozone budgets when air masses are subject to deep convective events. Some studies show an increase of up to 12% in the global ozone burden due to convective transport of gaseous precursors (Lawrence et al., 2003) while others show a decrease of 13 to 20% (Doherty et al., 2005; Lelieveld and Crutzen, 1994). In fact, all these studies highlight that convective transport of gaseous precursors exhibit a critical impact on tropospheric ozone budget. In the last two decades, some observational campaigns have been conducted worldwide to better understand the role of deep convection and its efficiency to transport gaseous species from the boundary layer to the upper troposphere (UT). Many studies over mid-latitudes (Dickerson et al., 1987; Hauf et al., 1995; Pickering et al., 1988) showed that deep convection could pump low tropospheric air to the tropopause region and proved effective transport of trace gases (e.g. CO, O₃; Ström et al., 1999; Fischer et al., 2003) even highly reactive and short-lived (e.g. VOC, Colomb et al., 2006; Stickler et al., 2006). More recently, a few campaigns above the tropical Amazonian forest (Warneke et al., 2001; Ganzeveld et al., 2008; Stickler et al.,

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2007) showed important influence of deep convection on VOC such as isoprene and formaldehyde. All these field observations reported that UT trace gas levels related to convective events were significantly higher than background levels. Generally, updrafts in deep convective systems have vertical speeds of 5 to 15 m s⁻¹ transporting air masses from low troposphere (LT) to the UT in about 30 min (Thompson et al., 1997; Houze, 2004). Mixing, chemical loss and deposition appear negligible during vertical transport for major species like CO, NO_x and VOC (methane, alkanes, alkenes, aromatics ...). Finally, deep convection acts as an additional source of reactive trace gases in the middle and upper troposphere by fast injections at various levels including cloud top, e.g. 10–15 km or more. Indeed, when rapidly transported to the UT, atmospheric lifetimes of reactive trace gases are longer (Dickerson et al., 1987; Poisson et al., 2000) due to temperature decrease and lower OH concentrations. As a consequence, trace gases will persist longer, leading to delayed O₃ formation (Dickerson et al., 1987). Gases with longer lifetimes (e.g. CO) can also travel far from source region due to higher wind speeds expanding the spatial scale of their influence on O₃ and HO_x radicals production. But, models to observations deviations have highlighted the lack of understanding of chemical and dynamical processes in tropical regions and the paucity of observations in these areas (Lelieveld and Crutzen, 1994; Lawrence et al., 2003; Doherty et al., 2005). Therefore, it is essential to collect new in-situ observation on atmospheric composition in tropical regions in order to better characterize on-going chemical and dynamical processes.

West Africa has never been exhaustively explored up to now. It is a particularly sensitive region because of its location in the tropical belt and its contrasted land-surface characteristics (from bare soils in the Sahel to dense vegetation in the tropical forest) as well as dynamical, chemical and human interactions that modulate significantly trace gases distribution and fate. West Africa is particularly sensitive during the monsoon season, when occurrence of mesoscale convective systems (MCS) is most recurrent (Houze, 2004; Zipser et al., 2006). The West African monsoon is due to several winds converging toward the Inter Tropical Convergence Zone (ITCZ) caused by the intense

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

solar heating of the equatorial zone. In the low layers (up to 3 km), a continental north-easterly wind (Harmattan), warm and dry, converges with a south-westerly wet and colder wind (monsoon flow). In the middle troposphere, a distinct layer is established between 3 and 4 km by the African Easterly Jet (AEJ) around 15° N which is an intense current reaching its utmost during the wet season. The AEJ allows westward air mass transport. In the UT, another westward current, the Tropical Easterly Jet (TEJ), flows above the Sahel at about 16 km altitude during the wet season (Lafore and Moncrieff, 1989; Nicholson, 2009).

The African Monsoon Multidisciplinary Analysis (AMMA) is an international program that aims to improve our knowledge and understanding of the West African monsoon and its variability. One of AMMA objectives is to determine the chemical composition of the middle and upper troposphere connected to deep convection events and to assess the ozone budget over West Africa (Mari and Prospero, 2005; Redelsperger et al., 2006). A Special Observation Period (SOP 2a2) took place in July and August 2006 during the monsoon season to fulfill these objectives. An overall review of the SOP is related in Reeves et al., 2009). Within this framework, the purpose of our study is (i) to characterize VOC vertical and latitudinal distribution in the tropical troposphere over West Africa (ii) to determine the impact of deep convection on VOC distribution and chemistry in the tropical UT and (iii) to characterize its spatial and temporal extensions. This work is based on airborne observations of trace gases and reactive VOC, collected on board the two French aircrafts, the ATR-42 and the Falcon-20 in August 2006. Various diagnostic tools (dynamical and chemical tracers, VOC tracer ratios, emission proxy, photochemical clocks) are set up and used together to investigate the impact of deep convection and its spatial and temporal characteristics. Section 2 illustrates briefly the campaign. Section 3 describes the instrumental set up. Section 4 reports VOC observations and their spatial distribution over the domain. Section 5 establishes a set of conditions by using CO, O₃ and relative humidity (RH) as convection indicators to distinguish non-perturbed conditions from those perturbed by convection. Section 6 assesses the impact of deep convection on trace gases and evaluates its

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

characteristics on the basis of VOC observations.

2 Field campaign and experimental strategy

SOP 2a2 took place in August 2006 during the monsoon season in West Africa. Measurements were performed by both French research aircraft ATR-42 and F-F20 based in Niamey (Niger) and operated by the French Service of Instrumented Aircraft for Environmental Research (SAFIRE). Flights were carried out between 2 and 21 August. The ATR-42 and F-F20 conducted 18 and 10 research flights, respectively, of 3 to 4 h each, from surface to 12 km. Flight details are reported in Table 1 and F-F20 flight tracks are shown in Fig. 1. Observations cover the region from 2 to 21° N and from 10° W to 7° E. The domain displays different surface types with a strong North-to-South vegetation gradient around 12° N. Above these latitudes, the surface presents almost bare soils to a sahelian type. Below 12° N, the surface presents an important tree coverage and vegetated surface. Main urban sites are Niamey (13.5° N, 2.1° E) in Niger and Cotonou (6.3° N, 2.4° E) in south Benin.

Flights fulfilled two objectives: (1) determination of the horizontal and vertical distribution of trace gases and (2) characterization of the air composition affected by MCS. Deploying two aircraft aimed at maximizing the horizontal and vertical geographical coverage of the region. The ATR-42 platform was set up to sample the LT up to 6 km and to perform flights before and after the passage of MCS. The F-F20 platform was used to sample the middle and upper troposphere up to 12 km and to explore MCS outflow undertaking several flight tracks at increasing distances from the cloud anvil. Flights were planned according to meteorological forecast reports carried out by AC-MAD (African Centre of Meteorological Application for Development) as well as MCS tracking using Meteosat-IR images. On the 11, 14, 15 and 17 August, four MCS passed over Niamey and were explored by the F-F20. On the 18 August, an MCS passed over Niamey. The convective region was explored by the ATR-42 and the F-F20 the day after (19 August). On the 13 and 16 August, F-F20 flights were performed in clear con-

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ditions as reference cases. Other flights by both aircrafts performed horizontal tracks down to Cotonou and over Gulf of Guinea.

3 Aircraft instrumentation

The French aircraft were both equipped with instrumentation to measure major trace gases: CO, O₃, NO_x, and VOC (formaldehyde and non-methane hydrocarbons (NMHC) from C₅–C₉ including alkanes, alkenes and aromatic compounds). Table 2 describes the instrumentation characteristics. NMHC were collected on cartridges using a new offline automatic sampler AMOVOC (Airborne Measurement Of Volatile Organic Compounds) developed at LISA (Bechara et al., 2008). AMOVOC was deployed simultaneously on both aircrafts. About 200 cartridges were collected using AMOVOC during SOP 2a2. AMOVOC performs 10 min integrated in-situ sampling on solid adsorbents (Carbosieve SIII, Carbopack B, Carbotrap C) at pre-selected times intervals. Sampling is performed during horizontal flight tracks at 200 mL min⁻¹ flow rate. Samples were shipped to the laboratory in Creteil, France and analyzed by TDAS/GC-MS (Chromtech/Varian). Details of analytical procedures are also given by Bechara et al., 2008). In addition, state parameters of temperature, pressure and aircraft position (altitude, latitude and longitude), irradiance and relative humidity were measured using the standard Safire aircraft instrument package. For most of the instruments, signals were recorded at 1 Hz frequency.

4 NMHC distribution

Air samples were collected throughout the tropospheric column from surface level up to 12 km. 18 NMHC (C₅ to C₉) comprising alkanes, alkenes and aromatics have been identified and quantified during AMMA. Collected NMHC cover three classes of reactivity (Carter, 1994). The most reactive compounds (due to OH radical) with a short

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

lifetime <1 day are isoprene, pentenes, xylenes and trimethylbenzenes. Lifetime of these compounds is especially governed by photochemistry. Compounds with intermediate lifetimes of 1 to 3 days are C5–C8 alkanes, toluene and ethylbenzene. These compounds can undergo photochemistry and transport concurrently. The least reactive compound with relatively long lifetime >3 days is benzene (lifetime ~9 days). Its concentration is primarily controlled by mixing and transport. Observed C5–C9 hydrocarbon concentrations ranged from 1.26 ppb (for isoprene) near the surface to values below detection limits (DL) (1–5 ppt). The most abundant NMHC and their mean values are: isoprene (0.17 ppb), toluene (0.16 ppb) and hexane (0.13 ppb) in the LT and ethylbenzene (0.18 ppb), hexane (0.13 ppb) and toluene (0.13 ppb) in the UT. Only 5% of the collected samples have values below DL.

4.1 NMHC latitudinal patterns

Figure 2 illustrates the north to south variability of some representative biogenic (isoprene) and anthropogenic NMHC (trans-2-pentene being the most reactive NMHC and benzene being the least reactive one; Carter, 1994). Land surface impact on NMHC mixing ratio variability is determined by considering data at altitudes below 2 km-altitude. Average mixing ratios and corresponding standard deviations over 1° latitudinal bins are reported in Fig. 2 using the log-normal distribution for statistical treatment. The number of samples in each latitude bin varies between 1 (at 7° N) and 32 (at 12° N). Standard deviation will then reflect the variability as well as the number of samples collected. Biogenic and anthropogenic compounds show an opposite pattern due to surface characteristics.

Isoprene concentrations are at constant level (about 0.20 ppb) between 14 and 11° N. Then, they show a clear increase between 11 and 7° N to about 0.40 ppb over the forest areas. At lower latitude, around Cotonou (6.3° N) and over the ocean, isoprene levels decrease to less than 0.10 ppb between 7 and 5° N. This trend clearly shows that biogenic isoprene is modulated by the vegetation gradient observed in West Africa.

Anthropogenic NMHC variability shows the highest concentrations values (around

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



0.11 ppb for benzene and 0.05 ppb for trans-2-pentene) near urban sites (Cotonou at 6.3° N and Niamey at 13.5° N). Mean levels decrease above the forest between 11 and 7° N reaching about 0.06 ppb for benzene and 0.02 ppb for trans-2-pentene. The trans-2-pentene decrease is more pronounced than benzene due to its higher reactivity.

5 Finally, LT content is tightly related to surface coverage. The contrasted surface coverage is a key factor influencing NMHC concentration gradients in West Africa.

4.2 NMHC vertical patterns

Vertical profiles for a selection of representative NMHC are plotted in Fig. 3. All observations as well as mean values in 1 km altitudinal bins are reported. The number of samples in each altitude bin varies between 4 and 51. Average mixing ratios and corresponding standard deviation in each altitude bin are calculated using log-normal distribution. The magnitude of standard deviation reflects the variability and the number of samples collected in each bin. Mixing ratios in air masses below 2 km-altitude show a large variability depicting the influence of surface emissions (Sect. 4.1). In the mid troposphere, mixing ratios tend to decrease with altitude. But, the vertical gradient is not of similar magnitude and shape for the different species.

Short-lived compounds exhibit a decrease with altitude (e.g. isoprene, trans-2-pentene). Relatively long-lived compounds (few days) show a homogeneous vertical profile (e.g. pentane, benzene, toluene, hexane in Fig. 3) attesting of the well-mixed column and the redistribution of surface emission in the higher layers.

A distinct layer is detected between 3 and 4 km. In this layer significant concentrations are measured. At this level, air masses are attributed to AEJ transporting air masses from the East (e.g. from Chad, Soudan). Another marked layer is met at about 9 km with high concentrations of pentane, benzene, toluene, hexane, octane, xylenes, ethylbenzene and trimethylbenzenes. These compounds that could originate from biomass burning (Hao et al., 1996; Karl, 2007) indicate biomass burning plumes that were crossed around Cotonou on 19 August. It is also consistent with CO enhancements (Mari et al., 2008; Ancellet et al., 2009). These plumes come from the

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Hemisphere where biomass burning is intense during this period.

In the UT (10–12 km), samples with high mixing ratios even for short-lived NMHC are encountered conferring a “C”-shaped profile to the NMHC vertical distribution. For instance, the reactive isoprene is detected at a maximum of 0.20 ppb on-board the F-F20 at 12 km-height even though it was thought to be present only near the surface where biogenic sources are important (Sect. 4.1). Moreover, short-lived aromatic compounds (1,2,3-trimethylbenzene, Fig. 3), show high concentrations in the UT compared to LT levels. Additional enrichment by long range transport from eastern countries is suspected as seen from back trajectories calculations by Ancellet et al. (2009). Such significant levels reveal that MCS might have influenced the free troposphere by uplifting NMHC from the lower troposphere. Although concentration levels generally decrease with altitude, significant levels were observed during this campaign, in the middle and upper troposphere. In fact, the tropical troposphere is thought to be frequently influenced by recent convective activity, transporting important levels of these species to the upper layers.

Finally, NMHC profile analysis highlights factors that concurrently influence NMHC distribution over West Africa: chemical loss, long range transport of polluted air masses, deep convection and biomass burning intrusion in the region.

5 Indicators used for classification of MCS impacted air masses

Sampled air masses in the UT have different origins (convective transport, biomass burning or long range transport) that could impact trace gases variability as revealed by NMHC observations in Sect. 4. Air masses classification is essential in order to distinguish between conditions perturbed by convection from those non-perturbed by convection. For that purpose, a multiple chemical and physical tracers approach has been taken using CO, O₃ and RH as indicators. Apart from their properties that will be discussed further, tracer selection criteria are: (i) high sampling time resolution (1–30 s) and (ii) atmospheric lifetime longer than MCS duration (2 months for CO and several

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

days for O₃) (Dickerson et al., 1987; Dessler, 2002; Folkins et al., 2002; Lawrence and Salzman, 2008).

Convective conditions are characterized by concurrent high CO, low O₃ and high RH in the air masses (Fig. 4, red lines). CO concentrations might reach 200 ppb, almost as large as those observed in LT, giving a “C”-shaped profile to CO vertical distribution. O₃ values are lower in MCS outflow (30 ppb) than outside it (50 ppb) because of the input of low-O₃ air masses from near the surface. O₃ follows an “S-shaped” profile in convective situations. RH is enhanced in the UT reaching values up to 80% as a signature of cloud presence as well as uplift of humid air masses from surface layers. These concomitant alterations in tracer levels point out important uplift of LT air by the MCS up to the UT and characterize convective air masses.

Average CO, O₃ and RH vertical patterns collected during all deployments of the ATR-42 and the F-F20 are discussed in details in Reeves et al. (2009) and Ancellet et al. (2009). Here, only average features are described. CO is produced primarily by incomplete combustion through urban areas and biomass burning emissions. Its major sink is the reaction with OH, followed by surface deposition (Hauglustaine et al., 1998; Granier et al., 2000). Because CO is destroyed through the troposphere, its concentrations are expected to decrease with altitude. In typical average background conditions during SOP 2a2, CO shows values of about 120±18 ppb in LT increasing to 200 ppb over the two main cities (Niamey, Cotonou) (Fig. 4). In upper layers (4–12 km), CO gradually decreases to 100±10 ppb.

Tropospheric O₃ is in situ produced in situ through NO_x, VOC and CO photochemistry and, to a lesser extent, due to stratospheric intrusions. The major O₃ sink is surface deposition. As a consequence, O₃ is expected to increase with altitude. During SOP 2a2, average LT O₃ concentrations show relatively little variation with values of 30±10 ppb. The lowest values (20 ppb) were detected above the forest region (7–11° N) where O₃ deposition is enhanced by vegetation. Highest values (70 ppb) were detected near the urban sites of Niamey and Cotonou. In upper layers (2–12 km), O₃ increases with altitude up to 60 ppb showing strong variability.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Water vapor as indicated by RH gives valuable information on cloud location. In average conditions, RH shows high values (70%) in LT particularly above the tropical forest influenced by the wet monsoon flow at latitudes south of 13° N. North of these latitudes, air masses are dryer (30%) due to the arid surface and the influence of the warm and dry Harmattan flow. With altitude, air mass layers get dryer up to 20% RH.

Non-convective conditions showing profiles as described above were encountered on 13 and 20 August 2006 (Fig. 4, black lines). Any shape deviation from average profiles can be considered as conditions perturbed by convection or other causes but concomitant variations of tracer vertical shapes are considered as a situation perturbed by convection (Fig. 4, red lines). A summary of the variation of convection indicators for each F-F20 flight is given in Table 3. Four cases were identified on 11, 14, 15 and 17 August 2006 corresponding to outflow explorations where the F-F20 flew in the active MCS. Air masses influenced by biomass burning intrusions have been also detected on the vertical profiles with high CO levels up to 180 ppb at 9 km on 19 August near Cotonou (Ancellet et al., 2009). These enhancements are associated with increased O₃ levels and lower RH showing that the use of a single tracer (e.g. CO) is not sufficient for convective case distinction.

Finally, the successful use of multiple chemical and physical indicators made possible a fine distinction between air masses in the cloud outflow perturbed by convection from those outside the outflow and not perturbed by convection (Table 3).

6 NMHC in the upper troposphere under convective and non-convective conditions

6.1 Evidence of deep convection impact on NMHC in the UT

To address impact of convective injections on NMHC distribution in the UT, we examined: (i) horizontal NMHC distribution, (ii) total NMHC content and (iii) surface NMHC emissions signature in the UT in convective vs. non convective conditions.

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(i) Horizontal NMHC distribution was examined in the UT along F-F20 MCS flight tracks (e.g. MCS of the 15 August 2006 in Fig. 5 for benzene, toluene and isoprene). NMHC observations are sparse and 10-min-integrated. We overlaid NMHC sampling points on CO, O₃ and RH data because of their higher measurement frequency (1 s). First, CO shows an important longitudinal positive gradient through the outflow with an increasing trend going up by 30%. Mean CO longitudinal gradient for the four MCS is 13±3 ppb.°E (e.g. 20 ppb.°E on the 15 August flight described in Fig. 5). Gradient calculations are based on observations on the 4 parallel legs made when exploring the MCS. Secondly, no O₃ gradient is observed in the outflow but O₃ concentrations are sensitive to cloud position. They are smaller inside the outflow (about 45 ppb) than outside it (about 60 ppb) because of the injection of low-O₃ and humid air masses from LT. The absence of trend in O₃ observations points out the dual nature of tropical convection. Convection impact on O₃ distribution almost shows a negative trend locally but fills the UT with O₃ precursors that will potentially lead to delayed O₃ production downwind the MCS (Folkens et al., 2002). Finally, RH values are also sensitive to MCS inflow and outflow position with higher values close to saturation inside the MCS (>80%). Compared to convection tracers, NMHC observations do not exhibit any meaningful trend. Values show great variability without any horizontal gradient.

(ii) While NMHC longitudinal distribution do not show a clear trend, total NMHC content is clearly affected by convection compared to non perturbed conditions. Box plots are reported for selected NMHC in Fig. 6 and sorted by non-convective, convective UT and LT observations. For all NMHC, mixing ratio distributions are more scattered in convective cases and similar to LT distribution. In non-convective cases, distribution is tighter. Average UT mixing ratios can double in convective outflow compared to non-perturbed air masses. Mixing ratios variability is different from one NMHC to another as well as from an MCS to another illustrating the heterogeneity of these systems. Nevertheless, NMHC concentrations are significantly enhanced by convective injection in the UT. Moreover these injections are dissimilar all over the convective cloud anvil.

(iii) Significant NMHC levels in convective UT are expected to come from surface

emissions. Since vegetation gradient and surface coverage are critical factors governing NMHC latitudinal composition (see Sect. 4.1), we investigated surface emissions influence on NMHC outflow composition. Satellite MCS tracking images provide information on the region where the MCS started to develop as well as its trajectory before being sampled. As MCS have westward motion, the latitude where they started to develop (called latitude zero here after) indicates the nature of the surface it passed through and is used as a proxy of emissions (Fig. 1). On 11 August, the MCS started to develop above Nigeria over a forested region and passed over Benin before being sampled around Niamey (FV48). On 14 August, the MCS started to develop east of Niamey above Sahel and moved westward to Niamey where it was explored (FV50). On 15 August, the MCS developed over a forested region in Benin and was explored south of Niamey (FV51). On 17 August, the MCS started to develop in the Air region north of Niamey above Sahel (FV53). Mean NMHC levels for isoprene and benzene in the MCS outflow are reported according to latitude zero in Fig. 7. Isoprene levels show exponential dependence on surface characteristics. Both MCS that developed over forested regions (FV48 and FV51) have isoprene content up to three times higher than the MCS that developed over the Sahel (FV50 and FV53). A different trend is observed for anthropogenic compounds such as benzene. Average benzene contents are rather homogeneous regarding the 4 MCS. Indeed, urban areas, which are benzene main sources in the domain, are limited and located at specific spots (Niamey 13°51 N–2.11° E, Cotonou 6.3° N–2.4° E, Lagos 6.4° N–3.4° E) while vegetation coverage follows a smooth north to south gradient. These observations highlight the influence of surface sources on the UT NMHC content mediated by convection.

The investigation of NMHC content in non-convective vs. convective conditions and its dependence on surface coverage reveals that deep convection increases NMHC content in the UT over West Africa.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

6.2 Photochemistry vs. transport regimes in West Africa UT for NMHC

Being reactive species, NMHC can be used as tracers of photochemical and transport processes governing trace gases variability. In particular, NMHC ratios with common sources and different atmospheric lifetimes provide further information on the degree of atmospheric processing in sampled air masses. NMHC ratios are then useful photochemical clocks to follow transport and ageing of air masses (Borbon et al., 2004).

We examined trans-2-pentene to benzene and toluene vs. benzene ratios from UT observations to differentiate regimes controlling NMHC distribution. Natural logarithms of trans-2-pentene/benzene vs. toluene/benzene ratios are plotted in Fig. 8. Trans-2-pentene is one of the most reactive measured NMHC (lifetime of few hours), toluene is of medium reactivity (lifetime of 2 days) and benzene is the least reactive NMHC (lifetime of 9 days). The least reactive NMHC (benzene) being at the denominator, the fresher air masses are, the greater the ratios are. Clearly, two sets of points are distinguished depending on non-convective situations (black) and convective situations (red). Each data set shows a linear relationship with correlation factor of 0.60 and 0.77, respectively. Data in non-convective situations are scattered along toluene/benzene axis and have a slope of 1.16. This agrees well with aged air masses composition with low trans-2-pentene mixing ratios close to DL after several days of photochemical processing. The scatter plot in the convective situation is more diffuse with greater data variability along toluene/benzene axis as well as along trans-2-pentene/benzene axis. Ratios display the highest ratios values and the slope exhibits a greater value (2.87) illustrating the freshness of sampled convective outflow enriched with reactive species such as trans-2-pentene. UT NMHC concentrations are thus governed by two distinct regimes that are affected by convective injections. Even with the complexity of the different processes in the upper tropospheric region, NMHC ratio enables a qualitative evaluation of the underlying photochemical processing and provides useful information on the air masses chemical and dynamical regimes.

ACPD

9, 20309–20347, 2009

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

6.3 Impact of deep convection on air mass reactivity

OH reactivity is used to evaluate the contribution of measured NMHC to photochemical activity. OH reactivity is also an indicator of the O₃ chemical production contribution. Here, estimated reactivity is based on NMHC measurements on board the ATR-42 and the F-F20. Individual NMHC reactivity ($R_{\text{OH},i}$) is calculated using Eq. (1):

$$R_{\text{OH},i} = k_{\text{OH},i} [\text{NMHC}_i] \quad (1)$$

where $k_{i,\text{OH}}$ is the OH reaction rate coefficients for every NMHC taken from Atkinson (2003); Atkinson et al. (2006) and calculated for LT and UT temperatures; $[\text{NMHC}_i]$ is the mixing ratio of the target observed NMHC.

Total NMHC reactivity ($R_{\text{OH},\text{total}}$) is derived from the sum of individual $R_{i,\text{OH}}$ using Eq. (2).

$$R_{\text{OH},\text{total}} = \sum R_{\text{OH},i} \quad (2)$$

While this approach gives a practical estimate of the contribution of measured NMHC to the atmospheric photochemistry; it does not explain the full atmospheric chemistry of the considered compounds.

UT air mass reactivity calculated for convective cases is compared to air masses reactivity for non-convective cases and for the LT (Fig. 10). In non-convective conditions, UT $R_{\text{OH},\text{total}}$ is $0.52 \pm 0.21 \text{ s}^{-1}$ while it raises in convective conditions up to $0.95 \pm 0.66 \text{ s}^{-1}$. Thus, air masses reactivity is enhanced by 40% in the UT due to injection of reactive NMHC. During convective events, UT $R_{\text{OH},\text{total}}$ is a similar value to the LT one which equals $0.82 \pm 0.53 \text{ s}^{-1}$.

The relative contribution of each NMHC, CO and O₃ to $R_{\text{OH},\text{total}}$ is evaluated according to altitude. The vertical profile of relative contributions to $R_{\text{OH},\text{total}}$ is reported in Fig. 9. Isoprene is the main reactant with OH at all altitudes, contributing up to half of the total reactivity (0.90 s^{-1} in LT (44%) and 1.00 s^{-1} in the UT (27%)). The sum of

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

other NMHC contributions is 30% in LT and 50% in the UT. As expected, O₃ and CO exhibit a negligible contribution of a few percent (<5%).

These results show that NMHC play an important role in air masses reactivity and UT photochemical potential which is expected to induce changes in ozone chemistry.

5 6.4 Fraction of low tropospheric air in fresh convective outflow

To obtain a quantitative evaluation on vertical transport impact and to assess to the extent to which the LT air affects the UT; we calculated the fraction f of LT air in the fresh convective outflow. To calculate f , we applied NMHC observations to Eq. (3) adapted from Bertram et al., 2007).

$$10 \quad f = \frac{[\text{UT}]_{\text{outflow}} - [\text{UT}]_{\text{bckgrd}}}{[\text{LT}] - [\text{UT}]_{\text{bckgrd}}} \quad (3)$$

where $[\text{UT}]_{\text{outflow}}$ is the mean mixing ratio of a selected specie in fresh convective outflow, $[\text{UT}]_{\text{bckgrd}}$ is the mean mixing ratio in non-convective UT and $[\text{LT}]$ is near-surface mean mixing ratio (below 2 km) according to the latitudinal gradient and MCS trajectory using “latitude zero” (cf. Sect. 6.1). The method presumes that photochemical loss is negligible during vertical transport, chemical lifetimes of species being longer or equivalent to vertical transport timescales (~30 min). First, we calculated f for each individual NMHC. Then we averaged f combining all the individual results for each of the 4 MCS (Fig. 11, left). The overall mean of f is about $40 \pm 15\%$ (Fig. 11). Results are consistent regarding the 4 MCS explored during SOP 2a2 as well as with other studies at mid-latitude (10 to 40% by Ray et al., 2004; 17% by Bertram et al., 2007) and in marine tropical regions (32 to 64% by Cohan et al., 1999). As expected, tropical deep convection above continent is more vigorous than at mid-latitude regions and weaker than marine tropical convection where larger humidity amounts are available.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

6.5 Vertical transport timescale during convective events

Vertical transport timescale estimation is based on an original approach using the photochemical processing equation with isoprene and benzene observations. Here, we assume that photochemical loss of isoprene is of similar timescale to that of a convective event. Therefore, isoprene (lifetime of few hours) is considered to be an efficient tracer of vertical transport. Consequently, a calculated time based on the photochemical processing equation can be attributed to the vertical pumping timescale. Depending on the reactivity of the species relative to OH, the photochemical age Δt can be determined by Eq. (4),

$$\Delta t = \frac{\ln \left[\frac{(\text{Isoprene/Benzene})_{\text{PBL}}}{(\text{Isoprene/Benzene})_{\text{UT}}} \right]}{(k_i - k_b) [\text{OH}]} \quad (4)$$

where [isoprene] and [benzene] are the mixing ratio of isoprene and benzene in LT and in the convective outflow, k_i and k_b are the OH reaction rate constants for isoprene and benzene, respectively ($k_i=1.0 \times 10^{-10} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$; $k_b=1.2 \times 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$; taken from Atkinson, 2003; Atkinson et al., 2006) and [OH] is OH radical mean concentration assumed to be $2 \times 10^6 \text{ molecules cm}^{-3}$ (Lawrence et al., 2003). Δt ranges from 12 to 36 min for the 4 MCS with an average of 25 ± 10 min (Fig. 11, right). Results are consistent with the expected timescale of vertical transport by deep convection (15–20 min) derived from typical vertical velocities ($5\text{--}15 \text{ m s}^{-1}$) (Fig. 11) and with modeling studies by Thompson et al. (1997) (30 min). The use of photochemical processing equation based on isoprene and benzene ratios is efficient to calculate vertical transport timescale in a convective system.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

7 Conclusions

Airborne measurements of tropospheric trace gases species collected by the two French aircrafts during AMMA SOP 2a2 in August 2006 are presented in this paper. These observations constitute a unique data set combining lower and upper tropospheric measurements over a large domain in West Africa. In particular, VOC (NMHC from C5 to C9) are reported from the boundary layer to the UT.

First, NMHC distributions were investigated. NMHC latitudinal distribution is closely connected to land surface characteristics over West Africa with an important north-to-south gradient going from bare soils over Sahel (North) to the tropical forest (South) with limited urban sites (Niamey and Cotonou). Biogenic compound concentrations (isoprene) exhibit an important north-to-south gradient following vegetation densification. Anthropogenic species concentrations are at maximum levels over urban sites and decrease over the forest. Vertical NMHC distributions show a “C-shaped” profile with significant UT concentrations due to convective injections and comparable to those observed near the surface, even for the most reactive NMHC.

Then, various diagnostics tools were set up and successfully applied to highlight the impact and characteristics of deep convection on NMHC. CO, O₃ and RH were used as simultaneous deep convection indicators to isolate convection impacted air masses. UT NMHC concentrations in WA appear to be controlled by two regimes: photochemistry and fast convective vertical transport. NMHC UT concentrations double when subject to convection and are dependent on MCS geographical origin, used as a proxy of surface type, indicating the vertical transport of near-surface air masses. As a consequence, UT total air masses reactivity regarding OH radical doubles in convective conditions (0.95 s⁻¹) compared to non-convective conditions (0.52 s⁻¹) and reaches values similar to LT reactivity (0.82 s⁻¹). Furthermore, isoprene plays an important role in total reactivity throughout the vertical tropospheric column, especially in convective outflow explaining 27% of UT total reactivity.

Finally, the fraction of LT air in convective outflow and vertical transport time scale

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

derived from NMHC observations were estimated to be $40\pm 15\%$ and 25 ± 10 min, respectively.

NMHC observations during AMMA reveal the potential impact of deep convection on UT chemistry. Even though ozone production was not directly observed because no flight was performed in aged outflow, post convective ozone formation is expected due to injection of reactive ozone precursors in the UT. Andrés-Hernández et al. (2009) show this ozone formation in MCS outflow during flights performed on-board the DLR-F20 within AMMA campaign and estimate O_3 production rates around 1 ppb h^{-1} in the MCS outflow.

Observations like those related in this paper are essential in order to assess the global impact of deep convection on ozone and other photooxidant production. Beyond characterizing local chemical effects, such observations build a database that can be used for relevant parameterization of mesoscale and global CTM models. In particular, deep convection spatial and temporal characteristics derived from NMHC observation in WA will be useful indicators for guiding model parameterizations.

Acknowledgement. This work is supported by the AMMA project. Based on a French initiative, AMMA was built by an international scientific group and is currently funded by a large number of agencies, especially from France, UK, US and Africa. It has been the beneficiary of a major financial contribution from the European Community's Sixth Framework Research Program. Detailed information on scientific coordination and funding is available on the AMMA International web site <http://www.amma-international.org>. We would like also to thank SAFIRE (Service des Avions Français Instrumentés pour la Recherche en Environnement) for deploying the research aircrafts.

References

Ancellet, G., Leclair de Bellevue, J., Mari, C., Nedelec, P., Kukui, A., Borbon, A., and Perros, P.: Effects of regional-scale and convective transports on tropospheric ozone chemistry revealed by aircraft observations during the wet season of the AMMA campaign, *Atmos. Chem. Phys.*, 9, 383–411, 2009, <http://www.atmos-chem-phys.net/9/383/2009/>.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Andrés-Hernández, M. D., Kartal, D., Reichert, L., Burrows, J. P., Meyer Arnek, J., et al.: Peroxy radical observations over West Africa during AMMA 2006: Photochemical activity in the outflow of convective systems, *Atmos. Chem. Phys.*, 9, 3681–3695, 2009, <http://www.atmos-chem-phys.net/9/3681/2009/>.

Atkinson, R.: Kinetics of the gas-phase reactions of OH radicals with alkanes and cycloalkanes, *Atmos. Chem. Phys.*, 3, 2233–2307, 2003, <http://www.atmos-chem-phys.net/3/2233/2003/>.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., et al.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, 2006, <http://www.atmos-chem-phys.net/6/3625/2006/>.

Bechara, J., Borbon, A., Jambert, C., and Perros, P. E.: New off-line aircraft instrumentation for non-methane hydrocarbon measurements, *Anal. Bioanal. Chem.*, 392, 865–876, 2008.

Bertram, T. H., Perring, A. E., Wooldridge, P. J., Crouse, J. D., Kwan, A. J., et al.: Direct measurements of the convective recycling of the upper troposphere, *Science*, 315, 816–820, 2007.

Borbon, A., Coddeville, P., Locoge, N., and Galloo, J. C.: Characterising sources and sinks of rural VOC in eastern France, *Chemosphere*, 57, 931–942, 2004.

Carter, W. P. L.: Development of ozone reactivity scales for volatile organic-compounds, *J. Air Waste Manag. Assoc.*, 44, 881–899, 1994.

Cohan, D. S., Schultz, M. G., Jacob, D. J., Heikes, B. G., and Blake, D. R.: Convective injection and photochemical decay of peroxides in the tropical upper troposphere: Methyl iodide as a tracer of marine convection, *J. Geophys. Res.-Atmos.*, 104(D4), 5717–5724, 1999.

Colomb, A., Williams, J., Crowley, J., Gros, V., Hofmann, R., et al.: Airborne measurements of trace organic species in the upper troposphere over Europe: the impact of deep convection, *Environ. Chem.*, 3, 244–259, 2006.

Dessler, A. E.: The effect of deep, tropical convection on the tropical tropopause layer, *J. Geophys. Res.*, 107(D3), 4033, doi:10.1029/2001JD000511, 2002.

Dickerson, R. R., Huffman, G. J., Luke, W. T., Nunnermacker, L. J., Pickering, K. E., et al.: Thunderstorms – an important mechanism in the transport of air-pollutants, *Science*, 235, 460–464, 1987.

Doherty, R. M., Stevenson, D. S., Collins, W. J., and Sanderson, M. G.: Influence of convective transport on tropospheric ozone and its precursors in a chemistry-climate model, *Atmos. Chem. Phys.*, 5, 3205–3218, 2005, <http://www.atmos-chem-phys.net/5/3205/2005/>.

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Fischer, H., de Reus, M., Traub, M., Williams, J., Lelieveld, J., et al.: Deep convective injection of boundary layer air into the lowermost stratosphere at midlatitudes, *Atmos. Chem. Phys.*, 3, 739–745, 2003, <http://www.atmos-chem-phys.net/3/739/2003/>.
- 5 Folkins, I., Braun, C., Thompson, A. M., and Witte, J.: Tropical ozone as an indicator of deep convection, *J. Geophys. Res.-Atmos.*, 107(D13), 4184, doi:10.1029/2001JD001178, 2002.
- Ganzeveld, L., Eerdekens, G., Feig, G., Fischer, H., Harder, H., et al.: Surface and boundary layer exchanges of volatile organic compounds, nitrogen oxides and ozone during the GABRIEL campaign, *Atmos. Chem. Phys.*, 8, 6223–6243, 2008,
10 <http://www.atmos-chem-phys.net/8/6223/2008/>.
- Granier, C., Pétron, G., Müller, J.-F., and Brasseur, G.: The impact of natural and anthropogenic hydrocarbons on the tropospheric budget of carbon monoxide, *Atmos. Environ.*, 34, 5255–5270, 2000.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates
15 of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 2006,
<http://www.atmos-chem-phys.net/6/3181/2006/>.
- Hao, W. M., Ward, D. E., Olbu, G., and Baker, S. P.: Emissions of CO₂, CO, and hydrocarbons from fires in diverse African savanna ecosystems, *J. Geophys. Res.*, 101(D19), 23577–
20 23584, 1996.
- Hauf, T., Schulte, P., Alheit, R., and Schlager, H.: Rapid vertical trace gas transport by an isolated midlatitude thunderstorm, *J. Geophys. Res.*, 100, 22957–22970, 1995.
- Hauglustaine, D. A., Brasseur, G. P., Walters, S., Rasch, P. J., Muller, J. F., Emmons, L. K., and Carroll, C. A.: MOZART, a global chemical transport model for ozone and related chemical
25 tracers 2. Model results and evaluation, *J. Geophys. Res.*, 103(D21), 28291–28335, 1998.
- Houze, R. A.: Mesoscale convective systems, *Rev. Geophys.*, 42, RG4003, doi:10.1029/2004RG000150, 2004.
- Karl, T. G., Christian, T. J., Yokelson, R. J., Artaxo, P., Hao, W. M., and Guenther, A.: The tropical forest and fire emissions experiment: method evaluation of volatile organic compound emissions measured by PTR-MS, FTIR, and GC from tropical biomass burning, *Atmos. Chem. Phys.*, 7, 5883–5897, 2007, <http://www.atmos-chem-phys.net/7/5883/2007/>.
- 30 Kesselmeier, J., Kuhn, U., Wolf, A., Andreae, M. O., Ciccioli, P., et al.: Atmospheric volatile organic compounds (VOC) at a remote tropical forest site in central Amazonia, *Atmos. Environ.*, 34, 4063–4072, 2000.

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Lafore, J. P., and Moncrieff, M. W.: A numerical investigation of the organization and interaction of the convective and stratiform regions of tropical squall lines, *J. Atmos. Sci.*, 46, 521–544, 1989.

5 Lawrence, M. G., and Salzmann, M.: On interpreting studies of tracer transport by deep cumulus convection and its effects on atmospheric chemistry, *Atmos. Chem. Phys.*, 8, 6037–6050, 2008, <http://www.atmos-chem-phys.net/8/6037/2008/>.

Lawrence, M. G., von Kuhlmann, R., Salzmann, M., and Rasch, P. J.: The balance of effects of deep convective mixing on tropospheric ozone, *Geophys. Res. Lett.*, 30, 1940, doi:10.1029/2003GL017644, 2003.

10 Lelieveld, J., and Crutzen, P. J.: Role of deep cloud convection in the ozone budget of the troposphere, *Science*, 264, 1759–1761, 1994.

Mari, C. H., Cailley, G., Corre, L., Saunois, M., Attie, J. L., Thouret, V., and Stohl, A.: Tracing biomass burning plumes from the Southern Hemisphere during the AMMA 2006 wet season experiment, *Atmos. Chem. Phys.*, 8, 3951–3961, 2008, <http://www.atmos-chem-phys.net/8/3951/2008/>.

15 Martin, R. V., Jacob, D. J., Logan, J. A., Bey, I., Yantosca, R. M., et al.: Interpretation of TOMS observations of tropical tropospheric ozone with a global model and in situ observations, *J. Geophys. Res.-Atmos.*, 107, 4351, doi:10.1029/2001JD001480, 2002.

20 Nedelec, P., Cammas, J. P., Thouret, V., Athier, G., Cousin, J. M., et al.: An improved infrared carbon monoxide analyser for routine measurements aboard commercial Airbus aircraft: technical validation and first scientific results of the MOZAIC III programme, *Atmos. Chem. Phys.*, 3, 1551–1564, 2003, <http://www.atmos-chem-phys.net/3/1551/2003/>.

Nicholson, S. E.: A revised picture of the structure of the “monsoon” and land ITCZ over West Africa, *Clim. Dynam.*, 32, 1155–1171, 2009.

25 Pickering, K. E., Dickerson, R. R., Huffman, G. J., Boatman, J. F., and Schanot, A.: Trace Gas Transport in the Vicinity of Frontal Convective Clouds, *J. Geophys. Res.*, 93(D1), 759–773, 1988.

Poisson, N., Kanakidou, M., and Crutzen, P. J.: Impact of non-methane hydrocarbons on tropospheric chemistry and the oxidizing power of the global troposphere: 3-dimensional modelling results, *J. Atmos. Chem.*, 36, 157–230, 2000.

30 Ray, E. A., Rosenlof, K. H., Richard, E. C., Hudson, P. K., Cziczo, D. J., et al.: Evidence of the effect of summertime midlatitude convection on the subtropical lower stratosphere from CRYSTAL-FACE tracer measurements, *J. Geophys. Res.-Atmos.*, 109, D18304,

doi:10.1029/2004JD004655, 2004.

Redelsperger, J. L., Thorncroft, C. D., Diedhiou, A., Lebel, T., Parker, D. J., and Polcher, J.: African monsoon multidisciplinary analysis – An international research project and field campaign, *B. Am. Meteor. Soc.*, 87, 1739–1746, 2006.

Reeves, C. E., Ancellet, G., Attie, J.-L., Bechara, J., Borbon, A., et al.: Chemical characterisation of the West Africa Monsoon during AMMA, *Atmos. Chem. Phys. Discuss.*, 9, 1585–1619, 2009, <http://www.atmos-chem-phys-discuss.net/9/1585/2009/>.

Saxton, J. E., Lewis, A. C., Kettlewell, J. H., Ozel, M. Z., Gogus, F., et al.: Isoprene and monoterpene measurements in a secondary forest in northern Benin, *Atmos. Chem. Phys.*, 7, 4095–4106, 2007, <http://www.atmos-chem-phys.net/7/4095/2007/>.

Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., et al.: Multi-model ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.-Atmos.*, 111(D8), D08301, doi:10.1029/2005JD006338, 2006.

Stickler, A., Fischer, H., Bozem, H., Gurk, C., Schiller, C., et al.: Chemistry, transport and dry deposition of trace gases in the boundary layer over the tropical Atlantic Ocean and the Guyanas during the GABRIEL field campaign, *Atmos. Chem. Phys.*, 7, 3933–3956, 2007, <http://www.atmos-chem-phys.net/7/3933/2007/>.

Stickler, A., Fischer, H., Williams, J., de Reus, M., Sander, R., et al.: Influence of summertime deep convection on formaldehyde in the middle and upper troposphere over Europe, *J. Geophys. Res.-Atmos.*, 111, D14308, doi:10.1029/2005JD007001, 2006.

Ström, J., Fischer, H., Lelieveld, J., and Schröder, F.: In situ measurements of microphysical properties and trace gases in two cumulonimbus anvils over western Europe, *J. Geophys. Res.*, 104, 12221–12226, 1999.

Thompson, A. M., Tao, W. K., Pickering, K. E., Scala, J. R., and Simpson, J.: Tropical deep convection and ozone formation, *B. Am. Meteor. Soc.*, 78, 1043–1054, 1997.

Warneke, C., Holzinger, R., Hansel, A., Jordan, A., Lindinger, W., et al.: Isoprene and its oxidation products methyl vinyl ketone, methacrolein, and isoprene related peroxides measured online over the tropical rain forest of Surinam in March 1998, *J. Atmos. Chem.*, 38, 167–185, 2001.

Zipsper, E. J., Cecil, D. J., Liu, C. T., Nesbitt, S. W., and Yorty, D. P.: Where are the most intense thunderstorms on earth?, *B. Am. Meteor. Soc.*, 87, 1057–1071, 2006.

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Details on ATR-42 and F-F20.

Date	ATR			F-F20		
	Flight no.	Objective	No. of VOC samples collected	Flight no.	Objective	No. of VOC samples collected
2 August 2006	AV38	Test flight	11			
	AV39	Test flight				
4 August 2006	AV40	Test flight				
5 August 2006	AV41	Pre-MCS				
6 August 2006	AV42	Post-MCS	6			
7 August 2006	AV43	Post-MCS				
8 August 2006	AV44	North-to-south exploration	6			
	AV45	North-to-south exploration	6			
9 August 2006	AV46	North-to-south exploration	3			
	AV47	North-to-south exploration	3			
11 August 2006	AV48	North-to-south exploration		FV48	MCS outflow exploration	4
	AV49	North-to-south exploration				
12 August 2006	AV50	Pre-MCS	4			
13 August 2006	AV51	North-to-south exploration	8	FV49	North-to-south exploration	12
14 August 2006	AV52	Pre-MCS	3	FV50	MCS outflow exploration	7
15 August 2006	AV53	Post-MCS	7	FV51	MCS outflow exploration	4
16 August 2006	AV54	North-to-south exploration	10			
17 August 2006	AV56	North-to-south exploration	10	FV53	MCS outflow exploration	7
17 August 2006	AV57	Post-MCS				
19 August 2006	AV58	Post-MCS	12	FV54	North-to-south exploration	4
				FV55	Post day +1 exploration	7
20 August 2006				FV56	North-to-south exploration	9
21 August 2006				FV57	Heat Low exploration	10

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Table 2. Aircraft instrumentation.

Compound	Platform	Instrument	Measurement technique	Time response	Detection limit	Precision	Accuracy	Reference
O ₃	ATR-42	Thermo-Electron TEI 49	UV absorption	3 s	1 ppb	1 ppb		SAFIRE http://www.safire.fr
	F-F20	MOZART Instrument		4 s	2 ppb	2 ppb	2%	
CO	ATR-42	Thermo-Electron 48CS	IR absorption	5 s	20 ppb	20 ppb		SAFIRE http://www.safire.fr Nedelec et al., 2003
	F-F20	MOZART Instrument		30 s	5 ppb	5 ppb	5%	
NMHC	ATR-42/ F-F20	AMOVOC/ TDAS/GC-MS	Solid sorbent cartridges	10 min	1–5 ppt	8–18%	1–5%	Bechara et al., 2008

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.

Table 3. Convection indicators variations for F-F20 flights. ↓: decreasing mixing ratios, ↑: increasing mixing ratios.

Date	Flight no.	Convection indicators		
		O ₃	CO	RH
11 August 2006	FV48	~ ↓	↑	↑
13 August 2006	FV49	↑	↓	↓
14 August 2006	FV50	↓	↑	↑
15 August 2006	FV51	~ ↓	↑	↑
17 August 2006	FV53	↓	~ ↑	↑
19 August 2006	FV54	↑	↑	↓
19 August 2006	FV55	↑	↓	↓
20 August 2006	FV56	↑	↓	↓
21 August 2006	FV57	↑	↓	↓

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.




Fig. 1. Explored area during AMMA SOP 2a2. The map reflects the land-surface coverage. Light yellow indicate sahelian type surface. Orange denotes semi-desert region. Light green indicates savanna. Dark green stands for tropical forest. Flights tracks performed by the F-F20 are displayed. The round markers are the locations where each explored MCS started to develop (called latitude zero). The diamond markers are the position where the MCS was explored. The arrows indicate the trajectory of the MCS.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.




Fig. 2. North-to-South variability profiles of biogenic and anthropogenic NMHC in the low troposphere (<2 km). Samples are collected on-board the ATR-42. top panel shows isoprene, bottom left panel shows trans-2-pentene and down right shows benzene. Open markers denote all observations. Solid markers represent the average value in each latitudinal bins of 1° latitude. Grey area represents standard deviations for all observations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.




Fig. 3. Vertical profiles of a selection of NMHC collected on both aircrafts during SOP 2a2. Open markers are all observations. Black markers are average values in altitude bins of 1 km. Grey area represents standard deviations for all observations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.




Fig. 4. Vertical profiles of CO, O₃ and RH used as convection indicators. Black lines (13 August flight) are typical variation in non-convective conditions. Red lines (14 August flight) are the vertical variations in convective conditions.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.




Fig. 5. Latitude vs. longitude distribution at UT level for CO (top left panel), O_3 (top right panel), RH (bottom left panel) in full line, and benzene (top left panel), isoprene (top right panel) and toluene (bottom left panel) in squares. Flight track is superimposed on the Meteosat satellite picture at 14:00 UTC in bottom right panel. The flight took place on 15 August between 12:35 UTC and 16:57 UTC.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.




Fig. 6. Selected NMHC variability in the low troposphere (LT), in non-convective UT (NC-UT) and in convective UT (C-UT). Lower and upper limits of the box correspond to 25% and 75% percentiles, bottom and top whiskers the 10% and 90%. Middle bars indicate the median and black dots the mean.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.




Fig. 7. Average UT content of isoprene (round marker) and benzene (squared marker) according to latitude zero. Bars indicate standard deviations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.





Fig. 8. Natural logarithms of trans-2-pentene to benzene vs. toluene to benzene ratios for F-F20 observations in the UT. Concentrations ratios are shown in black markers in non-convective conditions and in red markers during convective conditions. Lines show linear fits for each data set.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.

**Fig. 9.** Vertical profiles of O₃, CO and NMHC contribution to total reactivity R_{OH} .[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evidence of the
impact of deep
convection on
reactive VOC**

J. Bechara et al.




Fig. 10. Total NMHC reactivity according to OH radical in the boundary layer (LT, in red) and in the UT during non-convective (NC.UT in orange) and convective (C.UT, in yellow) situations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evidence of the impact of deep convection on reactive VOC

J. Bechara et al.




Fig. 11. Deep convection characteristics in West Africa. Left plot: Fraction of low tropospheric air in fresh convective outflow for each MCS. Vertical lines are error bars, mean fraction is in red, transparent grey area is standard deviation and dashed lines are minimum and maximum literature data comparison. *: Ray et al., 2004 in mid-latitude convection, **: Cohan et al., 1999 in marine tropical convection. Right plot: Vertical transport timescale for each MCS and theoretical time “Calc.” calculated considering a vertical speed of 15 m s^{-1} to reach 12 km-altitude. Red line indicates mean time, transparent grey area is the standard deviation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion