

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Biomass burning and
tropospheric ozone
during 2006 El Niño**

S. Chandra et al.

Effects of the 2006 El Niño on tropospheric ozone and carbon monoxide: implications for dynamics and biomass burning

S. Chandra^{1,2}, J. R. Ziemke^{1,2}, B. N. Duncan², T. L. Diehl^{1,2}, N. J. Livesey³, and
L. Froidevaux³

¹Goddard Earth Sciences and Technology Center, University of Maryland Baltimore County,
Baltimore, Maryland, USA

²NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³NASA Jet Propulsion Laboratory, Pasadena, California, USA

Received: 24 September 2008 – Accepted: 17 December 2008 – Published: 28 January 2009

Correspondence to: J. R. Ziemke (jerald.r.ziemke@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We have studied the effects of the 2006 El Niño on tropospheric O₃ and CO at tropical and sub-tropical latitudes measured from the OMI and MLS instruments on the Aura satellite. The 2006 El Niño-induced drought allowed forest fires set to clear land to burn out of control during October and November in the Indonesian region. The effects of these fires are clearly seen in the enhancement of CO concentration measured from the MLS instrument. We have used a global model of atmospheric chemistry and transport (GMI CTM) to quantify the relative importance of biomass burning and large scale transport in producing observed changes in tropospheric O₃ and CO. The model results show that during October and November both biomass burning and meteorological changes contributed almost equally to the observed increase in tropospheric O₃ in the Indonesian region. The biomass component was 4–6 DU but it was limited to the Indonesian region where the fires were most intense. The dynamical component was 4–8 DU but it covered a much larger area in the Indian Ocean extending from South East Asia in the north to western Australia in the south. By December 2006, the effect of biomass burning was reduced to zero and the observed changes in tropospheric O₃ were mostly due to dynamical effects. The model results show an increase of 2–3% in the global burden of tropospheric ozone. In comparison, the global burden of CO increased by 8–12%.

1 Introduction

El Niño and La Niña events are major sources of inter-annual and decadal variability in tropical tropospheric ozone (O₃) (e.g., Ziemke et al., 2003). The atmospheric effects of El Niño events are generally a change in convection in the tropical troposphere associated with an eastward shift of the warm SST anomaly and large scale Walker circulation. This shift results in an increase in tropospheric column O₃ in the western Pacific and a decrease in the eastern Pacific relative to non-El Niño years. The ef-

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fect of El Niño on specific humidity is usually opposite to that of column ozone (e.g., Chandra et al., 1998, 2007). During La Niña years, dynamical processes are largely reversed which results in a decrease (increase) of column ozone (specific humidity) in the western Pacific. The effects of El Niño on tropospheric composition have been extensively studied from both satellite and ground based measurements (e.g., Chandra et al., 1998; Fujiwara et al., 1999; Thompson et al., 2001) and by using global models of atmospheric chemistry and transport (e.g., Peters et al., 2001; Sudo and Takahashi, 2001; Chandra et al., 2002, 2003; Duncan et al., 2003; Zeng and Pyle, 2005; Doherty et al., 2006).

Recently Logan et al. (2008) studied the effects of the 2004 and 2006 El Niño events on tropospheric profiles of CO, O₃, and H₂O measured from the Tropospheric Emission Spectrometer (TES) on the Aura satellite. Their findings were generally consistent with the observed characteristics of O₃ and H₂O inferred from previous El Niño events. The changes during the 2004 El Niño in tropospheric O₃ and H₂O from TES are similar to those inferred from Ozone Monitoring Instrument (OMI) and Microwave Limb Spectrometer (MLS) flown on the same satellite (Chandra et al., 2007). The 2004 El Niño was a weak event compared to 2006 and significantly weaker compared to the El Niño of 1997–1998. This was reflected in the TCO (tropospheric column ozone) anomaly over the Indonesian region in the western Pacific as shown in Table 1 of Logan et al. (2008). For example, the TCO anomaly in November 1997 inferred from EP TOMS was 14.4 Dobson Units (DU; 1 DU=2.69×10²⁰ molecules-m⁻²). The corresponding values from TES for November 2004 and 2006 were 6.6 DU and 11.1 DU. Logan et al. (2008) attributed most of these changes to differences in the magnitude of CO emissions from fires in Indonesia which in these years were estimated to be 193 Tg, 24 Tg, and 82 Tg, respectively. They also suggested significant contribution from NO_x production due to lightning in late November and December 2006 when CO production due to large scale forest fires decreased significantly. The effects of forest fires in the Indonesian region during the 1997 El Niño was studied by Chandra et al. (2002) using the GEOS-CHEM global model of chemistry and transport. Their study

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Biomass burning and
tropospheric ozone
during 2006 El Niño**

S. Chandra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

suggested that about half of the increase in tropospheric column ozone in the Indonesian region was due to biomass burning and the other half due to dynamical effects. A similar conclusion was arrived by Sudo and Takahashi (2001) from their model study. Their conclusion about the contribution of biomass burning was indirect since they did not explicitly include the effect of biomass burning in their model study.

The combined OMI and MLS instruments on the Aura satellite have been providing near global measurements of TCO from August 2004 to present (e.g., Ziemke et al., 2006; Schoeberl et al., 2007) as discussed in Sect. 2. In addition the MLS instrument has been providing daily measurements of CO at several pressure levels in the troposphere and stratosphere (Livesey et al., 2008). The purpose of this paper is to study the effects of the 2006 El Niño on tropospheric composition measured from the OMI and MLS instruments on the Aura satellite and evaluate specific roles of the various processes using a global model of chemistry and transport. The global model used in this study is the NASA GMI CTM (Global Modeling Initiative Chemical Transport Model). It successfully simulates a wide range of observations of chemical constituents in the troposphere and stratosphere, including data collected by instruments on board satellites. We provide a brief overview of the relevant tropospheric processes in the GMI CTM in Sect. 3. In our study, the El Niño related changes in O₃ and CO fields based on OMI and MLS measurements are compared with corresponding changes based on GMI CTM simulation to delineate the relative importance of biomass burning and large scale transport.

In the following, Sect. 2 discusses the satellite data, Sect. 3 the GMI model, and Sect. 4 and 5 compare observed and measured O₃ and CO, respectively. Section 6 discusses the impact of biomass burning emissions over Indonesia during the 2006 El Niño, and Sect. 7 compares the strength and frequency of the 2006 and recent El Niño events with previous El Niño events. Finally, Sect. 8 provides a summary.

2 Measurements from OMI and MLS

OMI is one out of a total of four instruments onboard the Aura spacecraft which is flown in a sun-synchronous polar orbit at 705 km altitude with a 98.2° inclination. The spacecraft has an equatorial crossing time of 01:45 p.m. (ascending node) with around 98.8 min per orbit (14.6 orbits per day on average). OMI is a nadir-scanning instrument that for visible (350–500 nm) and UV wavelength channels (UV-1: 270–314 nm; UV-2: 306–380 nm) detects backscattered solar radiance to measure column ozone with near global coverage (aside from polar night latitudes) over the Earth with a resolution of 13 km×24 km at nadir. Besides ozone OMI also measures cloud-top pressure, aerosols and aerosol parameters, NO₂, SO₂, and other trace constituents in the troposphere and stratosphere (Levelt et al., 2006). Total ozone from OMI is derived from the TOMS version 8 algorithm. A description of this algorithm may be obtained from the TOMS V8 CD DVD ROM, or from the OMI Algorithm Theoretical Basis Document (ATBD) (from the webpage http://toms.gsfc.nasa.gov/version8/v8toms_atbd.pdf).

There are two standard OMI total ozone products. Here we use data from the collection 3 OMTO3 version 8.5 (v8.5) product that is based on the Total Ozone Mapping Spectrometer (TOMS) version 8 (v8) total O₃ algorithm. A description of this algorithm may be obtained from the TOMS v8 DVD or the OMI Algorithm Theoretical Basis Document (ATBD) from the TOMS web page http://toms.gsfc.nasa.gov/version8/v8toms_atbd.pdf. A main difference between v8 and v8.5 is that v8.5 uses measured cloud pressure whereas v8 and earlier versions used an infrared-measured climatology for cloud pressure. Another OMI algorithm based on the Differential Optical Absorption Spectroscopy Method (DOAS) technique gives similar estimates of total column ozone (Kroon et al., 2008b). These total column O₃ retrievals have been compared with ground-based data (McPeters et al., 2008) and aircraft-based measurements (Kroon et al., 2008a).

O₃ profiles from the Aura Microwave Limb Sounder (MLS) are used to estimate stratospheric column O₃ (SCO). The SCO is subtracted from OMI total column O₃

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

from OMI to yield tropospheric column O_3 and mean volume mixing ratio (e.g., Ziemke et al., 2006). The mean volume mixing ratio (units ppbv) is determined by taking tropospheric column O_3 (in Dobson Units, DU) and dividing this by pressure difference (in units hPa) between surface and tropopause, and then multiplying this by the factor 1.27. Validation of MLS v2.2 measurements is discussed by Froidevaux et al. (2008) and Livesey et al. (2008). The measurements of mean tropospheric O_3 from OMI/MLS and upper tropospheric CO from MLS are studied and compared with simulations from a global chemical transport model (discussed next).

3 NASA Global Modeling Initiative Chemical Transport Model

The GMI CTM is described and evaluated in Ziemke et al. (2006), Schoeberl et al. (2006), Strahan et al. (2007) and Duncan et al. (2007, 2008). The chemical mechanism includes 117 species, 322 chemical reactions, and 81 photolysis reactions to simulate tropospheric and stratospheric chemistry. It includes a detailed description of O_3 - NO_x -hydrocarbon chemistry, updated with recent experimental data. The chemical mass balance equations are integrated using the SMVGEAR II algorithm (Jacobson, 1995). Photolysis frequencies are computed using the Fast-JX radiative transfer algorithm (M. Prather, personal communication). The algorithm treats both Rayleigh scattering as well as Mie scattering by clouds and aerosol. The CTM simulates the radiative and heterogeneous chemical effects of sulfate, dust, sea-salt, organic carbon and black carbon aerosol on tropospheric photochemistry.

Biogenic emissions of isoprene and monoterpenes are calculated on-line as described in Guenther et al. (2006). The lightning NO_x parameterization fits the relationship between model-calculated convective mass fluxes and observed cloud-to-ground flash rates (D. Allen and K. Pickering, personal communication). Time-appropriate anthropogenic and biomass burning emissions include, surface emissions from industry/fossil fuel, biomass burning, biogenic and biofuel combustions and contributions from aircraft emissions. They are based on the Global Fire Emission Database, ver-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sion 2 (GFEDv2) and discussed in the GMI CTM papers listed above. Monthly total biomass burning of CO and NO_x are shown in Table 1 for the later part of 2006 when the Indonesian fires were most intense.

Table 1 indicates that the Indonesian fires increased the CO burden by 69 Tg during the three month period from September to November 2006. It is about 27% of the global CO burden (251 Tg) during the same period. The increase in NO_x due to Indonesian fires during the same period was 1.19 Tg compared to the global burden of about 19.4 Tg.

The meteorological fields that drive transport are from the Goddard Modeling and Assimilation Office (GMAO) GEOS-4 data assimilation system (GEOS-4-DAS) (Bloom et al., 2005). Features of the circulation, such as anticyclones, in the GMI CTM are realistically represented in the simulations because of the data constraints on meteorological analyses. The GEOS-4-DAS fields have been regrided to 42 vertical levels with a lid at 0.01 hPa. The horizontal resolution is 2.5° × 2° (longitude by latitude).

A validation of the global model used in this study is given in the Appendix. This analysis compares the zonal and seasonal variability of TCO derived from the model with observed variability from OMI/MLS.

4 Model comparison of El Niño related changes in O₃ with satellite measurements

As in Chandra et al. (2007) and Logan et al. (2008), we have chosen October, November and December 2005 as a baseline for estimating El Niño-related changes in these months in the preceding and the following year. Year 2005 was a neutral year from an El Niño perspective. Maps of tropospheric O₃ mean mixing ratio are shown in Fig. 1a for October 2004, 2005, and 2006. They show relative changes in ozone in the El Niño years (upper and lower panels) with respect to a non El Niño year (middle panel). All three panels show regions of minimum values near the dateline. However, both the upper and the lower panels show a greater eastward shift across the dateline which

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 results in an increase of about 6–15 ppbv (~ 4 –10 DU) over the western Pacific region and a decrease of smaller magnitude east of the dateline. The increase in O_3 mean mixing ratio is larger in October 2006 (lower panel) which is consistent with a relatively stronger El Niño event in 2006 compared to 2004. The model (Fig. 1b) captures most of the zonal characteristics of TCO shown in Fig. 1a, but it tends to underestimate the observed values by 5–10 ppbv over most of the region between ± 10 deg. (Fig. 1c). Outside this region the sign is reversed and the model is biased higher by about the same magnitude. These pattern shifts in the model and observation are comparable in November and December months (not shown) and are similar to those shown in Fig. 2 of Ziemke et al. (2000) for the 1997 El Niño event.

10 The El Niño related changes in ozone mean mixing ratio in 2006 are clearly discernible in Fig. 2a, b and c. These figures show differences in OMI/MLS O_3 (top panels) and the model O_3 (bottom panels) between years 2006 and 2005 for the months of October-December. The model and observed values are generally in good agreement except in the western Pacific region in October (Fig. 2a) where the model values are biased higher. This bias is also reflected in CO as indicated in Sect. 5 and is probably due to overestimation of the effect of biomass burning in the Indonesian region in October 2006. In general, observations and model indicate excellent agreement in characterizing both broad and small scale features including the dipole nature of ozone anomaly in the tropical Pacific region (Fig. 2a and c). The large negative anomaly over eastern Africa/Indian Ocean in December 2006 seen in the data (Fig. 2c) is a manifestation of dynamically induced changes and is well produced in the model. We have analyzed the model results to estimate the contributions of the upper (above 500 hPa) and the lower troposphere in producing the El Niño related changes in tropospheric ozone. In general, the contributions from the upper and the lower troposphere are almost equal. However, the large negative anomaly in December in the eastern Africa/Indian Ocean (Fig. 2c) comes mostly from the upper troposphere.

Biomass burning and tropospheric ozone during 2006 El NiñoS. Chandra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 Model comparison of CO with satellite measurements

The El Niño events of 2004 and 2006 caused droughts in the Indonesian region which allowed set fires to spread uncontrolled in both October and November of these years. The MLS measurements show a relatively larger increase in CO in 2006 as compared to 2004 (figure not shown). This is consistent with CO measurements from TES (Logan et al., 2008). The model results show CO to be well mixed in the free troposphere even though convection was greatly reduced in this region. During October 2006, the model shows an increase of about 200 ppbv in the Indonesian region over much of the free troposphere. Figure 3a–c show differences in upper tropospheric CO from MLS (top panels) and the model (bottom panels) between 2006 and 2005 for October–December. In October, both the model and measurements show elevated values of CO in 2006 over the Indonesian region in the western Pacific. One month later in November (Fig. 3b) upper tropospheric CO from both MLS and the model are greatly reduced. They become negligible by December as the fire subsides in this region. CO for both measurement and model exhibit very similar spatial and temporal patterns except in October 2006 (Fig. 3a). The model shows a sharp horizontal gradient with CO mixing ratio exceeding 200 ppbv in the Indonesian region. The MLS values of CO and horizontal gradients are significantly smaller, largely due to the coarser spatial resolution of MLS nadir measurements which were interpolated horizontally between orbital tracks. It is interesting to note that the large increase in CO in 2006 during October–November nearly disappears in December, yet O₃ in December in the western Pacific remains elevated (Fig. 2c). Logan et al. (2008) attribute some of these increases in O₃ due to lightning NO_x in the Indonesian region.

6 Impact of biomass burning during the 2006 El Niño

In Sects. 4 and 5 we analyzed the El Niño related changes in tropospheric O₃ and CO in 2006 using both the model and observations. In this section we analyze the

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



model results to estimate the relative importance of biomass burning and large scale transport in producing the El Niño related changes in tropospheric O₃. Our approach is similar to the one used by Chandra et al. (2002) in analyzing the effect of the 1997 El Niño on tropospheric O₃. The model was run in two modes: The first mode explicitly included the NO_x and CO emission rates associated with the Indonesian fires as given in Table 1. The results of these runs are discussed in Sects. 4 and 5. In the second mode, the model was run by excluding the contributions from the Indonesian fires. The model runs in this mode are used to assess the effects of large scale transport on tropospheric O₃ as shown in the bottom panels of Fig. 4a–c. The difference of the two runs gives the contribution from biomass burning as shown in the upper panels of Fig. 4a–c.

In October 2006 the increase in O₃ due to biomass burning is about 4–6 DU (Fig. 4a, upper panel). This increase is limited to a small region comprising the islands of Sumatra, Java and Borneo in the Indonesian region. It is comparable to 4–8 DU increase caused by changes in meteorological conditions (Fig. 4b, lower panel). The latter, however, extends over a much larger area in the Indian Ocean encompassing regions of the Indian subcontinent and south-east Asia in the north, to western Australia in the south. These patterns are essentially similar in November 2006 (Fig. 4b). In December, 2006, the O₃ increase due to biomass burning is reduced to zero and the dynamical component of the O₃ field is identical to Fig. 2c (lower panel).

Table 2 shows the model simulations of monthly averaged global tropospheric burden of CO, O₃ and NO_x with and without Indonesian fire emissions. The table shows that during October 2006 the Indonesian fires contributed about 36 Tg which was about 10.7% of the global budget of CO. The increase in CO during November and December were respectively 12.6% and 8.1% of the global budget. These changes are significant and suggest that the El Niño related fires in the Indonesian region may have contributed significantly to global pollution. In comparison the impact of fire was relatively small (<3%) on the global O₃ burden. Logan et al. (2008) have suggested NO_x production due to lightning as a contributing factor to the positive anomaly in TCO dur-

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing December 2006. The NO_x production due to lightning is implicitly included in the model calculations. In the tropical Pacific region (12°S – 12°N , 90°E – 170°E), lightning produced a total of 0.032 Tg NO in October 2006. This value increased to 0.0492 Tg NO in November and 0.0562 Tg NO in December 2006.

5 6.1 The El Niño of 2006: a long-term perspective

Though the main focus of this study is the El Niño event of 2006, such events are very frequent and may have significant impact on global pollution. According to the World Meteorological Organization (WMO) an El Niño (La Niña) event is identified when sea surface temperature (SST) in the Niño 3.4 region (5°S – 5°N , 120°W – 170°W) is at least 0.5°C above (below) normal when averaged over three consecutive months. The relation between ΔSST and ΔTCO (the difference in TCO between eastern and western Pacific) is shown in Fig. 5. The ΔTCO time series is derived by subtracting TCO in the eastern Pacific (15°S – 15°N , 135°W – 180°) from TCO in the western Pacific (15°S – 15°N , 95°E – 140°E). The time series is deseasonalized and smoothed with a three-month running average to be compatible with Niño 3.4 SST. TCO is derived from 15 29 years (1979–2008) of total column ozone measurements from Nimbus 7 TOMS, Earth Probe TOMS and OMI using the convective cloud differential (CCD) method (e.g., Ziemke et al., 2005). Figure 5 shows a very robust correlation between ΔSST and ΔTCO . Even though the 1997 El Niño produced largest perturbations in SST and 20 TCO in the tropical Pacific, El Niño events like 2006 are more common. They are also episodic. Before 1997 they occurred about every 3–4 years, but more recently they have occurred every 2 years. A frequent occurrence of El Niño, similar to 2006, has the potential of increasing the global pollution triggered by the dryness and the resulting forest fires in the Indonesian region.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



7 Summary and conclusion

We have studied the effects of the 2006 El Niño on tropospheric O₃, and CO at tropical and sub-tropical latitudes measured from the OMI and MLS instruments on the Aura satellite. The zonal characteristics of observed changes in these constituents are similar to those reported by Logan et al. (2008) based on measurements of these constituents from the TES instrument on the same satellite. During October and December 2006, both these studies revealed a dipole like structure in TCO in the tropics with an increase over the western Pacific region and a decrease over the eastern Pacific region. The dipole structure was weaker during November 2006.

The 2006 El Niño-induced drought allowed forest fires to spread rapidly during October and November in the Indonesian region. The effects of these fires are clearly seen in the enhancement of CO concentration measured from both the MLS and TES instruments. We have used a global model of atmospheric chemistry and transport (GMI CTM) to quantify the relative importance of biomass burning and large scale transport in producing observed changes in TCO. The model results show that during October and November both biomass burning and meteorological changes contributed almost equally to increases in TCO in the Indonesian region. The biomass component was 4–6 DU but it was limited to the Indonesian region where the fires were most intense. The dynamical component was 4–8 DU but it covered a much larger area in the western Pacific and Indian Ocean extending from South East Asia in the north to western Australia in the south. During December 2006 the effect of biomass burning was reduced to zero and the observed changes in TCO were mostly due to dynamical effects. Chandra et al. (2002) have obtained similar results for the 1997 El Niño using ozone measurements from Earth Probe TOMS and comparing them with the GEOS-CHEM model. The 1997 El Niño was significantly stronger than the 2006 El Niño and caused greater perturbations in TCO due to both forest fires and large scale transport. For the 2006 El Niño the model shows an increase of 2–3% in the global burden of tropospheric ozone. In comparison, the global burden of CO increased by 8–12%.

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This study shows that the 2006 El Niño can be characterized from long records of tropospheric O₃ and sea surface temperature as a moderate event, yet still the contribution to pollution from biomass burning emissions over Indonesia was substantial, both locally and also when evaluated on a global basis. In recent years El Niño events have occurred with greater frequency than in previous years dating back to 1979. Both the frequency and scale of emissions from biomass burning suggests that even relatively moderate El Niño events can be an important source of pollution in the troposphere.

Appendix A

Comparisons between OMI/MLS and model O₃

Ziemke et al. (2006) compared tropospheric ozone variability derived from OMI/MLS and the GMI CTM using earlier versions of both the model and measurements. As discussed in Sect. 2, the total ozone data used in this paper is based on the TOMS version 8.5 algorithm instead of version 8 as used by Ziemke et al. (2006). The model also uses different emission sources and meteorological fields. In the following we compare the temporal/seasonal and zonal variability of TCO fields derived from the model and the measurements under varying conditions.

Figure A1 compares the temporal and seasonal variations of the model and observations for three separate regions: American (left four panels), African (middle four panels), and Indonesian (right four panels). In each region latitudes extend from north to south with the top panel being northernmost. In northern latitudes tropospheric O₃ for the model and measurement maximizes around March–July. In southern latitudes largest amounts occur in September–November. In between at low latitudes the seasonal variability is small in comparison. In general the temporal variability in model and measurements are remarkably similar indicating the ability of the model to capture month-by-month variability in the data. All three top panels in Fig. A1 for the latitude band 30° N–35° N show offset differences of 5–10 DU (with GMI larger than OMI/MLS).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For the upper right panel in Fig. A1 (corresponding to east China) the model and measured O_3 in summer are comparable but the model values are larger than measured values by 5–10 DU in winter and spring months when STE is greatest.

Martin et al. (2002) and Chandra et al. (2002) compared the seasonal variability in tropospheric O_3 from the GEOS-Chem model and TOMS CCD measurements. A large discrepancy was found between the model and the CCD measurements over low latitudes in northern Africa including Abidjan (5° N, 4° W). The model indicated maximum tropospheric column O_3 around December–February during peak time of the biomass burning whereas TOMS showed greatest amounts around June–October at a time when there is little or no biomass burning in northern Africa. Our analysis shows that the OMI measurements agree much better with model than the TOMS measurements pertaining to this seasonal variability over Africa. The GMI model indicates that enhancements in tropospheric O_3 over northern Africa from the biomass burning lie almost entirely below 500 hPa and are highly localized to the relatively small region of burning emissions.

Figure A2 shows statistical comparisons of model and observations. The three panels in this figure show their zonal correlations (top panel), mean zonal offset difference (middle panel) and mean zonal RMS (lower panel). The grey shading in the top panel represents areas that are not statistically significant at 99.9% confidence level. Figure A2 shows that the positive correlation between the model and OMI/MLS TCO, extending out to mid-latitudes, drops to negative values in higher latitudes. Several factors could contribute to near-zero or negative cross-correlations. Aside from known model uncertainties which include STE, OMI retrievals become less sensitive in measuring tropospheric O_3 in high latitudes (i.e., high solar zenith angles) and have reduced sensitivity in measuring O_3 in the lower troposphere, especially for high solar zenith angles and large slant columns. The model and observations compare the best in the tropics and subtropics extending out to latitudes of about $\pm 30^\circ$ to $\pm 35^\circ$. This is the latitude range chosen for GMI and OMI/MLS measurements for studying the 2006 El Niño event.

Biomass burning and tropospheric ozone during 2006 El NiñoS. Chandra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Acknowledgements. The authors thank the Aura OMI and MLS instrument and algorithm teams for the extensive satellite measurements used in this study. The OMI instrument was built by Dutch-Finnish collaboration, and is managed by the Royal Netherlands Meteorological Institute (KNMI). The authors also thank the GMI processing group for their extensive efforts in producing the GMI CTM. Data used in this study were processed at NASA Goddard Space Flight Center. Funding for this research was provided in part by Goddard Earth Science Technology (GEST) grant NGC5-494.

References

- Bloom, S., da Silva, A., Dee, D., et al.: Documentation and validation of the Goddard Earth Observing System (GEOS) Data Assimilation System – Version 4, Technical Report Series on Global Modeling and Data Assimilation, 104606, 2005.
- Chandra, S., Ziemke, J. R., Min, W., and Read, W. G.: Effects of 1997–1998 El Niño on tropospheric ozone and water vapor, *Geophys. Res. Lett.*, 25, 3867–3870, 1998.
- Chandra, S. and Ziemke, J. R.: Tropical tropospheric ozone: Implications for dynamics and biomass burning, *J. Geophys. Res.*, 107(D14), doi:10.1029/2001JD000447, 2002.
- Chandra, S., Ziemke, J. R., and Martin, R. V.: Tropospheric ozone at tropical and middle latitudes derived from TOMS/MLS residual: Comparison with a global model, *J. Geophys. Res.*, 108(D9), 4291, doi:10.1029/2002JD002912, 2003.
- Chandra, S., Ziemke, J. R., Schoeberl, M. R., et al.: Effects of the 2004 El Niño on tropospheric ozone and water vapor, *Geophys. Res. Lett.*, 34, L06802, doi:10.1029/2006GL028779, 2007.
- Doherty, R. M., Stevenson, D. S., Johnson, C. E., et al.: Tropospheric ozone and El Niño–Southern Oscillation: Influence of atmospheric dynamics, biomass burning emissions, and future climate change, *J. Geophys. Res.*, 111, D19304, doi:10.1029/2005JD006849, 2006.
- Duncan, B. N., Martin, R. V., Staudt, A., et al.: Inter-annual and seasonal variability of biomass burning emissions constrained by satellite observations, *J. Geophys. Res.*, 108, D2, doi:10.1029/2002JD002378, 2003.
- Duncan, B. N., Strahan, S. E., Yoshida, Y., Steenrod, S. D., and Livesey, N.: Model study of the cross-tropopause transport of biomass burning pollution, *Atmos. Chem. Phys.*, 7, 3713–

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3736, 2007,

<http://www.atmos-chem-phys.net/7/3713/2007/>.

Duncan, B. N., West, J. J., Yoshida, Y., Fiore, A. M., and Ziemke, J. R.: The influence of European pollution on ozone in the Near East and northern Africa, *Atmos. Chem. Phys.*, 8, 2267–2283, 2008,

<http://www.atmos-chem-phys.net/8/2267/2008/>.

Froidevaux, L., Jiang, Y. B., Lambert, A., et al.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *J. Geophys. Res.*, 113(D15), D15S20, 2008.

Fujiwara, M., Kita, K., Kawakami, S., et al.: Tropospheric ozone enhancements during the Indonesian forest fire events in 1994 and in 1997 as revealed by ground-based observations, *Geophys. Res. Lett.*, 26, 2417–2420, 1999.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 2006, <http://www.atmos-chem-phys.net/6/3181/2006/>.

Jacobson, M.: Computaton of global photochemistry with SMVGear II, *Atmos. Environ.*, 29, 3541–2546, 1995.

Kroon, M., Petropavlovskikh, I., Shetter, R. E., Hall, S., Ullmann, K., Veefkind, J. P., McPeters, R. D., Browell, E. V., and Levelt, P.: OMI Total Ozone Column Validation with Aura-AVE CAFS Observations, *J. Geophys. Res.*, 113(D16), D15S13, doi:10.1029/2007JD008795, 2008a.

Kroon, M., Veefkind, J. P., Sneep, M., McPeters, R. D., Bhartia, P. K., and Levelt, P.: Comparing OMI-TOMS and OMI-DOAS total ozone column data, *J. Geophys. Res.*, 113(D16), D16S28, doi:10.1029/2007JD008798, 2008b.

Levelt, P. F., Hilsenrath, E., Leppelmeier, G. W., et al.: Science objectives of the Ozone Monitoring Instrument, *IEEE Trans. Geosci. Remote Sens.*, 44(5), 1199–1208, 2006.

Livesey, N. J., Filipiak, M. J., Froidevaux, L., et al.: Validation of Aura Microwave Limb Sounder O-3 and CO observations in the upper troposphere and lower stratosphere, *Geophys. Res.*, 113(D15), D15S02, 2008.

Logan, J. A., Megretskaia, I., Nassar, R., et al.: Effects of the 2006 El Niño on tropospheric composition as revealed by data from the Tropospheric Emission Spectrometer (TES), *Geophys. Res. Lett.*, 35, L03816, doi:10.1020/2007GL031698, 2008.

Martin, R. V., Jacob, D. J., Logan, J. A., et al.: Interpretation of TOMS observations of tropical tropospheric ozone with a global model and in-situ observations, *J. Geophys. Res.*,

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

107(D18), 4351, doi:10.1029/2001JD001480, 2002.

McPeters, R. D., Kroon, M., Labow, G. J., Brinkma, E., Balis, D., Petropavlovskikh, I., Veefkind, J. P., Bhartia, P. K., and Levelt, P.: Validation of the Aura Ozone Monitoring Instrument Total Column Ozone Product, *J. Geophys. Res.*, 113(D15), D15S14, doi:10.1029/2007JD008802, 2008.

Peters, W., Krol, M., Dentener, F. et al.: Identification of an El Niño-Southern Oscillation signal in a multiyear global simulation of tropospheric ozone, *J. Geophys. Res.*, 106(D10), 10398–10402, 2001.

Schoeberl, M. R., Duncan, B. N., Douglass, A., Waters, J., Livesey, N., Read, W., and Filipiak, M.: The Carbon Monoxide Tape Recorder, *Geophys. Res. Lett.*, 33, 12, L12811, doi:10.1029/2006GL026178, 2006.

Schoeberl, M. R., Ziemke, J. R., Bojkov, B., Livesey, N., Duncan, B. N., et al.: A trajectory-based estimate of the tropospheric ozone column using the residual method, *J. Geophys. Res.*, 112, D24S49, doi:10.1029/2007JD008773, 2007.

Strahan, S. E., Duncan, B. N. and Hoor, P.: Observationally-derived diagnostics of transport in the lowermost stratosphere and their application to the GMI chemistry transport model, *Atmos. Chem. Phys.*, 7, 2435–2445, 2007, <http://www.atmos-chem-phys.net/7/2435/2007/>.

Sudo, K. and Takahashi, M.: Simulation of tropospheric ozone changes during 1997–1998 El Niño: Meteorological impact on tropospheric photochemistry, *Geophys. Res. Lett.*, 28, 4091–4094, 2001.

Thompson, A. M., Witte, J. C., Hudson, R. D., et al.: Tropical tropospheric ozone and biomass burning, *Science*, 291, 2128–2132, 2001.

Zeng, G. and Pyle, J. A.: Influence of El Niño Southern Oscillation on stratosphere/troposphere exchange and the global tropospheric ozone budget, *Geophys. Res. Lett.*, 32, L01814, doi:10.1029/2004GL021353, 2005.

Ziemke, J. R., Chandra, S., and Bhartia, P. K.: A new NASA data product: Tropospheric and stratospheric column ozone in the tropics derived from TOMS measurements, *B. Am. Meteorol. Soc.*, 81, 580–583, 2000.

Ziemke, J. R., Chandra, S., and Bhartia, P. K.: A 25-year data record of atmospheric ozone from TOMS Cloud Slicing: Implications for trends in stratospheric and tropospheric ozone, *J. Geophys. Res.*, 110, D15105, doi:10.1029/2004JD005687, 2005.

Ziemke, J. R., Chandra, S., Duncan, B. N., Froidevaux, L., Bhartia, P. K., Levelt, P. F., and

Waters, J. W.: Tropospheric ozone determined from Aura OMI and MLS: Evaluation of measurements and comparison with the Global Modeling Initiative's Chemical Transport Model, J. Geophys. Res., 111, D19303, doi:10.1029/2006JD007089, 2006.

ACPD

9, 2735–2761, 2009

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Table 1. Emissions of CO and NO for 2006.

	CO (Tg)		NO (Tg)	
	Indonesia ^a	World ^b	Indonesia ^a	World ^b
July	5.9	92	0.10	7.3
August	16.6	103	0.29	7.4
September	18.3	99	0.32	6.8
October	44.2	73	0.76	6.3
November	6.3	79	0.11	6.3
December	0.2	103	0.00	7.1

^a Biomass burning emissions only.

^b Emissions from all sources, including fossil fuels, biofuels, and biomass burning, except biomass burning from Indonesia.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Monthly Averaged Global Tropospheric Burden of CO, O₃, and NO_x for the Model. Simulations with (“Indonesia”) and without (“No Indonesia”) the 2006 Indonesia fire emissions.

	Sep.	Oct.	Nov.	Dec.
CO (Tg)				
Indonesia	351	371	371	369
No Indonesia	336	335	329	341
ΔCO	15	36	42	28
ΔCO (%)	4.6%	10.7%	12.6%	8.1%
O ₃ (Tg)				
Indonesia	354	351	346	332
No Indonesia	347	343	337	324
ΔO ₃	8	8	10	8
ΔO ₃ (%)	2.2%	2.4%	2.8%	2.4%
NO _x (Tg N)				
Indonesia	0.115	0.121	0.129	0.133
No Indonesia	0.113	0.117	0.128	0.133
ΔNO _x	0.002	0.004	0.001	-0.000
ΔNO _x (%)	2.1%	3.0%	0.8%	-0.3%

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.




Fig. 1. (a) Tropospheric column ozone measured in DU from OMI/MLS for October 2004 (top panel), October 2005 (middle panel), and October 2006 (bottom panel). (b) Same as (a) but instead for the GMI model. (c) The difference (GMI minus OMI/MLS) between Fig. 1a and b.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.




Fig. 2. (a) Inter-annual difference (October 2006 minus October 2005) of tropospheric column ozone measured in DU from OMI/MLS (top panel) and the GMI model (bottom panel). (b) Same as (a) but instead for November. (c) Same as (a) but instead for December.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.




Fig. 3. (a) Inter-annual annual difference (October 2006 minus October 2005) of 215 hPa CO from MLS (top panel) and 227 hPa CO from the GMI model (bottom panel). All quantities represent volume mixing ratio in units ppbv. (b) Same as (a) but instead for November. (c) Same as (a) but instead for December.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Biomass burning and
tropospheric ozone
during 2006 El Niño**

S. Chandra et al.




Fig. 4. (a) Inter-annual difference (October 2006 minus October 2005) of GMI tropospheric column ozone (in DU) of the Indonesia biomass burning component (top panel) and the dynamically driven component (bottom panel). (b) Same as (a) but instead for November. (c) Same as (a) but instead for December.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.




Fig. 5. OMI/MLS time series Δ TCO (dark curve) plotted with Niño 3.4 time series Δ SST (light curve). Δ TCO is measured in DU while Δ SST is in Celsius degrees and was multiplied by a factor of three for scaling with Δ TCO. The Δ TCO time series is derived by subtracting TCO in the eastern Pacific (15° S– 15° N, 135° W– 180°) from TCO in the western Pacific (15° S– 15° N, 95° E– 140° E). The Δ TCO time series was deseasonalized and was also smoothed with a three-month running average to be compatible with the Niño 3.4 time series.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Biomass burning and tropospheric ozone during 2006 El Niño

S. Chandra et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. A1. Monthly time series comparisons of GMI (dotted curves) and OMI/MLS (solid curves) tropospheric column O₃ (in DU) for three separate regions: American (left four panels), African (middle four panels), and Indonesian (right four panels).

**Biomass burning and
tropospheric ozone
during 2006 El Niño**

S. Chandra et al.




Fig. A2. (top) Zonal correlations between OMI/MLS and GMI tropospheric column O_3 . (middle) Calculated mean zonal offset difference (GMI minus OMI/MLS) measured in DU between model and observation. (bottom) Calculated zonal RMS of difference (GMI minus OMI/MLS) in DU. For both model and observation, fields for similar months (January – December) from 2004 to 2006 were averaged together to derive mean 12-month latitude-longitude gridded distributions of tropospheric O_3 . All statistical calculations were done zonally for each latitude band using the 72 gridded longitudinal values from model and observation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)