

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

First steps towards the assimilation of IASI ozone data into the MOCAGE-PALM system

S. Massart¹, C. Clerbaux², D. Cariolle¹, A. Piacentini¹, S. Turquety², and J. Hadji-Lazaro²

¹URA CNRS/CERFACS No. 1875, 42 avenue Gaspard Coriolis, 31057 Toulouse, France

²Université Pierre et Marie Curie-Paris 6, CNRS 8190, LATMOS/IPSL, Paris, France

Received: 7 January 2009 – Accepted: 24 February 2009 – Published: 10 March 2009

Correspondence to: S. Massart (massart@cerfacs.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The Infrared Atmospheric Sounding Interferometer (IASI) is one of the five European new generation instruments carried by the polar-orbiting MetOp-A satellite. Data assimilation is a powerful tool to combine these data with a numerical model. This paper presents the first steps made towards the assimilation of the total ozone columns from the IASI measurements into a chemistry transport model. The IASI ozone data used are provided by an inversion of radiances performed at the LATMOS (Laboratoire Atmosphères, Milieux, Observations Spatiales). As a contribution to the validation of this dataset, the LATMOS-IASI data are compared to a four dimensional ozone field, with low systematic and random errors compared to ozonesondes and OMI-DOAS data. This field results from the combined assimilation of ozone profiles from the MLS instrument and of total ozone columns from the SCIAMACHY instrument. It is found that on average, the LATMOS-IASI data tends to overestimate the total ozone columns by 2% to 8%. The random observation error of the LATMOS-IASI data is estimated to about 6%, except over polar regions and deserts where it is higher. Using this information, the LATMOS-IASI data are then assimilated, combined with the MLS data. This first LATMOS-IASI data assimilation experiment shows that the resulting analysis is quite similar to the one obtained from the combined MLS and SCIAMACHY data assimilation.

1 Introduction

Onboarding a new generation of instruments with higher resolution in frequency, resolution and spectral coverage, the low-Earth orbits (LEO) satellites can perform very accurate observations of numerous parameters of the atmosphere: vertical profiles of temperature, humidity and trace gases measurements for example. Moreover, with their low orbital altitude, the LEO satellites can achieve a global coverage in less than one day. Recently, the LEO MetOp-A satellite, with the Infrared Atmospheric Sounding

ACPD

9, 6691–6737, 2009

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interferometer (IASI) onboard was launched. At an altitude of 817 km, MetOp-A takes about 100 min to complete one orbit. During that time, the Earth has rotated by around 25°. This means that a complete mapping of the Earth is obtained in about one day.

IASI is one of the five European new generation instruments onboard MetOp-A. It measures the infrared (IR) radiation emitted from the surface of the Earth and the atmosphere. Due to its very high resolution, data of excellent accuracy and resolution are obtained. These data are mainly atmospheric humidity and temperature profiles in the troposphere and lower stratosphere, as well as the concentration of the chemical components (methane, carbon monoxide, ozone) that play a key role in atmospheric chemistry. Aside the exploitation of IASI for improving short-term weather forecasting, this instrument can also be used to monitor the atmospheric content of the chemical species or to contribute to the improvement of short-term chemical weather forecasting systems.

The IASI footprint on the ground is about 12 km at nadir and 4 measurements are performed at the same time (every 50 km), but the vertical resolution is limited. In addition to total integrated columns, IASI also provides, depending on the considered species, additional information on their tropospheric and stratospheric contents. In contrast, most of the global Chemistry Transport Models (CTMs), have coarser horizontal resolution but more vertical levels. The assimilation of the IASI trace gases measurements by a CTM is therefore an efficient way to combine information brought by the instrument with the information brought by the model. Moreover, the assimilation provides global three dimensional distribution of the atmospheric chemical constituents on hourly time resolution.

Most of CTM systems assimilating atmospheric chemical constituents are based on satellite level 2 products that include vertical profiles of trace gases or partial and total column determination (Lahoz et al., 2007a). Thus, before attempting to assimilate the IASI data, quality products from the IASI spectra have to be elaborated. At the time this article is written, IASI is a new instrument and retrieval algorithms are still in an improving phase. The main objective of this work is therefore to obtain a cross-validation

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I ◀](#)[▶ I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of the IASI retrieval algorithm developed by the LATMOS (Turquety et al., 2004). This is first achieved by performing comparisons over a five month period (from August to December 2007) between the LATMOS-IASI level 2 ozone data and the ozone fields computed through the combined assimilation of SCIAMACHY and MLS ozone data into the MOCAGE-PALM system. In contrast to the usual direct comparison of different data sets, this methodology allows the computation of ozone fields at the time and the location of the observations, within the time and space resolutions of the used CTM. The computed ozone field is a good estimate of the true ozone distribution since it results from an assimilation process. Finally, the comparison with the field from the combined assimilation of SCIAMACHY and MLS data is better than a comparison with only one of them, since the combined assimilation gives accurate information through the whole atmospheric column.

The next sections describe the characteristics of the ozone data used (either assimilated or used for validation), the results from the combined assimilation of SCIAMACHY and MLS ozone data, and the validation of the corresponding analysis¹. Section 4 focuses on the determination of the error of the LATMOS-IASI ozone data in comparison with ozone analysis obtained from the assimilation of the SCIAMACHY and MLS data sets. Section 5 presents the first results of the assimilation of LATMOS-IASI ozone data within our MOCAGE-PALM system.

2 Ozone observations

The ozone data used for this study are either assimilated or used as independent source of information to evaluate outputs from the assimilations process. They come from both spacecraft and in situ instruments.

¹In this paper, an analysis will refer to a four dimensional (3D in space plus time) field, computed on the model grid, resulting from the assimilation process.

2.1 Ozone observations for the assimilations

2.1.1 LATMOS-IASI

A clear-sky retrieval scheme for the near real time inversion of the concentrations of ozone and other trace gases was developed at the LATMOS (Turquety et al., 2004). It includes the inversion module, based on a neural network approach, as well as an error analysis module. The efficiency of the algorithm was demonstrated by its application to the treatment of the atmospheric nadir measurements provided by the Interferometric Monitor for Greenhouse Gases (IMG)/ADEOS (Clerbaux et al., 2003).

Global distributions of tropospheric and total ozone column are now routinely extracted from the IASI radiances data (Clerbaux et al., 2009). Systematic validations of total columns and profiles retrievals, using ground-based and other satellite observations, are provided in Boynard et al. (2009). Preliminary results show that, compared to GOME-2, the LATMOS-IASI data overestimate the ozone total columns by about 6% at mid-latitudes with an excellent correlation of about 0.92. Over the polar regions and tropics, the correlation is found to be lower and the LATMOS-IASI data still overestimate the ozone content.

The current version of the LATMOS algorithm still encounters problems in case of observations recorded over icy and sandy surfaces. This is due to the lack of available auxiliary data related to surface emissivity and solar reflexion. They should be included in the METOP level 2 products, and should be implemented in the next version of the algorithm. In this paper, we chose not to use the observations recorded over icy and sandy surfaces that represents about 12% of the overall dataset. Both day and night measurements are used without any distinction and the total ozone columns are considered without averaging kernels.

During the period under study (153 days), there are 12 days without LATMOS-IASI observations due to some MetOp-A calibration operations. Otherwise, an average of about 135 000 retrievals of total ozone column are produced per day. With the aim of assimilating these data into a system with a $2^\circ \times 2^\circ$ global grid, we have to build super-

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observations. They are obtained by averaging all the observations measured at the same time (within a minute) in the corresponding $2^\circ \times 2^\circ$ cell. This procedure reduces the daily number of observations to an averaged value of the order of 21 500.

2.1.2 TOSOMI SCIAMACHY ozone columns

SCIAMACHY (Scanning Imaging Absorption Spectrometer for Atmospheric CHartography) is a scanning spectrometer whose primary objective is to perform global measurements of trace gases in the troposphere and in the stratosphere (Bovensmann et al., 1999). It is designed to measure sunlight transmitted, reflected and scattered by the Earth's atmosphere or surface in the ultraviolet, visible and near infrared wavelength region. With its moderate spectral resolution and a wide wavelength range, SCIAMACHY can measure many different trace gases despite their low concentrations. SCIAMACHY has three different viewing geometries: nadir, limb, and sun/moon occultations which yield total column values as well as vertical profiles of trace gases and aerosols in the stratosphere and in the troposphere. SCIAMACHY can observe the whole Earth with a coverage at the equator achieved within 3 days when using only nadir or limb modes.

In this study, we use the ozone columns retrieved from the SCIAMACHY spectra by the TOSOMI algorithm developed at the Royal Dutch Meteorological Institute (KNMI). The retrieval algorithm is an application to SCIAMACHY of the GOME algorithm TOGOMI (Valks and van Oss, 2003). This algorithm is based on the total ozone DOAS (Differential Optical Absorption Spectroscopy) algorithm developed for the OMI instrument (Veefkind et al., 2006). This is suitable as the OMI, SCIAMACHY and GOME instruments are very similar with respect to total ozone column retrieval using the DOAS method. The TOSOMI SCIAMACHY total ozone data (value and error) are available from <http://www.temis.nl>. They will be further refereed as SCIAMACHY data in this paper. Eskes et al. (2005) have presented comparisons between two years of SCIAMACHY data and co-located ground based measurements, and between 6 months of SCIAMACHY data and GOME retrievals. They found that SCIAMACHY ozone columns

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are, on average, 1.5% lower than the others data, without any strong dependence on geographical location.

In a similar way to what is done for LATMOS-IASI data, the SCIAMACHY ozone columns have been averaged for the assimilation, on the CTM 2° by 2° latitude-longitude grid and are used without taking account of the averaging kernels. During the period under study, there are 3 days without measurements. Otherwise, an average of 25 000 aggregated measurements of total ozone column is available each day.

2.1.3 Aura/MLS ozone profiles

The MLS instrument has been flying onboard the Aura satellite in a sun-synchronous polar orbit since August 2004. Vertical profiles of several atmospheric parameters are retrieved from the millimeter and sub-millimeter thermal emission measured at the atmospheric limb (Waters et al., 2006). Measurements are performed between 82° S and 82° N, with a long-track resolution that varies from about 165 km to nearly 300 km. For our study we have used the latest version (v2.2) of MLS ozone product. A detailed validation of this product and comparisons with other data sets are available in Froidevaux et al. (2008) and Livesey et al. (2008). They found that in the middle stratosphere, the MLS ozone data have small biases, with a precision of the order of 5%. The precision is similar in the upper troposphere and lower stratosphere but can reach 20% at 215 hPa.

The ozone observations assimilated in our study were screened according to their recommendations. We rejected the measurements with odd “Status” fields, with “Quality” fields lower than 1.2, and with “Convergence” fields greater than 1.8. We also restricted the data used to the pressure range from 215 hPa to 0.5 hPa, which represents measurements at 16 pressure levels per profile. One interesting feature of MLS is its ability to measure constituents in the Upper Troposphere-Lower Stratosphere (UTLS) where retrievals are made at 4 levels, namely 215 hPa, 147 hPa, 100 hPa and 68 hPa. Nevertheless, the precision is lower in this region with values from 5 to 100% below 100 hPa, from 20% to 30% at 100 hPa, while it is below 10% at lower pressure levels.

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The standard deviation of the observation error required by the assimilation algorithm is set to the specified observation error. The correlations between the measurements of a same profile are neglected. Froidevaux et al. (2008) showed that the averaging kernels' peaks are near unity for the levels we assimilate the data, that means we can use them without specifying the averaging kernels.

For the assimilation, in a manner consistent with the use of the other datasets, we have averaged the measurements that lie at a distance below 2° . It gives about 2000 measurements per day, and for the period in consideration, there are 16 days without Aura/MLS measurements.

2.2 Ozone observations for evaluation of the analyses

2.2.1 OMI-DOAS total ozone columns

Also onboard Aura, the Ozone Monitoring Instrument (OMI) is a nadir viewing imaging spectrograph that measures the solar radiation backscattered by the Earth's atmosphere and surface (Levelt et al., 2006). OMI combines the advantages of European ESA instruments GOME and SCIAMACHY with the characteristics of the NASA's TOMS instrument, that measures the complete spectrum in the ultraviolet/visible wavelength range with a very high spatial resolution and a daily global coverage. During one Aura orbit, OMI performs approximately 1650 measurements on the sunlit portion of the Earth. In the standard global observation mode, 60 across track ground pixels are measured simultaneously, the pixel size being $13\text{ km} \times 24\text{ km}$. These measurements cover a swath of approximately 2600 km wide on the surface, which enables measurements with a daily global coverage.

In this study, we use the OMI total ozone columns (data available from <http://www.temis.nl>) produced with the KNMI DOAS method (Veefkind et al., 2006). The OMI-DOAS total ozone columns have showed a globally averaged agreement better than 2% with the ground-based observations (Balis et al., 2007). The data shows no significant dependence on latitude except for the high latitudes of the Southern Hemisphere

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(SH) where it systematically overestimates the total ozone value by 3% to 5% .

2.2.2 Ozone sondes

Ozone soundings were obtained from the World Ozone and Ultraviolet Radiation Data Centre (WOUDC, <http://www.woudc.org/>) and the Southern Hemisphere Additional Ozonesondes project (SHADOZ, <http://croc.gsfc.nasa.gov/shadoz>, (Thompson et al., 2003). In this study, we use data from 23 locations mainly located in the latitude band from 30° S to 60° N (Fig. 1). Most of the stations are equipped with Electrochemical Concentration Cell (ECC) sondes. Four of them are equipped with Carbon-Iodide sondes and one with Brewer-Mast sondes.

As stressed by Geer et al. (2006), even if this data set may be somewhat heterogeneous, this heterogeneity is acceptable in order to obtain a wider coverage. According to laboratory comparisons of the three sonde types (Smit et al., 1998), the random variability of the overall data is estimated at about 10 to 15% in the UTLS and near 5% in the middle stratosphere where ozone concentrations are maximums.

3 Production of the ozone analysis

One aim is first to compare LATMOS-IASI total ozone column with data sets from other instruments. The way we perform this comparison is similar to what was completed in Massart et al. (2007). The LATMOS-IASI total ozone columns are compared to the columns computed from an ozone analysis. The analysis is obtained from the combined assimilation of MLS and SCIAMACHY data into the MOCAGE-PALM system. The next sections describe the assimilation system and the resulting the ozone analysis obtained for the five last months of 2007.

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1 Assimilation system

The assimilation system is similar to the one used in Massart et al. (2007). It is based on the MOCAGE-PALM system developed jointly by CERFACS and Météo-France in the framework of the FP5 European project ASSET (Lahoz et al., 2007b). The assimilation algorithm is a 3D-VAR, in the FGAT (first guess at appropriate time) variant (Fisher and Anderson, 2001). The system is based on the Météo-France comprehensive three-dimensional chemistry transport model (CTM) MOCAGE and the CERFACS PALM software (Buis et al., 2006). The CTM MOCAGE covers the planetary boundary layer, the free troposphere and the stratosphere. It provides a number of optional configurations with varying domain geometries and resolutions, as well as chemical and physical parameterization packages. MOCAGE is currently used for several applications, with recent examples in chemical weather forecasting (Dufour et al., 2004), chemistry-climate interactions (Teyssède et al., 2007) and intercontinental transport of ozone and of its precursors (Bousserez et al., 2007).

The first version of the MOCAGE-PALM assimilation system (Massart et al., 2005a,b), as it was originally implemented for the ASSET project, provided good quality ozone fields compared with ozonesondes and UARS/HALOE measurements with errors of the same order as those produced by several other assimilation systems (Geer et al., 2006). In order to improve our assimilation system, several changes have been recently made on the model resolution and on the characterization of the forecast error (Massart et al., 2007; Pannekoucke and Massart, 2008).

In this study, the domain geometry and resolution cover a global 2° by 2° horizontal grid with 60 level vertical levels from the surface up to 0.1 hPa. The meteorological forcing fields are provided by the operational European Centre for Medium-Range Weather Forecasts (ECMWF) numerical weather prediction model. For computing the ozone chemical fields, we adopted the linear ozone parameterization developed by Cariolle and Teyssède (2007) in its latest version 2. The forecast error covariance matrix of the MOCAGE-PALM assimilation system is split into a correlation matrix and a diago-

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nal matrix filled with the forecast error variance (square of the forecast error standard deviation). The correlation matrix is divided into a horizontal and a vertical operator, both modelled using a diffusion equation (Weaver and Courtier, 2001). The horizontal correlation is computed with a homogeneous length-scale of 2° at the equator (that corresponds to a distance of approximately 220 km). The vertical correlation is computed with a non-homogeneous length-scale.

In order to increase the consistency of our analysis, we added several diagnostics as discussed by Desroziers et al. (2005). Based on the combinations of observation minus background, observation minus analysis, and background minus analysis, they provide consistency check of an assimilation experiment. In our case, these diagnostics are used to compute two correction coefficients, one applied on the standard deviation of the forecast error, and the other applied on the standard deviation of the observation error. In theory, these coefficients are suitable for the assimilation period during which they are computed. In practice, they are computed over a daily period divided into 8 assimilation windows and are used for the next day. This procedure can be adopted due to the slow variation found in the standard deviation of the forecast and observation errors.

3.2 Numerical experiment

The period under study spans from first of August to the end of December 2007. Initial global ozone concentrations for 1 July 2007 were built from a July climatological ozone. A 15 days assimilation of MLS data from this initial state was performed to obtain coherent ozone concentrations for 15 July 2007. All the assimilation experiments start from this date. The computation of the diagnostics for the intercomparisons begins 1 August 2007 so that the fields have little-dependence on the initial condition. A simulation of MOCAGE in the free mode (without any assimilation) starting on 15 July 2007 is also performed in order to produce a reference from which the impact of the assimilation is evaluated.

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.

We assimilate the combined information from MLS and SCIAMACHY measurements. The major advantage in this combined assimilation is that the MLS data give constraints to the computed ozone profiles in the stratosphere and in the upper troposphere, whereas the SCIAMACHY data brings information on the integrated ozone columns. Before combining the two data sets, it is important to ensure coherence between these two sources of information. This is achieved by comparing the analyses of the two single instrument assimilation experiments of the two types of ozone measurements.

3.3 Separate MLS and SCIAMACHY assimilations

We first assimilated MLS and SCIAMACHY data in two separated experiments. For these experiments, the length-scale used for the vertical correlation model of the forecast error is set to a value of $0.35 \log(\text{hPa})$. The standard deviation of the background error is prescribed as a percentage of the ozone concentration, with a proportionality coefficient adjusted using the previously described a posteriori diagnostics. As illustrated in Fig. 2, the tuning of the correction coefficients for the standard deviation of the forecast and the observation errors, insures a good consistency of the assimilation system. All along the assimilation period, the minimum of the cost function remains close to half the number of observations. On average, the resulting observation error is approximately the same as the one specified in the data. The standard deviation of the forecast error is about 15% of the ozone concentration for the MLS assimilation, and 2% for the SCIAMACHY assimilation.

Even if the MLS data only provides ozone profiles, the total ozone columns computed from the MLS analysis, account for the information brought by the instrument. Therefore the total ozone columns computed from the MLS analysis and those computed from SCIAMACHY analysis can be legitimately compared. We computed for the whole period, the average and the standard deviation of the difference between the analysed SCIAMACHY total ozone columns and the analysed MLS ones. MLS having no measurements beyond 82°S and beyond 82°N , we assumed that the difference in

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

this polar regions is constant and equal to the one computed on 82° S and on 82° N for respectively the Southern and Northern polar regions. Figure 3 shows that there is a good consistency between the two sources of information, the averaged difference being mostly below 10 Dobson Unit (DU) in absolute value. The variation of the difference from one month to another is also below 10 DU. In the region between about 60° S and 20° N, for all months from August to December, the total ozone columns from SCIAMACHY analysis are lower than those from the MLS analysis, with little variations from one month to another. The monthly averaged differences are nearly systematically positives in the polar regions, with the SCIAMACHY analysis being larger than the MLS one. In the Southern Polar Region (SPR), the differences are larger for the three first months (August to October), while the maximum differences are found from October to November over the Northern Polar Region (NPR). Consequently, in order to remove the bias between the two data sets before performing a combined assimilation, we have corrected the SCIAMACHY data by subtracting the monthly differences between the SCIAMACHY and the MLS total ozone column analyses. In addition, since the diagnosed SCIAMACHY and MLS observation errors were of the same magnitude than the specified instrumental errors, we combined the MLS data and the corrected SCIAMACHY data without any change to the specified instrumental errors.

3.4 Forecast error parameterisation

In order to improve the quality of our combined analysis, we used a more efficient parameterisation of the forecast error. In our previous works, we assumed that the standard deviation of the forecast error is proportional to the ozone concentration, and that the length-scale for the vertical correlation of the forecast error is constant. In this study, following the work carried out by Pannekoucke and Massart (2008), we determined the standard deviation and the length-scale using an ensemble of assimilation runs. This method has previously proved to be efficient in the meteorological forecast framework (Belo Pereira and Berre, 2006). To build the ensemble, five sets of perturbed data are obtained using a random generation. For each observation of the reference

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data set, five observations are generated by a Gaussian random noise with a mean equal to the value of the reference observation and a standard deviation equal to the observation error. The monthly statistics are hence computed using four forecasts per day, given an ensemble of about 600 realisations for each 30 days period.

5 Figure 4 illustrates the standard deviation of the forecast error diagnosed for the whole period. It shows that on average the standard deviation is higher in the UTLS region with values between 3% and 5%. Below, in the troposphere, the standard deviation is higher over the polar regions than over the other regions of the globe. It is higher over the NPR in August and September, and higher over the SPR in December (figure not shown). In the middle stratosphere, the standard deviation is higher over the NPR
10 at the end of the period while it is higher over the SPR in August. In the upper stratosphere, the standard deviation is rather homogeneous. So, the assimilation process gives more confidence to the observations in the UTLS and in the polar regions.

Figure 5 shows the length-scale of the vertical correlation of the forecast error diagnosed for the entire period. The length-scale is nearly constant in the middle stratosphere, increases in the UTLS region, and decreases in the troposphere. In the upper stratosphere and mesosphere, it increases up to 0.5 hPa and decreases above that level (probably due to the top boundary effects). In terms of data assimilation, the information brought by an observation is spread over a larger vertical range if the length-scale is higher. So, the spread is more important in the upper stratosphere above
20 10 hPa, and in the troposphere but to a lower extent.

The seasonal and regional variability of the length-scale is higher in the upper stratosphere and mesosphere (see the shaded area in Fig. 5). Since our objective is mainly focused on tropospheric and stratospheric ozone, the optimisation of our system in the upper stratosphere and mesosphere is not crucial. In consequence, we adopted a vertical varying length-scale that is the same for the whole globe. This choice allows a
25 significant reduction of the numerical cost of the parameterisation.

**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4 Validation of the ozone analyses

The ozone analysis obtained by the combined assimilation of MLS and SCIAMACHY data as described previously, must be compared to independent data. The benefit brought by the assimilation is evaluated in comparison to the free simulation of MOCAGE. Since ozone total columns and profiles are assimilated, the evaluation is carried out by comparison with profiles from ozonesondes, and with total ozone columns from OMI-DOAS.

4.1 Comparison with ozonesondes

To compare ozonesonde measurements with the free model simulation and with the MLS plus SCIAMACHY analysis fields, all the observed profiles have been interpolated on a common vertical pressure grid. Then, for each interpolated observed profile, the co-located (in space and time) model profile is computed using the nearest neighbour for the spatial interpolation and a linear time interpolation. Following Geer et al. (2006), the difference in the vertical profiles is normalized by a combination of the Logan (1999) tropospheric climatology with the Fortuin and Kelder (1998) stratospheric climatology. The average and standard deviation of the normalized differences, expressed as percentages, are then computed over the globe and aggregated over latitude bands of 30 degrees, except for the equatorial region where they are averaged over a larger band of 60 degrees. The results are analysed only between the surface and the 5 hPa level since ozonesondes have large errors above that level. Unfortunately, for the period under study, ozonesondings were unavailable for the two latitude bands [60° S–30° S] and [60° N–90° N]. The number of ozonesonde profiles per latitude band used for the validation on the ozone analyses is given in Table 1.

Figure 6 shows the average difference between ozonesondes and the free simulation, and between ozonesondes and the combined MLS and SCIAMACHY analysis. It appears that the free simulation has a global bias below 20% in absolute value, with an overestimation of the ozone concentrations in the stratosphere, and an underestima-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion in the troposphere. The largest ozone overestimation is found in the SPR, between 250 hPa and 50 hPa. This overestimation is maximum around 150 hPa, and is more pronounced in November and December at the end of the ozone depletion period and the beginning of the ozone recovery (figure not shown). The model also overestimates the ozone by about 20%, each month, in the 30° N–60° N region between 100 hPa and 10 hPa. This apparent accumulation of ozone over the polar regions with the free simulation can be related to a too efficient equator to pole meridional circulation in the forcings of the model. Such a bias in the ECMWF wind analyses that force the CTM has been identified by Monge-Sanz et al. (2007) and Cariolle and Teysse re (2007). In the Equatorial region, the free model simulation tends to underestimate ozone particularly in the mid and upper troposphere (below 100 hPa). This results from the overestimation of the vertical wind velocities associated with the meridional circulation bias.

The combined assimilation of MLS and SCIAMACHY reduces these biases. In particular, in the stratosphere, between 150 hPa and 10 hPa, the difference between ozonesondes and the analysis is below 3% in absolute value. In the upper stratosphere above 10 hPa, since the standard deviation of the forecast error is low, observations have a smaller weight and the analysis remains close to the free simulation. In the UTLS, the standard deviation of the forecast error is larger and the analysis is driven towards the MLS data. (Jiang et al., 2007) reported that, compared to the ozonesondes, MLS products overestimate ozone at 215 hPa by about 20% at middle and high latitudes. Similarly, the assimilation of MLS data produces an ozone overestimation at this altitude range, especially over the SPR. Elsewhere in the UTLS and in the mid-troposphere, the information brought by the data is vertically spread and the biases are reduced.

Figure 7 shows the standard deviation of the difference between ozonesondes and the free simulation and between ozonesondes and the MLS plus SCIAMACHY analysis for the period under study. It appears that the random error between the model without assimilation and the ozonesondes is globally below 20% between 10 hPa and 100 hPa. This value is of the same magnitude as the measurement error. The error grows be-

**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

low 100 hPa and reaches about 50% at 250 hPa. The error is nearly the same for all latitudes, except over the SPR with slightly higher values, with no significant variations from one month to another. As expected, the analysis reduces the standard deviation. However, the reduction occurs mainly in the SPR between 200 hPa and 15 hPa. In particular, at 150 hPa the standard deviation decreases from 54% for the free simulation to 20% for the analysis. In other regions, the reduction covers mainly the 200 hPa to 15 hPa altitude range. A small increase of the standard deviation can be noticed in the troposphere.

4.2 Comparison with OMI-DOAS

The daily total ozone columns from OMI-DOAS were averaged on the 2° by 2° MOCAGE horizontal grid. The difference between this averaged OMI-DOAS data and the free simulation or the analysis, normalized by the OMI-DOAS ones, is computed each day at 12 UTC. Figure 8 shows the zonal average and the standard deviation of this difference.

The first panel of the Fig. 8 shows that the free simulation has an important bias in terms of total ozone columns compared to OMI-DOAS ones. The free simulation systematically overestimates the total ozone columns in the polar regions by 20% to 30%. This is consistent with the previous comparison with ozonesondes and it is related to the intensity of the meridional circulation. So, the biases of the free simulation are minimum at mid-latitudes, and increase again at low latitudes where the free simulation systematically underestimates the total ozone columns by 5% to 10%. This bias shows little variations from one month to another. The maximum value of 16.5% is reached in the middle of October around 10° N. In the SPR, the bias is lower in December, while it is lower in August over the NPR. The maximum biases are reached in the mid-September for the NPR with about -30% , and in the middle of November for the SPR with about -65% .

The computed total ozone columns from the analysis of the combined MLS and SCIAMACHY data agree well with the OMI-DOAS measurements. This is consistent

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with the low biased of the analysis compared to the ozonesondes in the altitude range where the largest fraction of the stratospheric ozone lies. Figure 8 shows that the analysis underestimates by less than 1.5% the total ozone columns between 30° S and 80°. It also shows that over the SPR, the analysis overestimates the ozone content by 3% to 8% with the largest contribution for the November month.

The lower panel of the Fig. 8 shows that the free simulation produces total ozone columns that deviate by 3% to 6% from the OMI-DOAS ones at most latitudes. The error is however the largest between 50° S and the South Pole. It peaks to 12.5% at 67° S. This error peak is present all over the period under study, with a maximum in September. Its location corresponds to the edges of the ozone hole, where ozone meridional gradients are maximums. The model error probably comes from a difficulty with the free simulation to locate the vortex edges with a sufficient accuracy.

Figure 8b indicates that the standard deviation of the analysis is reduced compared to the free simulation. It lies between 2 and 4% in most regions and represents an important improvement compared to the free simulation. In particular, the standard deviation computed with the analysis around 67° S is significantly reduced by a factor 2. The analysis reduces substantially the uncertainties around the edges of the Antarctic vortex. However, inside the Southern Polar vortex, there is no significant improvement in the statistics, especially for the month of November.

4.3 Synthesis on the MLS plus SCIAMACHY analysis

The combined assimilation of MLS and SCIAMACHY data provides ozone concentrations which fairly agree the independent ozonesondes measurements mainly in the stratosphere (with a difference below 3%). Given that the biggest fraction of the atmospheric ozone is located into the stratosphere, this implies that the analysed total ozone columns are slightly biased (predominantly below 1.5%) compared to the independent OMI-DOAS data.

In terms of random variability, the difference between the analysis and the ozonesondes is lower than 20% with an instrument accuracy around 5 to 15%. Concerning the

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



total ozone columns, the difference between the analysis and the OMI-DOAS data are mainly below 4%, the instrumental error being around 2%. This means that the random variability of our analysis is about 10% for the ozone concentrations and below 2% for the total ozone columns. Notwithstanding, the errors are higher at the edges of the Antarctic vortex and over the South Pole in November.

5 Assimilation of the LATMOS-IASI ozone data

The previous section showed that our ozone analysis provides a rather good four dimensional (3D in space plus time) representation of the true distribution with low systematic and random errors. It is thus pertinent to first compare the LATMOS-IASI data to this quality ozone fields before any attempt of assimilation.

5.1 Methodology

The comparison between the reference (not spatially averaged) LATMOS-IASI ozone data y_{iasI}^o and the combined MLS and SCIAMACHY analysis x_{m+s}^a is performed at the LATMOS-IASI time and space resolution. For each LATMOS-IASI data, we compute the co-located (in space and time) analysis using an observation operator \mathbf{H} that is a bi-linear spatial interpolation and a linear time interpolation between two successive analysed fields separated by 3 h. Afterwards, the difference between the LATMOS-IASI data and the co-located analysis, further referred as departure, is computed. This departure is then projected on the 2° by 2° horizontal model grid. Doing so, we obtained in each cell of the mesh, an ensemble of differences between LATMOS-IASI measurements and the co-located analysis. Figure 9 shows the repartition of the number of these departures per cell over the globe. This number varies from 200 to 3000 except in regions where the LATMOS-IASI data has been discarded due to insufficient knowledge of the emissivity properties of the ground.

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For each cell, we computed the average and the variance of the ensemble of the departures. The average represents the systematic error in terms of total column ozone between the LATMOS-IASI measurements and the combined MLS and SCIAMACHY analysis. To ensure coherence between the data and the model, this error has to be removed before assimilating LATMOS-IASI measurements. The variance of the departures gives information on the variance of the observation error. The departure is expressed as

$$\mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}_{m+s}^a = \mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}^t - \mathbf{H}(\mathbf{x}_{m+s}^a - \mathbf{x}^t), \quad (1)$$

where \mathbf{x}^t represents the true unknown ozone concentration and the linear observation operator \mathbf{H} is expressed as a matrix.

As the LATMOS-IASI observation error $\mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}^t$ and the MLS and SCIAMACHY analysis error $\mathbf{x}_{m+s}^a - \mathbf{x}^t$ are independent, the variance (denoted $\text{var}[\cdot]$) of the departure is given by the sum of the two variances,

$$\begin{aligned} \text{var}[\mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}_{m+s}^a] &= \text{var}[\mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}^t] \\ &+ \mathbf{H} \text{var}[(\mathbf{x}_{m+s}^a - \mathbf{x}^t)] \mathbf{H}^T, \end{aligned} \quad (2)$$

where \mathbf{H}^T is the transpose of the linear observation operator \mathbf{H} .

Then, the variance of the LATMOS-IASI observation error can be expressed as the difference between the variance of the departure and the variance of the MLS and SCIAMACHY analysis error (projected into the observation space),

$$\begin{aligned} \text{var}[\mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}^t] &= \text{var}[\mathbf{y}_{\text{iasi}}^o - \mathbf{H}\mathbf{x}_{m+s}^a] \\ &- \mathbf{H} \text{var}[(\mathbf{x}_{m+s}^a - \mathbf{x}^t)] \mathbf{H}^T. \end{aligned} \quad (3)$$

In order to estimate the variance of the LATMOS-IASI observation error, we estimated the variance of the MLS plus SCIAMACHY analysis error in a similar way to the estimation of the variance of the forecast error, using the ensemble of perturbed

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observations previously described. We found that the standard deviation of the analysis error is of 0.5% at the equator and increases to 1% at the Pole. This result agrees with the conclusions obtained with the combined assimilation of MLS and SCIAMACHY data. Consequently, we can assume that the variance of the forecast error is negligible compared to the variance of the LATMOS-IASI observation error. The variance of the LATMOS-IASI observation error is then simply obtained by computing the variance of the ensemble of departures.

To measure the benefit of assimilating LATMOS-IASI data instead of SCIAMACHY data, we can compare the mean and the standard deviation of the differences between the two analyses (MLS plus SCIAMACHY in one hand and MLS plus LATMOS-IASI in another hand) and the soundings or the OMI-DOAS measurements. Considering that the subscript $m+i$ refers to the combined assimilation of MLS and LATMOS-IASI, we compute \mathbf{d}_{m+i}^a and \mathbf{d}_{m+s}^a , where \mathbf{d}^a is the departure of the analysis,

$$\mathbf{d}^a = \mathbf{y}^o - \mathbf{H}\mathbf{x}^a, \quad (4)$$

with \mathbf{y}^o representing either OMI-DOAS or ozonesondes data and \mathbf{H} the corresponding observation operator.

The mean $E[\mathbf{d}^a]$ and the standard deviation $\text{Std}[\mathbf{d}^a]$ of these departures can next be computed. Finally, a gain \mathbf{g}_{ave} for the mean and a gain \mathbf{g}_{std} for the standard deviation can be expressed with

$$\begin{cases} \mathbf{g}_{\text{ave}} = 100 \frac{|E[\mathbf{d}_{m+s}^a]| - |E[\mathbf{d}_{m+i}^a]|}{(|E[\mathbf{d}_{m+s}^a]| + |E[\mathbf{d}_{m+i}^a]|)/2}, \\ \mathbf{g}_{\text{std}} = 100 \frac{\text{Std}[\mathbf{d}_{m+s}^a] - \text{Std}[\mathbf{d}_{m+i}^a]}{(\text{Std}[\mathbf{d}_{m+s}^a] + \text{Std}[\mathbf{d}_{m+i}^a])/2}. \end{cases} \quad (5)$$

When the difference \mathbf{d}_{m+i}^a between the independent observations and the analysis of MLS and LATMOS-IASI data is the same on average than the difference \mathbf{d}_{m+s}^a between the independent observations and the analysis of MLS and SCIAMACHY data, the gain \mathbf{g}_{ave} is null. When the difference \mathbf{d}_{m+i}^a is lower than the difference \mathbf{d}_{m+s}^a , the gain is positive, with an extreme value of 200% when $\mathbf{d}_{m+i}^a = 0$. On the opposite, the gain is

negative when the difference d_{m+i}^a is higher than the difference d_{m+s}^a . The behaviour of g_{std} is similar.

5.2 Statistics on the LATMOS-IASI data by comparison with ozone analysis

The time average of all the computed departures between the LATMOS-IASI data and our combined MLS and SCIAMACHY analysis, is given in the upper panel of Fig. 10. It shows that the systematic difference between the LATMOS-IASI data and the analysis lies between 2% and 8% which is consistent with the conclusions of Boynard et al. (2009).

The largest discrepancies are found at low and mid-latitudes. They could be the result of the presence of aerosols in large concentration whose signature is better seen in the IR sensor of the IASI instrument than in the SCIAMACHY sensor. At low and mid-latitudes, the LATMOS-IASI data overestimates the total ozone column by 6% on average. The maximum differences are reached over the Pacific and the Atlantic Oceans with values of 11% and 13% respectively while the difference is lower over the Indian Ocean. The overestimation of the LATMOS-IASI ozone data is also large in the vicinity of the desert regions of the North Africa and the Persian Gulf.

Over Europe and North America, there is a good agreement between the LATMOS-IASI data and our analysis. Nevertheless, the LATMOS-IASI data underestimates the total ozone over icy regions such as Siberia and Canada. Over the Arctic, the LATMOS-IASI data also underestimates the total ozone by about 6%, except over the north of Canada. In those regions the algorithm could still suffer from inadequate prescription of the ground emissivity.

The second panel of Fig. 10 shows the standard deviation of the ensemble of departures between the LATMOS-IASI data and our combined MLS and SCIAMACHY analysis. As explained previously, we assume that this standard deviation can be attributed to the standard deviation of the LATMOS-IASI observation error. Thus, the figure shows that the random observation error of the LATMOS-IASI data is about 6% except in polar regions and over specific regions. In particular, the observation error

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



can reach 10 to 15% over dry regions like Australia, North Africa to China and in a lesser extent, over the west coast of the United States, the west coast of the South Africa and the south part of the South America. The error also grows significantly over the polar regions and reaches 10% for the North and 40% for the South. This largest error seems to match the distribution of the ice caps.

5.3 LATMOS-IASI assimilation

To assimilate the LATMOS-IASI data, we first removed from the LATMOS-IASI super-observations the monthly averaged differences found between our previous analysis and the LATMOS-IASI data. The diagnosed standard deviation of the LATMOS-IASI observation error is used to set the variance of the observation error required by the assimilation system. In a similar manner to what was done with MLS and SCIAMACHY, we assimilate the combined MLS and LATMOS-IASI data sets. As this work is a first approach to the assimilation of the LATMOS-IASI data, we did not yet determine the most adequate standard deviation and length-scale of the vertical correlation of the forecast error. We simply used those determined for the combined assimilation of MLS and SCIAMACHY.

The difference in terms of ozone concentration between the two analyses is illustrated by Fig. 11. It shows that, on average, the two analyses differ in the troposphere, as expected, but also in the lower stratosphere, especially over the equatorial region and the SPR. Elsewhere, the differences are yet small. The main differences are found in the middle troposphere of the SPR, from 700 hPa to 200 hPa, where the combined MLS and LATMOS-IASI assimilation provides lower ozone concentrations than the combined MLS and SCIAMACHY one. As the previous comparison with ozonesondes has shown that the combined MLS and SCIAMACHY analysis overestimates ozone in this altitude range, the combined MLS and LATMOS-IASI analysis seem to bring significant improvements. This must be confirmed however by the analysis of the computed gains.

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The gain on the mean computed with the ozonesondes is displayed in Fig. 12. It shows that the gain is globally positive between 80 hPa and 10 hPa. As the difference between MLS plus SCIAMACHY analysis and ozonesondes is very low in this altitude range, it means that there is no real improvement or damage on the ozone bias in the stratosphere by assimilating LATMOS-IASI instead of SCIAMACHY. In this region, the analysis is mainly constrained by the good-quality MLS data, and the assimilation of total ozone columns has a small impact. Below, the assimilation of LATMOS-IASI reduces the tropospheric bias over the SPR, whereas the bias increases at mid-latitude of the Northern Hemisphere (NH), between 30° N and 60° N. Compared to the free simulation, the assimilation of SCIAMACHY data increases the bias in the troposphere over the SPR. As we imposed large observation errors to the LATMOS-IASI data in this region, the analysis is less constraining by their assimilation. The tropospheric ozone concentrations from the LATMOS-IASI analysis are then close to the ones of the free simulation, which are lower than those from the assimilation of SCIAMACHY data. A similar behaviour is found in the troposphere between 30° N and 60° N. The assimilations of SCIAMACHY or LATMOS-IASI data both decrease the bias compared to the free simulation. But the LATMOS-IASI data are less effective due to their larger errors.

In terms of standard deviation, Fig. 13 shows that there is a slight global improvement in the troposphere when assimilating LATMOS-IASI data instead of SCIAMACHY ones. The improvements are mainly visible over the SPR. The standard deviation computed with the MLS and LATMOS-IASI analysis is close to the one computed with the MLS analysis between 200 hPa and 500 hPa. Notwithstanding, below 500 hPa, the assimilation of the combined MLS and LATMOS-IASI data reduces the standard deviation compared to the combined MLS and SCIAMACHY analysis.

The gains on the mean and on the standard deviation, considering the OMI-DOAS total columns, are displayed on Fig. 14. This gain is positive over the SPR till 40° S. This comes from the contribution of the lower stratosphere and troposphere, as found previously. Elsewhere, the bias increases (plus 2% to 4%) with the assimilation of the

LATMOS-IASI data compared to the assimilation of the SCIAMACHY ones. As the observation error on the SCIAMACHY data is lower than the ones on the LATMOS-IASI data, the assimilation of the LATMOS-IASI data gives less constraints on the computed ozone columns. This explains why the bias with OMI-DOAS is increased while assimilating LATMOS-IASI instead of SCIAMACHY ones.

The gain on the standard deviation of the analysed total ozone column against OMI-DOAS data is mostly negative (bottom panel of Fig. 14). Between 40° S and 40° N, the gain is lower than -10% which signifies an increase of 10% on the standard deviation between the two analyses. Over the polar regions, the gain is slightly increasing.

6 Conclusions

For a period of five months which spans from August to December 2007, the MetOp-A/IASI initial dataset provided by the LATMOS inversion algorithm was compared to a four dimensional ozone field. This field results from the combined assimilation of ozone profiles from the MLS limb instrument and of total ozone columns from the SCIAMACHY nadir instrument. In comparison with ozone soundings and OMI-DOAS total ozone columns, we found that this analysed ozone field has low bias for the total columns (below 1.5%) and for vertical profiles in the stratosphere (below 3%). Its standard deviation is between 10% and 20% in the stratosphere and between 30% and 50% in the troposphere. Its standard deviation in terms of total ozone columns is mainly between 3% and 6%. Therefore, this analysed ozone field was chosen as a reference to compare with day/night LATMOS-IASI data. Thanks to this comparison, it is estimated that the LATMOS-IASI data tends to overestimate the total ozone columns by 2% to 8%, which is consistent with the findings of Boynard et al. (2009). The maximum overestimation is encountered over the equatorial region and in the vicinity of desert regions. The random observation error of the LATMOS-IASI data is about 6% except over Polar and dry regions where it is higher. This is related to the difficulty of the neural network algorithm to account for changing ground emissivity over sandy regions

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and its large weight given to mean climatological ozone profiles over polar regions.

A new set of LATMOS-IASI data was built removing the difference with the MLS and SCIAMACHY analysis, and specifying the standard deviation of their observation error. This LATMOS-IASI dataset was assimilated in combination with the MLS data. The combined MLS and LATMOS-IASI analysis agrees well with the MLS and SCIAMACHY one, with the main differences located in the troposphere and lower stratosphere over the SPR. Those differences are mainly driven by the model which is less constrained by the LATMOS-IASI data due to their lower precision (about 6%) compared to SCIAMACHY one (about 1%).

This work is first attempt to assimilate the IASI ozone data into a CTM. To further explore the potential of the IASI data, many improvements would be required. First, the quality of the retrievals can be increased in specific regions by an improved training of the neural network algorithm for these specific situations, e.g. with more detailed emissivity databases. In the near future, emissivity products should also be available as a METOP product. As the thermal contrast (difference of the temperature between the surface and the atmosphere) is larger during the night, daytime and nighttime IASI observations have different vertical sensitivities and accuracies (Clerbaux et al., 2009). It would therefore be interesting to diagnose and prescribe the random errors for the daytime data different from those for the night ones.

Secondly, in terms of assimilation, one major expected improvement would be the use of averaging kernels. Here also, the distinction between day and night data will be important. Using appropriate averaging kernels, a new ensemble of assimilation would be conducted in order to extract information of the covariance matrix of the forecast error. This is an important ingredient that can significantly improve our analysis system.

Finally, in order to make the best use of the whole information from the high horizontal IASI resolution, it would be interesting the increase the model grid resolution to avoid the use of super-observations. Going towards higher resolution, we also expect to gain in model accuracy due to a better representation of the horizontal gradients. Current model developments will allow us to use grid resolutions below 1° in the near future.

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. IASI has been developed and built under the responsibility of the Centre National d'Etudes Spatiales (CNES, France). It is flown onboard the Metop satellites as part of the EUMETSAT Polar System. The IASI L1 data are received through the EUMETCast near real time data distribution service.

5 The authors would like to thank Olivier Pannekoucke for his previous collaboration on the forecast error parameterisation and Irina Sandu for her advices on this paper.

The development of the MOCAGE-PALM assimilation system has benefit from the support of the Ether centre for atmospheric chemistry products and services (<http://munk.ipsl.jussieu.fr/>), and the French ADOMOCA project within the LEFE programs (<http://www.insu.cnrs.fr/co/lefe>).

10



The publication of this article is financed by CNRS-INSU.

References

15 Balis, D., Kroon, M., Koukouli, M. E., Brinksma, E. J., Labow, G., Veeffkind, J. P., and McPeters R. D.: Validation of Ozone Monitoring Instrument total ozone column measurements using Brewer and Dobson spectrophotometer ground-based observations, *J. Geophys. Res.*, 112, D24S46, doi:10.1029/2007JD008796, 2007. 6698

20 Belo Pereira, M. and Berre, L.: The use of an Ensemble approach to study the Background Error Covariances in a Global NWP model, *Mon. Weather Rev.*, 134, 2466–2489, 2006. 6703

Bousserez, N., Attié, J.-L., Peuch, V.-H., Michou, M., Pfister, G., Edwards, D., Emmons, L., Mari, C., Barret, B., Arnold, S. R., Heckel, A., Richter, A., Schlager, H., Lewis, A., Avery, M., Sachse, G., Browell, E. V., and Hair, J. W.: Evaluation of the MOCAGE chem-

ACPD

9, 6691–6737, 2009

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



istry transport model during the ICARTT/ITOP experiment, *J. Geophys. Res.*, 112, D10S42, doi:10.1029/2006JD007595, 2007. 6700

Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. H. P.: SCIAMACHY – Mission objectives and measurement modes, *J. Atmos. Sci.*, 56, 127–150, 1999. 6696

Boynard, A., Clerbaux, C., Coheur, P.-F., Hurtmans, D., Turquety, S., George, M., Hadji-Lazaro, J., Keim, C., and Mayer-Arnek, J.: Measurements of total and tropospheric ozone from the IASI instrument: comparison with satellite and ozonesonde observations, *Atmos. Chem. Phys. Discuss.*, submitted, 2009. 6695, 6712, 6715

Buis, S., Piacentini, A., and Déclat, D.: PALM: A Computational framework for assembling high performance computing applications, *Concurrency Computat.: Pract. Exper.*, 18(2), 247–262, 2006. 6700

Cariolle, D. and Teyssère, H.: A revised linear ozone photochemistry parameterization for use in transport and general circulation models: multi-annual simulations, *Atmos. Chem. Phys.*, 7, 2183–2196, 2007, <http://www.atmos-chem-phys.net/7/2183/2007/>. 6700, 6706

Clerbaux, C., Hadji-Lazaro, J., Turquety, S., Mège, G., and Coheur, P.-F.: Trace gas measurements from infrared satellite for chemistry and climate applications, *Atmos. Chem. Phys.*, 3, 1495–1508, 2003, <http://www.atmos-chem-phys.net/3/1495/2003/>. 6695

Clerbaux, C., Coheur, P.-F., Boynard, A., Clarisse, L., Hadji-Lazaro, J., Hurtmans, D., George, M., Pommier, M., Razavi, A., Turquety, S., and Wespes, C.: Monitoring of atmospheric composition using the thermal infrared IASI/METOP sounder, *Atmos. Chem. Phys. Discuss.*, submitted, 2009. 6695, 6716

Desroziers, G., Berre, L., Chapnik B., and Poli, P.: Diagnosis of observation, background and analysis-error statistics in observation space, *Q. J. Roy. Meteor. Soc.*, 131, 613, 3385–3396, 2005. 6701

Dufour, A., Amodei, M., Ancellet, G., and Peuch, V.-H.: Observed and modelled “chemical weather” during ESCOMPTE, *Atmos. Res.*, 74, 161–189, 2004. 6700

Eskes, H. J., van der A, R. J., Brinksma, E. J., Veeffkind, J. P., de Haan, J. F., and Valks, P. J. M.: Retrieval and validation of ozone columns derived from measurements of SCIAMACHY on Envisat, *Atmos. Chem. Phys. Discuss.*, 5, 4429–4475, 2005, <http://www.atmos-chem-phys-discuss.net/5/4429/2005/>. 6696

Towards the assimilation of IASI ozone data

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fisher, M. and Anderson, E.: Developments in 4D-Var and Kalman Filtering, ECMWF Technical Memorandum, 347, 38 pp., 2001. 6700
- Fortuin, J. P. F. and Kelder, H.: An ozone climatology based on ozonesonde and satellite measurements, *J. Geophys. Res.*, 103, 31, 709–734, 1998. 6705
- 5 Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Browell, E. V., Hair, J. W., Avery, M. A., McGee, T. J., Twigg, L. W., Sumnicht, G. K., Jucks, K. W., Margitan, J. J., Sen, B., Stachnik, R. A., Toon, G. C., Bernath, P. F., Boone, C. D., Walker, K. A., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W.,
10 Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *J. Geophys. Res.*, 113, D15S20, doi:10.1029/2007JD008771, 2008. 6697, 6698
- Geer, A. J., Lahoz, W. A., Bekki, S., Bormann, N., Errera, Q., Eskes, H. J., Fonteyn, D., Jackson, D. R., Juckes, M. N., Massart, S., Peuch, V.-H., Rharmili, S., and Segers, A.: The ASSET
15 intercomparison of ozone analyses: method and first results, *Atmos. Chem. Phys.*, 6, 5445–5474, 2006,
http://www.atmos-chem-phys.net/6/5445/2006/. 6699, 6700, 6705
- Jiang, Y. B., Froidevaux, L., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Bojkov, B.,
20 Leblanc, T., McDermid, I. S., Godin-Beekmann, S., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Perun, V. S., Schwartz, M. J., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Allaart, M., Andersen, S. B., Bodeker, G., Calpini, B., Claude, H., Coetzee, G., Davies, J., De Backer, H., Dier, H., Fujiwara, M., Johnson, B., Kelder, H., Leme, N. P., König-Langlo, G., Kyro, E., Laneve, G., Fook, L. S., Merrill, J., Morris, G., Newchurch, M., Oltmans, S., Parrondos, M. C.,
25 Posny, F., Schmidlin, F., Skrivankova, P., Stubi, R., Tarasick, D., Thompson, A., Thouret, V., Viatte, P., Vömel, H., von Der Gathen, P., Yela, M., and Zabolocki, G.: Validation of Aura Microwave Limb Sounder Ozone by ozonesonde and lidar measurements *J. Geophys. Res.*, 112, D24S34, doi:10.1029/2007JD008776, 2007. 6706
- Lahoz, W. A., Errera, Q., Swinbank, R., and Fonteyn, D.: Data assimilation of stratospheric constituents: a review, *Atmos. Chem. Phys.*, 7, 5745–5773, 2007a,
30 http://www.atmos-chem-phys.net/7/5745/2007/. 6693
- Lahoz, W. A., Geer, A. J., Bekki, S., Bormann, N., Ceccherini, S., Elbern, H., Errera, Q., Eskes, H. J., Fonteyn, D., Jackson, D. R., Khatatov, B., Marchand, M., Massart, S., Peuch, V.-H.,

**Towards the
assimilation of IASI
ozone data**S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rharmili, S., Ridolfi, M., Segers, A., Talagrand, O., Thornton, H. E., Vik, A. F., and von Clarmann, T.: The Assimilation of Envisat data (ASSET) project, *Atmos. Chem. Phys.*, 7, 1773–1796, 2007b,

<http://www.atmos-chem-phys.net/7/1773/2007/>. 6700

5 Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J. O. V., and Saari, H.: The Ozone Monitoring Instrument, *Geosci. Remote Sens.*, 44(5), 1093–1101, 2006. 6698

Livesey, N. J., Filipiak, M. J., Froidevaux, L., Read, W. G., Lambert, A., Santee, M. L., Jiang, J. H., Pumphrey, H. C., Waters, J. W., Cofield, R. E., Cuddy, D. T., Daffer, W. H., Drouin, B. J., Fuller, R. A., Jarnot, R. F., Jiang, Y. B., Knosp, B. W., Li, Q. B., Perun, V. S., Schwartz, M. J., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Avery, M., Browell, E. V., Cammas, J.-P., Christensen, L. E., Diskin, G. S., Gao, R.-S., Jost, H.-J., Loewenstein, M., Lopez, J. D., Nedelec, P., Osterman, G. B., Sachse, G. W., and Webster, C. R.: Validation of
10 Aura Microwave Limb Sounder O₃ and CO observations in the upper troposphere and lower stratosphere, *J. Geophys. Res.*, 113, D15S02, doi:10.1029/2007JD008805, 2008. 6697

15 Logan, J. A.: An analysis of ozonesonde data for the troposphere: Recommendations for testing 3-D models and development of a gridded climatology for tropospheric ozone, *J. Geophys. Res.*, 104, 16, 115–149, 1999. 6705

Massart, S., Cariolle, D., and Peuch, V.-H.: Vers une meilleure représentation de la distribution et de la variabilité de l’ozone atmosphérique par l’assimilation des données satellitaires, *C. R. Acad. Sci.*, 337, 1305–1310, 2005a. 6700

20 Massart, S., Manzoni, H., Cariolle, D., Peuch, V.-H., and Piacentini, A.: Validation of a 3D-Fgat assimilation of MIPAS ozone profiles in a global chemistry and transport model, Fourth WMO Symposium on Assimilation of Observations in Meteorology and Oceanography, Prague, Czech Republic, April 2005b. 6700

25 Massart, S., Piacentini, A., Cariolle, D., El Amraoui, L., and Semane, N.: Assessment of the quality of the ozone measurements from the Odin/SMR instrument using data assimilation, *Can. J. Phys.*, 85, 1209–1223, 2007. 6699, 6700

Monge-Sanz, B., Chipperfield, M.P., Simmons, A., and Uppala, S.: Mean age of air and transport in a CTM: Comparison of different ECMWF analyses, *Geophys. Res. Lett.*, 34, L04801, doi:10.1029/2006GL028515, 2007. 6706

30 Pannekoucke, O. and Massart, S.: Estimation of the local diffusion tensor and normalization for heterogeneous correlation modelling using a diffusion equation, *Q. J. Roy. Meteor. Soc.*,

**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

134, 635, 1425–1438, doi:10.1002/qj.288, 2008. 6700, 6703

Smit, H. G. J., Strater, W., Helten, M., Kley, D., Ciupa, D., Claude, H. J., Kohler, U., Hoegger, B., Levrat, G., Johnson, B., Oltmans, S. J., Kerr, J. B., Tarasick, D. W., Davies, J., Shitamichi, M., Srivastav, S. K., Vialle, C., and Velghe, G., JOSIE: The 1996 WMO International inter-comparison of ozonesondes under quasi flight conditions in the environmental simulation chamber at Julich, in: Proceedings of the XVIII Quadrennial Ozone Symposium, edited by: Bojkov, R. and Visconti, G., L'Aquila, Italy, September 1996, 971–974, 1998. 6699

Teyssède, H., Michou, M., Clark, H. L., Josse, B., Karcher, F., Olivie, D., Peuch, V.-H., Saint-Martin, D., Cariolle, D., Attié, J.-L., Nédélec, P., Ricaud, P., Thouret, V., van der A, R. J., Volz-Thomas, A., and Chéroux, F.: A new tropospheric and stratospheric Chemistry and Transport Model MOCAGE-Climat for multi-year studies: evaluation of the present-day climatology and sensitivity to surface processes, *Atmos. Chem. Phys.*, 7, 5815–5860, 2007, <http://www.atmos-chem-phys.net/7/5815/2007/>. 6700

Thompson, A.M., Witte, J.C., McPeters, R.D., Oltmans, S.J., Schmidlin, F.J., Logan, J.A., Fujiiwara, M., Kirchhoff, V. W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S., Ogawa, T., Johnson, B. J., Vömel, H., and Labow, G.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 1. Comparison with Total Ozone Mapping Spectrometer (TOMS) and ground-based measurements, *J. Geophys. Res.*, 108, 8238, doi:10.1029/2001JD000967, 2003. 6699

Turquety, S., Hadji-Lazaro, J., Clerbaux, C., Hauglustaine, D. A., Clough, S. A., Cassé, V., Schlüssel, P., and Mégie, G.: Operational trace gas retrieval algorithm for the Infrared Atmospheric Sounding Interferometer, *J. Geophys. Res.*, 109, D21301, doi:10.1029/2004JD004821, 2004. 6694, 6695

Valks, P. and van Oss, R.: TOGOMI Algorithm Theoretical Basis Document, Issue 1.2, TOGOMI/KNMI/ATBD/001, KNMI/ESA, November 2003. 6696

Veefkind, J. P., de Haan, J. F., Brinksma, E. J., Kroon, M., and Levelt, P. F.: Total Ozone from the Ozone Monitoring Instrument (OMI) Using the DOAS technique, *Geo. Rem. Sens.*, 44, 5, 1239–1244, doi:10.1109/TGRS.2006.871204, 2006. 6696, 6698

Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read, W. G., Siegel, P. H., Cofield, R. E., Filipiak, M. J., Flower, D. A., Holden, J. R., Lau, G. K., Livesey, N. J., Manney, G. L., Pumphrey, H. C., Santee, M. L., Wu, D. L., Cuddy, D. T., Lay, R. R., Loo, M. S., Perun, V. S., Schwartz, M. J., Stek, P. C., Thurstans, R. P., Boyles, M. A., Chandra, K. M., Chavez, M. C., Gun-Shing Chen, Chudasama, B. V., Dodge, R., Fuller, R. A., Girard, M.

- A., Jiang, J. H., Yibo Jiang, Knosp, B. W., LaBelle, R. C., Lam, J. C., Lee, K. A., Miller, D., Oswald, J. E., Patel, N. C., Pukala, D. M., Quintero, O., Scaff, D. M., Van Snyder, W., Tope, M. C., Wagner, P. A., and Walch, M. J.: The Earth observing system microwave limb sounder (EOS MLS) on the aura Satellite, IEEE T. Geosci. Remote, 44, 1075–1092, 2006. 6697
- 5 Weaver, A. and Courtier, P.: Correlation modelling on the sphere using a generalized diffusion equation, Q. J. Roy. Meteor. Soc., 127, 1815–1846, 2001. 6701

ACPD

9, 6691–6737, 2009

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I ◀](#)[▶ I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 1. Number of ozonesonde profiles used over the period August to December 2007 as a function of latitude band.

90° S–60° S	30° S–30° N	30° N–60° N
62	133	233

**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 1. Number of ozonesonde stations used for validation, by latitude (in 10° bins).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 2. Diagnostics of the SCIAMACHY assimilation for August 2007. **(a)** Values of the cost function at the minimum divided by the half number of observations. Values of the diagnosed **(b)** observation and **(c)** forecast error correction coefficients.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.




Fig. 3. Average difference (in DU) between the total ozone columns from the SCIAMACHY analysis minus the total ozone columns from the MLS analysis (black line) for the period 1 August 2007 to 31 December 2007. The shaded area represents the average plus and minus the standard deviation of the difference for the same period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 4. Zonal average of the August to December 2007 diagnosed ozone forecast error standard deviation (in %) as a function of pressure. The diagnostic is obtained from an ensemble of perturbed combined assimilations of MLS and SCIAMACHY data.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.




Fig. 5. Global average of the August to December 2007 diagnosed length-scale (in $\log(\text{hPa})$) of the vertical forecast correlation error, as a function of pressure. The diagnostic is obtained from an ensemble of perturbed combined assimilations of MLS and SCIAMACHY data. The shaded area represents the length-scale plus and minus its standard deviation calculated over the same period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.





Fig. 6. Mean of [ozonesondes – analysis] (black line) and [ozonesondes – free simulation] (dotted line) ozone profiles, normalised by climatology, for the period 1 August 2007 to 31 December 2007. The latitude band limits and the number of ozonesonde profiles used are given in the title of each sub-plot.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

**Fig. 7.** Same as Fig. 6 for the standard deviation.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 8. (a) Mean and (b) standard deviation of (OMI-DOAS-simulation)/OMI-DOAS total ozone column. All the plots are drawn for the period 1 August 2007 to 31 December 2007. The dotted line represents the free simulation, the black line represents the combined MLS and SCIAMACHY ozone analysis.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 9. Number of differences between the LATMOS-IASI measurements and the co-located combined MLS and SCIAMACHY analysis, computed on 2° by 2° cells, for the period 1 August 2007 to 31 December 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.




Fig. 10. Average **(a)** and standard deviation **(b)** of the difference in % between the LATMOS-IASI ozone data minus the co-located combined MLS and SCIAMACHY ozone analysis normalized by the IASI measurement, for the period 1 August 2007 to 31 December 2007.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 11. Zonal average of the difference in % between the time average ozone concentrations from the combined MLS and LATMOS-IASI analysis minus the ones from the combined MLS and SCIAMACHY analysis. The time average is performed on the period August to December 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.





Fig. 12. Gain in % on the mean of ozonesondes minus MLS plus LATMOS-IASI analysis compared to the mean of ozonesondes minus MLS plus SCIAMACHY analysis, for the period August to December 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Towards the
assimilation of IASI
ozone data**

S. Massart et al.

**Fig. 13.** Same as Fig. 12 for the standard deviation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Towards the
assimilation of IASI
ozone data**S. Massart et al.




Fig. 14. Gain in % on **(a)** mean and on **(b)** standard deviation of OMI-DOAS minus MLS and LATMOS-IASI analysis compared to OMI-DOAS minus MLS and SCIAMACHY analysis, for the period August to December 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)