9221

Atmos. Chem. Phys. Discuss., 9, 9221–9266, 2009 www.atmos-chem-phys-discuss.net/9/9221/2009/ © Author(s) 2009. This work is distributed under the Creative Commons Attribution 3.0 License.

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

Simulation of Mexico City plumes during the MIRAGE-Mex field campaign using the WRF-Chem model

X. Tie¹, S. Madronich¹, G. Li^{2,3}, Z. Ying^{1,4}, A. Weinheimer¹, E. Apel¹, and T. Campos¹

¹National Center for Atmospheric Research, Boulder, CO, USA ²Molina Center for Energy and the Environment, La Jolla, CA, USA ³Department of Atmospheric Sciences, Texas A&M, USA ⁴York University, Canada

Received: 19 March 2009 - Accepted: 23 March 2009 - Published: 8 April 2009

Correspondence to: X. Tie (xxtie@ucar.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.





Abstract

The quantification of tropospheric O_3 production in the Mexico City outflow is a major objective of the MIRAGE-Mex field campaign. We used a regional chemistry-transport model (WRF-Chem) to predict the distribution of O₃ and its precursors in Mexico City and the surrounding region during March 2006, and compared with in-situ aircraft mea-5 surement of O₃, CO, VOCs, NO_x, and NO_v concentrations. The comparison shows that the model is capable of capturing the timing/location of the measured city plumes, and the calculated variability along the flights is generally consistent with the measured results, showing a rapid enhancement of O_3 and its precursors when city plumes are detected. However, there are some notable differences between the calculated and 10 measured values, suggesting that, during transport from the surface of the city to the outflow plume, pollution levels are underestimated by about 0-25% during different flights. The calculated $O_3 - NO_y$, $O_3 - CO_z$, and $O_3 - NO_z$ correlations generally agree with the measured values, and the analysis of these correlations suggest that pho-

- tochemical O_3 production continues in the plume downwind of the city (aged plume), 15 adding to the O_3 already produced in the city and exported with the plume. The model is also used to quantify the contributions to OH reactivity from various compounds in the aged plume. This analysis suggests that oxygenated organics (OVOCs) have the highest OH reactivity and play important roles for the O_3 production in the aging plume. Furthermore, O_3 production per NO_y molecule consumed (O_3 production efficiency) is 20 more efficient in the aged plume than in the young plume near the city. The major con-
- tributor to the high O_3 production efficiency in the aged plume is the reaction $RO_2 + NO$. By contrast, the reaction of HO_2 +NO is rather uniformly distributed in the plume.

Introduction 1

In recent decades, air quality has deteriorated markedly in large cities of the developing 25 world. The export of air pollutants from urban to regional and global environments is of

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Figures** Þ١ Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

major concern because of wide-ranging potential consequences for human health and for cultivated and natural ecosystems, visibility degradation, weather modification, radiative forcing, and changes in tropospheric oxidation (self-cleaning) capacity. Rapidly increasing urbanization will be a major environmental driving force in the 21st century, affecting air quality on all scales – local, regional, and global.

The Atmospheric Chemistry Division (ACD) of the National Center for Atmospheric Research (NCAR) led an intensive field campaign called MIRAGE-Mex (Megacities Impact on Regional and Global Environment – Mexico City case study) in March 2006, as part of the MILAGRO (Megacities Initiative: Local And Global Research Observations)

- project. The goal of MIRAGE was to characterize the chemical/physical transformations and the ultimate fate of pollutants exported from urban areas, and to assess the current and future impacts of these exported pollutants on regional and global air quality, ecosystems, and climate. The campaign coordinated and integrated observations from ground stations, aircraft, and satellites, and provided a rich data base for improving regional and global medals of the transformations of aging urban
- ¹⁵ ing regional and global models of the transport and transformations of aging urban pollutants.

The Mexico City metropolitan area is one of the largest megacities in the world, covering 1500 km^2 with nearly 20 million inhabitants and including nearly 3.5 million vehicles. The city is at a high altitude (2240 m above sea level) and is in the tropics (10° N) and distance nearest time high color realisting intensity that leads to realist formed.

- (19° N), conditions permitting high solar radiation intensity that leads to rapid formation of ozone and secondary particulate matter. For example, hourly averaged ozone concentrations in the Mexico City Metropolitan Area (MCMA) exceed the Mexican air quality standard of 110 ppb frequently and often exceed twice the standard (Molina and Molina, 2002).
- The major focus of MIRAGE-Mex was to study the export of the urban plume to the surrounding area, and to study the evolution of chemical oxidants inside the plume. Tropospheric O_3 concentrations are functions of the chain lengths of NO_x ($NO_x=NO+NO_2$) and HO_x ($HO_x=OH+HO_2+RO_2$) radical catalytic cycles. Thus, O_3 formation in city plume is dependent on ambient levels of NO_x and volatile organic compounds (VOCs).

ACPD 9, 9221–9266, 2009					
Simulation of Mexico City plumes X. Tie et al.					
Title Page					
Abstract	Introduction				
Conclusions	References				
Tables	Figures				
14	►I				
•	F				
Back	Close				
Full Screen / Esc					
Printer-friendly Version					
Interactive Discussion					

BY

For example, Zaveri et al. (2003) found, in the 1999 Southern Oxidant study field campaign in Nashville, Tennessee, that the downwind O_3 concentrations in the Nashville plume are more sensitive to NO_x emissions than anthropogenic VOC emissions. By contrast, the study of Kleinman et al. (2003) suggested that O_3 production is VOC lim-

- ⁵ ited in the high-NO_x portions of the Philadelphia urban plume. Thielmann et al. (2002) suggested that at the rural area of Milan, Italy, ozone production per NO_x consumed is less efficient when the advected air masses originate from Milan. This lower efficiency is indicative of VOC-sensitive ozone production. A similar suggestion is also obtained in the New York City plume by Kleinman et al. (2000). From the different con-
- ¹⁰ clusions in different regions of the world one notices that the ozone formation in the city plume is strongly dependent on locations due to various ambient chemical conditions in different regions. In addition to ambient chemical conditions, ozone concentrations in a city plume are very sensitive to meteorological conditions, including extra-urban scale transport winds, vertical structures of wind fields and mixing processes, and mesoscale
- ¹⁵ convergence zones and transport processes. De Foy et al. (2006) suggested that the residence times in the Mexico City basin are less than 12 h with little carry-over from day to day and little recirculation of air back into the basin. Very efficient vertical mixing leads to a vertically diluted plume which, in April, is transported predominantly towards the Gulf of Mexico. Lei et al. (2008) studied the ozone production and response un-
- der different meteorological conditions in Mexico City, and Zhang et al. (2009) used the WRF-Chem model to study O₃ formation and compared model results to surface measurements. Mena-Carrasco et al. (2008) studied the regional impacts of Mexico City emissions on air quality and chemistry. Their study suggests that the influence of emissions from the Mexico City could lead to the enhancements in surface NO_x, CO, and O₃ concentrations to distances of about 200 km.

In this study, we use the WRF-Chem model to address scientific questions related to the evolution of ozone and its precursors in the Mexico City plume. We first intensively evaluate the model performance and then use the model to study ozone production in different chemical age regimes. The paper is organized as follows: In Sect. 2, we will

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Introduction Abstract Conclusions References **Figures** Þ١ Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

describe the MIRAGE-Mex flights and data as well as the regional chemical/dynamical model (WRF-Chem). In Sect. 3, the simulated chemical species will be compared to the measurements of CO, O_3 , NO_y , NO_z (NO_y - NO_x), NO_x , and VOC concentrations, so as to evaluate the performance of the model. Section 4 presents model-based estimates of O_3 evolution and OH reactivity related to O_3 production due to various processes, and the O_3 production efficiency in the city plume downwind of Mexico City.

2 Data and chemical model

2.1 Data

5

The MIRAGE-Mex field experiment obtained regional and large-scale measurements up to several hundred kilometers downwind of Mexico City using the C-130 aircraft. 10 During the field experiment, there were a total 12 flights which occurred in 2006 on 4 March (flight 1), 8 March (flight 2), 10 March (flight 3), 16 March (flight 5), 18 (flight March 6), 19 March (flight 7), 22 March (flight 8), 23 March (flight 9), 26 March (flight 10), 28 March (flight 11), and 29 March (flight 12), respectively. Among the flights, there were some short flights with a 3 h fly time, such as flight 10 and flight 11. There 15 were also some flights which were designed to fly over remote regions either to detect a long-range transport plume (more than 1000 km from the Mexico City) or to measure an intense biomass fire plume, such as the flights 7 and 9. In order to study the evolution of the Mexico City plume, we selected the flights which included Mexico City overpasses as well as downwind plume sampling, with flight duration of 7–10 h. With 20 these selection criteria, flights 2 (8 March), 3 (10 March), 6 (18 March), 8 (22 March),

11 (28 March), and 12 (29 March) are used in this study. Figure 1 shows the tracks for these 6 flights.

The meteorological conditions during this period were described in detail by Fast et al. (2007), and are briefly summarized here. On 8 March (flight 2), the outflow from the city exhibited a cycling pattern, first flowing over the eastern part of the city and

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Figures** Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

later turning southward. The C-130 flight route on this day was located just northeast of the city, followed by city overpasses during the middle of the mission. Thus, some parts of the city plume were detected by the flight. On 10 March (flight 3), the plume was predominantly transported toward the northeast from the city, and the flight was

- ⁵ located downwind in the plume. As a result, this flight measured the city plume during a long period of time, especially during the later flight times. On 18 March (flight 6), a trough extending from west to east across the US affected the position of the high pressure system over the Gulf of Mexico and led to variable wind directions over central Mexico. As a result, the flight detected the city plume when the flight passed over the
- city during later times of the flight. During the earlier part of the flight, the measured CO concentrations were very low, suggesting that background pollutants were detected during this period. On 22 March (flight 8), there was a plume transported toward the north and northeast from the city. The city plume was well detected by the flight during the middle of the flight, because the majority of the flight routes are located to the
- northeast of the city. On 28 and 29 March, there was a persistent high pressure center in central of Mexico, producing a northwestern wind. The flight routes of 28 March were mostly located upwind of the plume. As a result, the measured CO concentrations were generally low. Contrasted with other flights, the measured result during this day can be used as a background pollution state. The flight routes of 29 March were also located
 mostly far from the plume. However, during the middle of flight, the flight passed over
- 20 mostly far from the plume. However, during the middle of flight, the flight passed over the city, and high CO values were detected during this short period of time.

During the above 6 flights, O₃, CO, NO, NO₂, and NO_y concentrations were continuously measured with 1 s interval. Some volatile organic compounds (VOCs) were measured in-situ with high sampling frequency, while others were obtained more ²⁵ slowly via canister sampling. Some chemical species were unavailable on some flights. For example, O₃ was not measured on 18 March, and HCHO was not measured before 18 March. Archived data and related information are available at http://www-air.larc.nasa.gov/cgi-bin/arcstat-b.

ACPD 9, 9221–9266, 2009 Simulation of Mexico City plumes X. Tie et al.





2.2 WRF-Chem model

An important objective of this study is the intensive use of a regional chemical/dynamical model to compare with the measurements in order to evaluate and eventually improve the model. The evaluated model results are also used to study the chemical evolution of the Mexico City plume. The model used in this study is a recently developed regional chemical/transport model (WRF-Chem, version 2). The Weather Research and Forecasting (WRF) model is a mesocale numerical weather prediction system designed to serve both operational forecasting and atmospheric research needs. The effort to develop WRF has been a collaborative partnership, principally among the National Center for Atmospheric Research (NCAR), the National Oceanic and Atmospheric Administration (NOAA), the National Centers for Environmental Prediction (NCEP), the Forecast Systems Laboratory (FSL), the Air Force Weather Agency (AFWA), the Naval Research Laboratory, Oklahoma University, and the Federal Aviation Administration (FAA). The WRF model is a fully compressible and Euler nonhydro-

- static model. The detailed description of WRF model can be found in the WRF web-site http://www.wrf-model.org/index.php. In addition to dynamical calculations, a chemical model is coupled on-line with the WRF model (WRF-Chem). A detailed description of WRF-Chem is given by Grell et al. (2005), and is used here with some modifications in the chemical scheme (Tie et al., 2007a). The version of the model, as used
- in the present study, includes on-line calculation of dynamical inputs (winds, temperature, boundary layer, clouds etc.), transport (advective, convective, and diffusive), dry deposition (Wesely et al., 1989), gas phase chemistry, radiation and photolysis rates (Madronich and Flocke, 1999; Tie et al., 2003a), and surface emissions (including an on-line calculation of biogenic emission). The ozone formation chemistry is repre-
- sented in the model by the RADM2 (Regional Acid Deposition Model, version 2) gas phase chemical mechanism (Chang et al., 1989) which includes 158 reactions among 36 species. In this study, the model resolution measured 6×6 km in the horizontal direction, in a 900×900 km domain centered on Mexico City. The model ran from 1

ACPD

9, 9221-9266, 2009

Simulation of Mexico City plumes

X. Tie et al.





March to 30 March, 2006, and the results along flight tracks were selected for analysis. The hourly emissions for 17 chemical species, including SO_2 , CO, NO, and 14 different VOCs are shown in Table 1 of Tie et al. (2007a).

In previous studies, the WRF-Chem model was evaluated by comparing calculated

- O₃, CO, and NO_x to the ground measurement in the Mexico City (Tie et al., 2007a; Ying et al., 2009; Zhang et al., 2009). The model results showed that the magnitude and timing of simulated diurnal cycles of O₃, CO, and NO_x, and the maximum and minimum O₃ concentrations were generally consistent with surface measurements. The simulated chemical species concentrations compare favorably with the observations
 with differences between calculated and measured values of 10–30%. However, there was no systematic comparison of the model result with the MILAGBO field campaign
- was no systematic comparison of the model result with the MILAGRO field campaign, and will be a focus point in this study.

3 Model calculations

3.1 Transport evaluation for calculated CO at surface and in the plume

In order to evaluate model calculations of ozone production in the city plume, one important aspect is to understand the transport of the model from the surface and the planetary boundary layer (PBL) to the free troposphere. In this study, we use the calculated CO both at the surface and in the plume aloft to evaluate the model transport. The chemical lifetime of CO is of order of months, thus over the few days relevant here it can be considered as a passive tracer which is emitted from the surface, mixed in the PBL, and transported with prevailing winds.

3.1.1 Evaluation of calculated surface CO

The surface CO concentrations in the Mexico City have been monitored routinely since 1986 by the local air quality monitoring network (RAMA) (Molina and Molina, 2002).



In the first step of the evaluation, the calculated surface CO concentrations are compared with the measured CO concentrations of this network to get some insight of modeled CO emissions and transport processes in the PBL of Mexico City. Figure 2 shows the average CO concentrations in Mexico City during the 6 flight days (8, 10,

- ⁵ 18, 22, 28, and 19 in March). The comparison between calculated and measured spatial distribution of CO concentrations shows that the calculated CO concentrations are generally consistent with observations. For example, both the calculated and measured CO concentrations are higher in the center of the city with a value near 2 ppmv, and decreased in the edge of the city with a value of 1.0 ppmv. However, discrepan-
- cies between the calculation and measurement are also evident. The calculated CO concentrations are slightly higher than the measured values in the south of the city by about 0.2 to 0.4 ppmv. Figure 3 shows the comparison between calculated and measured diurnal cycles of CO concentrations averaged over all urban surface stations. Peak morning values of CO in both the calculated and measured results range from
- ¹⁵ 3 to 5 ppmv. The calculated minimum CO concentrations during noontime are also consistent with the measured concentrations with values of about 1 ppmv. However, it is evident that the calculated CO concentrations are overestimated in the evening. The overestimated values are highest on 8 March and lowest on 28 March, highlighting that there is a strong variability during different days. As suggested by Tie et al. (2007a),
- the diurnal variation of surface CO concentrations is mainly controlled by the daily variability of PBL height and emission of CO. In the early morning, emissions rapidly increase mainly due to an increase in mobile (vehicular) emissions, while the PBL is still shallow. As a result the emitted CO is confined near the surface, producing high CO concentrations. As morning progresses, the PBL height increases allowing rapid
- dilution of CO concentrations. In the evening, the peak CO concentrations occur again as a result of increase in emissions and a slowing of PBL mixing. Because there are no day to day variations of emission of CO in the model, the strong variation in different days is mainly due to the variability of different diurnal patterns of the PBL heights, suggesting that the decrease in the PBL height in the evening (esp. for 8 March) is

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Introduction Abstract Conclusions References **Figures** Þ١ Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion



faster in the model than the real situation. As a result, freshly emitted CO is retained in a shallow PBL, resulting in an overestimation of the surface CO concentrations. On 18 March, the calculated PBL height during evening time is better represented, and the overestimated surface CO concentrations are reduced. Overall, the calculated spatial distributions and magnitudes of surface CO concentrations are reduced.

distributions and magnitudes of surface CO concentrations are generally consistent with the measured values. The calculated diurnal variations are consistent with the measured results during morning and noontime, but overestimated during evening.

Figure 4 shows an overall comparison of daily average CO concentrations of the model with measured values. The result indicates the discrepancies (model – measurement) of CO surface concentrations range from -6 to +20%, with an average discrepancy of about +6%.

3.1.2 Evaluation of calculated CO plumes

The calculation of CO concentrations in the plumes is more complex than the calculation of its surface concentrations. The major determining factors for calculating ¹⁵ the surface concentrations are surface emissions and mixing in the PBL (Tie et al., 2007a). For calculating plume CO levels, addition factors must be considered including the transport of CO across the PBL and the calculation of the prevailing winds. The upper panels of Fig. 5 show that the calculated CO has a variability that is broadly consistent between model and observations for all the flights. For example, during the

- flight of 10 March, the city plume (high CO concentrations) is detected between 15 and 17 pm with a large variation. The calculated CO variability during this period is similar to the measured results, showing the capability of the model in capturing this high variable city plume. Another example is shown during the flight of 29 March, where both observations and model show evidence of high CO concentrations in the city plume for
- a short time (from 13:30 to 14:30 p.m.) with values rapidly increasing from background (about 100 ppbv) to about 600 ppbv. By contrast, the flight route of 28 March is mostly located upwind of the city and both calculation and measurement show very low CO concentrations, as no city plumes are detected. The flight of 28 March is included in

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Figures** Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

this study to get some insights regarding the background atmosphere.

In these 6 flights, city plumes are detected on 8, 10, 18, 22, and 29 March. To get more detailed estimates for the calculated city plume, in the following sections we will separate the concentrations into 2 categories in the 6 flights, including (a) the averaged

- values to represent the city plumes and (b) the averaged values to represent the background values in both the calculation and measurement during each of the flights. The criterion for a city plume encounter is either that a flight path passes directly over the city or through the downwind plumes as determined by rapidly increasing measured CO concentrations as indicated in Fig. 1. Otherwise, it is considered as background
- ¹⁰ condition. This criterion is also applied for other species (such as O_3 , NO_y , NO_x , and VOCs) which showed elevated values correlated with the CO plumes. Based on this separation, the lower panels of Fig. 5 show that the averaged CO concentrations in the city plumes range from 200 to 350 ppbv and the averaged background CO concentrations range from 100 to 150 ppbv. The calculated CO concentrations in both the city
- ¹⁵ plumes and in the backgrounds are systematically lower than the measured values. For example, the differences between the calculated and measured CO concentrations in the city plumes are -27, 2, -9, -24, and -25% for the flights on 8, 10, 18, 22, and 29 March, respectively. The averaged difference is -18%. As described before, the calculated surface CO concentrations are generally consistent with the measured val-
- ²⁰ ues, and even slightly higher than the measurements. The systematic underestimates of the CO concentrations in the city plume suggest that the model is underestimating transport processes (diffusion, vertical and horizontal transport) from the surface to the plume levels (sometimes in the PBL and sometimes outside of the PBL), and possibly overestimating mixing with cleaner background air. As a result, this underestimation
- is likely a systematic transport error for which some adjustment can be made. In the following sections, this transport error is applied to other chemical species that have a fairly long lifetime (such as O_3 , NO_y , NO_z , etc.) to get adjusted values. The adjusted values can reduce this transport error, and provide insights for other sources of errors (such as emission, etc.). An adjusted concentration for species (*X*) in the plumes

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Introduction Abstract Conclusions References **Figures** Þ١ Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

during different flights can be calculated by the following method:

 $X_{adi}(i) = X(i) * (1 + A(i)/100)$

where X(i) represents the modeled un-adjusted plume concentration of a chemical species (such as NO_v) during the *i*th flight (for example, flight of 8 March); $X_{adj}(i)$ represents adjusted concentration of the chemical species; A(i) represent an adjust-5 ment factor due to underestimates of the transport from the surface to the city plumes in different flights. According to the above study of CO, the factors are 27, -2, 9, 24, and 25% for the flights of 8, 10, 18, 22, and 29 March, respectively.

3.2 Evaluation of O₃ concentration during the flights

- The in-situ comparison of the calculated O_3 (and other species) to the measured values 10 along the flight tracks suggests that the model is capable of capturing the variability of O₃, showing a rapid enhancement when city plumes are detected. In the following sections, we will show the one by one comparison between the calculated and measured mean values in both the city plumes and in the backgrounds during different flights.
- This comparison illustrates the systematic bias of the calculations, and the correlations 15 between the calculations and measurements.

Figure 6 shows the one by one comparison for O_3 and other species. Table 1 gives the calculated and measured mean values in both the city plumes and backgrounds, and also correlations between the calculated and measured values, and ratio between

- the background and city values. The results suggest that the calculated O_3 is consis-20 tently underestimated in both the city plumes and in background compared to the measured O₃ concentrations. The averaged underestimates are 16% in the city plumes, and 7% in the background. One reason for the underestimates of O_3 concentrations is attributed to the underestimates of the transport processes. The effect of these under-
- estimates can be reduced by application of Eq. (1) to get adjusted O_3 concentrations. 25 However, CO is determined mostly by transport, but O₃ is also depends non-linearly on NO_x . Before the applying Eq. (1), the relationship between $D[CO](CO_{CAI} - CO_{MFA})$ and

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Introduction Abstract Conclusions References **Figures** Þ١ Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

(1)



D[O₃](O_{3CAL}-O_{3MEA}) needs to be examined. D[CO] and D[O₃] represent the discrepancies between the calculated and measured CO and O₃ during different flights. If the discrepancies in CO and O₃ are positively correlated, the underestimates of O₃ are at least partly attributable to the transport error. Figure 7 shows that D[CO] and D[O₃] are correlated to each other with a correlation coefficient of 0.7. Thus, the adjustment for O₃ concentrations can be applied by using Eq. (1). After this adjustment, the difference between the calculated and measured O₃ concentrations is improved (see Fig. 6) from about 16% to 1%, which suggests that the underestimates of O₃ concentrations in the city plumes is largely a result of the underestimates of the transport from the surface to the city plumes.

3.3 Evaluation of nitrogen species during the flights

The evaluated nitrogen species in this study include NO_x (NO+NO₂), NO_y (all nitrogen species), and NO_z (NO_y-NO_x). The model captures the timing/location of the city plume as well as the strong variability seen in the measurements (not show). However,
 the calculated NO_y concentrations are consistently smaller than the measured NO_y concentrations in both the city plumes and the backgrounds. The averaged underestimates are 40% in the city plumes, and 40% in the backgrounds (see Fig. 6 and Table 1). One of the reasons for the underestimates of the city plume concentrations is related to the large underestimates of the background values (by about 0.8 ppbv). Another
 reason results from the underestimation in the transport from the surface to the downwind plumes, as already discussed for CO. In order to take into account this transport underestimate, Eq. (1) is applied for the NO_y concentrations in the city plumes. With this consideration, the underestimates of NO_y concentrations in the city plumes reduce from 40 to 30%. Furthermore, if the underestimates of the background NO_y concentrations in the city plumes reduce from 40 to 30%.

trations (with an averaged value of 0.8 ppbv) is added in the calculated NO_y concentrations, the underestimates of plume NO_y concentrations decrease from 30 to about 20%. This remaining underestimate may then be mainly attributed to uncertainties in surface emissions of NO_x .



As suggested by Godowitch et al. (2008) and Fischer et al. (2003), NO_{z} ($NO_{v}-NO_{x}$) concentrations and the correlation between NO₇ and O₃ are good indicators for the rate of O₃ production in differently aged city plumes. Thus, the calculated NO₇ concentrations are evaluated by comparing to the measured NO₇ concentrations in the flights (shown in Fig. 6 and Table 1). The results suggest that the simulated NO_7 variability 5 is consistent compared with the measured results. High concentrations of NO₇ are seen in the city plumes, while during the flight of 28 March (background case) both the calculated and measured NO₂ concentrations are low with concentrations near 1.0 ppbv. However, the calculated NO₇ concentrations are consistently smaller than the measured concentrations in both the city plumes and backgrounds. The calculated 10 NO₇ concentrations in the city plumes are underestimated by 40%. Because NO₇ is a long-lived chemical species like NO_v, there is a possibility that the underestimate is also partially resulting from the underestimate of transport from the surface to the city plume. After the Eq. (1) is applied for the NO_z concentrations in the city plumes, the underestimates of NO₇ concentrations in the city plumes reduces to 28%. Furthermore, if the underestimate of the background NO₇ concentrations (with an averaged value of 0.7 ppbv) is added in the calculated NO₂ concentrations in the city plumes, the underestimate of NO_z concentrations reduces to 17%, which is generally the same error to

20

the NO_v calculations.

The calculated NO_x concentrations have a stronger variability than the calculated CO, O_3 , and NO_v . The enhancement of NO_x concentrations in the city plumes is much higher than the NO_v concentrations. For example, the ratios between background concentrations and city plume concentrations and are 0.2 and 0.1 for NO_v and NO_x, respectively. Unlike CO and NO_v, NO_x has a very short chemical lifetime (from hours

to days) in different altitudes (Ridley et al., 1996; Tie et al., 2007b). Because there are 25 rapid and complex chemical reactions inside the city plumes, the calculation of NO_x in the city plumes is more difficult than the less reactive species, such as CO and NO_v. As a result, a large variability occurs for the NO_x calculations. For example, during flight of 10 March, the calculated NO_x concentrations are close to the measured values

ACPD

9, 9221-9266, 2009

Simulation of Mexico **City plumes**

X. Tie et al.





with difference of 7%. However, during the flight of 29 March, the calculated NO_x concentrations are underestimated by 70% (not shown). In general, the calculated NO_x concentrations are consistently underestimated in all flights, with the averaged underestimate of 50%. The underestimate is partially due to the underestimate of surface emissions of NO_x as suggested by the NO_y analysis (about 20% underestimation of surface emissions). However, the large variability in calculated NO_x errors during different flights suggests that the calculated errors are also attributed to the complication of the interaction between fast chemical reactions and transport processes from the surface to the plumes.

10 3.4 Evaluation of VOC species during the flights

As suggested by Sillman (1995), VOCs are considered as the initiators for the ozone production process. In VOC-limited regions, the amounts of VOCs are a crucial factor in controlling the rate of O_3 production. Because there are different types of VOCs, the rate of ozone production is not only determined by the concentrations of VOCs, but also the rate of oxidation of VOCs. Most VOCs are first oxidized, mainly by OH with a rate coefficient of k_{OH+X} . The reactivity of OH with various VOC compounds (X), i.e. $k_{OH+X}(X)$, is a key parameter to determine the contributions of an individual VOC to the rate of O_3 production. For example, alkane species generally react with OH with slower rate constants, but the concentrations are normally high in cities. By contrast, aromatic species generally react with OH at factor rate constants, but the concentrations are normally high in cities.

- aromatic species generally react with OH at faster rate constants, but the concentrations are normally lower than alkanes. As a result, both the VOC species could be important for ozone productions in the city plumes. Altough VOC reactions are generally numerous and complex, the chemical scheme of WRF-Chem (RADM2) simplifies the VOC reactions into a relatively smaller set. For example, several alkane species
- are lumped into one alkane with a mean reaction rate. Thus, some assumptions are needed to match the calculated VOC species to measured VOC species. According to the availability and reliability of the VOC data collected in the aircraft, the following VOC species are evaluated in the model calculation, i.e. a surrogate alkane (alkanes)



with OH reaction coefficient less than 6.8×10^{-12}), an aromatic species (toluene), and oxygenated VOC species (HCHO and aldehydes).

The calculated toluene concentrations are in good agreement with the measured values (see Fig. 6). The enhanced concentrations in the city plume are well captured.

- The averaged calculated and measured toluene concentrations in the city plume are 2.2 and 2.1 ppby, respectively. The correlation coefficient between the calculated and measured values is 0.99 (see Table 1). Background concentrations are very low, suggesting that toluene originates mainly from urban emissions, and the high concentrations in the city plumes decrease rapidly due to fast reaction with OH. The ratio between
- the background and city is about 0.12. 10

As described above, the alkane in the WRF-Chem model is a surrogate compound with an OH reaction coefficient of less than 6.8×10^{-12} . As a result, the measured alkane species which have the reaction coefficient of $<6.8 \times 10^{-12}$ were summed for comparison with the surrogate alkane (Fig. 6 and Table 1). The results suggest that the

- calculated alkane is consistently higher than the measured values, showing a strong 15 urban enhancement. The averaged alkane concentrations for the calculations and measurements in the city plumes are 15 and 24 ppbv, respectively. However, the calculated background alkane concentrations are generally closer to the measured values, respectively 1.6 and 1.9 ppbv. The overestimated alkane concentrations in the city plumes are likely attributed to the overestimate of the surface emission of alkanes 20
- in the model, because this difference can not be explained by the transport adjustment or errors in background concentrations.

Unlike the toluene and alkanes, which are primarily emitted from the surface, the oxygenated VOCs (OVOCs, such as HCHO and aldehydes) are not only emitted, but also chemically produced in the plumes. As a result, the OVOCs could play relatively

25 more important roles in O₃ production in the downwind region of plumes. The result shows that the calculated HCHO concentrations are slightly underestimated during the flights of 18 and 22 March, but are overestimated during the flight of 29 March. The averaged concentrations for the calculations and the measurements are 3.5 and 3.8 ppbv

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Introduction Abstract Conclusions References **Figures** Þ١ Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

in the city plumes, with about 7% underestimate. The background concentrations are also slightly underestimated, with averaged concentrations of 0.6 and 0.8 ppbv for the calculations and measurements, respectively. The correlation coefficient between the calculated and measured values is relatively high, with a value of 0.90.

- ⁵ Another important OVOC species is aldehyde, for which the reaction rate with OH is higher than the rate of HCHO. In the city plumes, the measured aldehyde concentration is 3.1 ppbv, which is comparable with the measured value of HCHO (3.8 ppbv). With a higher reaction rate, aldehyde could play relatively a more important role in O₃ production in downwind region of plumes. The calculated aldehyde concentrations in
- the city plumes are overestimated compared to the measured values. However, the calculated background aldehyde concentrations are slightly underestimated compared with the measured values. The averaged concentrations for the calculations and the measurements are 3.1 and 4.1 ppbv in the city plumes, respectively. The averaged background concentrations are 0.51 and 0.70 ppbv for the calculations and measurements, respectively. The correlation between the calculated and measured values is
 - very good, with a coefficient of 0.99.

4 Analysis of the O₃ production in the plumes

The production of O₃ depends non-linearly on the concentrations of precursors (NO_x, CO, primary VOCs and secondary VOCs) (Kleinman et al., 2000). As a concentrated city plume leaves from city and the primary chemicals start to be diluted, the O₃ production rate and efficiency is expected to vary non-linearly. Furthermore, the concentrations of oxygenated VOCs (such as HCHO and aldehydes) are continuously produced and destroyed during outflow. Because the OVOCs are also important O₃ precursors (Singh et al., 1995; Apel et al., 2008), OVOC oxidations lead to additional O₃ production along plumes. In this study, we will focus on 3 issues regarding O₃ production

along the Mexico City plume; (1) what is the O_3 production behavior at different ages of the plume; (2) what is the O_3 production efficiency (molecules of O_3 produced per NO_x

AC 9, 9221–9	ACPD 9, 9221–9266, 2009					
Simulation of Mexico City plumes X. Tie et al.						
Title	Title Page					
Abstract	Introduction					
Conclusions	References					
Tables	Figures					
14	►I					
•	•					
Back	Close					
Full Scr	Full Screen / Esc					
Printer-friendly Version						
Interactive Discussion						

molecule consumed) in different plume ages; and (3) what are the roles of different O_3 precursors (such as CO, alkanes, OVOCs, etc.) in the O_3 production in different ages of the plumes. First, we define the chemical "age" of plume.

4.1 Chemical age of the city plume

⁵ In order to analyze the chemical characterizations in different chemical ages of the Mexico City plume, we define the chemical age of the city plume by

 $AG = -\ln(NO_x/NO_y)$

The conversion from NO_x to NO_y results mainly from the reactions of NO_x \rightarrow HNO₃ and NO_x \rightarrow organic nitrates. In addition, there is also an important pathway to convert NO_x to NO_y through the hydrolysis of N₂O₅ on the sulfate aerosols. However, this later reaction becomes important only when sulfate aerosol loading is high and solar radiation is low (e.g., at high latitudes or during nighttime) (Tie et al., 2003b; 2005). In Mexico City, the later reaction to convert NO_x to NO_y can be neglected. In some of the following discussion, we will define plumes as young, middle-aged, or old depending on whether the value of AG is 0–1, 1–2, or larger than 2, respectively.

4.2 O₃ and NO_x relationship

Ozone precursors such as CO, NO_x, and VOCs are found to be closely correlated to O₃ concentrations. First, we will show the O₃–NO_x correlation with different chemical ages (see Fig. 8). It shows that in the young plume, which is normally near the city area, O₃ concentrations are not very sensitive to NO_x concentrations. The ratio of Δ [O₃]/ Δ [NO_x] (the slope of the fits shown in Fig. 8) is only about 4.7. By contrast, O₃ concentrations are very sensitive to NO_x concentrations in the aged plume. The ratio of Δ [O₃]/ Δ [NO_x] rapidly increases to 30 in the aged plume, indicating that the slope of Δ [O₃]/ Δ [NO_x] is about 7 times higher in the aged plume than in the young plume. The calculated slope of Δ [O₃]/ Δ [NO_x] is consistent with the measured result. Because the aircraft

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Figures Tables** ►I. Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

(2)

measurements did not generally follow the city plumes, the measured results can be considered as the values of NO_x and O₃ measuring in different locations (Euclidian domain). The O₃-NO_x relationship in different chemical ages can also be considered as the relationships presenting for different spatial locations (such as nearby cities, remote areas). The analysis of O₃-NO_x relationship in this study suggests that the slope of Δ [O₃]/ Δ [NO_x] is lower in nearby city than in the remote areas.

4.3 O₃ and CO relationship

Parrish et al. (1998) studied O_3 -CO correlation and indicated that there were different slopes of $\Delta[O_3]/\Delta[CO]$ under different conditions (locations), with a larger slope of O_3 -CO correlation implying a higher O_3 production efficiency. In this study, the O_3 -CO correlation is shown in Fig. 9. The data in this study is somewhat different with the previous studies. For example, Parrish et al. (1998) studied the O_3 -CO correlation in 5 surface sites. By contrast, the data used in this study is collected from aircraft which is covered a larger spatial domain, including different chemical conditions (city plumes, backgrounds, young, and aged cases). In other words, the O_3 -CO correlation in this study can be considered a combined result from multiple local sites. The result of this study suggests that the O_3 -CO correlation is in a non-linear relationship. When the concentrations of CO are less than 400 ppbv, the measured slope of $\Delta[O_3]/\Delta[CO]$ is large, with a value of 0.175. While the concentrations of CO are greater than 400 ppbv,

- ²⁰ the measured slope reduced to 0.05. As we described before, the data used to calculate O_3 -CO correlation covers different aged plumes (as shown in Fig. 2). If we assume that the high CO concentration conditions represent young city air and low concentrations aged background conditions, the high slope of O_3 -CO correlation suggest that additional O_3 is produced in the aged plume. Otherwise, the slope of O_3 -CO
- ²⁵ correlation should be the same in different ages of plumes. This result is consistent with the study by Wood et al. (2008) who found that the ratio $\Delta[O_3]/\Delta[CO]$ increases with the age of plumes, suggesting that secondary VOCs are more efficient at producing O₃ in aged plumes. The WRF-Chem calculated ratio of $\Delta[O_3]/\Delta[CO]$ is consistent



with the measured result. However, the slope of the $\Delta[O_3]/\Delta[CO]$ in the young plume is overestimated. In the aged plume, the calculated slope is very similar to the measured value. Because this non-linearity relationship, the correlation coefficient between O_3 and CO is not very high, with values of 0.88 and 0.74 for measurement and calculation, respectively.

4.4 O₃ and NO_z relationship

5

 NO_{7} is the product of NO_{7} oxidation, and the rate of oxidation of NO_{7} to NO_{7} is a main factor to determine O_3 production efficiency which is defined by number of ozone molecules produced per molecule of nitrogen oxides consumed (Liu et al., 1987). Because O₃ production is also correlated to NO₇ in a non-linear way (Godowitch et al., 2008; Fischer et al., 2003), the slope of $\Delta[O_3]/\Delta[NO_7]$ provides useful information for the O₃ production efficiency in different chemical conditions. Figure 10 shows the calculated and measured O₃-NO₇ correlations. It shows that when NO₇ concentrations are high nearby the city area (young city plume) O_3 concentrations are not very sensitive to NO₂ concentrations. The ratio of $\Delta[O_3]/\Delta[NO_7]$ is about 3.3. The ratio increases 15 to 5.7 in the low NO_z regime (in the aged plume). According to the study by Zanis et al. (2007), large slope of the ratio of $\Delta[O_3]/\Delta[NO_7]$ indicate high ozone production efficiency. This study is consistent with the result of Zanis et al. (2007). The WRF-Chem calculation shows a similar result to the measurement in the aged plumes. However, the calculated ratio of $\Delta[O_3]/\Delta[NO_7]$ is generally overestimated in the young plumes. 20 A recent study by Shon et al. (2008) also suggests that the ozone production efficiency

4.5 The roles of different precursors for O₃ production in the plume

is higher in remote region than in urban area.

In order to study the causes of the different slopes of $NO_x - O_3$, $CO - O_3$, and $NO_z - O_3$ in

²⁵ different aged plumes, the WRF-Chem model is extensively applied to study the roles of ozone precursors on ozone production along the city plumes. The model has sev-

eral advantages in analyzing ozone production in different aged plumes. First, tracing plumes in the model can be considered as a Lagrangian way for studying chemistries coupled with dilution in plumes. Second, the individual effect of O_3 precursors on O_3 formation can be separately studied in the model.

- ⁵ The methodology used here is to track the city plume by following the high CO column concentrations from the city to downwind regions. For example, the column CO in the flight of 22 March is shown in Fig. 11. From Fig. 11, we see that the outflow of the city plume is predominantly toward the northeast. The cross section (altitude vs. downwind distance) of O_3 concentrations in the plume (lower panel of Fig. 11) suggests that
- ¹⁰ the pollutants are not limited to the PBL with plumes reaching about 3–4 km above the surface (the calculated PBL height in the model ranges 1.5 to 2 km on this day). The calculated O_3 concentrations in the plume show that there is a maximum concentration near the city (plume age near 1.6) with concentrations of more than 100 ppbv. There is also a secondary O_3 maximum at 300–400 km downwind of the city. At this location,
- ¹⁵ the O₃ concentrations have another maximum of 75 ppbv (the plume age is about 4.6). In order to estimate the causes for the ozone maximum in the aged plume, the reactivity of OH with various compounds *X*, i.e. $k_{OH+X}(X)$ is calculated. These chemical compounds include CO, slow reactive VOCs (alkanes), very reactive VOCs (alkenes and aromatic), and OVOCs (oxygenated VOCs including cresol, formaldehyde, ke-
- tones, glyoxal, methyl glyoxal, unsaturated dicarbonyl, peroxyacetyl nitrate, and organic nitrate). Figure 12 shows the reactivity of OH with these 4 types of compounds. The result shows that: (1) The CO reactivity is significant both in the city area and in the aged plume, but its relative importance in the urban area is less than the other types of reduced compounds. This is because although the OH reaction with CO is a ma-
- jor chemical destruction for CO, and the rate constant is relatively small. As a result, large amounts of CO are transported from the city to the downwind area. For example, the CO concentration is about 300 ppbv at 300 km along the plume (not shown), and the high CO concentrations in the aged plume result in the high OH reactivity in the aged plume. (2) The rate constants for OH+alkenes and OH+aromatics are

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Introduction Abstract Conclusions References **Figures** Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion

about 20–100 times higher than the rate constant for OH+CO. The fast rate constants lead to high OH reactivity in the city area, making these precursors the major ones for ozone production within Mexico City (Madronich, 2006; Tie et al., 2007a). As a result, a large portion of alkenes+aromatic is chemically destroyed near the city, and thus
less is transported downwind leading to the small OH reactivity in the aged plume. (3) OVOCs are mainly produced by chemical reactions in the outflow, including the reactions of OH with alkenes, aromatic, etc. and have important contributions to the OH reactivity near the city as well as in the aged plume. (4) The rate of alkane reaction with OH is relatively low compared to alkenes or aromatic. In Mexico City area, the alkane concentrations are lower than CO (Velasco et al., 2007; Tie et al., 2007a), although the rate of reaction with OH is higher than CO. As a result, alkanes contribute to the OH reactivity in both young and aged plumes, but are usually not the dominant species in either.

4.6 The O₃ production efficiency in the plume

The oxidization of CO and VOCs leads to the production of HO_2 an RO_2 radicals. Thus O_3 is produced in the presence of NO by following reactions:

$HO_2 + NO \rightarrow NO_2 + OH$	(R1)
$RO_2 + NO \rightarrow NO_2 + RO$	(R2)
$NO_2 + hv \rightarrow NO + O$	(R3)
$O + O_2 \rightarrow O_3$	(R4)

As a result, O_3 is produced through the (R1) to (R4) and the rate-determining steps are (R1) and (R2). The NO is re-generated in reaction (R3), so that it can participate in this O_3 -producing cycle more than once. At high NO_x, the main termination of this cycle is the reaction

 $NO_2 + OH \rightarrow HNO_3$



(R5)

The O_3 production efficiency (S) was defined by Liu et al. (1987) as

 $S = P(O_3)/L(NO_x)$

where $P(O_3)$ and $L(NO_x)$ are the production of O_3 and the loss of NO_x , respectively. According to the studies by Liu et al. (1987) and Lin et al. (1988), the advantage to use the O_3 production efficiency is that the variations in O_3 production and NO_x loss are often similar and therefore cancel each other. The O_3 production efficiency (*S*) is the ratio between O_3 production and NO_x loss, which reduces substantially the uncertainty in the estimated total regional O_3 production. However, significant nonlinearity still exists (Liu et al., 1987). Lin et al. (1988) used both absolute O_3 production and net O_3 production (including O_3 chemical production and the effect of transport) for the calculation of the O_3 production efficiency, and found that the O_3 production. In this study, we use the absolute O_3 production calculated from the reactions (R1) and (R2) to study the O_3 production efficiency along the city plume, which is expressed by

¹⁵
$$S = (P[HO_2] + P[RO_2])/(k_9 \times [NO_2][OH])$$

(3)

(5)

(6)

 $P[HO_2] = (k_5 \times [HO_2][NO])$

 $P[\mathrm{RO}_2] = (\Sigma k_{6i} \times [\mathrm{RO}_2]_i [\mathrm{NO}])$

25

Where $P[HO_2]$ and $P[RO_2]$ represent the O₃ production due to the oxidation of NO with HO₂ and RO₂, respectively. Note that there are several types of RO₂, and thus [RO₂]_{*i*} represents different RO₂ species, which are mainly produced by the oxidations of VOCs with OH reactions. The parameters of k_5 , k_{6i} , and k_9 are the rate coefficients for (R1), (R2), and (R5), respectively.

The calculated O_3 production efficiency (*S*) along the city plume is shown in Fig. 13a (upper panel). The O_3 production efficiency is higher in the aged plume with a maximum value of 60–80 and is lower near the city with a minimum value of 10, suggesting that the O_3 production is about 6 times more efficient in the aged plume than in the



young plume. This result explains that the secondary O_3 maximum in the aged plume shown in Fig. 11 results from the facts that: (1) There are high OH reactivities, largely due to the OVOCs oxidation, leading to the production of large amount RO_2 . (2) The O_3 production efficiency is much higher in the aged plume than in the young plume, leading to more production of O_3 molecules with each NO_x molecular being consumed in the aged plume. As a result, even though the NO_x concentrations are low, a fairly amount of O_3 is produced in the aged plumes. The high O_3 production efficiency is associated with lower concentrations of NO_x . As a result, the high O_3 production efficiency cannot be considered as high O_3 production rate. For example, the lower panel of Fig. 13a shows that although the production efficiency is lower in the city, the rate of O_3 production is highest (30 ppbv/h). By contrast, the rate of O_3 production is low in the aged plume (0.3 ppbv/h), although the production efficiency is highest in this region.

From Eq. (3), there are 2 important components in contributing to the O_3 production and production efficiency. One is through the HO₂+NO reaction ($P[HO_2]$), and another is through the RO₂+NO reaction ($P[RO_2]$). Because HO₂ and RO₂ result from different oxidations processes (e.g., CO oxidation produces only HO₂ radicals, while VOC oxidation produce both HO₂ and RO₂ radicals), it is important to understand the individual contributions of HO₂ and RO₂ to the O₃ production and production efficiency.

Figure 13b shows the individual contribution of $P[HO_2]$ and $P[RO_2]$ for the O₃ production and production efficiency.

$$S_{\text{HO}_2} = P[\text{HO}_2]/(k_9 \times [\text{NO}_2][\text{OH}])$$
(7)

$$S_{\text{RO}_2} = P[\text{RO}_2]/(k_9 \times [\text{NO}_2][\text{OH}])$$

The result shows that both the S_{HO_2} and S_{RO_2} are lower nearby the city (young plume) than the aged plume. For example, values of S_{HO_2} and S_{RO_2} are about 10 and 5 in the young plume and 20 to 50 in the aged plume, respectively. However, there is an important difference between the values of S_{HO_2} and S_{RO_2} . The high values of S_{RO_2} are mostly located in the aged plume (as shown in lower panel). By contrast, the

(8)

high values of S_{HO_2} are distributed throughout the entire plume. The wide distribution of S_{HO_2} can be explained by the fact that CO concentrations are widely distributed because of its long chemical lifetime. As a result, the HO₂ concentrations are more uniformly distributed, while the large RO₂ concentrations are mostly found in the plume. This result suggests that the high O₃ production efficiency in the aged plume (showing in Fig. 13a) is largely resulted from the RO₂+NO reactions and a small portion from the HO₂+NO reaction. Figure 13c shows the noontime concentrations of OH, HO₂, and RO₂ along the city plume. The results indicate that OH concentrations are lower insider the plume; HO₂ concentrations are more uniformly distributed; and RO₂ concentrations are enhanced inside the plume. This result suggests that the air pollutants emitted from the city intend to convert OH to RO₂ inside the plume. There is also clear indication that the largest enhancement of RO₂ occurs at aged plume region, which results in highest O₃ production efficiency in this region (see the lower panel of Fig. 13b).

5 Summary

- ¹⁵ The O₃ concentration and its evolution in the Mexico City plume during the MIRAGE-Mex field campaign were studied. The in-situ aircraft measurements of O₃, CO, VOCs, NO_x, and NO_y concentrations were compared to the calculations of a regional chemical/transport model (WRF-Chem). The model calculation was also used to interpret the measured results, and to understand the O₃ evolution in the downwind of the Mex-
- ico City plume. The comparisons between the calculations and measurements can be summarized by the following aspects: (1) The WRF-Chem model has a capability to capture the measured city plumes, and the calculated variability along the flight routes is generally consistent with the measured results, showing a rapid enhancement when city plumes are detected. (2) From the evaluation of CO concentrations (considered as
- a chemically inactive tracer) in both the surface measurements and the aircraft measurements, the transport from the surface of the city to the plume levels (ranging from 3 to 5 km) is underestimated by about 0 to 25% during 5 different flights. This contributes



to the underestimation of some long-lived chemical species (CO, O_3 , NO_y , etc.) in the plumes. (3) The uncertainties in calculating CO, O_3 , NO_x , VOC concentrations in the city plumes are -50 to +20%. After adjustment for the transport processes based on CO concentrations and background concentrations, the uncertainty range is reduced to

about -20 to +20% which is probably mostly related to the surface emission estimates, and other uncertain processes.

The results show that OH reactivity with CO is low in the city area (the young plume), but increases in the aged plume. This is because the OH reaction with CO is a major chemical destruction for CO, and the rate of reaction is low. As a result, large amounts of CO are transported from the city to the downwind area, resulting in ongoing OH reactivity. Alkenes and aromatic reaction rates with OH were high. As a result, the

fast reaction led to high OH reactivity in the city area, and low OH reactivity in the aged plume. OVOCs are mainly chemically produced by chemical reactions. Thus, OVOCs are continuously produced along the city plume, and have important contribu-

- ¹⁵ tions to the OH reactivity near the city as well as in the aged plume. Alkanes contribute to the OH reactivity in both young and old plumes, but usually are not the dominant species in either. The analysis of measurements (O_3 -CO, O_3 -NO_x, and O_3 -NO_z correlations) and the calculation of the O_3 production efficiency (*S*) showed that the ozone production was more efficient in the aged plume than in the young plume, indicating that substantial O_3 production occur downwind of the city, adding to the O_3 already
- produced in the city and exported with the plume.

10

Acknowledgement. The authors would like to thank George Grell and Steve Peckham at NOAA for providing us with the WRF-Chem model. We wish to thank Geoff Tyndall and Alan Fried for helpful discussions. The National Center for Atmospheric Research is sponsored by the National Science Foundation.

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Figures** Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion



References

- Apel, E. C., Brauers, T., Koppmann, R., et al.: Intercomparison of oxygenated volatile organic compound measurements at the SAPHIR atmosphere simulation chamber, J. Geophys. Res., 113, D20307, doi:10.1029/2008JD009865, 2008.
- ⁵ Brasseur, G., Orlando, J., and Tyndall, G.: Atmospheric Chemistry and Global Change, Oxford University Press, 1999.
 - Crutzen, P. J.: Physical and chemical processes which control the production, destruction, and distribution of ozone and some other chemically active minor constituents, GARP Publications Series, 16, 236–243, 1975.
- ¹⁰ Chameides, W. L. and Walker, J.: Time dependent photochemical model for ozone near the ground, J. Geophys. Res., 81, 413–420. 1976.
 - Chang, J. S., Binkowski, F. S., Seaman, N. L., McHenry, J. N., Samson, P. J., Stockwell, W. R., Walcek, C. J., Madronich, S., Middleton, P. B., Pleim, J. E., and Lansford, H. H.: The regional acid deposition model and engineering model, State-of-Science/Technology, Report
- 4, National Acid Precipitation Assessment Program, Washington DC, 1989.
- de Foy, B., Varela, J. R., Molina, L., and Molina, M.: Rapid ventilation of the Mexico City basin and regional fate of the urban plume, Atmos. Chem. Phys., 6, 2321–2335, 2006, http://www.atmos-chem-phys.net/6/2321/2006/.

Fast, J. D., de Foy, B., Acevedo Rosas, F., Caetano, E., Carmichael, G., Emmons, L., McKenna,

D., Mena, M., Skamarock, W., Tie, X., Coulter, R. L., Barnard, J. C., Wiedinmyer, C., and Madronich, S.: A meteorological overview of the MILAGRO field campaigns, Atmos. Chem. Phys., 7, 2233–2257, 2007,

http://www.atmos-chem-phys.net/7/2233/2007/.

Fischer, H., Kormann, R., Klüpfel, T., Gurk, C., Königstedt, R., Parchatka, U., Mühle, J.,

25 Rhee, T. S., Brenninkmeijer, C. A. M., Bonasoni, P., and Stohl, A.: Ozone production and trace gas correlations during the June 2000 MINATROC intensive measurement campaign at Mt. Cimone, Atmos. Chem. Phys., 3, 725–738, 2003, http://www.atmos-chem-phys.net/3/725/2003/.

Godowitch J. M., Hogrefe, C., and Rao, S. T.: Diagnostic analyses of a regional air qual-

 $_{\rm 30}$ ity model: Changes in modeled processes affecting ozone and chemical-transport indicators from NO_x point source emission reductions, J. Geophys. Res., 113, D19303, doi:10.1029/2007JD009537, 2008. **ACPD**

9, 9221–9266, 2009

Simulation of Mexico City plumes

X. Tie et al.





- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Wilczak, J., and Eder, B.: Fully coupled "online" chemistry within the WRF model, Atmos. Environ. 39, 6957–6975, 2005
- Kleinman, L. I., Daum, P. H., Imre, D. G., Lee, J. H., Lee, Y.-N., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J., and Newman, L.: Ozone production in the New York
- ⁵ City urban plume, J. Geophys. Res., 105, 14495–14512, 10.1029/2000JD900011, 2000.
 Kleinman L. I., Daum, P. H., Lee, Y.-N., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J. Hyde, P., Doskey, P., Rudolph, J., Fast, J., and Berkowitz, C.: Photochemical age determinations in the Phoenix metropolitan area, J. Geophys. Res., 108, 4096, doi:10.1029/2002JD002621, 2003.
- Lei, W., Zavala, M., de Foy, B., Volkamer, R., and Molina, L. T.: Characterizing ozone production and response under different meteorological conditions in Mexico City, Atmos. Chem. Phys., 8, 7571–7581, 2008,

http://www.atmos-chem-phys.net/8/7571/2008/.

15

Lin, X., Trainer, M., and Liu, S. C.: On the nonlinearity of the tropospheric ozone production, J. Geophys. Res., 93(D12), 15879–15888, doi:10.1029/88JD03750, 1988.

Liu, S. C., Trainer, M., Fenhsenfeld, F. C., Parrish, D. D., Williams, H., Fahey, D. W., Hubler, G., and Murphy, D.: Ozone production in the rural troposphere and the implications, for regional and global ozone distributions, J. Geophys. Res., 92, 4191–4207, 1987.

Madronich, S. and Flocke, S.: The role of solar radiation in atmospheric chemistry, in: Hand-

²⁰ book of Environmental Chemistry, edited by: Boule, P., Springer-Verlag, Heidelberg, 1–26, 1999.

Madronich, S.: Chemical evolution of gaseous air pollutants down-wind of tropical megacities: Mexico City case study, Atmos. Environ., 40, 6012–6018, 2006.

Mena-Carrasco, M., Carmichael, G. R., Campbell, J. E., Zimmerman, D., Tang, Y., Adhikary, B.,

D'allura, A., Molina, L. T., Zavala, M., García, A., Flocke, F., Campos, T., Weinheimer, A. J., Shetter, R., Apel, E., Montzka, D. D., Knapp, D. J., and Zheng, W.: Assessing the regional impacts of Mexico City emissions on air quality and chemistry, Atmos. Chem. Phys. Discuss., 8, 20283–20309, 2008,

http://www.atmos-chem-phys-discuss.net/8/20283/2008/.

- ³⁰ Molina, L. and Molina, M. (Eds.): Air Quality in the Mexico MegaCity: An Integrated Assessment, Kluwer Academic Publishers, Dordrecht, 2002.
 - Parrish, D. D., Trainer, M., Holloway, J. S., Yee, J. E., Warshawsky, M. S., Fehsenfeld, F. C., Forbes, G. L., and Moody, J. L.: Relationships between ozone and carbon monoxide at

ACPD							
9, 9221–9266, 2009							
Simulation of Mexico City plumes X. Tie et al.							
Title Page							
Abstract	Introduction						
Conclusions	References						
Tables	Tables Figures						
14	ia Pi						
•	•						
Back	Close						
Full Screen / Esc							
Printer-friendly Version							



surface sites in the North Atlantic region, J. Geophys. Res. 103(20), 13357-13376, 1998.

- Ridley, B. A., Dye, J. E., Walega, J. G., Zheng, J., Grahek, F. E., and Rison, W.: On the production of active nitrogen by thunderstorms over New Mexico, J. Geophys. Res., 101(D15), 20985–21006, 1996.
- ⁵ Shon Z. H., Madronich, S., Song, S.-K., Flocke, F. M., Knapp, D. J. Anderson, R. S., Shetter, R. E., Cantrell, C. A., Hall, S. R., and Tie, X.: Characteristics of the NO-NO₂-O₃ system in different chemical regimes during the MIRAGE-Mex field campaign, Atmos. Chem. Phys., 8, 7153–7164, 2008

Singh, H. B., Kanakidou, M., Crutzen, P. J., and Jacob, D. J.: High concentrations and photo-

chemical fate of oxygenated hydrocarbons in the global troposphere, Nature, 378, 50–54, 1995.

Sillman, S.: The use of NO_y, H₂O₂, and HNO₃ as indicators for ozone-NO_x-hydrocarbon sensitivity in urban locations, J. Geophys. Res., 100, 14175–14188, 1995.

Thielmann A., Prévôt, A. S. H., and Staehelin, J.: Sensitivity of ozone production de-

- rived from field measurements in the Italian Po basin, J. Geophys. Res., 107, 8194, doi:10.1029/2000JD000119, 2002.
 - Tie, X., Madronich, S., Walters, S., Rasch, P., and Collins, W.: Effect of Clouds on photolysis and oxidants in the troposphere, J. Geophys. Res., 108, 4642, doi:10.1029/2003JD003659, 2003a.
- ²⁰ Tie, X., et al.: Effect of sulfate aerosol on tropospheric NO_x and ozone budgets: Evidence during TOPSE, J. Geophys. Res., 108, 8364, doi:10.1029/2001D001508, 2003b.
 - Tie, X., Madronich, S., Walters, S., Edwards, D. P., Ginoux, P., Mahowald, N., Zhang, R. Y., Lou, C., and Brasseur, G.: Assessment of the global impact of aerosols on tropospheric oxidants, J. Geophys. Res., 110, D03204, doi:10.1029/2004JD005359, 2005.
- Tie, X., Madronich, S., Li, G. H., Ying, Z. M., Zhang, R., Garcia, A., Lee-Taylor, J., and Liu, Y.: Characterizations of chemical oxidants in Mexico City: A regional chemical/dynamical model (WRF-Chem) study, Atmos. Environ., 41, 1989–2008, 2007a.

Tie, X., Chandra, S., Ziemke, J. R., Granier, C., and Brasseur, G.: Satellite measurements of tropospheric column O₃ and NO₂ in Eastern and Southeastern Asia: Comparison with

- ³⁰ a global model (MOZART-2), J. Atmos. Chem., 56, 105–125, doi:10.1007/s10874-006-9045-7, 2007b.
 - Velasco, E., Lamb, B., Westberg, H., Allwine, E., Sosa, G., Arriaga-Colina, J. L., Jobson, B. T., Alexander, M. L., Prazeller, P., Knighton, W. B., Rogers, T. M., Grutter, M., Herndon, S. C.,

ACPD						
9, 9221–9266, 2009						
Simulation of Mexico City plumes X. Tie et al.						
Title Page						
Abstract	Introduction					
Conclusions	References					
Tables	Tables Figures					
14	۰					
•	•					
Back	Close					
Full Screen / Esc						
Printer-friendly Version						
Interactive Discussion						



Kolb, C. E., Zavala, M., de Foy, B., Volkamer, R., Molina, L. T., and Molina, M. J.: Distribution, magnitudes, reactivities, ratios and diurnal patterns of volatile organic compounds in the Valley of Mexico during the MCMA 2002 and 2003 field campaigns, Atmos. Chem. Phys., 7, 329–353, 2007,

5 http://www.atmos-chem-phys.net/7/329/2007/.

Wesely, M. L.: Parameterization of surface resistance to gaseous dry deposition in regionalscale numerical models, Atmos. Environ. 23, 1293–1304, 1989.

Ying, Z. M., Tie, X., and Li, G. H.: Sensitivity of ozone concentrations to diurnal variations of surface emissions in Mexico City: A WRF/Chem modeling study, Atmos. Environ, 43, 851– 859, 2009.

10

- Zanis, P., Ganser, A., Zellweger, C., Henne, S., Steinbacher, M., and Staehelin, J.: Seasonal variability of measured ozone production efficiencies in the lower free troposphere of Central Europe, Atmos. Chem. Phys., 7, 223–236, 2007, http://www.atmos-chem-phys.net/7/223/2007/.
- ¹⁵ Zaveri, R. A., Berkowitz, C. M., Kleinman, L. I., Springston, S. R., Doskey, P. V., Lonneman, W. A., and Spicer, C. W.: Ozone production efficiency and NO_x depletion in an urban plume: Interpretation of field observations and implications for evaluating O₃-NO_x-VOC sensitivity, J. Geophys. Res., 108, 4436, doi:10.1029/2002JD003144, 2003.

Zhang, Y., Dubey, M. K., and Olsen, S. C.: Comparisons of WRF/Chem simulations in Mexico

²⁰ City with ground-based RAMA measurements during the MILAGRO-2006 period, Atmos. Chem. Phys. Discuss., 9, 1329–1376, 2009,

http://www.atmos-chem-phys-discuss.net/9/1329/2009/.

ACPD

9, 9221-9266, 2009

Simulation of Mexico City plumes

X. Tie et al.





Table 1. Calculated and measured CO, O_3 , NO_y , NO_z , NO_x , VOCs averaged in the city plume and the background conditions in the flight tracks. Xcm, Xcc, Δ Xc(%), Xbm, Xbc, Δ Xb(%), Rcm, Ym, and Yc represent the mean measured, calculated concentrations, the difference (%) in the city plumes, the mean measured, calculated concentrations, the difference (%) in the backgrounds, the correlation R, and the ratios for the measured and calculated values between the backgrounds and city plumes.

	Mean concentrations (ppbv)			Correlation	Ratio (bckg/plm)				
	Xpm	Хрс	ΔXp(%)	Xbm	Xbc	ΔXb(%)	Rcm	Ym	Yc
CO	274	222	-18	129	88	-31	0.93	0.47	0.40
O ₃	93	78	-16	60	56	-7	0.92	0.65	0.71
NOv	9.7	5.3	-43	2.0	1.2	-41	0.90	0.21	0.22
NOz	6.0	3.6	-40	1.7	1.0	-40	0.71	0.28	0.28
NO _x	3.6	1.7	-52	0.35	0.17	-50	0.60	0.10	0.10
Alkane	15	24	+64	1.9	1.6	-16	0.97	0.12	0.07
TOL	2.19	2.2	+0.4	0.27	0.16	-40	0.99	0.12	0.07
HCHO	3.8	3.5	-7	0.8	0.6	-25	0.90	0.21	0.17
ALD	3.1	4.1	+32	0.7	0.51	-27	0.99	0.22	0.07

ACPD

9, 9221-9266, 2009

Simulation of Mexico City plumes

X. Tie et al.







Fig. 1. Flight route information for the flights of 8, 10, 18, 22, 28 and 29 March with 6 panels. Each panel includes upper (altitudes) and lower (horizontal) projections of flight routes. The black and red lines in the upper panels represent the flight altitudes ($km \times 10$) and CO concentrations (ppbv) along the flights. Rapid enhancements of CO concentrations indicate times when the city plume was detected. The colors in the lower parts show local time during the flights.







ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References Figures **Tables** ١ .∎. ► Close Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion











Fig. 4. Comparison of calculated (green) and measured (blue) surface CO concentrations averaged over each day. The red and yellow bars represent measured and calculated values averaged over all flight days.

ACPD

9, 9221-9266, 2009

Simulation of Mexico City plumes

X. Tie et al.







Fig. 5. Calculated (red lines) and measured (black lines) CO concentrations along the 6 flights (upper panel). The middle panel shows the averaged CO concentrations in the city plumes (see text for city plume definition), and the lower panel shows the averaged CO concentrations in the background conditions. The blue and green bars represent for the measured and calculated values during each flights. The red and yellow bars represent for the measured and calculated values averaged for the all flights.

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Figures** 4 Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion







Fig. 6. Comparison of observed (*x*-axis) and calculated (*y*-axis) concentration of CO, O_3 , NO_y , NO_z , NO_x , toluene, alkanes, HCHO, and aldehydes along C-130 flight tracks. Each point represents the averaged values in the city plume or background air during each flight. The blue line and dots represent adjusted values by using Eq. (1).



Fig. 7. Comparison of model-measurement differences for CO, $D[CO]=CO_{CAL}-CO_{MEA}$, and $D[O_3]=O_{3CAL}-O_{3MEA}$. Each point represents the averaged values in the city plume or background air during each flight.

ACPD 9, 9221-9266, 2009 Simulation of Mexico **City plumes** X. Tie et al. **Title Page** Abstract Introduction Conclusions References **Tables Figures** .∎. ►I. ► Close Back Full Screen / Esc **Printer-friendly Version** Interactive Discussion



Fig. 8. Measured (upper panel) and calculated (lower panel) $O_3 - NO_x$ correlation with different ages of the plume. The black, blue, and red (dots and lines) present the young age (AG-1), middle age (AG-2), and old age (AG-3) plumes, respectively. The lines used the first-order polynomial fits for measured and modeled results.

Fig. 9. Measured (black dots and line) and calculated (blue dots and line) O_3 -CO correlation. The red circle represent the flights are roughly in the city plume regime, and green circle represents the flights are roughly in the background regime. The lines used the second-order polynomial fits for measured and modeled results.

Fig. 10. Same as Fig. 9 except for $O_3 - NO_z$ correlation.

ACPD

9, 9221-9266, 2009

Simulation of Mexico City plumes

X. Tie et al.

Fig. 11. Calculated plume track for the flight of 22 March (upper panel). The contour lines are the calculated CO column concentrations from the surface to 2 km at 12 pm, and the cross dots represent the downwind plume track. The lower panel shows the O_3 (ppbv) cross-section of the plume following the downwind plume track, with two O_3 maxima shown for different chemical ages.

Fig. 12. Calculated OH reactivities (min⁻¹) for CO, alkanes, alkenes+aromatic, and oxygenated VOCs (including cresol, formaldehyde, ketones, glyoxal, methyl glyoxal, unsaturated dicarbonyl, peroxyacetyl nitrate, and organic nitrate) at noontime for the flight of 22 March.

Interactive Discussion

Fig. 13a. Calculated O_3 production efficiency (upper panel) and the rate of O_3 production (ppbv/h) along the plume track at noontime for the flight of 22 March.

Interactive Discussion

Interactive Discussion

ACPD

Mar-22 12PM OH Ca HO2 RO2 0.0 0.1 0.2 0.3 0.5 1.0 10.0 15.0 20.0 30.0 50.0 pptv

ACPD

9, 9221–9266, 2009

Simulation of Mexico City plumes

X. Tie et al.

