

Interactive
Comment

Interactive comment on “First multi-year occultation observations of CO₂ in the MLT by ACE satellite: observations and analysis using the extended CMAM” by S. R. Beagley et al.

M. López-Puertas (Referee)

puertas@iaa.es

Received and published: 10 June 2009

General comments

The paper presents observations of CO₂ (and CO) in the mesosphere and lower thermosphere (MLT) as measured by ACE (using the solar occultation technique) and analysed with the CMAM model. This represents a new dataset of CO₂ measurements in the upper atmosphere which is very important given the scarcity of global CO₂ measurements in this atmospheric region. The paper also includes a sensitivity study of CO₂ abundances to several model parameters which is very illustrative for understanding the CO₂ distribution. The conclusions reached about the processes responsible for

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the CO₂ distribution are, however, rather speculative and probably the dynamics aspects should be further checked before those speculations are put forward.

The CO₂ dataset presented represents a significance advance to our knowledge of the MLT region and therefore worth to be published. However, there are some major points which need to be addressed before the paper can be accepted for publication.

1. The paper does not discuss completely all important work previously done on this subject. Some important references are missing and the discussion of previous work is not fully accurate at some passages.

2. The paper lacks in a thorough and deep description and discussion of ALL errors affecting the retrieved CO₂ field. Sentences as " .. and such error has not been taken account" should not be acceptable and the 2K perturbation temperature error does not seem to be realistic. This is very important since when reaching conclusions about the processes controlling CO₂ in the MLT region from data/model comparisons, the errors in the former have to be taken into account and, depending on their magnitude, the conclusions might differ.

3. The retrieval inversion of pressure/temperature and CO₂ performed as a joint retrieval in a first step but carried out in a second step retrieving only CO₂, poses some problems from the information content point of view. Since the same spectral windows are used for retrieving temperature and CO₂, both parameters should be retrieved simultaneously (using an adequate constraint) and I cannot see the point of why a second step retrieval, retrieving only CO₂, is necessary.

4. The CO₂ distributions shown in Fig. 1 seems to have significant artifacts due to the sampling and averaging (over one month). This fact is mentioned in p. 11562, l. 26 and ff. This point should be included when describing the distributions in this figure, and discuss in more detail how it affects the shown distributions. The fact that such plots have several discontinuities in latitude probably makes this point even more important. For example, is the sharp altitude shift in February at about 15°N caused by

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

this artifact?

5. I am not fully convinced that the uncertainties in the dynamical processes are not large enough for explaining the data/model discrepancy and that new chemical loss processes for CO₂ are needed to be invoked.

6. About the presentation of the paper. I think that it could be significantly shortened. There are many figures which do not add new information and whose discussion make the paper lengthy and repetitive. I do not see any point in presenting the sensitivity study for all cases as zonal mean as well as profiles for several latitudes, and then for different months (April and August), e.g., figures 4, 7, 9 and 10). Also, much of what is said in the "Discussion" section is repeated from the previous section. I suggest to merge the "Results" and "discussion" sections or omit several paragraphs in the "Discussion" section already described before.

7. Including so many panels in each figure make them, at least in the printed version, really unreadable. This technical problem has to be solved, otherwise the features discussed in the text are hardly discernible in the figures.

Specific Comments

In the specific comment below I normally quote the paragraph in question from the manuscript and then comment on it.

INTRODUCTION

P. 11553, l. 16-18. "(Kaufmann et al., 2002). They found that the CO₂ volume mixing ratio (VMR) deviated from a well mixed state, which we will call the "knee", around 70 km."

The fact that CO₂ deviates from the well mixed value much lower, at about 80 km, that models have predicted before, was shown much earlier than Kaufmann et al. by the SAMS, ATMOS and ISAMS measurements (See Lopez-Puertas and Taylor, 1989; Rinsland et al., 1992; Lopez-Puertas et al., 1998b; and Zaragoza et al., 2000).

C1646

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



See also the discussion on this topic in Lopez-Puertas et al., 2000, A review of CO₂ and CO abundances in the middle atmosphere, in *Atmospheric Science Across the Stratopause*, Amer. Geophys. Union, Geophys. Monograph 123, 83-100, 2000. This paper reviews all CO₂ and CO measurements at that time and might be useful for the reader to mention it in the introduction.

P. 11554, l. 10-13. "There is also some evidence that highly vibrationally excited hydroxyl molecules affect the CO₂ asymmetric stretch mode (Kumer et al., 1978)."

This has been evidenced more clearly from SABER measurements very recently, showing that the efficiency in transferring vibrational energy from OH to $\tilde{\nu}_2$ N₂ is about a factor of 3 larger than previously thought, see, e.g., López-Puertas et al., JGR, 109, 09307, 2004, 10.1029/2003JD004383. The authors might want to include this more updated reference.

P. 11554, l. 12-14. "As stated by Kaufmann et al. (2002) the O(1D) excitation mechanism and the non-LTE model parameters constitute the most important uncertainties of retrieved CO."

Edwards et al., JGR, 101, 26577, 1996, 10.1029/96JD02133 were able, from CLAES CO₂ 10 μm measurements, to give rather small constraints on the excitation of CO₂(001) from O(1D).

P. 11554, l. 18-20. "However, for solar occultation measurements the absorption only depends on the CO₂ density, the kinetic temperature and the pressure and not on the vibrational excitation of the CO₂ molecules".

This assertion is generally true only if fundamental transitions (e.g. those with the ground vibrational state as the lower level of the transition) are involved. For hot transitions, NLTE can be significant and actually this has been used for retrieving non-LTE populations, see, e.g., Rinsland et al., JGR, 1992. Being even more rigorous, non-LTE can be important in occultation measurements even for fundamental transitions,

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment

through the vibrational partition function, see, Edwards et al., J. Quant. Spectros. Radiat. Transfer, 59, 423, 1998. This situation does actually take place for ALL CO₂ transitions at altitudes above around 100 km in our atmosphere. An extended discussion of this topic is given in Lopez-Puertas and Taylor, Non-LTE Radiative Transfer in the Atmosphere. World Scientific, Singapore/New Jersey/London/Hong Kong, Sec. 8.10, 2001.

P. 11554, l. 25-27. "... and in the Atmospheric Laboratory for Applications and Science (ATLAS) 1, 2 and 3 missions (Kaye and Miller, 1996)."

I have not seen any CO₂ measurements retrieved from the ATLAS missions spectra in that reference. Has CO₂ been actually retrieved from those missions? To my knowledge no retrieved CO₂ have been published from these missions. I suggest to remove that sentence and the reference.

P. 11555, l. 20. "The vertical resolution is $\approx 3-4$ km" In line 20 (P. 11557) is mentioned that the vertical sampling can be as much as 6 km. Maybe some more details should be given here. Are there particular months for which the vertical resolution is best/worst?

P. 11555-6, l. 22-27 and 1-7. the p/T and CO₂ retrieval.

It is not clear for me how the retrieval was performed. I have understood that p/T and CO₂ were retrieved jointly in a first step but with a strong regularization on CO₂. Then, in a second step, the retrieved p/T from the first step was used to retrieve CO₂. In the two steps the same micro-windows were used for p/T and CO₂. If this is what has been done, I think the method is not appropriate. You cannot use the result of the first retrieval in a MW as a priori for the second retrieval from the same MW of the same measurement, because then, the a priori and the measurement are no longer statistically independent, and the usual retrieval formalism does no longer hold. Maybe I did not understand correctly. I think this should be described in more detail and/or clarified.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment

P. 11556, l. 5-15. It would be very useful to list the CO₂ transitions used in the retrievals. Does the temperature information comes from the rotational distribution of the lines? Are there enough lines to cover the range from optically thin to moderate for the whole altitude range (50-120 km)? No need to include hot bands? It would be useful to roughly indicate which bands were used for different altitudes.

P. 11556, l. 25-29. "As reported by McLandress et al. (2006), the meridional wind in the CMAM is characterized by summer-to-winter \bar{w} in the mesosphere and winter-to-summer \bar{w} in the lower thermosphere, between 100 and 120 km. The former is a feature of the thermally indirect circulation driven primary by non-orographic gravity wave drag (GWD), whereas the meridional wind reversal in the lower thermosphere is a direct result of the resolved wave drag."

Is this meridional wind patten, with two opposite circulations, below and about 100 km, a generally accepted result of 2D and 3D models or is it present only in the CMAM model?

Could the authors explain in some more details for the non-dynamicist readers the origin of these two circulation branches: What does "thermally indirect" circulation mean? Could they explain the differences between "non-orographic gravity wave drag" and "resolved wave drag"?

P. 11556 (bottom and 11557) "The meridional CO₂ distribution for the solstice months appears to be consistent with the large-scale circulation exhibited by the extended Canadian Middle Atmosphere Model (CMAM)"

It is really difficult to get the picture of the circulation predicted by the model from the maps in Fig. 1. First, because they are very small. Second, because of the limited latitude range. Third, because of the presence of the sampling artifacts which which cannot be easily distinguished from actual features. For example, the authors mention, l. 5, "... the January CO₂ data for the austral subpolar region appear to indicate the up-welling cell up to about 85 km (5.10–3hPa) and downwelling cell in the region

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

above," The upwelling cell is clear, although it should be mentioned that it is about 50N (at higher latitudes, $\sim 70\text{N}$, it is not so clear and confusing, artifact?) but with respect to the descending cell, do the authors mean the only point at $3\text{e-}3$ mb? It is really hard to see it. The authors should use larger figures, or maybe a zoom of the figures, or draw circles around the features or refer to the latitudes more specifically to show the features more clearly.

It is really hard to see these features at June/July. How does the CO_2 values at the top altitudes compare with the errors of the measurements?

The evaluation of the error budget is incomplete and seems rather optimistic. The propagation of the temperature error in the CO_2 retrieval, estimated by just shifting the temperature profile by 2K, is underestimated. The abstract of the ACE validation paper by Sica et al, ACP, 2008 state that "There is evidence of a systematic high bias (roughly 3–6 K) in the ACE-FTS temperatures in the mesosphere ..."

Further, I cannot see any scientific justification for estimating the CO_2 error as the differences between the CO_2 retrieved with a strong regularization and the CO_2 derived including the temperature from the first step (If I did understand right). I suggest that only one p/T CO_2 joint retrieval should be done and the errors (noise) will come up from the retrieval. If it is used a retrieval grid different from the measurements grid, then some kind of regularization should be applied which will impact the noise error and the vertical resolution (averaging kernel).

P. 11557, l. 23-24 "Above the mesopause, temperature changes rapidly with altitude, and a temperature uncertainty of 2 K is possibly an underestimate."

According to the ACE temperature validation paper, the temperature is underestimated at the mesopause and below. The temperature mapping in CO_2 should be revised and quantified at all altitudes where the CO_2 is presented. Spectroscopic errors and other systematic instrumental and forward model errors should also be quantified.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

It is not clear which is the noise error of a single CO₂ profile. This should be given and then an estimate of the values shown in the zonal mean, when some averaging are performed. I suggest to include a table listing the errors from the different sources, random and systematic at several altitudes covering the CO₂ range.

As new processes occurring in the atmosphere are proposed, this requires a good estimate of the CO₂ error to be considered when comparing with models results.

P. 11559, l. 17-19 and ff. Comparison of CO₂ is done for a particular set of CO₂ profiles, e.g. CRISTA, SABER and rockets. Is there any rationale of why selecting those measurements? There are other measurements, e.g., SAMS, ISAMS and ATMOS, see e.g. Lopez-Puertas et al., 2000 cited above. Why have not them been included? About the rocket profile, this differs from that compiled by Wintersteiner et al. (see Wintersteiner et al. 1992, and Lopez-Puertas et al, 2000). It might be useful for the reader to know about distinct those rocket profiles are. Also, the CO₂ retrieval from SABER, version 1.06, is very preliminary and was not validated. This should also be stated in the manuscript to warn the reader.

P. 11560, l. 1-10. Discussion about the altitude of the knee.

It might be useful to have in mind the errors in CO₂ when comparing among different measurements. E.g. SABER being about 20 ppmv larger than ACE above the knee. This is 6%, which is within ACE errors (and probably SABER errors too). The rocket profile compiled by Wintersteiner et al. (1992) has the "knee" around 90 km, higher than the rocket profile shown here.

P. 11560, l. 11-13. "ACE measures the ground state of CO₂ and, hence, provides more reliable information on the CO₂ abundance."

I'm not in favor of this kind of sentences. It is true that ACE CO₂ is less affected by non-LTE processes but the "goodness" of the CO₂ profile should be judged on the overall error budget. I suggest that the authors state the errors and, if they want to compare,

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

mention that is smaller/larger(?) than in other measurements.

P. 11560, l. 15. "Since CMAM is a climate model we cannot compare with the same dates .." You can, but it does not make sense.

P. 11560, l. 18-20. "The ACE-FTS gives a reasonable latitudinal coverage in April and from Fig. 4 we see that the overall structure exhibited by the model for the standard scenario, A, is similar to that of the observations. However, the measurements appear to have more structure with latitude."

It is mentioned (see above) that because of averaging over one month, the data includes some artifacts, shown as latitudinal structure. How much of the model/data discrepancy is due to these artifacts? This should be commented. Also, the best way to see if the model reproduce the measurements is to make a difference plot. I suggest that, instead of showing so many model zonal means for different conditions in Fig. 4, they authors include only the measurement plot, the model plot (whatever run with probably best fit), and a difference plot. The behavior of the different model runs are better shown along with profiles (as in Figs. 5). Having less panels might also help in showing larger figures.

Also, probably it would be best for showing any agreement/disagreement between data/model to compare for a solstice month (not April which is near equinox and exhibit less latitudinal gradients).

Stating that model and data give "SIMILAR" results is somehow in contradiction with the line 23 below where it is discussed about the disagreement between both. This is confusing and needs some rewriting.

P. 11560, l. 21 "fall-off of CO₂ mixing ratio with height for the ACE measurements is clearly seen to occur at lower altitudes than for the model results in the control run, scenario A."

This difference is seen more clearly in profile plots (not in zonal means). Certainly this

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

is NOT seen CLEARLY in this figure.

P. 11561, l. 15 "... transporting CO₂ up the vertical gradient." ... transporting CO₂ upwards.?

Figure 5. I suggest to remove the panel for 3°N. It does not add any new information to that for 30°N. This would help in enlarging the figure and reducing redundant text, e.g., (p. 11562, l. 4-6). Also, I suggest to include in the measured profile the CO₂ error bars (and remove Fig. 2).

P. 11562, l. 8. The role of K_{zz} on the CO₂ and CO distributions was already studied by Lopez-Puertas et al. (2000). A reference here might be useful for the reader.

P. 11562, l. 9 and ff. I would write these sentences in a easier way. Replace: "Viewed as an experiment on the role of GWD this suggests ... mid-latitude regions" by: "The reason of CO₂ being diffused upwards to higher altitudes in the polar regions is that the impact of turbulence generated by unresolved gravity wave breaking in the model mixing is most important in this region than in the tropics and mid-latitude regions (see Fig. 6)."

P. 11562, l. 19. "... that the CMAM simulation of CO provides a reasonable representation of CO."

- At 30-40 N there is a significantly larger descent in the data than in the model - The increase at 90°N is not in the model. Are these features within 30% errors?

P. 11562, l. 27-28. "... descent over the pole. However, this is an artifact due to the sampling limitations of the ACE experiment during a time of strong descent."

Are the features above also caused by these artifacts? This should be discussed earlier in the description of the data and the description of figures 1, 4 and 7.

P. 11563, l. 5. Insert a "," after "turned off"

P. 11563, l. 5-8 The CO obtained for case C for 30°N and 80°S (Figs. 8) seems in

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



rather good agreement with measurements; except for 80°S below 65 km, but in this case there is no other better model simulation.

P. 11563, l. 18. I suggest to remove also the panel for 3°N in Fig. 8 and the corresponding discussion (which is very similar than for panel 8a) in the text.

Could not be both CO₂ and CO measurements reasonably explained using a low K_{zz}? It is true that CO₂ is overestimated but it might be within the measured CO₂ error bars. If so, there it will be no need to invoke any new mechanism for sequestering CO₂.

Fig. 7. As for Fig. 5, I think there is no need to show zonal mean for all simulations. The differences of the sensitivity study are seen more clearly in profiles figures, as Fig. 8. The same applies to Figs. 9 and 10, which are similar to figures 5 and 7 except for August, instead of April.

P. 11564, l. 18-20. "We note that the slopes of the ACE contours in the polar region are affected by sampling as for April as can also be seen in Jin et al. (2008)."

This should be discussed much earlier in the paper (see comments above).

P. 11565, l. 11-24. Lopez-Puertas et al. (2000) were able to reproduce such a low "knee" in the CO₂ profiles by using reasonable K_{zz}. The problem with overestimation of CO₂ above about 80 km in the previous version of the 2D Garcia and Solomon model was that molecular diffusion was not properly taken into account. Probably this was the same reason for TIME-GCM not being able to reproduce such CO₂ profiles. In any case, the low CO₂ profiles of ATMOS, ISAMS, and SABER (similar to that of ACE) can be well reproduced, with no overestimation above around 75-80 km by the Whole Atmosphere Community Climate model (WACCM) (R. Garcia and D. Marsh, personal communication).

Part of the discussion in this section has already been described in the previous section, when describing the sensitivity study. I suggest to shorten this section in that sense.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

P. 11567, l. 14-17 "However, there is some justification for higher temperatures being used. This is because the vibrational levels of CO₂ are non-thermally excited in the mesosphere so that daytime vibrational temperatures are higher than the kinetic temperature (e.g., Lopez-Puertas and Taylor, 2001)".

The CO₂ 15 μm levels, mainly the CO₂(010), is very close to LTE up to about 100 km. Furthermore, the number of CO₂ molecules in this level (and in higher levels) represents only about 1% of the total number of CO₂ molecules, most of which are in the ground state. I suggest to remove that paragraph. It does make sense to consider the temperature as low at 200 K in that region.

P. 11568. l.9. "In the control model simulation, scenario A, CMAM CO is up to a factor of two too low above 0.01 hPa (80 km) but it is not clear how serious a disagreement this is."

I do not understand this sentence. Is it not clear that a factor of 2 is a large disagreement? (Unless the measurements have a similar error).

P. 11568. As before, there is quite a lot of repetition from previous section.

P. 11568, l.25 "We note that the agreement between model and measurements is improved somewhat but that the model CO₂ still remains too high. "

How high in comparison with ACE CO₂ error?

P. 11569, l.7-9 "As is clear from above results the most reasonable scenario for agreement between the ACE observations and CMAM simulations is with an increased J value ..."

This is questionable. One could argue that this gives a very large overestimation of CO (see Figs. 8 a and c).

P. 11569, l. 19-22 "An interesting feature of such a phenomenon is that it will be sporadic, and its effect and its effects will vary from season to season with varying dust

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

amounts which might account for the variation required in the “enhanced” J (CO₂) to account for the observations in April and August.”

If sporadic, how can it account for a global and constant CO₂ depletion? Is there any hint that the dust amounts changed from April to August and in the correct sense? Wouldn't we expect a larger meteoritic ablation in August? Ins this is the right sense?

P. 11569, l. 23-27 and Fig. 11 I can't see the need for including the figures for temperatures. Possible temperature differences are not invoked to explain the CO₂/CO measurements/model differences or any comparison is made among modeled and measured temperatures. I suggest to remove this paragraph and Fig. 11.

The error bars of Fig. 2 are already shown in Fig. 3. I suggest to remove Fig. 2.

Table 1, scenario B. Insert a "," after "factor of 5"

Interactive comment on Atmos. Chem. Phys. Discuss., 9, 11551, 2009.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

