

***Interactive comment on “Measurements of volatile organic compounds in the middle of Central East China during Mount Tai Experiment 2006 (MTX2006): observation of regional background and impact of biomass burning” by J. Suthawaree et al.***

**J. Suthawaree et al.**

deutschf@tmu.ac.jp

Received and published: 13 November 2009

General comments: Based on VOCs measurements on top of Mount Tai, the manuscript conducted interesting evaluation about the sources of VOCs in the middle of Central East China region. The methodologies are well accepted. The work could be helpful due to the lack of VOCs data in this area.

Response The authors appreciate the reviewer comments. We will try our best to  
C7084

improve the manuscript according to the reviewer suggestions and comments.

Specific comments: 1. To enhance the introduction and discussion of the manuscript, the Authors are suggested to use more recent researches in China on VOCs, e.g. Source profiles ( Liu et al., AE, 2008, 42: 6247-6260), source apportionments for both urban and rural site in PRD ( Liu et al., AE, 2008, 42: 6261-6274), VOCs tracers for biomass burning (Xie et al., AE, 2008, 42: 6000-6010), and the VOCs in Beijing city (Shao et al, JGR, 2009, DOI:10.1029/2008/2008JD010863).

Response 1. Thank you very much for your suggestions. Suggested references are added to the manuscript in introduction and discussion sections.

2. The discussion in line 18-21, page 16719, the authors need to be more explicit about chemical regimes of ozone formation, I guess the model results indicate that VOCs-limited was true for urban areas, was it the same for remote site (as Mount Tai ) and the whole region?

Response 2. The model results provided by Kanaya et al. (2009) reveal NO<sub>x</sub> limited regime for Mount Tai. The sentence is revised to be more concise and explicit.

3. It could be better if the analysis for the sources of VOCs have support of local (or regional information), the ERs comparison with national inventory was not that convincing. As a background site, the discussion reveals the impact of biomass burning, I wonder if there were any local fires during the measurements.

Response 3. The authors also concerned the issue pointed out by reviewer and try the best to find local and regional scale information for comparison. Nonetheless, to the best of authors' knowledge, there is still very limit information on ERs in the region for C<sub>2</sub>Cl<sub>4</sub>, CH<sub>3</sub>Cl and CH<sub>3</sub>Br. For ethane and propane and benzene, comparisons of ERs were made with the regional scale information provided by Carmichael et al. (2003). The impact of local biomass burning is believed to be at the minimum owing to the local of the site on the mountain top. Nonetheless, long-range or mid-range transport

of emission from biomass burning was found to impact VOC mixing ratio observed at this site.

#### References

Carmichael, G. R., Tang, Y., Kurata, G., Uno, I., Streets, D., Woo, J.-H., Huang, H., Yienger, J., Lefer, B., Shetter, R., Blake, D., Atlas, E., Fied, A., Apel, E., Eisele, F., Cantrell, C., Avery, M., Barrick, J., Sachse, G., Brune, W., Sandholm, S., Kondo, Y., Singh, H., Talbot, R., Bandy, A., Thornton, D., Clarke, A., and Heikes, B.: Regional-scale chemical transport modeling in support of the analysis of observations obtained during the TRACE-P experiment, *J. Geophys. Res.*, 108 (D21), 8823, doi:10.1029/2002JD003117, 2003.

Kanaya, Y., Pochanart, P., Liu, Y., Li, J., Tanimoto, H., Kato, S., Suthawaree, J., Inomata, S., Taketani, F., Okuzawa, K., Kawamura, K., Akimoto, H., and Wang, Z.: Rates and regimes of photochemical ozone production over Central East China in June 2006: a box model analysis using comprehensive measurements of ozone precursors, *Atmos. Chem. Phys.*, 9, 7711-7723, 2009.

Liu, Y., Shao, M., Fu, L., Lu, S., Zeng, L., and Tang, D.: Source profiles of volatile organic compounds (VOCs) measured in China: Part I, *Atmos. Environ.*, 42(25), 6247-6260, 2008.

Liu, Y., Shao, M., Lu, S., Chang, C. C., Wang, J. L., and Fu, L.: Source apportionment of ambient volatile organic compounds in the Pearl River Delta, China: Part II, *Atmos. Environ.*, 42(25), 6261-6274, 2008.

Shao, M., Lu, S., Liu, Y., Xie, X., Chang, C., Huang, S., and Chen, Z.: Volatile organic compounds measured in summer in Beijing and their role in ground-level ozone formation, *J. Geophys. Res.*, 114, D00G06, doi:10.1029/2008JD010863, 2009.

Xie, X., Shao, M., Liu, Y., Lu, S., Chang, C., and Chen, Z.: Estimate of initial isoprene contribution to ozone formation potential in Beijing, China, *Atmos. Environ.*, 42, 6000-

C7086

6010, 2008.

---

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 9, 16715, 2009.

C7087