

Interactive  
Comment

## ***Interactive comment on “Scanning electron microscopy and molecular dynamics of surfaces of growing and ablating hexagonal ice crystals” by W. C. Pfalzgraff et al.***

**W. C. Pfalzgraff et al.**

nesh@pugetsound.edu

Received and published: 19 November 2009

We are grateful to the reviewer for suggesting that we take a closer look at geometrical aspects of this work. We now think there are two pyramidal prismatic facets, the predominant being the well-known 28-degree (101 $\bar{1}$ ) facet. The 14-degree has Miller-Bravais indices (202 $\bar{1}$ ), we believe. More details are given in our online comment to reviewer #1.

Regarding the stability (or transience) of the pyramidal prismatic facets, we rarely see a crystal that does not exhibit some development of this facet. We are currently trying to gather quantitative information about their incidence and extent, and whether those

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



properties depend on growth conditions (temperature, and ambient pressure to the extent that the VP-SEM allows).

As for the preference for prismatic attachment over basal attachment (to the copper substrate), our working hypothesis is that the crystal spacing of the copper substrate offers a better match to the prismatic facet.

Regarding the time dependence of relevant thermodynamic parameters, we have looked at the potential energy over time (Fig. 1 of this response). A feature around 5 ns – a reduction in potential energy – suggests (perhaps) a surface freezing process. We have not yet designed an approach that would allow us to pin down such structural changes, however. Perhaps a count of hydrogen bonds, or tracking the dipole correlation function? In our minds, a larger question is, what simulation scale is needed to reproduce the growth features seen in the SEM, and what approach might let us get to that scale?

---

Interactive comment on Atmos. Chem. Phys. Discuss., 9, 20739, 2009.


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



[Interactive  
Comment](#)

**Fig. 1.** Potential energy of the free-standing nanocolumn.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)