

Interactive comment on “The impact of MM5 and WRF meteorology over complex terrain on CHIMERE model calculations” by A. de Meij et al.

Anonymous Referee #1

Received and published: 6 February 2009

General Comments

It is an interesting paper providing some answers on the effect of meteorology on air pollution. It includes results (model output and measurements) not shown before and some explanations are given on the agreement or disagreement between model output and observations. On the other hand, all the conclusions concerning model performance (both in terms of meteorology and air quality) are rather uncertain because the measurements were limited; there was only a small number of measuring stations and they only provided surface measurements. Therefore, one cannot get information on the variation of the model performance throughout the model domain and there can be no comparison of the vertical structure of the atmosphere (which plays a key role in air pollution applications) with observations. Some more specific comments on this paper

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



follow in the next section.

Specific comments

In the Abstract and in the Concluding Remarks sections (lines 6 and 22 respectively) it is mentioned that "The analysis shows that the performance of both models is similar, however some small differences are still noticeable". It should be clarified that this refers to surface meteorological data only. As mentioned afterwards, "The PBL height by WRF meteorology is a factor 2.8 higher at noon in January than calculated by MM5"; this is a substantial (and not just noticeable) difference in the simulated vertical structure of the atmosphere.

Another finding reported in the Abstract, line 20, is that "changing the Noah Land Surface Model for the 5-layer soil temperature model, the calculated monthly mean PM10 concentrations increase by 30%"; It is not specified (in the Abstract) which model this statement refers to (MM5 or WRF).

In the first paragraph of the introduction (line 8) it is mentioned that "chemistry-transport-dispersion models (CTMs) have the advantage that they can be used to complement monitoring data, assess the effects of future changes in aerosol and aerosol precursor emissions"; CTMs also treat gaseous pollutants (and so does the paper). Therefore this statement should refer to gaseous pollutants too.

Both the third and fourth paragraphs of the Introduction talk about the fact that pollutant concentrations and their uncertainties depend on meteorology, chemistry and emissions. In this sense, they overlap and should be merged into one concise paragraph.

In page 2322 line 5 there should be a reference supporting that Po Valley is one of the most polluted, industrialized and heavily populated areas in Europe.

In page 2323, line 10 it is mentioned that PM10 and O3 "are prevailing";

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



the most in Po Valley. The term “prevailing”; here is unclear. Does it mean that they have larger concentrations/more adverse health effects than other pollutants for example?

Four monthly simulations are mentioned in the last paragraph of the methodology section (page 2323). It should be specified if there is nudging of the simulations towards observations or not. Also, there are comparisons with yearly data. Where there yearly model runs as well?

Citation should be added to the definition of the planetary boundary layer in line 15, page 2328. That is Stull, 1988.

In line 14, the term “absolute temperature”; can easily be confused with the absolute temperature scale in Kelvin (K) where zero is the absolute zero. Since the discussion of temperature is in degrees C, the term “temperature”; instead of “absolute temperature”; is more suitable here. Moreover, in the paragraph under this title, in all subsequent discussions of temperature and in Table 3a, the statistic Relative Bias (RB) is used for model evaluation purposes. This statistic takes unreasonably large values when the average observed temperature is close to 0°C and becomes infinity if the average observed temperature is equal to 0°C, even if the model temperature estimates are very good and differ from observations only by a few tenths of a degree. Therefore such values do not necessarily represent a bad model performance. They are the result of the fact that in the Celsius temperature scale, zero is defined as the temperature where water freezes. It is suggested that RB is either not used for temperature or that its calculation is based on the absolute temperature scale, after the conversion of temperature in degrees Kelvin.

In the first paragraph of page 2335 where wind direction is discussed, in all subsequent discussions on wind direction and in Table 3d a number of statistics such the mean, bias, coefficient of determination etc. have been calculated. One should note that circular data such as wind direction should be treated in a different manner. Common

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

evaluation statistics such as the ones calculated here do not apply. To bring a simple example on this, the arithmetic mean of 0 and 360 is 180 although 0 is apparently a better choice as a mean value. Moreover, the terms "underestimation" and "overestimation"; (e.g. p. 2335, line 5) are meaningless here. An "underestimation" of a measurement of 10 wind direction by 6 yields a value of 4. An "underestimation" of a value of 10 by 12 yields 358 which can then be seen as a (large) overestimation. The whole statistical analysis of wind direction should be redone accounting for the particularities of circular data. Further explanations on this can be found in a number of publications such as the "Statistical analysis of circular data" by N.I. Fisher which, in general, is a good starting point for the study of this field.

Concerning rain, certain statistics have been calculated in order to evaluate the prediction of the correct amount of rain by MM5 and WRF. However, it is probably even more important for the prediction of rain to know whether a model captured the event of precipitation during a certain day or not. For this purpose hit ratio statistics should be calculated and reported for rain.

It is mentioned in page 2340, line 12 that relative humidity is in general overestimated by MM5. By looking at the yearly bias values in Table 3b one cannot conclude this. Yearly averages of relative humidity are slightly overestimated by WRF indeed.

In line 18 of the same page, "For the winter period WRF gives higher temperatures"; higher than what, MM5 or observations?

Since section 4.2.1 refers to winter 2005, the statement in line 7 page 2343 that the underestimation of PM10 is because of overestimation of the relative humidity by the two meteorological models, is not true. Examination of Table 3b shows that in winter 2005 relative humidity was underestimated. Besides, this is also mentioned in the second paragraph of section 4.1.4 (line 17 page 2340).

In line 15, page 2344 a comparison between the "cloud cover"; and the

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

“cloud attenuation” between MM5 and WRF is done. In order to get an idea of the sensitivity of PBL height to cloud cover or cloud attenuation, it would be useful to quantify the differences in those quantities between the two models. Moreover, MM5 does not predict cloud cover and cloud attenuation. Has there been some diagnostic procedure to estimate those quantities? Which one? An alternative solution is to make a quantitative comparison of cloud liquid water which is a prognostic variable of MM5.

In lines 16-21, page 2344, the differences in cloud attenuation between the two models is attributed to the differences in cloud liquid water. In line (22) of the same page, it is stated that more cloud liquid water in WRF is the result of more cloud attenuation. The cause-and-effect relationship between cloud liquid water and cloud attenuation is reversed in those statements. This is a contradiction which should be clarified.

In page 2353, lines 21-23 some findings concerning the model performance in predicting ammonium and nitrate are reported. Given the uncertainties in the measurements of ammonium and nitrate, one cannot draw reliable conclusions on the agreement of the model with measurements. Therefore this part should be excluded from the conclusions section.

In page 2355, line 28; low inversion heights are presented as a cause of stagnant conditions. If by “stagnant conditions” the author means low wind and weak vertical mixing, low inversion heights are rather a result, not a cause.

In page 2356, lines 6 and 7, it is mentioned among the findings of this study that gas and aerosol concentrations have a non-linear dependence on the meteorological conditions. In this paper, although some monotonic relationships between meteorological variables and concentrations of air pollutants were highlighted, there was no investigation on a possible (non-)linearity of those relationships. Therefore there is not enough evidence in the present paper to conclude a non-linear dependence. However, this statement on the non-linearity can remain if references are added (e.g. the references

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

in page 2321, line 23).

In page 2356, lines 8-12, apart from the improvements in meteorology and the emissions, updated chemical mechanisms is also a key requirement for better air quality models.

Results in Table 3a have two decimal digits. Temperature measurements typically have an accuracy of one decimal digit (which means three significant digits for temperatures T such that $|T| \geq 10^\circ\text{C}$ and two significant digits for temperatures T such that $|T| < 10^\circ\text{C}$). If this is the case with measurements from the ARPA networks, the calculated statistics should be rounded accordingly. The same process should be applied to tabulated model data because measurements are involved in the process of modeling by data assimilation in the examined model runs or in the models which produced the initial/boundary conditions. Similar considerations about the number of decimal or significant digits apply in all of the presented results.

In tables 4, 5, 6 and 7 only the mean values of model and measured pollutant concentrations are reported. It is suggested that more statistical measures are calculated in order to get a more complete picture of the CHIMERE performance and its dependence on the meteorological input. For example, the statistics used in the meteorological evolution could also be used in the air quality evaluation. In addition, the factor of two statistic could be calculated. This is defined as the fraction of model values which lie between 50% and 200% of the corresponding measurements and it is particularly useful for the evaluation of CTMs.

In the wind rose on the left-hand side of Figure 2, there are some wind directions (such as the eastern) where not a single observation was recorded throughout a year. This result is rather suspicious and the wind rose should be reexamined.

Technical corrections

In line 27, page 2322 the use of 'such as' implies that there were

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



more meteorological parameters which were evaluated in the study. This is not true so Δ ; such as Δ ; should be removed.

Many web references are included in the paper. Given that web pages change over time, the date of last access should be added to all of them. For example, instead of (<http://aqm.jrc.it/citydelta>), (<http://aqm.jrc.it/citydelta>, last accessed 06.06.2006).

Page 2344, line 3: Δ ; surface Δ ; should probably be replaced by Δ ; sensible Δ ;

Page 2344, line 11: Δ ; observed Δ ; should be replaced by Δ ; estimated Δ ;

Page 2344, line 18: Δ ; hydrometer Δ ; should be replaced by Δ ; hydrometeor Δ ;

Page 2344, line 22: Δ ; is Δ ; should probably be replaced by Δ ; in Δ ;

The text from line 24 page 2348 until line 3 page 2349 overlaps with the text in lines 16-22 page 2331. Instead of repeating the reader can be directed to the chapter where this content appeared first (3.1). The same applies to lines 1-7 in page 2350; their content has already been presented in section 4.2.4.

Interactive comment on Atmos. Chem. Phys. Discuss., 9, 2319, 2009.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)