

**The effect of future ambient air pollution on human mortality to 2100 using output
from the ACCMIP model ensemble – SUPPLEMENTAL MATERIAL**

Raquel A. Silva, J. Jason West, Jean-François Lamarque, Drew T. Shindell, William J. Collins, Stig Dalsoren, Greg Faluvegi, Gerd Folberth, Larry W. Horowitz, Tatsuya Nagashima, Vaishali Naik, Steven T. Rumbold, Kengo Sudo, Toshihiko Takemura, Daniel Bergmann, Philip Cameron-Smith, Irene Cionni, Ruth M. Doherty, Veronika Eyring, Beatrice Josse, I. A. MacKenzie, David Plummer, Mattia Righi, David S. Stevenson, Sarah Strode, Sophie Szopa and Guang Zeng

1 Air pollutant ambient concentrations

The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) included contributions from several modeling groups. While up to 14 models reported ozone concentrations (depending on the scenario), only up to 6 models reported species used in the calculation of PM_{2.5} concentrations, and up to 4 models reported their own estimate of PM_{2.5} concentrations (Table S1).

Figure S1 shows ten world regions used for all regional calculations presented below.

Global population-weighted differences (Future year – 2000) in ozone and PM_{2.5} concentrations for the different models are shown in Tables S2 and S3, respectively, while regional multi-model average differences are shown in Figures S2 and S3. For both pollutants, metrics are consistent with the underlying epidemiological studies for the health impact assessment:

- Seasonal (6-month) average of daily 1-hr maximum ozone concentration;
- Annual average PM_{2.5} concentration.

PM_{2.5} concentration is estimated using the sum of PM_{2.5} species mass mixing ratios reported by six models:

$$\text{Estimated PM}_{2.5} = \text{BC} + \text{OA} + \text{SO}_4 + \text{SOA} + \text{NO}_3 + \text{NH}_4 + 0.25 * \text{SS} + 0.1 * \text{Dust},$$

where BC – Black Carbon, Dust, OA – (Primary) Organic Aerosol corrected to include species other than carbon, SO₄ - Sulfate, SOA – Secondary Organic Aerosol, and SS – Sea Salt, following Fiore et al. (2012) and Silva et al. (2013). The factors 0.25 and 0.1 are intended to indicate the fractions of sea salt and dust that are in the PM_{2.5} size fraction.

2 Future Population and Baseline Mortality Rates

Figure S4 shows future total and exposed population in 2030, 2050 and 2100 estimated from International Futures (IFs) country-level population per age group, used in the health impact assessment, as well as United Nations (UN) and Representative Concentration Pathway scenarios (RCPs) totals as context.

Figure S5 shows baseline mortality rates for chronic Respiratory diseases (RESP, ICD-9¹ BTL: B347), ischemic heart disease (IHD, ICD-9: 410-414), cerebrovascular disease (STROKE, ICD-9: 430-435, 437.0-437.2, 437.5-437.8), chronic obstructive pulmonary disease (COPD, ICD-9: 490-492.8, 494, 496) and lung cancer (LC, ICD-9 BTL: B101) estimated from IFs country-level mortality rates of cardiovascular diseases, chronic respiratory diseases and malignant neoplasms. Here we show average values for the exposed population (adults age 25 and older), but we used age distributed values for IHD and STROKE in the premature mortality calculation to align with available relative risks of exposure for these diseases.

¹ ICD-9 - International Classification of Diseases, revision 9.

3 Detailed results

Table S4 shows the multi-model average global future ozone premature mortality, including uncertainty for RCP8.5, while Table S5 shows the multi-model average across ten world regions. Table S6 shows the multi-model average global PM_{2.5} mortality (IHD+STROKE+COPD+LC), including uncertainty for RCP8.5, while Table S7 shows the multi-model average across ten world regions. The multi-model average corresponds to the average of estimates given by the available models for each scenario/period.

Tables S8 and S9 show the global burden on mortality of ozone and PM_{2.5} concentrations in 2000 relative to 1850, using present-day population and baseline mortality rates, and in 2030, 2050 and 2100 for all RCPs relative to 1850, using future population and baseline mortality rates. Also shown are two alternative cases for global burden calculation, using: A) 2000 concentrations relative to 1850 and present-day population but future baseline mortality rates; B) 2000 concentrations relative to 1850 but future population and baseline mortality rates.

Figures S6 and S7 show future global and regional air pollution-related premature mortality per million people in 2030, 2050 and 2100, for all RCPs relative to 2000.

References

Cameron-Smith, P., Lamarque, J.-F., Connell, P., Chuang, C., and Vitt F.: Toward an Earth system model: atmospheric chemistry, coupling, and petascale computing, *J. Phys.: Conf. Ser.* 46, 343-350, doi:10.1088/1742-6596/46/1/048, 2006.

Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C.,

Sitch, S., Totterdell, I., Wiltshire, A. and Woodward, S.: Development and evaluation of an Earth-System model-HadGEM2, *Geosci Model Dev*, 4, 1051-75, 2011.

Donner, L. J., Wyman, B. L., Hemler, R. S., Horowitz, L. W., Ming, Y., Zhao, M., Golaz, J. C., Ginoux, P., Lin, S. J., Schwarzkopf, M. D., Austin, J., Alaka, G., Cooke, W. F., Delworth, T. L., Freidenreich, S. M., Gordon, C. T., Griffies, S. M., Held, I. M., Hurlin, W. J., Klein, S. A., Knutson, T. R., Langenhorst, A. R., Lee, H. C., Lin, Y. L., Magi, B. I., Malyshev, S. L., Milly, P. C. D., Naik, V., Nath, M. J., Pincus, R., Ploshay, J. J., Ramaswamy, V., Seman, C. J., Shevliakova, E., Sirutis, J. J., Stern, W. F., Stouffer, R. J., Wilson, R. J., Winton, M., Wittenberg, A. T. and Zeng, F. R.: The Dynamical Core, Physical Parameterizations, and Basic Simulation Characteristics of the Atmospheric Component AM3 of the GFDL Global Coupled Model CM3, *J Climate*, 24, 3484-519, 2011.

Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S., Eyring, V., Folberth, G. a., Ginoux, P., Horowitz, L. W., Josse, B., Lamarque, J.-F., MacKenzie, I. a., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S. T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T. and Zeng, G.: Global air quality and climate, *Chem Soc Rev*, 41(19), 6663, doi:10.1039/c2cs35095e, 2012.

Institute for Health Metrics and Evaluation (IHME): Global Burden of Disease Study 2010 (GBD 2010) Results by Cause 1990-2010 - Country Level, Seattle, United States, 2013.

Jöckel, P., Tost, H., Pozzer, A., Bruhl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., Van Aardenne, J. and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESy1: consistent simulation of ozone from the surface to the mesosphere, *Atmos Chem Phys*, 6, 5067-104, 2006.

Josse, B., Simon, P., and Peuch, V.-H.: Rn-222 global simulations with the multiscale CTM MOCAGE, *Tellus*, 56B, 339–356, 2004.

Koch D., Schmidt G. A. and Field C. V.: Sulfur, sea salt, and radionuclide aerosols in GISS ModelE, *J Geophys Res*, 111, D06206, 2006.

Lamarque J.-F., Kyle G. P., Meinshausen M., Riahi K., Smith S. J., van Vuuren D. P., Conley A. J. and Vitt F.: Global and regional evolution of short-lived radiatively-active gases and aerosols in the Representative Concentration Pathways, *Clim Change*, 109, 191-212, 2011.

Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): overview and description of models, simulations and climate diagnostics, *Geosci. Model Dev.*, 6, 179-206, doi:10.5194/gmd-6-179-2013, 2013.

- Naik, V., Horowitz, L. W., Fiore, A. M., Ginoux, P., Mao, J., Aghedo, A. M. and Levy, H.: Impact of preindustrial to present-day changes in short-lived pollutant emissions on atmospheric composition and climate forcing, *J Geophys Res Atmos*, 118, 8086–8110, doi:10.1002/jgrd.50608, 2013.
- Oman L. D., Ziemke J. R., Douglass A. R., Waugh D. W., Lang C., Rodriguez J. M. and Nielsen J. E.: The response of tropical tropospheric ozone to ENSO, *Geophys Res Lett*, 38, 2011.
- Righi, M., Eyring, V., Gottschaldt, K.-D., Klinger, C., Frank, F., Jöckel, P. and Cionni, I.: Quantitative evaluation of ozone and selected climate parameters in a set of EMAC simulations, *Geosci. Model Dev.*, 8, 733–768, doi:10.5194/gmd-8-733-2015, 2015.
- Scinocca J. F., McFarlane N. A., Lazare M., Li J. and Plummer D.: Technical Note: The CCCma third generation AGCM and its extension into the middle atmosphere, *Atmos Chem Phys*, 8, 7055-74, 2008.
- Shindell D. T., Pechony O., Voulgarakis A., Faluvegi G., Nazarenko L., Lamarque J-F, Bowman K., Milly G., Kovari B., Ruedy R., Schmidt, G. A.: Interactive ozone and methane chemistry in GISS-E2 historical and future climate simulations, *Atmos Chem Phys*, 13, 2653–89, 2013.
- Silva, R. A., West, J. J., Zhang, Y., Anenberg, S. C., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S., Skeie, R., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V., Josse, B., MacKenzie, I. A., Plummer, D., Righi, M., Stevenson, D. S., Strode, S., Szopa, S. and Zeng, G.: Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change, *Environ Res Lett*, 8, 034005, doi:10.1088/1748-9326/8/3/034005, 2013.
- Skeie R. B., Berntsen T. K., Myhre G., Tanaka K., Kvalevag M. M. and Hoyle C. R.: Anthropogenic radiative forcing time series from pre-industrial times until 2010, *Atmos Chem Phys*, 11, 11827-57, 2011.
- Stevenson D. S., Doherty R. M., Sanderson M. G., Collins W. J., Johnson C. E. and Derwent R. G.: Radiative forcing from aircraft NOx emissions: Mechanisms and seasonal dependence, *J Geophys Res*, 109, 2004.
- Szopa S., Balkanski Y., Schulz M., Bekki S., Cugnet D., Fortems-Cheiney A., Turquety S., Cozic A., Deandreis C., Hauglustaine D., Idelkadi A., Lathiere J., Lefevre F., Marchand M., Vuolo R., Yan N. and Dufresne J.-L.: Aerosol and ozone changes as forcing for climate evolution between 1850 and 2100, *Climate Dynamics*, doi: 10.1007/s00382-012-1408-y, 2012.
- Teyssedre H., Michou M., Clark H. L., Josse B., Karcher F., Olivie D., Peuch V. H., Saint-Martin D., Cariolle D., Attie J. L., Nedelec P., Ricaud P., Thouret V., Van Der A. R. J., Volz-Thomas A. and Cheroux F.: A new tropospheric and stratospheric Chemistry and Transport Model MOCAGE-Climat for multi-year studies: evaluation of the present-day climatology and sensitivity to surface processes, *Atmos Chem Phys*, 7, 5815-60, 2007.

Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Rose, S. K.: The representative concentration pathways: An overview, *Clim Change*, 109, 5–31, doi:10.1007/s10584-011-0148-z, 2011.

Watanabe S., Hajima T., Sudo K., Nagashima T., Takemura T., Okajima H., Nozawa T., Kawase H., Abe M., Yokohata T., Ise T., Sato H., Kato E., Takata K., Emori S. and Kawamiya M.: MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments, *Geosci Model Dev*, 4, 845-72, 2011.

Zeng G., Pyle J. A. and Young P. J.: Impact of climate change on tropospheric ozone and its global budgets, *Atmos Chem Phys*, 8, 369-87, 2008.

Zeng G., Morgenstern O., Braesicke P. and Pyle J. A.: Impact of stratospheric ozone recovery on tropospheric ozone and its budget, *Geophys Res Lett*, 37, L09805, 2010.

Tables and Figures

Table S1 – Models that reported ozone, PM_{2.5} species and PM_{2.5} (mmrpm2p5) concentrations for ACCMIP, with type of ozone output (h – hourly, m – monthly) and number of reported PM_{2.5} species.

Model	Institution	Contact	Ozone	PM _{2.5}	References
CESM-CAM-superfast	LLNL	Dan Bergmann, Philip Cameron-Smith	h	-	Lamarque et al., 2013; Cameron-Smith, et al., 2006
CMAM	CCCMA, Environment Canada	David Plummer	h	-	Scinocca et al., 2008
EMAC	DLR, Germany	Veronika Eyring, Irene Cionni, Mattia Righi	m	-	Jöckel et al., 2006 Righi et al., 2015
GEOSCCM	NASA GSFC	Sarah Strode	h	-	Oman et al., 2011
GFDL-AM3	NOAA GFDL	Vaishali Naik, Larry Horowitz	h	8, mmrpm2p5	Donner et al., 2011; Naik et al., 2013
GISS-E2-R	NASA-GISS	Drew T. Shindell Greg Faluvegi	h	8, mmrpm2p5	Koch et al., 2006; Shindell et al., 2013
HadGEM2	Hadley Centre Met Office, UK	William Collins, Gerd Folbert, Steven Rumbold	m	6	Collins et al., 2011
LMDzORINCA	IPSL-LSCE, France	Sophie Szopa	m	-	Szopa et al., 2012
MIROC-CHEM	NIES-JAMSTEC-NagoyaU- KyushuU, Japan	Tatsuya Nagashima, Kengo Sudo, Toshihiko Takemura	h	6, mmrpm2p5	Watanabe et al., 2011
MOCAGE	MeteoFrance, France	Beatrice Josse	h	-	Josse et al., 2004; Teyssedre et al., 2007
NCAR-CAM3.5	NCAR	Jean-François Lamarque	h	6, mmrpm2p5	Lamarque et al., 2011, 2012
OsloCTM2	CICERO and Univ. Oslo, Norway	Stig Dalsoren, Ragnhild Skeie	m	8	Skeie et al., 2011
STOC-HadAM3	University of Edinburgh, UK	Ian MacKenzie, Ruth Doherty, David Stevenson	m	3 (not used)	Stevenson et al., 2004
UM-CAM	NIWA, New Zealand	Guang Zeng	h	-	Zeng et al., 2008, 2010

Table S2 – Global population-weighted differences (Future year – Hist. 2000) in ozone concentrations (ppb) for the 14 models in 2030, 2050, 2100 for the four RCPs. Pollutant concentrations are weighted by exposed population (adults aged 25 and older) in each future year. Models with the symbol * reported only monthly average ozone concentrations.

Models	2030				2050				2100			
	RCP2.6	RCP4.5	RCP6.0	RCP8.5	RCP2.6	RCP4.5	RCP6.0	RCP8.5	RCP2.6	RCP4.5	RCP6.0	RCP8.5
CESM-CAM-superfast^(a)				3.7								9.7
CMAM	-1.2	2.1		4.4					-9.0	-6.1		9.6
GEOSCCM											-4.7	
GFDL-AM3	0.5	4.7	1.3	9.0	-1.3	1.6	2.3	5.7	-8.7	-9.6	-8.3	5.1
GISS-E2-R	0.2	4.2	1.5	12.0	0.1	0.2	2.9	5.6	-5.1	-11.7	-7.6	2.9
MIROC-CHEM	-0.5		0.9	6.9	-2.3		1.6	4.1	-8.1		-8.5	1.3
MOCAGE	2.7		1.5	15.2					-9.4		-11.9	1.5
NCAR-CAM3.5	-2.5	0.9	-2.1	3.7					-11.7	-11.2	-11.4	0.9
UM-CAM	-1.4	2.5		7.1					-8.9	-7.3		3.9
CICERO-OsloCTM2*	0.0	2.8		8.2					-9.3	-9.5		4.2
EMAC*		4.0		9.5						-8.9		5.9
HadGEM2*	-0.9	-0.1		0.5					-7.7	-3.7		13.6
LMDzORINCA*	-1.7	1.7	0.2	7.2	-8.7	-4.7	-2.8	0.7	-9.9	-9.0	-8.9	3.6
STOC-HadAM3*	0.7			11.0					-10.3			3.5

^(a) CESM-CAM-superfast reported concentrations for RCP2.6 and RCP 6.0, but the simulations for these scenarios used an inconsistent SST file and are not a matched set with the other simulations, so they were not considered here.

Table S3 – Global population-weighted differences (Future year – Hist. 2000) in PM_{2.5} concentrations (estimated as a sum of reported species) (µg/m³) for the 6 models in 2030, 2050, 2100 for the four RCPs. Pollutant concentrations are weighted by exposed population (adults aged 25 and older) in each future year.

Models	2030				2050				2100			
	RCP2.6	RCP4.5	RCP6.0	RCP8.5	RCP2.6	RCP4.5	RCP6.0	RCP8.5	RCP2.6	RCP4.5	RCP6.0	RCP8.5
GFDL-AM3	0.1	1.1	0.03	1.9	-3.4	-2.3	1.4	-1.0	-5.6	-6.3	-4.8	-3.3
GISS-E2-R	-2.0	-1.3	0.9	0.1	-4.4	-4.5	0.8	-3.1	-5.1	-5.9	-5.1	-4.0
NCAR-CAM3.5	-0.4	0.01	-0.03	1.3					-5.7	-6.4	-4.9	-3.9
MIROC-CHEM	0.2		0.5	1.0	-2.7		1.3	-1.5	-3.9		-3.3	-2.1
CICERO-OsloCTM2	2.6			3.8					-3.3			-0.4
HadGEM2	0.5	0.9		1.7					-3.5	-4.6		-3.9

Table S4 – Global respiratory premature ozone mortality in 2030, 2050 and 2100 for all RCPs (considering the change in future ozone concentrations relative to 2000 concentrations), showing the multi-model average (deaths/year) for RCP2.6, RCP4.5 and RCP6.0 deterministic estimates and the empirical mean with 95% CI in parenthesis for RCP8.5 probabilistic estimates (including uncertainty in the RRs and across models). These results correspond to Figure 1. All numbers are rounded to three significant digits.

2030	
RCP2.6	11,900
RCP4.5	100,000
RCP6	71,200
RCP8.5	264,000 (-39,300 , 648,000)
2050	
RCP2.6	-450,000
RCP4.5	-360,000
RCP6	441,000
RCP8.5	246,000 (-59,600 , 556,000)
2100	
RCP2.6	-1,020,000
RCP4.5	-917,000
RCP6	-718,000
RCP8.5	316,000 (-187,000 , 1,380,000)

Table S5 – Premature ozone-related respiratory mortality in ten world regions relative to 2000 concentrations: (a) 2030, (b) 2050, (c) 2100, showing the multi-model average (deaths/year) of the deterministic results. All numbers are rounded to three significant digits.

(a) 2030				
Region	RCP2.6	RCP4.5	RCP6.0	RCP8.5
North America	-17,000	-12,500	-10,900	-8,200
South America	-2,710	-500	-3,260	1,840
Europe	-8,870	-5,590	-7,190	-880
Former Soviet Union	-2,200	-1,030	-1,600	660
Africa	2,100	6,440	-3,520	9,020
India	52,900	82,000	-6,440	124,000
East Asia	-11,300	25,700	103,000	127,000
Southeast Asia	2,980	5,010	4,890	5,980
Australia	-280	-120	-100	20
Middle East	-3,630	930	-3,700	7,460

(b) 2050

Region	RCP2.6	RCP4.5	RCP6.0	RCP8.5
North America	-85,500	-70,200	-52,100	-41,100
South America	-8,180	-4,910	-8,920	8,530
Europe	-49,400	-40,000	-34,800	-16,600
Former Soviet Union	-9,760	-6,390	-5,710	-440
Africa	13,100	16,600	-5,520	30,500
India	154,000	290,000	32,200	256,000
East Asia	-439,000	-514,000	518,000	3,830
Southeast Asia	900	-21,300	19,600	-9,920
Australia	-1,260	-590	-490	250
Middle East	-24,800	-9,930	-21,100	18,300

(c) 2100

Region	RCP2.6	RCP4.5	RCP6.0	RCP8.5
North America	-104,000	-66,300	-111,000	-21,100
South America	-19,800	-20,200	-25,900	7,950
Europe	-44,600	-24,900	-41,600	2,390
Former Soviet Union	-12,500	-8,180	-11,100	1,290
Africa	51,100	-16,000	-49,400	128,000
India	-230,000	-267,000	-125,000	292,000
East Asia	-509,000	-383,000	-241,000	-99,700
Southeast Asia	-65,000	-71,400	-28,000	-21,000
Australia	-1,620	-990	-1,510	790
Middle East	-83,400	-58,100	-83,200	29,800

Table S6 – Global premature PM_{2.5} mortality (IHD+Stroke+COPD+LC) in 2030, 2050 and 2100 for all RCPs (considering the change in future PM_{2.5} concentrations relative to 2000 concentrations), showing multi-model average (deaths/year) for RCP2.6, RCP4.5 and RCP6.0 deterministic estimates and the empirical mean with 95% CI in parenthesis for RCP8.5 probabilistic estimates (including uncertainty in the RRs and across models). These results correspond to Figure 4. All numbers are rounded to three significant digits.

2030	
RCP2.6	-258,000
RCP4.5	-289,000
RCP6	-169,000
RCP8.5	17,200 (-386,000 , 661,000)
2050	
RCP2.6	-1,670,000
RCP4.5	-1,760,000
RCP6	16,700
RCP8.5	-1,210,000 (-1,730,000 , -835,000)
2100	
RCP2.6	-1,930,000
RCP4.5	-2,390,000
RCP6	-1,760,000
RCP8.5	-1,310,000 (-2,040,000 , -174,000)

Table S7 – Premature PM_{2.5} mortality (IHD+Stroke+COPD+LC) in ten world regions: (a) 2030, (b) 2050, (c) 2100, showing the multi-model average (deaths/year) of the deterministic results. All numbers are rounded to three significant digits.

(a) **2030**

Region	RCP2.6	RCP4.5	RCP6.0	RCP8.5
North America	-77,800	-83,500	-59,700	-77,100
South America	-570	-6,100	-6,960	-6,290
Europe	-153,000	-152,000	-137,000	-176,000
Former Soviet Union	-119,000	-82,000	-101,000	-116,000
Africa	35,100	31,800	-10,200	46,100
India	150,000	176,000	-2,690	245,000
East Asia	-90,100	-137,800	151,000	86,000
Southeast Asia	27,800	-30,700	36,200	-430
Australia	-560	-180	-440	-30
Middle East	-30,700	-4,430	-37,400	-7,230

(b) 2050

Region	RCP2.6	RCP4.5	RCP6.0	RCP8.5
North America	-106,000	-114,000	-104,000	-107,000
South America	-7,550	-9,550	-6,720	-7,940
Europe	-198,000	-187,000	-193,000	-200,000
Former Soviet Union	-144,000	-158,000	-154,000	-156,000
Africa	40,200	46,000	-21,100	66,100
India	-6,540	97,000	152,000	308,000
East Asia	-1,050,000	-1,200,000	356,000	-906,000
Southeast Asia	-113,000	-193,000	52,500	-182,000
Australia	-370	-390	-250	-240
Middle East	-81,200	-47,800	-64,300	-47,200

(c) 2100

Region	RCP2.6	RCP4.5	RCP6.0	RCP8.5
North America	-105,000	-128,000	-116,000	-110,000
South America	-15,600	-21,300	-12,800	-15,000
Europe	-104,000	-110,000	-112,000	-103,000
Former Soviet Union	-75,200	-109,000	-111,000	-97,500
Africa	111,000	-68,100	-107,000	147,000
India	-531,000	-606,000	-315,000	62,700
East Asia	-886,000	-926,000	-673,000	-882,000
Southeast Asia	-153,000	-250,000	-103,000	-202,000
Australia	30	-850	-770	-440
Middle East	-168,000	-176,000	-209,000	-127,000

Table S8 – Global burden on mortality of ozone concentrations in the present-day for 2000 concentrations relative to 1850, showing multi-model average and 95% CI including uncertainty in RR and across models (deaths/year), and in 2030, 2050 and 2100 for all RCPs relative to 1850, showing multi-model averages (deaths/year) given by the deterministic values. Also shown, future burdens using (Case A) 2000 concentrations relative to 1850 and present-day population but future baseline mortality rates and (Case B) 2000 concentrations relative to 1850 but future population and baseline mortality rates. These results are plotted in Figure 7. All numbers are rounded to three significant digits.

	2000	2030	2050	2100
Present-day	382,000 (121,000 to 728,400)			
RCP2.6		756,000	1,840,000	1,170,000
RCP4.5		775,000	1,990,000	1,090,000
RCP6.0		891,000	2,600,000	1,570,000
RCP8.5		972,000	2,460,000	2,360,000
Case A		569,000	1,540,000	1,490,000
Case B		735,000	2,090,000	2,040,000

Table S9 – Global burden on mortality of PM_{2.5} concentrations in the present-day for 2000 concentrations relative to 1850, showing multi-model average and 95% CI including uncertainty in RR and across models (deaths/year), and in 2030, 2050 and 2100 for all RCPs relative to 1850, showing multi-model averages (deaths/year) given by the deterministic values. Also shown, future burdens using (Case A) 2000 concentrations relative to 1850 and present-day population but future baseline mortality rates and (Case B) 2000 concentrations relative to 1850 but future population and baseline mortality rates. These results are plotted in Figure 8. All numbers are rounded to three significant digits.

	2000	2030	2050	2100
Present-day	1,700,000 (1,300,000 to 2,100,000)			
RCP2.6		2,360,000	1,820,000	948,000
RCP4.5		2,440,000	1,870,000	559,000
RCP6.0		2,640,000	3,500,000	1,350,000
RCP8.5		2,620,000	2,250,000	1,550,000
Case A		1,590,000	1,440,000	1,230,000
Case B		2,620,000	3,310,000	2,880,000




Figure S1 – Ten world regions




Figure S2 – Regional population-weighted difference in ozone concentrations (ppb) in 2030, 2050 and 2100 relative to 2000. (FSU – Former Soviet Union)




Figure S3 – Regional population-weighted difference in PM_{2.5} concentrations (µg/m³) in 2030, 2050 and 2100 relative to 2000.




Figure S4 – Present-day and future population (millions of people) showing global totals for exposed population (adults 25 and older) from Landscan 2011 (2010) and IFs (2030, 2050, 2100), as well as total population for the RCP scenarios for 2030 and 2100 (Van Vuuren et al., 2011) and for UN Population Prospects 2012 medium fertility scenario for 2030, 2050 and 2100. Also shown are regional exposed populations for IFs.

Sources:

- Oak Ridge National Laboratory (ONRL) - LandScan 2011 Global Population Dataset, <http://spruce.lib.unc.edu.libproxy.lib.unc.edu/content/gis/LandScan/>. Data retrieved on 12/05/2012.
- Web-Based IFs - The International Futures (IFs) modeling system, version 6.54., www.ifs.du.edu. Data retrieved on 07/2012.
- United Nations, Department of Economic and Social Affairs, Population Division (2013). World Population Prospects: The 2012 Revision. <http://esa.un.org/wpp/Excel-Data/population.htm>. Data retrieved on 12/03/2013.




Figure S5 – Global and regional average present-day and future baseline mortality rates (deaths per 1000 people per year) for RESP, IHD, STROKE, COPD and LC, for adults aged 25 and older from the Global Burden of Disease Study 2010 mortality dataset (IHME, 2013) and IFs (2030, 2050, 2100). The IHD and Stroke averages are shown for illustration only, since the mortality estimates are obtained using baseline mortality rates per 5-year age group.

Sources:

- Web-Based IFs - The International Futures (IFs) modeling system, version 6.54., www.ifs.du.edu. Data retrieved on 07/2012.
- IHME (2013). Data retrieved from 12/2013 to 03/2014.




Figure S6 – Future ozone respiratory mortality per million people for all RCP scenarios in 2030, 2050 and 2100, showing the multi-model regional average (deaths/year per million people) in ten world regions (Figure S1) and globally, for future air pollutant concentrations relative to 2000 concentrations.




Figure S7 – Future PM_{2.5} mortality (IHD+STROKE+COPD+LC) per million people for all RCP scenarios in 2030, 2050 and 2100, showing the multi-model regional average (deaths/year per million people) in ten world regions (Figure S1) and globally, for future air pollutant concentrations relative to 2000 concentrations.