The Rate of Equilibration of Viscous Aerosol Particles

Simon O'Meara¹, David O. Topping^{1,2}, Gordon McFiggans¹

¹Centre for Atmospheric Science, School of Earth, Atmospheric & Environmental Sciences, University of Manchester, Manchester, M13 9PL, United Kingdom

5 ²National Centre for Atmospheric Science (NCAS), University of Manchester, Manchester, M13 9PL, United Kingdom

Correspondence to: Gordon McFiggans (g.mcfiggans@manchester.ac.uk)

Abstract. The proximity of atmospheric aerosol particles to equilibrium with their surrounding condensable vapours can substantially impact their transformations, fate and impacts and is the subject of vibrant research activity. In this study we first compare equilibration timescales estimated by three different models for diffusion through aerosol particles to assess any sensitivity to choice of model framework. Equilibration times for diffusion coefficients with varying dependencies on composition are compared for the first time. We show that even under large changes in the saturation ratio of a semi-volatile component (e_s) of 1-90% predicted equilibration timescales are in agreement, including when diffusion coefficients vary

- with composition. For condensing water and a diffusion coefficient dependent on composition, a plasticising effect is observed, leading to a decreased estimated equilibration time with increasing final e_s . Above 60% final e_s maximum equilibration times of around 1 s are estimated for comparatively large particles (10 µm) containing a relatively low diffusivity component (1x10⁻²⁵ m²s⁻¹ in pure form). This, as well as other results here, questions whether particle-phase
- diffusion through water-soluble particles can limit hygroscopic growth in the ambient atmosphere. In the second part of this study, we explore sensitivities associated with the use of particle radius measurements to infer diffusion coefficient dependencies on composition using a diffusion model. Given quantified similarities between models used in this study, our
- 20 results confirm considerations that must be taken into account when designing such experiments. Although quantitative agreement of equilibration timescales between models is found, further work is necessary to determine their suitability for assessing atmospheric impacts, such as their inclusion in polydisperse aerosol simulations.

1 Introduction

10

Recent attention on the phase state of atmospheric particles has motivated questions about the means to model diffusion through them. It had been conventionally assumed that particles possess a liquid phase state, such that timescales of diffusion were much less than their atmospheric residence times. However, several recent studies present evidence that particles can exist in an amorphous solid state (Smith et al., 2002; Murray and Bertram, 2008; Virtanen et al., 2010; Vaden et al., 2011). Viscosities for amorphous solid particles will be higher than for liquid ones, resulting in lower condensed phase diffusion coefficients and potentially limiting the rate of gas-particle partitioning for condensing or evaporating components. For such particles it is important to critically assess models that attempt to predict or infer the effects of diffusion limitations in order to report findings with confidence.

Fick's first and second laws of diffusion state that the rate of transport of a given component through a given area is proportional to the concentration gradient normal to the area. The Fickian diffusion coefficient (D_i) is the proportionality constant between the diffusive flux and the concentration gradient (Eq. (1)).

Recent attempts at modelling diffusion through particles have centred on Fick's second law, which in spherical coordinates is:

$$\frac{\partial C_i(r,t)}{\partial t} = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{D_i \partial C_i(r,t)}{\partial r} \right),\tag{1}$$

where C is the concentration of species i, r is the radius from the particle centre and t is time. Fick's second law is applied 10 when the concentration gradient, and therefore flux, changes with time and distance, i.e. non-steady state. This study analyses and compares three approximations of Eq. (1) used to model diffusion through particles:

i) The ETH model presented by Zobrist et al. (2011), based on the Euler forward step method;

5

ii) The 'kinetic multi-layer model of gas-particle interactions in aerosol and clouds' (KM-GAP) (Shiraiwa et al., 2012), based on coupled differential equations;

15 iii) The Fick's Second Law solved by Partial Differential Equation model (hereafter referred to as Fi-PaD), a formulation of which was used in Smith et al. (2003).

A description of each model is provided in the method below. For a system with D_i independent of composition, it has been reported that Fi-PaD and KM-GAP give very similar results (Shiraiwa et al., 2010). To our knowledge however, no detailed comparison of all three approaches, including cases of D_i dependent on composition, has yet been published.

- 20 Despite this, a recent study by Lienhard et al. (2015) linked the impact of particulate viscosity on ice nucleation using a composition dependent D_i . A critical review of these models is intended to guide those with an interest in simulating particle evolution inside instruments, chamber experiments, and the ambient atmosphere. For non-equilibrium viscous particles, diffusivity (along with other properties such as volatility) determines the temporal evolution of particle composition and size- and number-distributions (Zaveri et al., 2014). These are key factors determining aerosol impact on climate and health,
- 25 therefore the choice of diffusion model could have far-reaching consequences (Pöschl, 2005). In addition to differences in modelled particle size and composition change, inappropriate choice of model formulation and assumptions therein could lead to differences in inferred properties, such as diffusion coefficients from single particle levitation measurements (Lienhard et al., 2014; Zobrist et al., 2011).

The numerical methods employed by all three models involve discretisation in time and space. However, subtle 30 differences in how they define concentration gradients may induce variations in estimated diffusion rate. Therefore, it is expected that any differences in rate will increase with greater heterogeneity in the concentration-radius profile, i.e. an increasingly steep diffusion front. Such fronts have been observed when water and glassy organics diffuse through one another (Nowakowski et al., 2015). It is currently unclear which of the models investigated here, if any, is suitable to such a situation, given the paucity of experimental data available.

Indeed, the Fickian framework may not be appropriate for some systems; in polymer studies it is well known that non-Fickian diffusion occurs for many examples of liquids diffusing through glassy polymers (Thomas and Windle, 1982;

- 5 Kee et al., 2005), and in such systems a narrow diffusion front is often observed (e.g. Thomas and Windle, 1982). Non-Fickian diffusion results from structural changes following diffusion and the resultant composition change. It arises when the rate of deformation is comparable to that of diffusion (Crank 1975). Alternative models have been proposed, such as the free volume model (He et al., 2006; Price et al., 2014) and the Maxwell-Stefan model (Krishna and Wesselingh, 1997). The aim of this study, however, is to compare the estimated equilibration timescales of the Fickian diffusion models that are used
- 10 in atmospheric aerosol science and, in turn, assess sensitivities of derived diffusion coefficients in such particles. In most test cases below the diffusing semi-volatile component has the self-diffusion coefficient of water at room temperature, and the resulting diffusion timescales are most relevant to water and water-soluble particles, however, the findings regarding consistency between models are applicable to components with self-diffusion coefficients across the investigated range $(2x10^{-9} 1x10^{-25} \text{ m}^2\text{s}^{-1})$.

15 2 Method

2.1 Model Description

The ETH model, KM-GAP and Fi-PaD used the same representation of an aerosol particle: it was assumed spherical, and split into concentric shells. A comparatively thin surface shell was assumed to equilibrate instantly with the gas-phase in all simulations (for the purpose of comparing particle-phase diffusion models and gaining insight into the limitation imposed by

- 20 particle-phase diffusion on mass-transfer this assumption is reasonable). The initial concentration profile in bulk shells (those below the surface) was homogeneous, and in equilibrium with the initial gas phase concentration. Figure 1a demonstrates how the particle is represented in a 2-D view. Fig. 1b illustrates the concentration-radius profiles of a semivolatile component at several time steps using the ETH model in the case of an instantaneous increase in saturation ratio from 1 to 90% when the diffusion coefficient is independent of composition. In contrast, Fig. 1c shows the same information but when the diffusion coefficient has a logarithmic dependence on composition and the self-diffusion
- coefficients of the two components are very different, that of the non-volatile $(D_{nv}^0) = 1 \times 10^{-21} \text{ m}^2 \text{s}^{-1}$ and for the semi-volatile $(D_{sv}^0) = 2 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$. The "diffusion front" is clear in this example and arises from the very different diffusion coefficient values in neighbouring shells that result from variations in shell composition.

The *e*-folding time for the difference in concentration of the semi-volatile component at the surface $([sv]_{eq})$ and of 30 its average concentration across the particle bulk $(\overline{[sv]}_b)$ was used as a metric for diffusion time by Zaveri et al. (2014). It is readily transferable to other studies, and is the chosen metric for diffusion timescale here. The ratio of the concentration difference in the surface and bulk-average of the semi-volatile component at any time (t) to that difference at t = 0 is:

$$Q = \frac{|[sv]_{eq} - \overline{[sv]}_{b,t\geq 0}|}{|[sv]_{eq} - \overline{[sv]}_{b,t=0}|}.$$
(2)

The *e*-folding time was therefore the time taken for *Q* to increase/decrease by a factor of e^1 . Figures 1b and 1c therefore demonstrate the concentration-radius profiles at several steps between Q = 1, which occurs at t = 0, and $Q = e^{-1}$, which marks the *e*-folding time. Comparing *e*-folding times between models strictly only tests model consistency at this particular stage of diffusion and not before this. However, *e*-folding time agreement would indicate agreement at previous times (and future ones), because the underlying equations are identical. For reassurance on this, concentration-radius profiles at times prior to the *e*-folding time were compared. All models were run on a 12 core Intel Core i7 processor with a speed of 3.2 ID GHz.

Fick's first law, which assumes a constant concentration gradient and therefore flux, with time, is the basis of the ETH model. However, following the Euler forward step method, if time steps are sufficiently short this model should be able to capture changes in concentration gradient, and therefore flux, and therefore replicate the second law. The flux between shells is thus found by:

15
$$J_{bk,bk-1,i} = -A_{bk}D_{bk,bk-1,i} \frac{([i]_{bk}-[i]_{bk-1})}{0.5(\delta_{bk}+\delta_{bk-1})},$$
 (3)

where *J* is the flux (mol s⁻¹) between bulk (*b*) shell numbers *k* and k - 1 (ascending from k = 1 for the near-surface shell to k = n at the centre). If k = 1, then k - 1 is the surface layer (*s*). *A* is the surface area of the shell's outer surface. $D_{bk,bk-1,i}$ is the diffusion coefficient at the shell boundary and here is found using one of the dependencies on composition given in Sect. 2.3. [*i*] is concentration of component *i* at the shell centre and δ is shell width. Figure 1a demonstrates how

20 these terms relate to the physical representation of the particle. From Eq. (3) the change in number of moles in a shell is found by:

$$\Delta N_{bk,i} = (J_{bk,bk-1,i} - J_{bk+1,bk,i})\Delta t, \tag{4}$$

where Δt is the time interval (setting of Δt is described in Sect. 2.2). The version of the ETH model we have written was tested against the model output in Zobrist et al. (2011), and found to replicate their results accurately (Fig. A1 of the appendix for the replica plot).

In KM-GAP the number of moles of a component in a shell is found by integrating the following coupled ordinary differential equations (ode) with respect to time:

$$\frac{dN_{s,i}}{dt} = (J_{b1,s,i} - J_{s,b1,i}),\tag{5}$$

$$\frac{dN_{b1,i}}{dt} = (J_{s,b1,i} - J_{b1,s,i}) + (J_{b2,b1,i} - J_{b1,b2,i}), \tag{6}$$

$$30 \quad \frac{dN_{bk,i}}{dt} = (J_{bk+1,bk,i} - J_{bk,bk+1,i}) + (J_{bk-1,bk,i} - J_{bk,bk-1,i}), (k = 2, \dots, n-1),$$
(7)

25

. . .

$$\frac{dN_{bn,i}}{dt} = (J_{bn-1,bn,i} - J_{bn,bn-1,i}).$$
(8)

The flux is found by:

10

30

$$J_{bk,bk\pm1,i} = K_{bk,bk\pm1,i}[i]_{bk}A_{bk},$$
(9)
where K is the transport rate coefficient (m s⁻¹):

$$K_{bk,bk\pm1,i} = \frac{2D_{bk,bk\pm1,i}}{(\delta_{bk}+\delta_{bk\pm1})},$$
(10)

5 where δ is shell width. $D_{bk,bk\pm 1,i}$ is the diffusion coefficient at the shell boundary (Sect. 2.3). Note that in Eqs. (9) and (10), if k = 1, then k - 1 is the surface shell (s).

Equations (5)-(10) were solved in Matlab software using the ode23tb numerical solver, which has an adaptive time step. It was found that the solver became increasingly unstable as the gradient of D_i with r increased, thus error tolerances (given in the appendix) were increased appropriately. ode23tb uses a Runge-Kutta method of two stages: a trapezoidal rule followed by a backward differentiation formula stage.

Fi-PaD treats Eq. (1) as an initial-boundary problem, with initial conditions:

$$C_{bk,i}(r < R_p, 0) = C_{i,eq^0},\tag{11}$$

$$C_{s,i}(R_p, 0) = C_{i,eq},$$
(12)

where eq represents the equilibrium condition. Equation (11) states that initial concentrations in the bulk shells are in 15 equilibrium with the original e_s value, whilst Eq. (12) states that the initial concentration at the surface is in equilibrium with the new e_s . The boundary conditions were:

$$\frac{\partial N_i(0,t)}{\partial t} = 0,\tag{13}$$

$$\frac{\partial C_i(R_p,t)}{\partial t} = 0,\tag{14}$$

where R_p is the particle radius. Equation (13) states that there is no flux at the centre of the particle and Eq. (14) states that

- 20 the concentration of components at the surface is constant. For Fi-PaD the numerical solver pdepe in Matlab software was used. The solver uses the method of lines, which discretises the problem in space to gain a system of ordinary differential equations that are then solved using the numerical solver ode15s in Matlab. ode15s is similar to ode23tb in that both are designed for stiff systems, however, ode15s has a high order of accuracy for a given error tolerance. The default error tolerances for pdepe were found to provide stable solutions across the range of parameter spaces used here; the contrast to
- the variable error tolerances used in KM-GAP is attributed to the difference in the accuracy of their ode solvers.

2.2 Particle Representation

Particles were assumed to consist of two components: a non-volatile (nv) and semi-volatile (sv), which were assigned the molar mass and density of sucrose and water, respectively. In general, components with relatively high molar masses are expected to have comparatively low diffusion coefficients (Haynes, 2015). To test the effect of using a high molar mass component against using sucrose on equilibration times, a molar mass (M) of 700 g mol⁻¹ (M of sucrose = 342.296 g mol⁻¹)

and density (p) of 2.0 x10³ kg m⁻³ (p of sucrose = 1.5805×10^3 kg m⁻³) was assigned to the non-volatile component and its



self-diffusion coefficient was set relatively low: $1.0 \times 10^{-25} \text{ m}^2 \text{ s}^{-1}$. When the saturation ratio of the semi-volatile component (e_s) increased from 1-90% the *e*-folding times for all three models increased between 13-16% from those using sucrose values (since the molar volume of the non-volatile component increased a decreased semi-volatile component concentration was required to attain equilibrium, leading to a decreased concentration gradient for the same change in e_s). Since these changes to *e*-folding times are similar across the models and are for a comparatively large change in e_s , our conclusions are expected to be applicable to a broad range of component M and *p* values.

All models assumed ideality for most simulations (see later) so that at equilibrium the value of e_s equalled the mole fraction of the semi-volatile component in the condensed phase. While estimates of accommodation coefficients for semi-volatiles cover a wide range, for the purposes of this study we have held it constant at unity (as has been found reasonable

10 for that of water vapour on liquid water in multiple studies, e.g. Kolb et al. (2010)). Assuming ideality, the volume of a component was equal to the product of its number of moles (*N*) and molar volume. The volume of a shell was therefore given by:

$$V_{bk} = N_{nv,bk} \left(\frac{M_{nv}}{p_{nv}}\right) + N_{sv,bk} \left(\frac{M_{sv}}{p_{sv}}\right),\tag{15}$$

where M is molar mass and p is density.

5

To accurately simulate the size change in particles resulting from condensation (growth) or evaporation (shrinkage), at the end of each time interval, shell volumes were recalculated using the new values of $N_{i,bk}$. In KM-GAP and Fi-PaD, a maximum change to the particle radius of 0.1% was allowed per time step; if the radius change exceeded 0.1% the interval was iteratively shortened until the change was acceptable. Decreasing this maximum acceptable change did not change *e*folding times significantly (<2% for both KM-GAP and Fi-PaD when a maximum radius change of 0.01 % was used

- 20 instead), thus it was considered sufficient to account for volume change. For the ETH model, it has been recommended that to ensure model stability, the number of moles inside any shell should not change by more than 2% over a single time interval (Zobrist et al. 2011). The same condition was used here because values below 2% did not change predicted *e*-folding times significantly (<1% change when maximum change in number of moles was 0.01% instead).</p>
- Bulk shells (those below the surface) were initially set to have equal widths. The surface shell represents the sorption layer, where transfer between the condensed and gas-phase occurs. Since the surface shell is contained within the initial particle diameter, the width should be sufficiently thin to not significantly affect the *e*-folding time, i.e. one must not decrease the width of bulk shells such that diffusion is accelerated. A factor of 1×10^{-3} of the particle radius was chosen to calculate the surface shell width because using lower factors resulted in no significant change to estimated *e*-folding time.
- During condensation the surface shell expands; however, since this shell simulates the boundary between the shell and the 30 gas-phase it should remain comparatively thin. Therefore, if the surface shell grew to double its initial width, it was reduced back to its initial width by transferring the excess volume to the near-surface shell, or, if this near-surface shell had a width greater than the total radius divided by the number of shells, the transferred material was used to make a new near-surface shell. The concentration of components in the transferred material was the same as in the surface shell (i.e. at equilibrium

with the gas phase). This approach had potential to introduce numerical diffusion by decreasing the distance for diffusion in the case of introducing a new shell and decreasing the concentration gradient in the case of transfer.

To gain an indication of whether numerical diffusion influenced one model more than another, e-folding times were found with this approach (transfer on) and without it (transfer off). For the latter case the surface shell was allowed to grow 5 without adjustment, leading to an unrealistically wide shell and comparatively longer equilibration times, but eliminating the possibility of numerical diffusion. If numerical diffusion affected one model more than another we would expect the difference in *e*-folding times between the transfer on and transfer off cases to vary between them. However, there was no substantial difference between models: for a change in e_s of 1-90 % all models had an increase of 20-30 % in e-folding times from the transfer on to the transfer off case; and for a change in e_s of 60-80% the increase was between 6-10%. These differences are negligible in comparison to the several orders of magnitude change in e-folding times seen across the range

- 10
- of non-volatile component diffusivity used below.

During evaporation the width of the surface shell decreased and the mass of non-volatile component in the surface shell tended toward zero. If the surface shell decreased below a factor of 1×10^{-1} of its initial width it was returned to its initial width by transferring a sufficient volume from the near-surface shell. The concentration of components in the

- 15 transferred material was equal to that in the surface shell, thus the concentration in the surface shell was maintained and any excess semi-volatile component was presumed to evaporate. Similarly, if the near-surface shell shrank to below a factor of 1x10⁻¹ of the initial width of the surface shell, then the two shells were coalesced into a new surface shell at equilibrium concentration. It was found that decreasing the width at which transfer and coalescence were invoked led to a decrease and convergence of predicted *e*-folding times, indicating decreasing numerical diffusion (which could occur due to steepening of
- the concentration gradient through either coalescence or transfer). A decrease of no more than 1% was seen across models 20 and changes in e_s when using lower factors than 1×10^{-1} of the initial width of the surface shell, thus this factor was concluded to be sufficiently low to effectively prevent numerical diffusion.

2.3 D Dependence

At any point in the particle the diffusion coefficient of both components was the same, i.e., we assumed symmetrical 25 diffusion coefficients, which is valid for an ideal binary mixture (Wesselingh and Bollen, 1997). We compared models using three functions of D_i:

i) D_i independent of the semi-volatile mole fraction (x_{sv}) and therefore fixed throughout the simulation;

ii) D_i with a logarithmic dependence on semi-volatile mole fraction, which has been observed for ideal systems by Vignes (1966):

 $D_{i}(x_{sv}) = D_{sv}^{0} D_{nv}^{0}^{(1-x_{sv})},$ 30 (16)

where D_{sv}^0 is the self-diffusion coefficient of the semi-volatile component and D_{nv}^0 is the self-diffusion coefficient of the nonvolatile component;

iii) D_i with a sigmoidal dependence on x_{sv} , which was observed for the citric acid-water system by Lienhard et al. (2014):

$$D_i(x_{sv}) = D_{sv}^{0} {}^{x_{sv}\alpha} D_{nv}^{0} {}^{(1-x_{sv}\alpha)},$$
(17)

where \propto is a correction parameter given by:

 $\ln(\alpha) = (1 - x_{sv})^2 [C + 3D - 4D(1 - x_{sv})].$ ⁽¹⁸⁾

5 Where the values of *C* and *D* were chosen as -3.105 and 3.300 respectively. These provided a relatively steep "cliff-edge" sigmoidal dependence and therefore a substantial variation from the logarithmic dependence, enabling a test of consistency between models across a wide range of dependencies. Examples of these dependencies are shown in Fig. 2.

For the latter two cases $D_i(x_{sv})$ was calculated within the numerical solvers of KM-GAP and Fi-PaD, whilst for the ETH model it was calculated at the start of each time step. x_{sv} at a shell boundary was found using the arithmetic mean concentration of the semi-volatile component across the bounding shells.

In the first part of the study we compare the equilibration timescales estimated by models when the diffusion coefficient is constant and when it follows the logarithmic and sigmoidal dependencies on composition given above. Self-diffusion coefficients of the non-volatile and semi-volatile components range between that of water at room temperature, $2.0 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$ (Starr et al., 1999), and a comparatively low value of $1.0 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$, which according to the Stokes-Einstein

15 relationship between diffusivity and viscosity, is representative of a glassy material (Debenedetti and Stillinger, 2001). *e*folding times are found for several changes in the vapour-phase saturation ratio of a semi-volatile component, and across a range of particle sizes and differences in the self-diffusion coefficient of components.

Finally, we present an example of the differences in modelled particle size change with time when different dependencies of D_i on composition are assumed, thereby providing guidance on the most effective experimental procedure

20 for inference of diffusion coefficient dependencies on composition. For actual inferences one would preferably have good knowledge of the system's deviation from ideality. In an attempt to replicate a real system, we therefore use the estimation for water activity and density as a function of sucrose weight fraction presented in Zobrist et al. (2011). The initial and surface shell water activity were set equal to the initial and current gas-phase saturation ratio of water (the saturation ratio changed with time), respectively, with the accommodation coefficient of water assumed to be one.

25 3 Results

30

Numerical convergence of *e*-folding times was observed with increasing spatial resolution for all three models due to improved resolution of concentration gradients and therefore changes in D_i (when dependent on composition) and flux with space. *e*-folding times showed an exponential relationship with shell number (e.g. Fig. A2 of the appendix), thus the criteria for shell number was that at which the *e*-folding time was within 10% of the asymptote. Generally as the gradient of D_i with particle radius increased, the shell number increased to maintain convergence (Table A2 of the appendix). However, increasing the shell number increases the possibility of accumulating significant round-off error, in addition to requiring

greater computer time. The round-off error at the chosen resolution was investigated by halving the number of significant



numbers assigned to variable values. The difference in predicted e-folding times between the two precisions was found to be negligible, with a maximum of 2%, indicating that round-off error was not a substantial source of inaccuracy.

Zobrist et al. (2011) reported requiring up to several thousand shells in the ETH model to resolve concentration gradients. However, we found that using of the order of hundreds gave convergence for the cases with steepest concentration

- 5 gradients (Fig. A2 of the appendix). The difference in required shell resolution between the studies could be due to differences in D_i dependence on composition. Using the Matlab software it was found that computational time for the case of diffusion coefficient independent of composition was quickest, gradually increasing as the steepness of the diffusion coefficient dependence on composition increased, largely due to the greater spatial resolution. For D_i independent of composition the ETH model took of the order 1 s to reach the e-folding state while KM-GAP and Fi-PaD were of the order
- 10^2 s. For a steep diffusion coefficient dependence, the chosen example was the logarithmic dependence, with $D_{nv}^0 = 1 \times 10^{-25}$ 10 m^2s^{-1} and $D_{sv}^0 = 2x10^{-9} m^2s^{-1}$ and e_s instantaneously increased from 1-90%: to reach *e*-folding states the ETH model took of the order 10^2 s while both KM-GAP and Fi-PaD took of the order 10^4 s.

In the first model comparison, e-folding times were found when D_i was independent of x_{sv} . For a complete analysis of model output, initial particle diameters $(D_{p,t=0})$ were varied between $1x10^{-5}$ m and $1x10^{-8}$ m, which covers most of the size range observed in the ambient atmosphere (Seinfeld and Pandis, 2006) and D_i ranged between 2.0×10^{-9} m²s⁻¹ and 15 $1.0 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$. *e*-folding times were found across this parameter space for a change in e_s of 1-90% and 90-1% for all three models. This relatively large change in e_s was chosen to create a large concentration gradient, as this would most likely induce disagreement between models. However, all models agreed very well across the whole range of particle size and D_i (Fig. A3).

In the next case D_i varied logarithmically with mole fraction of the semi-volatile, between a maximum of 2.0×10^{-9} 20 $m^2 s^{-1}$ at $x_{sv} = 1$ and a minimum given by D_{nv}^0 (i.e. D_i at $x_{sv} = 0$). D_{nv}^0 ranged between 2.0×10^{-9} and $1.0 \times 10^{-25} m^2 s^{-1}$. Contour plots of e-folding times as a function of D_{nv}^0 and $D_{p,t=0}$ and for a 1-90% and a 90-1% change in e_s are shown in Figs. 3a and 3b, respectively.

For both changes in e_s there is good agreement of e-folding times between all models, with a maximum variation of 10 %, which is well within the uncertainty caused by varying degrees of numerical convergence and potential numerical 25 diffusion. Diffusion times are much shorter than in the constant D_i case due to the high diffusivity of the semi-volatile component. Fig. 3a shows that even when starting with a glassy particle, if the saturation ratio of a plasticising semi-volatile component increases sufficiently, the e-folding state can be reached in less than 1 s. For the decreasing e_s used in Fig. 3b a low diffusivity outer casing will form on the particle, impeding diffusion and evaporation. However, Fig. 3b shows that if a particle initially of water-like diffusivity is quickly dried, the e-folding state is reached within 10 s, even when the non-30

volatile component has a relatively low diffusivity. *e*-folding times for 1-90% and 90-1% changes in e_s were also found using the sigmoidal dependence of D_i on x_{sv} ;

the results are given in Figs. 3c and 3d, respectively. In the 90-1% case an unpractical computer time (>12 hours) was



required to attain numerical convergence at low values of D_{nv}^0 , therefore the minimum D_{nv}^0 is 1×10^{-20} m²s⁻¹. For this relatively large change in e_s the sigmoidal dependence induces a steeper diffusion front than the logarithmic dependence. Despite this, the models show good agreement here also. In the 1-90% case, a maximum variation in *e*-folding times of 5% is seen while for 90-1% this value is 30 %. This latter variation is between KM-GAP and the other two models and is greater than expected from different degrees of numerical convergence. However, given the gradual divergence of the *e*-

- 5 greater than expected from different degrees of numerical convergence. However, given the gradual divergence of the *e*-folding isolines in Fig. 3d, we do not attribute the discrepancy to model framework differences, but to an insufficient shell resolution in KM-GAP. Diffusion is quicker using the sigmoidal dependence than the logarithmic dependence, particularly for the 90-1% scenario. This is explained by the higher D_i values at x_{sv} >0.5 (Fig. 2).
- e_s changes more realistic of the atmosphere were also tested. Results for 60-80% and 80-60% (Fig. A4) are similar to those for 1-90% and 90-1% for their respective D_i dependency; there is good model agreement, and across the $D_{p,t=0}$ and D_{nv}^0 range and for both dependencies, *e*-folding time is less than 1 s. Results for 10-20% and 20-10%, given in Fig. 4, also show agreement between models. For both dependencies diffusion is much slower than in the 1-90% and 60-80% simulations, approaching 1 ky at low D_{nv}^0 and high $D_{p,t=0}$. This shows that at low saturation ratios of semi-volatile component, gas-particle partitioning can be limited by condensed-phase diffusion in viscous particles.
- 15 *e*-folding times between models were also found to be in good agreement for these changes in e_s when D_{nv}^0 was fixed at $1.0 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$ and D_{sv}^0 was varied between $1.0 \times 10^{-25} - 2 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$. As discussed, the agreement between models in estimating *e*-folding times indicates that the estimated profiles of concentration with particle radius prior to the *e*-folding state are consistent between models because the underlying equations are the same. By comparing concentration-radius profiles at various stages of diffusion we indeed found good model agreement across all cases. In Fig. 5 we show the 20 example of the logarithmic dependence of D_i on x_{sv} , an instantaneous change in saturation ratio of 1-90% and with $D_{nv}^0 = 1 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$ and $D_{sv}^0 = 2 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$. At several times preceding and including *e*-folding time the concentrationradius profiles are in good agreement.

In the final part of this study the estimated temporal profile of particle radius was compared between the sigmoidal and logarithmic D_i dependencies. We have used the water activity and density dependence on sucrose weight fraction as 25 described in Zobrist et al. (2011) for the sucrose-water system in an attempt to replicate a non-ideal system. The ETH model was employed, though the results above indicate that KM-GAP and Fi-PaD would produce identical profiles. For the inference of D_i dependency from radius measurements the signal to noise ratio is minimised by inducing a large change in radius relative to the measurement accuracy over a time-span that is large compared to the measurement frequency.

Taking the case of water as the semi-volatile component, from Fig. 3 it is clear that for certain values of D_{nv}^0 and certain changes in e_s attaining a large ratio of equilibrium time to measurement frequency may be difficult, even if the change in radius is large. Indeed, the radius-time profiles in Figs. 6 and 7 for instantaneous changes in e_s and a $D_{nv}^0 = 1 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$ confirm that for changes with a high final e_s , significant radius change is estimated to occur over less than 1 s, while the measurement frequency reported in the studies of Zobrist et al. (2011) and Lienhard et al. (2014) is approximately 15 s.

Nevertheless, for the e_s change of 1-90% in Fig. 5, there is a notable difference in the radius profiles between the dependencies. Despite having lower D_i at low x_{sv} , the radius change from the sigmoidal dependence is more rapid than the logarithmic, indicating that the D_i at higher x_{sv} has a dominating effect on the profile. The inference of D_i dependency using such a large change in e_s is therefore poorly constrained for lower x_{sv} . For better constraint smaller changes in e_s are

- 5 required, such as those used in Lienhard et al. (2014). An example of the radius profiles following incremental changes in e_s , $D_{nv}^0 = 1 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$, $D_{sv}^0 = 2 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$ and using both dependencies is shown in Fig. 7. This plot demonstrates the need for consideration of the time a given e_s is maintained in measurement experiments, since the difference in the equilibrium timescales between the e_s increments covers several orders of magnitude. Indeed, over low changes in e_s such as between 1-10%, equilibration time may be too long to be practical for gaining a useful measurement of radius change. It is worthwhile
- 10 to note that the rate of change of e_s over an increment is preferably much greater than the rate of equilibration, as this provides the greatest potential for a clear signature of the D_i dependence and therefore greatest constraint on inference.

4 Discussion and Conclusion

The results above show that despite variations in their numerical methods, all three Fickian-based diffusion models tested here: the ETH model, KM-GAP and Fi-PaD give good agreement of estimated *e*-folding timescales over a wide range of changes to the saturation ratio of the semi-volatile component and over a wide range of differences in the self-diffusion coefficient of the semi-volatile and non-volatile components. Furthermore, there is good agreement between models when different dependencies of diffusion coefficient on composition are used. This result has not been reported before to our knowledge and verifies consistency between existing Fickian diffusion models. The maximum disagreement in *e*-folding times for results gained with satisfactory shell resolution is 10%, which is within the uncertainty generated by varying

- 20 degrees of numerical convergence and potential numerical diffusion. The consistency in modelled concentration-radius profiles at times preceding and including the *e*-folding state (Fig. 5) shows that if used for a polydisperse aerosol population, the models would give agreement in changes to the size distribution. In addition, if the diffusing component were reactive the rate of particle-phase reaction would depend on its concentration; therefore model agreement in concentration-radius profiles would give consistent reaction rates across the particle (which in turn could affect diffusion rate).
- 25 Using the three diffusion models as described above and with the spatial resolutions presented in the appendix, the ETH model takes approximately two orders of magnitude less computer time than Fi-PaD or KM-GAP for a given diffusion scenario. With the models giving consistent estimates of diffusion, the ETH model therefore appears to be favourable.

The *e*-folding times given in Fig. 3 for changes in e_s of 1-90% and 90-1%, and in Fig. A4 for changes of 60-80% and 80-60%, show that for a semi-volatile component with water-like (at room temperature) diffusivity, given a sufficiently high starting/finishing e_s , attainment of the *e*-folding state is effectively instant compared to residence times in the atmosphere and chamber experiments. This is due to the plasticising effect of water (and applies to any semi-volatile component with a sufficiently high self-diffusion coefficient). At lower values of e_s diffusion time can be much longer (Fig.

4)), consistent with measurement studies (e.g. Zobrist et al., 2011 and Lienhard et al., 2014). The question therefore arises that for a given D_{sv}^0 , at what e_s can equilibration be assumed instant? Figure 7 indicates that for water condensing at room temperature equilibration time is less than 1 s when the final e_s is greater than 50% for the sigmoidal dependence used here and when it is greater than 60% for the logarithmic dependence. These results indicate no limitation on mass transfer of

5

water from particle-phase diffusion at high relative humidity and at ambient temperature, and therefore no impediment to the formation of cloud droplets. Experimental results from Lienhard et al. (2014) and Zobrist et al. (2011) indicate that this is also true down to ~250 K.

For a hygroscopicity tandem differential mobility analyser (HTDMA), which has a typical residence time of 20-25 s (Zardini et al. 2008) and e_s change of <10-~90%, Figs. 3 and 5 show that equilibration is attained even when sampling 10 relatively large (~1x10⁻⁵ m) particles containing components of relatively low diffusivity (self-diffusion coefficients ~1x10⁻²⁵ m²s⁻¹).

Note, however, that we have not considered extreme dependencies of D_i on composition. If, for example, a very high mole fraction of water were required before the "cliff-edge" in the sigmoidal dependence (Fig. 2)) was reached, longer diffusion times than those shown here would be expected.

15

Regarding the inference of diffusion coefficient dependence on composition from particle radius measurements, we have shown that incremental changes in e_s provide the best constraint, and note that changes should occur over a short time compared to the equilibration time. The consistency between the diffusion models shown here indicates that the choice of model does not affect the accuracy of the inferred dependence (as long as sufficient spatial and temporal resolution is used). In a follow-up study we intend to investigate the implementation of composition-dependent D_i in the Model for Simulating

- 20 Aerosol Interactions and Chemistry (MOSAIC) (Zaveri et al., 2008). MOSAIC is used for chamber and ambient studies and can therefore include, among other factors, multiple components, chemistry and volatility. Furthermore, it can model polydisperse aerosol, providing insight into how composition-dependent diffusion coefficients affect the evolution of size distributions.
- As mentioned, Fickian-diffusion is, strictly speaking, limited to ideal-systems. Thus, for cases where dissolution occurs, for example, the employed or derived diffusion coefficients are actually effective values of D_i . As we mention briefly in the introduction, numerous alternative theories to Fickian diffusion exist. Although an analysis of such frameworks is beyond the scope of this study, a similar critical analysis may be useful in the future when data from more complex multicomponent systems exist.

Appendix A

30 To validate our version of the ETH model Figure 3d of Zobrist et al. (2011) was reproduced using our version of the model and the relative humidity measurements presented in their Fig. 3a. Note that in reproducing this figure the dependence of diffusion coefficient on water activity given in Zobrist et al. (2011) was used. Furthermore, water activity and density estimated as a function of sucrose weight fraction, as described in Zobrist et al. (2011), was used. Our reproduction is given in Fig. A1.

Figure A2 shows the convergence of *e*-folding times with increasing shell number for the ETH model. Results are for self-diffusion coefficients of the semi-volatile and non-volatile components of 2.0×10^{-9} m²s⁻¹ and 1.0×10^{-25} m²s⁻¹, respectively, with a logarithmic dependence of D_i on composition and change to the vapour-phase saturation ratio of the semi-volatile of 1-90% (Fig. A2a) and 90-1% (Fig. A2b). These cases were chosen because they are expected to have the strongest concentration and diffusion coefficient gradients through the particle (compared to other cases in this study) and should therefore require greatest spatial resolution. The exponential fits in Fig. A2 were obtained using Igor Pro software.

The absolute tolerances that were required to attain stability in the KM-GAP model are given in Table A1. The 10 tolerance was dependent on the self-diffusion coefficient of the non-volatile and the initial particle diameter. The relative tolerance was kept fixed at 1.0×10^{-12} . These tolerances were used for changes to the semi-volatile saturation ratio of 1-90% and 90-1%. Since these represent the largest changes in saturation ratio used in this study, the tolerances in Table A1 are conservative values for all other cases presented in the study.

The number of shells used for each model is given in Table A2. The optimum number of shells required for acceptable numerical convergence was found to be dependent on the change in semi-volatile saturation ratio and the difference in the self-diffusion coefficients of the components.

Results for model estimates of *e*-folding times when the diffusion coefficient was kept constant are given in Fig. A3. Good agreement can be seen between all three models across the values of constant diffusion coefficient and initial particle diameter. In Fig. A4 are the *e*-folding times for changes of 60-80 and 80-60% in the saturation ratio of the semi-volatile component using the logarithmic and sigmoidal dependencies of diffusion coefficient on semi-volatile component

mole fraction.

20

5

Author Contributions

S.O. wrote the model codes, ran the simulations, created plots and tables and wrote the manuscript. D.O.T. and G.M. provided substantial input to the method, model development and manuscript.

25 Acknowledgments

The Natural Environment Research Council has funded this work through the Ph.D. studentship of S. O'Meara, grant number NE/K500859/1, and through grant number NE/J02175X/1.

References

15

20

25

Crank, J.: The Mathematics of Diffusion (2nd Edition), Clarendon Press, Oxford, 1975.

Debenedetti, P.G., and Stillinger, F.H.: Supercooled Liquids and the Glass Transition, Nature, 410, 259-267, 2001.

Haynes, W.M. (Ed.): CRC Handbook of Chemistry and Physics, 96th edn., Internet Version 2016, www.hbcpnetbase.com., 2015.

He. X., Fowler, A., and Toner, M.: Water Activity and Mobility in Solutions of Glycerol and Small Molecular Weight Sugars: Implication for Cryo- and Lyopreservation, J. Appl. Phys., 100(074702), doi: 10.1063/1.2336304, 2006.

Kee, D. D., Liu, Q., and Hinestroza, J.: Viscoelastic (non-Fickian) Diffusion, Can. J. Chem. Eng., 83, 913-929, 2005.

 Kolb, C. E., Cox, R. A., Abbatt, J. P. D., Ammann, M., Davis, E. J., Donaldson, D. J., Garrett, B. C., George, C., Griffiths, P.
 T., Hanson, D. R., Kulmala, M., McFiggans, G., Pöschl, U., Riipinen, I., Rossi, M. J., Rudich, Y., Wagner, P. E., Winkler, P. M., Worsnop, D. R., and O'Dowd, C. D.: An Overview of Current Issues in the Uptake of Atmospheric Trace Gases by Aerosols and Clouds, Atmos. Chem. Phys., 10(21), 10561–605, 2010.

Krishna, R., and Wesselingh, J.A.: The Maxwell-Stefan Approach to Mass Transfer, Chem. Eng. Sci., 52(6), 861–911, 1997.

Lienhard, D. M., Huisman, A. J., Bones, D. L., Te, Y.F., Luo, B. P., Krieger, U. K., and Reid, J. P.: Retrieving the Translational Diffusion Coefficient of Water from Experiments on Single Levitated Aerosol Droplets, Phys. Chem. Chem. Phys., 16, 16677–83, 2014.

Lienhard, D. M., Huisman, A. J., Krieger, U. K., Rudich, Y., Marcolli, C., Luo, B. P., Bones, D. L., Reid, J. P., Lambe, A. T., Canagaratna, M. R., Davidovits, P., Onasch, T. B., Worsnop, D. R., Steimer, S. S., Koop, T., and Peter, T.: Viscous organic aerosol particles in the upper troposphere: diffusivity-controlled water uptake and ice nucleation?, Atmos. Chem. Phys., 15, 13599-13613, doi:10.5194/acpd-15-13599-2015, 2015.

- Murray, B. J, and Bertram, A. K.: Inhibition of Solute Crystallisation in Aqueous H(+)-NH(4)(+)-SO4(2-)-H2O Droplets, Phys. Chem. Chem. Phys., 10(22), 3287–3301, 2008.
- Price, H.C., Murray, B.J., Mattsson, J., O'Sullivan, D., Wilson, T.W., Baustian, K.J., and Benning, L.G.: Quantifying Water Diffusion in High-Viscosity and Glassy Aqueous Solutions Using a Raman Isotope Tracer Method, Atmos. Chem. Phys., 14, 3817-3830, 2014.
- Ulrich, P.: Atmospheric Aerosols: Composition, Transformation, Climate and Health Effects, Angew. Chem. Int. Edit., 44, 7520–40, 2005.
- Seinfeld, J. H., and Pandis, S. N.: Atmospheric Chemistry and Physics., 2nd ed., John Wiley & Sons Inc., Hoboken, NJ, USA, 2006.

- Shiraiwa, M, Pfrang, C., and Pöschl, U.: Kinetic Multi-Layer Model of Aerosol Surface and Bulk Chemistry (KM-SUB): The Influence of Interfacial Transport and Bulk Diffusion on the Oxidation of Oleic Acid by Ozone, Atmos. Chem. Phys., 10, 3673–91, 2010.
- Shiraiwa, M., Pfrang, C., Koop, T., and Pöschl, U.: Kinetic Multi-Layer Model of Gas-Particle Interactions in Aerosols and Clouds (KM-GAP): Linking Condensation, Evaporation and Chemical Reactions of Organics, Oxidants and Water, Atmos. Chem. Phys., 12(5), 2777–94, 2012.
 - Smith, G. D., Woods, E., DeForest, C.L., Baer, T., and Miller, R.E.: Reactive Uptake of Ozone by Oleic Acid Aerosol Particles: Application of Single-Particle Mass Spectrometry to Heterogeneous Reaction Kinetics, J. Phys. Chem. A, 106, 8085–95, 2002.
- 10 Smith, G. D., Woods, E., Baer, T., and Miller, R.E.: Aerosol Uptake Described by Numerical Solution of the Diffusion -Reaction Equations in the Particle, J. Phys. Chem. A, 107, 9582–87, 2003.
 - Starr, F. W., Sciortino, F., and Stanley, H. E.: Dynamics of Simulated Water under Pressure, Phys. Rev. E, 60(6), 6757–68, 1999.

Thomas, N.L., and Windle, A.H.: A Theory of Case II Diffusion, Polymer, 23, 529-542, 1982.

15 Vaden, T. D., Imre, D., Beranek, J., Shrivastava, M., and Zelenyuk, A.: Evaporation Kinetics and Phase of Laboratory and Ambient Secondary Organic Aerosol, P. Natl. Acad. Sci. USA, 108(6), 2190–95, 2011.

Vignes, A.: Diffusion in Binary Solutions, Industrial & Engineering Chemistry Fundamentals 5(2), 189-99, 1966.

Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An Amorphous Solid State of Biogenic Secondary Organic Aerosol Particles, Nature, 467(7317), 824–27, 2010.

Wesselingh, J. A., and Bollen, A. M.: Multicomponent Diffusivities from the Free Volume Theory, T. I. Chem. Eng.-Lond., 75(A), 590–602, 1997.

Zaveri, R.A., Easter, R. C., Fast, J.D., and Peters, L.K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), J. Geophys. Res., 113(D13204), doi:10.1029/2007JD008782, 2008.

Zaveri, R. A., Easter, R. C., Shilling, J. E., and Seinfeld J. H.: Modeling Kinetic Partitioning of Secondary Organic Aerosol and Size Distribution Dynamics: Representing Effects of Volatility, Phase State, and Particle-Phase Reaction, Atmos. Chem. Phys., 14, 5153–81, 2014.

5

20

25

Zardini, A. A., Sjogren, S., Marcolli, C., Krieger, U. K., Gysel, M., Weingartner, E., Baltensperger, U., and Peter, T.: A Combined Particle trap/HTDMA Hygroscopicity Study of Mixed Inorganic/organic Aerosol Particles., Atmos. Chem. Phys., 8, 5589–5601, 2008.

Zobrist, B., Soonsin, V., Luo, B. P., Krieger, U. K., Marcolli, C., Peter, T., and Koop, T.: Ultra-Slow Water Diffusion in Aqueous Sucrose Glasses, Phys. Chem. Chem. Phys., 13(8), 3514–26, 2011.

Table A1. Absolute tolerances used in KM-GAP whilst the self-diffusion coefficient of the semi-volatile component was held constant at 2.0×10^{-9} m²s⁻¹ and the saturation ratio of the semi-volatile component was increased and decreased from 1-90% and 90-1%. The absolute tolerance required for stability depended on the self-diffusion coefficient of the non-volatile component (D_{nv}^0) and the initial particle diameter ($D_{p,t=0}$).

	$D_{\mathrm{p,t=0}}$ (m)				
$D_{nv}^0 (\mathrm{m}^2 \mathrm{s}^{-1})$	1.0x10 ⁻⁵	1.0x10 ⁻⁶	1.0×10^{-7}	1.0x10 ⁻⁸	
$1.0 \times 10^{-8} - 1.0 \times 10^{-14}$	1.0×10^{-12}	1.0×10^{-13}	$1.0 \mathrm{x} 10^{-14}$	1.0×10^{-15}	
1.0x10 ⁻¹⁶	1.0×10^{-13}	$1.0 \mathrm{x} 10^{-14}$	1.0×10^{-15}	1.0×10^{-16}	
1.0x10 ⁻¹⁸	$1.0 \mathrm{x} 10^{-14}$	1.0×10^{-15}	1.0×10^{-16}	1.0×10^{-17}	
1.0x10 ⁻²⁰	1.0×10^{-15}	$1.0 \mathrm{x} 10^{-16}$	1.0×10^{-17}	$1.0 x 10^{-18}$	
1.0x10 ⁻²²	$1.0 \mathrm{x} 10^{-16}$	1.0×10^{-17}	$1.0 x 10^{-18}$	1.0×10^{-19}	
1.0x10 ⁻²⁴	1.0×10^{-17}	$1.0 x 10^{-18}$	1.0×10^{-19}	1.0×10^{-20}	
1.0×10^{-26}	1.0×10^{-18}	1.0x10 ⁻¹⁹	1.0×10^{-20}	1.0x10 ⁻²¹	

Table A2. Number of shells used in each model for each change in the vapour-phase saturation ratio of the semi-volatile component (Δe_s) and for different values of non-volatile component self-diffusion coefficient (D_{nv}^0).

	Δe_s							
$D_{nv}^0 (\mathrm{m}^2 \mathrm{s}^{-1})$	1-90%	90-1%	60-80%	80-60%	10-20%	20-10%		
	ETH model # shells							
1.0x10 ⁻⁸ - 1.0x10 ⁻¹⁴	40	40	40	40	40	40		
1.0x10 ⁻¹⁶ - 1.0x10 ⁻²⁶	300	300	40	40	40	40		
	KM-GAP #	KM-GAP # shells						
1.0x10 ⁻⁸ - 1.0x10 ⁻¹²	40	40	40	40	40	40		

1.0×10^{-14}	60	60	40	40	40	40	
1.0x10 ⁻¹⁶	100	100	40	40	40	40	
1.0x10 ⁻¹⁸	200	200	40	40	40	40	
1.0×10^{-20}	250	250	100	100	40	40	
1.0x10 ⁻²²	270	270	100	100	40	40	
1.0x10 ⁻²⁴	300	300	100	100	40	40	
1.0×10^{-26}	330	330	100	100	40	40	
	Fi-PaD # shells						
1.0x10 ⁻⁸ - 1.0x10 ⁻¹⁴	40	40	40	40	40	40	
1.0x10 ⁻¹⁶ - 1.0x10 ⁻²⁶	300	300	80	80	40	40	



Figure 1. a) a schematic of a particle split into shells, as used in the diffusion models, with shell boundaries represented by lines, and symbols relating to those used in the model equations (α_i , X_i and e_s are the accommodation coefficient, particle-

- 5 phase mole fraction and vapour-phase saturation ratio of the semi-volatile, respectively). The relative width of the surface shell is shown larger than that used in models for clarity. b) and c): the concentration-radius profiles estimated using the ETH model at various times during diffusion for a semi-volatile component diffusing inwards with an initial mole fraction of 0.01 and equilibrium mole fraction of 0.90 (instantaneous change in saturation ratio of 1-90% assuming ideality), where t_e is the *e*-folding time. Note that the axes are relative, and normalised by the total radius (R_p) of the particle on the abscissa and
- by the equilibrium concentration ($[i]_{eq}$) on the ordinate. b) is for the diffusion coefficient independent of composition and c) is for the diffusion coefficient with a logarithmic dependence on mole fraction ($D_{nv}^0 = 1 \times 10^{-25}$, $D_{sv}^0 = 2 \times 10^{-9}$ m²s⁻¹).





Figure 2. Example dependencies of D_i on the mole fraction of the semi-volatile component. For the constant case both components have a value of $2x10^{-9}$ m²s⁻¹ and for the other cases the self-diffusion coefficient of the semi-volatile component is set to $2x10^{-9}$ m²s⁻¹ and that of the non-volatile is $1x10^{-25}$ m²s⁻¹.



Figure 3. *e*-folding time contour plots for different changes to the saturation ratio of the semi-volatile component (Δe_s) and different diffusion coefficient dependencies: a) Δe_s = 1-90% logarithmic dependence, b) Δe_s = 90-1%, logarithmic dependence, c) Δe_s = 1-90%, sigmoidal dependence, and d) Δe_s = 90-1% sigmoidal dependence. D⁰_{nv} is the diffusion coefficient at a semi-volatile mole fraction of 0, while D⁰_{sv} (diffusion coefficient at a semi-volatile mole fraction of 1) was fixed at 2.0x10⁻⁹ m²s⁻¹.



Figure 4. *e*-folding time contour plots for different changes to the saturation ratio of the semi-volatile component (Δe_s) and different diffusion coefficient dependencies: a) $\Delta e_s = 10{\text -}20\%$ logarithmic dependence, b) $\Delta e_s = 20{\text -}10\%$, logarithmic dependence, c) $\Delta e_s = 10{\text -}20\%$, sigmoidal dependence, and d) $\Delta e_s = 20{\text -}10\%$ sigmoidal dependence. D_{nv}^0 is the diffusion coefficient at a semi-volatile mole fraction of 0, while D_{sv}^0 (diffusion coefficient at a semi-volatile mole fraction of 1) was fixed at $2.0 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$.



Figure 5. The concentration ([*sv*])-radius (*r*) profiles of the semi-volatile component at times (t) preceding and including the *e*-folding time (t_e) as estimated by the three models given in the legend, for the case of logarithmic dependence of D_i on x_{sv} , an instantaneous increase in e_s from 1-90% and with $D_{nv}^0 = 1 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$ and $D_{sv}^0 = 2 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$. Note that for clarity the radius axis begins at 2×10^{-6} m and not 0 m.



Figure 6. The radius (R_p) change with time for a single particle subject to the changes in e_s shown by the orange curve and right vertical axis. Here $D_{sv}^0 = 2x10^{-9} \text{ m}^2 \text{s}^{-1}$ and $D_{nv}^0 = 1x10^{-25} \text{ m}^2 \text{s}^{-1}$ using the D_i dependencies given in the legend.



Figure 7. a) Radius (R_p) change with time for a single particle experiencing the changes in saturation ratio of the semivolatile component (e_s) shown by the orange curve (allied with the right vertical axis), for $D_{sv}^0 = 2x10^{-9}$ m²s⁻¹ and $D_{nv}^0 = 1x10^{-25}$ m²s⁻¹ and using the D_i dependencies given in the legend. b), c), d), e): time intervals of a) over select changes in e_s (as shown by their orange curve and right vertical axes).



Figure A1. The water mole fraction as a function of time and distance through a single particle using the ETH model and the same inputs as for Fig. 3 of Zobrist et al. (2011).



Figure A2. The *e*-folding time convergence with increasing shell number for the ETH model: change in the semi-volatile component saturation ratio in a) was 1-90% and in b) was 90-1%. The self-diffusion coefficient of the semi-volatile was $2.0 \times 10^{-9} \text{ m}^2 \text{s}^{-1}$ and that of the non-volatile was $1.0 \times 10^{-25} \text{ m}^2 \text{s}^{-1}$. Orange curves are the exponential best fits.



Figure A3. *e*-folding times (isolines) for the three models given in the legend. Where D_{con}^0 is the constant diffusion coefficient used throughout the simulation. $D_{p,t=0}$ is the initial particle diameter. In a) the saturation ratio of the semi-volatile is increased from 1-90% instantaneously, whilst in b) it is decreased from 90-1% instantaneously.



Figure A4. *e*-folding time contour plots for different instantaneous changes in the saturation ratio of the semi-volatile component (Δe_s) and different diffusion coefficient dependencies: a) $\Delta e_s = 60-80\%$ and logarithmic dependence, b) $\Delta e_s = 80-60\%$ and logarithmic dependence, c) $\Delta e_s = 60-80\%$ and sigmoidal dependence, and d) $\Delta e_s = 80-60\%$ and sigmoidal dependence. Models are given in the legend.