FINAL RESPONSE

The replies to all referee comments can be found below together with the revised manuscript. The changes made in the manuscript are marked in yellow. Note that in addition to the changes mentioned in the replies, Fig. 3 was slightly changed. This did not, however, change the results significantly.

Reply to Referee #1

We thank Anonymous Referee #1 for their helpful comments. We have answered to the comments below. The bold text is quoted from the referee's comments, and the text in italics has been added to the manuscript.

GENERAL COMMENTS

The paper explores the differences in estimated growth rates for sub-3nm particles by three different methods. The work seems solid and thorough, and it is of publishable quality. However, I'm unsure if the interest is broad enough for ACP as opposed to e.g. Aerosol Science & Technology. I'd recommend either GMD or AMT, though the paper is a modelling study that is of interest to observationalists, so it sits between those journals. I'll leave it up to the editor to make a judgement call. If the paper is to be published in ACP, I have several minor comments that should be addressed.

We believe that our paper fits into the scope of ACP (as also agreed by Referee #2) and is relevant for the readership of the journal, as the methods discussed in the paper are commonly used for analyzing experimental data on new particle formation both in the atmosphere and in chamber experiments. Experimentally determined growth rates are used to characterize particle formation and growth events, and, thus, they are routinely utilized in the analysis of particle distribution data (e.g. Kulmala et al., 2012). Moreover, growth rates extracted from measurements are often compared to those determined by theoretical or modeling approaches. Therefore, assessing the differences between different growth rate definitions is useful also for modelers, and, in general, for all researchers working with atmospheric particle data.

SPECIFIC COMMENTS

P2 L11: "CGR" doesn't correspond to the previous text. Can you define it? Is it "condensational growth rate"?

Yes, in this paper CGR is used as the abbreviation for the growth rate determined based on irreversible vapor condensation, and the letters stand for condensational growth rate. This has now been added to the text.

P3 L6-7: "AGR" and "FGR", same as above.

AGR is used as the abbreviation for the growth rate obtained with the appearance time method, and the letters stand for appearance time growth rate. FGR is used as the abbreviation for the growth rate corresponding to the molecular fluxes, i.e. flux-equivalent growth rate. These are now clarified in the text.

P4 L7-8 and throughout: I'm somewhat confused by the different sinks in this paper. Is the loss due to coagulation with larger particles included here? Is the loss due to coagulation between the model-resolved clusters included elsewhere when discussing the "loss coefficient"?

We agree that the processes included in the model are not explained very clearly. The model includes all the possible collisions between the clusters (up to the clusters consisting of 70 molecules), and

also an additional, time-independent external sink. The external sink can be thought to represent the loss of the clusters due to the coagulation onto pre-existing larger particles in the atmosphere. The losses discussed in the paper always refer to the external sink, characterized by the loss coefficient of the vapor monomer (in 1/s). The loss frequencies of the clusters are somewhat lower than that of the monomer (see Eq. (8)). The role of coagulation between the modeled clusters in the cluster growth dynamics is characterized by the non-monomer fraction of the flux.

To clarify the processes included in the model, we now added the following sentences in the beginning of the Sect. 2.2 (page 6, line 16):

We simulated the time-evolution of cluster concentrations in a one-component system using the Atmospheric Cluster Dynamics Code (ACDC; McGrath et al., 2012; Olenius et al., 2014). The model included the production of monomers, all the possible collision and evaporation processes between different clusters, and the losses of clusters due to an external sink.

Eqn 1: What about the source by a collision of two clusters that are not in the previous bin, but where the sum of their molecules would put the resultant particle in the current bin?

It is true that Eq. (1) considers only the flux originating from the previous $\lim J_{i-1,i}$ although the collisions between clusters belonging to $\lim < i-1$ may also contribute to the flux into $\lim i$. This contribution is in practice notable only when the concentrations of large clusters are high, or the bin widths very narrow. In the simulations of this work, all possible collisions between clusters were considered but the studied conditions were selected so that the contribution of collisions from $\lim < i-1$ to the flux into $\lim i$ was negligible for all size classes *i*. We added the following remark in Sect. 2.1 (page 4, line 8):

In situations with high concentrations of large clusters, the overall flux into bin i may contain the contributions of fluxes from smaller bins < i-1. This makes the analysis of the dynamics more complex and such situations are not addressed in this study.

Sect 2.2: Would it have made sense to do a simulation set where you simulation every cluster size from 1-70, so you don't have any numerical issues? This would be a useful comparison.

Actually, every cluster size from 1 to 70 was simulated. Only when calculating the growth rates and fluxes, the clusters were grouped into the size bins. This was done as the growth rate of a single cluster is ambiguous when clusters grow not only by monomer additions but also by cluster-cluster collisions. To make this clear, we modified a sentence in the Sect. 2.2 (page 7, line 4):

After simulating the time-evolution of the discrete cluster concentrations, the clusters were grouped into size bins containing an equal number of cluster (in most cases ten), for which fluxes and growth rates were determined.

Furthermore, we want to point out that the previous study by Olenius et al. (2014) discusses the case where clusters grow only by monomer attachments, and thus the growth rates can be unambiguously determined with the resolution of one molecule.

P7 L4: Why is the concentration in units cm-3 s-1?

This was a typographical error that has now been corrected. The correct unit is cm⁻³.

P7 L30: What values of monomer sources were used earlier (I think only steady state concentrations were given)?

The monomer source rates used in the different simulations are now presented in Table 1.

Reply to Referee #2

We thank Anonymous Referee #2 for their helpful comments. We have answered to the comments below. The bold text is quoted from the referee's comments, and the text in italics has been added to the manuscript.

GENERAL COMMENTS

In their manuscript, Kontkanen et al. present a set of numerical simulations of a one-component system, with the purpose to study the effect of cluster-cluster collisions on the growth of particles, and also the impact of different definitions and methods of the growth rate. The study is interesting, and gives a good overview of the different ways of looking at fresh particle growth, and the author's approach brings interesting new viewpoints to recent studies of nanoparticle formation in the atmosphere. The work is quite theoretical, and no comparison to actual experiments is made (as the studied one-component system would be quite difficult to produce). I think that the author's work is useful and interesting, and also in the scope of ACP. In my opinion the manuscript could be published in ACP, but I would appreciate it if some more information on the simulations and their analysis could be provided. I have listed my concerns below.

* I did not fully follow how the simulations were performed. Were all inputs kept constant over the whole time? The authors state that there was an initial cluster distribution corresponding to a low monomer concentration. I addition, it is stated that a 'steady-state monomer concentration' is reached at some point. This suggest that the aerosol formation and growth proceeds until some kind of a steady state is reached. If it is so, this has at least the following implications for the data analysis:

The model inputs, i.e. the monomer source rate, the external loss, temperature and the vapor properties, were kept constant during one simulation. The initial cluster distribution, corresponding to the monomer concentration of 5×10^5 cm⁻³, was used so that the simulations would correspond to the real situation in, for example, the atmosphere or a chamber experiment where the concentration of condensing vapor (e.g. sulfuric acid) is higher than zero already before the actual new particle formation event. As concluded by the referee, the simulations were continued until a steady state was reached.

a) from which part of the process are the various size-dependent growth rates determined? For AGR, the GR is necessarily from the start, but for the other GRs some other times could be used. Was the steady-state situation used? What was the criterion for steady state? Were the different growth rates determined at different time points within one simulation? Were the growth rates time-dependent within one simulation?

AGR was obtained by differentiating (t_{app} , D_p)-data, where the appearance time of a size bin, t_{app} , was defined as the time at which the concentration of the bin reaches 50% of the total increase in the concentration (see Sect. 2.1.2). AGR thus yields one time-independent value for each size bin in one simulation. FGR and CGR, on the other hand, were time-dependent in the simulations, as FGR depends on the time evolution of the fluxes, and CGR on the time evolution of the monomer concentration. Therefore, both FGR and CGR changed during one simulation until the steady state was reached. We determined FGR and CGR at the mean appearance times of consecutive size bins, and at the steady state. This is already mentioned on page 8, although we now slightly modified the sentence to be more exact. As the main aim of this study was to compare the different growth rates with each other, only the values obtained at the appearance times were presented in the paper. However, the steady-state results were qualitatively similar. The criterion for the steady state was that

the relative changes in the concentrations of the monomer and clusters as a function of time were negligible (less than 10^{-4} %) when the integration was continued for several minutes.

b) A 'steady state' in which everything stays constant is rarely seen in nature, partly because the growing particles contribute to a growth in the sink and therefore the Q/S fraction also changes. This could, for example, introduce a coupling between the formation rate and the AGR of FGR, because high formation rate causes a fast-increasing sink quickly to the distribution. I understand that the sink in the simulations was not dynamic, and therefore this effect is omitted. Do the authors have an idea how this affects the results? In any case, it would be useful to have a description of the time-dependent developing distribution, for example using a figure. The same figure could then be used to give a graphical explanation of the derivation of the different growth rates.

The external sink, corresponding to the loss of clusters due to coagulation onto large particles in the atmosphere, was indeed kept constant in our simulations. The sink of the vapor monomer and small clusters due to collisions with the simulated larger clusters (up to the clusters containing 70 molecules), naturally increased during the simulation when the cluster concentrations were increasing. On the other hand, the clusters growing beyond the size of 70 molecules were removed from the simulation, and therefore did not contribute to the sink. This means that in our simulations the sink remained lower than if the sink due to the particles larger than 70 molecules was taken into account, and thus the concentrations of vapor monomer and clusters could increase higher. However, as we are here interested only in the beginning of a new particle formation event (when sub-3nm clusters are formed), this effect is likely not very big, as it takes several hours before the nanometer sized clusters can reach sizes large enough to contribute significantly to the total sink. Thus, the coupling between the particle formation and the sink is likely to become significant mainly in very rapid particle formation bursts, where both the initial formation and the growth occur fast. Following the referee's suggestion, we now added a figure showing the time evolution of the cluster distribution in one simulation into the Appendix (Fig. A2 in the revised manuscript).

* A big question in particle formation research is the size dependence of growth in particle formation. From the results in this paper (Figs 1d, 2d, 3d, and 4d), it seems that for this one-component case, AGR and CGR tend to decrease with size while FGR increases with size. Is this a result that could be considered general? Is it possible that this changes in a true, multi-component case?

CGR decreases in all cases with the increasing cluster diameter, which results directly from its definition (see Eq. (7)). The size dependencies of AGR and FGR, on the other hand, are not affected only by the substance properties, but also by the ambient conditions, as well as the size classification, i.e. the resolution of the analysis (see Eqs (3) and (4)). The results for a multi-component system with an evaporation profile similar to our one-component substance can be expected to qualitatively resemble the results presented here. However, the size-dependencies of AGR and FGR observed in our simulations cannot be generalized for arbitrary substances and environments, as they are affected by the evaporation profile of the substance, and the time evolution of the cluster distribution, determined by the time evolution of the sources and sinks. For example, in the atmosphere the monomer source rate is usually time-dependent, which may affect the size-dependence of the growth rates (see Olenius et al. (2014) where this case is discussed).

To clarify this, we added the following sentence in the Sect. 3.1 (page 9, line 24):

It needs to be noted that the size-dependencies of AGR and FGR observed here cannot be generalized for arbitrary substances and environments, because they are affected by the vapor properties as well as the ambient conditions.

* Finally, I think that the discussion in the conclusions could be improved. In the last paragraph, (p. 14, L 23) the authors state that "conventional methods used to determine particle formation rates from growth rates may give estimates far from their true values". Based on earlier discussion, it seems that the authors bring forward the idea that not a single 'true' value exists (for example, the growth rate can mean the AGR, FGR or CGR). In the study of particle formation, finding the values for the formation rate J and the growth rate GR has for some time been a key objective. It would be useful if the authors could discuss here which of the given growth rates is the true value meant.

We agree that the sentence quoted by the referee was not entirely clear, and it has now been slightly changed. The aim of the statement is not to suggest that any of these growth rates would be the true growth rate of particles. Indeed, in the cases discussed in this study, where clusters grow not only by monomer collisions but also by cluster-cluster collisions, it is not possible to give one, unambiguous value for the growth rate of clusters of a specific size. With the "true value", we refer to the true value of the particle formation rate J, i.e. the flux of particles past a certain size, that is always unambiguously defined. We want to point out that determining a formation rate from a measured growth rate (here AGR) utilizing Eq. (5), as is often done, may lead to values far from the real particle flux. This issue is discussed, for example, in the end of the Sect. 3.1.

SPECIFIC COMMENTS

P7 L33: Regarding the monomer concentration ranges, I understand the limitation of the lower level. However, the upper level for the monomer concentration is less justified. This is a case of selecting a too long time step; reducing the time step in the differential equation solving would present usable results. In any case, the limits for the usability of the methods should be given.

Actually, the upper limit for the monomer concentration is not determined by the time step of the simulations, but the width of the size bins, for which the growth rates are determined. As clarified in the end of the Sect. 2.2 (page 8, line 2), if the monomer concentration was set to a too high value, concentrations of large clusters become so high that a significant fraction of the flux from a certain size bin may end up not only in the next size bin but also in the size bins larger than that. This would complicate the growth dynamics, and the method used to calculate the flux-equivalent growth rate (see Sect. 2.1.1) would not be exactly valid (see also the reply to a comment on Eq. (1) by Referee #1). Also, determining AGR from the appearance times of two consecutive bins (Eq. (4)) would not be justified if significant fractions of the flux from bin *i* would end up also in bins > i+1. As we are interested in situations where the clusters grow mainly by collisions of vapor monomer and small clusters, we have not addressed cases where the self-coagulation of the population becomes significant. We have now added a reference to Table 1 when discussing the limitations of the methods (page 8). In general, the limits for the validity of these methods are determined by the combination of all the parameters in the simulations (e.g. monomer source rate, the magnitude of the external sink, the properties of model substance) and the size bin width used in the analysis, and are thus systemspecific.

P19, Table 1: For reproducibility, it is not enough to give the final monomer concentration, but also the monomer source rate is important information. I think that the monomer source rates used to obtain the final concentrations in Table 1 should be given also in Table 1.

As the referee suggests, we now added the monomer source rates used in different simulations into Table 1.

Growth of atmospheric clusters involving cluster-cluster collisions: comparison of different growth rate methods

J. Kontkanen¹, T. Olenius², K. Lehtipalo^{1,3}, H. Vehkamäki¹, M. Kulmala¹ and K. E. J. Lehtinen⁴

¹Department of Physics, University of Helsinki, 00014 Helsinki, Finland

²Department of Environmental Science and Analytical Chemistry (ACES) and Bolin Centre for Climate Research, Stockholm University, 10691 Stockholm, Sweden

³Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen, Switzerland

⁴Finnish Meteorological Institute and Department of Applied Physics, University of Eastern Finland, 70211 Kuopio, Finland

Correspondence to: J. Kontkanen (jenni.kontkanen@helsinki.fi)

- 10 Abstract. We simulated the time evolution of atmospheric cluster concentrations in a one-component system where clusters grow not only by condensation of monomers, but where cluster-cluster collisions also significantly contribute to the growth of the clusters. Our aims were to investigate the consistency of the growth rates of sub-3 nm clusters determined with different methods, and the validity of the common approach to use them to estimate particle formation rates. We compared the growth rate corresponding to particle fluxes (FGR), the growth rate derived from the appearance times of clusters (AGR) and the
- 15 growth rate calculated based on irreversible vapor condensation (CGR). We found that the relation between the different growth rates depends strongly on the external conditions and the properties of the model substance. The difference between the different growth rates was typically highest at the smallest, sub-2nm sizes. FGR was generally lower than AGR and CGR; at the smallest sizes the difference was often very large, while at sizes larger than 2 nm, the growth rates were closer to each other. AGR and CGR were in most cases close to each other at all sizes. The difference between the growth rates was generally
- 20 lower in conditions where cluster concentrations were high, and evaporation and other losses thus less significant. Furthermore, our results show that the conventional method used to determine particle formation rates from growth rates may give estimates far from the true values. Thus, care must be taken not only in how the growth rate is determined but also in how it is applied.

1 Introduction

5

Atmospheric new particle formation has been observed to occur frequently in various environments around the world (Kulmala et al., 2004). The process has been estimated to significantly contribute to the global concentrations of cloud condensation nuclei (CCN), and thus affect the Earth's climate (Spracklen et al. 2008; Merikanto et al., 2009). The primary quantity characterizing new particle formation events is the particle formation rate, which is defined, for any size, as the flux of particles growing past that size (Kulmala et al., 2004). For determining this flux, the particle growth rate (GR) is commonly used (Kulmala et al., 2012).

- 30 With respect to analyzing and quantifying new particle formation events, GR has had several different interpretations and uses. Theoretically, GR for a given particle is straightforward to define: it is the rate at which the particle diameter changes at a given moment in time. However, as this growth is caused by random collisions of vapor molecules, GR can vary a lot in time and from particle to particle. Especially, all particles of the same size and chemical composition do not grow at exactly the same rate, as is inherently assumed in, for example, the condensational growth term in the standard version of the continuous
- 35 aerosol general dynamic equation (GDE; e.g. Seinfeld and Pandis, 2006). Still, a mean size-dependent value can be derived for GR, resulting in the well-known expressions for the free-molecular and continuum regimes of condensational growth, as well as various interpolations for the transition regime (see e.g. Seinfeld and Pandis, 2006). These expressions have been used and are convenient when trying to estimate vapor concentrations from observed GR or vice versa (e.g. Dal Maso et al, 2005; Nieminen et al., 2010). In this article, such a growth rate is called the growth rate based on irreversible vapor condensation,
- 40 abbreviated as CGR (condensational growth rate). Another important use of GR lies in relating it to the dynamics of the

evolving size distribution as the population of particles undergoes condensational growth. It is used in this context especially when estimating so-called survival rates, i.e. the fraction of particles that are not scavenged by background particles, instrument walls or other sinks while growing to a certain size. In this case, it is natural to define GR at a specific diameter by the flux *J* of particles growing past the given diameter (e.g. Wang et al., 2013; Olenius et al., 2014). In this article, such a growth rate is

5 called the flux-equivalent growth rate FGR.

To study the first steps of new particle formation, the growth rates of small, sub-3nm particles have been deduced from experimental data using various methods. The CGR method has been applied to specific measurement conditions by using the observed concentrations of precursor vapors in the calculation (Nieminen et al., 2010). The growth rates of charged particles have been derived from ion spectrometer data by following the time evolution of the concentration maximum (Hirsikko et al.,

- 10 2005; Manninen et al., 2009; Yli-Juuti et al., 2011). Measuring sub-3 nm electrically neutral particles is challenging, and therefore their growth rates have been indirectly deduced from the time lag between the rise in sulfuric acid concentration and the increase in the concentration of 3 nm particles (Weber et al., 1997; Sihto et al., 2006). However, recent instrumental development has enabled the detection of neutral clusters with mobility diameters of down to ~1 nm by using instruments such as the DEG-SMPS (Jiang et al., 2011) and the PSM (Particle Size Magnifier; Vanhanen et al., 2011). Consequently, the growth
- 15 rates in the sub-3 nm size range have been assessed, for example, by fitting the size distributions measured with a DEG-SMPS to the GDE (Kuang et al., 2012), or by determining the times at which the onset of new particle formation is detected with CPCs with different cut-off sizes (Riccobono et al., 2012). Recently, the growth rates of sub-3nm particles and molecular clusters have been determined from the appearance times of clusters in different size bins measured by scanning the PSM (Kulmala et al., 2013; Lehtipalo et al., 2014) or from the appearance times of specific clusters detected by mass spectrometers
- 20 (Lehtipalo et al. 2014; 2016). The appearance time has generally been defined as the time at which the concentration in the size bin, or the signal intensity of the cluster, reaches 50% of its total increase. In this article, the growth rate derived from the appearance times is referred to as AGR (appearance time growth rate).

To investigate the validity of the appearance time method, Lehtipalo et al. (2014) applied the method to particle size distribution data simulated with an aerosol dynamics model. They found that the growth rates determined from the appearance times were

- 25 close to the average condensational growth rates used as input in the simulation. On the other hand, Olenius et al. (2014) took a different approach to assess the AGR method by using a cluster kinetics model that does not inherently assume any growth rates but simulates the evolution of the cluster population via discrete collisions and evaporations of molecules. They compared the growth rates obtained with the appearance time method (AGR) to the growth rates corresponding to the molecular fluxes (FGR) and concluded that AGR was higher than FGR in the studied conditions. The difference was largest for the smallest
- 30 clusters and was often strongly affected by the ambient conditions. Although Olenius et al. (2014) showed that AGR and FGR may not be equal, they concentrated on an ideal situation where the growth proceeds only by monomer collisions and evaporations. In reality, there are situations where collisions between two clusters may contribute significantly to the growth (Lehtipalo et al., 2016) and they should, therefore, be taken into account when calculating the flux-equivalent growth rate. Furthermore, Olenius et al. (2014) used a resolution of a single molecule in their analysis, which is not possible when analyzing
- 35 experimental particle size distributions.

As GR can been interpreted and determined from experimental data in many different ways, it is essential to compare the results obtained with different methods. In this study, we compare the three above mentioned growth rate definitions, FGR, CGR and AGR, by applying them to modelled particle size distribution data. We use the same dynamic model as Olenius et al. (2014) to simulate the time evolution of cluster concentrations in a one-component system. As opposed to the simulations

40 done by Olenius et al. (2014), in our model system a significant part of clusters' growth proceeds via collisions of small clusters in addition to monomer attachments. Because the growth rate of a single cluster is ambiguous in this case, we group the clusters into size bins for which we calculate the growth rates. This also makes our analysis resemble the analysis of measured particle size distributions with a size resolution poorer than one molecule. We estimate AGR as in Lehtipalo et al. (2014), FGR analogously with Olenius et al. (2014), and CGR directly from the vapor monomer concentration (Nieminen et al., 2010). Our aim is to answer the following questions: 1) How important are cluster-cluster collisions for the growth of the cluster population in different conditions in our model system? 2) How consistent with each other are the flux-equivalent growth rate,

- 5 the growth rate derived from the appearance times of the clusters, and the growth rate calculated based on irreversible vapor condensation? 3) How valid is the conventional method used to estimate particle formation rates from growth rates? We examine these questions in different conditions by varying the saturation vapor pressure of the vapor, the vapor source rate and the magnitude of an external sink reducing the vapor and clusters. The simulated conditions correspond to the typical conditions observed during new particle formation in, for example, a boreal forest. In most of the simulations the size-
- 10 dependent evaporation rate is set to decrease monotonically with increasing cluster size, corresponding to increasing cluster stability. However, we also test a different evaporation profile in order to study the effect of elevated concentrations of stable small clusters on the growth of the cluster population. Furthermore, we investigate how the size resolution, i.e. the width of the size bins, affects the results.

15 2 Methods

2.1 Determining the growth rates

For the growth rate analysis, the clusters were grouped into size bins so that each bin contains an equal number of cluster sizes, i.e. in linear volume space the bins are of equal width. The time evolution of the total cluster concentration C_i in a certain size bin *i* can be described with the following equation:

20
$$\frac{dC_i}{dt} = J_{i-1,i} - J_{i,i+1} - S_i,$$
 (1)

where $J_{i-1,i}$ is the flux coming to size bin *i* from the previous bin *i*-1, $J_{i,i+1}$ is the flux from bin *i* to bin *i*+1 and $S_i = \sum S_j$ is the total external sink for size bin *i*, summed over all cluster sizes *j* that belong to bin *i*. In situations with high concentrations of large clusters, the overall flux into bin *i* may contain the contributions of fluxes from smaller bins < *i*-1. This makes the analysis of the dynamics more complex and such situations are not addressed in this study.

25 Equation 1 can be obtained directly by integrating the continuous general dynamic equation (GDE; Friedlander, 1977) for aerosols, including only the growth and sink terms. If traditional continuous approach is used and clusters are assumed to grow synchronously by condensation, we can write

$$J_{i,i+1} = n \cdot \text{GR}\Big|_{\text{at the boundary between bins } i \text{ and } (i+1)}$$
(2)

and similarly for $J_{i-1,i}$ (Lehtinen et al., 2007). Here *n* is the number concentration distribution function dC/dd_p and GR is the 30 diameter growth rate of the clusters dD_p/dt .

In principle, it seems straightforward to combine Eqs (1) and (2) to obtain a method to determine growth rates from size distribution data. However, the possible contribution of larger clusters to growth, and the need to somehow approximate n and GR (at bin boundaries) complicate the task.

2.1.1 Flux-equivalent growth rate (FGR)

35 Here we follow the Eulerian approach used by Olenius et al. (2014) and referred to as the flux-equivalent growth rate (FGR). The method is based on defining GR by Eq. (2) even if the original underlying assumptions of Eq. (2) would not be valid. Furthermore, *n* is approximated with the value at the lower side of the bin boundary, resulting in

$$FGR_{i,i+1} = \frac{J_{i,i+1}}{n_i} = \Delta D_{p,i} \frac{J_{i,i+1}}{c_i},$$
(3)

where $\Delta D_{p,i}$ is the width of bin *i* in diameter space. These assumptions and approximations are generally made when treating experimental, size bin classified distribution data by using Eqs (1) and (2).

To obtain the net flux $J_{i,i+1}$ between size bins *i* and *i*+1, we first determine the fluxes between different cluster sizes with the following method: a) each collision between cluster sizes *p* and *q* grows *p* by the addition of the molecules in *q* if *p* > *q*,

- 5 contributing to the flux between p and p+q. For q, such a collision is treated as a sink. b) If the colliding clusters are of the same size (p = q), one of them is considered to grow and the other to be lost. In addition, to obtain the net flux between cluster sizes p and q, monomer evaporations and cluster fissions are taken into account. After calculating the net fluxes between each pair of cluster sizes, the net flux $J_{i,i+1}$ can be obtained by summing up the individual collision–evaporation fluxes crossing the boundary between the bins. Note that when using this method it must be ensured that the growth is dominated by collisions of
- small enough clusters, and the bins are wide enough, so that the growth from bin i to the bins larger than i+1 does not occur.

2.1.2 Growth rate from appearance times of clusters (AGR)

25

30

One possible way to assess growth rates from the time evolution of a particle distribution is based on the times at which concentrations in different size bins reach their maximum (Lehtinen and Kulmala, 2003). This is convenient for cases like nucleation bursts where there is a growing mode of particles. However, in cases where the system approaches a time-

15 independent steady-state this method obviously does not work. Here we investigate a method to obtain GR from appearance times of clusters (AGR) in the size bins, by defining the appearance time $t_{app,i}$ for the size bin with the mean diameter $D_{p,i}$ as the time at which the concentration of the bin reaches 50% of the total increase in the concentration in that size bin (Lehtipalo et al., 2014). Then, AGR is obtained from the numerical differentiation of the (t_{app} , D_p)-data:

$$AGR_{i,i+1} = \frac{D_{p,i+1} - D_{p,i}}{t_{app,i+1} - t_{app,i}}.$$
(4)

20 In recent experimental studies, AGR has been determined by applying a linear fit to the (t_{app}, D_p) -data over a range of several instrumental size classes. Because the aim of this work is to examine the particle flux and growth rate as a function of size, AGR is here determined separately for each size bin as in Eq. (4).

As one of the main reasons to estimate GR from particle size distribution dynamics is to estimate particle flux (or formation rate) J, we also compare the different methods in terms of fluxes. Thus, similarly to Eq. (3), we define a 'cluster appearance time flux' as follows:

$$J_{\text{app},i,i+1} = \frac{\text{AGR}_{i,i+1}}{\Delta D_{\text{p},i}} C_i$$
(5)

In the analysis of experimental particle size distribution data, Eq. (5) is generally used to calculate the particle flux towards larger sizes from measured or calculated growth rates (Kulmala et al., 2012). Thus, comparing the flux J_{app} to the real flux J, which is directly obtained from the simulations, provides information on how the use of Eq. (5) affects the conclusions of the data analysis.

In addition to the change in the mass diameter of the clusters, we determine FGR and AGR also as the change in the number of molecules, N_{mols} . These are obtained by replacing the width of the size bin $\Delta D_{\text{p},i}$ in diameter space in Eqs (3) and (4) with the width of the bin ΔN_{mols} in number of molecules space.

2.1.3 Growth rate by assuming irreversible vapor condensation (CGR)

35 The kinetic hard-spheres collision rate between a vapor molecule with diameter D_{mon} and a cluster with diameter D_p is given by

$$K_{\rm kin} = \frac{\pi}{4} (D_{\rm p} + D_{\rm mon})^2 (\bar{c}_{\rm p}^{\ 2} + \bar{c}_{\rm mon}^{\ 2})^{1/2} (C_{\rm mon} - C_{\rm e}), \tag{6}$$

where \bar{c} is the thermal speed, C_{mon} is the vapor concentration and C_{e} is the vapor concentration corresponding to the equilibrium vapor pressure over the cluster. When analyzing experimental data, it is usually assumed that saturation vapor

pressure $p_{sat} = 0$, which means that the vapor is assumed to condense irreversibly. In this case, the growth rate of a cluster as a change of mass diameter is obtained from (Nieminen et al., 2010):

$$CGR = \frac{\gamma}{2\rho} \left(1 + \frac{D_{\text{mon}}}{D_{\text{p}}}\right)^2 \left(\frac{3k_{\text{B}}T}{\pi}\right)^{1/2} \left(\frac{1}{m_{\text{p}}} + \frac{1}{m_{mon}}\right)^{1/2} m_{\text{mon}} C_{\text{mon}}.$$
(7)

Here ρ is the condensed phase density, and m_p and m_{mon} are the masses of the cluster and the vapor molecule. γ is a correction

5 factor that needs to be added if CGR is calculated in continuum regime (see Nieminen et al., 2010). When applying Eq. (7) in this study for calculating CGR for different size bins, we use the diameter and mass of the cluster at the upper limit of each size bin for D_p and m_p .

In addition to the fact that CGR calculated from Eq. (7) takes into account only monomer collisions and no evaporation, the essential difference between CGR and FGR is the perspective from which the growth is studied. CGR corresponds to the

10 traditional Lagrangian approach, where the growth of an individual cluster between different size bins is followed. FGR, on the other hand, corresponds to the Eulerian approach, where the net flux between adjacent size bins is studied. See Olenius et al. (2014) for further discussion about the differences between these approaches.

2.2 Simulations

We simulated the time evolution of cluster concentrations in a one-component system using the Atmospheric Cluster Dynamics

- 15 Code (ACDC; McGrath et al., 2012; Olenius et al., 2014). The model included the production of monomers, all the possible collision and evaporation processes between different clusters, and the losses of clusters due to an external sink. The model substance was assumed to consist of spherical molecules and clusters with the properties of sulfuric acid: a molecular mass (m_{mon}) of 98.08 amu, a liquid density (ρ) of 1830 kg m⁻³ and a surface tension (σ) of 0.05 N m⁻¹. However, the saturation vapor pressure of the model substance was lowered from that of sulfuric acid to decrease the evaporation rate of clusters. This
- 20 qualitatively mimics the stabilization of sulfuric acid clusters by base molecules, such as ammonia or amines (e.g. Kurtén et al., 2008). The simulations included clusters with 1 to 70 molecules; the clusters growing larger than that were assumed to be stable and removed from the simulation. The Gibbs free energy of formation of the clusters was calculated from the classical one-component liquid droplet model to obtain a qualitatively realistic evaporation profile. In the majority of the simulations the Gibbs free energy profile had a single maximum and no minima as a function of cluster size. This corresponds to a
- 25 monotonically increasing stability with the increasing cluster size in the studied size range. In addition, a set of simulations was performed using a free energy profile with lowered formation free energies for the smallest clusters (see Fig. A1 in Appendix). This corresponds to a system with elevated concentrations of small stable clusters, similarly as in the simulation study by Vehkamäki et al. (2012) and possibly also in the atmosphere (Kulmala et al., 2013). Collision rates between clusters were obtained from Eq. (6) with $C_e = 0$, and cluster evaporation rates were calculated from the Gibbs free energies of formation
- 30 of the clusters (e.g. Ortega et al., 2012). The external losses were assumed to depend on the cluster size according to (Lehtinen et al., 2007):

$$S(D_{\rm p}) = S(D_{\rm mon}) \times \left(\frac{D_{\rm p}}{D_{\rm mon}}\right)^b \tag{8}$$

Here D_{mon} is the diameter of a monomer. The exponent *b* was set to 1.6, in which case Eq. (8) corresponds to the typical sizedependency of losses caused by background aerosol particles in a boreal forest (Lehtinen et al., 2007). In all the simulations

35 the temperature was set to 278 K. The initial cluster concentrations were set to a steady-state distribution at a monomer concentration of 5×10^5 cm⁻³. After simulating the time evolution of the discrete cluster concentrations, the clusters were grouped into size bins containing an equal number of clusters (in most cases ten), for which fluxes and growth rates were determined.

A summary of the performed simulations is presented in Table 1. In the first four simulation sets, the Gibbs free energy profile 40 had one maximum and no minima. In the first simulation set, the effect of monomer concentration was studied: a constant source of monomer was assumed so that the final steady-state monomer concentration was 10⁶-10⁷ cm⁻³. These monomer concentrations are of the same order of magnitude as sulfuric acid concentrations observed during new particle formation in a boreal forest (Kulmala et al., 2013). The reference loss coefficient $S(D_{mon})$, describing the external sink, was set to 10^{-3} s⁻¹, which is of the order of magnitude of the loss of clusters onto pre-existing particles in a boreal forest (Dal Maso et al., 2005) or walls in a chamber experiment. A saturation vapor pressure of 2×10^{-10} Pa was used to study the situation where small clusters significantly contribute to the growth of the cluster population.

In the second simulation set, the aim was to investigate the effect of cluster stability on the growth of the cluster population by varying the saturation vapor pressure, to which the evaporation rates are directly proportional, from 1.5×10^{-10} Pa to 1×10^{-9} Pa. The final steady-state monomer concentration was set to 5×10^{6} cm⁻³.

In the third simulation set, the effect of the magnitude of the external sink was studied by setting the loss coefficient to 0.7 $\times 10^{-3}-2\times 10^{-3}$ s⁻¹. The monomer source rate was set to 5.5×10^3 cm⁻³ s⁻¹, which produces the steady-state monomer concentration of 5×10^6 cm⁻³ when the loss coefficient is 10^{-3} s⁻¹ and the saturation vapor pressure is 2×10^{-10} Pa.

In the fourth simulation set, we studied how the width of the size bins affects the growth rates by varying the size bin width from 5 to 14 clusters. Furthermore, as we wanted to compare our results directly with the results of Olenius et al. (2014) who used an ideal precision of one molecule in their simulations, we performed additional simulations with a cluster population

that grows only by monomer additions. In these simulations, we set the saturation vapor pressure of the model substance to 1×10^{-9} Pa, and allowed only monomer collisions and evaporations in our system.

The fifth simulation set was otherwise identical with the first simulation set, but the Gibbs free energy profile was different: a negative term of $90 \times (\exp(-(n_{mols} - 1)/4.2) - \exp(-(n_{mols} - 1)/4.5))$ was added to the classical expression for the free energy in order to decrease the formation free energies of the smallest clusters while keeping the free energies of larger clusters unchanged. The purpose of this simulation set was to see the effect of elevated concentrations of stabilized small clusters on

20 unchanged. The purpose of this simulation set was to see the effect of elevated concentrations of stabilized small clusters on the growth of the cluster population.
Finally, we also performed an additional set of simulations by varying the monomer source rate and the seturation varying

Finally, we also performed an additional set of simulations by varying the monomer source rate and the saturation vapor pressure simultaneously. The monomer source rate was varied between 1×10³ cm⁻³ s⁻¹ and 5×10⁴ cm⁻³ s⁻¹ and the saturation vapor pressure between 1×10⁻⁹ Pa and 1.5×10⁻¹⁰ Pa in different simulations. The loss coefficient was set to 10⁻³ s⁻¹. A Gibbs free energy profile containing a maximum and no minima was assumed.

- It should be noted that the studied ranges of different parameters, summarized in Table 1, were selected so that our analysis methods were valid in the simulated conditions. If the monomer concentration was set to a too low value, or the saturation vapor pressure and loss factor were too high, the concentration of clusters in the largest size bins would not increase in the simulation and determining growth rates would not be reasonable. On the other hand, if the monomer concentration was very
- 30 high, or the saturation vapor pressure and the loss coefficient very low, the concentrations of large clusters may become so high that a significant fraction of the flux from a certain size bin would end up not only in the next size bin but also in the size bins larger than that. In this case, the method that we use to calculate the flux-equivalent growth rate would not be valid.

3 Results and discussion

5

We determined the collision-evaporation fluxes between different size bins (J_{true}) and the fluxes calculated from the appearance 35 times (J_{app} ; see Eq. (5)) from all the simulations. We also calculated how large fraction of the flux J_{true} from each size bin is due to the collision and evaporation processes involving two clusters compared to the total flux including also monomer collisions and evaporations; hereinafter this is referred to as the non-monomer fraction of the flux. Then, we determined different growth rates (AGR, FGR and CGR) based on Eqs. (3), (4) and (7) for all the size bins. The growth rates were determined both with respect to the change in the number of molecules of cluster (denoted with the subscript N) and the change

40 in the cluster mass diameter (denoted with the subscript D). Therefore, the figures presenting the size dependency of the growth rates are shown using two different definitions for the bin size. The ratios of the different growth rates and the fluxes from

different size bins are shown as a function of the number of molecules, with the corresponding diameters presented on the upper *x*-axis. In all the figures, the fluxes originating from a certain size bin and the growth rates of that bin are plotted at the upper limit of the size bin. J_{true} , FGR and CGR were determined both at the mean appearance times of consecutive size bins and at the final steady state. The figures are, though, presented only for the appearance time case, the results for the steady

5 state being qualitatively similar. An example of the time evolution of the cluster distribution in one simulation is shown in Fig.
 A2 in Appendix.

Sections 3.1 and 3.2 discuss the results of the simulations where the free energy profile was assumed to have a single maximum and no minima, and the effects of the monomer concentration (3.1), the saturation vapor pressure, i.e. cluster stability (3.2), and the magnitude of the external sink (3.2) on the fluxes and growth rates were studied. Section 3.3 focuses on the effect of

10 the width of the size bins on the results. In Sect. 3.4 the simulations with a different free energy profile, leading to elevated concentrations of small stable clusters, are discussed. Finally, Sect. 3.5 presents the results of the simulations where the monomer source rate and the saturation vapor pressure were simultaneously varied.

3.1 Effect of monomer concentration

In the first simulation set, the steady-state monomer concentration was varied to see how it affects the growth of the cluster population. When the monomer source rate, and thus also the steady-state monomer concentration, increases, the non-monomer fraction of the flux becomes higher (Fig. 1a) as the relative number of clusters compared to monomers increases. At $C_{mon} = 10^6$ cm⁻³ and at $C_{mon} = 5 \times 10^6$ cm⁻³, the non-monomer fraction ranges from 2% to 9% and from 10% to 18%, and is highest in the smallest size bin. At $C_{mon} = 10^7$ cm⁻³, the non-monomer fraction varies between 28% and 40% and is highest in the size bin of 21–30-mers. The observed size-dependency of the non-monomer fraction of the flux is likely mainly due to the size-20 dependency of cluster concentrations and their losses.

- In Fig. 1b the true collision-evaporation fluxes from each size bin (J_{true} ; solid line) and the fluxes calculated from the appearance times (J_{app} ; dashed line) are presented. Both J_{true} and J_{app} increase with the increasing steady-state monomer concentration. This is due to higher cluster concentrations and, in the case of J_{app} , shorter time between the appearances of adjacent clusters (Δt_{app}). Furthermore, J_{app} and J_{true} decrease with increasing cluster size because of the decreasing cluster concentrations, with
- 25 the most prominent decrease observed for the lowest monomer concentration. At a low vapor concentration the relative role of the external sink becomes more significant, and therefore, the relative decrease in the cluster concentrations accumulates more strongly with the increasing cluster size.

Figures 1c and 1d present the different growth rates as a function of the number of molecules in the cluster and the cluster diameter. FGR is shown as solid lines, AGR as dashed lines, and CGR as dotted lines. All growth rates are generally higher

- 30 when the steady-state monomer concentration is higher. This is due to higher values of fluxes in the case of FGR and shorter time between the appearances of adjacent size bins (Δt_{app}) in the case of AGR. For CGR the dependency on the monomer concentration follows directly from Eq. (7). From Fig. 1c we can also see that FGR_N increases with the number of molecules in the cluster, which is caused by a relatively stronger decrease in the concentration of clusters with size compared to the decrease in the fluxes (see Eq. (3)). AGR_N also generally increases with the number of molecules in the cluster due to
- 35 decreasing Δt_{app} , but has a minimum in the size bin of 21–30-mers at the lowest monomer concentration. As explained by Olenius et al. (2014) this may be caused by the time evolution of the evaporation fluxes from large clusters to small clusters: these fluxes are lowest at the appearance times of the clusters in the smallest size bin, which may increase the AGR of the small sizes. CGR_N also increases with the number of molecules in the cluster. Furthermore, Fig. 1d shows that FGR_D increases with the cluster diameter, although the increase is very weak in the simulation with the lowest monomer concentration. On the
- 40 other hand, AGR_D decreases with the cluster diameter because the change in the diameter as a result of the addition of one molecule becomes smaller with the increasing cluster size. Finally, CGR_D decreases with the cluster diameter according to Eq.

(7). It needs to be noted that the size-dependencies of FGR and AGR observed here cannot be generalized for arbitrary substances and environments, because they are affected by the vapor properties as well as the ambient conditions.

We also studied the ratio of AGR to FGR (solid line in Fig. 1e), and the ratios of CGR to FGR (dotted line in Fig. 1e) and AGR to CGR (Fig 1f). AGR is higher than FGR at all sizes, their ratio depending strongly on the steady-state monomer concentration

- and the size bin. The AGR to FGR ratio generally increases with decreasing monomer concentration, reaching the highest values at $C_{\text{mon}} = 10^6 \text{ cm}^{-3}$. However, at the largest size bins the ratio is slightly lower at $C_{\text{mon}} = 5 \times 10^6 \text{ cm}^{-3}$ than at $C_{\text{mon}} = 10^7 \text{ cm}^{-3}$, which is due to the size-dependency of the external sink. The AGR to FGR ratio is highest at the smallest size bin (~10²– 10¹⁰ depending on the monomer concentration) and lowest at the largest size bin (~1.4–4). The CGR to FGR ratio behaves generally in a similar way as the AGR to FGR ratio, being slightly closer to one at the highest monomer concentration. Thus,
- 10 it seems that FGR of the smallest clusters can be significantly lower than AGR and CGR, especially at low monomer concentrations. This is caused by the fact that when calculating FGR, the flux from the size bin is divided by the mean value of the size distribution function in that bin ($C_i/\Delta D_{p,i}$ in Eq. (3)), while, theoretically, it should be the value at the bin boundary (see also Vuollekoski et al., 2012). This assumption affects the results most clearly in the smallest size bin, where the concentration decreases very fast as the function of the cluster size (see Fig A4 in Appendix) and the largest contribution to
- 15 the total concentration comes from the vapor monomer. For this same reason, using Eq. (5) to calculate J_{app} from AGR often results in too high values compared to the real particle flux. On the other hand, AGR and CGR are close to each other at all sizes; their ratio ranges from 0.8 to 4 at $C_{mon} = 10^6$ cm⁻³, from 0.7 to 1.0 at $C_{mon} = 5 \times 10^6$ cm⁻³, and from 0.9 to 1.8 at $C_{mon} = 10^7$ cm⁻³ (Fig. 1f). The similarity of AGR and CGR is rather surprising when considering the very different definitions of these growth rates (see Sect. 2.1), and the fact that AGR is affected by all possible collision and evaporation processes between the
- 20 clusters, while CGR is derived considering only the condensation of single molecules.

3.2 Effect of saturation vapor pressure and external sink

In the second simulations set the effect of cluster evaporation rate was studied by varying the saturation vapor pressure. When the saturation vapor pressure is lowered from 1×10^{-9} Pa to 1.5×10^{-10} Pa, the non-monomer fraction of the flux from the smallest size bin increases from 7% to 23% (Fig. 2a). In the largest size bin the non-monomer fraction varies between 2% and 15%.

- This shows that if the saturation vapor pressure is low, and therefore evaporation fluxes small, the cluster concentrations may rise so high that non-monomer collisions have a considerable effect on the growth of a cluster population. The collision-evaporation fluxes for all size bins (J_{true} ; solid line in Fig. 2b) are higher when the saturation vapor pressure is low, which is due to higher cluster concentrations. Similarly, the flux derived from appearance times (J_{app} ; dashed line in Fig. 2b) generally increases with decreasing saturation vapor pressure, which is due to higher concentrations and shorter time
- 30 between the appearance times of different size bins (Δt_{app}). The flux-equivalent growth rate FGR increases when saturation vapor pressure is lowered because of larger fluxes (Figs 2c and 2d). Except for the smallest size bin, AGR is also higher with the lower saturation vapor pressures due to shorter Δt_{app} . In the smallest size bin AGR is highest when the saturation vapor pressure is highest because the small clusters reach their appearance time faster in this case. CGR is also slightly higher when the saturation vapor pressure is lower. This may seem
- 35 illogical as CGR depends only on the monomer concentration, which is the same at the appearance time of the monomer in all the simulations. However, similar to FGR, CGR is determined at the mean appearance times of consecutive size bins, and not at the appearance time of the monomer. Thus, the differences in CGR are caused by differences in the appearance times of the size bins with varying saturation vapor pressures. Figure 2c also shows that FGR_N and CGR_N increase with the number of molecules in the cluster with all the saturation vapor pressures. AGR_N increases as a function of size with the lower saturation
- 40 vapor pressures of 2×10^{-10} Pa and 1.5×10^{-10} Pa, but has a minimum in the size bin of 21–30-mers when saturation vapor pressure is 1×10^{-9} Pa. This may result from the time development of the evaporation fluxes, as discussed in Sect. 3.1. FGR_D increases

with the cluster diameter, and AGR_D and CGR_D decrease with the cluster diameter, regardless of the saturation vapor pressure (Fig. 2d).

The ratios of AGR and FGR (solid line in Fig. 2e) and CGR and FGR (dotted line in Fig. 2e) depend strongly on the saturation vapor pressure and the size bin. Still, with all three saturation vapor pressures AGR and CGR are higher than FGR at all sizes.

- 5 In the smallest size bin, the AGR to FGR ratio varies between 10^3 and 10^9 increasing with the saturation vapor pressure. The CGR to FGR ratio behaves similarly as the AGR to FGR ratio, but it is slightly higher at the largest sizes when $p_{sat} = 1 \times 10^{-9}$ Pa. Altogether, FGR gives clearly lower growth rates than AGR and CGR for the smallest clusters, with the differences increasing when the saturation vapor pressure, and thus also evaporation fluxes, become larger. However, in the largest size bin AGR and CGR are close to FGR: the AGR to FGR ratio ranges from 1.4 to 1.7 and the CGR to FGR ratio from 1.3 to 2.7.
- Furthermore, AGR and CGR are close to each other with all the saturation vapor pressures (Fig. 2f). When $p_{sat} = 1 \times 10^{-9}$ Pa, the AGR to CGR ratio varies between 0.5 and 1.3, being highest in the smallest size bin. With $p_{sat} = 2 \times 10^{-10}$ Pa and $p_{sat} = 1.5 \times 10^{-10}$ Pa, the AGR to CGR ratio ranges from 0.7 to 1.0 and from 0.8 to 1.2, increasing with increasing cluster size. In the third simulation set, the effect of the external sink on the growth of cluster population was studied by varying the value

of the loss coefficient from 0.7×10^{-3} s⁻¹ to 2×10^{-3} s⁻¹. Lowering the loss coefficient seems to have similar effects on the results

- 15 as lowering the saturation vapor pressure (see Fig. A3 in Appendix). This results from the fact that in both cases the number of clusters increases, and therefore the cluster collisions become more important relative to evaporation or other losses. When the loss coefficient is lowered, the non-monomer fraction of the flux increases, J_{true} and J_{app} get higher values, and FGR and CGR increase, as expected. AGR also increases, except for the smallest size bin, where AGR is higher with a higher loss coefficient due to shorter Δt_{app} . The AGR to FGR ratio increases with the loss coefficient: for instance, in the smallest size bin
- 20 the AGR to FGR ratio ranges from 700 to 10⁶, and in the largest size bin the ratio is between 1.4 and 2.2. The AGR to CGR ratio, on the other hand, varies from 0.6 to 1.7 in the smallest size bin and from 1.0 to 1.2 in the largest size bin. The highest values of the ratio are obtained with the highest loss coefficient.

3.3 Effect of size resolution

In the fourth simulation set, the width of size bins was varied too see how the size resolution affects the growth rates. When

the size bins are wider, the non-monomer fraction of the flux at a certain size is higher (Fig. 3a). This is partly due to the size dependency of the non-monomer fraction, and partly due to the differences in the appearance times of bins with different widths, as the values are determined at the appearance times.

The collision-evaporation fluxes (J_{true} ; solid line in Fig. 3b) are not greatly affected by the size bin width as the flux from the size bin originates mostly from the largest clusters of that bin. Therefore, the small differences in J_{true} obtained with different

size bin widths are mainly caused by the differences in the appearance times of bins with different widths. On the other hand, the flux calculated from the appearance times (J_{app} ; dashed line in Fig. 3b) at a certain size becomes lower when the size bin width is decreased. This results from the decrease in the mean value of the size distribution function of the bin ($C_{i}/\Delta D_{p,i}$ in Eq. (5)) used for calculating J_{app} .

The flux-equivalent growth rate FGR at a certain size, on the other hand, increases when the bin width is decreased (Figs 3c and 3d), due to the lower mean value of the size distribution function of the bin ($C_i/\Delta D_{p,i}$ in Eq. (3)). Also, CGR becomes slightly higher when the bin width is decreased. Furthermore, AGR is also higher with narrower size bins as then the size bin width is relatively higher compared to Δt_{app} than with wider size bins (see Eq. (4)).

The relation of different growth rates to each other is also affected by the width of the size bins (Figs 3e and 3f). The AGR to FGR ratio gets higher values when the size bins are wider. In the smallest size bin the ratio is 10^3-10^4 depending on the bin

40 width, and in the largest size bin the ratio is correspondingly 1.2–1.7. The CGR to FGR ratio is slightly higher than the AGR to FGR ratio at small sizes and lower at large sizes. The ratio of AGR to CGR, on the other hand, slightly increases with decreasing bin size. In the smallest size bin the ratio is 0.6–1.0 depending on the bin width, while in the largest size bin the

ratio is ~1.0 in all cases. Altogether, high size resolution seems beneficial when using FGR to describe the growth of the cluster population, or when calculating particle fluxes from growth rates utilizing Eq. (5).

In order to compare our results directly with those of Olenius et al. (2014), we performed additional simulations with the saturation vapor pressure of 1×10^{-9} Pa and allowing only monomer collisions and evaporation in our system. In this case the

5 AGR to FGR ratios with different bin widths become higher compared to the simulations where cluster collisions contribute to the growth. The CGR to AGR ratio does not change as significantly.

3.4 Effect of stable small clusters

In the fifth simulation set a different cluster free energy profile was used to study the effect of elevated concentrations of stable small clusters on the growth of the population. The contribution of non-monomer collisions to the fluxes between different

- 10 size bins is significantly increased by the stabilization of small clusters (Fig. 4a, see also Fig. 1a for a comparison). In the smallest size bin the growth mainly proceeds by non-monomer collisions: the non-monomer fraction of the flux is 56–71% with different monomer concentrations. In the largest size bin, the non-monomer fraction depends strongly on the steady-state monomer concentration: the fraction is 15% with the lowest monomer concentration and 62% with the highest monomer concentration.
- The collision-evaporation fluxes (J_{true} ; solid line in Fig. 4b) and the fluxes derived from the appearance times of clusters (J_{app} ; dashed line in Fig. 4b) also increase in the presence of stabilized small clusters (see Fig. 1b for a comparison). Correspondingly, FGR and AGR are higher, while CGR does not change significantly (Figs 4c and 4d, see Figs 1c and 1d for a comparison). The ratios of AGR to FGR (solid line in Fig. 4e) and CGR to FGR (dotted line in Fig. 4e) are lower when there are small stable clusters present (see Fig. 1e for a comparison). This is clear especially at small sizes, indicating that FGR increases there more
- 20 than AGR or CGR due to the elevated concentrations of small clusters. The increase of FGR in the smallest size bin can be explained by a slower decrease of the concentration as a function of the cluster size in the presence of small stable clusters (see Fig A4 in Appendix). The AGR to FGR ratio varies between 10^2 and 10^8 in the smallest size bin and between 1.5 and 3.5 in the largest size bin, being highest with the lowest monomer concentration. The CGR to FGR behaves similarly to the AGR to FGR ratio; the most notable difference is that the CGR to FGR ratio is below one (0.6–0.7) at the largest sizes when C_{mon} =
- 25 5×10^6 cm⁻³ and $C_{\text{mon}} = 10^7$ cm⁻³. On the other hand, the presence of small stable clusters increases the AGR to CGR ratio slightly (Fig. 4f, see Fig. 1f for a comparison). The AGR to CGR ratio varies between 0.8 and 6 at $C_{\text{mon}} = 10^6$ cm⁻³, between 1.2 and 2.0 at $C_{\text{mon}} = 5 \times 10^6$ cm⁻³, and 1.8 and 4.2 at $C_{\text{mon}} = 10^7$ cm⁻³.

3.5 Combined effect of external conditions and the properties of model substance

To see the combined effect of external conditions and the properties of model substance on the growth of clusters, an additional 30 set of simulations was performed by varying monomer source rate and saturation vapor pressure simultaneously. A Gibbs free energy profile containing a maximum and no minima was assumed. Figure 5a shows the non-monomer fraction of the flux from the smallest size bin (solid lines) and from the largest size bin (dashed lines) in all these simulations. The ratio of the monomer source rate to the loss coefficient (Q/S), which largely determines how the system behaves, is presented on the *x*axis, and the color of the line shows the saturation vapor pressure. The non-monomer fraction of the flux increases with

35 increasing Q/S and decreasing saturation vapor pressure. With the highest saturation vapor pressure, the non-monomer fraction ranges from 6% to 21% in the smallest size bin and from 1% to 17% in the largest size bin, while with the lowest saturation vapor pressure, the ratios are 10%–53% and 3%–44% in the smallest and largest bin, respectively.

The ratio of AGR to FGR and AGR to CGR in different simulations is presented for the smallest size bin (solid lines) and the largest size bin (dashed lines) in Figs 5b and 5c. In the smallest size bin, the AGR to FGR ratio decreases with increasing Q/S

40 and the decreasing saturation vapor pressure. In the largest size bin, however, the ratio has a minimum at $Q/S = 5 \times 10^6 \text{ cm}^{-3}$ with the lowest saturation vapor pressures, and at highest Q/S-values the ratio is lowest when $p_{\text{sat}} = 10^{-9}$ Pa. In the smallest size

bin the AGR to FGR ratio ranges from 500 to 10^{16} and from 100 to 10^{9} with the highest and lowest saturation vapor pressures, respectively; in the largest size bin the corresponding ranges for the ratio are $1.4-10^{4}$ and 2.2-3.3. These results show that, depending on the external conditions and the properties of model substance, FGR can be several orders of magnitude lower than AGR, especially at the smallest sizes. This is related to the fact that in the smallest size bin FGR is not the best quantity

- 5 for describing the cluster growth rate (see the last paragraph of Sect. 3.1). On the other hand, if one wishes to use the growth rate to estimate the particle fluxes, the fluxes J_{app} calculated from AGR using Eq. (5) are in these cases significantly too high. AGR and CGR are considerably closer to each other compared to FGR and AGR or CGR. In the smallest size bin the AGR to CGR ratio varies from 0.5 to 9.5 and from 1.3 to 3.3 with the highest and lowest saturation vapor pressures, respectively, and in the largest size bins the corresponding ranges for the ratios are 1.3–1.6 and 0.8–3.
- Finally, we also studied the total concentration of clusters (2–70-mers) in different simulations. Figure 5d shows that the concentration of clusters increases with increasing Q/S and the decreasing saturation vapor pressure. With the highest saturation vapor pressure the steady state cluster concentration varies from 1.9×10^3 cm⁻³ to 2.9×10^6 cm⁻³, and with the lowest saturation vapor pressure from 1.2×10^4 cm⁻³ to 4.3×10^6 cm⁻³. When comparing Figs 5a–5d, we may conclude that in the conditions where the concentration of clusters becomes high, and thus their collisions become more important relative to
- 15 evaporation and other losses, the contribution of non-monomer collisions to the growth of clusters becomes significant. Furthermore, in these conditions growth rates determined with different methods tend to be closer to each other than in the conditions where cluster concentrations are lower.

4 Conclusions

We used a dynamic model to simulate the time evolution of cluster concentrations in a system where cluster-cluster collisions

- 20 significantly contribute to the growth of clusters. More specifically, we studied how consistent the flux-equivalent growth rate (FGR), the growth rate derived from the appearance times of the clusters (AGR), and the growth rate calculated based on irreversible vapor condensation (CGR) are with each other in different, atmospherically relevant, conditions.
- In majority of the simulations the Gibbs free energy of formation of the clusters was assumed to have a single maximum and no minima, which corresponds to the increasing stability of clusters with increasing cluster size. In most of these simulations
- 25 FGR was lower than AGR and CGR. The difference was highest, often several orders of magnitude, in the smallest size bin (at ~1.2 nm). This results from the very low value of FGR at the smallest sizes, caused by the approximations made in its derivation. In the largest size bin (at ~2.2 nm), FGR was closer to AGR and CGR. The difference between FGR and AGR or CGR was observed to decrease in conditions where cluster concentrations are high and thus evaporation and other losses are less important, i.e. when the monomer source rate is high, when the saturation vapor pressure is low and when the external
- 30 losses of clusters are low. Furthermore, in these conditions a higher fraction of the flux was found to be due to cluster-cluster collisions than in the conditions with lower cluster concentrations. Finally, it was observed that AGR and CGR are typically clearly closer to each other than to FGR; their difference was often very small, and within the factor of 10 in all the simulations. This is rather surprising as AGR is affected by all possible collision and evaporation processes between the clusters, while CGR is derived considering only the condensation of single molecules.
- 35 In one simulation set, a different free energy profile was used, leading to elevated concentrations of stable small clusters, which could correspond to the situation in the atmosphere. In this case, a significantly higher fraction of the growth was due to clustercluster collisions than in other simulations. Furthermore, the growth rates of clusters were higher and the different growth rates were closer to each other than in the simulations without stable small clusters.

Moreover, the used size resolution, i.e. the size bin width, was observed to affect the relation between the different growth

40 rates. Generally, the difference between the different growth rates increased with increasing size bin width. Thus, when determining growth rates from measured particle size distributions, a size resolution as high as possible should be used.

Altogether, our results demonstrate that different approaches to determine the growth rates of nanometer-sized clusters may give different values depending on the ambient conditions, the properties of the condensing vapor and the clusters, and the size resolution used in the analysis. Especially at the smallest, sub-2nm sizes, the differences between growth rates deduced with different methods can be significant. Our results also indicate that the conventional method used to determine particle formation rates based on growth rates may give estimates far from the true particle fluxes. This should be kept in mind when

applying these methods to measured particle size distributions, and utilizing the results in particle formation event analyses.

Acknowledgements

This research was supported by the Academy of Finland Centre of Excellence program (grant no. 272041), the European Research Council (ERC) projects ATM-NUCLE (grant no. 227463), MOCAPAF (grant no. 257360), and ATMOGAIN (grant no. 278277), and the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie

(grant no. 656994).

5

10

References

Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P., and Lehtinen, K. E. J.: Formation and growth of fresh atmospheric aerosols: Eight years of aerosol size distribution data from SMEAR II, Hyytiälä, Finland, Boreal Environ.

15 Res., 10, 323–336, 2005.

Friedlander, S. K.: Smoke, dust and haze – fundamentals of aerosol dynamics, John Wiley and Sons, New York, 1977.

Hirsikko, A., Laakso, L, Hõrrak, U., Aalto, P. P., Kerminen, V.-M., and Kulmala, M.: Annual and size dependent variation of growth rates and ion concentrations in boreal forest, Boreal Environ. Res., 10, 357–369, 2005.

Jiang, J. K., Zhao, J., Chen, M. D., Eisele, F. L., Scheckman, J., Williams, B. J., Kuang, C. A., and McMurry, P. H.: First
Measurements of Neutral Atmospheric Cluster and 1–2 nm Particle Number Size Distributions During Nucleation Events, Aerosol Sci. Technol., 45, II–V, doi:10.1080/02786826.2010.546817, 2011.

Kuang, C., Chen, M., Zhao, J., Smith, J., McMurry, P. H., and Wang, J.: Size and time-resolved growth rate measurements of 1 to 5 nm freshly formed atmospheric nuclei, Atmos. Chem. Phys., 12, 3573-3589, doi:10.5194/acp-12-3573-2012, 2012.

Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V.-M., Birmili, W., and McMurry, P. H.:
Formation and growth rates of ultrafine atmospheric particles: A review of observations, J. Aerosol Sci., 35, 143–176, 2004.

Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V.-M.: Measurement of the nucleation of atmospheric aerosol particles, Nat. Protoc., 7, 1651–1667, doi:10.1038/nprot.2012091, 2012.

Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger,
S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä , M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M. V, Smith, J. N., Ehn, M., Mentel, T. F., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., and Worsnop, D. R.: Direct observations of atmospheric aerosol nucleation, Science, 339, 943–946, doi:10.1126/science.1227385, 2013.

Kurtén, T., Loukonen, V., Vehkamäki, H., and Kulmala, M.: Amines are likely to enhance neutral and ion-induced sulfuric acid-water nucleation in the atmosphere more effectively than ammonia, Atmos. Chem. Phys., 8, 4095-4103, doi:10.5194/acp-8-4095-2008, 2008.

Lehtinen, K. E. J. and Kulmala, M.: A model for particle formation and growth in the atmosphere with molecular resolution in size, Atmos. Chem. Phys., 3, 251-257, doi:10.5194/acp-3-251-2003, 2003.

5

10

Lehtinen, K. E. J., dal Maso, M., Kulmala, M., and Kerminen, V.–M.: Estimating nucleation rates from apparent particle formation rates and vice versa: Revised formulation of the Kerminen–Kulmala equation, J. Aerosol Sci., 38, 988–994, 2007.

Lehtipalo, K., Leppä, J., Kontkanen, J., Kangasluoma, J., Franchin, A., Wimmer, D., Schobesberger, S., Junninen, H., Petäjä, T., Sipilä, M., Mikkilä, J., Vanhanen, J., Worsnop, D. R., and Kulmala, M.: Methods for determining particle size distribution and growth rates between 1–3 nm using the Particle Size Magnifier, Boreal Environ. Res., 19, 215–236, 2014.

Lehtipalo, K., Rondo, L., Kontkanen, J., Schobesberger, S., Jokinen, T., Sarnela, N., Kürten, A., Ehrhart, S., Franchin, A., Nieminen, T., Riccobono, F., Sipilä, M., Yli-Juuti, T., Duplissy, J., Adamov, A., Ahlm, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., Dommen, J., Downard, A. J., Dunne, E. M., Flagan, R. C., Guida, R., Hakala, J., Hansel, A., V. –M., Keskinen, H., Kim, Jud, W., Kangasluoma, J., Kerminen, J., Kirkby, J., Kupc, A., Kupiainen-Määttä, Laaksonen, Lawler, M. J., Leiminger, O., A., М., Mathot, S., Olenius, Т.,

- 15 Ortega, I. K., Onnela, A., Petäjä, T., Praplan, A., Rissanen, M. P., Ruuskanen, T., Santos, F. D., Schallhart, S., Schnitzhofer, R., Simon, М., Smith, J. N., Tröstl, J., Tsagkogeorgas, G., Tomé, A., Vaattovaara, Ρ., Vehkamäki, Н., Vrtala, A. Е., Wagner, P. Е., Williamson, C., Carslaw, P. M., Donahue, N. М., Wimmer, D., Winkler, Virtanen, A., Κ. S., Baltensperger,
- 20 U., Riipinen, I., Curtius, J., Worsnop, D. R., and Kulmala, M.: The effect of acid-base clustering and ions on the growth of atmospheric nano-particles, Nature Communications, 2016, in review.

McGrath, M. J., Olenius, T., Ortega, I. K., Loukonen, V., Paasonen, P., Kurtén, T., Kulmala, M., and Vehkamäki, H.: Atmospheric Cluster Dynamics Code: a flexible method for solution of the birth-death equations, Atmos. Chem. Phys., 12, 2345-2355, doi:10.5194/acp-12-2345-2012, 2012.

25 Manninen, H. E., Nieminen, T., Riipinen, I., Yli-Juuti, T., Gagné, S., Asmi, E., Aalto, P. P., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Charged and total particle formation and growth rates during EUCAARI 2007 campaign in Hyytiälä, Atmos. Chem. Phys., 9, 4077-4089, doi:10.5194/acp-9-4077-2009, 2009.

Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, Atmos. Chem. Phys., 9, 8601–8616, doi:10.5194/acp-9-8601-2009, 2009.

30 Nieminen, T., Lehtinen, K. E. J., and Kulmala, M.: Sub-10 nm particle growth by vapor condensation – effects of vapor molecule size and particle thermal speed, Atmos. Chem. Phys., 10, 9773-9779, doi:10.5194/acp-10-9773-2010, 2010.

Olenius, T., Riipinen, I., Lehtipalo, K., and Vehkamäki, H.: Growth rates of atmospheric molecular clusters based on appearance times and collision–evaporation fluxes: Growth by monomers, J. Aerosol Sci., 78, 55–70, 2014.

Ortega, I. K., Kupiainen, O., Kurtén, T., Olenius, T., Wilkman, O., McGrath, M. J., Loukonen, V., and Vehkamäki, H.: From
quantum chemical formation free energies to evaporation rates, Atmos. Chem. Phys., 12, 225-235, doi:10.5194/acp-12-225-2012, 2012.

Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics: from air pollution to climate change, 2nd Edn., John Wiley & Sons, Inc., Hoboken, NJ, 2006.

Sihto, S.-L., Kulmala, M., Kerminen, V.-M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A., and Lehtinen, K. E. J.: Atmospheric sulphuric acid and aerosol formation: implications from

5 atmospheric measurements for nucleation and early growth mechanisms, Atmos. Chem. Phys., 6, 4079-4091, doi:10.5194/acp-6-4079-2006, 2006.

Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I., Merikanto, J., Mann, G. W., Chipperfield, M. P., and Wiedensohler, A.: Contribution of particle formation to global cloud condensation nuclei concentrations, Geophys. Res. Lett., 35, L06808, doi:10.1029/2007GL033038, 2008.

10 Wang, J., McGraw, R. L., and Kuang, C.: Growth of atmospheric nano-particles by heterogeneous nucleation of organic vapor, Atmos. Chem. Phys., 13, 6523-6531, doi:10.5194/acp-13-6523-2013, 2013.

Vanhanen, J., Mikkilä, J., Sipilä, M., Manninen, H. E., Lehtipalo, K., Siivola, E., Petäjä, T., and M., K. Particle Size Magnifier for Nano-CN Detection. Aerosol Sci. Technol. 45:533–542, 2011.

Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measurements of new particle formation and ultrafine particle growth rates at a clean continental site, J. Geophys. Res., 102, 4375–4385, 1997.

Vehkamäki, H., McGrath, M. J., Kurtén, T., Julin, J., Lehtinen, K. E. J., and Kulmala, M.: Rethinking the application of the first nucleation theorem to particle formation, J. Chem. Phys., 136, 094107, doi:10.1063/1.3689227, 2012.

Vuollekoski, H., Sihto, S.-L., Kerminen, V.-M., Kulmala, M., and Lehtinen, K. E. J.: A numerical comparison of different methods for determining the particle formation rate, Atmos. Chem. Phys., 12, 2289-2295, doi:10.5194/acp-12-2289-2012, 2012

Yli-Juuti, T., Nieminen, T., Hirsikko, A., Aalto, P. P., Asmi, E., Hõrrak, U., Manninen, H. E., Patokoski, J., Dal Maso, M., Petäjä, T., Rinne, J., Kulmala, M., and Riipinen, I.: Growth rates of nucleation mode particles in Hyytiälä during 2003–2009: variation with particle size, season, data analysis method and ambient conditions, Atmos. Chem. Phys., 11, 12865-12886, doi:10.5194/acp-11-12865-2011, 2011.

25

20

15

Simulation set 1: Varying the monomer source rate			
Monomer source rate (cm ⁻³ s ⁻¹)	1.0×10 ³	5.5×10 ³	1.8×10 ⁴
Steady-state monomer concentration (cm ⁻³)	1×10^{6}	5×10^{6}	1×10^{7}
Loss coefficient (s ⁻¹)	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹
Saturation vapor pressure (Pa)	2×10 ⁻¹⁰	2×10 ⁻¹⁰	2×10 ⁻¹⁰
Size bin width (molecules)	10	10	10
Simulation set 2: Varying the saturation vapor pressure			
Monomer source rate (cm ⁻³ s ⁻¹)	<mark>5.8×10³</mark>	5.5×10^{3}	5.1×10 ³
Steady-state monomer concentration (cm ⁻³)	5×10^{6}	5×10^{6}	5×10^{6}
Loss coefficient (s ⁻¹)	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹
Saturation vapor pressure (Pa)	1.5×10 ⁻¹⁰	2×10 ⁻¹⁰	1×10 ⁻⁹
Size bin width (molecules)	10	10	10
Simulation set 3: Varying the loss coefficient			
Monomer source rate (cm ⁻³ s ⁻¹)	<mark>5.5×10³</mark>	5.5×10 ³	5.5×10 ³
Steady-state monomer concentration (cm ⁻³)	6.4×10^{6}	5×10^{6}	2.6×10^{6}
Loss coefficient (s ⁻¹)	7×10 ⁻⁴ s ⁻¹	10 ⁻³ s ⁻¹	2×10 ⁻³ s ⁻¹
Saturation vapor pressure (Pa)	2×10 ⁻¹⁰	2×10 ⁻¹⁰	2×10 ⁻¹⁰
Size bin width (molecules)	10	10	10
Simulation set 4: Varying the size bin width			
Monomer source rate (cm ⁻³ s ⁻¹)	5.5×10 ³	5.5×10 ³	5.5×10 ³
Steady-state monomer concentration (cm ⁻³)	5×10^{6}	5×10^{6}	5×10^{6}
Loss coefficient (s ⁻¹)	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹
Saturation vapor pressure (Pa)	2×10 ⁻¹⁰	2×10 ⁻¹⁰	2×10 ⁻¹⁰
Size bin width (molecules)	5	10	14
Simulation set 5: Varying the monomer source rate with a different Gibbs free energy profile			
Monomer source rate (cm ⁻³ s ⁻¹)	1.1×10^{3}	<mark>9.6×10³</mark>	5.5×10 ⁴
Steady-state monomer concentration (cm ⁻³)	1×10^{6}	5×10^{6}	1×10^{7}
Loss coefficient (s ⁻¹)	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹	10 ⁻³ s ⁻¹
Saturation vapor pressure (Pa)	2×10 ⁻¹⁰	2×10 ⁻¹⁰	2×10 ⁻¹⁰
Size bin width (molecules)	10	10	10

ie 1. Summary of the performed simulations.



Figure 1. The effect of steady-state monomer concentration (shown as different colors) on quantities describing cluster growth: a) the non-monomer fraction of flux from each size bin, b) the true collision-evaporation flux from each size bin (J_{true} ; solid line) and the fluxes calculated from the appearance times of clusters (J_{app} ; dashed line), c) and d) the flux-equivalent growth rate (FGR; solid line), the growth rate derived from the appearance times of clusters (AGR; dashed line), and the growth rate calculated based on irreversible vapor condensation (CGR; dotted line), e) the ratios of AGR to FGR (solid line) and CGR to FGR (dotted line), f) the ratio of AGR to CGR.

5



Figure 2. The effect of saturation vapor pressure (shown as different colors) on quantities describing cluster growth: a) the non-monomer fraction of flux from each size bin, b) the true collision-evaporation flux from each size bin (J_{true} ; solid line) and the fluxes calculated from the appearance times of clusters (J_{app} ; dashed line), c) and d) the flux-equivalent growth rate (FGR; solid line), the growth rate derived from the appearance times of clusters (AGR; dashed line), and the growth rate calculated based on irreversible vapor condensation (CGR; dotted line), e) the ratios of AGR to FGR (solid line) and CGR to FGR (dotted line), f) the ratio of AGR to CGR.



Figure 3. The effect of size bin width (shown as different colors) on quantities describing cluster growth: a) the non-monomer fraction of flux from each size bin, b) the true collision-evaporation flux from each size bin (J_{true} ; solid line) and the fluxes calculated from the appearance times of clusters (J_{app} ; dashed line), c) and d) the flux-equivalent growth rate (FGR; solid line), the growth rate derived from the appearance times of clusters (AGR; dashed line), and the growth rate calculated based on

irreversible vapor condensation (CGR; dotted line), e) the ratios of AGR to FGR (solid line) and CGR to FGR (dotted line), f) the ratio of AGR to CGR.

5



Figure 4. The effect of steady-state monomer concentration (shown as different colors) on quantities describing cluster growth in the presence of stable small clusters: : a) the non-monomer fraction of flux from each size bin, b) the true collisionevaporation flux from each size bin (J_{true} ; solid line) and the fluxes calculated from the appearance times of clusters (J_{app} ; dashed line), c) and d) the flux-equivalent growth rate (FGR; solid line), the growth rate derived from the appearance times of clusters (AGR; dashed line), and the growth rate calculated based on irreversible vapor condensation (CGR; dotted line), e)

5

the ratios of AGR to FGR (solid line) and CGR to FGR (dotted line), f) the ratio of AGR to CGR.



Figure 5. The results of the simulations with different saturation vapor pressures (p_{sat} ; shown as different colors) and different ratios of monomer source rate to the loss coefficient (Q/S; shown on the *x*-axis) in the smallest size bin (solid line) and in the largest size bin (dashed line): a) the non-monomer fraction of the flux, b) the ratio of the appearance time growth rate (AGR) to the flux-equivalent growth rate (FGR), c) the ratio of the appearance time growth rate to the growth rate calculated based on irreversible vapor condensation (CGR), d) the steady-state concentration of all clusters (2–70-mers).



Figure A1. a) The classical Gibbs free energy profile with one maximum and no minima (solid lines) and with an additional negative term corresponding to the stabilization of the smallest clusters (dashed lines) at different steady-state monomer
concentrations (shown as different colors). b) Evaporation profiles corresponding to the two Gibbs free energy profiles (note that the evaporation rate is independent of the vapor concentration).



Figure A2. The time evolution of the cluster distribution during the first 5 hours of one of the simulations, where the steady state monomer concentration was set to 5×10^6 cm⁻³, the loss coefficient to 10^{-3} s⁻¹, and the saturation vapor pressure to 2×10^{-10} Pa. The concentrations of the vapor monomer and the individual clusters are shown as the lines with colors from light blue

5 to pink. The different size bins, for which the growth rates were determined, are shown as the thick lines with colors from black to light brown. The black circles mark the appearance times of each size bin.



5

Figure A3. The effect of loss coefficient (shown as different colors) on quantities describing cluster growth: a) the nonmonomer fraction of flux from each size bin, b) the true collision-evaporation flux from each size bin (J_{true} ; solid line) and the fluxes calculated from the appearance times of clusters (J_{app} ; dashed line), c) and d) the flux-equivalent growth rate (FGR; solid line), the growth rate derived from the appearance times of clusters (AGR; dashed line), and the growth rate calculated based on irreversible vapor condensation (CGR; dotted line), e) the ratios of AGR to FGR (solid line) and CGR to FGR (dotted line), f) the ratio of AGR to CGR.



Figure A4. Cluster distribution at the mean appearance time of the two smallest size bins (solid lines) and at the final steady state (dashed lines) when a) the Gibbs free energy profile has one maximum and no minima b) the Gibbs free energy profile has an additional negative term corresponding to the stabilization of the smallest clusters. The colors show the steady state monomer concentration.