

1 **Size-resolved aerosol composition at an urban and a rural site in**  
2 **the Po Valley in summertime: implications for secondary aerosol**  
3 **formation**

4 S. Sandrini<sup>1</sup>, D. van Pinxteren<sup>2</sup>, L. Giulianelli<sup>1</sup>, H. Herrmann<sup>2</sup>, L. Poulain<sup>2</sup>, M.C. Facchini<sup>1</sup>, S.  
5 Gilardoni<sup>1</sup>, M. Rinaldi<sup>1</sup>, M. Paglione<sup>1</sup>, B.J. Turpin<sup>3</sup>, F. Pollini<sup>1</sup>, N. Zanca<sup>1</sup>, S. Decesari<sup>1</sup>

6 1 Institute for Atmospheric Sciences and Climate (ISAC), National Research Council (CNR), 40129 Bologna, Italy

7 2Leibniz-Institut für Troposphärenforschung (TROPOS), 04318 Leipzig, Germany

8 3 - Environmental Science and Engineering, Gillings School of Global Public Health, University of North Carolina at  
9 Chapel Hill – NC 27599-7431 - USA

10

11 Correspondence to: Silvia Sandrini (S.Sandrini@isac.cnr.it)

1 **Abstract.** The aerosol size-segregated chemical composition was analyzed at an urban (Bologna) and a rural site (San  
2 Pietro Capofiume) in the Po Valley, Italy, during June and July 2012, by ion-chromatography (major water soluble ions  
3 and organic acids) and evolved gas analysis (total and water soluble carbon), to investigate sources and mechanisms of  
4 secondary aerosol formation during the summer. A significant enhancement of secondary organic and inorganic aerosol  
5 mass was observed under anticyclonic conditions with recirculation of planetary boundary layer air, but with substantial  
6 differences between the urban and the rural site. The data analysis, including a Principal Component Analysis (PCA) on  
7 the size-resolved dataset of chemical concentrations, indicated that the photochemical oxidation of inorganic and  
8 organic gaseous precursors was an important mechanism of secondary aerosol formation at both sites. In addition at the  
9 rural site a second formation process, explaining the largest fraction (22%) of the total variance, was active at night-  
10 time, especially under stagnant conditions. Nocturnal chemistry in the rural Po Valley was associated with the  
11 formation of ammonium nitrate in large accumulation mode (0.42 – 1.2  $\mu\text{m}$ ) aerosols favored by local thermodynamic  
12 conditions (higher relative humidity and lower temperature compared to the urban site). Nocturnal concentrations of  
13 fine nitrate were, in fact, on average five times higher at the rural site than in Bologna. The water uptake by this highly  
14 hygroscopic compound under high RH conditions provided the medium for increased nocturnal aerosol uptake of water  
15 soluble organic gases and possibly also for aqueous chemistry, as revealed by the shifting of peak concentrations of  
16 secondary compounds (water soluble organic carbon (WSOC) and sulfate) toward the large accumulation mode (0.42 –  
17 1.2  $\mu\text{m}$ ). Contrarily, the diurnal production of WSOC (proxy for secondary organic aerosol) by photochemistry was  
18 similar at the two sites but mostly affected the small accumulation mode of particles (0.14 – 0.42  $\mu\text{m}$ ) in Bologna, while  
19 a shift to larger accumulation mode was observed at the rural site. A significant increment in carbonaceous aerosol  
20 concentration (for both WSOC and water insoluble carbon (WINC)) at the urban site was recorded mainly in the quasi-  
21 ultrafine fraction (size range 0.05-0.14  $\mu\text{m}$ ) indicating a direct influence of traffic emissions on the mass concentrations  
22 of this range of particles.

23

24

25

## 1. Introduction

26 The knowledge of the size-segregated chemical composition of atmospheric aerosols, i.e. the chemical composition as a  
27 function of particle size, is key to the understanding of several important characteristics of particles such as optical  
28 properties, hygroscopicity and reactivity, which affect the atmospheric radiation budget, cloud formation and human  
29 health. Moreover, the size distribution of inorganic and organic components reflects their origin, hence encloses a  
30 wealth of information about aerosol formation mechanisms and atmospheric processing, including secondary formation  
31 (Seinfeld and Pandis, 1998).

32 Accumulation mode particulate matter mostly originates from aerosol accretion via gas-to-particle conversion of  
33 oxidized vapors (Seinfeld and Pandis, 1998). However, accumulation mode particles can be selectively scavenged by  
34 clouds and fogs which, in absence of precipitation, produce chemically-processed particles upon evaporation of water  
35 (Meng and Seinfeld, 1994). The resulting size distribution is bimodal with the small mode accounting for secondary  
36 aerosols produced uniquely by gas-to-particle conversion (“condensation mode”) and the large mode containing  
37 particles that underwent cloud processing (“droplet mode”) (Hering and Friedlander, 1982; John et al., 1990).  
38 Therefore, knowledge about the concentrations of aerosol organic and inorganic compounds in size-segregated aerosol  
39 samples provides information on the nature of secondary formation processes. Measurements of the size-segregated  
40 chemical composition of aerosols are traditionally performed using multi-stage impactors followed by off-line chemical

1 analysis. Recently, the development of online mass spectrometric techniques offered new opportunities for size-  
2 segregated chemical observations with a much greater time-resolution with respect to impactors (Jimenez et al., 2003).  
3 However, aerosol mass spectrometers (AMS) generally suffer of poor sensitivity for thermally refractory compounds  
4 and could not be deployed for the analysis of coarse particles ( $>2.5 \mu\text{m}$ ) chemical composition. Therefore, multistage  
5 impactors are still unsurpassed in terms of the number of chemical determinations that can be performed on the samples  
6 and the range of particle sizes that can be probed. An example of comparison of aerosol chemical measurements  
7 performed using a five-stage impactor and AMS is provided by our previous study focusing on the 2009 field campaign  
8 in the Po Valley, Italy (Decesari et al., 2014).

9 The Po Valley is a region in Europe characterized by high levels of pollution, due to the concurrent high density of  
10 anthropogenic sources and its orographic and meteorological characteristics particularly unfavourable for pollutant  
11 dispersion. In particular, several studies have shown how this area is dominated by secondary material during the  
12 summer (Crosier et al., 2007), with a large presence of secondary inorganic aerosols (SIA). On an annual average, SIA  
13 accounts for 40% of  $\text{PM}_{10}$  mass at the urban site of Bologna (Matta et al., 2003; Putaud et al., 2010) while a 50%  
14 contribution has been evaluated at the rural site of San Pietro Capofiume over shorter observation periods during winter  
15 and summer (Carbone et al., 2010). Crosier et al., (2007) observed that most of the time, during the summer, aerosol  
16 over the Po Valley was composed of regional ammonium sulfate and organic material, while under anticyclonic  
17 conditions, with recirculation of air over the region, the composition was dominated by ammonium nitrate close to  
18 ammonia emission sources.

19 The present study focuses on the chemical size-resolved composition of aerosol determined by two 5-stage low-pressure  
20 Berner impactors during a field campaign performed in the summer 2012 in the Po Valley. With respect to the previous  
21 campaigns in this area, this is the first experiment where a two-site approach was used. One-month long intensive  
22 observations were performed at an urban site (Bologna, BO) and at a nearby rural site (San Pietro Capofiume, SPC).  
23 The goal of this paper is to provide insights into factors controlling the variability of aerosol composition and to explore  
24 possible formation pathways of secondary compounds in this region during the summer under different meteorological  
25 conditions and air mass history. The two-site approach was adopted to estimate the contribution and composition of  
26 rural background particles with respect to the urban contribution, according to the Lenschow perspective (Lenschow et  
27 al., 2001), based on the assumption that PM concentration measured at an urban location is the result of the addition of  
28 a regional background, urban contribution given by the sources inside the agglomeration, and road traffic for near  
29 streets sites.

30 Source attribution was addressed through the analysis of the time series of the main aerosol species together with  
31 meteorological parameters, and by the statistical approach of principal component analysis (PCA) applied to the  
32 different size classes as independent variables. The study also takes into account the comparison between an urban and  
33 a rural site to assess the impact of traffic and other urban sources on the regional background, to explore differences in  
34 secondary aerosol formation resulting from different meteorological conditions, and to assess the regional and local  
35 variability of secondary aerosol formation processes. Finally, the fact that sampling was performed separately during  
36 day and night, allowed analysis of the concentrations of aerosol constituents together with the dynamics of the boundary  
37 layer (Gietl et al., 2008).

38

## 39 **2. Experimental**

### 40 **2.1 Sampling sites**

1 Size-segregated aerosol sampling was performed during the PEGASOS field campaign in the Po Valley (Italy), from  
2 June 12 to July 9, 2012, at the urban site of Bologna (BO - 44° 29' N, 11° 20' E, 54 m a.s.l.) and at the rural site of San  
3 Pietro Capofiume (SPC - 44°39' N, 11° 37' E, 11 m a.s.l.), 30 km northeast from the city of Bologna. Both sites are  
4 located in the eastern part of the Po Valley (Fig. 1). Bologna is a city of 400.000 inhabitants, the most populous in the  
5 southern Po Valley, with a surrounding area characterized by widespread agricultural and industrial activities and by the  
6 presence of several high-traffic roads. Sampling was performed in the northern outskirts of Bologna, on the roof of the  
7 Institute of Atmospheric Sciences and Climate of the National Research Council, at about 20 m above ground. San  
8 Pietro Capofiume is a rural site characterized by a flat terrain and by croplands extending in all directions, and can be  
9 considered an ideal receptor site for regional-scale air pollution in the Po Valley. In SPC aerosol samplers were  
10 positioned on a platform at about 8 m above ground level.

11 The west-east orientation of the valley favours westerly or easterly circulations, hence from either the inner Po Valley  
12 (including Lombardy and Milan) or the Adriatic Sea. In the presence of SW winds, SPC can result downwind of BO  
13 and an influence from the urban area cannot be excluded. Under the same conditions, a possible inflow of marine air  
14 from the Ligurian Sea is also possible.

15

## 16 **2.2 Aerosol sampling**

17 Two five-stage Berner impactors (flow rate 80 L min<sup>-1</sup>) with 50% particle cutpoints at 0.14, 0.42, 1.2, 3.5, and 10 µm  
18 aerodynamic diameter ( $D_p$ ) were used at the two sites. The particles were collected on aluminium and Tedlar foils.  
19 Aluminium foils for carbonaceous aerosol analysis were placed on each stage of the impactor with Tedlar half foils for  
20 ion-chromatographic analysis placed on top of them, covering 50% of the aluminium substrates (Matta et al., 2003). A  
21 12-hour time resolution was adopted for sampling, one night-time and one daytime sample collected every day, from  
22 21:00 to 09:00 LT and from 09:00 to 21:00 LT respectively. However, it is worth noting that nocturnal sampling  
23 actually included several hours of light: from dawn (which in this period of the year occurs at approximately 5:30 am)  
24 to 9 am.

25

## 26 **2.3 Analytical measurements**

27 The Tedlar substrates were extracted in 10 mL of mQ water for 30 minutes in an ultrasonic bath. The extracts were  
28 analysed by ion-chromatography for the quantification of water-soluble inorganic species and organic acids (acetate,  
29 formate, methansulfonate, oxalate). A TOC analyzer (Shimadzu, Japan) was used for the determination of water-soluble  
30 organic carbon (WSOC). An aliquot of each sample is analyzed for total carbon (TC) by combustion at 680°C in the  
31 presence of a catalyst, while another aliquot is acidified in a reaction vessel to determine IC, which is decomposed to  
32 CO<sub>2</sub>. In both cases the evolved CO<sub>2</sub> is measured by a non-dispersive infrared gas analyser (NDIR) and the organic  
33 fraction (WSOC) is obtained by difference between TC and IC.

34 Fractions of the aluminium foils were used for the quantification of total carbon (TC) by evolved gas analysis with a  
35 Multi N/C2100 analyser (Analytik Jena, Germany) equipped with a module for solid samples. Portions of the  
36 aluminium foils were exposed to increasing temperature (up to 950°C) in pure oxygen carrier gas. Under these

1 conditions all carbonaceous matter (organic, carbonate and elemental carbon) is converted to CO<sub>2</sub> (Gelencser et al.,  
2 2000) and TC is measured as total evolved CO<sub>2</sub> by a non-dispersive infrared (NDIR) analyser.  
3 Aluminium foils in SPC were also used for the gravimetric determination of aerosol mass by weighing the substrates  
4 before and after sampling on a UMT-2 microbalance with a reading precision of 0.1 µg and a standard deviation of ca.  
5 1%.  
6

## 7 **2.4 Back-trajectory calculation**

8 Air mass back trajectories are a useful tool when studying the aerosol composition as a function of the air mass history.  
9 For every hour during each sampling interval (12 hours during night and 12 hours during day) 96 h back trajectories  
10 arriving at 500 m a.g.l. were calculated by the HYSPLIT model (HYbrid Single-Particle Lagrangian Integrated  
11 Trajectory, version 4) (Draxler and Rolph, 2003) in the ensemble mode, using input field from the global 1° GDAS  
12 archive (<http://www.arl.noaa.gov/ss/transport/archives.html>). Ensembles of 27 trajectories for a given starting time for all  
13 possible off-sets in X, Y, and Z dimensions (ca. 250 m off-set in Z, one grid cell off-set in X and Y) were calculated for  
14 every hour, resulting in a total of 324 trajectories describing the air-mass history of each 12h sample. In addition to the  
15 endpoints of the trajectories other HYSPLIT output parameters (sunflux, mixing layer depth) were stored and averaged  
16 along the trajectories (sunflux) and at the receptor site (sunflux and mixing layer depth) during the sampling intervals.  
17 Even though the mixing layer height provided by the HYSPLIT model might be quite inaccurate, the trend of this  
18 parameter was taken as a proxy of the boundary layer dynamics for the campaign. In addition, residence time indices  
19 (RTIs) were calculated by GIS analysis, reflecting the time the sampled air masses resided above certain land cover  
20 categories (water, natural vegetation, agricultural lands, bare areas and urban areas). Details of this method are given in  
21 van Pinxteren et al., (2010).  
22  
23

## 24 **2.5 Aerosol Liquid Water Content calculation**

25 Hourly aerosol liquid water content (ALWC) was calculated by the online versions of the Extended Aerosol Inorganic  
26 Model III (E-AIM, <http://www.aim.env.uea.ac.uk/aim/aim.php>; (Clegg et al., 1998); (Wexler and Clegg, 2002)). The  
27 inorganic concentrations for sulfate, ammonium, nitrate, sodium and chloride measured by two HR-TOF-AMS, placed  
28 respectively in BO and in SPC during the campaign, were used as inputs in conjunction with RH, while temperature  
29 was kept fixed at 298.15 by this model. The size segregated concentrations of the inorganic components collected by  
30 the Berner impactors could not be used for this purpose, due to their low temporal resolution (12 hours), resulting in a  
31 flattening of RH and T values averaged over the sampling time, which prevented accurate calculation of ALW on  
32 particles. Particulate water retained by polar organic matter was neglected in the calculations, because the inclusion of  
33 ionic organic compounds (oxalic, glutaric and maleic carboxylic acids) had been shown to play only a minor role in  
34 water uptake during the campaign (Hodas et al., 2014). The ALWC calculated from AMS relates to the fine fraction (<  
35 1 µm diameter) of aerosol.  
36

## 37 **2.6 Principal Component Analysis**

38 Source categories for the 5 impactor size intervals of particulate matter were studied by means of Principal Component  
39 Factor Analysis (PCA) using the XLStat software (Addinsoft, version 2013.2.04). Besides the concentrations of particle

1 constituents chloride, sulfate, nitrate, sodium, ammonium, magnesium, calcium and WSOC, the database for PCA also  
2 included the modelled meteorological parameters from the HYSPLIT model listed in the previous section, the measured  
3 meteorological parameters temperature and relative humidity and the modelled residence times from GIS analysis.  
4 The values below detection limits were replaced by half the respective value of detection limit (Farnham et al., 2002) in  
5 the final dataset.

6 Since a prerequisite of PCA is the normal distribution of the variables used in the analysis, the normal distribution of  
7 concentration data has been checked by the Shapiro-Wilk normality test. Data that were not normally distributed were  
8 log-10 transformed before the analysis.

9 The orthogonal transformation method with Varimax rotation of Principal Components was applied to redistribute the  
10 variance in order to generate more interpretable factor loadings and scores (Vandeginste, 1998). As there are no defined  
11 criteria for the number of factors which are used in the Varimax rotation, we performed several PCAs with varying  
12 numbers of rotated factors (4–9) and judged on the interpretability of the results by trying to assign a physical meaning  
13 to the extracted factors. The number of rotated factors was regarded too high if factors showed very low contribution to  
14 the overall variance and no distinct physical meaning. Contrarily the number was regarded too low if previously  
15 resolved sources were now folded into one principal component. The most reasonable results were obtained by rotating  
16 the first 6 factors.

17

### 18 **3. Results and discussion**

19

#### 20 **3.1 Back trajectory patterns**

21 The PEGASOS summer campaign was characterized by the occurrence of different meteorological patterns, with a first  
22 part characterized by days of very perturbed weather followed by stable anticyclonic conditions, and a second part  
23 experiencing more variable meteorological conditions. During the study period, weak westerly breezes affected the  
24 sampling sites at night and in morning hours, while short, intense easterly breezes flowed during the late afternoon (Fig.  
25 2 in Wolf et al. (2015)). However, in days of stronger synoptic forcing, a wind pattern characterized by strong easterly  
26 winds, persisting all the day, was observed (especially in the second half of June), while a wind pattern characterized by  
27 SW winds from the Apennines was also common in June and accounted for most of the days in July until the end of the  
28 campaign. An overview of the main air mass transport patterns intersecting the area during the campaign, from back-  
29 trajectory analysis, is reported in a parallel paper (Decesari et al., in preparation). A hierarchical cluster analysis  
30 (Dorling et al., 1992) of the obtained back trajectories was performed to reduce the number of “origins of air masses”,  
31 appointing every calculated individual back trajectory to the most appropriate cluster, roughly corresponding to a  
32 specific synoptic situation. At each step of the process, the appropriate number of clusters was selected by looking at  
33 the variations of the total spatial variance (TSV - defined as the sum of the squared distances between the endpoints of  
34 the single trajectory and the mean of the trajectories in that cluster). The optimum number of clusters was selected in  
35 correspondence with the number after which the TSV did not vary substantially. The analysis led to identification of  
36 five main patterns affecting the Po Valley during the experimental campaign. A map with the mean trajectories for each  
37 cluster and the corresponding percentage of occurrence for trajectories calculated over 96 h and arriving at 500 m  
38 a.g.l. is shown in Fig. 2.

39 Clusters 1 and 3, defined respectively “WEST low” and “EAST low” according to their low travelling altitude (below  
40 1000 m a.g.l.) and prevalent direction, were characterized by short trajectory lengths, and corresponded to a higher

1 residence time of air masses in the basin. Cluster 1, in particular, had the highest occurrence and accounted for 38% of  
2 the total trajectories. Both included a smaller number of very short trajectories which were defined respectively as  
3 “WEST-low local“ and “EAST-low local“, occurring during the days characterized by stagnant conditions and low  
4 wind speed, and were associated with the accumulation of pollutants.

5  
6

### 7 **3.2 Bulk PM10 aerosol composition**

8 The size cut between fine and coarse particles in the Berner impactor size distributions is set to a 1.2  $\mu\text{m}$  aerodynamic  
9 diameter (i.e., the size cut between the 3<sup>rd</sup> and the 4<sup>th</sup> impactor stage), therefore in this study  $\text{PM}_{1.2}$  and  $\text{PM}_{1.2-10}$   
10 represent the fine and the coarse aerosol fractions, respectively. Fig. 3 shows the time series of  $\text{PM}_{1.2}$  and  $\text{PM}_{1.2-10}$  mass  
11 concentrations and of the contribution of  $\text{PM}_{1.2}$  and  $\text{PM}_{1.2-10}$  for the rural site SPC, together with air mass categories and  
12 back trajectory length. The aerosol mass was not available for BO. Air mass categories indicate the prevalent direction  
13 of airmasses during each sampling day.

14 During the campaign, the days showing the lowest aerosol mass concentrations were characterized by the longest  
15 trajectories, which corresponded to air masses transported over long-range from the North Atlantic Ocean (WEST), at  
16 the beginning and at the end of the sampling period. The aerosol mass increased from 15 to 18 June, with an enhanced  
17 contribution of the  $\text{PM}_{1.2}$  fraction on the total  $\text{PM}_{10}$ , together with a sharp decrease of the trajectories length, following  
18 the onset of an anticyclonic period with low wind and air stagnation over the Po Valley. An episode of Saharan dust  
19 transport was observed during a period of air transport from south, starting on 19 June at 5 km height with maximum on  
20 20 June when it reached the PBL, resulting in an increase of  $\text{PM}_{10}$  mass at ground level (Bucci et al., in preparation).  
21 The highest contributions of  $\text{PM}_{1.2}$  to  $\text{PM}_{10}$  were observed most of the times when 4-day trajectories were very short (<  
22 1500 km). During the first, persistent stagnation period, lasting from 16 to 19 June, the  $\text{PM}_{1.2}$  contribution to  $\text{PM}_{10}$  was  
23 the highest, with maxima during the night, peaking at 67% of total  $\text{PM}_{10}$  mass on 17 June.

24 Table 1 lists the concentrations of the aerosol chemical constituents, separately for  $\text{PM}_{1.2}$  and  $\text{PM}_{1.2-10}$  fractions and for  
25 day and night samples. All the chemical species concentrations were higher at the urban compared to the rural site but  
26 with small differences in most of the cases. The significantly higher concentrations of fine TC in BO can result from a  
27 higher contribution of elemental carbon (EC) and organic carbon (OC) from vehicular traffic at the urban site. Coarse-  
28 mode  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  also occurred at higher concentrations at the urban site, suggesting a source from road dust  
29 resuspension for these mineral elements. The only exception was represented by the nocturnal concentrations of fine-  
30 mode nitrate, on average five times higher at SPC compared to BO, and so for the counter-ion ammonium.

31 The concentrations of many species were higher at night compared to daytime (Tab. 1). The nocturnal enhancement can  
32 be due either to the accumulation of pollutants in a shallow boundary layer or to enhanced formation/emission of  
33 specific aerosol species at night-time. Fig. 4 shows the average  $\text{PM}_{10}$  composition at the urban and at the rural site  
34 separately for day and night, with indication of the percentage contributions of each species on the total mass of the  
35 measured compounds. The change in  $\text{PM}_{10}$  chemical composition between day and night at the rural site indicates that  
36 the enhanced mean concentrations found at night were not purely an effect of atmospheric dynamics but were impacted  
37 by chemical processes that led to the formation of specific aerosol compounds, especially ammonium nitrate, in the dark  
38 hours of the day. The lower average nocturnal temperatures (22°C versus 25°C), and the higher average nocturnal  
39 relative humidity (70% versus 48%) of SPC with respect to BO probably played an important role.

1 Water insoluble carbon (WINC) was calculated as the difference between TC and WSOC ( $WINC = TC - WSOC$ )  
2 (Matta et al., 2003). In order to obtain the mass of water-soluble organic material (WSOM) and water insoluble  
3 carbonaceous material (WINCM) conversion factors were applied to include elements different from carbon in the  
4 organic molecules. Two distinct factors were used respectively for the soluble and the insoluble fraction of organic  
5 carbon. WSOC was multiplied by 1.9 at the rural site and by 1.7 at the urban site, applying the two factors derived from  
6 AMS measurements performed at the two sites during the same campaign (Gilardoni et al., 2014). WINC was  
7 multiplied by 1.2 as found in the literature (Zappoli et al., 1999).

8 The total mass concentration of the chemical species determined on impactor samples averaged  $12.2 \mu\text{g m}^{-3}$  at the urban  
9 and  $8.2 \mu\text{g m}^{-3}$  at the rural site during daytime and 15.1 and 15.9 at night-time, respectively. It is interesting to note how  
10 in daytime the aerosol loading was higher at the urban compared to the rural site, indicating a higher contribution from  
11 urban sources, though the chemical composition was to some extent homogeneous. By contrast at night the aerosol  
12 mass was similar at the two sites but the chemical composition was different, with an enrichment of ammonium nitrate  
13 at the rural site, probably also favoured by the high ammonia concentrations from agricultural sources, which during the  
14 campaign were measured only at the rural site (Sullivan et al., 2015)

15 The measured mass of  $PM_{10}$  at the urban site consisted on average of 42% and 40% secondary inorganic aerosol (SIA)  
16 in daytime and night-time respectively. At the rural site, SIA represented 46% in daytime and 50% at night, due to a  
17 higher contribution from fine-mode ammonium nitrate. Carbonaceous matter was the dominant fraction at the urban site  
18 with 48% and 49% of the measured mass in daytime and night-time, respectively, and with the soluble fraction  
19 accounting for 26% and 27% of the mass in the two cases.

20 Finally, it is worth to remind that the above results are sensitive to sampling artifacts which can affect aerosol collection  
21 with low-pressure impactors (e.g., evaporative losses of semivolatile compounds). Figures S8 and S9 report the  
22 comparison between SIA measurements with the Berner impactor at SPC and co-located measurements using a  
23 different off-line system (a HiVol sampler) and an on-line method (HR-ToF-AMS). The results show that the Berner  
24 impactor observations are generally in line with the parallel measurements and especially that the main features of the  
25 time trend (e.g., the sharp diurnal variations in nitrate concentrations) are reproduced by all the instruments.

26  
27

### 28 **3.3 Size-resolved aerosol composition**

29 The time-series of size-resolved sulfate, nitrate and WSOC concentrations are shown in Fig. 5 (similar plots for  
30 additional chemical components and relative humidity are reported in the Supplementary Material, Figs. S1-S7). The  
31 figure highlights significant differences between the urban and the rural site in the formation of secondary inorganic and  
32 organic aerosol, especially during the two periods of stagnant conditions, i.e. 16-19 June and 5-7 July, which favored  
33 the accumulation of aerosol compounds from local sources. During the first of such events, the sulfate concentrations  
34 increased in BO and SPC to a similar extent in daytime, while higher concentrations were measured in SPC at night.  
35 Fine-mode nitrate, consisting primarily in ammonium nitrate, was virtually absent during the day throughout the  
36 campaign, particularly in BO, as a consequence of the high summer temperatures, which favored the thermal  
37 decomposition of  $NH_4NO_3$  into gas phase ammonia and nitric acid. Higher fine-mode nitrate concentrations were  
38 instead measured in SPC, reaching high levels at night. For both sulfate and nitrate the most prominent enhancements

1 affected the accumulation mode, i.e. size bins 2 (0.14 – 0.42  $\mu\text{m}$ ) and especially 3 (0.42 – 1.2  $\mu\text{m}$ ) of the impactor,  
2 which correspond to small and large accumulation mode.

3 Aerosol WSOC exhibited higher concentrations under stagnant conditions similarly to SIAs. During the first stagnation  
4 event, in particular, higher nocturnal concentrations in the accumulation mode were measured in SPC compared to BO  
5 (Fig. 5). Conversely, WSOC concentrations were in general higher in BO than in SPC in the quasi-ultrafine (0.05 – 0.14  
6  $\mu\text{m}$ ) fraction, with an average concentration of 0.39  $\mu\text{g m}^{-3}$  (17% of the total WSOC in the  $\text{PM}_{10}$  fraction) compared to  
7 0.19  $\mu\text{g m}^{-3}$  (11% of total WSOC in  $\text{PM}_{10}$ ). Since secondary formation is believed to represent the major source of  
8 WSOC in the absence of biomass burning (Weber et al., 2007), the quasi-ultrafine WSOC excess at the urban site could  
9 result from the increased condensation of secondary products on the large surface area of a higher number of very small  
10 particles found at the urban site. In addition to freshly nucleated particles, in fact, this aerosol fraction, twice as high in  
11 number at urban sites compared to rural areas (Westerdahl et al., 2005), includes particles released by anthropogenic  
12 sources such as combustion emissions from vehicular traffic. In addition, this quasi-ultrafine mode excess of WSOC  
13 mass could also reflect a direct contribution from anthropogenic primary emissions (Zhang et al., 2012).

14 On 5 and 6 July, the Po Valley was again influenced by stagnant conditions, with low wind speeds and short air-mass  
15 trajectories. However, the stagnation event in early July was considerably shorter than that of 16–19 June. During such  
16 event, high concentrations of fine-mode sulfate were simultaneously observed in BO and SPC with the 6 July  
17 experiencing the highest diurnal concentration of the whole campaign (4.9  $\mu\text{g m}^{-3}$  in  $\text{PM}_{10}$  in BO). Nitrate similarly  
18 increased at both sites at night, persisting in daytime in BO only, and reaching the diurnal maximum of 4.2  $\mu\text{g m}^{-3}$  on  
19  $\text{PM}_{10}$ . Similarly, WSOC increased at both sites, though more significantly in BO, where it reached the highest average  
20 diurnal concentration on 6 July. This day was characterized by the highest daytime relative humidity (47.8% and 60.4%  
21 in BO and SPC respectively, averaged over the time span of Berner impactor samplings) and a relatively lower  
22 temperature (27°C at both sites). Clouds were present on the sampling sites and a light rain was recorded in the  
23 afternoon. Such conditions apparently favored the formation of secondary organic (WSOC) and inorganic aerosol  
24 compounds in the Po Valley basin.

25 Fig. 6 shows the size-segregated concentrations of sulfate, nitrate, ammonium and WSOC during one day characterized  
26 by background conditions (15 June) and one day characterized by stagnant conditions (18/6), separately for diurnal and  
27 nocturnal hours. On the background day (panels a and c), fine particles in BO exhibited a maximum in the speciated  
28 aerosol mass in the small accumulation mode (0.14 – 0.42 $\mu\text{m}$ ), with a relative increase in the large accumulation mode  
29 (0.42 – 1.2 $\mu\text{m}$ ) at night. The speciated aerosol mass in SPC was almost evenly distributed between the two modes at  
30 night and day. Nitrate was always present in the coarse mode, where non-volatile nitrate salts can form through reaction  
31 of gaseous nitric acid with alkaline soil particles or resuspended dust (Harrison and Pio, 1983; Laskin et al., 2005).  
32 Under stagnant conditions (panels b and d) the speciated particle mass concentrations increased but the peak in the size  
33 distribution, which for BO was again observed in the small accumulation mode, shifted to the large accumulation mode  
34 in SPC both in daytime and at night. The nocturnal increase of large accumulation mode particulate matter  
35 concentration was much more evident at the rural site (SPC) than at the urban site (BO), mostly because of an increase  
36 of ammonium nitrate (showing more than 3 times higher concentrations with respect to background conditions), but  
37 also accompanied by increases in WSOC and sulfate concentrations. The SIA mass (i.e. the sum of sulfate, nitrate and  
38 ammonium) reached 67% of the total  $\text{PM}_{1.2}$  mass in the night of 16 June, during the first event of stagnation. The  
39 increase of SIA (and, to a lesser extent, of WSOC) in large accumulation mode aerosols under stagnant conditions was

1 therefore the main process modulating the impactor size distribution of submicron aerosol during the experiment. Our  
2 data seem to exclude that this shift in diameter of SIA-containing aerosols from the small to the large accumulation  
3 mode size range is due to impactor sampling artifacts (see Supplementary material SI-1, Figure S10).

4

### 5 **3.4 Secondary aerosol formation under stagnant conditions**

#### 6 **3.4.1 Secondary inorganic components**

7 The processes responsible for the accumulation of sulfate, nitrate and secondary organic aerosol (SOA) on stagnation  
8 days were further investigated by analyzing the relationship between the (size-segregated) concentrations of secondary  
9 inorganic compounds and relative humidity (RH), as well as with aerosol liquid water content (ALWC). SPC was  
10 characterized by higher levels of RH than Bologna, especially at night. During the campaign RH and temperature  
11 profiles at the two sites showed substantial day-night variations as a consequence of diabatic processes at the surface  
12 (nocturnal radiative cooling versus daytime heating from solar irradiation). As a consequence, the highest RH occurred  
13 at both sites just before dawn, from 4 to 5 am, and the lowest in the afternoon from 1 to 3 pm. During the diurnal  
14 sampling periods (from 9am to 9pm) hourly RH ranged from 15 to 66% in BO and from 19% to 73% in SPC, while  
15 during the nocturnal sampling periods (from 9pm to 9am) RH ranged from 24% to 82% in BO and from 35% to 92% in  
16 SPC. A plot of RH averaged on the time intervals of the Berner Impactor is reported in the Supplementary Material  
17 (Fig. S7).

18 Since the equilibrium constant for the reaction of  $\text{NH}_4\text{NO}_3$  formation is both RH and temperature dependent (Stelson  
19 and Seinfeld, 1982), thermodynamic conditions were more favorable in SPC than in BO for the existence of condensed  
20 phase nitrate. Both temperature and RH affect the equilibrium of  $\text{NH}_4\text{NO}_3$ , but the changes in RH were more marked.  
21 During the campaign the hourly nocturnal temperatures and corresponding relative humidity of deliquescence (RHD)  
22 for ammonium nitrate and ammonium sulfate (in brackets) ranged from 12°C (69.3% for ammonium nitrate and 80.7%  
23 for ammonium sulfate) to 30°C (58.6% for ammonium nitrate and 79.5 for ammonium sulfate) in SPC and from 16°C  
24 (67.1% for ammonium nitrate and 80.3 for ammonium sulfate) to 30°C in BO (Watson et al. (1994); E-AIM model,  
25 <http://www.aim.env.uea.ac.uk/aim/model2/mod2rhwh.php>). Nocturnal periods of time with RH greater than the RHD of  
26 hygroscopic salts (Watson et al., 1994)) were considerably longer in SPC than in BO, and therefore deliquesced  
27 particles were common at the rural and very rare at the urban site. This was confirmed by the simultaneous  
28 enhancements of the aerosol liquid water content (ALWC), calculated from hourly averaged AMS data at the two sites  
29 (see the experimental section) by the E-AIM model (Fig. 7). While the ALWC in SPC showed a consistent average  
30 diurnal trend, with a maximum before dawn when both RH and the concentrations of hygroscopic salts were the highest  
31 (Hodas et al., 2014), the diurnal variation in Bologna was less marked, and night-time ALWC concentrations were one  
32 order of magnitude smaller than at the rural station (Fig. 7). Considering the Berner impactor sampling periods, ALWC  
33 averaged  $0.33 \mu\text{g m}^{-3}$  in daytime in BO (min 0, max  $11.2 \mu\text{g m}^{-3}$ ) and  $0.76 \mu\text{g m}^{-3}$  at night (min 0, max  $16.6 \mu\text{g m}^{-3}$ ),  
34 while in SPC it averaged  $0.21 \mu\text{g m}^{-3}$  during day (min 0, max  $10.4 \mu\text{g m}^{-3}$ ), and  $6.55 \mu\text{g m}^{-3}$  during night conditions  
35 (min 0, max  $59.3 \mu\text{g m}^{-3}$ ). The 12-h averaged ALWC data show concentrations increasing rapidly for RH above 60%  
36 (Figure 8). There is a considerable variability in ALWC levels that must be attributed to the availability of hygroscopic  
37 material in the aerosol, as particulate water in subsaturated condition is a function not only of RH but also of the molar  
38 concentration of dissolved material in the aerosol.

1 Figure 8 (panels b,c) shows the relation between nitrate and sulfate concentrations separately in the small accumulation  
2 mode (condensation mode, CD) and in the large accumulation mode (droplet mode, DL) versus ALWC for the two  
3 sites. The correlation between particulate nitrate and ALWC was strong for both modes in SPC where ALWC levels  
4 above  $1 \mu\text{g m}^{-3}$  were frequent.

5 Our data are in agreement with the findings of Hodas et al., (2014) indicating that particulate nitrate was the primary  
6 driver of ALWC observed at night in the rural Po Valley in summertime. The same study suggested that ALWC  
7 enhanced the particle phase partitioning of water soluble organic gases and provided a medium for aqueous phase  
8 organic reactions that can form SOA. More generally, ALWC can favor – and sometimes can be necessary for – the  
9 formation of SOA and SIA in large accumulation mode aerosols (the droplet mode). Figure 8c shows that no  
10 relationship was observed between sulfate and the local estimates of ALWC at either site, with the exception of a  
11 moderate positive correlation for the droplet mode in SPC but not in BO. These results indicate that most sulfate was  
12 formed at the regional scale during this study (through cloud chemistry in the larger accumulation mode or gas phase  
13 chemistry in the smaller accumulation mode). However a small but non-zero increment of droplet-mode sulfate  
14 concentration in SPC can be attributed to the greater ALWC characterizing the rural site with respect to the urban site.

15 As a final remark, in this discussion, we attributed the high concentrations of ammonium nitrate at the rural site to the  
16 more favorable thermodynamic conditions respect to the urban site. This process was amplified by a positive feedback  
17 of ammonium nitrate itself that, by increasing the ALWC in the aerosol phase, further promotes the uptake of  
18 precursors ( $\text{NH}_3$ ,  $\text{HNO}_3$ ,  $\text{N}_2\text{O}_5$ ) from the gas-phase. We do not consider here the effect of the different distribution of  
19 gaseous precursors of ammonium nitrate in the Po Valley, especially of ammonia which is enriched in the rural areas  
20 (Sullivan et al., 2015), which will be object of a future study.

21

### 22 3.4.2 Carbonaceous species

23 The distribution of the carbonaceous components of aerosol (WSOC,  $\text{WINC} = \text{TC} - \text{WSOC}$ ) over the Berner impactor  
24 size intervals, with emphasis on accumulation mode particles (condensation and droplet modes) is hereby discussed.  
25 The dependence of WSOC on aerosol liquid water content (ALWC) is also investigated analogously to the case of SIA  
26 treated in the previous section.

27  $\text{PM}_{10}$  WSOC concentrations in this study ranged from  $0.13$  to  $4.6 \mu\text{g C m}^{-3}$  (average:  $2.1 \mu\text{g C m}^{-3}$ ) in BO and from  $0.28$   
28 to  $5.2 \mu\text{g C m}^{-3}$  (average:  $1.6 \mu\text{g C m}^{-3}$ ) in SPC. On average, 79% and 77% of this  $\text{PM}_{10}$  WSOC concentration was in  
29 the fine ( $\text{PM}_{1.2}$ ) fraction in BO and SPC respectively, with only slightly higher contributions at night than in daytime at  
30 both sites.

31 The water-soluble fraction of total carbon (WSOC/TC) in  $\text{PM}_{1.2}$  was 52% on average in BO and 61% in SPC. The  
32 WSOC fraction in SPC was comparable to what observed at other rural sites, e.g. 57% in  $\text{PM}_1$  in K-pusztá (Hungary)  
33 during the summer (Krivacsy et al., 2001). The greater water-soluble fraction of carbon found at SPC with respect to  
34 BO is in line with literature results, showing higher WSOC fractions in rural areas as a consequence of the concurrent  
35 higher input of SOA and the reduced fraction of insoluble carbonaceous particles from traffic sources (Weber et al.,  
36 2007).

1 The left panels in Fig. 9 show the linear regressions of WSOC versus TC in submicron particles (impactor stages 1 to 3)  
2 at the two sites, separately for day and night conditions. At the urban site, a higher slope was observed in daytime ( $0.71$   
3  $\pm 0.10$  at 95% confidence level) than at night-time ( $0.57 \pm 0.08$ ), pointing to the effect of a daytime source for WSOC  
4 (consistent with photochemical SOA formation). In daytime, the WSOC fraction of TC in BO overlaps well with that  
5 observed in SPC ( $0.72 \pm 0.05$ ). Contrary to BO, however, WSOC fractions in SPC were similar between night and day  
6 ( $0.70 \pm 0.07$  at night). Fig. 9 shows, in fact, that WSOC and TC occurred in similar proportions in daytime between  
7 SPC and BO, but with smaller concentrations in SPC (which were therefore “diluted” with respect to BO). In addition,  
8 carbonaceous aerosol concentrations increased at night in SPC and in similar proportions between WSOC and WINC,  
9 hence producing an aerosol with different characteristics in SPC with respect to BO.

10 The correlation of WSOC with a non-volatile SIA component (sulfate) in  $PM_{1.2}$  is shown in the right panels of Fig.9. A  
11 good correlation ( $R^2 = 0.7$ ) was observed at both stations in daytime, suggesting that WSOC shared a photochemical  
12 source with sulfate. The correlation in BO was much smaller at night ( $R^2 = 0.4$ ) than in daytime, which is expected,  
13 because particulate organic compounds have multiple sources other than photochemistry. Interestingly, the correlation  
14 between WSOC and sulfate remained high (0.7) at night in SPC, pointing to a common nocturnal source for WSOC and  
15 sulfate at the rural site.

16 Fig. 10 shows, for the two sites, the size-resolved concentration time series of WSOC and WINC for quasi-ultrafine  
17 mode ( $0.05 - 0.14 \mu m$ ), condensation mode ( $0.14 - 0.42 \mu m$ ) and droplet mode ( $0.42 - 1.2 \mu m$ ) particles. The quasi-  
18 ultrafine fraction provided the smallest contribution to aerosol mass, but with significantly higher concentrations at the  
19 urban compared to the rural site for both WSOC and WINC. This feature was observed also in previous studies (Sardar  
20 et al., 2005; Snyder et al., 2010; Zhang et al., 2012). The urban excess of WINC witnesses the effect of local emissions  
21 of insoluble primary carbonaceous particles. On average WSOC accounted for 52% of quasi-ultrafine TC in SPC and  
22 only 42% in BO, with the lower WSOC fraction at the urban site caused by the higher concentrations of water insoluble  
23 carbon. An urban increment for WSOC in quasi-ultrafine particles can be observed, although smaller than for WINC,  
24 and can be explained by local sources of fresh SOA and by condensation on a greater number of ultrafine particles.  
25 WINC in this size range displayed nocturnal excess compared to WSOC, particularly in BO, without a clear relation  
26 with the trajectory lengths. We argue that the nocturnal peaks in quasi-ultrafine WINC concentrations could be related  
27 to the morning traffic rush hours, which, during the summer, had a maximum at 8-9 am, before the break-up of the  
28 nocturnal boundary layer and were therefore included in the nocturnal sampling periods. The evening traffic rush time  
29 between 7 and 8 pm had apparently a minor effect, since occurring while the boundary layer was still well mixed.

30 The distribution of carbonaceous fractions in the accumulation mode (condensation and droplet modes) showed that  
31 WSOC was in general dominant over WINC, more at the rural than at the urban site. The average WSOC/TC ratio in  
32 the condensation mode was 65% in SPC and 60% in BO, while showing a greater difference between the two stations in  
33 the droplet mode (66% in SPC and 56% in BO). The WSOC concentrations were inversely related to trajectory length,  
34 indicating an effect of stagnation on oxidized organic aerosol production. Interestingly, the increase of WSOC levels  
35 during the stagnation periods did not affect the same size fractions to the same extent at the two sites. During the first  
36 episode, the maximum daytime WSOC concentration in the condensation mode, recorded on 19 June, corresponded to a  
37 185% increase in BO and 150% in SPC compared to 15 June, the last day under the influence of North Atlantic  
38 circulation before the onset of the local recirculation. The enhancement of WSOC in daytime for droplet mode particles  
39 amounted to 150% in BO and 195% in SPC. Therefore, the accumulation of water-soluble organic compounds occurred

1 approximately to the same extent at the two sites in daytime, but the increase was more marked in droplet mode  
2 particles in SPC and in the condensation mode in BO. The maximum nocturnal WSOC concentration, found on June 18,  
3 corresponded to an enhancement in the condensation mode of 140% and 325% in BO and SPC, respectively, with  
4 respect to the background conditions of 15 June. The same increase in the droplet mode amounted to 115% (BO) and  
5 440% (SPC). Therefore, the change in WSOC concentrations between background and stagnating conditions was more  
6 heterogeneous between sites for nocturnal samples than for the diurnal ones, which is expected because the atmosphere  
7 is much more stratified at night and atmospheric composition at ground level is more impacted by local conditions. In  
8 this case, a nocturnal enhancement of WSOC concentrations during the first stagnation period occurred only in SPC,  
9 with a maximum in the droplet mode.

10 During the second stagnant period, on 5-6 July, WSOC increased considerably in daytime in BO in both size ranges,  
11 while only a small increase was observed in SPC and limited to the droplet mode.

12 The behavior of accumulation mode WSOC after the onset of stagnating conditions was therefore reversed during the  
13 (short) July event with respect to the first episode of 16 – 20 June, with a marked increase in the droplet mode occurring  
14 in BO in July while interesting the SPC site in June (especially at night). The days (and nights) of maximum increase of  
15 droplet mode WSOC were in fact those showing the highest submicron nitrate concentrations, and were always humid  
16 days (or nights). Fig. 8d shows that WSOC was positively correlated with ALWC only in the droplet mode and only in  
17 SPC. The lack of correlation for BO samples can be explained by the very short duration of the humid stagnation period  
18 in July. Clearly, the increase of WSOC in droplet mode aerosols in the stagnation periods was not homogeneous in the  
19 Po Valley and was associated locally to the presence of deliquesced particles. These findings indicate that the  
20 enrichment of WSOC was contributed by aqueous processes, including condensation in the aerosol liquid water, which  
21 were active preferentially during colder night-time hours. Fig. 8 c,d also shows that the behavior of WSOC reflects that  
22 of sulfate in SPC for droplet mode aerosols, which explains the good correlation between WSOC and sulfate for night-  
23 time samples observed only at the rural site where ALWC was high (Fig. 9 right panels).

24

### 25 **3.5 Principal component analysis**

26 In the previous section, we focused on the time trends and size distributions of major carbonaceous and inorganic ionic  
27 species and we concluded that at least two secondary formation processes were active in the Po Valley: a first, probably  
28 photochemical, active throughout the campaign at both stations affecting the concentrations of all species and  
29 particularly in the condensation mode during stagnation periods; and a second one associated with deliquesced particles,  
30 and selectively important for nitrate (both condensation and droplet mode) and to a lesser extent for sulfate and WSOC  
31 in droplet mode particles. In this section, we will extract source information from all the chemical dataset and from  
32 ancillary information. Principal component analysis (PCA) was used to analyse the variability of the main variables of  
33 interest (the concentration of main SIAs and WSOC) in conjunction with the variability of the concentrations of minor  
34 species and tracers as well as of the physical parameters of the atmosphere. Six principal components were retained for  
35 interpretation of the SPC and BO datasets, explaining respectively 79% and 77% of the total variance. Increasing the  
36 number of factors progressively rises the explained variance, but each additional factor contributes only a small  
37 fractional increase and interpretation of the additional factors is also challenging. For the above reasons, a 6-factor  
38 solution was chosen as the best one. The results of the PCA are summarized in Table 2 as factor loadings, which

1 represent the correlation of each variable with each factor, and hence suggest possible sources, formation mechanisms  
2 and source regions. For better clarity, loadings with absolute values below 0.2 ( $|x| \leq 0.2$ ) were omitted and only those  
3 with absolute values larger than 0.6 ( $|x| \geq 0.6$ ) were considered “high” (in bold in the table).

4 The first rotated component (RC1) explained the largest fraction of the dataset variance (22%) in SPC. It involved high  
5 loadings of nitrate and ammonium for all the size classes. Among the meteorological parameters, the factor was  
6 strongly positively correlated with RH (0.82), and negatively with temperature (-0.85). Thus RC1 describes the local  
7 meteorology and the night-time condensation of ammonium nitrate in large accumulation mode particles (0.42 – 1.2  
8  $\mu\text{m}$ ). Ammonium nitrate condensation in SPC was moderately positively correlated with droplet mode sulfate, WSOC  
9 and oxalate, compounds (sulfate and oxalate) that share a source in aqueous secondary formation processes. A similar  
10 source was identified in the Bologna dataset in RC6 which, compared to SPC, explained only a smaller fraction (9%) of  
11 the total variance. The relationship with temperature and relative humidity was weaker in BO than in SPC, though in the  
12 same direction, and no clear relation was observed with droplet mode WSOC and sulfate

13 RC2 was the second most important factor in SPC, explaining 15% of the total variance in the dataset. Sulfate, oxalate,  
14 ammonium and WSOC were the species which correlated most with this factor, especially in the small (0.14 – 0.42  $\mu\text{m}$ )  
15 and large (0.42 – 1.2  $\mu\text{m}$ ) accumulation mode. This factor was positively correlated with the sunflux integrated along  
16 the airmass trajectory. This factor can therefore account for SIA and SOA photochemical production. An analogous  
17 factor in BO (RC2) explained a similar amount of total variance (17%). A positive correlation (0.6) with relative  
18 humidity was observed in BO, which was not significant in SPC. The significant contribution represented by this source  
19 on the total variance described for the two datasets highlights the importance of regional-scale secondary aerosol  
20 formation processes for the Po Valley environment.

21 The third factor in SPC is RC3, accounting for 14% of the total variance. This rotated component showed high loadings  
22 particularly for  $\text{Ca}^{2+}$  but also for  $\text{Mg}^{2+}$  in size bins 3 to 5, corresponding to particles from 0.42  $\mu\text{m}$  to 10  $\mu\text{m}$ . Oxalate  
23 also showed a significant loading in size bin 5 (3.5 – 10  $\mu\text{m}$ ), which suggested the uptake of gas-phase carboxylic acids  
24 by mineral particles, in agreement with past observations (Laongsri and Harrison, 2013; van Pinxteren et al., 2014).  
25 This factor is analogous to RC4 in the BO dataset, which explains 11% of the total variance, and was attributed to road  
26 dust resuspension. High loadings in BO were shifted toward larger diameters of particles (1.2 to 10  $\mu\text{m}$ ).

27 The fourth rotated component in SPC (RC4), which explained another 12% of the total variance, displayed the highest  
28 loadings for magnesium in coarse mode particles and sodium in size bins 3 to 5 (from 0.42 to 10  $\mu\text{m}$ ). Sulfate in size  
29 bin 5 (3.5 – 10  $\mu\text{m}$ ) was also moderately correlated with this factor, and nitrate too, but to a lesser extent. This factor  
30 can be interpreted as a contribution from seasalt components. The corresponding rotated component in BO was RC1,  
31 which explained 19% of the total variance and, compared to SPC, displayed higher loadings for sulfate and nitrate in  
32 size bin 4 (1.2 to 3.5  $\mu\text{m}$ ).

33 The fifth rotated component (RC5) in SPC, explaining 8% of the total variability, only included high loadings for  
34 residence times indexes (RTI), without any relevant correlation with other parameters. This factor was identified both in  
35 SPC and in BO (RC5), explaining respectively 8% and 10% of the total variance. While this factor does not represent  
36 an aerosol source, it indicates that during this campaign the impact of air mass history (long-range transport) was likely  
37 small as compared to other impacts such as day/night variability or local impacts.

1 Finally the last rotated component, which explained an additional 8% of the total variance in SPC (RC6) and 11% in  
2 BO (RC3), contained high loadings only in the first size range (0.05 – 0.14  $\mu\text{m}$ ) for sulfate, ammonium, and to a lesser  
3 extent WSOC. The significant loading of sulfate and ammonium in quasi-ultrafine particles, together with a moderate  
4 positive correlation with solar radiation and temperature, suggests a possible source for this component in ultrafine  
5 particle nucleation. The role of these chemical species in the formation of new particles and their  
6 condensation/coagulation on smaller particulate matter are well-known, as well as the influence of solar radiation on  
7 this aerosol generation process (Hamed et al., 2010). The negative relation with relative humidity confirmed that this  
8 source was active during the day, when relative humidity was at minimum.

9

## 10 **Discussion and conclusions**

11 The PCA results indicate that several factors determined the variability in the size-segregated chemical composition in  
12 the region during the PEGASOS Po Valley field campaign, but each of them affected preferentially specific size  
13 intervals with an overall effect of shaping the aerosol mass distribution at the two sites. Two factors corresponding to  
14 seasalt and mineral dust with absorbed nitrate regulated the concentrations and composition of coarse particles ( $\text{PM}_{1.2-10}$ ),  
15 while only one factor was found to determine an enrichment of ammonium sulfate in the quasi-ultrafine range.  
16 Finally, the variability in composition of accumulation mode aerosol could be reduced to two factors, with one related  
17 to regional-scale photochemical formation of SOA and SIA, and a second one more dependent on local conditions at  
18 surface level and causing a nocturnal increase of SOA and SIA in the droplet mode. However, if the factor for  
19 photochemical secondary aerosols was equally represented at the two sites, the other one (for night-time condensation)  
20 was much more important at the rural station (SPC) than in BO. The effect on a simple non-volatile SIA component,  
21 sulfate, is exemplified in the scatter plot in Fig. 11. A good correlation ( $R^2 = 0.9$ ) was indeed observed between the  
22 accumulation mode sulfate concentrations at the two sites in daytime (Fig. 11), with only a slight dilution (-12%) of the  
23 concentrations at the background site with respect to the urban site, which is expected for an aerosol component which  
24 is typically associated to regional-scale photochemical pollution. By contrast, the correlation between the concentration  
25 trends at the two sites is much lower at night, especially as regards the droplet mode, where significantly higher sulfate  
26 concentrations occurred in SPC compared to BO during the stagnant days from 16 to 20 June. Clearly, stagnating  
27 conditions and the onset of thermal inversions at night favored a partial “chemical segregation” of air masses in the  
28 surface atmospheric layers within the Po Valley, and the size-segregated chemical composition evolved separately at  
29 the urban sites close to the Apennine foothills with respect to the rural areas in the inner Po Valley during dark hours.  
30 Specifically, rural areas were characterized by the presence of ammonium nitrate and by ALWC levels above  $10 \mu\text{g m}^{-3}$ .  
31 Deliquesced aerosols could host aqueous phase formation of sulfate (via reaction of  $\text{SO}_2$  with  $\text{H}_2\text{O}_2$ ).

32 Similarly to SIAs in the droplet mode, water-soluble products of volatile organic compound (VOC) oxidation could  
33 readily be taken up by deliquesced particles in SPC at night. A meaningful fraction of the newly formed (1–3 h old)  
34 WSOC mass, in fact, has been shown to possess similar semi-volatile properties to  $\text{NH}_4\text{NO}_3$  (Hennigan et al., 2008;  
35 Wilson et al., 2006) and can rapidly partition to aerosol water or cloud/fog droplets. The nature of the nocturnal  
36 enrichment of WSOC in the droplet mode, depending on the reactivity in the aqueous phase, can be described by either  
37 a reversible mechanism (condensation of water soluble organic compounds triggered by the change in RH and ALWC)  
38 or an irreversible reaction (oxidation of VOCs or OVOCs with production of stable compounds). Hodas et al. (2014),

1 based on measurements performed during the same campaign in SPC, observed an exponential decrease in gas phase  
2 glyoxal concentrations with increasing ALW, and a local nocturnal production of aqueous SOA was indeed observed by  
3 parallel near real-time WSOC sampled with a Particle into Liquid Sampler (PILS) at the same site (Sullivan et al.,  
4 2015). The analysis of impactor samples provides only a few clues to disentangle the two effects. The only two organic  
5 markers for SOA that were quantitatively determined in all samples were oxalate and methanesulfonate (MSA). The  
6 robust correlation between WSOC, oxalate and sulfate, both in daytime and at night-time in SPC, indicates that the  
7 accumulation of particulate polar organic compounds contributed to the (irreversible) production of stable (oxidized)  
8 species. Oxalate is generally known to share with sulfate an important aqueous-phase oxidation pathway (Sorooshian et  
9 al., 2006). MSA is a more specific marker than oxalate, being related to the atmospheric processing of dimethylsulfide  
10 (DMS) whose emissions are unevenly distributed on the Earth surface, and can be intense in biogenically-rich marine  
11 waters. Our data showed indeed that fine-mode MSA was maximum in the days between 26 June and 1 July,  
12 characterized by an easterly or a south-southwesterly circulation, bringing marine air masses into the Po Valley basin.  
13 Such increase from long-range transport affected particularly the size intervals 0.05 – 0.14  $\mu\text{m}$  and 0.14 – 0.42  $\mu\text{m}$ .  
14 MSA concentrations in the droplet mode (0.42 – 1.2  $\mu\text{m}$ ) showed instead an enhancement at night under stagnant  
15 conditions, similarly to ammonium nitrate (see Supplementary Material), particularly marked in SPC during both  
16 episodes and in BO only during 5-6 July. The increase of MSA in droplet mode particles under stagnating conditions  
17 points to a DMS (or other reduced sulfur species) source other than from the marine boundary layer. In inland areas  
18 DMS has sometimes been reported as dominantly from terrestrial sources (vegetation and soils) and anthropogenic  
19 sources (manure and livestock), with higher temperatures and solar radiation enhancing its emission (Perraud et al.,  
20 2015). These findings suggest that the VOCs participating to the formation of WSOC in the Po Valley also included  
21 organic compounds emitted by agricultural activities or even by natural sources, and that ALWC in the atmospheric  
22 nocturnal surface layer acted as a medium for their formation during summer time.

23 In conclusion, the characteristics of the size-segregated aerosol compositions and its variability at a rural and an urban  
24 background site in the Po Valley could be explained by a limited number of factors reflecting main physico-chemical  
25 processes and/or transport patterns in the atmosphere. For accumulation mode particles in particular, our analysis points  
26 to two main processes: (1) The photochemical SIA and SOA which occur at comparable concentrations between BO  
27 and SPC and are particularly evident in daytime hours when the lower atmosphere is well mixed, and indicating that a  
28 major fraction of background submicron aerosol concentrations in the Po Valley actually originates from regional-scale  
29 sources, which can extend over vast continental areas (see also Fig. S12 in Decesari et al. (2014)). This has implications  
30 for air quality mitigation, because this photochemical component is expected to show little sensitivity to local-scale  
31 (city-level) regulations. (2) Nocturnal SIA and SOA whose formation is enhanced in the shallow, cool and humid  
32 boundary layer and mediated by the presence of aerosol liquid water. Such component of the rural background aerosol  
33 appears more volatile (hence labile) and more heterogeneously distributed across the Po Valley, with the inner part  
34 (where most agricultural activity reside) acting as a source region respect to its southern periphery (more urbanized).  
35 The rural background concentration level is therefore variable, somewhat “tilted” horizontally across the Valley, at least  
36 for half of the day. These results represent an example of a limitation of the classical Lenschow model.

37

38

39

## 1 Acknowledgements

2 This research was conducted as part of the “Supersito” Project, supported by Emilia Romagna Region and Regional  
3 Agency for Prevention and Environment (ARPA Emilia Romagna) under Deliberation Regional Government n. 428/10.  
4 The work was also made possible by the European Commission under the Framework Programme 7 (FP7) projects  
5 PEGASOS (Grant Agreement 265148), BACCHUS (Grant Agreement 603445) and by the CNR Joint Lab Project Air-  
6 Sea Lab, which are highly acknowledged.

7

## 8 References

- 9 Bucci, S., F. Cairo, P. Cristofanelli, S. Decesari, J. Groess, and F. Fierli, in preparation, Transport regimes  
10 analysis over the Po Valley during summer 2012: impacts on Planetary Boundary Layer variability  
11 and aerosol content. In preparation.
- 12 Carbone, C., S. Decesari, M. Mircea, L. Giulianelli, E. Finessi, M. Rinaldi, S. Fuzzi, A. Marinoni, R. Duchi, C.  
13 Perrino, T. Sargolini, M. Varde, F. Sprovieri, G. P. Gobbi, F. Angelini, and M. C. Facchini, 2010, Size-  
14 resolved aerosol chemical composition over the Italian Peninsula during typical summer and winter  
15 conditions: *Atmospheric Environment*, v. 44, p. 5269-5278.
- 16 Clegg, S. L., P. Brimblecombe, and A. S. Wexler, 1998, Thermodynamic model of the system  $H^+-NH_4^+-SO_4^{2-}$ -  
17  $NO_3^-H_2O$  at tropospheric temperatures: *Journal of Physical Chemistry A*, v. 102, p. 2137-2154.
- 18 Crosier, J., J. D. Allan, H. Coe, K. N. Bower, P. Formenti, and P. I. Williams, 2007, Chemical composition of  
19 summertime aerosol in the Po Valley (Italy), northern Adriatic and Black Sea: *Quarterly Journal of*  
20 *the Royal Meteorological Society*, v. 133, p. 61-75.
- 21 Decesari, S., J. Allan, C. Plass-Duelmer, B. J. Williams, M. Paglione, M. C. Facchini, C. O'Dowd, R. M. Harrison,  
22 J. K. Gietl, H. Coe, L. Giulianelli, G. P. Gobbi, C. Lanconelli, C. Carbone, D. Worsnop, A. T. Lambe, A.  
23 T. Ahern, F. Moretti, E. Tagliavini, T. Elste, S. Gilge, Y. Zhang, and M. Dall'Osto, 2014, Measurements  
24 of the aerosol chemical composition and mixing state in the Po Valley using multiple spectroscopic  
25 techniques: *Atmospheric Chemistry and Physics*, v. 14, p. 12109-12132.
- 26 Decesari, S., M. C. Facchini, S. Sandrini, M. Paglione, S. Gilardoni, M. Rinaldi, P. Cristofanelli, F. Cairo, G. P.  
27 Gobbi, V. Poluzzi, A. Morgillo, G. Bonafè, T. Mentel, A. Kindler-Schaar, H. Manninen, L. Poulain, F.  
28 Wolf, U. Baltensperger, L. Ganzeveld, E. Nemitz, and S. Pandis, in preparation, The 2012 PEGASOS-  
29 SUPERSITO Po Valley campaign: an overview. In preparation.
- 30 Dorling, S. R., T. D. Davies, and C. E. Pierce, 1992, CLUSTER-ANALYSIS - A TECHNIQUE FOR ESTIMATING THE  
31 SYNOPTIC METEOROLOGICAL CONTROLS ON AIR AND PRECIPITATION CHEMISTRY - RESULTS FROM  
32 ESKDALEMUIR, SOUTH SCOTLAND: *Atmospheric Environment Part a-General Topics*, v. 26, p. 2583-  
33 2602.
- 34 Draxler, R. R., and G. D. Rolph, 2003, HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory)  
35 Model access via NOAA ARL READY Website (<http://www.arl.noaa.gov/ready/hysplit4.html>). NOAA  
36 Air Resources Laboratory, Silver Spring, MD. (2003).
- 37 Farnham, I. M., A. K. Singh, K. J. Stetzenbach, and K. H. Johannesson, 2002, Treatment of nondetects in  
38 multivariate analysis of groundwater geochemistry data: *Chemometrics and Intelligent Laboratory*  
39 *Systems*, v. 60, p. 265-281.
- 40 Gelencser, A., T. Meszaros, M. Blazso, G. Kiss, Z. Krivacsy, A. Molnar, and E. Meszaros, 2000, Structural  
41 characterisation of organic matter in fine tropospheric aerosol by pyrolysis-gas chromatography-  
42 mass spectrometry: *Journal of Atmospheric Chemistry*, v. 37, p. 173-183.
- 43 Gietl, J. K., T. Tritscher, and O. Klemm, 2008, Size-segregated analysis of PM10 at two sites, urban and rural,  
44 in Munster (Germany) using five-stage Berner type impactors: *Atmospheric Environment*, v. 42, p.  
45 5721-5727.
- 46 Gilardoni, S., P. Massoli, L. Giulianelli, M. Rinaldi, M. Paglione, F. Pollini, C. Lanconelli, V. Poluzzi, S. Carbone,  
47 R. Hillamo, L. M. Russell, M. C. Facchini, and S. Fuzzi, 2014, Fog scavenging of organic and inorganic  
48 aerosol in the Po Valley: *Atmospheric Chemistry and Physics*, v. 14, p. 6967-6981.

1 Hamed, A., W. Birmili, J. Joutsensaari, S. Mikkonen, A. Asmi, B. Wehner, G. Spindler, A. Jaatinen, A.  
2 Wiedensohler, H. Korhonen, K. E. J. Lehtinen, and A. Laaksonen, 2010, Changes in the production  
3 rate of secondary aerosol particles in Central Europe in view of decreasing SO<sub>2</sub> emissions between  
4 1996 and 2006: *Atmospheric Chemistry and Physics*, v. 10, p. 1071-1091.

5 Harrison, R. M., and C. A. Pio, 1983, Size-differentiated composition of inorganic atmospheric aerosols of  
6 both marine and polluted continental origin: *Atmospheric Environment*, v. 17, p. 1733-1738.

7 Hennigan, C. J., M. H. Bergin, J. E. Dibb, and R. J. Weber, 2008, Enhanced secondary organic aerosol  
8 formation due to water uptake by fine particles: *Geophysical Research Letters*, v. 35.

9 Hering, S. V., and S. K. Friedlander, 1982, Origins of aerosol sulfur size distributions in the Los-Angeles Basin:  
10 *Atmospheric Environment*, v. 16, p. 2647-2656.

11 Hodas, N., A. P. Sullivan, K. Skog, F. N. Keutsch, J. L. Collett, S. Decesari, M. C. Facchini, A. G. Carlton, A.  
12 Laaksonen, and B. J. Turpin, 2014, Aerosol Liquid Water Driven by Anthropogenic Nitrate:  
13 Implications for Lifetimes of Water-Soluble Organic Gases and Potential for Secondary Organic  
14 Aerosol Formation: *Environmental Science & Technology*, v. 48, p. 11127-11136.

15 Jimenez, J. L., J. T. Jayne, Q. Shi, C. E. Kolb, D. R. Worsnop, I. Yourshaw, J. H. Seinfeld, R. C. Flagan, X. F.  
16 Zhang, K. A. Smith, J. W. Morris, and P. Davidovits, 2003, Ambient aerosol sampling using the  
17 Aerodyne Aerosol Mass Spectrometer: *Journal of Geophysical Research-Atmospheres*, v. 108.

18 John, W., S. M. Wall, J. L. Ondo, and W. Winklmayr, 1990, MODES IN THE SIZE DISTRIBUTIONS OF  
19 ATMOSPHERIC INORGANIC AEROSOL: *Atmospheric Environment Part a-General Topics*, v. 24, p.  
20 2349-2359.

21 Krivacsy, Z., A. Hoffer, Z. Sarvari, D. Temesi, U. Baltensperger, S. Nyeki, E. Weingartner, S. Kleefeld, and S. G.  
22 Jennings, 2001, Role of organic and black carbon in the chemical composition of atmospheric  
23 aerosol at European background sites: *Atmospheric Environment*, v. 35, p. 6231-6244.

24 Laongsri, B., and R. M. Harrison, 2013, Atmospheric behaviour of particulate oxalate at UK urban  
25 background and rural sites: *Atmospheric Environment*, v. 71, p. 319-326.

26 Laskin, A., T. W. Wietsma, B. J. Krueger, and V. H. Grassian, 2005, Heterogeneous chemistry of individual  
27 mineral dust particles with nitric acid: A combined CCSEM/EDX, ESEM, and ICP-MS study: *Journal of*  
28 *Geophysical Research-Atmospheres*, v. 110.

29 Lenschow, P., H. J. Abraham, K. Kutzner, M. Lutz, J. D. Preuss, and W. Reichenbacher, 2001, Some ideas  
30 about the sources of PM<sub>10</sub>: *Atmospheric Environment*, v. 35, p. S23-S33.

31 Matta, E., M. C. Facchini, S. Decesari, M. Mircea, F. Cavalli, S. Fuzzi, J. P. Putaud, and A. Dell'Acqua, 2003,  
32 Mass closure on the chemical species in size-segregated atmospheric aerosol collected in an urban  
33 area of the Po Valley, Italy: *Atmospheric Chemistry and Physics*, v. 3, p. 623-637.

34 Meng, Z. Y., and J. H. Seinfeld, 1994, ON THE SOURCE OF THE SUBMICROMETER DROPLET MODE OF URBAN  
35 AND REGIONAL AEROSOLS: *Aerosol Science and Technology*, v. 20, p. 253-265.

36 Perraud, V., J. R. Horne, A. S. Martinez, J. Kalinowski, S. Meinardi, M. L. Dawson, L. M. Wingen, D. Dabdub,  
37 D. R. Blake, R. B. Gerber, and B. J. Finlayson-Pitts, 2015, The future of airborne sulfur-containing  
38 particles in the absence of fossil fuel sulfur dioxide emissions: *Proceedings of the National Academy*  
39 *of Sciences of the United States of America*, v. 112, p. 13514-13519.

40 Putaud, J. P., R. Van Dingenen, A. Alastuey, H. Bauer, W. Birmili, J. Cyrus, H. Flentje, S. Fuzzi, R. Gehrig, H. C.  
41 Hansson, R. M. Harrison, H. Herrmann, R. Hitzemberger, C. Hueglin, A. M. Jones, A. Kasper-Giebl, G.  
42 Kiss, A. Kousa, T. A. J. Kuhlbusch, G. Loeschau, W. Maenhaut, A. Molnar, T. Moreno, J. Pekkanen, C.  
43 Perrino, M. Pitz, H. Puxbaum, X. Querol, S. Rodriguez, I. Salma, J. Schwarz, J. Smolik, J. Schneider, G.  
44 Spindler, H. ten Brink, J. Tursic, M. Viana, A. Wiedensohler, and F. Raes, 2010, A European aerosol  
45 phenomenology-3: Physical and chemical characteristics of particulate matter from 60 rural, urban,  
46 and kerbside sites across Europe: *Atmospheric Environment*, v. 44, p. 1308-1320.

47 Sardar, S. B., P. M. Fine, and C. Sioutas, 2005, Seasonal and spatial variability of the size-resolved chemical  
48 composition of particulate matter (PM<sub>10</sub>) in the Los Angeles Basin: *Journal of Geophysical*  
49 *Research-Atmospheres*, v. 110.

50 Seinfeld, J. H., and S. N. Pandis, eds., 1998, *Atmospheric Chemistry and Physics, from Air Pollution to*  
51 *Climate Change*: New York, John Wiley and Sons, 1326 p.

1 Snyder, D. C., A. P. Rutter, C. Worley, M. Olson, A. Plourde, R. C. Bader, T. Dallmann, and J. J. Schauer, 2010,  
2 Spatial variability of carbonaceous aerosols and associated source tracers in two cities in the  
3 Midwestern United States: *Atmospheric Environment*, v. 44, p. 1597-1608.

4 Sorooshian, A., V. Varutbangkul, F. J. Brechtel, B. Ervens, G. Feingold, R. Bahreini, S. M. Murphy, J. S.  
5 Holloway, E. L. Atlas, G. Buzorius, H. Jonsson, R. C. Flagan, and J. H. Seinfeld, 2006, Oxalic acid in  
6 clear and cloudy atmospheres: Analysis of data from International Consortium for Atmospheric  
7 Research on Transport and Transformation 2004: *Journal of Geophysical Research-Atmospheres*, v.  
8 111.

9 Stelson, A. W., and J. H. Seinfeld, 1982, RELATIVE-HUMIDITY AND TEMPERATURE-DEPENDENCE OF THE  
10 AMMONIUM-NITRATE DISSOCIATION-CONSTANT: *Atmospheric Environment*, v. 16, p. 983-992.

11 Sullivan, A. P., N. Hodas, B. J. Turpin, K. Skog, F. N. Keutsch, S. Gilardoni, M. Paglione, M. Rinaldi, S. Decesari,  
12 M. C. Facchini, L. Poulain, H. Herrmann, A. Wiedensholer, E. Nemitz, M. M. Twigg, and J. L. J. Collett,  
13 2015, Evidence for ambient dark aqueous SOA formation in the Po Valley, Italy: *Atmos. Chem. Phys.*  
14 *Discuss.*, v. 15, p. 36.

15 van Pinxteren, D., E. Brüeggemann, T. Gnauk, K. Mueller, C. Thiel, and H. Herrmann, 2010, A GIS based  
16 approach to back trajectory analysis for the source apportionment of aerosol constituents and its  
17 first application: *Journal of Atmospheric Chemistry*, v. 67, p. 1-28.

18 van Pinxteren, D., C. Neusuess, and H. Herrmann, 2014, On the abundance and source contributions of  
19 dicarboxylic acids in size-resolved aerosol particles at continental sites in central Europe:  
20 *Atmospheric Chemistry and Physics*, v. 14, p. 3913-3928.

21 Vandeginste, B. G. M., 1998, Data mining of water quality data by chemometrical methods: *Monitoring of*  
22 *Water Quality: the Contribution of Advanced Technologies*, p. 49-53.

23 Watson, J. G., J. C. Chow, F. W. Lurmann, and S. P. Musarra, 1994, AMMONIUM-NITRATE, NITRIC-ACID, AND  
24 AMMONIA EQUILIBRIUM IN WINTERTIME PHOENIX, ARIZONA: *Journal of the Air & Waste*  
25 *Management Association*, v. 44, p. 405-412.

26 Weber, R. J., A. P. Sullivan, R. E. Peltier, A. Russell, B. Yan, M. Zheng, J. de Gouw, C. Warneke, C. Brock, J. S.  
27 Holloway, E. L. Atlas, and E. Edgerton, 2007, A study of secondary organic aerosol formation in the  
28 anthropogenic-influenced southeastern United States: *Journal of Geophysical Research-*  
29 *Atmospheres*, v. 112.

30 Westerdahl, D., S. Fruin, T. Sax, P. M. Fine, and C. Sioutas, 2005, Mobile platform measurements of ultrafine  
31 particles and associated pollutant concentrations on freeways and residential streets in Los  
32 Angeles: *Atmospheric Environment*, v. 39, p. 3597-3610.

33 Wexler, A. S., and S. L. Clegg, 2002, Atmospheric aerosol models for systems including the ions H<sup>+</sup>, NH<sub>4</sub><sup>+</sup>,  
34 Na<sup>+</sup>, SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, Cl<sup>-</sup>, Br<sup>-</sup>, and H<sub>2</sub>O: *Journal of Geophysical Research-Atmospheres*, v. 107.

35 Wilson, W. E., B. D. Grover, R. W. Long, N. L. Eatough, and D. J. Eatough, 2006, The measurement of fine-  
36 particulate semivolatile material in urban aerosols: *Journal of the Air & Waste Management*  
37 *Association*, v. 56, p. 384-397.

38 Wolf, R., I. El Haddad, M. Crippa, S. Decesari, J. G. Slowik, L. Poulain, S. Gilardoni, M. Rinaldi, S. Carbone, F.  
39 Canonaco, R. J. Huang, U. Baltensperger, and A. S. H. Prevot, 2015, Marine and urban influences on  
40 summertime PM<sub>2.5</sub> aerosol in the Po basin using mobile measurements: *Atmospheric*  
41 *Environment*, v. 120, p. 447-454.

42 Zappoli, S., A. Andracchio, S. Fuzzi, M. C. Facchini, A. Gelencser, G. Kiss, Z. Krivacsy, A. Molnar, E. Meszaros,  
43 H. C. Hansson, K. Rosman, and Y. Zebuhr, 1999, Inorganic, organic and macromolecular  
44 components of fine aerosol in different areas of Europe in relation to their water solubility:  
45 *Atmospheric Environment*, v. 33, p. 2733-2743.

46 Zhang, X., Z. Liu, A. Hecobian, M. Zheng, N. H. Frank, S. Edgerton, and R. J. Weber, 2012, Spatial and  
47 seasonal variations of fine particle water-soluble organic carbon (WSOC) over the southeastern  
48 United States: implications for secondary organic aerosol formation: *Atmospheric Chemistry and*  
49 *Physics*, v. 12, p. 6593-6607.

50

51

1

2 **Table 1 – Fine (PM<sub>1,2</sub>), coarse (PM<sub>1,2-10</sub>) and PM<sub>10</sub> mean concentrations (in µg m<sup>-3</sup>, from integrated impactor**  
 3 **mass size-distributions) for main inorganic ions, water-soluble organic carbon (WSOC) and total carbon (TC)**  
 4 **separately for day (D) and night (N) samples during the PEGASOS summer campaign.**

5


	PM <sub>1,2</sub>				PM <sub>1,2-10</sub>				PM <sub>10</sub>			
	BO (urban)		SPC (rural)		BO (urban)		SPC (rural)		BO (urban)		SPC (rural)	
	D	N	D	N	D	N	D	N	D	N	D	N
<b>Sulfate</b>	2.1	2.6	1.7	2.3	0.27	0.32	0.16	0.28	2.3	2.8	1.8	2.6
<b>Nitrate</b>	0.24	0.47	0.32	2.40	1.6	1.7	0.85	1.5	1.8	2.1	1.2	3.9
<b>Chloride</b>	0.023	0.043	0.005	0.041	0.17	0.27	0.05	0.14	0.13	0.27	0.06	0.19
<b>Ammonium</b>	0.83	1.1	0.70	1.5	0.05	0.07	0.02	0.15	0.88	1.1	0.69	1.6
<b>Magnesium</b>	0.01	0.01	0.01	0.01	0.06	0.07	0.04	0.04	0.07	0.08	0.04	0.05
<b>Calcium</b>	0.07	0.07	0.06	0.07	0.50	0.48	0.35	0.41	0.56	0.53	0.40	0.47
<b>Sodium</b>	0.04	0.05	0.03	0.02	0.37	0.48	0.20	0.22	0.41	0.53	0.23	0.24
<b>WSOC</b>	1.4	1.9	0.94	1.6	0.54	0.53	0.28	0.44	1.9	2.4	1.2	2.1
<b>TC</b>	2.5	3.5	1.3	2.2	1.5	1.5	0.8	1.4	4.1	5.2	2.1	3.5

6

1 **Table 2 – Factor loadings of PCA after Varimax rotation. Only absolute values larger than 0.2 are shown.**  
 2 **Absolute values larger than 0.60 are considered significant and printed in bold. The number beside each**  
 3 **chemical species in the first column indicates the impactor stage.**  
 4


Site Rotated component Assigned source	SPC						BO					
	RC1	RC2	RC3	RC4	RC5	RC6	RC1	RC2	RC3	RC4	RC5	RC6
	conden sation	Photo chem. SIA+ SOA	dust	Sea salt	RTI var.	gasSIA+ gasSOA	Sea salt	Photo chem. SIA+ SOA	gasSIA+ gasSOA	resusp.	RTI var.	conden sation
Nitrate_1	<b>0.81</b>			-0.24	0.24							<b>0.69</b>
Sulfate_1	-0.32	0.28				<b>0.79</b>		0.26	<b>0.84</b>			
Oxalate_1	-0.34		0.25	-0.24	0.45	0.37	-0.53	0.24	0.42			
Ammonium_1		0.21				<b>0.84</b>			<b>0.79</b>		0.29	
WSOC_1		0.23	0.35	-0.4	0.29	0.46	-0.45	0.37	0.41	0.3	0.38	
Nitrate_2	<b>0.79</b>	0.36						0.28				<b>0.84</b>
Sulfate_2		<b>0.83</b>		0.3			0.42	<b>0.75</b>	0.22			
Oxalate_2	0.28	<b>0.81</b>	0.31	-0.22				<b>0.71</b>	0.43			0.34
Sodium_2				0.25	-0.23	0.33	0.43	-0.21	0.56	0.23	-0.29	
Ammonium_2	0.56	<b>0.69</b>				0.24	0.39	<b>0.77</b>				0.38
WSOC_2	0.3	<b>0.71</b>	0.46					<b>0.72</b>	0.43			0.3
Nitrate_3	<b>0.82</b>	0.4					0.21	0.36	-0.25			<b>0.78</b>
Sulfate_3	0.36	<b>0.79</b>					0.32	<b>0.86</b>				
Oxalate_3	0.45	<b>0.73</b>	0.32					<b>0.76</b>				
Sodium_3	-0.27		0.21	<b>0.84</b>			<b>0.87</b>					
Ammonium_3	<b>0.67</b>	<b>0.67</b>					0.29	<b>0.81</b>				0.32
Magnesium_3	-0.24	0.22	<b>0.8</b>	0.33			<b>0.62</b>		0.24	0.42		0.28
Calcium_3			<b>0.92</b>							0.5	-0.37	0.27
WSOC_3	0.5	<b>0.75</b>	0.32					<b>0.86</b>				0.23
Chloride_4	0.57			0.32		-0.29	0.51	0.22	-0.45			0.31
Nitrate_4	<b>0.82</b>			0.43			<b>0.79</b>	0.28		0.28		0.33
Sulfate_4	<b>0.69</b>	0.32		0.39	-0.23		<b>0.75</b>	0.45		0.24	-0.3	
Oxalate_4		0.5	0.35	0.38	-0.23		0.31	0.24	0.24	0.36	-0.37	
Sodium_4				<b>0.94</b>			<b>0.9</b>		-0.21			
Ammonium_4	<b>0.89</b>	0.27					0.34	0.3		0.24		<b>0.7</b>
Magnesium_4			0.57	<b>0.77</b>			<b>0.88</b>	0.21		0.29		
Calcium_4			<b>0.93</b>				0.41		<b>0.64</b>	-0.28		0.29
WSOC_4	0.47	0.31	0.52					<b>0.67</b>		0.42		-0.28
Chloride_5	0.57	0.29		0.23			0.47		<b>-0.66</b>	0.26		
Nitrate_5	0.56	0.37	0.37	0.46			0.54			<b>0.65</b>		
Sulfate_5	0.47		0.33	<b>0.64</b>	-0.3		<b>0.69</b>			<b>0.6</b>	-0.22	
Oxalate_5		0.29	<b>0.64</b>	0.21				0.29		0.55	-0.46	
Sodium_5		-0.23		<b>0.75</b>			<b>0.71</b>		-0.5	0.32		
Ammonium_5	<b>0.8</b>	0.29				0.26		0.23		0.54		0.5
Magnesium_5	0.29		<b>0.6</b>	<b>0.66</b>			<b>0.69</b>		-0.31	0.56		
Calcium_5			<b>0.87</b>			0.24	0.33			<b>0.89</b>		
WSOC_5	0.32	0.3	<b>0.65</b>	0.21				0.48		0.55		-0.27
RT_waterandice					<b>-0.95</b>							<b>-0.93</b>
RT_naturalveg					<b>0.7</b>	-0.47						<b>0.82</b>
RT_agriculture					<b>0.81</b>	0.39	-0.33					<b>0.84</b>
RT_urbanareas					<b>0.9</b>							<b>0.91</b>
RT_bareareas	0.43					<b>0.64</b>	-0.53			0.29		0.25
Sunflux_alongtraj		<b>0.68</b>				0.5	-0.22	0.24	<b>0.68</b>		-0.25	
Temperature	<b>-0.85</b>					0.3		-0.29	<b>0.69</b>		-0.26	-0.23
RH	<b>0.82</b>					-0.31		<b>0.6</b>	-0.49			0.34
Explained variance (%)	22	15	14	12	8	8	19	17	11	11	10	9
Cumulative variance (%)	22	37	51	63	71	79	19	36	47	58	68	77

5



1  
2 Fig. 1 – Location of the sampling sites in the Po Valley.


1



2


3 Fig. 2 - Map with average trajectories for each obtained cluster with, in brackets, corresponding percentage of  
4 occurrence. The numbers outside the brackets identify each cluster. Figure refers to 96h trajectories arriving at  
5 500 m a.g.l.

1




2

3 **Fig. 3 - Time series of PM<sub>1,2</sub> and PM<sub>10</sub> mass concentrations (in µg m<sup>-3</sup>) and of the PM<sub>1,2</sub> to PM<sub>10</sub> ratio (%) for**  
4 **SPC. 4-days (96h) back trajectory length (km) is plotted superimposed to the graph, while air mass**  
5 **classification (in colors) is reported on top of it. The samples are labelled according to collection starting**  
6 **date, with “D” and “N” denoting respectively daytime and night-time samples.**




1  
2  
3  
4  
5  
6


Fig. 4 – Average day and night PM<sub>10</sub> composition at BO and SPC during the campaign. “Other ions” include chloride, nitrite, potassium and magnesium. WSOM stands for water soluble organic matter, while WINCM for water insoluble carbonaceous matter. Average mass, ALWC, RH and temperature are indicated below each pie. The size of each pie chart is proportional to the total measured mass reported.



1  
2 **Figure 5 – Time series of sulfate, nitrate and WSOC size-segregated concentrations in BO and SPC. Please note**  
3 **the different scale for nitrate and WSOC in BO.**




1  
 2 **Fig. 6 - Size-resolved aerosol composition for BO and SPC during day (top) and night (bottom) respectively**  
 3 **during one day characterized by background conditions (15/6) and during one day under stagnant conditions**  
 4 **(18/6) (notice the different scale for concentrations in the bottom right panel).**  
 5



1


2 Fig. 7 – Boxplot of Average diurnal variation of Aerosol Liquid Water Content (ALWC) with superimposed  
 3 diurnal variation of Relative Humidity (RH) and Aerosol Liquid Water Content (ALWC) during the campaign.  
 4 Please note the different scale of ALWC at the two sites. The shaded area denotes  $\pm 1$  standard deviation for  
 5 ALWC concentrations. The labels q1 and q3 in the boxplot respectively denote the first and the third quartiles.



1


2 **Fig. 8 – Relationship between ALWC and RH% (a), and between nitrate (b), sulfate (c) and WSOC (d) in the**  
 3 **condensation (CD, left panel) and in the droplet (DL, right panel) mode of particles, and ALWC averaged over**  
 4 **the sampling periods of the Berner Impactors.**

5



1  
2  
3  
4

Fig. 9 - Linear regressions between WSOC and TC (left) and WSOC and  $\text{SO}_4^{2-}$  (right) in the size intervals: 1) 0.05 – 0.14  $\mu\text{m}$ , 2) 0.14 – 0.42  $\mu\text{m}$ , and 3) 0.42 – 1.2  $\mu\text{m}$  in daytime and at night for BO (top) and SPC (bottom).




1

2

3 Fig. 10 – Time-series of WSOC and WINC (=TC-WSOC) concentrations in the size intervals: 1) 0.05 – 0.14 μm,  
 4 2) 0.14 – 0.42 μm, and 3) 0.42 – 1.2 μm. 4 days back-trajectories length is superimposed to each graph. Results  
 5 for the urban (BO) and for the rural (SPC) station are shown.

6



1

2

3 Fig. 11 – Scatterplot of sulfate concentration at SPC vs BO during day and night for the impactor stages 1 (0.05 –  
 4 0.14  $\mu\text{m}$ ), 2 (0.14 – 0.42  $\mu\text{m}$ ) and 3 (0.42 – 1.2  $\mu\text{m}$ ). The regression lines are referred to the diurnal (gray line) and  
 5 nocturnal (black line) concentrations in the three stages as a whole. Condensation and droplet mode samples for  
 6 the stagnant nights 16-19 June are filled in red.

7

8

9