

Replies to Editor comments/suggestions

Thanks for your revised version, which now has been seen by the two original referees again. One referee is satisfied with your changes, but the second referee (while agreeing that you have made substantial progress) still has important issues with your manuscript. Therefore I would ask you to revise the paper again taking the comments by the referee into account.

Reply: We wish to thank editor for going through the manuscript and allowing us to revise again for the betterment. We have taken care of all the suggestions made by the reviewer.

A few comments also from my side: The major remaining point seems to be the message conveyed by Figure 6 and the supporting evidence from the MLS observations (which have a limited spatial resolution). For example, in Fig. 6 you show 5 downward arrays but is this structure really reflected in Figs. 3 and 5?

Reply: Kindly note that the diagram shown in the figure 6 (now figure 7) is a typical structure of the cyclone but not exactly what we observed in the Figures 3 or 4. Figures 3 and 4 show the climatology of cyclone centred composite of the Ozone and Water vapour obtained from 16 cyclones. Our main interest to show a schematic picture is to convey the message on how the tropopause parameters vary and from where the mass flux will occur. Ozone comes down from the lower stratosphere due to subsidence at the centre of the cyclone and water vapour enters in to the lower stratosphere from the side bands due to anti-cyclonic circulation. Up and down arrows show the updrafts and downdrafts, where we can regularly observe in the cyclone structure.

There was one mistake i.e., representing the ozone coming from the side ways of the cyclone, which is rectified in the revised manuscript.

Further, you have added more observations and you have nicely provided background information in tables. However, the central figures 3 and 4 seem unchanged. Is this true? Or do you have to update the figures after the inclusion of the new data?

Reply: Kindly note that we have used same data set in the current version of the manuscript. Please go through Table 1 where we have mentioned all the cyclones during our initial submission itself. Thus, there is no change in the data set used in the current version and old version. Only difference is we show them according to the season in the revised manuscript.

Finally, there are a lot of technical/typesetting issues to be resolved in the revised version, as listed in the review. I have also noted that there is a typo on page 13, "Wie" should be "Wei" and the quantity "F" should be in italics in the text. Please also abbreviate the journal in reference Wei (1987).

Reply: We wish to inform that there was some problem in one of our computers (MS word) which is creating additional space. In the revised version of the manuscript, we have taken care of these typos. If this problem still persists, we will take help of production department of ACP. We have rectified the mistake in reference mentioned above.

In the revised version please make sure that the changes made are clearly noticeable (see also review).

Reply: Kindly note that we have incorporated all the suggestion made by the reviewer earlier with track changes and was uploaded. Our sincerely apology for not able to see the changes and we don't know what happened exactly.

Replies to Anonymous Referee #2 comments/suggestions

General comments:

The paper is much improved compared to the previous ACPD version. Unfortunately, the submitted version of the paper with 'change track' does not show the changes related to the ACPD Version. Therefore, it is very difficult to identify all the changes. The manuscript has still some shortcomings. Therefore, I recommend some revisions to address the comments listed below before publication by ACP.

Reply: The authors wish to thank the reviewer for his/her thorough review of the manuscript and for offering recommendations to improve the manuscript content. We have provided point-by-point replies to the reviewer's comments and information on how we have handled the revised manuscript. Kindly note that we have incorporated all the suggestion made by the reviewer earlier with track changes and was uploaded. Our sincerely apology for not able to see the changes and we don't know what happened exactly.

Presentation quality

The presentation quality is better than in the last version, however there are still plenty of redundant blank characters and some grammar issues. I recommend careful proof-reading by the production department of ACP.

Reply: Thank you very much for your appreciation. We wish to inform that there was some problem in one of our computers (MS word) which is creating additional space. In the revised version of the manuscript, we have taken care of these typos. If this problem still persists, we will take help of production department of ACP.

Specific comments:

1. Introduction:

p. 3, line 55: remove blank character 'water vapor- poor' or use 'dry'

Reply: Removed.

p. 3, line 65: 'Increase of water vapor in the LS region will leads to troposphere warming and stratospheric cooling might be due to lose ozone ...' Water has major consequences for the radiative balance and heat transport in the atmosphere. Enhanced ozone loss is a secondary effect of increasing water vapor.

Reply: We have changed this sentence in the revised manuscript as suggested.

p. 3, line 67: remove blank character '... Change, 2007) . Even ...'

Reply: Removed.

p. 3, line 67: 'Solomon et al. (2010) reported the relation between global warming and lower stratospheric water vapor.' Unspecific statement: please add some details

Reply: We have revised this statement with better clarity in the revised manuscript.

p. 3, line 71: 'long term' to 'long-term' (?)

Reply: Corrected.

p. 5, line 101-102: '...and also reported on the impact of the TCs on in the UTLS region on the regional scales.' What does this mean?

Reply: We have re-written this statement with better clarity.

p. 5, line 103: insert blank character 'RavindraBabu'

p. 5, line 114: remove blank character 'Aura- Microwave'

Reply: Removed.

2. Data and Methodology

p. 5, line 119: remove blank character 'Aura -MLS'

p. 6, line 127: remove '(1)' ??

Reply: Removed.

p. 6, line 129: What is the meaning of 'best track data'?

Reply: This is the IMD observed tropical cyclones best track data. We changed it in the revised manuscript.

p. 6, line 140: remove 'the' before 'Table 2'

Reply: Removed.

p. 7, line 149: 'We have 94 ± 21 mean MLS profiles for each cyclone' That means you use all MLS profiles for one cyclone (e.g. O3B) for all days of the cyclone period (4days) within 1000 km from the cyclone center. Is that correct? A figure showing the position of the MLS tracks (profiles) in the cyclone-centered coordinate system would be very helpful to see the data coverage.

Reply: Yes, we have used all MLS profiles during cyclone period (all cyclone days). The MLS overpasses have been separated with respect to cyclone centre for each day of the individual cyclone. As suggested by the reviewer, we added one more figure (as figure 2) to show the total MLS profiles around the cyclone-center used in the present study.

p. 7, line 164: 'Large convection around the eye and ...' Please add 'eye of the cyclone'

Reply: Added.

3. Results and discussion

Just for understanding: Figure 2 is for the period 2002-2013 and Figure 3 for 2007-2013. Why do you use different time periods?

Reply: Figure 2 shows the Cyclone centered – composite of mean difference in the tropopause parameters between climatological mean obtained by using GPS RO data from the year 2002-2013 and individual tropopause parameters observed during cyclones that have occurred during 2007-2013 respectively.

In general, if we want to represent any parameter climatologically, data length should contain at least one solar cycle so that while making composite, all the dominant oscillations like SAO, AO, QBO, ENSO, solar cycle etc., will be removed and remaining will represent true background. Thus, we make use of all GPS RO data available since 2002 (11 years). We also checked by considering data only during 2007-2013 as a background but could not see any major difference. This is mainly because of less number of GPS RO data available before 2007.

Compared to the previous version of the manuscript cyclones during the monsoon season (O3B, PHET, Gonu) are in addition included in the new version of the paper (20% of all profiles). Therefore, you use a different set of MLS data for calculating Fig. 3 and 4 in the new and old version of the paper. Is that correct? I am wondering why Figure 3 and 4 are exactly the same in the new and old version of the paper. I would expect some differences.

151 **Reply: We have used same data set in the current version of the manuscript. Please go**
152 **through Table 1 where we have mentioned all the cyclones during our initial submission**
153 **itself. Thus, there is no change in the data set used in the current version and old**
154 **version. Only thing is we show them according to the season in the revised manuscript.**

155 **4. Summary and conclusions**

156 I still recommend to adapt Fig. 6. The diagram does not describe the eastwest asymmetry
157 found in Figs- 3-5. From Fig.3, it is not clear that upward transport of high ozone values from
158 the lower stratosphere into the troposphere occurs outside of the cyclone center (red arrows)
159 as shown in Fig. 6.

160 **Reply: It is our mistake to represent the ozone coming from the side ways of the cyclone**
161 **which is rectified in the revised manuscript. Please note that the diagram shown in the**
162 **figure 6 (now figure 7) is a typical structure of the cyclone but not exactly what we**
163 **observed in the Figures 3 or 4. Figures 3 and 4 show the climatology of cyclone centred**
164 **composite of the Ozone and Water vapour obtained from 16 cyclones. Our main interest**
165 **to show a schematic picture is to convey the message on how the tropopause parameters**
166 **vary and from where the mass flux will occur. Ozone comes down from the lower**
167 **stratosphere due to subsidence at the centre of the cyclone and water vapour enters in**
168 **to the lower stratosphere from the side bands due to anti-cyclonic circulation. Up and**
169 **down arrows show the updrafts and downdrafts, where we can regularly observe in the**
170 **cyclone structure.**

171
172 Figures and Tables:

173 Table 2: Please add time unit for column 'total sustained' and 'sub-stained time with
174 maximum intensity'. The year (or date) of the cyclone occurrence would be in addition a
175 useful information.

176 **Reply: We have added time units and year of the cyclones occurred as suggested in the**
177 **revised manuscript.**

178
179 **We once again thank the reviewer for going through the manuscript carefully and**
180 **offering potential solutions to further improve the manuscript content.**

181
182
183 ---END---

**Effect of tropical cyclones on the Stratosphere-Troposphere Exchange
observed using satellite observations over north Indian Ocean**

M. Venkat Ratnam^{1*}, S. Ravindra Babu², Siddarth Shankar Das³, GhouseBasha¹⁴, B.V.
Krishnamurthy⁵⁴ and B.Venkateswararao²

¹National Atmospheric Research Laboratory (NARL), Gadanki, India.

²Jawaharlal Nehru Technological University, Hyderabad, India.

³Space Physics Laboratory (SPL), VSSC, Trivandrum, India.

⁴~~Masdar Institute of Science and Technology, Abu Dhabi, UAE.~~

⁵~~CEBROSS~~⁴CEBROSS, Chennai, India.

*vratnam@narl.gov.in, 08585-272123 (phone), 08585-272018 (Fax)

Abstract

Tropical cyclones play an important role in modifying the tropopause structure and dynamics as well as stratosphere-troposphere exchange (STE) processes in the Upper Troposphere and Lower Stratosphere (UTLS) region. In the present study, the impact of cyclones that occurred over the North Indian Ocean during 2007-2013 on the STE processes is quantified using satellite observations. Tropopause characteristics during cyclones are obtained from the Global Positioning System (GPS) Radio Occultation (RO) measurements and ozone and water vapor concentrations in the UTLS region are obtained from Aura-Microwave Limb Sounder (MLS) satellite observations. The effect of cyclones on the tropopause parameters is observed to be more prominent within 500 km from the centre of the tropical cyclone. In our earlier study, we have observed decrease (increase) in the tropopause altitude (temperature) up to 0.6 km (3K) and the convective outflow level increased up to 2 km. This change leads to a total increase in the tropical tropopause layer (TTL) thickness of 3 km within the 500 km from the centre of cyclone. Interestingly, an enhancement in the ozone mixing ratio in the upper troposphere is clearly noticed within 500 km from cyclone centre, whereas the enhancement in the water vapor in the lower stratosphere is more significant on south-east side extending from 500-1000 km away from the cyclone centre. The cross-tropopause mass flux for different intensities of cyclones are estimated and found that the mean flux from the stratosphere to the troposphere for cyclonic storms is $0.05 \pm 0.29 \times 10^{-3} \text{ kg m}^{-2}$ and for very severe cyclonic storms it is $0.5 \pm 1.07 \times 10^{-3} \text{ kg m}^{-2}$. More downward flux is noticed in the north-west and south-west side of the cyclone centre. These results indicate that the cyclones have significant impact in effecting the tropopause structure, ozone and water vapor budget and consequentially the STE in the UTLS region.

(Keywords: Tropical cyclone, tropopause, ozone, water vapor, STE processes.)

1. Introduction

The tropical cyclones with deep convective synoptic scale systems persisting for a few days to week, and play an important role on the mass exchange between the troposphere and the stratosphere, and vice versa (Merril, 1998; Emmanuel, 2005). They transport large amount of water vapor, energy and momentum to the upper troposphere and lower stratosphere (UTLS) region (Ray and Rosenlof, 2007). Cyclones provide favorable conditions for entry of the water vapour-rich and ozone-poor air from surface to the lower stratosphere (LS) and water vapor-poor dry and ozone-rich air from the LS to the upper troposphere (UT) leading to the stratosphere-troposphere exchange (STE) (Romps and Kuang 2009; Zhan and Wang, 2012; Vogel et al., 2014). These exchanges occur mainly around the tropopause and change the thermal and chemical structure of the UTLS region. The concentration of the water vapour transported from troposphere to stratosphere is controlled by the cold temperatures present at the tropopause and this is a major factor in the STE (Fueglistaler et al., 2009). As a consequence, the STE events play an important role in controlling the ozone in the UTLS region, which will affect the radiation budget of the Earth atmosphere (Intergovernmental Panel on Climate Change, 1996).

Water vapour has major consequences for the radiative balance and heat transport in the atmosphere. Enhanced ozone loss is a secondary effect of increasing water vapor. Increase of water vapor in the LS region will leads to troposphere warming and stratospheric cooling might be due to lose ozone(Rind and Lonergan, 1995; Forster and Shine, 1999; Dvortsov and Solomon, 2001; Forster and Shine, 2002; Myhre et al., 2007; Intergovernmental Panel on Climate Change, 2007). Even very small changes in lower stratospheric water vapor could affect the surface climate (Riese et al., 2012). Soloman et al. (2010) reported on the role of stratospheric water vapor in for the global warming, the relation between global warming and lower stratospheric water vapor. LS water vapor plays an

important role on the distribution of ozone in the lower stratosphere (Shindell, 2001). It is important contributor for [long-term](#) change in the LS temperatures (Maycock et al., 2014).

In general, most of the air enters into the stratosphere over the tropics (Brewer, 1949; Dobson, 1956). As suggested by Newell and Gould-Stewart (1981), Bay-of-Bengal (BoB) is one of the active regions where troposphere air enters into the stratosphere. It is also one of the active regions for the formation of deep convection associated cyclones which contains strong updrafts. Earlier studies have shown a close relationship between cyclones and moistening of the upper troposphere (Wang et al., 1995; Su et al., 2006; Ray and Rosenlof, 2007).

Several studies have been carried out related to water vapor, ozone transport as well as STE processes around the UTLS region during cyclones. Koteswaram (1967) described the thermal and wind structure of cyclones in the UTLS region with the major findings of cold core persisting just above the 15 km and the outflow jets very close to the tropopause. Penn (1965) reported enrichment in ozone and warmer air situated above the tropopause over the eye region during hurricane Ginny. Danielsen (1993) reported on troposphere-stratosphere transport and dehydration in the lower tropical stratosphere during cyclone period. Baray et al. (1999) studied the STE during cyclone Marlene and they observed maximum of ozone change at 300 hPa level. Zou and Wu (2005) observed the variations of columnar ozone in different stages of hurricane by using satellite measurements. Bellevue et al. (2007) observed increase in ozone concentration in the upper troposphere during Tropical Cyclone (TC) event. Significant contribution of cyclones on hydration of the UT is reported by Ray and Rosenlof (2007) and injection of tropospheric air into the low stratosphere due to overshooting convection by cyclones is reported by Romps and Kuang (2009). Das (2009) and Das et al. (2016) have studied the stratospheric intrusion into troposphere during the passage of cyclone by using Mesosphere-Stratosphere-Troposphere (MST) Radar

observations. Strong enhancement of ozone in the upper troposphere is observed during TCs over BoB (Fadnavis et al., 2011). The increased ozone levels in the boundary layer as well as near surface by as much as 20 to 30 ppbv due to strong downward transport of ozone in the tropical convection is also observed (Betts et al., 2002; Sahu and Lal, 2006; Grant et al., 2008). Cairo et al. (2008) reported that the colder temperatures are observed in the Tropical Tropopause Layer (TTL) region during cyclone Davina and also reported on the impact of the TCs on the UTLS structure and dynamics region at on the regional scales. A detailed review on the effect of TCs on the UTLS can be found in same report. Recently, Ravindra Babu et al. (2015) reported the effect of cyclones on the tropical tropopause parameters using temperature profile obtained from Constellation Observing System for Meteorology, Ionosphere and Climate (COSMIC) Global Position System Radio Occultation (GPS-RO) measurements. Many studies have been carried out on the role of extra tropical cyclones on the STE (for example Reutter et al., 2015 and references therein) though the quantitative estimates of STE provided by these case studies varied considerably. However, the vertical and horizontal variation of ozone and water vapor in the UTLS region and cross-tropopause flux quantification during cyclones over north Indian Ocean is not well investigated.

In the present study, we investigate the spatial and vertical variations of ozone and water vapor in the UTLS region for all the cyclones occurred over north Indian Ocean during 2007 to 2013 by using Aura-Microwave Limb Sounder (MLS) satellite observations. The effect of cyclones on the tropopause characteristics is also presented using COSMIC GPS-RO measurements. We also present the cross-tropopause mass flux estimated for each of the cyclones.

2. Data and Methodology

In the present study, we used Aura-MLS water vapor and ozone measurements (version 3.3) provided by the Jet Propulsion Laboratory (JPL). The version 3.3 was released

308 | in January 2011 and this updated version has change in the vertical resolution. The vertical
309 | resolution of the water vapor is in the range 2.0 to 3.7 km from 316 to 0.22 hPa and along
310 | track horizontal resolution varies from 210 to 360 km for pressure greater than 4.6 hPa. For
311 | ozone, vertical resolution is ~2.5 km and the along track horizontal resolution varies between
312 | 300 and 450 km (Livesey et al., 2011). The Aura MLS gives around 3500 vertical profiles per
313 | day and it crosses the equator at ~1:40 am and ~1:40 pm local time. For calculating the cross-
314 | tropopause mass flux, we used ERA-Interim winds obtained during cyclone period. (4)

315 | We have taken the cyclone track information data from India Meteorological
316 | Department (IMD) tropical cyclones observed best track data from year 2007-2013. During
317 | this period, around 50 cyclones have formed over the north Indian Ocean. Due to the
318 | considerable variability of cyclone life-cycles, for the present study we selected only 16
319 | cyclones that lasted for more than 4 days. The tracks of all the cyclones used for the present
320 | study are shown in Figure 1. Table 1 shows the classification of the cyclones over the North
321 | Indian Ocean. The TCs over the north Indian ocean are classified in to different categories by
322 | IMD based on their maximum sustained wind speed. There are classified as : (1) low pressure
323 | when the maximum sustained wind speed at the sea surface is < 17 knots (32
324 | km/hr), (2) depression (D) at 17–27 knots (32–50 km/hr), (3) deep depression (DD) at 28–33
325 | knots (51–59 km/hr), (4) cyclonic storm (CS) at 34– 47 knots (60–90 km/hr), (5) severe
326 | cyclonic storm (SCS) at 48–63 knots (90–110 km/hr), (6) very severe cyclonic storm (VSCS)
327 | at 64–119 knots (119–220 km/hr), and (7) super cyclonic storm (SuCS) at > 119 knots (220
328 | km/hr) (Pattnaik and Rama Rao, 2008). ~~The~~ Table 2 shows the different cyclones used in the
329 | present study and their maximum intensity, sustained time, and sustained time for peak
330 | intensity period of the each cyclone. The mean sustained time for cyclones that occurred
331 | during pre-monsoon, monsoon and post-monsoon seasons is 85.5 ± 52.4 hours, ~~for monsoon~~
332 | ~~season is~~ 122 ± 46.5 and ~~for post-monsoon season is~~ 112.6 ± 29.47 hours, respectively. Out

of the 16 cyclones, 4 cyclones (CS-1, SCS-2 and VSCS-1) formed during pre-monsoon season, 3 cyclones formed during monsoon season (CS-1, VSCS-1 and SuCS-1) and 9 cyclones (CS-1, SCS-2, and VSCS-6) formed during post-monsoon season (Table 2). Depressions and deep depressions are not considered. The total available MLS profiles for each cyclone that are used in the present study are listed in ~~the~~ Table 2. We have 94 ± 21 mean MLS profiles for each cyclone used in the present study and when segregated season wise, there are 108 ± 6 , 99 ± 21 and 88 ± 23 during monsoon, pre-monsoon and post-monsoon season, respectively. The available total MLS profiles for each cyclone vary with respect to sustained period of the cyclone and overall we have 1517 MLS profiles within 1000 km from the cyclone centre from all the 16 cyclones (Figure 2b). Since there are (temporal) limitations in the satellite measurements, ~~thus~~ mean cross-tropopause flux is estimated only for those cases of the cyclones that lasted for more than 4 days. However, our quantification of the cross-tropopause flux will not be affected by this limitation as earlier studies revealed that the maximum STE occurs during mature to peak stage of cyclone. Details on the selection of 16 cyclones are presented in Ravindra Babu et al. (2015). ~~The tracks of all the cyclones used for the present study are shown in~~ Figure 1, ~~and~~ different colors indicate different categories of the cyclones.

2.1. Tropopause characteristics observed during cyclones

As mentioned earlier, in the tropical region the amount of water vapor transported into the lower stratosphere from the troposphere is controlled by the cold tropical tropopause temperatures (Fueglistaler et al., 2009). Large convection around the eye [of the cyclone](#) and strong updrafts near the eye-walls transports large amount of water vapor into the lower stratosphere through the tropopause. In this way, cyclones will affect the tropopause structure (altitude/temperature). Thus, before quantification of STE, we show the tropopause characteristics observed during the TCs. We used post-processed products of level 2 dry

358 temperature profiles with vertical resolution around 200 m provided by the COSMIC Data
359 Analysis and Archival Center (CDAAC) for estimating the tropopause parameters during
360 cyclones period from 2007-2013. COSMIC GPS-RO is a constellation of six microsatellites
361 equipped with GPS receivers (Anthes et al., 2008). We also used CHALLENGING Minisatellite
362 Payload (CHAMP) GPS-RO data that are available between the years 2002 to 2006 and
363 COSMIC data from 2007-2013 for getting background climatology of tropopause parameters
364 over the north Indian Ocean.

365 Climatological mean of all the tropopause parameters are obtained by combining
366 GPS-RO measurements obtained from CHAMP and COSMIC (2002-2013). The tropopause
367 parameters include cold-point tropopause altitude (CPH) and temperature (CPT), lapse rate
368 tropopause altitude (LRH) and temperature (LRT) and the thickness of the tropical
369 tropopause layer (TTL), defined as the layer between convective outflow level (COH) and
370 CPH and are calculated for each profile of GPS-RO collected during the above mentioned
371 period. First, we separated the available RO profiles with respect to distance away from the
372 cyclone centre around 1000 km for individual cyclone for each day of the respective cyclone.
373 After separating, we calculated the tropopause parameters as mentioned above for each RO
374 profile. Total number of occultations used in the present study is shown in Figure 2(a). Then
375 we separated the tropopause parameters with respect to the different cyclone intensity. After
376 estimating the tropopause parameters for all the 16 TCs with respect to different intensity,
377 cyclone-centre composite of all tropopause parameters is obtained. After careful analysis, it is
378 found that there is no much variation in the tropopause parameters observed between D and
379 DD, and between CS and SCS, and thus they are combined to DD and CS, respectively. To
380 quantify the effect of the TCs on the tropopause characteristics, the climatological mean is
381 removed from the individual tropopause parameters. The climatological mean tropopause
382 parameters is estimated from the temperature profiles obtained by using GPS-RO data from

2002-2013. We also calculated the difference of tropopause parameters for different cyclone intensities (Figures are not shown). Figure 2-3 shows the cyclone centered – composite of mean difference in the tropopause parameters (CPH, LRH, CPT, LRT, COH and TTL thickness) between climatological mean (2002-2013) and individual tropopause parameters observed during cyclones (irrespective of cyclone intensity) and the more detailed results on effect of TCs on the tropopause variations and mean temperature structure in UTLS region during TCs can be found in Ravindra Babu et al. (2015). We have reported that the CPH (LRH) is lowered by 0.6 km (0.4 km) in most of the areas within the 500 km radius from the cyclone centre and the temperature (CPT/LRT) is more or less colder or equal to the climatological values from the area around 1000 km from the cyclone centre. Note that effect of cyclone can be felt up to 2000 km but since the latitudinal variation also comes into picture when we consider 2000 km radius, we restrict our discussion related to variability within 1000 km from the cyclone centre. COH (TTL thickness) has increased (reduced) up to 2 km within 500 km from the cyclones and in some areas up to 1000 km. Note that this decrease in TTL thickness is not only because of pushing up of the COH but also due to decrease of CPH. From the above results, we concluded that the tropical tropopause is significantly affected by the cyclones and the effect is more prominent within 500 km from the cyclone centre. These changes in the tropopause parameters are expected to influence water vapor and ozone transported in the UTLS region during cyclones.

3. Results and discussion

3.1. Ozone variability in the UTLS region during cyclones

To see the variability and the transport of ozone during the passage of cyclones, we investigated the spatial and vertical variability of ozone in the UTLS region using MLS satellite observations. As mentioned in Section 2.1, we also separated the MLS profiles based on the distance from the TC centre for each day of the individual cyclone. From all the 16

cyclones cases, we separated the available MLS profiles with respect to distance from the cyclone centre around 1000 km and also we separated the MLS profiles with respect to different intensities of the cyclones. Figure 3-4 shows the normalized cyclone centered – composite of mean ozone mixing ratio (OMR) observed during cyclones (irrespective of cyclone intensity) at 82hPa, 100hPa, 121hPa, and 146 hPa pressure levels during 2007-2013. Note that we have reasonable number of MLS profiles (1517) from 16 cyclones to generate the meaningful cyclone-centre composite of ozone. Black circles are drawn to show distances 250 km, 500 km, 750 km and 1000 km away from cyclone center. Since large variability in OMR is noticed from one pressure level to other, we normalized the values to the highest OMR value at a given pressure level. The highest OMR values at 82 hPa, 100 hPa, 121 hPa and 146 hPa pressure levels is 0.38 ppmv, 0.28 ppmv, 0.19 ppmv and 0.13 ppmv, respectively. Large spatial variations in the OMR are observed with respect to the cyclone centre. At 82 hPa, higher OMR (~0.4 ppmv) in the South-West (SW) side up to 1000 km and comparatively low OMR values (~0.2 ppmv) are noticed in the north of the cyclone centre. At 100 hPa, an increase in the OMR (~0.2 ppmv) near the cyclone centre within 500 km is clearly observed. This enhancement in OMR extends up to 146 hPa and is more prominent slightly in the western and eastern side of the cyclone. In general, the large subsidence located at the top of the cyclone centre is expected to bring lower stratospheric ozone to the upper troposphere. This might be the reason for the enhancement of ozone in the cyclone centre within 500 km. Earlier several studies have reported that the intrusion of the stratospheric air in to the troposphere due to the subsidence in the eye region (Penn, 1965; Baray et al., 1999; Das et al., 2009; Das et al., 2015). The present results also support this aspect that the detrainment of ozone reached to the 146 hPa might be due to strong subsidence. Interestingly, an enhancement in OMR in south east side at 121 hPa but is not either at 100 hPa or at 146 hPa can be noticed which need to be investigated further. Thus, in

433 | general, higher ozone concentrations are observed in cyclone_centre_within 500 km and
434 | slightly aligned to the western side of the cyclone centre.

435 | In order to quantify the impact of cyclones_on UTLS ozone more clearly we have
436 | obtained anomalies by subtracting the mean cyclone-centered ozone observed during
437 | cyclones_from the background climatology of UTLS ozone that is calculated by using the
438 | total available MLS profiles from 2007-2013. Figure 3-4(e-h) shows the normalized mean
439 | difference of cyclone-centered ozone obtained after removing the background climatology
440 | values for different pressure levels shown in Figure 3-4(a-d). The maximum difference in
441 | OMR for corresponding normalized value at 82 hPa, 100 hPa, 121 hPa and 146 hPa pressure
442 | levels is -0.089 ppmv, -0.19 ppmv, -0.09 ppmv and -0.06 ppmv, respectively. Enhancement
443 | in the OMR (~0.1 ppmv) up to 1000 km from the cyclone_centre is observed at 82 hPa.
444 | Interestingly, at 100 hPa_OMR is more or less uniform_throughout 1000 km from the cyclone
445 | centre_except ~500 km radius from the centre_where significant increase of OMR (~0.2
446 | ppmv) is observed. This increase_in the OMR is within 500 km from cyclone_centre_and
447 | extends up to 121 hPa. However, enhancement in OMR at 146 hPa extends up to 1000 km
448 | but distributed towards eastern and western sides of cyclone_centre. Thus, it is clear that the
449 | detrainment of lower stratospheric ozone will reach up to 146 hPa during cyclone period due
450 | to presence of strong subsidence in the cyclone_centre.We also calculated the cyclone-centre
451 | composite of ozone based on different cyclone intensities such as DD, SCS and VSCS. After
452 | carefully going through them, we have found that this detrainment of ozone reaching up to
453 | 146 hPa is more in the higher intensity period of the TCs. We do not know what happens
454 | below this pressure level due to limitation in the present data, however, studies (Das et al.,
455 | 2015; Jiang et al., 2015) have shown that LS ozone can reach as low as boundary layer during
456 | cyclones. It will be interesting to see the variability in the water vapor as large amount of it is
457 | expected to cross the tropopause during the cyclone period and reach lower stratosphere.

3.2. Water vapor variability in the UTLS region during cyclones

As mentioned earlier, enormous amount of water vapor is expected to be pumped from lower troposphere to the upper troposphere and even it can penetrate into the lower stratosphere during cyclones. To see the linkage between tropopause variability and the transport of water vapor during cyclones, we investigated the horizontal and vertical variability of water vapor in the UTLS region using MLS satellite observations. Figure 4 shows the normalized cyclone centered – composite of mean water vapor mixing ratio observed during cyclones (irrespective of cyclone intensity) at 82hPa, 100hPa, 121hPa, and 146 hPa pressure levels observed by MLS during 2007-2013. Black circles are drawn to shown the 250 km, 500 km, 750 km and 1000 km away from cyclone center. The highest Water Vapor Mixing Ratio (WVMR) values for corresponding normalized value at 82 hPa, 100 hPa, 121 hPa, and 146 hPa pressure levels is 4.44 ppmv, 4.49 ppmv, 6.9 ppmv and 16.03 ppmv, respectively. Significantly higher WVMR values are noticed extending from 500 km up to 1000 km from the cyclone centre at 121 (~6.5 ppmv), 146 hPa (~15 ppmv) levels with more prominence in the eastern side of the cyclone centre. Comparatively low values are noticed in the centre of the cyclone, especially at 121 hPa. These results are comparing well with higher WVMR observed in the eastern side of cyclones over Atlantic and Pacific Oceans (Ray and Rosenlof, 2007). These results also compare well with those reported by Ravindra Babu et al. (2015) where they used GPS-RO measured relative humidity and found enhancement in RH in the eastern side of the centre in the upper troposphere (10-15 km) over north Indian Ocean. The higher WVMR values are observed in the eastern side of the cyclone centre might be due to the upper level anti-cyclonic circulation over the cyclones. It is interesting to note that high WVMR lies not at the centre but extend from 500 to 1000 km from the centre of cyclone. The WVMR show high at 121 and 146 hPa than at 100 and 82 hPa. It seems less water vapor has been transported to 100 and 82 hPa from below. As we

483 know, water vapor mostly origin from lower troposphere and decreasing with height. So
484 vertical transport of water vapor from the lower troposphere to the UTLS may lead to water
485 vapor enhanced at 121 and 146 hPa and some time it reaches to higher altitudes. The higher
486 WVMR presented at 100 and 82 hPa levels show the signature of the tropospheric air
487 entering even in to the lower stratosphere during cyclones.

488 | In order to quantify the impact of cyclones on the UTLS water vapor more clearly, we
489 | have obtained anomalies by subtracting the mean cyclone-centered water vapor observed
490 | during cyclones from the background climatology mean of UTLS water vapor. Figure 4-5(e-
491 | h) shows the normalized mean difference of the cyclone-centered WVMR obtained after
492 | removing the background climatology values for different pressure levels shown in Figure 4
493 | 5(a-d). The maximum difference in WVMR for corresponding normalized values at 82 hPa,
494 | 100 hPa, 121 hPa, and 146 hPa pressure levels is -0.44 ppmv, -0.81 ppmv, -2.55 ppmv and -
495 | 9.09 ppmv, respectively. More than 7 ppmv differences are observed at 146 hPa within the
496 | 1000 km from the centre and at 121 hPa difference of ~ 2 ppmv is noticed extending up to
497 | 2000 km (figure not shown) in the eastern side of the centre. At 100 hPa and 82 hPa levels,
498 | the increase in the WVMR is ~0.8 and ~0.6 ppmv, respectively, and the enhancement is more
499 | observed in the NE side of the cyclone centre. Thus, a clear STE is evident during the
500 | cyclone over north Indian Ocean where a clear enhancement in the water vapor (ozone) in the
501 | lower stratosphere (upper troposphere) is observed. For quantifying the amount of STE, we
502 | calculated the cross-tropopause mass flux for each cyclone by considering the spatial extent
503 | within the 500 km from the cyclone centre and results are presented in the following sub-
504 | section.

505 3.3. Cross tropopause flux observed during cyclones

506 | We adopted method given by WeiWie (1987) to estimate the cross tropopause mass
507 | flux, F . F is defined as:

Formatted: Font: Italic

Formatted: Font: Italic

$$F = \frac{1}{g} \left(-\omega + V_h \cdot \nabla P_{tp} + \frac{\partial P_{tp}}{\partial t} \right) = \left(-\frac{\omega}{g} + \frac{1}{g} V_h \cdot \nabla P_{tp} \right) + \frac{1}{g} \frac{\partial P_{tp}}{\partial t} = F_{AM} + F_{TM} \quad (1)$$

where ω is the vertical pressure-velocity, V_h is the horizontal vector wind, P_{tp} is the pressure at the tropopause, g is the acceleration due to gravity, F_{AM} is the air mass exchange due to horizontal and vertical air motions, F_{TM} is the air mass exchange due to tropopause motion.

The wind information is taken from ERA-Interim, and the tropopause temperature and pressure within 500 km from the cyclone centre is estimated from COSMIC GPS-RO measurements (Ravindra Babu et al., 2015). These values are considered for the maximum intensity day for each of the 16 cyclones and the respective cross tropopause flux is estimated. Since the above mentioned results showed that the higher OMR values are observed in the west and NW side and more water vapor is located at the eastern side of the cyclone centre, we separated the area into 4 sectors with respect to cyclone centre as C1 (NW side), C2 (NE side), C3 (SW side), and C4 (SE side), respectively as shown in Figure 34(a). List of cyclones used in the present study with their names, cyclone intensity (CI), centre latitude, centre longitude, minimum estimated central pressure on their peak intensify day are provided in Table 3. The total flux F (equation 1) depends on the air mass exchange due to horizontal and vertical air motion (F_{AM}), and the air mass exchange due to tropopause motion itself (F_{TM}). Since number of COSMIC GPS-RO measurements are not sufficient to estimate the second term (F_{TM}) for each event, we calculated only the first part of the equation (F_{AM}) individually for each of cyclone with respect to different sectors mentioned above and the values are presented in Table 3. However, we roughly estimated the contribution of second term by assuming change in the tropopause pressure by 0.5 hPa increase (decrease) within 6 hr and could see cross-tropopause flux for CS is $0.25 \pm 0.07 \times 10^{-3} \text{ kgm}^{-2} \text{ s}^{-1}$ ($-0.36 \pm 0.07 \times 10^{-3} \text{ kgm}^{-2} \text{ s}^{-1}$) and for VSCS it is $-0.24 \pm 0.3 \times 10^{-3} \text{ kgm}^{-2} \text{ s}^{-1}$ ($-0.85 \pm 0.3 \times 10^{-3} \text{ kgm}^{-2} \text{ s}^{-1}$). If there is change in the tropopause pressure by 1 hPa increase (decrease), the flux for CS is

Formatted: Font: Italic

0.55±0.07×10⁻³ kgm⁻²s⁻¹ (-0.66±0.07×10⁻³ kgm⁻²s⁻¹) and for VSCS it is 0.06±0.3×10⁻³ kgm⁻²s⁻¹ (-1.16±0.3×10⁻³ kgm⁻²s⁻¹).

Figure 5-6 shows the cross-tropopause flux estimated in each sector from the centre of the cyclone for the different cyclone intensities (estimated based on the cyclone centre pressure). Red lines show the best fit. It clearly shows that the downward flux is always more in C1 and C3 sectors, whereas C2 sector show more upward flux. The flux itself varies with the cyclone intensity and it is found that the increase in downward flux as the cyclone centre pressure decreases particularly for C1 and C3 sectors. Whereas, in C4 sector, increase in the upward flux is seen as the cyclone intensity increases but always upward in C2 sector, irrespective of the cyclone intensity. The second term (in equation 1) itself corresponds the air mass exchange from the tropopause motion and generally during cyclone period there is an ~400 m difference in tropopause altitude (LRH) within 500 km from the centre of the cyclone (Figure 23). Thus, the spatial and temporal variation of the tropopause during the cyclones itself is very important for to decide the flux as downward or upward. Interestingly, C1 and C3 sectors of cyclone show dominant downward mean flux and C2 and C4 sectors show dominant upward mean flux with the values of 0.4±0.4×10⁻³ kgm⁻², 1.2±1.0×10⁻³ kgm⁻², 0.2±0.1×10⁻³ kgm⁻² and 0.12±0.3×10⁻³ kgm⁻², respectively. These results strongly support our findings of higher ozone in the NW and SW sides and higher water vapor in the NE side of the cyclone centre. The mean flux is observed to vary with the intensity of the cyclone. Mean flux for the severe cyclonic storms (CS) is -0.05±0.29×10⁻³ kgm⁻² whereas for very severe cyclonic storms (VSCS) it is -0.5±1.07×10⁻³ kgm⁻². Reutter et al. (2015) reported the upward and downward mass fluxes across the tropopause are more dominant in a deeper cyclones compared to a less intense cyclones over the North Atlantic. Our results are comparable with their results with the averaged mass flux of the stratosphere to troposphere as 0.3×10⁻³ kgm⁻² s⁻¹ (340 kgkm⁻² s⁻¹) in the vicinity of cyclones over the North Atlantic Ocean. They also

557 reported that the more transport across the tropopause occurred in the west side of the
558 cyclone centre during intensifying and mature stages of the cyclones over the North Atlantic
559 region.

560 **4. Summary and conclusions**

561 In this study, we have investigated the vertical and spatial variability of ozone and
562 water vapor in the UTLS region during the passage of cyclones occurred between 2007 and
563 2013 over the North Indian Ocean by using Aura-MLS satellite observations. In order to
564 make quantitative estimate of the impact of cyclones on the ozone and water vapor budget in
565 the UTLS region, we removed the mean cyclone-centre ozone and water vapor from the
566 climatological mean calculated using MLS data from 2007 to 2013. We estimated the mean
567 cross-tropopause flux for each of the cyclones on their peak intensity day. The main findings
568 are summarized below.

- 569 1. Lowering of the CPH (0.6 km) and LRH (0.4 km) values with the coldest CPT and
570 LRT (2–3 K) within a 500 km radius from the cyclone centre is noticed. Higher (2
571 km) COH leading to the lowering of TTL thickness (~3 km) is clearly observed
572 (Ravindra Babu et al., 2015).
- 573 2. The impact of cyclones on ozone and the tropopause (altitude/temperature) is more
574 prominent within 500 km from the cyclone centre, whereas it is high from 500 km to
575 1000km in case of water vapor.
- 576 3. Detrainment of ozone is highest in the cyclone centre (within 500 km from the centre)
577 due to strong subsidence over top of the cyclone centre and this detrained ozone
578 reaches as low as 146 hPa level (~13-14 km).
- 579 4. The detrainment of ozone is more in the higher intensity period (SCS or VSCS) of the
580 cyclone compared to the low intensity (D or DD).

5. Interestingly, significant enhancement in the lower stratospheric (82 hPa) water vapor is noticed in the east and southeast side from the cyclone_centre.
6. Dominant downward [upward] cross-tropopause flux is observed in C1 (NW) and C3 (SW) [C2 (NE) and C4 (SE)] sectors of the cyclone.

Figure 7 shows the typical ~~assumption of~~ structure (not to scale) of the TC along with convective towers, updrafts, downdrafts. ~~Figure 6 depicts which and it depicts~~ above mentioned tropopause variability with respect to cyclone centre results in the form of the schematic diagram. The tropopause altitude (CPH) is lowered by 0.6 km within 500 km from the centre of the cyclone. The convective out flow level (COH) slightly pushes up (~2 km) within 500 km from the centre of the cyclone but not exactly in the centre. Thus, a decrease of about 3 km in the TTL thickness is observed within the 500 km from the cyclone_centre. Cyclone includes eye that extends from few km to 10's of kilometers. Strong convective towers with strong updrafts extending up to the tropopause in the form of spiral bands extending from 500 to 1000 km are present. Strong water vapor transport in to the lower stratosphere (82 hPa) while pushing up the COH is observed around these spiral bands in the present study. Between these spiral bands equal amount of subsidence is expected with strong subsidence existing at the centre of the cyclone. Significant detrainment of ozone present above or advected from the surroundings is observed reaching as low as 146 hPa at the cyclones_centre. Thus, it is clear that ozone reaches upper troposphere from lower stratosphere through the centre of the cyclone, whereas water vapor transport in to the lower stratosphere will happen from the 500 to 1000 km from the cyclones_centre. Since more intense cyclones are expected to occur in a changing climate (Kuntson et al., 2010), the amount of water vapor and ozone reaching to the lower stratosphere and upper troposphere, respectively, is expected to increase thus affecting complete tropospheric weather and climate. Future studies should focus on these trends.

606 **Acknowledgements:** We would like to thank COSMIC Data Analysis and Archive Centre
607 (CDAAC) for providing GPS-RO data used in the present study through their FTP site
608 (<http://cdaac-www.cosmic.ucar.edu/cdaac/products.html>). The provision of tropical cyclone
609 best track data used in the present study by IMD through their website
610 (<http://www.imd.gov.in/section/nhac/dynamic/cyclone.htm>) and Aura-MLS observations
611 obtained from the GES DISC through their ftp site (<https://mls.jpl.nasa.gov/index-eos-mls.php>) is highly acknowledged. This work is supported by Indian Space Research
612 Organization (ISRO) through CAWSES India Phase-II Theme 3 programme. The authors
613 would like to thank the Editor Dr. Rolf Müller, and two anonymous reviewers whose
614 comments helped considerably in improving the quality of this paper

Formatted: Justified, Space After: 10 pt, Line spacing: Double, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: 12 pt

617 **References:**

- 618 Anthes, R. A., Bernhardt, P. A., Chen, Y., Cucurull, L., Dymond, K. F., Ector, D., Healy, S.
619 B., Ho, S.-H., Hunt, D. C., Kuo, Y.-H., Liu, H., Manning, K., McCormick, C., Meehan, T.
620 K., Randel, W. J., Rocken, C., Schreiner, W. S., Sokolovskiy, S. V., Syndergaard, S.,
621 Thompson, D. C., Trenberth, K. E., Wee, T.-K., Yen, N. L., and Zeng, Z.: The
622 COSMIC/Formosat/3 mission: Early results, *B. Am. Meteorol. Soc.*, 89, 313–333, 2008.
- 623 Baray, J. L., Ancellet, G., Radriambelo T., and Baldy, S.: Tropical cyclone Marlene and
624 stratosphere-troposphere exchange, *J. Geophys. Res.*, 104, 13,953–13,970,
625 doi:10.1029/1999JD900028-1999.
- 626 Bellevue, J., Baray, J. L., Baldy, S., Ancellet, G., Diab, R. D., and Ravetta, F.: Simulations of
627 stratospheric to tropospheric transport during the tropical cyclone Marlene event, *Atmos.*
628 *Environ.*, 41, 6510–6526, doi:10.1016/j.atmosenv.2007.04.040, 2007.
- 629 Betts, A. K., Gatti, L. V., Cordova, A. M., Silva Dias, M. A. F., and Fuentes, J. D.: Transport
630 of ozone to the surface by convective downdrafts at night, *J. Geophys. Res.*, 107, 8046,
631 doi:10.1029/2000JD000158, 2002.
- 632 Brewer, A. W.: Evidence for a world circulation provided by the measurements of helium
633 and water vapor distribution in the stratosphere. *Quarterly Journal of Royal Meteorological*
634 *Society.*, 75, 351–363, doi:10.1002/qj.49707532603-1949.
- 635 Cairo, F., Buontempo, C., MacKenzie, A. R., Schiller, C., Volk, C. M., Adriani, A., Mitev,
636 V., Matthey, R., Di Donfrancesco, G., Oulanovsky, A., Ravegnani, F., Yushkov, V., Snels,
637 M., Cagnazzo, C., and Stefanutti, L.: Morphology of the tropopause layer and lower
638 stratosphere above a tropical cyclone: a case study on cyclone Davina (1999), *Atmos.*
639 *Chem. Phys.*, 8, 3411–3426, doi:10.5194/acp-8-3411-2008, 2008.
- 640 Danielsen, E. F.: In situ evidence of rapid, vertical, irreversible transport of lower
641 tropospheric air into the lower tropical stratosphere by convective cloud turrets and by

larger-scale upwelling in tropical cyclones, *J. Geophys. Res.*, 98, 8665–8681, doi:
10.1029/92JD02954-1993.

Das, S. S.: A new perspective on MST radar observations of stratospheric intrusions into
troposphere associated with tropical cyclone. *Geophys. Res. Lett.*, 36, L15821, doi:
10.1029/2009GL039184-2009.

Das, S.S., Ratnam, M. V., Uma, K.N., Subrahmanyam, K.V., Girach, I.A., Patra, A.K.,
Aneesh, S., Suneeth, K.V., Kumar, K.K., Kesarkar, A.P., Sijikumar, S., and Ramkumar,
G.: Influence of Tropical Cyclone on Tropospheric Ozone: Possible Implication, *Atmos.*
Chem. Phys. Discuss., 15, 19305-19323, 2015.

Das, S.S., Ratnam, M.V., Uma, K. N., Patra, A. K., Subrahmanyam, K. V., Girach, I. A.,
Suneeth, K.V., Kumar, K. K., and Ramkumar, G.: Stratospheric intrusion into the
troposphere during the tropical cyclone Nilam (2012), *Q. J. Royal Meteor. Soc.*, doi:
10.1002/qj.2810, 2016.

Dobson, G. M. B.: Origin and Distribution of the Polyatomic Molecules in the Atmosphere,
Royal Society of London Proceedings Series A, 236, 187–193,
doi:10.1098/rspa.1956.0127, 1956.

Dvortsov, V. L., and Solomon, S.: Response of the stratospheric temperatures and ozone to
past and future increases in stratospheric humidity, *J. Geophys. Res.*, 106, 7505 – 7514,
2001.

Emanuel, K. A.: Increasing destructiveness of tropical cyclones over the past 30 years,
Nature, 436, 686–688, doi:10.1038/nature03906, 2005.

Fadnavis, S., Berg, G., Buchunde, P., Ghude, S. D., and Krishnamurti, T. N.: Vertical
transport of ozone and CO during super cyclones in the Bay of Bengal as detected by
Tropospheric Emission Spectrometer, *Environ. Sci. Pollut. R.*, 18, 301–315,
doi:10.1007/s11356-010-0374-3, 2011.

Formatted: Justified

667 Forster, P.M. de F., and Shine, K. P.: Stratospheric water vapour changes as a possible
 668 contributor to observed stratospheric cooling, *Geophys. Res. Lett.*, 26, 3309-3312, 1999.
 669 Forster, P. M. And Shine, K. P.: Assessing the climate impacts of trends in stratospheric
 670 water vapour. *Geophys. Res. Lett.* 29: 1086–1089, doi: 10.1029/2001GL013909-2002.
 671 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Fu, I., Folkins, Q., and Mote, P. W.:
 672 Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009.
 673 Grant, D. D., Fuentes, J. D., DeLonge, M. S., Chan, S., Joseph, E., Kucera, P., Ndiaye, S. A.,
 674 and Gaye, A. T.: Ozone transport by mesoscale convective storms in western Senegal,
 675 *Atmos. Environ.*, 42, 7104–7114, doi:10.1016/j.atmosenv.2008.05.044, 2008.
 676 IPCC 1996: IPCC, *Climate Change 1995 – The Science of Climate Change*, Contribution of
 677 Working Group I to the Second Assessment Report, section 2 edited by: Houghton, J. T.,
 678 MeiraFilho, L. G., Callander, B. A., Harris, N., Kattenberg, A., and Maskell, K., University
 679 Press, Cambridge, 572 pp., 1996.
 680 Jiang, Y.C., Zhao, T.L., Liu, J., Xu, X.D., Tan, C.H., Cheng, X.H., Bi, X.Y., Gan, J.B., You,
 681 J.F., and Zhao, S.Z.: Why does surface ozone peak before a typhoon landing in
 682 southeast China? *Atmos. Chem. Phys.*, 15, 13331–13338, 2015
 683 Koteswaram, P.: On the structure of hurricanes in the upper troposphere and lower
 684 stratosphere, *Mon. Weather Rev.*, 95, 541–564, 1967.
 685 Knutson, T.R., John, L., McBride, Johnny Chan, Kerry Emanuel, Greg Holland, Chris
 686 Landsea, Isaac Held, James P. Kossin, Srivastava, A.K., and Masato Sugi: Tropical
 687 cyclones and climate change, *Nature Geosci.*, 3, 157 – 163, 2010.
 688 Livesey, N., Read, W. G., Frovidaux, L., Lambert, A., Manney, G. L., Pumphrey, H. C.,
 689 Santee, M. L., Schwartz, M. J., Wang, S., Cofield, R. E., Cuddy, D. T., Fuller, R. A.,
 690 Jarnot, R. F., Jiang, J. H., Knosp, B. W., Stek, P. C., Wagner, P. A., and Wu, D. L.: Earth

691 Observing System (EOS) Aura Microwave Limb Sounder (MLS) Version 3.3 Level 2 data
692 quality and description document, JPL D-33509, JPL publication, USA, 2011.

693 Maycock, A. C., Joshi, M.M., Shine, K.P., Davis, S.M and Rosenlof, K.H.: The potential
694 impact of changes in lower stratospheric water vapour on stratospheric temperatures over
695 the past 30 years. *Quart. J. Roy. Meteor. Soc.*, 140, 2176–2185, doi:10.1002/qj.2287-
696 2014.

697 Merrill, R. T. (1988), Characteristics of the upper-tropospheric environmental flow around
698 hurricanes, *J. Atmos. Sci.*, 45, 1665–1677, doi:10.1175/1520-
699 0469(1988)045<1665:COTUTE>2.0.CO;2.

700 Myhre, G., Nilsen, J. S., Gulstad, L., Shine, K. P., Rognerud, B., and Isaksen, I. S. A.:
701 Radiative forcing due to stratospheric water vapour from CH₄ oxidation, *Geophys. Res.*
702 *Lett.*, 34, L01807, doi:10.1029/2006gl027472, 2007.

703 Newell, R. E., and Gould-Stewart, S.: A stratospheric fountain, *Journal of Atmospheric*
704 *Science.*, 38, 2789–2796, doi:10.1175/1520-0469-1981.

705 Pattnaik, D. R. and Rama Rao, Y. V.: Track Prediction of very severe cyclone “Nargis” using
706 high resolution weather research forecasting (WRF) model, *J. Earth Syst. Sci.*, 118, 309–
707 329, 2008.

708 Penn, S.: Ozone and temperature structure in a Hurricane, *J. Appl. Meteorol.*, 4, 212–216,
709 1965.

710 RavindraBabu, S., VenkatRatnam, M., Basha, G., Krishnamurthy, B.V. and Venkateswara
711 Rao, B.: Effect of tropical cyclones on the tropical tropopause parameters observed using
712 COSMIC GPS RO data. *Atmos. Chem. Phys.*, 15, 10239-10249, doi: 10.5194/acp-15-
713 10239-2015.

714 Ray, E. A. and Rosenlof, K. H.: Hydration of the upper troposphere by tropical cyclones, *J.*
715 *Geophys. Res.*, 112, D12311, doi:10.1029/2006JD008009, 2007.

716 Reutter, P., Škerlak, B., Sprenger, M., and Wernli, H.: Stratosphere-troposphere exchange
 717 (STE) in the vicinity of North Atlantic cyclones. *Atmos. Chem. Phys.*, 15, 10939–10953,
 718 2015.

719 Riese, M., F. Ploeger, A. Rap, B. Vogel, P. Konopka, M. Dameris, and Forster, P.: Impact of
 720 uncertainties in atmospheric mixing on simulated UTLS composition and related radiative
 721 effects, *J. Geophys. Res.*, 117, D16305, doi:10.1029/2012JD017751-2012.

722 Rind, D., and Lonergan, P.: Modeled impacts of stratospheric ozone and water vapor
 723 perturbations with implications for high-speed civil transport aircraft. *J. Geophys.*
 724 *Res.*, **100**, 7381-7396, doi:10.1029/95JD00196- 1995.

725 Romps, D. M. and Kuang, Z. M.: Overshooting convection in tropical cyclones, *Geophys.*
 726 *Res. Lett.*, 36, L09804, doi:10.1029/2009GL037396, 2009.

727 Sahu, L. K. and Lal, S.: Changes in surface ozone levels due to convective downdrafts over
 728 the Bay of Bengal, *Geophys. Res. Lett.*, 33, L10807, doi:10.1029/2006GL025994, 2006.

729 Shindell, D.T.: Climate and ozone response to increased stratospheric water vapor. *Geophys.*
 730 *Res. Lett.* 28, 1551-1554, 2001.

731 IPCC, 2007a: Climate Change 2007: The Physical Science Basis. Contribution of Working
 732 Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate
 733 Change [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B.M.Tignor and H.L.
 734 Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York,
 735 NY, USA, 996 pp.

736 Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J.,
 737 and Plattner, G.-K.: Contributions of Stratospheric Water Vapor to Decadal Changes in the
 738 Rate of Global Warming, *Science*, 327, 1219–1223, 2010. Stenke, A. and Grewe, V.:
 739 Simulation of stratospheric water vapor trends: impact on stratospheric ozone
 740 chemistry. *Atmos. Chem. Phys.*, 5, 1257–1272, doi: 10.5194/acp-5-1257-2005.

741 Su, H., Read, W.G., Jiang, J. H., Waters, J. W., Wu, D. L., and Fetzer, E. J.: Enhanced
 742 positive water vapor feedback associated with tropical deep convection: New evidence
 743 from Aura MLS. *Geo. Research Letters.*, 33, L05709, doi: 10.1029/2005GL025505-2006.
 744 Vogel, B., Günther, G., Müller, R., Groß, J.-U., Hoor, P., Krämer, M., Müller, S., Zahn, A.,
 745 and Riese, M.: Fast transport from Southeast Asia boundary layer sources to northern
 746 Europe: rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon
 747 anticyclone, *Atmos. Chem. Phys.*, 14, 12745-12762, doi:10.5194/acp-14-12745-2014,
 748 2014.
 749 Wei, M. Y.: A new formulation of the exchange of mass and trace constituents between the
 750 stratosphere and troposphere. *Journal of Atmospheric Science.*, 44(20), 3079–3086,
 751 doi:10.1175/1520-0469-1987.
 752 Zhan, R. and Wang, Y.: Contribution of tropical cyclones to stratosphere–troposphere
 753 exchange over the northwest Pacific: estimation based on AIRS satellite retrievals and
 754 ERA-Interim data, *J. Geophys. Res.*, 117, D12112, doi: 10.1029/2012.
 755 Zou, X., and Y. Wu.: On the relationship between Total Ozone Mapping Spectrometer
 756 (TOMS) ozone and hurricanes. *J. Geophys. Res.*, 110, D06109, doi:
 757 10.1029/2004JD005019-2005.

Figure captions:

Figure 1. Tropical cyclone tracks of different categories (cyclonic storm (CS, blue color), severe cyclonic storm (SCS, orange color), very severe cyclonic storm (VSCS, red color) and super cyclonic storm (SuCs, magenta color)) that occurred over North Indian Ocean during 2007 - 2013.

Figure 2. Cyclone-centred composite of total available (a) COSMIC GPS RO occultations and (b) MLS profiles obtained from all the 16 cyclones that are used in the present study.

Figure 23. Cyclone centered – composite of mean difference in the tropopause parameters between climatological mean (2002-2013) and individual tropopause parameters observed during cyclones (irrespective of cyclone intensity) in (a) CPH (km), (b) LRH (km), (c) CPT (K), (d) LRT (K), (e) COH (km) and (f) TTL thickness (km). Black circles are drawn to show the 250 km, 500 km, 750 km and 1000 km away from cyclone center.

Figure 34. Normalized cyclone centered – composite of mean ozone mixing ratio observed during cyclones (irrespective of cyclone intensity) at (a) 82hPa, (b) 100hPa, (c) 121hPa, (d) 146 hPa levels by MLS during 2007-2013. (e) to (h) same as (a) to (d) but for normalized mean difference in the ozone mixing ratio between climatological mean (2007-2013) and individual events. Black circles are drawn to show the 250 km, 500 km, 750 km and 1000 km away from cyclone center. Sectors showing C1 (NW), C2 (NE), C3 (SW) and C4 (SE) are also shown in (a).

Figure 45. Same as Fig. 43, but for water vapor mixing ratio.

Figure 56. Cross-tropopause flux estimated in the (a) C1 (NW), (b) C2 (NE), (c) C3 (SW), and (d) C4 (SE) sectors from the centre of cyclone for different cyclone intensities (estimated based on cyclone centre pressure). Red lines show the best fit.

Figure 67. Schematic diagram showing the variability of CPH (brown color line) and COH (magenta color line) with respect to the centre of cyclone. Spiral bands of convective

towers reaching as high as COH are shown with blue color lines. Light blue (red) color up (down) side arrow shows the up drafts (downdrafts/subsidence). Thickness of the arrows indicates the intensity.

Table captions:

Table1.Classification of cyclonic systems over the north Indian Ocean.

Table 2. Tropical cyclones occurred during different seasons, cyclone name, cyclone Intensity (CI), cyclone period, total sustained time, Sustained time with maximum intensity and total number of available MLS profiles

Table 3. Cyclone name, cyclone Intensity (CI), centre latitude, centre longitude, estimated central pressure and estimated cross-tropopause mass flux with respect to cyclone centre for C1 (NW side), C2 (NE side), C3 (SW side) and C4 (SE side), respectively.

796 **Tables:**

797 **Table1.**IMD classification of cyclonic systems over the north Indian Ocean.

Intensity of the system	Maximum sustained surface winds (knots) at sea (1 knot =0.5144 m/s)
Low pressure area	<17
Depression	17–27
Deep depression (DD)	28–33
Cyclonic storm (CS)	34-47
Severe cyclonic storm (SCS)	48-63
Very severe cyclonic storm (VSCS)	64–119
Super cyclonic storm (SuCS)	>119

798

799

800

801

802

803

804

805

806

807

808

809

810 **Table 2.** Tropical cyclones occurred during different seasons, cyclone name, cyclone
811 Intensity (CI), cyclone period, total sustained time, Sustained time with maximum intensity
812 and total number of available MLS profiles.

Season	Cyclone Name	Cyclone Intensity (CI)	Cyclone Period (days)	Total Sustained time (hours)	Sustained Time with maximum intensity (hours)	Total available MLS profiles
Monsoon (JJA)	03B(2007)	CS	>4	75	6	104
	PHET (2010)	VSCS	>4	168	42	116
	Gonu (2007)	ScCS	>4	123	72	105
Pre-Monsoon (MAM)	Mahasen(2013)	CS	>4	24	24	119
	Aila (2009)	SCS	4	72	9	79
	Laila (2010)	SCS	4	96	27	82
	Nargis (2008)	VSCS	>4	150	87	118
Post-Monsoon (SON)	Nilam (2012)	CS	>4	102	36	52
	Jal (2010)	SCS	4	99	30	75
	Helen (2013)	SCS	4	78	30	72
	Giri (2010)	VSCS	4	66	15	65
	Phailin (2013)	VSCS	>4	147	66	111
	Leher (2013)	VSCS	>4	114	36	111
	SIDR (2007)	VSCS	>4	138	72	114
Winter (DJF)	Madi (2013)	VSCS	>4	150	36	104
	Thane (2011)	VSCS	>4	120	36	90

Formatted Table

813

814

815

816 **Table 3.** Cyclone name, cyclone Intensity (CI), centre latitude, centre longitude, estimated
817 central pressure and estimated cross-tropopause mass flux with respect to cyclone centre
818 for C1 (NW side), C2 (NE side), C3 (SW side) and C4 (SE side), respectively.

819


					Flux @500km			
Cyclone	CI	Centre Latitude	Centre Longitude	Estimated Central Pressure (hPa)	C1	C2	C3	C4
03B	CS	23.5	66	986 (25Jun2007)	-0.013	0.661	-0.603	-0.258
Aila	SCS	22	88	968 (25May2009)	1.90E-04	0.191	-0.299	-0.072
Helen	SCS	16.1	82.7	990 (21Nov2013)	0.025	0.216	-0.095	-0.11
Jal	SCS	11	84	988(6Nov2010)	0.025	0.384	-0.4	-0.218
Laila	SCS	14.5	81	986 (19May2010)	-0.012	0.123	-0.352	-0.299
Mahasen	CS	18.5	88.5	990 (15May2013)	-0.006	0.354	-0.473	-0.256
Nilam	CS	11.5	81	990 (31Oct2012)	0.016	0.313	-0.274	-0.097
Nargis	VSCS	16	94	962 (2May2008)	-0.828	0.094	-1.946	0.384
Giri	VSCS	19.8	93.5	950 (22Oct2010)	-0.518	0.022	-0.823	0.032
Gonu	SuCS	20	64	920 (4Jun2007)	-0.502	0.123	-2.563	0.37
Lehar	VSCS	13.2	87.5	980 (26Nov2013)	-0.55	0.119	-2.019	0.411
Madi	VSCS	13.4	84.7	986 (10Dec2013)	-0.375	0.054	-1.449	0.352
Phailin	VSCS	18.1	85.7	940 (11Oct2013)	-0.9	0.179	-2.576	0.479
Phet	VSCS	18	60.5	964 (2Jun2010)	-1.058	0.203	-2.698	0.559
SIDR	VSCS	19.5	89	944 (15Nov2007)	-0.493	0.066	-0.926	0.231
Thane	VSCS	11.8	80.6	970 (29Dec2011)	-1.272	0.356	-2.979	0.558

820

821

822 **Figures:**

823




824

825 **Figure 1.** Tropical cyclone tracks of different categories (cyclonic storm (CS, blue color),
826 severe cyclonic storm (SCS, orange color), very severe cyclonic storm (VSCS, red color)
827 and super cyclonic storm (SuCs, magenta color)) that occurred over North Indian Ocean
828 during 2007 - 2013.

829

830



831

832

833

Figure 2. Cyclone-centred composite of total available (a) COSMIC GPS RO occultations and (b) MLS profiles obtained from all the 16 cyclones that are used in the present study.




Figure 23. Cyclone centered – composite of mean difference in the tropopause parameters between climatological mean (2002-2013) and individual tropopause parameters observed during cyclones (irrespective of cyclone intensity) in (a) CPH (km), (b) LRH (km), (c) CPT (K), (d) LRT (K), (e) COH (km) and (f) TTL thickness (km). Black circles are drawn to show the 250 km, 500 km, 750 km and 1000 km away from cyclone center (taken from Ravindra Babu et al., ACP, 2015).





Figure 34. Normalized cyclone centered – composite of mean ozone mixing ratio observed during cyclones (irrespective of cyclone intensity) at (a) 82 hPa, (b) 100 hPa, (c) 121 hPa, (d) 146 hPa levels by MLS during 2007-2013. (e) to (h) same as (a) to (d) but for normalized mean difference in the ozone mixing ratio between climatological mean (2007-2013) and individual events. Black circles are drawn to show the 250 km, 500 km, 750 km and 1000 km away from cyclone center. Sectors showing C1 (NW), C2 (NE), C3 (SW) and C4 (SE) are also shown in (a).




851

852 | **Figure 45.** Same as Fig. 43, but for water vapor mixing ratio.

853

854



855

856

857

858

859


860

861

862

863

Figure 56. Cross-tropopause flux estimated in the (a) C1 (NW), (b) C2 (NE), (c) C3 (SW), and (d) C4 (SE) sectors from the centre of cyclone for different cyclone intensities (estimated based on cyclone centre pressure). Red lines show the best fit.



Formatted: Font: (Default) Times New Roman, 12 pt

Figure 67. Schematic diagram showing the variability of CPH (brown color line) and COH (magenta color line) with respect to the centre of cyclone. Spiral bands of convective towers reaching as high as COH are shown with blue color lines. Light blue (red) color up (down) side arrow shows the up drafts (downdrafts/subsidence). Thickness of the arrows indicates the intensity.