

1 **Effects of 20-100 nanometre particles on liquid clouds in the clean**
2 **summertime Arctic**

3
4 W. R. Leaitch, A. Korolev, A. A. Aliabadi
5 Environment Canada, Toronto, Canada

6
7 J. Burkart, M. Willis, J.P.D. Abbatt
8 Department of Chemistry, University of Toronto, Toronto, Canada

9
10 H. Bozem, P. Hoor
11 Institute of Atmospheric Physics, University of Mainz, Mainz, Germany

12
13 F. Köllner, J. Schneider
14 Particle Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany

15
16 A. Herber, C. Konrad
17 Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

18
19 R. Brauner
20 Jade University, Elsfleth, Germany

21
22 Date: July 28 2016

23
24 Correspondence to Richard.Leaitch@Canada.ca

25
26

27 **Abstract.** Observations addressing effects of aerosol particles on summertime Arctic clouds are
28 limited. An airborne study, carried out during July, 2014 from Resolute Bay, Nunavut, Canada,
29 as part of the Canadian NETCARE project, provides a comprehensive in-situ look into some
30 effects of aerosol particles on liquid clouds in the clean environment of the Arctic summer.
31 Median cloud droplet number concentrations (CDNC) from 62 cloud samples are 10 cm^{-3} for
32 low-altitude cloud (clouds topped below 200 m) and 101 cm^{-3} for higher-altitude cloud (clouds
33 based above 200 m). The lower activation size of aerosol particles is $\leq 50 \text{ nm}$ diameter in about
34 40% of the cases. Particles as small as 20 nm activated in the higher-altitude clouds consistent
35 with higher supersaturations (S) for those clouds inferred from comparison of the CDNC with
36 cloud condensation nucleus (CCN) measurements. Over 60% of the low-altitude cloud samples
37 fall into the CCN-limited regime of Mauritsen et al. (ACP, 2011) within which increases in
38 CDNC may increase liquid water and warm the surface. These first observations of that CCN-
39 limited regime indicate a positive association of the liquid water contents (LWC) and CDNC, but
40 no association of either the CDNC or LWC with aerosol variations. Above the Mauritsen limit,
41 where aerosol indirect cooling may result, changes in particles with diameters from 20 nm to 100
42 nm exert a relatively strong influence on the CDNC. Within this exceedingly clean environment,
43 as defined by low CO and low concentrations of larger particles, the background CDNC are
44 estimated to range between 16 cm^{-3} and 160 cm^{-3} , where higher values are due to activation of
45 particles $\leq 50 \text{ nm}$ that likely derive from natural sources. These observations offer the first wide-
46 ranging reference for the aerosol cloud albedo effect in the summertime Arctic.

47 **1. Introduction**

48

49 Mass concentrations of the atmospheric aerosol in the Arctic are higher during winter than in
50 summer due to differences in transport of anthropogenic particles and wet scavenging (e.g.
51 Barrie, 1986; Stohl, 2006). Atmospheric chemistry and aerosol-cloud Arctic research has largely
52 focussed on the springtime. The winter to summer transition offers the opportunity to examine
53 changes in chemistry as the sun rises over the polluted polar atmosphere (e.g. Barrie et al., 1988)
54 and to study impacts of anthropogenic aerosol on the Arctic solar radiation balance (e.g. Law and
55 Stohl, 2007; Quinn et al., 2008). Greater-than-expected warming of the Arctic (e.g. Christensen
56 et al., 2013) and rapidly diminishing Arctic sea ice extent (e.g. Maslanik et al., 2011) have drawn
57 considerable attention to the role of anthropogenic and biomass burning particles as warming
58 agents for the Arctic (e.g. Law and Stohl, 2007; Quinn et al., 2008; Shindell et al., 2008; Brock
59 et al., 2011; Jacob et al., 2010; UNEP, 2011; Stohl et al., 2013). Recent evidence indicates that
60 the net impact of aerosol particles on the Arctic over the past century has been one of cooling
61 rather than warming (Najafi et al., 2015).

62 Low-level liquid water clouds are frequent in the sunlit Arctic summer (e.g. Intrieri et al.,
63 2001), and these clouds can have a net cooling effect (e.g. Brenner et al., 2001; Garret et al.,
64 2004; Lubin and Vogelmann, 2010; Zhao and Garrett, 2015; Zamora et al., 2015;). Knowledge
65 of the influence of the atmospheric aerosol on climatic aspects of these clouds is complicated by
66 the relatively large potential differences in the albedo of the underlying surface (e.g. Herman,
67 1977; Lubin and Vogelmann, 2010) and the fact that the Arctic is relatively free of
68 anthropogenic influence in summer, which means that particles from natural sources can be the
69 most significant nuclei for cloud droplets. Those sources shift the number distribution toward

70 particles smaller than 100 nm (e.g. Heintzenberg and Leck, 1994; Ström et al., 2003;
71 Heintzenberg et al., 2006; Engvall et al., 2008; Tunved et al., 2013; Leaitch et al., 2013;
72 Heintzenberg et al., 2015). Particles smaller than 100 nm are often dismissed as being too small
73 to nucleate cloud droplets due to the assumption that the cooling mechanisms are too slow to
74 generate the supersaturation (S) required to activate the smaller particles in Arctic liquid clouds
75 (e.g. Garret et al., 2004; Lubin and Vogelmann, 2010; Browse et al., 2014; Zhao and Garrett,
76 2015). That assumption may lead to reduced estimates from natural feedbacks to climate and
77 increased estimates of aerosol indirect forcing from anthropogenic sources. Lohmann and Leck
78 (2005) hypothesized the need for highly surface-active particles to explain CCN activity at S less
79 than 0.3%. However, cloud S is also strongly constrained by the concentrations of particles
80 larger than 100 nm, and in the clean summertime Arctic environment with relatively low
81 concentrations of particles larger than 100 nm, there is some evidence that higher S may be
82 achieved and smaller particles activated (e.g. Hudson et al., 2010; Korhonen et al., 2010; Leaitch
83 et al., 2013). Further, the suggestion that the minima between 50 and 100 nm in Arctic particle
84 size distributions results from cloud processing implies consistent activation sizes less than 100
85 nm (Heintzenberg et al., 2015). The effect of the background aerosol on liquid clouds has been
86 identified as one of the most important factors for reducing uncertainty in the aerosol cloud
87 albedo effect (Carslaw et al., 2013). Moreover, the effectiveness of particles smaller than 100 nm
88 for cloud droplet nucleation is a large factor in that uncertainty.

89 Effects of pollution on clouds may also lead to warming, but a reference to clean clouds
90 is still required (e.g. Garrett et al., 2009). Mauritsen et al. (2011) modeled cloud radiative
91 forcing for low clouds using CCN number concentrations derived from shipborne observations
92 over the Arctic Ocean (Tjernström et al., 2004; Tjernström et al., 2014). They found the impact

93 from changes in CCN for ultra-low values ($< 10 \text{ cm}^{-3}$), where CCN concentrations are equivalent
94 to model CDNC, results in a net warming due to associated longwave changes, whereas for
95 concentrations greater than 10 cm^{-3} CCN increases are estimated to produce a net atmospheric
96 cooling. This CCN concentration threshold is referred to here as the "Mauritsen limit", although
97 this value of 10 cm^{-3} is not a universal limit (Mauritsen et al., 2011). In the clean summertime
98 Arctic, knowledge of the natural aerosol and its influence on cloud microphysics is critical to the
99 assessment of aerosol effects on Arctic climate.

100 Past studies of Arctic aerosols and clouds have emphasized the areas of the Beaufort and
101 Chukchi Seas (e.g. Hobbs and Rango, 1998; Curry et al., 2001 and references therein; Lohmann
102 et al., 2001; Yum and Hudson, 2001; Peng et al., 2002; Wylie and Hudson, 2002; Earle et al.,
103 2011; Lance et al., 2011; Jouan et al., 2014; Klingebiel et al., 2014). Most of those studies have
104 focused on springtime when the aerosol can be influenced by anthropogenic or biomass burning
105 sources. As well, there has been considerable interest in mixed-phase clouds in the lower Arctic
106 troposphere (e.g. Shupe et al., 2004; Sandvik et al., 2007; Morrison et al., 2012), but a notable
107 lack of in-situ aerosol observations in combination with liquid water clouds over the summertime
108 Arctic. Among the studies that have considered in-situ aerosol measurements and summertime
109 Arctic clouds, Zamora et al. (2015) examined the efficiency of biomass burning (BB) plumes on
110 indirect forcing. They estimated half of the possible maximum forcing from these plumes,
111 mostly due to the reduction in cloud-base S by higher concentrations of larger particles that
112 control water uptake. Shupe et al. (2013) discussed some differences between clouds coupled
113 and uncoupled to the surface. They did not conduct in-situ cloud microphysics observations, their
114 and vertical aerosol characterizations were constrained to particles $>300 \text{ nm}$. Hobbs and Rango
115 (1998) found that droplets in June low clouds over the Beaufort Sea occasionally contained drops

116 as large as 35 μm diameter. They also found that cloud-top CDNC correlated significantly with
117 cloud base “aerosols”. They suggested that cloud-top entrainment did not control CDNC,
118 although there may be times when entrainment influences Arctic CDNC (e.g. Klingebiel et al.,
119 2014).

120 Motivated by limited knowledge of aerosol effects on summertime Arctic clouds and
121 particle activation details, the Canadian Network on Climate and Aerosols: Addressing Key
122 Uncertainties in Remote Canadian Environments (NETCARE - <http://www.netcare-project.ca/>),
123 conducted airborne aerosol and cloud observations during July, 2014 in the area around Resolute
124 Bay, Nunavut, Canada. The observations from this study are used here to characterize CDNC,
125 LWC, and the volume-weighted mean droplet diameter (VMD). Further, aerosol particle size
126 distributions (5 nm and larger; CCNC(0.6%)) from outside of clouds are compared with droplet
127 number concentrations from inside of clouds. Specifically, in the indicated sections, the
128 following questions are addressed.

- 129 1) Given the scarcity of data, what are the characteristics of clouds in the summertime Arctic,
130 and do clouds near the surface have characteristics different from those aloft? (Sect. 3.2)
- 131 2) What are the sizes of particles that act as nuclei for cloud droplets? This will allow a closer
132 connection between aerosol processes, particle sizes and climate effects? (Sect. 3.3)
- 133 3) What is the relationship between droplet size and droplet number? In particular, what is the
134 aerosol influence on cloud below the Mauritsen-limit, and is it possible to assess a
135 background influence of the aerosol on clouds in the Arctic summer? (Sect. 3.4)

136

137 **2. Methodologies**

138

139 The instrument platform was the Alfred Wegener Institute (AWI) Polar 6 aircraft, a DC-3
140 aircraft converted to a Basler BT-67 (see Herber, A., Dethloff, K., Haas, C., Steinhage, D.,
141 Strapp, J. W., Bottenheim, J., McElroy, T. and Yamanouchi, T.; POLAR 5 - a new research
142 aircraft for improved access to the Arctic, ISAR-1, Drastic Change under the Global Warming,
143 Extended Abstract, pp. 54-57, 2008).

144

145 **2.1 Instrumentation**

146

147 The following measurements are relevant to this discussion:

- 148 a) Particle number concentrations >5 nm diameter were measured with a TSI 3787 water-
149 based ultrafine condensation particle counter (UCPC), sampling at a flow rate of 0.6 L
150 min^{-1} . Hereafter, these measurements are referred to as N5.
- 151 b) Aerosol particle size distributions from 20 nm to 100 nm (45 s up scans and 15 s down
152 scans) were measured using a Brechtel Manufacturing Incorporated (BMI) Scanning
153 Mobility System (SMS) coupled with a TSI 3010 Condensation Particle Counter (CPC).
154 The sheath and sample flows were set to 6 L min^{-1} and 1 L min^{-1} . BMI software was used
155 to process these distributions.
- 156 c) Aerosol particle size distributions from 70 nm to 1 μm were measured using a Droplet
157 Measurement Technology (DMT) Ultra High Sensitivity Aerosol Spectrometer (UHSAS)
158 that detect particles using scattering of 1054 nm laser light (e.g. Cai et al., 2008).
- 159 d) CCNC(0.6%) were measured using a DMT CCN Model 100 counter operating behind a
160 DMT low pressure inlet at approximately 650 hPa. For the nominal water S of 1%, the
161 effective S at 650 hPa was found to be 0.6% as discussed below. This S was held

162 constant throughout the study for greater measurement stability, improved response, and
163 to examine the hygroscopicity of smaller particles.

164 e) Droplet size distributions from 2-45 μm were measured with a Particle Measuring
165 Systems (PMS) FSSP-100. This FSSP-100 had been modified with new tips to reduce
166 shattering artifacts (Korolev et al., 2011). It was mounted in a canister under the port-side
167 wing. The CDNC, VMD and LWC are calculated from the measured droplet
168 distributions.

169 f) Two-dimensional cloud particle images from about 50 μm to 800 μm were measured
170 using a PMS 2DC Grey-scale probe. These observations are used here only to ensure the
171 absence of the ice phase. This 2DC-Grey was also modified with new tips to reduce
172 shattering artifacts (Korolev et al., 2011). It was mounted in a canister beside the FSSP-
173 100.

174 g) Carbon monoxide (CO) is used here as a relative indicator of aerosol influenced by
175 pollution sources and as a potential tracer for aerosol particles entering cloud. CO was
176 measured with an Aerolaser ultra-fast carbon monoxide monitor model AL 5002 based
177 on VUV-fluorimetry, employing the excitation of CO at 150 nm. This instrument was
178 modified such that in-situ calibrations could be conducted in flight.

179

180 Details of the instrument calibration and evaluations are given in the Supplement (S1).

181

182 **2.1 State parameters and Winds**

183

184 State parameters and meteorological measurements were made with an AIMMS-20,
185 manufactured by Aventech Research Inc. This instrument consists of three modules: 1) an Air
186 Data Probe that measures the three-dimensional aircraft-relative flow vector (true air speed,
187 angle-of-attack, and sideslip), temperature and relative humidity, and includes a three-axis
188 accelerometer pack for turbulence measurement; 2) an Inertial Measurement Unit that consists of
189 three gyros and three accelerometers providing the aircraft angular rate and acceleration; 3) a
190 Global Positioning System for aircraft 3D position and inertial velocity. Horizontal and vertical
191 wind speeds were measured with accuracies of 0.50 and 0.75 m/s, respectively. However, the
192 vertical resolution was insufficient to measure gusts in the sampled clouds. The accuracy and
193 resolution for temperature measurement are 0.30 and 0.01 C. The accuracy and resolution for
194 relative humidity measurement are 2.0 and 0.1 %. The sampling frequency is 1 Hz.

195

196 **2.2 Inlets**

197

198 Aerosol particles were sampled through a shrouded inlet diffuser (diameter 0.35 cm at intake
199 point), which is the same inlet discussed by Leitch et al. (2010). For the airspeeds during this
200 study, particle transmission by the inlet is near unity for particles from 20 nm to <1 μm . The
201 intake was connected inside the cabin to a 1.9 cm OD stainless steel manifold off of which
202 sample lines were drawn to the various instrument racks using angled inserts. Total flow at the
203 intake point was approximately isokinetic at 55 L min⁻¹ based on the sum of flows drawn by the
204 instrumentation (35 L min⁻¹) and the measured manifold exhaust flow. The manifold exhaust
205 flowed freely into the back of the cabin such that the intake flow varied with aircraft TAS and
206 the manifold was not significantly over pressured.

207 CO was sampled through a separate inlet consisting of a 0.40 cm OD Teflon tube using
208 aircraft forward motion to push air into the line in combination with a rear-facing 0.95 cm OD
209 Teflon exhaust line that reduced the line pressure. The continuously measured sample flow was
210 approximately 12 L min⁻¹.

211

212 **2.3 Data Analysis Approach**

213

214 Eleven research flights were conducted from Resolute Bay, Nunavut (74°40'48"N 94°52'12"W)
215 from July 4, 2014 to July 21, 2014, inclusive. These measurements were associated with two
216 distinct weather regimes. During period 1 (July 4-12), weather conditions around Resolute Bay
217 were affected by an upper low (Supplement Fig. S4). The wind speeds at 500 hPa were mostly
218 light and variable. The surface (1000 hPa) was dominated by weak high-pressure with generally
219 clear skies, light winds, and occasional scattered to broken stratocumulus. Low-cloud or fog was
220 at times present in association with open water, and the air was relatively clean as discussed
221 below. There was a transition period from July 13-16 when flights were not possible due to fog
222 at Resolute Bay. During period 2 (July 17-21), the area came under the influence of a deep low
223 pressure system to the south (Supplement Fig. S5) that brought more wind and higher cloud. The
224 air was not as clean as during period 1, based on the measured aerosol mass and CO
225 concentrations (see Table 1). This was possibly due to transport of BB aerosol from the
226 Northwest Territories; further discussion in Section 2.3.1. Based on the bulk Richardson number
227 and data from radiosondes, Aliabadi et al. (2015) estimated boundary-layer heights at 254 m
228 (± 155 m) across the study.

229 A summary of all flight tracks is shown in Fig. 1. Flights mostly consisted of vertical
230 profiles and low level transits over ice, water and melt ponds that contributed to the formation of
231 low cloud, where low cloud is defined here as cloud tops below 200 m-msl. Higher level cloud
232 was also sampled during the profiles and transits. The polynyas that were sampled over are
233 shown in the top center of each panel of Fig. 2. Cloud was sampled on eight of the 11 flights,
234 more frequently during period 1 because of overall better visual contrast between clouds and
235 surfaces. Furthermore, period 2 was marked by the presence of the Canadian Coast Guard Ship
236 Amundsen in Lancaster Sound (bottom center of each Fig. 2 panel) when the flights were
237 focused on sampling of the ship's plume (e.g. Aliabadi et al., 2016).

238 All aerosol number concentrations are given in terms of standard atmospheric pressure
239 and temperature (STP: 1 atm and 15°C). The CDNC are also referenced to STP where
240 comparisons are made with the aerosol number concentrations. Number concentrations of
241 particles larger than 100 nm (N100) are taken from the UHSAS. All data, except the SMS, are 1
242 second averages that represent a sampling path length of 60–80 m. Size distributions between 20
243 and 100 nm are from the SMS are 1-minute averages. Except for the Fig. S3 example, all particle
244 number concentrations smaller than 100 nm are from the SMS. Nx-100 refers to the number
245 concentration within the interval “x-100” where x ranges between 20 and 90. Values of Nx with
246 $x < 100$ are derived from the sum of Nx-100 (SMS) + N100 (UHSAS).

247 Clouds were sampled during a flight whenever possible, mostly by ascending or
248 descending through them. It was not possible to sample below the low-altitude cloud bases.
249 Most clouds were liquid phase, based on the 2DC-Grey images of cloud particles $>50 \mu\text{m}$, and
250 only liquid phase clouds are discussed here. In addition, none of the liquid clouds exhibited
251 detectable precipitation, except that droplets in a couple of the lowest altitude clouds were very

252 low in number and relatively large in size (30-40 μm); considering the settling speeds of such
253 droplets, they may be viewed as precipitating. The higher-altitude clouds were either stratus or
254 stratocumulus, whereas the low-level clouds were fog or stratus. Although still light, turbulence
255 appeared to be the greatest in the July 7 stratocumulus. Cloud droplet sizes are represented by the
256 volume-weighted mean diameter (VMD), which has the property that the VMD can be used with
257 CDNC to calculate LWC.

258 The pre-cloud aerosol for the HA clouds is mostly derived from averages of values
259 collected within about 50 m of cloud base when a cloud base was visible and achievable. In some
260 cases, as discussed in Section 2.3.1 and 2.3.2, the pre-cloud aerosol concentrations include
261 contributions from above cloud (July 19) or are from similar or lower altitudes in the clear air
262 upwind of the cloud. For the aerosol measurements made with the 1-minute averaged number
263 concentrations from the SMS, values from further below-cloud are necessary in some cases.
264 These values are however consistent with the 1-second aerosol measurements closer to cloud
265 base.

266 Every possible liquid cloud was sampled along a flight path, and some cloud layers were
267 sampled more than once. That will bias the sample numbers to clouds of greater spatial extent.
268 However, that bias is appropriate from a climate perspective since cloud extent is a major factor
269 for the impacts of clouds on climate. A total of 62 liquid water cloud samples, or averages of
270 individual cloud penetrations, were averaged with the constraint that the mean LWC is $> 0.01 \text{ g}$
271 m^{-3} . The samples are integrations over periods ranging from 11 to 1000 seconds with a median
272 sample time of 65 seconds that is equivalent to a horizontal path length of about four kilometers.
273 In sections 2.3.1 and 2.3.2, a range of detailed examples are used to show how the aerosol
274 observations relate to the cloud observations for the higher-altitude (HA) cloud (clouds based

275 above 200 m) and low-altitude (LA) cloud (clouds topped below 200 m), and to 1) demonstrate
276 how the pre-cloud aerosol concentrations were assessed for the 62 samples and 2) note where
277 effects of entrainment may be a factor and how multiple cloud layers are considered. At 200 m
278 or below, the LA clouds were in the boundary layer, in flight indistinguishable from the surface
279 (i.e. some were possibly fog). Thus, sampling below such clouds was not possible due to
280 proximity to the surface. Besides cloud microphysics, the only in-cloud measurements
281 considered valid are the CO and thermodynamics. For completeness, the aerosol measurements
282 within cloud are included in the plots associated with sections 2.3.1 and 2.3.2, but such
283 measurements, including the CCN, are unreliable due to issues of drying and partial drying
284 associated with the inlet and a particular instrument as well as droplet shattering on the inlet (e.g.
285 Hudson and Frisbie [1991] and Hallett and Christensen [1984]). The in-cloud aerosol
286 measurements are not part of the subsequent analysis.

287

288 **2.3.1 Higher Altitude (HA) Cloud Examples**

289

290 Four examples of profiles through HA clouds are shown in Fig. 3. There are two panels for each
291 profile: the left-hand panel shows CO, CDNC and particle number concentrations (N5, Nx-100,
292 N100, CCNC(0.6%)); the right-hand panel shows temperature, equivalent potential temperature
293 (θ_e), LWC and VMD. The temperatures, θ_e and VMD are scaled as indicated.

294 July 7 Case (Fig. 3 a, b): One of several similar profiles through a stratocumulus layer on
295 during the transits to and from the polynyas north of Resolute Bay. The CDNC (at STP) are
296 relatively constant with altitude while LWC and VMD both increase steadily with altitude. These
297 features characterize cloud formation by lifting of air, and they indicate the cloud droplets were

308 nucleated on particles from below cloud base. The cloud top is relatively sharply capped by a
309 temperature inversion of about 2°C at 2350 m, and the particle profiles along with θ_e and CO are
300 relatively constant below cloud base. There is no indication that entrainment, based on the LWC
301 profile, does anything other than reduce the CDNC. Within cloud, the number concentrations of
302 larger particles (N100) is reduced due to nucleation scavenging; although such particles are not
303 completely eliminated as smaller droplets can enter the inlet and dry in the sampling lines.
304 Smaller particles can be artificially increased in cloud due to the shattering of larger droplets on
305 the aerosol intake (e.g. Hudson, 1993), as indicated by the increase in the corresponding N5
306 higher in cloud. Thus, in-cloud aerosol measurements are shown here only for completeness, but
307 they are not used in the subsequent analysis. CDNC range up to 265 cm⁻³ and the mean value is
308 199 cm⁻³. Below cloud base, the N5, N20-100, N30-100, N50-100, N100 and CCNC(0.6%) are
309 approximately 235 cm⁻³, 167 cm⁻³, 145 cm⁻³, 94 cm⁻³, 67 cm⁻³ and 117 cm⁻³, respectively. The
310 below-cloud N20 of 234 cm⁻³ approximately equals the N5 offering confidence in terms of
311 number concentration closure. The N30 (N30-100 + N100) compare most closely with the mean
312 CDNC leading to the conclusion that on average cloud droplets nucleated on particles down to
313 about 30 nm. Based on the maximum CDNC, it is possible that particles as small as 20 nm
314 contributed to the CDNC in this cloud; for 20 nm particles of ammonium sulphate to activate,
315 Köhler equilibrium theory indicates that S in the cloud bases would have had to reach above
316 1.5%.

317 July 17 Case (Fig. 3 c, d): The maximum and mean CDNC (STP), of about 75 cm⁻³ and
318 55 cm⁻³, respectively, are lower while the VMD peak of 20 µm is higher compared with the July
319 7 profile. The LWC are generally similar between July 7 and 17, except that there are more breaks in
320 the July 17 profile. Many of those breaks are due to the aircraft passing through the edges of

321 cloud during this profile. The inversion topping the cloud is weaker and the LWC peak occurs
322 further from cloud top in the July 17 case versus the July 7 case. That LWC feature in
323 combination with the general CO increase, beginning at about 660 m, suggests that the erosion of
324 cloud top by entrainment went deeper into the July 17 cloud. Above 660m, the CDNC also
325 decrease, suggesting the higher concentrations of N50-100, N100 and CCN above cloud relative
326 to below cloud did not enhance the CDNC. Continuity from at least 100 m below cloud base is
327 indicated by the CO and θ_e profiles, and the N50 approximates the mean CDNC and possibly
328 maximum CDNC. The CCNC(0.6%) are 30-40 cm^{-3} below cloud, indicating a S higher than
329 0.6%. The comparison between the July 7 and 17 cases is a specific example of the potential
330 importance of smaller particles for the cloud albedo effect.

331 July 19 Case (Fig. 3 e, f): The July 19 profile includes two cloud layers, one from 1200-
332 1400 m and a second from 1400-1500 m. The layer separation appears in the CO concentrations,
333 which are approximately uniform through the lower layer and increasing in the upper layer. The
334 CO levels of 100+ ppbv in this case are among the highest observed during this study. Transport
335 patterns suggest that BB contributed to this aerosol (Köllner et al., Pollution in the summertime
336 Canadian High Arctic observed during NETCARE 2014: Investigation of origin and
337 composition, in Geophysical Research Abstracts, 17, EGU2015-5951, European Geophysical
338 Union General Assembly 2015, Vienna, Austria, 2015). The mean CDNC (STP) in the lower
339 and upper layers are 239 cm^{-3} and 276 cm^{-3} respectively. The VMD reached 15 μm in the lower
340 layer. The VMD are overall smaller and decrease with altitude in the upper layer, consistent with
341 the lower LWC and higher CDNC. In the upper layer, the CDNC increase from cloud bottom to
342 near cloud top consistent with the increase in aerosol from below the layer to above the layer.
343 The N50 and N100 estimated for the lower (upper) layer are 269 (334) cm^{-3} and 197 (221) cm^{-3}

344 respectively, where the upper layer values are an average of the aerosol at 1400 m and just above
345 cloud top. Thus, on average the CDNC in both layers are approximated by the activation of
346 particles sized between 50 and 100 nm, and the maximum CDNC is approximated by the
347 activation of 50 nm particles. The CCNC(0.6%) are slightly below the N100, which would be
348 consistent with the lower hygroscopicity of BB particles. Comparison of below-cloud
349 CCNC(0.6%) with CDNC suggests cloud S above 0.6%.

350 July 20 Case (Fig. 3 g and h): This is a more complex cloud with substantial LWC
351 variations that suggest three cloud layers. The values of mean CDNC at STP are 45 cm^{-3} , 49 cm^{-3}
352 and 65 cm^{-3} in the upper, middle and lower layers respectively. The VMD reach about $20 \text{ }\mu\text{m}$ in
353 the lower layer and $26 \text{ }\mu\text{m}$ in the upper layer with the lower CDNC. These layers are relatively
354 stable with CO and θ_e increasing slightly from below the cloud to above the top cloud layer.
355 N50 just below the lower layer approximately equals CDNC in that layer. It is more difficult to
356 estimate the pre-cloud aerosol for the middle and upper layers, but particles at least as small as
357 50 nm were apparently activated. For the summary statistics, the respective pre-cloud N100,
358 N50 and CCNC(0.6%) are estimated at 24 cm^{-3} , 44 cm^{-3} and 24 cm^{-3} for the upper cloud layer,
359 32 cm^{-3} , 52 cm^{-3} and 32 cm^{-3} for the middle layer and 34 cm^{-3} , and 66 cm^{-3} and 35 cm^{-3} for the
360 lower layer. Comparison of the CCNC(0.6%), which are in approximately the same
361 concentration as the N100, and CDNC suggests S near or in excess of 0.6%.

362

363 **2.3.2 Low-Altitude (LA) Examples**

364

365 July 5 and July 7 Cases: The two examples in Fig. 4 are for cloud or fog over the polynyas north
366 of Resolute Bay on July 5 and 7. Four cloud samples were collected on July 5 at altitudes below

367 200 m. The time series in Fig. 4a covers the period of collection of the two lowest samples:
368 16:18:02-16:21:57 at 130 m and 16:39:35-16:40:18 at 88 m. In the air upwind of the cloud or
369 fog, the N100, N30 and CCNC(0.6%) are estimated at 3 cm^{-3} , $10\text{-}14 \text{ cm}^{-3}$ and 5 cm^{-3} . The mean
370 values of the CDNC of 2.8 cm^{-3} at 130 m and 0.7 cm^{-3} at 88 m are explained by the N100 and S
371 less than 0.6%. The maximum CDNC of 12 cm^{-3} at 130 m suggests the activation of smaller
372 particles, possibly as small as 30 nm, and S exceeding 0.6% perhaps due to some uplift
373 influenced by orographic features north of the north polynya. At 88 m, the mean VMD (not
374 shown) was $29 \text{ }\mu\text{m}$ and ranged up to $35 \text{ }\mu\text{m}$ giving those droplets potential to deposit over an
375 hour or more, thereby potentially transferring water from the polynya to the downwind ice. On
376 July 7, cloud or fog was present below 120 m and thicker towards the north edge of the north
377 polynya and again to the north over the ice. Seven samples were identified over the period 16:06-
378 16:29 based on the LWC above 0.01 g m^{-3} . The CDNC are overall higher than on July 5 with
379 sample averages ranging from 4 cm^{-3} to 13 cm^{-3} ; the one-second CDNC are as high as 34 cm^{-3}
380 and the mean VMD (not shown) range from $19.6 \text{ }\mu\text{m}$ to $22.8 \text{ }\mu\text{m}$. The CO mixing ratio is slightly
381 higher within the cloud (81 ppbv) than above (79 ppbv); although this difference may not be
382 significant. In the air nearly free of cloud and below 120 m, the N100 are $4\text{-}5 \text{ cm}^{-3}$, the N50 are
383 $8\text{-}11 \text{ cm}^{-3}$ and the N20 are variable between 17 cm^{-3} and 130 cm^{-3} ; CCN are unavailable for this
384 part of the flight. Mean values of CDNC/N100 and CDNC/N50 for seven cloud samples are 4.8
385 and 1.0, respectively, indicating that on average particles of about 50 nm were activated in this
386 LA cloud. Based on the overall relationship between CCNC(0.6%) and N50, which is discussed
387 in section 3.3, the mean S in the LA cloud of July 7 is estimated at 0.6%. Comparison with the
388 maximum CDNC suggests that particles as small as 20 nm may have participated in the
389 nucleation of droplets.

390 July 8 Case: Fig. 5 shows a time series of altitude, CO, N100, N80-100, N90-100,
391 CCNC(0.6%) and CDNC from the sampling above and in the low cloud over Lancaster Sound
392 on July 8. The cloud over the open water of the Sound is visible in the satellite picture in Fig. 2b.
393 Cloud was also present over the ice to the west, but it was much thinner and reached only to
394 about 150 m above the surface. Over the water, the cloud was sampled as high as 230 m by
395 descending into it down to about 150 m between 17:27 UT and 17:43 UT as shown in Fig. 5.
396 Observations in profiles from two of five samples are shown in Fig. 6. This cloud deepened as
397 the aircraft approached the ice edge from over the water, and thinned abruptly over the ice with
398 tops below 150 m as shown in Fig. 5 (time 17:47). The thicker cloud was associated with a shift
399 in wind direction to more southerly suggesting an influence of the Prince Regent Inlet and
400 surrounding terrain on the flow as well as possibly circulations influenced by the water-ice
401 transition. The cloud layer was relatively stable and the θ_e profiles suggest a surface heat sink
402 (Fig. 6a). Profiles of LWC and VMD in Fig. 6 (b, c) do not show increases with altitude
403 characteristic of vertical mixing, such as for some of the HA clouds (Fig. 3); the change in the
404 VMD per 50 m increase in height is about $1.7 \mu\text{m}$ for the well mixed cloud of July 7 (Fig. 3 a, b),
405 whereas it is about $0.2 \mu\text{m}$ per 50 m for the LA cloud of flight 8 in Fig. 6. The CO mixing ratio
406 shows little variation with time and altitude. The pre-cloud aerosol concentrations are more
407 difficult to assess. Based on concentrations just above the cloud, particles $>90 \text{ nm}$ explain the
408 CDNC. Based on the concentrations downwind at 150 m (approximately 17:47), activation of
409 particles $>80 \text{ nm}$ is needed to explain the CDNC. The CCNC(0.6%) are about 129 cm^{-3}
410 downwind and between 157 cm^{-3} and 234 cm^{-3} just above cloud. It is concluded that in this case
411 the droplets likely nucleated on particles mostly larger than 80-95 nm and the S in the clouds
412 were less than 0.6%. For the purposes of summary statistics discussed next, the N100, N50 and

413 CCNC(0.6%) have been selected as an average of the downwind and immediately above cloud
414 concentrations: 73 cm^{-3} , 319 cm^{-3} and 168 cm^{-3} , respectively.

415

416 **3. Summary Observations and Discussion**

417

418 Summary statistics for the cloud and aerosol samples are discussed in 3.1, the microphysics of
419 low-altitude and higher-altitude clouds are contrasted in 3.2, particle activation is summarized in
420 3.3 and in section 3.4 the relationship between VMD and CDNC is used to consider the
421 transition of aerosol indirect effects from potential warming to potential cooling. All analyses
422 are based on the 62 cloud samples discussed in section 2.3. The LA cloud subset is comprised of
423 24 samples and the HA cloud subset consists of 38 samples.

424

425 **3.1 Summary of mean observations**

426

427 The mean and median values of the microphysical properties of the cloud and pre-cloud aerosols
428 as well as the altitudes and temperatures derived from the 62 cloud samples are given in Table 1,
429 separated between periods 1 and 2. Values of the CDNC and the LWC are given relative to in-
430 situ volumes as well as STP. As discussed above, the pre-cloud CCNC(0.6%), N50, and N100
431 are averages of those values collected within about 50 m of cloud base where a cloud base was
432 clear and achievable. In other cases the pre-cloud CCNC(0.6%), N50 and N100 are the values at
433 the similar or lower altitudes in the clear air upwind of the cloud, except in the case of July 8
434 when the pre-cloud aerosol is based on the measurements in the area downwind plus those
435 immediately above cloud. The CCNC(0.6%) samples in Table 1 are limited to 44 due to
436 instrument problems, all of which occurred during the early part of July 7.

437 Cloud liquid water paths (LWP) are estimated for 36 of the samples when a complete
438 profile between cloud base and cloud top was possible. The LWP are shown at the bottom of
439 Table 1. Of the 36 LWP estimates, 34 are above 200 m, and the mean and median altitudes are
440 1044 m and 862 m, respectively. Not included in the summary statistics are the samples from
441 July 8 shown in Figs. 5 and 6. For the minimum altitude reached in that cloud, the LWP ranged
442 from 12 to 25 and thus the total LWP for that cloud exceeded 25.

443 During period 1, the median sampling altitude is lower and the temperatures are slightly
444 below freezing compared with just above freezing during period 2. The CO mixing ratios are
445 overall low and at approximately background values during period 1. The median CDNC are
446 higher during period 1 than period 2, but the mean values are similar. The CDNC compare more
447 closely with the N50 during period 1, while during period 2 the CDNC are about equally
448 between the N50 and N100. The CCNC(0.6%) equated with particles between 50 nm and 100
449 nm during period 1, whereas during period 2 they were closer to the N100 values. The reduction
450 in particle hygroscopicity during period 2 may be due to an increased presence of organics in the
451 aerosol during that time (Willis et al., 2016).

452

453 **3.2 Comparison of LA and HA cloud**

454

455 The LA clouds were close to the surface, and all associated with open water; some or all may be
456 technically fogs. They may be formed by advection of warmer moist air over a cooler surface
457 (the July 8 LA cloud that moved from Baffin Bay westward along Lancaster Sound was likely
458 dominated by that process), by radiation cooling or by the passage of very cold air over a warm
459 moist surface. The latter, also known as sea smoke, is the likely explanation for the clouds over

460 the polynyas; also, it is possible that there was an advection component associated with the sea
461 smoke moving from the polynyas over the ice surfaces. In general, the LA clouds are associated
462 with low-level horizontal advection and heat and water exchange with the underlying ice or
463 water surface. In contrast, vertical motions are responsible for some of the HA clouds, and none
464 of the HA clouds interact so closely with the underlying surface. Due to those differences, the
465 characteristics of the LA and HA clouds are considered separately. Table 2 shows the mean and
466 median values for the samples separated between LA and HA clouds; vertical profiles of CDNC,
467 LWC and VMD samples are shown in Supplement Fig. S7. On average, the LA samples have
468 lower CDNC and higher VMD compared with the HA cases, and the LA clouds are activating on
469 larger particles relative to the HA clouds (e.g. CDNC/N50). The values of the
470 CDNC/CCNC(0.6%) indicate that the S are <0.6% for the LA clouds and close to 0.6% for the
471 HA clouds.

472 Variations in LWC are correlated with those of CDNC for the LA samples (Fig. 7a). The
473 coefficient of determination (R^2) rises from 0.57 to 0.98 if the one LA point at (137, 0.032) is
474 removed. In contrast, the correlation of the LWC with the CDNC for the HA samples is low
475 ($R^2=0.12$). There is no correlation of the LWC with the VMD for the LA points ($R^2=0.04$), and
476 for the HA clouds there is a modest correlation of LWC with MVD ($R^2=0.26$). Variations in
477 LWC with VMD within a cloud system are consistent with lifting of air from below, i.e.
478 nucleation of droplets at cloud base followed by their growth with increasing altitude, such as the
479 case shown in Fig. 3a and 3b. Variations of LWC with VMD can also result from homogeneous
480 mixing (i.e. entrainment of dry air that reduces LWC by partial evaporation of droplets without
481 reducing CDNC). The strong dependence of the variations in LWC with those of the CDNC in
482 the LA clouds may reflect changes in rate of cooling, collision-coalescence or inhomogeneous

483 mixing along the cloud transport pathway. For example, increases in the rate of cooling within or
484 between clouds will increase condensation rates, and potentially S, resulting in increased LWC
485 and CDNC. Changes in collision-coalescence will affect the CDNC and LWC in similar ways:
486 more collision-coalescence, lower CDNC and lower LWC due to precipitation. Inhomogeneous
487 mixing, the entrainment of dry air parcels into a cloud without mixing with the cloud droplets,
488 will reduce the CDNC averaged across the cloud and at the same time reduce the mean LWC.
489 Changes in the aerosol that are interactive with some of the cloud processes may contribute to
490 the CDNC and potentially the LWC through their influence on collision-coalescence.

491 The LWC-CDNC correlation is identifiable for individual flights with sufficient LA
492 samples: four flights, comprising 20 of the 24 LA samples, had three or more points as shown in
493 Fig. 8. The regressions for each of the July 7, 8 and 17 cases are approximately linear, and the
494 respective mean VMD are 20.8 μm , 18.8 μm and 18.2 μm . The mean LWC are 0.05 g m^{-3} , 0.3 g
495 m^{-3} and 0.07 g m^{-3} . The VMD are relatively close together confirming similarities in the
496 relationships, even if not purely linear. For comparison, the mean VMD for the July 5 samples is
497 29.2 μm and the LWC is 0.02 g m^{-3} , which indicates that the July 5 case does not fit the linear
498 relationship shown in Fig. 8. The reasons behind the similarity of the VMD for the July 7, 8 and
499 17 are unknown, but it occurs despite the varied pre-cloud N50 and N100: N50 range of 5-272
500 cm^{-3} ; N100 range of 1.1-73 cm^{-3} . The consistencies among the three flights for greatly differing
501 aerosol and CDNC imply a much smaller role for the aerosol in terms of the LWC. The
502 distributions of droplets extend above 20 μm in these cases, but few are of sufficient size to
503 initiate collision-coalescence (about 30 μm) (e.g. Rosenfeld et al., 2001) unless some fall out
504 already had occurred. Greater temporal and spatial coverage are needed to assess the
505 microphysical processes in these clouds.

506

507 **3.3 Particle Activation Sizes**

508

509 Here, the sizes and CCN activity of particles that acted as nuclei for cloud droplets are examined.

510 The CDNC are plotted versus N100 in Fig. 9a, separated between LA and HA samples. The

511 CDNC are most often higher than the N100 and more so for the HA samples, which indicates

512 that particles smaller than 100 nm activated in most cases and most often in the HA clouds. The

513 mean and median values of CDNC(STP)/N100 are 2.2 and 1.8 for all 62 samples, and the 30th

514 percentile of the CDNC/N100 is 1.2, which means that in about 70 % of the cases droplets

515 nucleated on particles significantly smaller than 100 nm. Fig. 9a can be compared with the

516 results of Hegg et al. (2012) who showed a linear fit of CDNC to N100 for marine stratocumulus

517 with a slope of 0.72 for which the N100 in 94% of the samples was $>150 \text{ cm}^{-3}$. Here, a slope

518 larger than unity is indicated, and the N100 are $<100 \text{ cm}^{-3}$ in 90% of the samples. The

519 comparison indicates that relationships derived for higher concentration environments do not

520 necessarily apply to those of lower concentration environments. In the clean environment often

521 found in the Arctic during summer, the absence of larger particles may lower water uptake rates

522 during droplet nucleation, which will increase the S, enabling cloud droplets to nucleate on

523 smaller particles; the absence of larger particles may also help increase the concentrations of

524 smaller particles in the Arctic during summer, by promoting new particle formation through a

525 reduced condensation sink (e.g. Tunved et al., 2013; Leaitch et al., 2013). The CDNC are plotted

526 against the N50 in Fig. 9b showing that the mean activation size of the HA clouds was often

527 close to 50 nm. The median value of CDNC/N50 is 0.78 for all samples indicating that, based on

528 the averaged CDNC, cloud droplets nucleated on particles near or smaller than 50 nm about 40%

529 of the time. That percentage will increase if particle activation is considered relative to the
530 maximum CDNC.

531 The mean and median values of the CCNC(0.6%) associated with all cloud samples (84
532 cm^{-3} and 47 cm^{-3}) are generally consistent with previous Arctic CCNC measurements. For
533 example, during the summer above 85°N , Martin et al. (2011) measured a mean CCNC at 0.73%
534 S of 47 cm^{-3} with a standard deviation of 35 cm^{-3} , Yum and Hudson (2001) measured CCNC at
535 0.8% S below 1700 m over the Beaufort Sea during May, 1998 that ranged from 41 cm^{-3} to 290
536 cm^{-3} , and Radke et al. (1976) measured a mean CCNC at 1% S of 90 cm^{-3} in June near Barrow,
537 Alaska. Considering the median values of CDNC/CCNC(0.6%) for the LA and HA samples
538 (Table 2) and the slopes of linear regressions of CDNC versus CCNC(0.6%) (Fig. 10a), the
539 average inferred S for the HA clouds is about 0.6%, consistent with the overall activation of
540 smaller particles in those clouds. The mean S inferred for the LA clouds is significantly lower
541 than 0.6%. Based on the activation of a 90 nm particle (July 8 case; CCNC(0.6%) of 168 cm^{-3} in
542 Fig. 10a) of low-moderate hygroscopicity, a reasonable estimate is 0.3% for the mean of the LA
543 clouds with some higher values indicated by the points near a CCNC(0.6%) of 25 cm^{-3} in Fig.
544 10a. The S for these clean clouds are in contrast to polluted marine environments for which
545 estimates for these types of clouds are 0.2% or less (e.g. Modini et al., 2015). Consistent with the
546 present results, Hudson et al. (2010) found that effective S in marine stratus tended to increase
547 with a decrease in the CCNC, and for CCNC smaller than about 200 cm^{-3} their effective S ranged
548 between 0.3% and 1.2%.

549 Variations in the measured CCNC(0.6%) are explained well by variations in smaller
550 (N50) and larger (N100) particles as shown in Fig. 10b. The slopes of the power-law fits, for

551 which the exponents are both close to unity, indicate that the CCNC(0.6%) at 0.6% S on average
552 fall between 50 nm and 100nm.

553

554 **3.4 Aerosol Influences on Warming to Cooling**

555

556 The relationship between the VMD and CDNC shown in Fig. 11 exhibits a scattered but clear
557 tendency for smaller VMD with increasing CDNC. The solid black curve is a reference line
558 based on the study-mean LWC of 0.12 g m^{-3} (Table 1); points falling above or below the black
559 curve have higher or lower LWC, respectively. The vertical dashed green line represents our best
560 estimate of the Mauritsen limit below which Mauritsen et al. (2011) showed the cloud may
561 produce a net warming for an increase in the CDNC. The net warming is a consequence of an
562 increase in longwave absorption due to an increase in the LWC, where the latter results from a
563 reduction in deposition for the smaller droplets associated with increased CDNC. A value of 16
564 cm^{-3} is our best estimate of the Mauritsen limit for this data set because all points with CDNC
565 below that value fall well below the mean LWC, therefore offering greater potential for changes
566 in the CDNC to increase the LWC. Above the estimated Mauritsen limit, an increase in CDNC
567 may produce a net cooling due to the cloud albedo effect, since at that point the longwave
568 forcing does not change significantly as the effects of deposition are minimized and the cloud
569 effectively behaves as a black body.

570 The aerosol influence on clouds with CDNC below the Mauritsen-limit is considered in
571 section 3.4.1. In section 3.4.2, the potential background influence of the aerosol on clouds with
572 CDNC above the Mauritsen-limit is examined.

573

574 **3.4.1 Below the Mauritsen limit**

575
576 Seventeen of the 62 samples fall at or below our best estimate of the Mauritsen limit. Fifteen of
577 those 17 samples are from LA clouds with median pre-cloud N50 and N100 estimates of 8.2 cm^{-3}
578 and 3.0 cm^{-3} respectively. The lower number concentrations contribute to overall larger VMD;
579 although some of the points below the estimated Mauritsen limit have VMD values much less
580 than $20 \text{ }\mu\text{m}$. Increases in small particles, potentially from particle nucleation or fragmentation
581 (e.g. Leck and Bigg, 1999 and 2010), are hypothesized to increase the CDNC thereby enhancing
582 longwave warming by these clouds, at least until the CDNC exceed the estimated Mauritsen
583 limit. The LA points from the July 5 and the July 7 cases, identified in Fig. 11, offer one insight.
584 The median CDNC for July 5 is six times lower than the July 7 CDNC: 1.3 cm^{-3} and 7.8 cm^{-3} , for
585 July 5 and 7, respectively. The median N50 are 6 cm^{-3} and 8.3 cm^{-3} for July 5 and 7,
586 respectively, and the median N100 are 3 cm^{-3} and 2.2 cm^{-3} for July 5 and July 7, respectively.
587 The CDNC are similar to N50 in the July 7 case, but lower than both the N50 and N100 in the
588 July 5 case indicating that the aerosol was not a limiting factor in the July 5 case. Consistent with
589 the discussion in section 3.2, all 15 LA points show a correlation of LWC with the CDNC
590 ($R^2=0.57$), but correlations of CDNC with N50 and N100 are weak: $R^2=0.19$ and 0.06 ,
591 respectively. The CCN are not used here because only seven points with CCNC(0.6%) are
592 available; the seven do, however, correlate well with the N50. If the limit of 10 cm^{-3} of
593 Mauritsen et al. (2011) is applied, reducing the number of points to 12, the assessment does not
594 change: the LWC-CDNC correlation improves slightly and the correlations of the CDNC with
595 the N100 and the N50 weaken.

596 The LWC do not correlate with either the N50 or the N100 (Supplement Fig. 8). In this
597 low CDNC environment, where cloud droplets may grow large enough to be gravitationally

598 removed from the cloud without the support of collision-coalescence, the absence of a positive
599 correlation of either the CDNC or LWC with the aerosol indicates that small changes in the
600 aerosol did not contribute significantly to the changes in the LWC. Variations in other
601 processes, such as mixing or the rate of cooling, may be responsible for the correlation of CDNC
602 and LWC. It can be argued that some aerosol must exist for these clouds to form, but these
603 observations show no association of changes in either the CDNC or LWC with changes in the
604 aerosol.

605

606 **3.4.2 Background aerosol influence on clouds**

607

608 Above the estimated Mauritsen limit, the general reduction in the VMD with the CCNC(0.6%)-
609 associated increase in CDNC reflects the impact of increases in aerosol on clouds. In Fig. 11,
610 samples are identified between those associated with lower CO (green circles; <81 ppbv, the
611 median CO value of all samples) and those with highest CO (red circles; >90 ppbv); six samples
612 have no CO measurement and the remaining points have CO falling within 81-90 ppbv. Five of
613 the seven higher-CO samples are from the July 19 case (e.g. Fig. 3e, 3f) that has been linked with
614 BB (Köllner et al., 2015; reference above), and the highest CDNC point (273 cm^{-3} ; no CO
615 measurement) is also from July 19 and likely influenced by BB. The higher-CO samples cover a
616 range of CDNC from 16 cm^{-3} to at least 238 cm^{-3} with CO reaching up to 113 ppbv. The higher
617 CO samples are associated with larger particles ($N_{50}/N_{100}=1.5$), consistent with a BB influence,
618 compared with the lower CO samples ($N_{50}/N_{100}=3.2$). These values for BB fall at the low end
619 of the observations from Zamora et al. (2015), but their CO concentrations are much higher than
620 those measured in this study. The lower-CO samples may be dominated by regional biogenic

621 emissions (Willis et al., 2016). The lower- and higher-CO points overlap over a CDNC range of
622 16 cm^{-3} to 160 cm^{-3} , consistent with the range of pre-industrial CDNC from global models of 30
623 cm^{-3} to 140 cm^{-3} (Penner et al., 2006; Korhonen et al., 2008). In this clean environment, the
624 contributions from 20-100 nm particles have a broad impact on the range of CDNC, affirming
625 the large uncertainty associated with estimating a baseline for the cloud albedo effect discussed
626 by Carslaw et al. (2013).

627

628 **4. Summary and Conclusions**

629

630 Aerosol particle size distributions, CCNC(0.6%) at 0.6% water S, carbon monoxide (CO) and
631 cloud microphysics were measured from an airborne platform based out of Resolute Bay,
632 Nunavut from July 4 to July, 21, 2014 as one part of the Canadian NETCARE project. The
633 flights were conducted over ice and water surfaces from about 60 m above the surface to about
634 6000 m. Sixty-two (62) cloud-averaged samples were derived, each constrained for the mean
635 $\text{LWC} > 0.01 \text{ g m}^{-3}$ or the cloud threshold used here. The analysis separates the cloud samples
636 between 24 low-altitude (LA: $< 200 \text{ m}$) samples and 38 higher altitude (HA: $> 200 \text{ m}$) samples as
637 well as situations of lower and higher CO and observations above and below the Mauritsen et al.,
638 (2011) CCNC(0.6%) (or CDNC) limit.

639 The median pre-cloud N100 of 33 cm^{-3} and the median CO mixing ratio of 81 ppbv
640 indicate that the aerosols supporting the sampled clouds were relatively clean, and particularly
641 during the first part of the study many of the aerosol particles may have been derived from
642 regional natural sources. The median CDNC at STP is 10 cm^{-3} for the LA clouds (24 samples)
643 and 101 cm^{-3} for the HA clouds (38 samples), which correspond with the median pre-cloud N50

644 of 11 cm^{-3} for the LA samples and 133 cm^{-3} for the HA samples. The lower sizes of particles
645 activated in cloud varied from about 20 nm to above 100 nm. In 40% of cases, the average lower
646 size of activation was 50 nm or smaller. Overall, smaller particles were activated more often in
647 the HA clouds. Variations in particle chemistry will induce some variance in these results, but
648 because activation diameters are estimated starting with larger particles and moving to smaller
649 sizes, changes in chemistry only offer the possibility of activation of particles still smaller than
650 estimated here; although that would have to occur at the expense of larger particles.

651 From the median values of CDNC/CCNC(0.6%) (1.2 for the HA clouds and 0.6 for the
652 LA clouds) and the linear regression of CDNC and CCNC(0.6%), it is inferred that the average S
653 were approximately 0.6% for the HA clouds and 0.3% for the LA clouds. Higher estimates will
654 be obtained if the maximum CDNC are taken into consideration rather than the mean CDNC.
655 The relatively high S for these clean Arctic stratus and stratocumulus have similarities with the
656 observations of Hudson et al. (2010) for relatively clean stratus off the coast of California.

657 In 17 cases, 15 of which are LA clouds, the CDNC fell at or below the CCN limit
658 discussed by Mauritsen et al. (2011), which is estimated here as 16 cm^{-3} . These are the first
659 collection of simultaneous observations of the microphysics of aerosols and clouds in this unique
660 regime in which the net radiative impact of increases in the CDNC is hypothesized to be
661 warming due to changes in the LWC. The LWC of the points below the Mauritsen limit all fall
662 below the study-mean LWC, and the LWC increases with the CDNC. Neither the CDNC nor the
663 LWC are positively correlated with the pre-cloud aerosol (N50 or N100). In this environment of
664 low cloud or fog and ultra-low CDNC, variations in cloud processes such as mixing or the rate of
665 cooling may be responsible for the correlation of CDNC and LWC. These observations show no

666 association of changes in either the CDNC or LWC with changes in the aerosol within the
667 Mauritsen limit.

668 Forty-five cloud samples with CDNC above the Mauritsen limit exhibit a clear influence
669 of changing aerosol. The cloud microphysics for the clouds formed in cleaner air (smaller
670 particles and lower CO: <81 ppbv) overlap with clouds formed in what was likely more polluted
671 air (larger particles and higher CO: >90 ppbv) covering a CDNC range of 16-160 cm⁻³. It is
672 concluded that 20-100 nm particles from natural sources can have a broad impact on the range of
673 CDNC in clean environments, affirming a large uncertainty in estimating a baseline for the cloud
674 albedo effect.

675

676

677 *Acknowledgements.* The complete data set is available from the NETCARE web site by
678 contacting Richard Leitch (Richard.Leitch@ec.gc.ca) or Jon Abbatt
679 (jabbatt@chem.utoronto.ca). A spreadsheet containing the details of the 62 samples discussed
680 here is included with the supplement. The authors acknowledge a large number of people for
681 their contributions to this work. We thank Kenn Borek Air, in particular Kevin Elke and John
682 Bayes for their skillful piloting that facilitated these cloud observations. We are grateful to John
683 Ford, David Heath and the U of Toronto machine shop, Jim Hodgson and Lake Central Air
684 Services in Muskoka, Jim Watson (Scale Modelbuilders, Inc.), Julia Binder and Martin
685 Gerhmann (AWI), Mike Harwood and Andrew Elford (EC), for their support of the integration
686 of the instrumentation and aircraft. We thank Mohammed Wasey for his support of the
687 instrumentation during the integration and in the field. We are grateful to Carrie Taylor (EC),
688 Bob Christensen (U of T), Kevin Riehl (Kenn Borek Air), Lukas Kandora, Manuel Sellmann and
689 Jens Herrmann (AWI), Desiree Toom, Sangeeta Sharma, Dan Veber, Andrew Platt, Anne Marie
690 Macdonald, Ralf Staebler and Maurice Watt (EC), Kathy Law and Jennie Thomas (LATMOS)
691 for their support of the study. We thank the Biogeochemistry department of MPIC for providing
692 the CO instrument and Dieter Scharffe for his support during the preparation phase of the
693 campaign. We thank the Nunavut Research Institute and the Nunavut Impact Review Board for
694 licensing the study. Logistical support in Resolute Bay was provided by the Polar Continental
695 Shelf Project (PCSP) of Natural Resources Canada under PCSP Field Project #218-14, and we
696 are particularly grateful to Tim McCagherty and Jodi MacGregor of the PCSP. Funding for this
697 work was provided by the Natural Sciences and Engineering Research Council of Canada
698 through the NETCARE project of the Climate Change and Atmospheric Research Program, the
699 Alfred Wegener Institute and Environment Canada.

700 **References**

- 701
- 702 Aliabadi, A. A., Staebler, R. M., de Grandpré, J., Zadra, A., and Vaillancourt, P. A.: Comparison
703 of Estimated Atmospheric Boundary Layer Mixing Height in the Arctic and Southern Great
704 Plains under Statically Stable Conditions: Experimental and Numerical Aspects, *Atmos-Ocean.*,
705 In Press, doi:10.1080/07055900.2015. 1119100, 2015.
- 706
- 707 Aliabadi, A.A., J. L. Thomas, A. Herber, R. M. Staebler, W. R. Leitch, K. S. Law, L. Marelle, J.
708 Burkart, M. Willis, J. P. D. Abbatt, H. Bozem, P. Hoor, F. Köllner, J. Schneider, and M.
709 Lévassieur. Ship emissions measurement in the Arctic from plume intercepts of the Canadian
710 Coast Guard Amundsen icebreaker from the Polar 6 aircraft platform. *Atmos. Chem. Phys.*
711 *Discuss.*, doi: 10.5194/acp-2015-1032, 2016.
- 712
- 713 Barrie, L.A.: Arctic air pollution: An overview of current knowledge, *Atmos. Environ.*, 20, 643–
714 663, doi:10.1016/0004-6981(86)90180-0, 1986.
- 715
- 716 Barrie, L.A., Bottenheim, J.W., Schnell, R.C., Crutzen, P.J., and Rasmussen, R.A.: Ozone
717 depletion and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*,
718 334, 138-141, 1998.
- 719
- 720 Brenner, T.C., Curry, J.A., and Pinto, J.O.: Radiative transfer in the summertime Arctic, *J.*
721 *Geophys. Res.*, 106, 15173-15183, 2001.
- 722
- 723 Brock, C.A., Cozic, J., Bahreini, R., Froyd, K.D., Middlebrook, A.M., McComiskey, A.,
724 Brioude, J., Cooper, O.R., Stohl, A., Aikin, K.C., de Gouw, J.A., Fahey, D.W., Ferrare, R.A.,
725 Gao, R.-S., Gore, W., Holloway, J.S., Hubler, G., Jefferson, A., Lack, D.A., Lance, S., Moore,
726 R.H., Murphy, D.M., Nenes, A., Novelli, P.C., Nowak, J.B., Ogren, J.A., Peischl, J., Pierce,
727 R.B., Pilewskie, P., Quinn, P.K., Ryerson, T.B., Schmidt, K.S., Schwarz, J.P., Sodemann, H.,
728 Spackman, J.R., Stark, H., Thomson, D.S., Thornberry, T., Veres, P., Watts, L.A., Warneke, C.,
729 and Wollny, A.G.: Characteristics, sources, and transport of aerosols measured in spring 2008
730 during the aerosol, radiation, and cloud processes affecting Arctic climate (ARCPAC) project,
731 *Atmos. Chem. Phys.*, 11, 2423–2453, 2011, doi:10.5194/acp-11-2423-2011, 2011.
- 732
- 733 Browse, J., Carslaw, K.S., Mann, G.W., Birch, C.E., Arnold, S.R., and Leck, C.: The complex
734 response of Arctic aerosol to sea-ice retreat, *Atmos. Chem. Phys.*, 14, 7543-7557,
735 doi:10.5194/acp-14-7543-2014, 2014.
- 736
- 737 Burkart, J., Willis, M., Köllner, F., Schneider, J., Bozem, H., Hoor, P.,
738 Ghahremaninezhadgharelar, R., Wentworth, G., Norman, A.L., Brauner, R., Konrad, C., Herber,
739 A., Leitch, R., and Abbatt, J.P.D.: Evidence for New Particle Formation in the Summertime
740 Arctic near Resolute Bay, Nunavut, Canada, presented at the 7th Symposium on Aerosol–Cloud–
741 Climate Interactions, American Meteorological Society, Phoenix, Arizona, 2015.
- 742
- 743 Cai, Y., Montague, D.C., Mooiweer-Bryan, W., Deshler, T.: Performance characteristics of the
744 ultra-high sensitivity aerosol spectrometer for particles between 55 and 800 nm: Laboratory and
745 field studies, *J. Aerosol Sci.*, 39, 759 – 769, 2008.

746
747 Carslaw, K.S., Lee, L.A., Reddington, C.L., Pringle, K.J., Rap, A., Forster, P.M., Mann, G.W.,
748 Spracklen, D.V., Woodhouse, M.T., Regayre, L.A., and Pierce, J.R.: Large contribution of
749 natural aerosols to uncertainty in indirect forcing. *Nature*, 503, doi:10.1038/nature12674, 2013.
750
751 Christensen, J. H. et al.: Chapter 14, in *Climate Change 2013, The Physical Science Basis*
752 (editors Stocker, T. F. et al.), Intergovernmental Panel on Climate Change, Cambridge Univ.
753 Press, Cambridge, U.K, 2013.
754
755 Curry, J.A.: Introduction to special section: FIRE Arctic clouds experiment, *J. Geophys. Res.*,
756 106, 14,985-14,987, 2001.
757
758 Earle, M. E., Liu, P.S.K., Strapp, J.W., Zelenyuk, A., Imre, D., McFarquhar, D.M., Shantz, N.C.,
759 and Leaitch, W.R.: Factors influencing the microphysics and radiative properties of liquid-
760 dominated Arctic clouds: Insight from observations of aerosol and clouds during ISDAC (2008),
761 *J. Geophys. Res.*, 116, D00T09, doi:10.1029/2011JD015887, 2011.
762
763 Engvall, A-C, Krejci, R., Strom, J., Treffeisen, R., Scheele, R.: Changes in aerosol properties
764 during spring-summer period in the Arctic troposphere, *Atmos. Chem. Phys.*, 8, 445–462,
765 doi:10.5194/acp-8-445-2008, 2008.
766
767 Garrett, T. J., Zhao, C., Dong, X., Mace, G.G., Hobbs, P.V.: Effects of varying aerosol regimes
768 on low-level Arctic stratus, *Geophys. Res. Lett.*, 31, L17105, doi:10.1029/2004GL019928, 2004.
769
770 Garrett, T. J., Maestas, M.M., Krueger, S.K., and Schmidt C.T.: Acceleration by aerosol of a
771 radiative-thermodynamic cloud feedback influencing Arctic surface warming, *Geophys. Res.*
772 *Lett.*, 36, L19804, doi:10.1029/2009GL040195, 2009.
773
774 Hallett, J., and L. Christensen: The splashing and penetration of raindrops into water, *J. Rech.*
775 *Atmos.* 18, 226-242, 1984.
776
777 Heintzenberg, J., and Leck C.: Seasonal-variation of the atmospheric aerosol near the top of the
778 marine boundary layer over Spitsbergen related to the Arctic sulphur cycle, *Tellus B*, 46, 52–67,
779 doi.: 10.1034/j.1600-0889.1994.00005.x, 1994.
780
781 Heintzenberg, J., Leck, C., Birmili, W., Wehner, B., Tjernström, M., and Wiedensohler, A.:
782 Aerosol number-size distributions during clear and fog periods in the summer high Arctic: 1991,
783 1996, and 2001, *Tellus*, 58B, 41–50, 2006.
784
785 Heintzenberg, J., Leck, C., and Tunved, P.: Potential source regions and processes of aerosol in
786 the summer Arctic, *Atmos. Chem. Phys.*, 15, 6487–6502, doi:10.5194/acp-15-6487-2015, 2015.
787
788 Hegg, D.A., Covert, D.S. Jonsson, H.H., and Woods, R.K.: A simple relationship between cloud
789 drop number concentration and precursor aerosol concentration for the regions of earth’s large
790 marine stratocumulus decks, *Atmos. Chem. Phys.*, 12, 1229–1238, doi:10.5194/acp-12-1229-
791 2012, 2012.

792
793 Herman, G.F.: Solar radiation in summertime Arctic stratus clouds. *J. Atmos. Sci.*, 34, 1423-
794 1432, 1977.
795
796 Hobbs, P.V. and Rango, A.L.: Microstructures of low and middle-level clouds over the Beaufort
797 Sea, *Q. J. R. Meteorol. Soc.*, 124, 2035-2071, 1998.
798
799 Hudson, J.G. and P.R. Frisbie: Cloud condensation nuclei near marine stratus. *J. Geophys. Res.*,
800 96, 20,795-20,808, 1991.
801
802 Hudson, J.G.: Cloud condensation nuclei near marine cumulus. *J. Geophys. Res.*, 98, 2693-2702,
803 1993.
804
805 Hudson, J.G., Noble, S., and Jha, V.: Stratus cloud S, *Geophys. Res. Lett.*, 37, L21813,
806 doi:10.1029/2010GL045197, 2010.
807
808 Intrieri, J. M., Shupe, M.D., Uttal, T., and McCarty, B.J.: An annual cycle of Arctic cloud
809 characteristics observed by radar and lidar at SHEBA, *J. Geophys. Res.*, 107, C10, 8030, doi.
810 10.1029/2000JC000423, 2002.
811
812 Jacob, D.J., Crawford, J. H., Maring, H., Clarke, A.D., Dibb, J.E., Emmons, L.K., Ferrare, R.A.,
813 Hostetler, C.A., Russell, P.B., Singh, H.B., Thompson, A.M., Shaw, G.E., McCauley, E.,
814 Pederson, J.R., and Fisher J.A.: The Arctic research of the composition of the troposphere from
815 aircraft and satellites (ARCTAS) mission: Design, execution, and first results, *Atmos. Chem.*
816 *Phys.*, 10, 5191–5212, doi:10.5194/acp-10-5191-2010, 2010.
817
818 Jouan, C., Pelon, J., Girard, E., Ancellet, G., Blanchet, J.P., and Delanoë, J.: On the relationship
819 between Arctic ice clouds and polluted air masses over the North Slope of Alaska in April 2008,
820 *Atmos. Chem. Phys.*, 14, 1205–1224, doi:10.5194/acp-14-1205-2014, 2014.
821
822 Klingebiel, M., de Lozar, A., Molleker, S., Weigel, R., Roth, A., Schmidt, L., Meyer, J., Ehrlich,
823 A., Neuber, R., Wendisch, M., and Borrmann, S.: Arctic low-level boundary layer clouds: In situ
824 measurements and simulations of mono- and bimodal supercooled droplet size distributions at
825 the top layer of liquid phase clouds, *Atmos. Chem. Phys.*, 15, 617–631, doi:10.5194/acp-15-617-
826 2015, 2015.
827
828 Korhonen, H., K. S. Carslaw, D. V. Spracklen, D. A. Ridley, and J. Ström: A global model study
829 of processes controlling aerosol size distributions in the Arctic spring and summer, *J. Geophys.*
830 *Res.*, 113, D08211, doi:10.1029/2007JD009114, 2008.
831
832 Korolev, A.V., Emery, E.F., Strapp, J.W., Cober, S.G., Isaac, G.A., Wasey, M., and Marcotte,
833 D.: Small ice particles in tropospheric clouds: fact or artifact? Airborne icing instrumentation
834 evaluation experiment, *B. Am. Meteorol. Soc.*, 92, 967–973, 2011.
835
836 Lance, S., Shupe, M.D., Feingold, G., Brock, C.A., Cozic, J., Holloway, J.S., Moore, R.H.,
837 Nenes, A., Schwarz, J.P., Spackman, J.R., Froyd, K.D., Murphy, D.M., Brioude, J., Cooper,

838 O.R., Stohl, A., and Burkhardt J.F.: Cloud condensation nuclei as a modulator of ice processes in
839 Arctic mixed-phase clouds, *Atmos. Chem. Phys.*, 11, 8003–8015, doi:10.5194/acp-11-8003-
840 2011, 2011.

841

842 Law, K.S., and Stohl, A.: Arctic air pollution: origins and impacts, *Science*, 315, 1537–1540,
843 2007.

844

845 Leaitch W. R., Lohmann, U., Russell, L.M., Garrett, T., Shantz, N.C., Toom-Sauntry, D., Strapp,
846 J.W., Hayden, K.L., Marshall, J., Wolde, M., Worsnop, D.R., Jayne, J.T.: Cloud albedo increase
847 from carbonaceous aerosol, *Atmos. Chem. Phys.*, 10, 7669–7684, doi:10.5194/acp-10-7669-
848 2010, 2010.

849

850 Leaitch, W.R., Sharma, S., Huang, L., Macdonald, A.M., Toom-Sauntry, D., Chivulescu, A., von
851 Salzen, K., Pierce, J.R., Shantz, N.C., Bertram, A., Schroder, J., Norman, A.-L., and Chang
852 R.Y.-W.: Dimethyl sulphide control of the clean summertime Arctic aerosol and cloud,
853 *Elementa: Science of the Anthropocene*, 1, 000017, doi: 10.12952/journal.elementa.000017,
854 2013.

855

856 Leck C., and Bigg, E.K.: Aerosol production over remote marine areas - A new route, *Geophys.*
857 *Res. Lett.*, 26, 3577–3580, 1999.

858

859 Leck C., and Bigg, E. K.: New particle formation of marine biological origin, *Aerosol Sci.*
860 *Technol.*, 44, 570–577, doi:10.1080/02786826.2010.481222, 2010.

861

862 Lohmann, U., Humble, J., Leaitch, R., Isaac, G., and Gultepe, I.: Simulations of ice clouds
863 during FIRE.ACE using the CCCMA single column model, *J. Geophys. Res.*, 106, 15123–15138,
864 2001.

865

866 Lohmann and Leck? Lohmann, U. and Leck, C.: Importance of submicron surface active organic
867 aerosols for pristine Arctic clouds, *Tellus B*, 57, 261–268, 2005.

868

869 Lubin, D. and Vogelmann, A.M.: Observational quantification of a total aerosol indirect
870 effect in the Arctic, *Tellus*, 62B, 181–189, DOI: 10.1111/j.1600-0889.2010.00460.x, 2010.

871

872 Martin, M., Chang, R.Y.-W., Sierau, B., Sjogren, S., Swietlicki, E., Abbatt, J.P.D., Leck, C., and
873 Lohmann, U.: Cloud condensation nuclei closure study on summer arctic aerosol, *Atmos. Chem.*
874 *Phys.*, 11, 11335–11350, doi:10.5194/acp-11-11335-2011, 2011.

875

876 Maslanik, J., Stroeve, J., Fowler, C., and Emery, W.: Distribution and trends in Arctic sea ice age
877 through spring 2011, *Geophys. Res. Lett.*, 38, L13502, doi:10.1029/2011gl047735, 2011, 2011.

878

879 Mauritsen, T., Sedlar, J., Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau,
880 B., Persson, P.O.G., Brooks, I.M., and Swietlicki E.: An Arctic CCN-limited cloud-aerosol
881 regime, *Atmos. Chem. Phys.*, 11, 165–173, doi:10.5194/acp-11-165-2011, 2011.

882

883 Modini, R.L., Frossard, A.A., Ahlm, L., Russell, L.M., Corrigan, C.E., Roberts, G.C., Hawkins,
884 L.N., Schroder, J.C., Bertram, A.K. Zhao, R., Lee, A.K.Y., Abbatt, J.P.D., Lin, J., Nenes, A.,
885 Wang, Z., Wonaschütz, A., Sorooshian, A., Noone, K.J., Jonsson, H., Seinfeld, J.H.,
886 Toom-Saunty, D., Macdonald, A.M., and Leaitch W.R.: Primary marine aerosol-cloud
887 interactions off the coast of California, *J. Geophys. Res. Atmos.*, 120, 4282–4303,
888 doi:10.1002/2014JD022963, 2015.

889

890 Morrison, H., de Boer, G., Feingold, G., Harrington, J., Shupe, M.D., and Sulia K.: Resilience of
891 persistent Arctic mixed-phase clouds, *Nat. Geosci.*, 5, 11–17, doi:10.1038/ngeo1332, 2012.

892

893 Najafi, M. R., Zwiers, F.W. and Gillett N.P.: Attribution of Arctic temperature change to
894 greenhouse-gas and aerosol influences. *Nature Climate Change*, doi: 10.1038/NCLIMATE2524,
895 2015.

896

897 Peng, Y., Lohmann, U., and Leaitch, R.: The cloud optical depth – cloud droplet effective radius
898 relationship for clean and polluted clouds from RACE and FIRE.ACE. *J. Geophys. Res.*, 107,
899 4106, 10.1029/2000JD000281, 2002.

900

901 Penner J.E., Quaas, J., Storelvmo, T., Takemura, T., Boucher, O., Guo, H., Kirkevåg, A.,
902 Kristjánsson, J.E., and Seland, Ø.: Model intercomparison of indirect aerosol effects, *Atmos.*
903 *Chem. Phys.*, 6, 3391–3405. doi:10.5194/acp-6-3391-2006, 2006.

904

905 Quinn, P.K., Bates, T.S., Baum, E., Doubleday, N., Fiore, A.M., Flanner, M., Fridlind, A.,
906 Garrett, T.J., Koch, D., Menon, S., Shindell, D., Stohl, A., and Warren S.G.: Short-lived
907 pollutants in the Arctic: Their climate impact and possible mitigation strategies, *Atmos. Chem.*
908 *Phys.*, 8, 1723–1735, doi:10.5194/acp-8-1723-2008, 2008.

909

910 Radke et al.,

911

912 Rosenfeld, D., Rudich, Y., and Lahav, R.: Desert dust suppressing precipitation: A possible
913 desertification feedback loop, *Proc. Natl. Acad. Sciences*, 98, 5975–5980, 2001.

914

915 Sandvik, A., Biryulina, M., Kvamsto, N., Stamnes, J., and Stamnes, K.: Observed and simulated
916 microphysical composition of Arctic clouds: Data properties and model validation, *J. Geophys.*
917 *Res.*, 112, D05205, doi:10.1029/2006JD007351, 2007.

918

919 Shindell, D.T., Chin, M., Dentener, F., Doherty, R.M., Faluvegi, G., Fiore, A.M., Hess, P., Koch,
920 D.M., MacKenzie, I.A., Sanderson, M.G., Schultz, M.G., Schulz, M., Stevenson, D.S., Teich, H.,
921 Textor, C., Wild, O., Bergmann, D.J., Bey, I., Bian, H., Cuvelier, C., Duncan, B.N., Folberth, G.,
922 Horowitz, L.W., Jonson, J., Kaminski, J.W., Marmor, E., Park, R., Pringle, K.J., Schroeder, S.,
923 Szopa, S., Takemura, T., Zeng, G., Keating, T.J., and Zuber, A.: A multi-model assessment of
924 pollution transport to the Arctic, *Atmos. Chem. Phys.*, 8, 5353–5372, doi:10.5194/acp-8-5353-
925 2008, 2008.

926

927 Shupe, M.D., Kollias, P., Matrosov, S.Y., and Schneider, T.L.: Deriving mixed-phase cloud
928 properties from Doppler radar spectra, *J. Atmos. Ocean. Technol.*, 21, 660-670, 2004.

929
930 Shupe, M.D., Persson, P.O.G., Brooks, I.M., Tjernström, M., Sedlar, J., Mauritsen, T., Sjogren,
931 S., and Leck, C.: Cloud and boundary layer interactions over the Arctic sea ice in late summer,
932 *Atmos. Chem. Phys.*, 13, 9379–9400, doi:10.5194/acp-13-9379-2013, 2013.
933
934 Stohl A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys. Res.*,
935 111, D11306, doi:10.1029/2005JD006888, 2006.
936
937 Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V.P., Kopeikin, V.M., and
938 Novigatsky, A.N.: Black carbon in the Arctic: the underestimated role of gas flaring and
939 residential combustion emissions, *Atmos. Chem. Phys.*, 13, 8833–8855, doi:10.5194/acp-13-
940 8833-2013, 2013.
941
942 Ström, J., Umegård, J., Tørseth, K., Tunved, P., Hansson, H.-C., Holmém, K., Wismann, V.,
943 Herber, A., König-Langlo, G.: One year of particle size distribution and aerosol chemical
944 composition measurements at the Zeppelin Station, Svalbard, March 2000–March 2001, *Physics
945 and Chemistry of the Earth* 28 (2003) 1181–1190, 2003.
946
947 Tjernström, M., Leck, C., Persson, P.O.G., Jensen, M.L., Oncley, S.P., and Targino, A.: The
948 summertime Arctic atmosphere: Meteorological measurements during the Arctic Ocean
949 experiment 2001 (AOE-2001), *B. Am. Meteor. Soc.*, 85, 1305–1321, 2004, 2004.
950
951 Tjernström, M., Leck, C., Birch, C.E., Bottenheim, J.W., Brooks, B.J., Brooks, I.M., Bäcklin, L.,
952 Chang, R.Y.-W., de Leeuw, G., Di Liberto, L., de la Rosa, S., Granath, E., Graus, M., Hansel,
953 A., Heintzenberg, J., Held, A., Hind, A., Johnston, P., Knulst, J., Martin, M., Matrai, P.A.,
954 Mauritsen, T., Müller, M., Norris, S.J., Orellana, M.V., Orsini, D.A., Paatero, J., Persson,
955 P.O.G., Q. Gao, Rauschenberg, C., Ristovski, Z., Sedlar, J., Shupe, M.D., Sierau, B., Sirevaag,
956 A., Sjogren, S., Stetzer, O., Swietlicki, E., Szczodrak, M., Vaattovaara, P., Wahlberg, N.,
957 Westberg, M., and Wheeler, C.R.: The Arctic summer cloud ocean study (ASCOS): Overview
958 and experimental design, *Atmos. Chem. Phys.*, 14, 2823–2869, doi:10.5194/acp-14-2823-2014,
959 2014.
960
961 Tunved, P., Ström, J., Krejci R.: Arctic aerosol life cycle: linking aerosol size distributions
962 observed between 2000 and 2010 with air mass transport and precipitation at Zeppelin station,
963 Ny-Alesund, Svalbard, *Atmos. Chem. Phys.*, 13, 3643-3660, doi:10.5194/acp-13-3643-2013,
964 2013.
965
966 United Nations Environment Programme, Near-term Climate Protection and Clean Air Benefits:
967 Actions for Controlling Short-Lived Climate Forcers, UNEP, Nairobi, Kenya,
968 <http://www.unep.org/publications/ebooks/SLCF/>, 2011.
969
970 Wylie, D., and J.G. Hudson: Effects of long range transport and clouds on cloud
971 condensation nuclei in the Springtime Arctic, *J. Geophys. Res.*, 107, 4318,
972 doi:10.1029/2001JD000759, 2002.
973

974 Yum, S.S., and Hudson, J.G.: Vertical distributions of cloud condensation nuclei spectra over the
975 springtime Arctic Ocean, *J. Geophys. Res.*, 106, 15045–15052, doi:10.1029/2000JD900357,
976 2001.
977
978 Zamora, L.M., Kahn, R.A., Cubison, M.J., Diskin, G.S., Jimenez, J.L., Kondo, Y., McFarquhar,
979 G.M., Nenes, A., Thornhill, K.L., Wisthaler, A., Zelenyuk, A., and Ziemba L.D.: Aircraft-
980 measured indirect cloud effects from biomass burning smoke in the Arctic and subarctic, *Atmos.*
981 *Chem. Phys. Discuss.*, 15, 22823–22887, doi:10.5194/acpd-15-22823-2015, 2015, 2015.
982
983 Zhao, C. and Garrett, T.J.: Effects of Arctic haze on surface cloud radiative forcing. *Geophys.*
984 *Res. Lett.* 10.1002/2014GL062015, 2015.
985
986

987 **Table 1.** Summary of averaged cloud observations with LWC>0.01 g m⁻³ for study periods 1 and
 988 2. Values without parentheses are referenced to ambient volumes and values in parentheses are
 989 referenced to STP. 5;95 are the 5th and 95th percentiles.

<u>Measurement</u>	Period 1 (July 5-11): 35 samples; 1.2 hours in cloud			Period 2 (July 11-21): 27 samples; 0.4 hours in cloud		
	<u>Mean</u>	<u>Median</u>	<u>5;95</u>	<u>Mean</u>	<u>Median</u>	<u>5;95</u>
Altitude (m-msl)	920	178	88;2272	1011	835	97;2608
Temperature (°C)	-1.9	-0.4	-6.5;2.2	+1.2	+2.2	-4.9;3.5
CDNC (STP) (cm ⁻³)	75 (85)	93 (91)	1.1;154 (1.1;185)	73 (83)	52 (55)	13;228 (14;265)
LWC (STP) (g m ⁻³)	0.12 (0.13)	0.10 (0.12)	0.014;0.32 (0.013;0.32)	0.12 (0.13)	0.12 (0.13)	0.025;0.26 (0.024;0.31)
VMD (µm)	17.2	18.7	9.9;30.0	15.0	14.5	9.1;21.4
CCNC(0.6%) (cm ⁻³): (17 P-1; 27 P-2)	90	120	2;168	81	43	18;227
N50 (cm ⁻³)	113	134	4.8;319	126	68	29;334
N100 (cm ⁻³)	35	47	1.3;73	81	31	13.8;274
CDNC(STP)/CCNC(0.6%)	0.75	0.56	0.18;1.50	1.18	1.22	0.47;1.87
CDNC(STP)/N50	0.82	0.90	0.16;1.40	0.73	0.68	0.28;1.08
CDNC(STP)/N100	2.78	2.63	0.28;7.94	1.37	1.25	0.58;2.15
CCNC(0.6%)/N50	0.64	0.63	0.50;0.84	0.64	0.64	0.52;0.87
CCNC(0.6%)/N100	1.92	1.79	0.67;3.11	1.27	1.0	0.75;2.28
CO (ppbv)	79	80	77;81	90	87	81;108
LWP (g m ⁻²); (13 P-1; 23 P-2)	30	27	1.5;4	22	13	1.0;70.5

Table 2. Summary of averaged observations for low-altitude (LA) and higher-altitude (HA) clouds. Values without parentheses are referenced to ambient volumes and values in parentheses are referenced to STP. 5, 95 are the 5th and 95th percentiles.

<u>Measurement</u>	LA (<200m): 24 samples; 0.89 hours in cloud			HA (>200m): 38 samples; 0.72 hours in cloud		
	<u>Mean</u>	<u>Median</u>	<u>5:95</u>	<u>Mean</u>	<u>Median</u>	<u>5:95</u>
Altitude (m-msl)	129	127	79;178	1485	1481	457;2391
Temperature (°C)	+0.6	+0.2	-2.5;2.9	-1.2	+0.9	-6.5;2.7
CDNC (STP) (cm ⁻³)	31 (30)	11 (10)	1;106 (1;102)	101 (118)	91 (101)	28;211 (31;245)
LWC (STP) (g m ⁻³)	0.10 (0.10)	0.05 (0.05)	0.01;0.34 (0.01;0.33)	0.13 (0.15)	0.13 (0.15)	0.04;0.25 (0.04;0.30)
VMD (µm)	20.7	20.1	14.6;31	13.4	12.5	9.1;19.4
CCNC(0.6%) (cm ⁻³); (16 LA; 28 HA)	74	24	2;184	90	58	21;217
N50 (cm ⁻³)	91	11	4.2;319	136	133	41;334
N100 (cm ⁻³)	26	4	1.3;73	73	47	20;232
CDNC(STP)/CCNC(0.6%)	0.61	0.57	0.18;1.3	1.2	1.2	0.6;1.9
CDNC(STP)/N50	0.61	0.44	0.14;1.5	0.91	0.93	0.5;1.3
CDNC(STP)/N100	2.3	1.4	0.35;9.0	2.1	1.9	0.7;3.7
CCNC(0.6%)/N50	0.66	0.71	0.52;0.7	0.68	0.64	0.5;0.9
CCNC(0.6%)/N100	1.8	1.6	0.96;2.6	1.5	1.1	0.8;3.4
CO (ppbv)	81	80	78;82	86	83	77;107

Figure. Captions

Figure 1. Compilation of the flight tracks. All flights originated from Resolute Bay (74°40'48"N 94°52'12"W).

Figure 2. Satellite images from July 5 when LA clouds were sampled over the two polynyas to the north and from July 8 when LA clouds were sampled along Lancaster Sound (July 8). Lancaster Sound is cloud free on July 5 and mostly covered by cloud on July 8. Resolute Bay is marked with a "X". Images are courtesy of NASA Worldview: <https://earthdata.nasa.gov/labs/worldview/>.

Figure 3. Four examples of profiles through HA clouds. a) Case from July 7 showing CO, CDNC, CCNC(0.6%) and particle number concentrations, where Nx-100, N100 and N5 are for particles sized between "x" nm and 100 nm, >100 nm and >5 nm respectively. b) Case from July 7 showing LWC, VMD, θ_e and temperature, where VMD, θ_e and temperature have been scaled as indicated in the legend. c) As in a), but case from July 17 and without N5. d) As in b), but case from July 17. e) As in a), but case from July 19. f) As in b) but case from July 19. g) As in a) but case from July 20 and without N5. h) as in b), but case from July 20. The CDNC are all referenced to STP, and θ_e is given in degrees Centigrade before scaling.

Figure 4. Time series during the sampling of low (LA) cloud or fog over the polynyas north of Resolute Bay. a) July 5 time series showing CO, CDNC, CCNC(0.6%) and particle number concentrations, where N30-100 is for particles sized between 30 nm and 100 nm and N100 is for particles sized >100 nm. b) July 7 time series showing CO, CDNC and particle number concentrations, where N20-100, N50-100 and N100 are for particles sized between 20 nm and 100 nm, between 50 nm and 100 nm and >100 nm respectively. CCNC(0.6%) measurements are unavailable for this period on July 7. Wind direction and relative position of polynyas are indicated in both panels. CDNC are referenced to STP.

Figure 5. Time series of altitude, CO, N80-100, N90-100, N100, CCNC(0.6%) and CDNC from low cloud (LA) cloud sampling over Lancaster Sound on July 8. The cloud was deeper over the open water of the Sound (see satellite picture in Fig. 2b). Over the ice to the west, the cloud was not as deep and could not be sampled. Segments over water and ice are indicated at the top of the figure.

Figure 6. Profiles down into cloud showing a) θ_e , b) LWC and c) VMDData for periods 17:27-17:29 UT and 17:38-17:39 UT during July 8. d) shows CDNC, N100, CO and CCNC(0.6%) for the 17:27-17:29 UT profile, and e) shows CDNC, N100, CO and CCNC(0.6%) for the 17:38-17:39 UT profile.

Figure 7. The LWC plotted as a function of the CDNC (a) and VMD (b) for the LA (orange) and HA (blue) samples. Linear regressions for each of the LA and HA samples are also plotted, and the coefficients of determination are given in the legends.

Figure 8. As in Fig. 7a, but only for four identified LA cases (July 5, 7, 8 and 11). Linear regressions for each set of samples are also plotted, and the coefficients of determination are given in the legends.

Figure 9. Plots of CDNC versus a) N100 and b) N50. Points are identified between LA (yellow) and HA (blue) samples, and the 1:1 lines are for reference.

Figure 10. a) CDNC plotted versus the CCNC(0.6%) measured at 0.6% supersaturation; points are identified between LA (yellow) and HA (blue) samples, and linear regressions through the origin are shown. b) CCNC(0.6%) plotted versus N50 and N100; power law fits to each are provided for reference.

Figure 11. The mean VMD of all cloud samples plotted versus the CDNC. All CDNC are referenced to the in-situ pressure. The dashed vertical green line represents the “CCN-limited” division discussed by Mauritsen et al (2011) and estimated here as 16 cm^{-3} . The solid black line is another reference showing the relationship between VMD and CDNC for a constant LWC: the study mean LWC of 0.12 g m^{-3} (Table 1). Samples with higher CO ($>90 \text{ ppbv}$) are identified by the open red circles. Also highlighted for the discussion are LA samples from July 5 (red dots) and July 7 (orange dots). The median CDNC are 1.3 cm^{-3} and 7.8 cm^{-3} , for July 5 and 7, respectively; the N50 are 6 cm^{-3} and 8.3 cm^{-3} for July 5 and 7, respectively; the N100 are 3 cm^{-3} and 2.2 cm^{-3} for July 5 and July 7, respectively.




Figure 1. Compilation of the flight tracks. All flights originated from Resolute Bay (74°40'48"N 94°52'12"W).




Figure 2. Satellite images from July 5 when LA clouds were sampled over the two polynyas to the north and from July 8 when LA clouds were sampled along Lancaster Sound (July 8). Lancaster Sound is cloud free on July 5 and mostly covered by cloud on July 8. Resolute Bay is marked with a “X”. Images are courtesy of NASA Worldview: <https://earthdata.nasa.gov/labs/worldview/>.




Figure 3. Four examples of profiles through HA clouds. a) Case from July 7 showing CO, CDNC, CCNC(0.6%) and particle number concentrations, where Nx-100, N100 and N5 are for particles sized between “x” nm and 100 nm, >100 nm and >5 nm respectively. b) Case from July 7 showing LWC, VMD, θ_e and temperature, where VMD, θ_e and temperature have been scaled as indicated in the legend. c) As in a), but case from July 17 and without N5. d) As in b), but case from July 17. e) As in a), but case from July 19. f) As in b) but case from July 19. g) As in a) but case from July 20 and without N5. H) as in b), but case from July 20. The CDNC are all referenced to STP, and θ_e is given in degrees Centigrade before scaling.




Figure 4. Time series during the sampling of low (LA) cloud or fog over the polynyas north of Resolute Bay. a) July 5 time series showing CO, CDNC, CCNC(0.6%) and particle number concentrations, where N30-100 is for particles sized between 30 nm and 100 nm and N100 is for particles sized >100 nm. b) July 7 time series showing CO, CDNC and particle number concentrations, where N20-100, N50-100 and N100 are for particles sized between 20 nm and 100 nm, between 50 nm and 100 nm and >100 nm respectively. CCNC(0.6%) measurements are unavailable for this period on July 7. Wind direction and relative position of polynyas are indicated in both panels. CDNC are referenced to STP.




Figure 5. Time series of altitude, CO, N80-100, N90-100, N100, CCNC(0.6%) and CDNC from low cloud (LA) cloud sampling over Lancaster Sound on July 8. The cloud was deeper over the open water of the Sound (see satellite picture in Fig. 2b). Over the ice to the west, the cloud was not as deep and could not be sampled. Segments over water and ice are indicated at the top of the figure.




Figure 6. Profiles down into cloud showing a) θ_e , b) LWC and c) VMDData for periods 17:27-17:29 UT and 17:38-17:39 UT during July 8. d) shows CDNC, N100, CO and CCNC(0.6%) for the 17:27-17:29 UT profile, and e) shows CDNC, N100, CO and CCNC(0.6%) for the 17:38-17:39 UT profile.




Figure 7. The LWC plotted as a function of the CDNC (a) and VMD (b) for the LA (orange) and HA (blue) samples. Linear regressions for each of the LA and HA samples are also plotted, and the coefficients of determination are given in the legends.




Figure 8. As in Fig. 7a, but identifying the specific LA cases of July 5, 7, 8 and 17. Linear regressions for each set of samples are also plotted, and the coefficients of determination are given in the legends. The slopes are significant at a 95% confidence level within $\pm 30\%$ for July 7 and within 60% for July 8. The slopes in the July 5 and 17 cases are not significant at a 95% confidence level.





Figure 9. Plots of CDNC versus a) N100 and b) N50. Points are identified between LA (yellow) and HA (black asterisk) samples, and the 1:1 lines are for reference.





Figure 10. a) CDNC plotted versus the CCNC measured at 0.6% supersaturation; points are identified between LA (yellow) and HA (blue) samples, and linear regressions through the origin are shown; the CCNC(0.6%) points are limited to 44 of the 62 total, due to problems with the CCN measurement; the 44 are split 16 and 28 between LA and HA,. b) CCNC(0.6%) (44 points) plotted versus N50 and N100; power law fits to each are provided for reference.




Figure 11. The mean VMD of all cloud samples plotted versus the CDNC. All CDNC are referenced to the ambient pressure. The dashed vertical green line represents the “CCN-limited” division discussed by Mauritsen et al (2011) and estimated here as 16 cm^{-3} . The solid black line is another reference showing the relationship between VMD and CDNC for a constant LWC: the study mean LWC of 0.12 g m^{-3} (Table 1). Samples with higher CO ($>90 \text{ ppbv}$) are identified by the open red circles. Also highlighted for the discussion are LA samples from July 5 (blue dots) and July 7 (orange dots).