

Author Response to Reviewer #1

We thank reviewer #1 for their constructive suggestions. We reproduce the comments here in italics.

5 *Page 3, Line 27. These simulations have a very coarse spatial resolution (4x50). The authors recognise this weakness*

Due to computational limits, we use a coarse spatial resolution. This means we are calculating radiative effects using aerosol concentrations and meteorological variables averaged over quite large (about 400 km) spatial scales. As aerosol and meteorological factors vary on much smaller spatial scales, the coarse spatial resolution likely introduces some errors. This 10 may be more important for the AIE, as cloud reflectance has a nonlinear relationship with CDNC. Despite this, we do not expect these errors to qualitatively change our results and conclusions.

15 *The model evaluation using AERONET AOD demonstrates that including the waste combustion emission source does not degrade the model. The authors could consider using measured BC or OC mass concentrations from regions heavily impacted by waste combustion as an additional evaluation of the model.*

This is an interesting suggestion. We did have interest in including measurements of BC/OC mass concentrations; however, robust measurements of mass concentration (especially speciated mass concentrations) are rare in developing countries (which is where most waste combustion emissions are located). One possibility is the SPARTAN network (<http://spartan-network.org/>), which measures PM_{2.5} concentrations in developing countries. Unfortunately, there are not enough 20 measurements available to achieve robust statistics. We would be interested in hearing about other possible datasets.

Page 6, line 31: Suggest reword to remove the word “trend”.

25 We agree. We are replacing the word “trend” with “variability”.

Author Response to Reviewer #2

We thank reviewer 2 for their constructive response. We reproduce reviewer comments here in italics.

30 *General comment on results: As the authors point out, many assumptions about the domestic waste emissions have large uncertainties. Since this is one of the first studies that investigates climate impacts of waste-combustion emissions, it would be useful to formulate explicitly in the conclusions, which aspects of the emission information would be most useful to improve in future work*

35 Thank you for this suggestion and we feel this will be a useful addition to the discussion. We have added the following lines to the conclusions:

40 “There is little information on emission size distribution and optical properties from waste combustion. Better knowledge of these parameters along with continued validation of emission mass fluxes will reduce model uncertainty.”

General comment on presentation: Some phrases are rather colloquial, and I suggest going through the paper and rephrase them.

45 Thank you for suggesting these corrections. We have made a number of edits following the reviewer’s examples.

p. 2, line 7: “. . . how the particles are mixed together”: do you mean mixed together within the population or mixed together within one particle?

We mean mixed together within a single particle. We have rephrased the manuscript to read:

“the manner of mixing of different species within a single particle”

5 *p.2, second paragraph: Scattering and absorption in general also depends strongly on the morphology of the particles. Assuming different variants of spherical particles may not reflect reality very well.*

This is a good point. We do assume spherical particles (a common assumption in global modeling studies). This will introduce some errors. We have added the following lines:

10 “In this study, we assume spherical particles, which is not perfectly realistic for fresh combustion aerosol, and these details of particle shape may alter optical properties.”

p. 2, line 25: define kappa.

15 We have added the following lines:
“Particle hygroscopicity can be described with the hygroscopicity parameter, kappa, which is the ratio of the number of moles of solute per dry volume to the moles of water per volume of pure water (Petters and Kreidenweis, 2007).”

20 *p. 3-4: What about sea salt and dust emissions?*

Sea salt emissions follow the scheme of Jeagle et al. (2011), and dust emissions follow the DEAD scheme of Zender et al. (2003). We have added this statement and references to the methods section.

25 *p. 3-4: Does the chemistry model include ammonium nitrate? If not, can you say something how this might impact the conclusions?*

30 Ammonium nitrate is not included in the TOMAS microphysics; however, it is included in the “bulk” GEOS-Chem aerosol setup, which runs concurrently with TOMAS. We are interested in incorporating ammonium nitrate into TOMAS in the future.

p. 5: Direct radiative effect calculation: are the six different optical assumptions applied to all BC containing particles, regardless of their source (i.e. not only to the waste combustion particles)?

35 This is a good question, and we have added a clarifying sentence to the manuscript. The optical assumptions are applied to all BC containing particles. We do not track particles by emission source after they have been emitted into the model (sources are emitted separately, but once emitted they enter the same tracer). Given the coarse spatial and temporal resolution of our model, it is fair to assume some degree of mixing at the model length scale. In general, BC from different sources (fossil fuel vs. biomass burning) may mix at different timescales due to co-emitted species. Our optical assumptions 40 are therefore idealized cases.

p. 9, line 29/30: the minus sign was separated from the number due to a line break. Make sure that this doesn't happen.

In the final typeset version, we will look for this error.

45 *p. 8, line 18: typo: dominant*

Thank you.

Figure 2: To draw the conclusion that the BASE run and the WASTE_OFF run are different, you need to show the differences in slope and r2 are statistically significant. If it turns out that they are indeed different, please comment if the effect size is meaningful (i.e. of practical importance).

5 The differences are not statistically significant. We included this figure first to demonstrate that including this emission source does not degrade the model comparison, and second to show TOMAS has skill at reproducing observed aerosol optical properties (this is the first GEOS-Chem-TOMAS paper that uses GEOS-Chem version 10). We have added the line:

10 “While these changes are not statistically significant, we note that GEOS-Chem-TOMAS has some skill at reproducing
observed aerosol optical properties, and including this inventory does not degrade model comparison”

The aerosol radiative effects of uncontrolled combustion of domestic waste

5 John K. Kodros¹, Rachel Cucinotta², David A. Ridley³, Christine Wiedinmyer⁴, Jeffrey R. Pierce¹

¹Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado, USA

²University of North Carolina at Charlotte, Charlotte, North Carolina, USA

³Department of Civil & Environmental Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

10 ⁴National Center for Atmospheric Research, Boulder, Colorado, USA

Correspondence to: J. K. Kodros (jkodros@atmos.colostate.edu)

Abstract. Open, uncontrolled combustion of domestic waste is a potentially significant source of aerosol; however, this aerosol source is not generally included in many global emissions inventories. To provide a first estimate of the aerosol radiative impacts from domestic-waste combustion, we incorporate the Wiedinmyer et al. (2014) emissions inventory into GEOS-Chem-TOMAS, a global chemical-transport model with online aerosol microphysics. We find domestic-waste combustion increases global-mean black carbon and organic aerosol concentrations by 8% and 6%, respectively, and by greater than 40% in some regions. Due to uncertainties regarding aerosol optical properties, we estimate the globally averaged aerosol direct radiative effect to range from -5 mW m^{-2} to -20 mW m^{-2} ; however, this range increases to -40 mW m^{-2} to $+4 \text{ mW m}^{-2}$ when we consider uncertainties in emission mass and size distribution. In some regions with significant waste combustion, such as India and China, the aerosol direct radiative effect may exceed -0.4 W m^{-2} . Similarly, we estimate a cloud-albedo aerosol indirect effect of -13 mW m^{-2} , with a range of -4 mW m^{-2} to -49 mW m^{-2} due to emission uncertainties. In the regions with significant waste combustion, the cloud-albedo aerosol indirect effect may exceed -0.4 W m^{-2} .

25 1 Introduction

Open, uncontrolled combustion of domestic waste occurs on a global scale, potentially emitting significant amounts of black carbon (BC) and organic aerosol (OA); however, this source is not commonly considered in modeling studies estimating aerosol radiative impacts (Christian et al., 2010; Mohr et al., 2008; Wiedinmyer et al., 2014). Uncontrolled combustion of domestic waste refers to the intentional disposal of trash in an open fire either at home or at a dump (Wiedinmyer et al., 30 2014). This practice is common in developing countries in both rural and urban areas where municipal waste collection is insufficient; however, open combustion of domestic waste also occurs in rural developed countries (Christian et al., 2010; Wiedinmyer et al., 2014). The proximity of waste-combustion emissions to residences has created concerns for both air

quality and climate. In a separate study, we estimate the global burden of mortalities due to chronic exposure of waste-combustion PM_{2.5} (Kodros et al., 2016). Here we provide a first estimate of the aerosol radiative impacts of domestic-waste combustion.

Carbonaceous aerosol from combustion sources can impact climate in several ways; the best understood of these 5 being the direct radiative effect (DRE) and cloud-albedo aerosol indirect effect (AIE) (Boucher et al., 2013). The DRE refers to scattering and absorption of incoming solar radiation by aerosols (Charlson et al., 1992). The magnitude and relative amount of scattering and absorption is dependent on particle composition and size, as well as the manner of mixing of different species within a single particle (Bond et al., 2006; Bond et al., 2013). BC has a strong absorbing component, while 10 OA is either considered as completely scattering or having some absorbing component in the shorter visible and UV wavelengths (i.e. brown carbon) (Andreae and Gelencsér, 2006; Hammer et al., 2015; Wang et al., 2014), although these optical properties of organic aerosol are an open area of research. Additionally, scattering and absorption efficiencies of carbonaceous particles depend on the ratio of particle diameter to wavelength of radiation. For solar radiation, atmospheric aerosol typically have peak efficiencies between 100 nm to 1 μm , though this will vary with composition (Seinfeld and Pandis, 2012). Finally, absorption tends to increase in an internally mixed population (all particles in a size range contain an 15 identical mixture of all chemical species) compared to an externally mixed population (different chemical species exist in separate particles) (Chung and Seinfeld, 2005; Jacobson, 2001). In this study, we assume spherical particles, which is not realistic for fresh combustion aerosol. Details of particle shape may alter optical properties.

The cloud-albedo AIE refers to altering the reflectivity of clouds by changing the cloud droplet number concentration (CDNC) (Twomey, 1974). The ability for OA and BC cloud condensation nuclei (CCN) to activate into cloud 20 droplets is a strong function of particle size. In typical atmospheric conditions, OA and BC (when mixed with hydrophilic species) typically act as CCN when they have diameters larger than 40-100 nm (Petters and Kreidenweis, 2007). The number of particles in this size range is dependent on primary emissions as well as nucleation, condensation, and coagulation (Pierce and Adams, 2009). As these microphysical processes are often inter-connected, the AIE response to an emission source may 25 be non-linear (e.g. additional primary emissions may reduce nucleation and condensation growth while increasing coagulation) (Bauer et al., 2010; Butt et al., 2016; D'Andrea et al., 2015). The AIE is also dependent on particle hygroscopicity (Petters and Kreidenweis, 2007). Particle hygroscopicity can be described with the hygroscopicity parameter, kappa, which is the ratio of the number of moles of solute per dry volume to the moles of water per volume of pure water (Petters and Kreidenweis, 2007). While combustion aerosol tend to have lower hygroscopicity (kappa for OA=~0.1 and BC= 0.0), mixing with inorganic species, such as sulfate and sea salt, and chemical aging of the organics to more-hygroscopic 30 species allows for activation at smaller diameters.

Waste combustion is an often-overlooked emission source, especially in developing countries (Christian et al., 2010). Wiedinmyer et al. (2014) provided a new global inventory for the uncontrolled combustion of domestic waste, estimating global BC and OC emissions of 0.6 and 5.1 Tg yr⁻¹, respectively. This estimate is an order of magnitude greater than earlier estimates of emissions from uncontrolled waste combustion in Bond et al. (2004) (which focused on combustion-

Jack 5/12/2016 5:00 PM

Deleted: how the

Jack 5/12/2016 5:00 PM

Deleted: particles are mixed together

aerosol emissions in general, not just from waste combustion). Amount of annual waste burned was the largest difference between the inventories, where Bond et al. (2004) estimated a total of 33 Tg and Wiedinmyer et al. (2014) predicted 970 Tg. Further, Wiedinmyer et al. (2014) considered waste combustion in both rural and urban areas in developing and rural areas in developed countries. Many global inventories, including Emissions Database for Global Atmospheric Research (EDGAR, 5 Janssens-Maenhout et al. 2010), do not include estimates of uncontrolled waste-combustion emissions. In addition to carbonaceous emissions, Wiedinmyer et al. (2014) estimated 0.5 Tg yr⁻¹ of sulfur-dioxide (SO₂) and 1.1 Tg yr⁻¹ of ammonia (NH₃). These gas-phase species may increase nucleation and condensation rates, thereby increasing particle number and mass concentration through secondary aerosol formation.

There are several uncertainties related to emissions in the Wiedinmyer et al. (2014) inventory. First, the reported 10 uncertainty in the Akagi et al. (2011) emission factors for BC and OC are 42% and 93%. This is in addition to uncertainties in the mass of waste generated (i.e. estimates of fuel use). Second, the Wiedinmyer et al. (2014) inventory does not include information on emission size distribution. The emitted particle size distribution varies with fuel type and flaming condition, and goes through rapid aging in the first few hours after emission on spatial scales not resolved in global models (Janhäll et al., 15 2010; Sakamoto et al., 2016). Finally, this inventory does not include information on the optical properties of organic aerosol. Absorptive OA (or brown carbon, BrC) has been observed in some wood and agricultural waste fuels, under certain flaming conditions (Chen and Bond, 2010; McMeeking et al., 2014; Saleh et al., 2013); however, we are not aware of studies determining these aerosol properties for domestic-waste combustion. In this manuscript, we explore uncertainties in total emission mass and size distribution (affecting both DRE and AIE) as well as optical properties (affecting just DRE).

To our knowledge, this is the first paper to quantify the global aerosol radiative impacts from uncontrolled 20 combustion of domestic waste. In Section 2, we discuss the model set-up (2.1), methods for model evaluation (2.2), and calculation of radiative effects (2.3-2.4). In Section 3, we show comparisons to ground observations of model simulations with and without waste-combustion emissions. In Section 4, we discuss changes in modeled aerosol mass (4.1), size-resolved aerosol number concentration (4.2), direct radiative effect (4.3), and cloud-albedo aerosol indirect effect (4.4). In Section 5, we share our conclusions.

25 2 Methods

2.1 Model overview

To calculate aerosol number, mass, and size distributions from domestic-waste combustion, we use the Goddard Earth Observing System global chemical-transport model (GEOS-Chem) version 10.01 coupled with TwO Moment Aerosol Sectional (TOMAS) microphysics scheme (Adams and Seinfeld, 2002). We use GEOS-Chem with 47 vertical layers and a 30 4°x5° horizontal resolution. This version of TOMAS uses 15 size sections ranging from 3 nm to 10 μm with tracers for sulfate, sea salt, OA, BC, and dust (Lee and Adams, 2012; Y. Lee et al., 2013). GEOS-Chem is driven using reanalysis meteorology fields from GEOS5 (<http://gmao.gsfc.nasa.gov>). Since meteorology is prescribed, the addition of waste-

Jack 5/14/2016 1:02 PM

Deleted: look at

combustion emissions does not feedback into weather patterns. All simulations are for the year 2010, with 1 month of model spin-up that is not included in analysis.

We use a ternary nucleation scheme involving water, sulfuric acid, and ammonia following the parameterization of Napari et al. (2002), scaled with a global tuning factor of 10^{-5} (Jung et al., 2010; Westervelt et al., 2013). In addition, we use a binary nucleation scheme (sulfuric acid and water) when ammonia mixing ratios are less than 1 pptv (Vehkamaki et al., 2002). Secondary organic aerosol (SOA) sources include a biogenic monoterpene component of 19 Tg yr⁻¹ as well as an anthropogenically enhanced component of 100 Tg yr⁻¹ (we do not know if the anthropogenically enhanced SOA is due to anthropogenic VOCs or an enhancement of biogenic SOA due to anthropogenic influences on chemistry) (D' Andrea et al., 2013; Spracklen et al., 2011). SOA is condensed irreversibly onto existing aerosol at the time of emission of the parent compounds using 10% of monoterpene emissions for the biogenic component and 0.2 Tg-SOA per Tg-CO for the anthropogenic component (CO emissions are used as a proxy for the spatial and temporal distribution of anthropogenically enhanced SOA).

In all simulations we use the following emission inputs. We use fossil-fuel combustion anthropogenic emissions from the Emissions Database for Global Atmospheric Research (EDGAR; Janssens-Maenhout et al. 2010) version 4. EDGAR emissions are overwritten in the United States by the Environmental Protection Agency 2011 National Emissions Inventory (NEI2011; <http://www3.epa.gov/ttn/chief/>), in Canada by Criteria Air Contaminants (CAC; <http://www.ec.gc.ca/inrp-npri/>), in Mexico by Big Bend Regional Aerosol and Visibility Study (BRAVO; Kuhns et al. 2005), in Europe by the Cooperative Programme for Monitoring and Evaluation of the Long-Range Transmission of Air Pollutants in Europe (EMEP; Vestreng et al. 2009), and in Asia by the Streets inventory (Streets et al. 2003). Black and organic carbon (OC) emissions from biofuel and fossil-fuel combustion are from Bond et al. (2007) and biomass burning from Global Fire Emissions Database version 3 (GFED3; van der Werf et al. 2010) with three-hourly scale factors. [Sea salt aerosol emissions follow the scheme of Jaeglé et al. \(2011\)](#), while dust emissions are based on the DEAD scheme (Zender et al., 2003).

To estimate the impact of waste-combustion emissions, we include the Wiedinmyer et al. (2014) inventory to the above emissions set-up. Table 1 describes all simulations used in this study. Our BASE simulation uses total BC, OC, SO₂, and NH₃ mass as in Wiedinmyer et al. (2014). We emit 1% of SO₂ as aerosol-phase sulfate (SO₄) and use a fixed OA to OC ratio of 1.8. We scale bulk BC and OA mass into size sections assuming a single lognormal mode with a geometric number-mean diameter (GMD) of 100 nm and a standard deviation of 2. We note that Wiedinmyer et al. (2014) does not include information on the emission size distribution, and this is a significant source of uncertainty (L. Lee et al., 2013). The spatial distribution of emissions is shown in Figure 1 for OA, BC, and NH₃. The emissions of SO₂ show a similar spatial pattern, but with a factor of 2 lower in magnitude than NH₃. The regions with the largest particle emissions are eastern China, northern India, eastern Europe, and Mexico. To test the sensitivity of aerosol radiative effects to waste-combustion emission size distribution, we include two sensitivity simulations, SIZE200 and SIZE30. Here, we conserve total emission mass, but increase the GMD from 100 nm to 200 nm and decrease the GMD from 100 nm to 30 nm, respectively. These distributions

are chosen to reflect the range of reported values from fresh and aged fossil fuel and biomass burning (Ban-Weiss et al., 2010; Carrico et al., 2010; Janhäll et al., 2004; Sakamoto et al., 2015). In Kodros et al. (2015), we also tested the sensitivity of aerosol radiative effects to uncertainty in the standard deviation of the emission size distribution and find the results are qualitatively similar to uncertainties in the GMD (a narrowing of the emission size distribution has a similar effect as to 5 | decreasing the value of the GMD, and visa versa), and so we do not include that here. In PRIMARY, we emit only aerosol-phase BC and OA waste combustion, and remove emissions of gas-phase SO₂ and NH₃, from this source. This simulation will test the contribution of nucleation and condensation from gas-phase species emitted from waste combustion. Finally, we test the sensitivity of aerosol effects to a factor of 2 uncertainty in total waste-combustion emission mass (HIGHMASS and 10 | LOWMASS). The factor of 2 uncertainty is similar to uncertainties listed in emission factors in Akagi et al. (2011). We compare these simulations to a control simulation with no waste-combustion emissions (WASTE_OFF).

Jack 5/14/2016 1:03 PM
Deleted: making the GMD smaller

2.2 Aerosol optical depth comparison

We compare model aerosol optical depth (AOD) with measurements made with the Aerosol Robotic Network (AERONET). AERONET is a network of ground-based remote sensing instruments aimed at providing global, long-term measurements of aerosol properties. AERONET uses a sun photometer to measure AOD at several wavelengths in 15 minute intervals when 15 | the sun is not obscured by clouds (Holben et al., 1998).

To compare to AERONET measurements, we output GEOS-Chem-TOMAS size-resolved mass and number concentrations at three-hour intervals for the BASE and WASTE_OFF simulations. We calculate model AOD using refractive indices from the Global Aerosol Database (Koepke et al., 1997) and published Mie code (Bohren and Huffman, 1983). In this calculation, we only consider an external mixing-state assumption, though we note the extinction optical depth 20 | (what we are referring to as AOD) of an internal mixture is very similar (though the individual scattering and absorption optical depths are different). We average AERONET AOD into three-hour intervals and match measured time and locations with model AOD. We restrict observed AOD to contain greater than six measurements within each three-hour time period, thus removing intervals where measurements were limited due to clouds or instrument error. We also restrict our comparison to only include time periods where the observed AOD is greater than 0.01 to avoid detection-limit issues, as this is the 25 | uncertainty limit of AERONET measurements (Holben et al., 1998). We then compare the time average of the co-sampled sites that match these criteria.

Jack 5/16/2016 2:18 PM
Deleted: 3

2.3 Direct radiative effect calculation

We calculate the all-sky direct radiative effect (DRE) using GEOS-Chem-TOMAS monthly averaged aerosol mass and number concentrations and refractive indices from GADS. We calculate aerosol optical depth, single scatter albedo, and 30 | asymmetry parameter using Mie code published in Bohren and Huffman (1983). We use these aerosol optical properties and monthly mean albedo and cloud fraction from GEOS5 as inputs to the offline version of the Rapid Radiative Transfer Model

Jack 5/14/2016 1:04 PM
Deleted: 6

for GCMs (RRTMG; Iacono et al., 2008), implemented in GEOS-Chem (Heald et al., 2014), that considers longwave and shortwave and treats clouds with the Monte Carlo independent column approximation (McICA; Pincus et al., 2003).

We estimate the DRE for six different optical assumptions: a core-shell morphology with and without absorptive OA, an external mixture with a constant absorption enhancement factor of 1.5 (ext*1.5) with and without absorptive OA, a 5 homogeneous internal mixture, and an external mixture. These optical assumptions are discussed in detail in Kodros et al. (2015) (see Table 1 in Kodros et al. (2015) and Table 3 in this paper) [and are applied to all particles, regardless of emission source](#). In all cases, we assume the particles are spherical. Briefly, the homogeneous internal mixture assumes all particles in a size bin have the same composition, and the refractive index of the particle is a volume-weighted average of the individual species. The core-shell mixture assumes black carbon forms a spherical core and the other hydrophilic species form a 10 homogeneously mixed shell around the core. The external mixture assumes black carbon exists as a separate population from the other homogeneously mixed species. Finally, in ext*1.5 we simply multiply the absorption predicted in the external mixture by a constant enhancement factor of 1.5. This is done in some models to estimate the enhanced absorption observed in core-shell mixtures (such as Wang et al., 2014). In the BrC calculations, we estimate absorptive OA (or brown carbon) 15 using the modeled OA to BC ratio and the parameterization in Saleh et al. (2014). In all other calculations, we assume that OA is near-purely scattering throughout the visible spectrum.

We note several important limitations in our treatment of aerosol optics. First, mixing state may vary regionally and temporally, yet here we apply each mixing state globally and at all times. Second, it has been suggested that the optical properties of OA may change with photolysis, yet here our optical properties are held fixed (Lee et al., 2014; Zhao et al., 2015). Therefore, these mixing states should be considered as ideal cases. Finally, we use monthly mean concentrations and 20 properties as well as monthly mean cloud properties. Despite these limitations, we feel this method is sufficient for a first estimate of the DRE from waste-combustion emissions.

2.4 Cloud-albedo aerosol indirect effect calculation

We use monthly mean GEOS-Chem-TOMAS mass and number concentrations and the activation parameterization of Abdul-Razzak and Ghan (2002) to calculate cloud droplet number concentrations (CDNC). We assume a constant updraft 25 velocity of 0.5 m s^{-1} . For the AIE, we assume aerosol species are mixed internally within each size bin, and calculate kappa, the hygroscopicity parameter, as a volume-weighted average of the individual species (Petters and Kreidenweis, 2007). As a control, we assume an effective cloud drop radii of $10 \mu\text{m}$ and then perturb this value by taking the ratio of CDNC with and without waste-combustion emissions to the one-third power (as in Rap et al., 2013; Scott et al., 2014). To simulate the 30 change in the top-of-the-atmosphere radiative flux due to changes in effective cloud drop radii, we use RRTMG with monthly mean cloud fraction, temperature, pressure, and liquid water content from GEOS5. Again, we note that the use of monthly mean aerosol and meteorology fields is a limitation in this method, yet we feel it is sufficient for a first estimate.

3 Model Results

3.1 Comparison to observations

Figure 2 shows comparisons of time-averaged model AOD with AERONET measured AOD at all available sites for the BASE (a-c) and WASTE_OFF (d-f) simulations at 380 nm, 500 nm, and 1020 nm wavelengths (one dot for each site). After 5 we apply the criteria described in Section 2.2, we have 173 sites at 380 nm, 193 sites at 500 nm, and 211 sites at 1020 nm.

The model is generally capable of capturing the variability in measured AOD (in WASTE_OFF at 500 nm slope=0.94 and log mean bias=-0.18). Given the coarse spatial resolution of the model, the large variability is expected ($r^2=0.59$), as more- and less-polluted areas may exist within the same model gridbox (~400 km length scale). Including waste-combustion emissions slightly improves the model comparison in all three metrics and at all wavelengths (with the exception of the slope 10 at 1020 nm). While these changes are not statistically significant, we note that GEOS-Chem-TOMAS has some skill at reproducing observed aerosol optical properties, and including this inventory does not degrade model comparison

Jack 5/12/2016 3:13 PM

Deleted: trend

3.2 Changes in mass concentrations from domestic-waste combustion

Figure 3 shows the annual-mean percent change in boundary-layer OA, BC, NH₃, and PM_{2.5} in the BASE-WASTE_OFF simulations, and global-mean OA and BC changes are in Table 2. Combustion of domestic waste under the BASE emission 15 assumptions increases BC and OA concentrations in the global mean by 8.5% and 5.6%, respectively. Regionally, domestic-waste combustion increases surface BC concentrations by 10-20% throughout much of Asia, northern Africa, and South and Central America. The largest increases in BC (greater than 40%) occur in central Mexico (i.e Mexico City), Turkey, northern Egypt (i.e. Cairo), Sri Lanka, and Papua New Guinea, as well as increases of 30-40% in Pakistan, Morocco, and Central America. Despite greater OA emissions, the relative change of OA from domestic-waste combustion is generally less than 20 BC due to large OA sources from biofuel combustion and secondary organic aerosol in waste-combustion regions. Waste combustion does increase OA concentrations by more than 40% in Morocco, Turkey, Egypt and the eastern portions of the Middle East. Increases in BC and OA lead to increases in surface PM_{2.5} concentrations of 1-5% over most land masses outside of North America and Australia, and 5-10% in source regions.

Figure 3 also shows the percent change in gas-phase NH₃. Waste combustion contributes increases in NH₃ of 1-10% 25 in Mexico, Central and South America, and Africa. These are regions of significantly lower NH₃ concentrations compared to Asia and Europe in our simulations, which have negligible relative changes to NH₃. In our model, NH₃ may contribute to particle nucleation and growth, which may alter particle size distributions (but have little effect on PM_{2.5} mass). These changes are discussed in the next section. Relative changes in SO₄ mass generally follow the spatial pattern of OA; however, these changes are less than 1%, and so are not shown.

30 Table 2 shows the global, annual-mean percent change in boundary-layer BC and OA mass for BASE and the sensitivity simulations compared to WASTE_OFF (thus isolating the contribution of domestic-waste combustion to BC and OA). The largest changes to modeled BC and OA mass come when emission BC and OA mass is doubled (HIGHMASS)

and halved (LOWMASS). In these simulations, the model response to changes in emission mass is close to linear. In simulations, PRIMARY, SIZE30, and SIZE200 total BC and OA emission mass is held constant. The BC and OA mass distributions in SIZE30 and SIZE200 are shifted to smaller and larger sizes, respectively, which may lead to changes in deposition rates; however, these changes prove to be minor (Table 2). The global-mean relative change in BC and OA mass 5 from domestic-waste combustion ranges from 4.23-16.9% and 2.80-11.2% given the factor of 4 uncertainty in emission mass tested here.

3.3 Changes to size-resolved number concentrations

Modeled percent changes to annual-mean size-resolved number concentrations for the BASE-WASTE_OFF comparison are shown in Figure 4 and the global averages in Table 2. Here we consider the following size classes: all particles with a 10 diameter greater than 10 nm (N10), 40 nm (N40), 80 nm (N80), and 150 nm (N150). N40, N80, and N150 are proxies for climate-relevant particles. In the global mean for our BASE assumptions, waste combustion leads to a larger increase in N80 (1.04%) and N150 (1.16%) than in N10 (0.86%) and N40 (0.90%). This is due to particle emissions represented as a single lognormal mode centered around 100 nm as well as a microphysical feedback (discussed below).

Regionally, the sign and magnitude of changes to particle number concentration varies with size class. Waste 15 combustion tends to increase particle number concentration in each size class in Mexico, Central and South America, and the Pacific Islands. Conversely, waste combustion decreases N10 and N40 in much of Europe and Asia. This discrepancy is caused by a feedback in aerosol microphysics (discussed in detail in Westervelt et al. 2014, Pierce and Adams et al. 2009, Kodros et al. 2015). Briefly, emitted primary accumulation-mode particles (in BASE this is centered around 100 nm) increases particle surface area, thus increasing the sink for condensation vapors (e.g H₂SO₄, NH₃, organics). This increased 20 condensation sink reduces nucleation rates, reduces growth rates, and increases coagulation scavenging of nucleation mode particles. This effect is predominant in Europe and Asia and for smaller size cutoffs (N10 and N40) with a significant contribution of nucleation mode particles (such that waste combustion emissions cannot compensate for the reduced nucleation and growth). The regions of Central and South America, and Africa have lower concentrations of condensable 25 vapor than Europe and Asia in WASTE_OFF, and so this feedback is less dominant. In addition, the Americas and Africa have a larger fractional increase in NH₃ concentrations, which aid in nucleation and growth (Figure 3).

Figure 5a shows the change in global-mean particle number distribution for each of the sensitivity simulations relative to WASTE_OFF. In the BASE-WASTE_OFF comparison, particle number concentration decreases for sizes less than 80 nm (due to the feedback described above), and increases for sizes greater than 80 nm (due to primary emissions) (see 30 Table 2 for integrated values). The emission mass scale factor tends to scale the total waste combustion number distribution up or down, while maintaining the general shape of the distribution. The HIGHMASS simulation has a larger increase in accumulation-mode particles as well as a stronger suppression of nucleation/growth and thus a larger decrease in nucleation-mode particles relative to BASE. Meanwhile, LOWMASS scales in the opposite direction. In the PRIMARY simulation, we remove gas-phase NH₃ and SO₂ emissions from waste combustion, which contributes to particle nucleation and

Jack 5/14/2016 11:33 AM

Deleted: te

condensational growth. The removal of these condensable vapors results in a larger suppression of nucleation and growth relative to BASE, causing a larger decrease in sub-80nm particles as well as fewer particles in the 80-200 nm range. When only primary BC and OA from waste combustion are emitted, the global-mean N10 decreases (-0.17%, Table 2). Nucleation and condensational growth is a less significant contributor to particles greater than 200 nm, and so PRIMARY and BASE
5 have the same distribution at these larger sizes.

Altering the GMD of the emissions size distribution leads to the largest changes in modeled distributions. In SIZE200, we increase emission GMD while keeping total mass constant. Assuming a constant density, this necessitates a smaller number flux of emitted particles (as larger particles have more mass). The reduced particle number concentration relative to BASE allows for nucleation through emitted NH₃ and SO₂ to increase the number of sub-80 nm particles;
10 however, increases in global-mean particle number concentration are less than 1% in all size classes (Table 2). Conversely, global-mean particle number concentration increases by greater than 4% in all size classes in the SIZE30 simulation. The increased particle number concentration does lead to a stronger suppression of nucleation; however, as emission GMD is now 30 nm, particle number concentration changes become positive at 20 nm (compared to 80 nm for BASE). Across all
15 simulations, our particle number concentration is most sensitive to decreasing the value of the GMD of the assumed size distribution. Thus, understanding near-source, sub-grid-scale aging, which takes primary combustion particles from diameters around 30 nm to larger sizes (e.g. Sakamoto et al. 2016), is very important in determining the size-distribution effects of waste combustion. Across all sensitivity simulations, we find a range in the global-mean particle number concentration changes due to waste combustion of -0.2 to 4.0% for N10, 0.2 to 1.8% in N40, 0.5 to 5.4% in N80, and 0.4 to 4.4% in N150.

20 3.4 Aerosol direct radiative effect

Table 3 and Figure 6 show the global-mean all-sky DRE from waste combustion for the six different mixing-state assumptions for the BASE-WASTE_OFF comparison as well as other sensitivity simulations. Under the BASE assumptions, the global-mean DRE is negative for all mixing states, ranging from -5 mW m⁻² to -20 mW m⁻². Globally, the OA to BC ratio from waste-combustion emissions is around 14 to 1, leading to increased scattering and a predominantly negative or cooling
25 DRE. The relative emission mass of OA and BC is a source of uncertainty in Wiedinmyer et al. (2014). In Kodros et al. (2015), we found that the sign of the DRE is sensitive to the OA to BC ratio, where increasing the relative amount of BC leads to more positive (less negative) DRE. As in previous studies, the DRE becomes more positive when BC is assumed to be mixed internally (represented as core-shell, homogenous, and ext*1.5) as opposed to externally mixed and when OA is assumed to absorb as brown carbon.

30 Figure 7 shows maps of annual-mean all-sky DRE from waste combustion for all mixing states for BASE-WASTE_OFF. Waste combustion causes a largely negative DRE in much of Asia, Eastern Europe, and the Americas. In regions with bright surfaces (high albedo), such as the Sahara desert or Greenland ice sheet, the aerosol mixture has a high-enough absorbing component to decrease planetary albedo and cause a positive or warming DRE. The largest-magnitude

Jack 5/16/2016 2:32 PM
Deleted: our

Jack 5/16/2016 2:32 PM
Deleted: s

DRE from waste combustion occurs in eastern China (ranging from -0.4 to -0.1 W m^{-2}), as well as northern India and Eastern Europe (-0.3 to -0.05 W m^{-2}).

Across all simulations and mixing-state combinations, the global-mean DRE from waste combustion ranges from -40 mW m^{-2} to $+4 \text{ mW m}^{-2}$ (Table 3). In simulations HIGHMASS and LOWMASS, the factor-of-2 uncertainty in emission mass leads to approximately a factor-of-2 scale in DRE. In PRIMARY, the removal of the emission gas-phase condensable vapors results in a reduction of sub-200 nm particles, which does not greatly affect the volume distribution (Figure 5b). As a result, the global-mean DRE in PRIMARY is similar to that in BASE. The slightly more negative DRE in the internal mixtures (core-shell and homogenous) may be a result of smaller shell diameters due to reduced condensation. Both SIZE200 and SIZE30 result in less negative DRE. In SIZE200, the mass distribution is moved to larger sizes resulting in decreases in mass scattering and absorption efficiencies (the size distribution in the BASE simulation has nearly optimal mass scattering and absorption efficiencies). However, the fractional decrease in scattering efficiency is greater than that for absorption, resulting in a lower single-scattering albedo and an overall decrease in the magnitude of cooling. In SIZE30, the mass distribution is shifted to smaller sizes, which again lowers mass scattering and absorption efficiencies with a greater decrease in scattering relative to absorption. Again, this lowers the single-scattering albedo and decreases the magnitude of cooling.

3.5 Cloud-albedo aerosol indirect effect

Figure 8 shows the cloud-albedo AIE from domestic-waste combustion in the BASE-WASTE_OFF simulation. Waste-combustion emissions increase CDNC, and thus cloud reflectivity leading to a negative AIE in Central and South America, Africa, and the Pacific Islands. These are regions where the microphysical feedback suppressing nucleation and growth is less dominant, and waste combustion leads to increases in N40 (Figure 4). Conversely, despite increases in aerosol mass from waste combustion in Asia and Europe, the AIE is zero or slightly positive. This is caused by feedbacks in aerosol and cloud microphysics. The 'suppression of nucleation and growth' feedback (described earlier) leads to decreases in N40 and thus CDNC. In addition, heavy anthropogenic pollution in India and China results in a strong competition for water vapor. Increases in N80 and N150 (Figure 4) decrease the maximum supersaturation achieved in updrafts, which further limits droplet activation. As a result of the competing regional effects, the global-mean AIE for BASE-WASTE_OFF is -13 mW m^{-2} (Table 3).

The global, annual-mean AIE for all simulations is listed in Table 3 and plotted in Figure 6. In the global mean, the AIE ranges from -4 mW m^{-2} to -49 mW m^{-2} based on all emission uncertainties. In general, AIE is stronger in simulations with greater particle-number increases from waste combustion (SIZE30, HIGHMASS) and weakest in simulations with smaller particle-number increases (SIZE200, PRIMARY, LOWMASS). The largest-magnitude AIE occurs in the SIZE30 simulation (-49 mW m^{-2}) due to the large increase in particle number concentration (Table 2 and Figure 5). The lowest magnitude AIE occurs in the PRIMARY simulation, where condensable vapors from waste-combustion emissions are

Jack 5/16/2016 2:36 PM

Deleted: dominate

removed, leading to fewer particles in the 40-200 nm size range relative to BASE (Figure 5). This implies a non-trivial contribution of nucleation and growth to CCN concentrations from the NH_3 and SO_2 emissions from trash combustion.

4. Conclusions

In this paper, we use a global chemical-transport model to provide a first estimate of aerosol concentrations and radiative impacts from open, uncontrolled combustion of domestic waste. Using emissions from Wiedinmyer et al. (2014), we find increases in global-mean boundary-layer BC and OA mass of 8.5% and 5.6%, with regional increases exceeding 40%, relative to simulations without waste-combustion emissions. In our base simulations, we estimate domestic-waste combustion emissions lead to a global-mean DRE -5 to -20 mW m^{-2} (range due to optical assumptions) and an AIE of -13 mW m^{-2} . In areas where open waste-combustion occurs, the resulting aerosol radiative impacts can range from -0.1 to -0.4 W m^{-2} . On a global scale, we find minor improvements of model AOD compared to observed AERONET AOD when including waste-combustion emissions.

We include several sensitivity simulations to explore uncertainties in waste-combustion emission size distribution, gas-phase species (leading to secondary aerosol), and emitted particulate mass. In the global, annual mean, we estimate a range in the DRE of +4 to -40 mW m^{-2} and in the AIE of -4 to -49 mW m^{-2} . We find altering the waste-combustion emission size distribution while keeping total mass constant (SIZE200 and SIZE30) leads to substantial changes in the modeled size distribution, thus affecting the AIE (-8 to -49 mW m^{-2}) and to a lesser extent the DRE (+4 to -18 mW m^{-2}). In PRIMARY, we emit only primary BC and OA particles from waste combustion (and not particle-precursor gases). In this simulation, waste combustion leads to a stronger suppression of nucleation and growth, smaller increases in particle number concentration, and a lower magnitude AIE (-4 mW m^{-2}). Thus, the particle-precursor gases emitted from waste-combustion are important contributors to new-particle formation and growth, and they contribute to the negative AIE from waste-combustion emissions. We introduce a factor of 2 uncertainty in emission mass in HIGHMASS and LOWMASS. We find this leads to a near-linear increase in BC and OA mass and DRE (-2.5 mW m^{-2} to -40 mW m^{-2}). Scaling emission mass up/down also increases/decreases emitted particle number concentration, leading to a range of AIE of -7 to -24 mW m^{-2} . There is little information on emission size distribution and optical properties from waste combustion. Better knowledge of these parameters along with continued validation of emission mass fluxes will reduce model uncertainty.

Our estimates for the aerosol radiative effects from waste combustion are relatively small compared to the total aerosol burden; however, they are of similar magnitude to recent estimates of radiative impacts from biofuel combustion (-0.06 to +0.08 W m^{-2} for the DRE and -0.05 to +0.01 W m^{-2} for the AIE), another emission source associated with developing countries (Bond et al., 2013; Butt et al., 2016; Kodros et al., 2015; Lacey et al., 2015). Based on our estimates, it appears that the aerosol radiative effects from waste combustion likely have an overall cooling effect.

In a separate study (Kodros et al., 2016), we estimated that this waste-combustion aerosol leads to 284,000 (95% CI: 188,000-404,000) premature deaths globally each year. If waste-combustion aerosol is reduced to protect human health,

this may also lead to positive radiative forcing (warming tendency) due to a reduction of the waste-combustion-aerosol cooling. A reduction in waste combustion would also lead to a change in greenhouse gases, although these greenhouse-gas estimates are out of the scope of this paper due to the complex nature of greenhouse-gas emissions from waste decomposition and/or recycling. The greenhouse-gas changes due to choice of waste-removal method is an important topic
5 to be studied in the future.

10

Acknowledgements

This research has been supported by a grant from the U.S. Environmental Protection Agency's Science to Achieve Results (STAR) program through grant #83543801. Although the research described in the article has been funded by the U.S. Environmental Protection Agency's STAR program, it has not been subjected to any EPA review and therefore does not
15 necessarily reflect the views of the Agency, and no official endorsement should be inferred. David Ridley was funded through National Aeronautics and Space Administration grant: NASA NN14AP38G.

References

Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation 3. Sectional representation, *J. Geophys. Res.*, 107(D3), 4026, doi:10.1029/2001JD000483, 2002.
20 Adams, P. J. and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, *J. Geophys. Res.*, 107(D19), 4370, doi:10.1029/2001JD001010, 2002.

Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D. and Wennberg, P. O.:
25 Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11(9), 4039–4072, doi:10.5194/acp-11-4039-2011, 2011.

Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys. Discuss.*, 6, 3419–3463, doi:10.5194/acpd-6-3419-2006, 2006.
30 Ban-Weiss, G. A., Lunden, M. M., Kirchstetter, T. W. and Harley, R. A.: Size-resolved particle number and volume emission factors for on-road gasoline and diesel motor vehicles, *J. Aerosol Sci.*, 41, 5–12, doi:10.1016/j.jaerosci.2009.08.001, 2010.

Bauer, S. E., Menon, S., Koch, D., Bond, T. C. and Tsigaridis, K.: A global modeling study on carbonaceous aerosol
35 microphysical characteristics and radiative effects, *Atmos. Chem. Phys.*, 10, 7439–7456, doi:10.5194/acp-10-7439-2010, 2010.

Bohren, C. F. and Huffman, D. R.: Absorption and scattering of light by small particles., 1983.

5 Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H. and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109(D14), D14203, doi:10.1029/2003JD003697, 2004.

Bond, T. C., Habib, G. and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, *J. Geophys. Res.*, 111(D20), D20211, doi:10.1029/2006JD007315, 2006.

10 Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G. and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850–2000, *Global Biogeochem. Cycles*, 21(2), doi:10.1029/2006GB002840, 2007.

15 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G. and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res. Atmos.*, 118(11), 5380–5552, doi:10.1002/jgrd.50171, 2013.

20 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kermenian, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Sathesh, S. K., Sherwood, S., Stevens, B. and Zhang, X. Y.: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.

25 Butt, E. W., Rap, A., Schmidt, A., Scott, C. E., Pringle, K. J., Reddington, C. L., Richards, N. A. D., Woodhouse, M. T., Ramirez-Villegas, J., Yang, H., Vakkari, V., Stone, E. A., Rupakheti, M., S. Praveen, P., G. van Zyl, P., P. Beukes, J., Josipovic, M., Mitchell, E. J. S., Sallu, S. M., Forster, P. M. and Spracklen, D. V.: The impact of residential combustion emissions on atmospheric aerosol, human health, and climate, *Atmos. Chem. Phys.*, 16(2), 873–905, doi:10.5194/acp-16-873-2016, 2016.

30 Carrico, C. M., Petters, M. D., Kreidenweis, S. M., Sullivan, A. P., McMeeking, G. R., Levin, E. J. T., Engling, G., Malm, W. C. and Collett, J. L.: Water uptake and chemical composition of fresh aerosols generated in open burning of biomass, *Atmos. Chem. Phys.*, 10(11), 5165–5178, doi:10.5194/acp-10-5165-2010, 2010.

35 Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Hansen, J. E. and Hofmann, D. J.: Climate forcing by anthropogenic aerosols., *Science*, 255, 423–430, doi:10.1126/science.255.5043.423, 1992.

40 Chen, Y. and Bond, T. C.: Light absorption by organic carbon from wood combustion, *Atmos. Chem. Phys.*, 10(4), 1773–1787, doi:10.5194/acp-10-1773-2010, 2010.

Christian, T. J., Yokelson, R. J., Cárdenas, B., Molina, L. T., Engling, G. and Hsu, S.-C.: Trace gas and particle emissions from domestic and industrial biofuel use and garbage burning in central Mexico, *Atmos. Chem. Phys.*, 10(2), 565–584, doi:10.5194/acp-10-565-2010, 2010.

45 Chung, S. H. and Seinfeld, J. H.: Climate response of direct radiative forcing of anthropogenic black carbon, *J. Geophys. Res.*, 110, D11102, doi:10.1029/2004JD005441, 2005.

50

D'Andrea, S. D., Häkkinen, S. a. K., Westervelt, D. M., Kuang, C., Levin, E. J. T., Kanawade, V. P., Leaitch, W. R., Spracklen, D. V., Riipinen, I. and Pierce, J. R.: Understanding global secondary organic aerosol amount and size-resolved condensational behavior, *Atmos. Chem. Phys.*, 13(22), 11519–11534, doi:10.5194/acp-13-11519-2013, 2013.

5 D'Andrea, S. D., Acosta Navarro, J. C., Farina, S. C., Scott, C. E., Rap, A., Farmer, D. K., Spracklen, D. V., Riipinen, I. and Pierce, J. R.: Aerosol size distribution and radiative forcing response to anthropogenically driven historical changes in biogenic secondary organic aerosol formation, *Atmos. Chem. Phys.*, 15(5), 2247–2268, doi:10.5194/acp-15-2247-2015, 2015.

10 Hammer, M. S., Martin, R. V., van Donkelaar, A., Buchard, V., Torres, O., Ridley, D. A. and Spurr, R. J. D.: Interpreting the Ultraviolet Aerosol Index observed with the OMI satellite instrument to understand absorption by organic aerosols: implications for atmospheric oxidation and direct radiative effects, *Atmos. Chem. Phys. Discuss.*, 15(19), 27405–27447, doi:10.5194/acpd-15-27405-2015, 2015.

15 Heald, C. L., Ridley, D. A., Kroll, J. H., Barrett, S. R. H., Cady-Pereira, K. E., Alvarado, M. J. and Holmes, C. D.: Contrasting the direct radiative effect and direct radiative forcing of aerosols, *Atmos. Chem. Phys.*, 14(11), 5513–5527, doi:10.5194/acp-14-5513-2014, 2014.

20 Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I. and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sens. Environ.*, 66(1), 1–16, doi:10.1016/S0034-4257(98)00031-5, 1998.

25 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A. and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.*, 113(D13), D13103, doi:10.1029/2008JD009944, 2008.

Jacobson, M. Z.: Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols., *Nature*, 409(6821), 695–7, doi:10.1038/35055518, 2001.

30 Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B. and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from *in situ* and remote sensing observations, *Atmos. Chem. Phys.*, 11(7), 3137–3157, doi:10.5194/acp-11-3137-2011, 2011.

Janhäll, S., M. Jonsson, Å., Molnár, P., A. Svensson, E. and Hallquist, M.: Size resolved traffic emission factors of submicrometer particles, *Atmos. Environ.*, 38(26), 4331–4340, doi:10.1016/j.atmosenv.2004.04.018, 2004.

35 Janhäll, S., Andreae, M. O. and Pöschl, U.: Biomass burning aerosol emissions from vegetation fires: particle number and mass emission factors and size distributions, *Atmos. Chem. Phys.*, 10(3), 1427–1439, doi:10.5194/acp-10-1427-2010, 2010.

Janssens-Maenhout, A., Petrescu, A., Muntean, M., and Blujdea, V.: Verifying Greenhouse Gas Emissions: Methods to Support International Climate Agreements, The National Academies Press, Washington, DC., available at: http://www.nap.edu/openbook.php?record_id=12883 (last access: 17 March 2014), 2010.

40 Jung, J., Fountoukis, C., Adams, P. J. and Pandis, S. N.: Simulation of *in situ* ultrafine particle formation in the eastern United States using PMCAMx-UF, *J. Geophys. Res.*, 115(D3), D03203, doi:10.1029/2009JD012313, 2010.

Kodros, J. K., Scott, C. E., Farina, S. C., Lee, Y. H., L'Orange, C., Volckens, J. and Pierce, J. R.: Uncertainties in global aerosols and climate effects due to biofuel emissions, *Atmos. Chem. Phys.*, 15(15), 8577–8596, doi:10.5194/acp-15-8577-2015, 2015.

Kodros, J. K., Wiedinmyer, C., Ford, B., Cucinotta, R., Gan, R., Magzamen, S., Pierce, J.R.: Global burden of mortalities due to chronic exposure to ambient PM_{2.5} from open combustion of domestic waste, submitted to Environ. Health Perspect., 2016.

5 Kopke, P., Hess, M., Schult, I., and Shettle, E. P.: Global Aerosol Data Set, Max Planck Inst. für Meteorol., Hamburg, Germany, 1997.

Kuhns, H., Knipping, E. M. and Vukovich, J. M.: Development of a United States–Mexico Emissions Inventory for the Big Bend Regional Aerosol and Visibility Observational (BRAVO) Study, *J. Air Waste Manage. Assoc.*, 55(5), 677–692, 10 doi:10.1080/10473289.2005.10464648, 2005.

15 Lacey, F. and Henze, D.: Global climate impacts of country-level primary carbonaceous aerosol from solid-fuel cookstove emissions, *Environ. Res. Lett.*, 10(11), 114003 [online] Available from: <http://stacks.iop.org/1748-9326/10/i=11/a=114003>, 2015.

Lee, H. J. J., Aiona, P. K., Laskin, A., Laskin, J. and Nizkorodov, S. A.: Effect of solar radiation on the optical properties and molecular composition of laboratory proxies of atmospheric brown carbon., *Environ. Sci. Technol.*, 48(17), 10217–26, doi:10.1021/es502515r, 2014.

20 Lee, L. a., Pringle, K. J., Reddington, C. L., Mann, G. W., Stier, P., Spracklen, D. V., Pierce, J. R. and Carslaw, K. S.: The magnitude and causes of uncertainty in global model simulations of cloud condensation nuclei, *Atmos. Chem. Phys.*, 13(17), 8879–8914, doi:10.5194/acp-13-8879-2013, 2013.

25 Lee, Y. H. and Adams, P. J.: A Fast and Efficient Version of the TwO-Moment Aerosol Sectional (TOMAS) Global Aerosol Microphysics Model, *Aerosol Sci. Technol.*, 46(6), 678–689, doi:10.1080/02786826.2011.643259, 2012.

Lee, Y. H., Pierce, J. R. and Adams, P. J.: Representation of nucleation mode microphysics in a global aerosol model with sectional microphysics, *Geosci. Model Dev.*, 6, 1221–1232, doi:10.5194/gmd-6-1221-2013, 2013.

30 McMeeking, G. R., Fortner, E., Onasch, T. B., Taylor, J. W., Flynn, M., Coe, H. and Kreidenweis, S. M.: Impacts of nonrefractory material on light absorption by aerosols emitted from biomass burning, *J. Geophys. Res. Atmos.*, 119(21), 12,272–12,286, doi:10.1002/2014JD021750, 2014.

35 Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M. and Jimenez, J. L.: Characterization of Primary Organic Aerosol Emissions from Meat Cooking, Trash Burning, and Motor Vehicles with High-Resolution Aerosol Mass Spectrometry and Comparison with Ambient and Chamber Observations, *Environ. Sci. Technol.*, 43(7), 2443–2449, doi:10.1021/es8011518, 2009.

40 Napari, I., Kulmala, M. and Vehkamäki, H.: Ternary nucleation of inorganic acids, ammonia, and water, *J. Chem. Phys.*, 117, 8418–8425, doi:10.1063/1.1511722, 2002.

Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7(8), 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.

45 Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation and primary emission rates, *Atmos. Chem. Phys.*, 9(4), 1339–1356, doi:10.5194/acp-9-1339-2009, 2009.

Pincus, R., Barker, H. W., and Morcrette, J. J.: A fast, flexible, approximate technique for computing radiative transfer in inhomogeneous cloud fields, *J. Geophys. Res.-Atmos.*, 108, 4376, doi:10.1029/2002jd003322, 2003.

50

Rap, A., Scott, C. E., Spracklen, D. V., Bellouin, N., Forster, P. M., Carslaw, K. S., Schmidt, A. and Mann, G.: Natural aerosol direct and indirect radiative effects, *Geophys. Res. Lett.*, 40, 3297–3301, doi:10.1002/grl.50441, 2013.

Sakamoto, K. M., Allan, J. D., Coe, H., Taylor, J. W., Duck, T. J. and Pierce, J. R.: Aged boreal biomass-burning aerosol size distributions from BORTAS 2011, *Atmos. Chem. Phys.*, 15(4), 1633–1646, doi:10.5194/acp-15-1633-2015, 2015.

Sakamoto, K. M., Stevens, R. G., Pierce, J. R.: The evolution of biomass-burning aerosol size distributions due to coagulation: dependence on fire and meteorological details and parameterization, submitted to *Atmos. Chem. Phys.* 2016.

10 Saleh, R., Hennigan, C. J., McMeeking, G. R., Chuang, W. K., Robinson, E. S., Coe, H., Donahue, N. M. and Robinson, a. L.: Absorptivity of brown carbon in fresh and photo-chemically aged biomass-burning emissions, *Atmos. Chem. Phys.*, 13(15), 7683–7693, doi:10.5194/acp-13-7683-2013, 2013.

15 Saleh, R., Robinson, E. S., Tkacik, D. S., Ahern, A. T., Liu, S., Aiken, A. C., Sullivan, R. C., Presto, A. A., Dubey, M. K., Yokelson, R. J., Donahue, N. M. and Robinson, A. L.: Brownness of organics in aerosols from biomass burning linked to their black carbon content, *Nat. Geosci.*, 7(August), 647–650, doi:10.1038/NGEO2220, 2014.

20 Scott, C. E., Rap, a., Spracklen, D. V., Forster, P. M., Carslaw, K. S., Mann, G. W., Pringle, K. J., Kivekäs, N., Kulmala, M., Lihavainen, H. and Tunved, P.: The direct and indirect radiative effects of biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 14(1), 447–470, doi:10.5194/acp-14-447-2014, 2014.

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons., 2012.

25 Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, A. and Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos. Chem. Phys.*, 11, 12109–12136, doi:10.5194/acp-11-12109-2011, 2011.

30 Streets, D. G., Bond, T. .., Carmichael, G. R., Fernandes, S. .., Fu, Q., He, D., Klimont, Z., Nelson, S. .., Tsai, N. Y., Wang, M. .., Woo, J.-H. and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *J. Geophys. Res.*, 108(D21), 8809, doi:10.1029/2002JD003093, 2003.

Twomey, S.: Pollution and the planetary albedo, *Atmos. Environ.*, 8(12), 1251–1256 [online] Available from: <http://www.sciencedirect.com/science/article/pii/0004698174900043> (Accessed 11 December 2014), 1974.

35 Van Der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., Defries, R. S., Jin, Y. and Van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

40 Vehkamäki, H.: An improved parameterization for sulfuric acid–water nucleation rates for tropospheric and stratospheric conditions, *J. Geophys. Res.*, 107(D22), 4622, doi:10.1029/2002JD002184, 2002.

Vestreng, V., Ntziachristos, L., Semb, A., Reis, S., Isaksen, I. S. A. and Tarrasón, L.: Evolution of NO_x emissions in Europe with focus on road transport control measures, *Atmos. Chem. Phys.*, 9(4), 1503–1520, doi:10.5194/acp-9-1503-2009, 2009.

45 Wang, X., Heald, C. L., Ridley, D. a., Schwarz, J. P., Spackman, J. R., Perring, a. E., Coe, H., Liu, D. and Clarke, a. D.: Exploiting simultaneous observational constraints on mass and absorption to estimate the global direct radiative forcing of black carbon and brown carbon, *Atmos. Chem. Phys.*, 14(20), 10989–11010, doi:10.5194/acp-14-10989-2014, 2014.

Westervelt, D. M., Pierce, J. R., Riipinen, I., Trivitayanurak, W., Hamed, A., Kulmala, M., Laaksonen, A., Decesari, S. and Adams, P. J.: Formation and growth of nucleated particles into cloud condensation nuclei: model-measurement comparison, *Atmos. Chem. Phys.*, 13(15), 7645–7663, doi:10.5194/acp-13-7645-2013, 2013.

5 Westervelt, D. M., Pierce, J. R. and Adams, P. J.: Analysis of feedbacks between nucleation rate, survival probability and cloud condensation nuclei formation, *Atmos. Chem. Phys.*, 14(11), 5577–5597, doi:10.5194/acp-14-5577-2014, 2014.

10 Wiedinmyer, C., Yokelson, R. J. and Gullett, B. K.: Global emissions of trace gases, particulate matter, and hazardous air pollutants from open burning of domestic waste., *Environ. Sci. Technol.*, 48(16), 9523–30, doi:10.1021/es502250z, 2014.

15 Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B. and Lin, J.-T.: [Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, Atmos. Chem. Phys., 11\(7\), 3137–3157, doi:10.5194/acp-11-3137-2011](#), 2011. Zender, C. S.: Mineral Dust Entrainment and Deposition (DEAD) model: Description and 1990s dust climatology, *J. Geophys. Res.*, 108(D14), 4416, doi:10.1029/2002JD002775, 2003.

15 Zhao, R., Lee, A. K. Y., Huang, L., Li, X., Yang, F. and Abbatt, J. P. D.: Photochemical processing of aqueous atmospheric brown carbon, *Atmos. Chem. Phys.*, 15(11), 6087–6100, doi:10.5194/acp-15-6087-2015, 2015.

Table 1. Description of simulations

Simulation	GMD ^a	Mass Scale Factor	Species
BASE	100 nm	1.0	primary and secondary
SIZE200	200 nm	1.0	primary and secondary
SIZE30	30 nm	1.0	primary and secondary
PRIMARY	100 nm	1.0	primary BC and OA only
HIGHMASS	100 nm	2.0	primary and secondary
LOWMASS	100 nm	0.5	primary and secondary
WASTE_OFF	N/A	0.0	None

^aGMD = geometric number-mean diameter

Table 2. Percent change in global, annual-mean N10, N40, N80, N150, BC mass, OA mass from domestic-waste combustion (i.e. relative to WASTE_OFF simulation).

Simulation	N10 [%]	N40 [%]	N80 [%]	N150 [%]	BC [%]	OA [%]
BASE	0.86	0.90	1.04	1.16	8.46	5.60
SIZE200	0.86	0.55	0.46	0.36	8.36	5.53
SIZE30	4.02	5.43	5.42	4.40	8.07	5.14
PRIMARY	-0.17	0.232	0.52	0.89	8.47	5.61
HIGHMASS	1.68	1.77	2.06	2.31	16.9	11.2
LOWMASS	0.44	0.45	0.53	0.59	4.23	2.80

Table 3. Global-mean all-sky direct radiative effect and cloud-albedo aerosol indirect effect relative to the WASTE_OFF simulation for all sensitivity simulations and mixing-state assumptions

Simulation	Direct Radiative Effect [mW m ⁻²]						AIE [mW m ⁻²]
	Core-Shell, BrC	Ext*1.5, BrC	Homogenous	Core-Shell	Ext*1.5	External	
BASE	-5.0	-8.7	-11	-13	-17	-20	-13
SIZE200	+3.2	-6.0	-3.7	-8.7	-16	-18	-8.3
SIZE30	+4.1	+3.7	-1.5	-1.6	-2.3	-5.8	-49
PRIMARY	-5.7	-9.3	-12	-14	-17	-20	-4.0
HIGHMASS	-10	-17	-22	-26	-34	-40	-24
LOWMASS	-2.5	-4.3	-5.6	-6.5	-8.7	-10	-6.7




Figure 1. Emission fluxes from the open combustion of domestic waste for (a) organic aerosol, (b) black carbon, and (c) NH_3 from Wiedinmyer et al. (2014).




Figure 2. One-to-one plots of time-average AOD for BASE (a,c,e) and WASTE_OFF (b,d,f) compared to AERONET observations at 380 nm (a,b), 500 nm (c,d), and 1020 nm (e,f). The values in the box give the slope of the linear regression (m), the correlation (r^2), and the log-mean bias (LMB), while the black dashed line represents the 1:1 line and the red line is the result of the linear regression.




Figure 3. Percent changes in annually averaged boundary-layer (a) OA, (b) BC, (c) NH_3 , and (d) $\text{PM}_{2.5}$ mass due to domestic-waste combustion (BASE-WASTE_OFF).




Figure 4. Annual-mean percent change in boundary-layer (a) N10, (b) N40, (c) N80, (d) N150 for domestic-waste combustion (BASE-WASTE_OFF).




Figure 5. The change in global-mean (a) number distribution and (b) volume distribution for the different sensitivity studies relative to the WASTE-OFF simulation.




Figure 6. Global-mean all-sky direct radiative effect (DRE) and cloud-albedo aerosol indirect effect (AIE) for the baseline (BASE), aerosol-phase emissions only (PRIMARY), larger (SIZE200) and smaller (SIZE30) geometric mean diameter, and doubled (HIGHMASS) and halved (LOWMASS) emission mass relative to the WASTE_OFF simulation. The gold bars represent the DRE for the core-shell mixing state, while the error bars show the range due to the other optical assumptions.




Figure 7. The all-sky direct radiative effect for domestic-waste combustion (BASE-WASTE_OFF) assuming a (a) core-shell with brown carbon, (b) homogenous internal, (c) core-shell, (d) external with brown carbon and an absorption enhancement factor of 1.5, (e) external with an absorption enhancement factor of 1.5, and (f) external mixing state.





Figure 8. The cloud-albedo aerosol indirect effect difference between BASE-WASTE_OFF.