

Interactive comment on “STUDY OF AFRICAN DUST WITH MULTI-WAVELENGTH RAMAN LIDAR DURING THE “SHADOW” CAMPAIGN IN SENEGAL” by I. Veselovskii et al.

Anonymous Referee #1

Received and published: 11 March 2016

This paper presents case studies of multi-wavelength lidar measurements of dust including lidar ratio and angstrom exponent, adding to the published literature of case studies for dust lidar measurements. In addition, the paper applies a microphysical inversion scheme to retrieve volume concentration, effective radius and real refractive index; however, these inversions are affected by known problems with the assumptions which may affect the results. For instance, the authors use simulations to make a useful demonstration that backscatter angstrom exponent is sensitive to wavelength dependence of the imaginary refractive index, and that neglecting this spectral dependence can result in large errors in retrieved quantities. The authors also point out potential difficulties with the spheroidal model used to represent the optical properties of the

C1

dust particles and show that this prevents the effective use of depolarization measurements in the retrieval. I find this paper to be useful given the importance of determining dust optical and microphysical properties, and in particular due to the importance of understanding the limitations of the tools currently available for determining them (and for motivating increased effort in improving those tools if possible).

However this is not the first discussion of either of these sources of retrieval bias. There's a definite conclusion that spectral dependence of the imaginary refractive index needs to be taken into account, but it's not shown how to do that in this paper. It's also not clear that the results in this study really justify the use of the spheroidal model. Instead of providing solutions to these difficulties, I believe the authors' aim is to show that some quantities can still be determined with acceptable accuracy even given these major difficulties. The biggest issue I have with this manuscript is that this premise is not clearly spelled out and is not well supported. I could be convinced that this may be true in some circumstances, but I do not find enough supporting evidence in the manuscript, or enough information to determine under what circumstances it could be true. More discussion and supporting evidence for this assertion is needed.

More information about the uncertainties in the measurements and retrievals is also needed, including error bars for all quantities at different altitudes, and also information about the vertical resolution of the measurements and retrievals.

SPECIFIC COMMENTS

Page 4, line 13. Three references are given for the statement “For typical dust PSDs the AERONET model provides lidar and depolarization ratios which agree reasonably well with observed values”. I feel that these citations are misleading. The first paper, Wiegner et al. 2009, is the only one of these three that actually addresses the AERONET model. They find many limitations to the model and find that sometimes the agreement is good and sometimes quite poor. Sometimes they can improve the agreement only by shifting the refractive index away the measured values, which to my

C2

mind calls into question whether the agreement should be considered “reasonable”. I can find nothing at all about the AERONET model in the second paper, Esselborn et al. 2009. The third paper, by Tesche et al. 2009, has no new information and only quotes Wiegner et al. 2009 plus another paper which at the time was a future paper. I don't know if this future paper was published, but if so then this manuscript should probably be using that reference rather than a secondary reference.

Page 4, line 14. What kind of results did these “first attempts” find?

Page 5, line 1-2. “more measurements are needed”. I certainly agree with this, but I think you have even more strongly motivated the need for improved particle modeling, both by your introduction and in the results and analysis that follow. You might consider discussing this significant need in the conclusion section.

Page 5, line 3 ff. When first encountering this paragraph I was not familiar enough with dust climatology in Africa to understand why a field mission would be designed for this time of year. It would be helpful if you'd consider adding a sentence or two about the meteorological/climatological conditions, with references, somewhere near this paragraph in order to help other readers who are unfamiliar. For example, the explanation of the Harmattan mechanism and references that were at the beginning of what was section 4.3 of the original manuscript would be fine.

Page 6, line 4 ff. Are the micropulse lidar measurements used in this paper? If not, then probably remove this section.

Page 7, line 6. Uncertainty in depolarization is estimated as plus or minus 15%. Since depolarization ratio is often expressed as a percentage, “15%” could mean either absolute or relative error. Please resolve the ambiguity.

Page 7, line 14. What is the vertical resolution of the backscatter, extinction, and depolarization ratio measurements? Not the range bin size but the effective vertical resolution: that is, how many independent measurements are in one of these profiles?

C3

Section 3. I found this analysis combining the aerosol and wind measurements, plus back-trajectories, to be interesting. However, I also have some confusion about how the lidar properties reflect the characterization of these airmasses. Adding additional analysis such as was done with the other three case studies in sections 4 and 5 would really help to strengthen the logical cohesion of the paper, to better understand this observation, and also perhaps to improve the diversity of the case studies presented.

I am specifically confused about the near uniformity of the particle depolarization ratio and lidar ratio in the observations from 15-16 April. Even where the airmass is being described as “continentalized maritime” the particle depolarization ratio is in excess of 25%, perhaps implying a significant amount of dust even in this layer. Likewise, the lidar ratio appears to be quite high everywhere, much higher than would be expected for marine aerosol. Any comments about this?

Page 9, line 3 and Figures 1 and 4. It's confusing that the labels for the layers in Figure 4 don't match the labels A,B,C,D from figure 1. When “Layer B” is mentioned in the text in the discussion of figure 4, it's not clear whether this is Figure 1's layer B or Figure 4's layer B.

Page 9, line 1 “during the first part of the observation period, it is lower”. Please quantify. It's difficult to read quantitative results off the color charts, so it's important to have numbers in the text.

Page 9, line 15-17, and Fig 5-7. Selections are made to guarantee major dust contribution. I think this choice to filter for only dust cases makes these figures extremely confusing. It would be much more helpful to have the full data set in the timeseries of Figures 5-7, with the observations that are dominated by dust indicated by shading or some other way. There is discussion about the variability of EAE and lidar ratio being due to dust episodes, but since only dust cases are shown, it's impossible to see the full range of variability. And it's confusing to attribute the variability within the displayed data set to dust episodes when everything shown is a dust episode. It also makes it

C4

impossible to compare the lidar-observed values in Figures 6 and 7 with the AERONET time series in Figure 5, since Figure 5 shows the complete time-series. This mismatch undercuts the usefulness of including the AERONET timeseries. For example, there's no very clear correlation in Figure 6 between the extinction and the angstrom exponents as there is in the AERONET time series, but I'm not sure if it's really less correlated, or if it just appears that way because of the missing data. Also, Figure 6 and Figure 7 do not include the same number of points. Since both are supposed to be selected by the same criteria, this seems like a simple oversight rather than intentional, but it adds to the difficulty in comparing and interpreting the data in these figures.

Figure 6-7, continued: please clarify the error bars. One error bar is shown for each lidar time series for the whole period, except LR355 which has none. Are the uncertainties the same for every observation? What is the uncertainty for LR355? Describe the error bars in the caption. Are they random or systematic, one sigma or two, etc.?

Section 4.2. It seems that all the case studies are pure dust, and all the depolarization ratios shown in all figures are quite high, except where the extinction and backscatter are lower perhaps implying there is not much aerosol at all. It would be useful to see a contrasting case, if one exists. If there is no case with significant aerosol that is not dominated by dust, then a case with a smaller dust fraction (perhaps the case from Figure 4) would still be good to see. Without getting to see some dynamic range of the measurements (again, except where the signal is much lower), it's less convincing that the measurements and retrievals are accurate. On the other hand, it may be that there just is no other aerosol type present at that time and place at high enough concentration to make much difference to the measurements. A fuller discussion of the instrument accuracy and uncertainties, including complete error bars on the figures, would help with understanding which variability is significant.

Page 11, line 8. This is a good point, and the following demonstration is useful. But BAE is a non-monotonic function of particle radius even for spectrally constant complex refractive index and spherical particles (i.e. just from Mie modeling). Is spectral depen-

C5

dence of the imaginary part of the refractive index the only way to achieve significantly negative BAE?

Page 11, line 12. Does the real part have a spectral dependence as well?

Page 11, continued. The large negative BAE signature is present in this case but not the previous case. Is this discussion meant to imply that there is spectral dependence in the imaginary refractive index only in some of these cases? What is the explanation for different dust layers having strong spectral dependence of the imaginary refractive index in some cases but not others?

Page 12, line 2. I agree the study demonstrates the importance of accounting for spectral dependence of the imaginary refractive index. Is it possible to do this using values of EAE and BAE more similar to the measurements for this case, to demonstrate more conclusively that this measurement case is in a regime where this effect is significant?

Page 12, lines 10-17. The upper layer where the intensive properties are different appears to have fairly small extinction and backscatter, meaning less signal. Error bars showing the uncertainties in the upper layers would make the attribution of this variability to differences in aerosol type more convincing.

Figures 9, 11, 14, 15, 16. Again, more details about the error bars would be good. Include error bars on all quantities at more than just one altitude and describe them in the captions. The text suggests that the uncertainties vary; the figures should reflect this.

Page 12, line 18. "The relative humidity on 10 April was higher than on 13, 29 March". Can you show RH for the other two cases also? Or at least quantify the values for the earlier cases in the text.

Page 13, lines 10-11. Please specify, is this the same version of the algorithm you are using in this study?

Page 13, line 13. Consider using a different acronym. It's difficult to remember later if

C6

“S” is spheres or spheroids. Maybe NSVF for “non-spherical volume fraction”.

Page 14, line 4. It would be useful to show the AERONET comparison on the figures.

Page 14, 2-12. What is the AERONET non-spherical fraction? Does this support the use of the assumption of 100% non-spherical fraction?

Page 14, line 17 and Figure 18. “the main features of the particle volume size distribution”. Can you be more specific about how much information is provided by inversion about the size distribution (how many bins or coefficients) and how it's obtained? These size distributions seem surprisingly detailed for an inversion of just 5 pieces of information. I suppose this is probably explained in an earlier paper, but a brief description of the inversion (probably in a new section between section 2 and 3) would still be helpful here in this paper.

Page 14, line 14. If the retrieved imaginary refractive index is unreliable due to errors associated with a faulty assumption in the retrieval, what evidence is there that the other retrieved variables are trustworthy? While I believe they may be, it doesn't seem obvious that this must be so. The question certainly deserves more discussion, if any of the retrieval results are to be considered useful.

Page 15, line 14. “Assuming that results obtained using 3-beta plus 2-alpha data are more representative of the actual values...” Similar to the previous comment: this seems like a big assumption and very important to the analysis of the results. If the measured depolarization ratio can't be reproduced by the spheroidal model but we want to believe the results of the inversion of backscatter and extinction only, then we need to be convinced that the spheroidal model can at least correctly determine backscatter and extinction. It seems that backscatter would be of particular concern, since, as pointed out in the introduction “the spheroid model was not specifically designed for lidar applications where scattering in the backward direction is considered.” Indeed the introduction suggests that previous studies found that the spheroid model also leads to errors in refractive index. Is there additional analysis that can be done to

C7

demonstrate the correctness of the inversion of the 3-beta plus 2-alpha data using the spheroid model or to better characterize the errors?

Page 15 and Figure 19. What is the effect of these two experiments on other retrieved quantities, like the volume concentration or the real refractive index?

Page 17, line 3. One of the conclusions is that for small enough depolarization the 3-backscatter + 2-extinction + 1-depolarization inversion permits the spherical/non-spherical fraction to be estimated. This isn't part of the analysis of the paper and there is no real support for it here; it might be better to delete it. If it is kept, then besides supporting it with further analysis, it would also be good to clarify whether this gives more complete or more accurate information than the spherical/non-spherical separation that has been practiced for lidar measurements for over a decade (Sugimoto and Lee, 2006; Tesche et al. 2009)

Figure 1, caption. What quantity is the “lidar signal”?

Figure 3, caption and annotation. It would be useful to explain the correspondence between the four trajectories and the regions A,B,C,D from Figure 1.

TECHNICAL COMMENTS

Page 16, line 20. “Somehow” = “somewhat”

Figure 4, the labels on the color axis are too small.

REFERENCES

Sugimoto, N., and C. H. Lee (2006), Characteristics of dust aerosols inferred from lidar depolarization measurements at two wavelengths, *Appl Optics*, 45(28), 7468-7474.

Tesche, M., A. Ansmann, D. Müller, D. Althausen, R. Engelmann, V. Freudenthaler, and S. Groß (2009), Vertically resolved separation of dust and smoke over Cape Verde using multiwavelength Raman and polarization lidars during Saharan Mineral Dust Experiment 2008, *J. Geophys. Res.*, 114(D13), D13202.

C8

