

The response to the reviewers' comments is in italic.

Anonymous Referee #1 Received and published: 31 January 2017

The manuscript uses ground- and satellite-based retrievals of cloud fraction, cloud liquid and ice water content and cloud phase profiles from lidar and radar to compare their performance at two Arctic sites: Barrow and Eureka. They propose to merge ground and satellite retrievals of cloud fraction to compensate for their inherent limitations: issues for CloudSat and CALIPSO to detect low-level clouds versus issues for surface based measurements to detect high clouds. I do recommend major revisions as there are some issues with the presentation of the results and the actual content of the conclusions.

*We appreciate the reviewer's valuable comments. The manuscript becomes better with revisions in response to reviewer's comments and suggestions.*

1. The method section needs some extensive work, because the explanations are currently confusing and insufficient. I have detailed the problems in the specific comments below. Are monthly means calculated and used throughout? This is never explicitly said.

*Changes have been made in the method section in the revised manuscript in response to reviewer's suggestions. Details can be found in the response to reviewer's specific comments below. Monthly means are calculated and used throughout, and this is specified in the revised manuscript.*

2. The detectability issue with CloudSat and CALIPSO for low level clouds is not new, there are already a number of papers that discuss this, e.g. Kay and Gettelman 2009, or Huang et al. (JCLI, 2012, doi: 10.1175/JCLI-D-11-00131.1). The real novelty of this paper is 1) to give an estimate/magnitude to this deficiency and 2) inspect the consequences when looking at the annual cycle of cloud cover in the Arctic. This should be made more prominent.

*The references the reviewers suggested have been added in the revised manuscript with correspondent discussion. The reviewer also summarized the novelty of our work well, and we highly appreciated that and have included that in the revised manuscript.*

3. The authors have decided to separate the results from Barrow from Eureka. Why is this? Are the two sites giving different results other than differing climatologies?

*We actually spent quite some time figuring out the best way to present the results, either separating by different physical parameters, e.g. cloud amount, cloud phase, and cloud water content, or by different locations, e.g. Barrow and Eureka. We then decided to go with the latter for clearer presentation. The climatologies at these two sites are not the same, so we do not think the content are redundant.*

4. Although a blended product is a good idea, because of the good performance overall of the surface-based observations (even if less high clouds are detected, the differences with the satellite based observations are small, possibly because of the location and type of clouds). I wonder if such a product is that needed for these two locations. It might be of more use if done for the tropics.

*We totally agree with the reviewer that such a blended product might be more useful in the tropics. We would like to argue that such products may be as valuable in the polar regions as*

*they are in the tropics because of the ubiquitous low-level clouds in the polar regions, and lack of detection capability from CloudSat and CALIPSO. Such discussions have been added in the revised manuscript.*

Specific comments:

1. The title is awkward: shouldn't "observations" be "observatories"? or add "sites" at the end.

*In the title, "observations" was changed to "observatories".*

2. Line 28, page 2: Here, and elsewhere, the authors refer to CloudSat&CALIPSO as "space-based radar-lidar" which makes it quite general when one could imagine that other (future) radars and lidars might have different sensitivities and consequently issues/ strengths. If for example the characteristics of the Earthcare mission instruments will be such that they will experience the same problems, then this should be said. Otherwise it would be better in the introduction to say that when referring to "space-based radar-lidar" the authors mean CloudSat and CALIPSO.

*Responding to reviewer's comment, the following text has been added in the revised manuscript. "Space-based radar and lidar in this paper refer to existing instruments, i.e. Cloud Profiling Radar (CPR) onboard the CloudSat and the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard the Cloud-Aerosol lidar and Infrared Pathfinder Satellite Observation (CALIPSO). However, the conclusions will likely be valid for the space-based radar and lidar instruments in the foreseeable future, i.e. the ATmospheric backscatter LIDar (ATLID), and the CPR onboard the EarthCARE mission (Heliere et al. 2007)."*

Section 2:

3. What is the temporal resolution of the profiles, surface and satellite based, when they are compared? Monthly means? Does it mean that the surface profiles are accumulated over a month and then cloud fraction calculated using a cloud mask? Please explain.

*The temporal resolution in the comparison is monthly. All surface profiles in a month are accumulated for calculation of monthly means. This has been added in the revised manuscript.*

4. Throughout the manuscript, please specify whether the lowest levels are identified about the surface or above mean sea level (which presumably is rather close at the two sites? This is not specified).

*The lowest levels are identified above the mean sea level. This has been added in the revised manuscript, "Monthly means are calculated for both surface observations and for the space-based sensors. All heights are above the mean sea level. All surface profiles in a month are accumulated for calculation of monthly means.", and "The vertical resolution of the calculated monthly means is interpolated to 100 m to be consistent with those from surface observations." In the last paragraph of section 2.*

5. Page 3, line 17: when introducing VFM, please specify which resolution, vertical or horizontal? Depending on which the 1/3, 1 and 5 km refer to, then specify the other resolution. This might help understand the method described on page 4 (see point 10 below)

*The following text has been added in the revised manuscript, "The Vertical Feature Mask (VFM) from CALIPSO's CALIOP provides cloud vertical distribution in up to 10 vertical layers at 5 km*

and 1 km horizontal resolutions, and up to 5 vertical layers at 1/3 km horizontal resolution (Vaughan et al. 2009). The vertical resolution is 30 m below 8.2 km, and 60 m between 8.2 and 20.2 km. A Selective Iterated Boundary Location (SIBYL) scheme is applied to detect all features within a given scene. Strongly scatter features, e.g. stratus clouds, can be identified in a single laser pulse, with the 1/3 km horizontal resolution, and these features are then removed in order to detect any surrounding aerosol layers. Weakly scattering features, e.g. thin cirrus clouds, are detected with the average of several laser pulses, e.g. 5 km horizontal resolution, for higher signal-to-noise ratio (Vaughan et al. 2005). Compared to the 1 km resolution data, the 5 km resolution product can identify weaker cloud features using an iterative multi-resolution averaging scheme (Vaughan et al. 2009). Combination of the cloud layer products at 5 km and 1/3 km provides a complete vertical distribution of clouds from CALIPSO (Vaughan et al. 2009, Vaughan et al. 2005).”.

6. When using GEOPROF, the authors choose the CPR\_cloud\_mask variable to be above 20 for a range bin to be cloudy. What is the convention in GEOPROF-LIDAR? How does this choice affect the results?

*The threshold in the GEOPROF-LIDAR is also 20. In the revised manuscript, we added, “This threshold is the same as that used in the 2B-GEOPROF-LIDAR (Mace et al. 2009, Mace et al. 2009). A false positive detection of 5% is estimated with this threshold in the 2B-GEOPROF-LIDAR (Mace et al. 2009)” The impact of the choice on the results is beyond the scope of this study.*

7. Line 20, page 4: here the authors specify that the satellite based profiles are selected if found within 50 km from the sites. Given the narrow swath and polar orbit, how many orbits per month actually fulfill this condition of at least one profile within 50 km? Do “6000 total sample numbers” and “1500 total sample numbers” refer to the total number of profiles?

*The text in the revised manuscript has been changed as “The monthly mean sample number of the satellite sensors is a function of latitude in the Arctic, with the fewest at 60° N, gradually increasing to a maximum around 80° N (Liu 2015). Both factors are reflected in the large number of samples at Eureka, with over 6000 total samples per month from June 2006 to December 2010 at Eureka, and around 1500 total samples at Barrow per month from middle February 2008 to December 2010.”.*

8. Page 4, lines 21-23: this sentence is confusing, maybe a simple schematic would help visualize what you mean? What is the original vertical resolution of each product?

*The vertical resolution is 30 m below 8.2 km, and 60 m between 8.2 and 20.2 km. The vertical resolution of 2B-GEOPROF and 2B-GEOPROF-lidar are at 240 m. We added this information in the revised manuscript.*

*A schematic would be great. But we did not figure out a way to make a simple schematic. So, we re-wrote the description to calculate the mean cloud vertical distribution.*

9. Lines 23-25 page 4: this sentence does not make any sense, what is a “cloud case number”? again maybe a schematic would help. Then at the end of the sentence “in a selected time period” refers to a month?

*A schematic would be great. But we did not figure out a way to make a simple schematic. So, we re-wrote the description to calculate the mean cloud vertical distribution as the following in the revised manuscript “Vertical profiles of all these products within 50 km of the two Arctic atmospheric observation sites, Barrow and Eureka, are extracted and archived. The cloud fraction vertical distribution at a resolution of 30 m is calculated as follows. The mean cloud fraction at each vertical level is calculated as the ratio of number of profiles with cloud detected at this vertical level to the total profile numbers. The cloud vertical distribution from CALIPSO at 1/3 km and 5 km are calculated first, then combined as the mean of the cloud fractions from CALIPSO 1/3 km and 5 km at each vertical level. This combined product is referred as CALIPSO 5 km, provides a complete vertical distribution of clouds from CALIPSO, and is shown in section 3. To compare, the vertical profiles of cloud fraction from CALIPSO at 1/3 km and 1 km are also combined, and shown in section 3. The combined product is referred as CALIPSO 1 km. For cloud microphysical property vertical distribution, the mean cloud phase frequency at each vertical level is calculated as the ratio of numbers of profiles with each phase to the total profile numbers. Mean cloud water content for ice (liquid) phase at each vertical level is calculated as the mean values of water content from all available ice (liquid) cloud retrievals at that level. For deriving these statistics, ice in any type of cloud (ice and mixed phase) is included, while liquid in any type of cloud (liquid and mixed phase) is included. After this step, the vertical resolution of all products is 30 m. Total cloud (ice cloud, liquid cloud, mixed phase cloud) amounts are also calculated, as the ratio of number of profiles with cloud (ice cloud, liquid cloud, mixed phase cloud) detected in any layer to the total number of profiles”.*

10. Page 4, Lines 25 onward on how the CALIPSO profiles are dealt with: again a schematic might help, as well as a clear explanation of what the horizontal and vertical resolution of these profiles are, and what it means to combine the 1/3 and 1 or 5 km products. Finally, what is the final vertical resolution of all of the products (CloudSat alone, CALIPSO alone, combined and surface)? Also why use both the 1/3 and 1km combination and the 1/3 and 5 km combination?

*Please see response to comment #4 and #9. As stated in the manuscript, it would be meaningful to see how combined 1/3 km and 1 km compares to combined 1/3 km and 5 km. The comparison of combined 1/3 km and 5 km shows more complete description, as we expected.*

*As in the response to comment #9, the vertical resolution is 30 m. We then interpolated to 100 m to be consistent with and compared to those from surface observations. These have been added in the revised manuscript.*

11. Page 5, last paragraph of section 2: are the surface products only selected when coincident with an A-train orbit? And, most importantly, are the profiles to be used in section 3 monthly means/accumulations??

*All surface profiles in a month are included in the monthly mean calculation. This is specified in the revised manuscript. “Monthly means are calculated for both surface observations and for the space-based sensors. All heights are above the mean sea level. All surface profiles in a month are accumulated for calculation of monthly means.”*

12. Figure 1, 2, 7, 9, 10 and 11: the color bar covers 0-50% but from the text cloud fractions exceed this value at low levels it seems. Why not use the full range of available values?

*Figures 1,2,7,9,10 have been updated in the revised manuscript. The color range extends to 0-80% for Barrow, and 0-60% for Eureka. We also tried extending to 0-100% for both stations, and the details in the figures were not shown as well.*

13. How is the “monthly mean total cloud amount” calculated for each instrument? (e.g. line 28, page 6)

*The following text has been added in the revised manuscript “Total cloud (ice cloud, liquid cloud, mixed phase cloud) amounts are also calculated, as the ratio of number of profiles with cloud (ice cloud, liquid cloud, mixed phase cloud) detected in any layer to the total number of profiles”.*

14. Figures 4, 5, 6 need to be redone with either thicker lines or (better) in color, to help distinguish between the different lines. It is really hard to read these as they are.

*Figure 4 and 5 have been updated with lines in color in the revised manuscript. We think the lines in Figure 6 are clear, so we did not update Figure 6.*

15. Page 7, sentence on lines 3-4: this is awkward, since you’ve already explained that the surface products were described in Shupe (2007, 2011), why not skip this first sentence and add reference to these two studies in the next sentence.

*Revised as the reviewer suggested.*

16. Page 9, line 10: “Major differences” between what? Barrow and Eureka or surface and satellite?

*This paragraph has been revised as the following, “Vertical distributions of ice cloud, liquid cloud, and mixed phase cloud at Eureka from space-based observations show similar patterns above 1 km as those from surface observations (Figure 9). The major differences between surface and space-based observations in the cloud vertical distributions at Eureka (Figure 8d, 8e, 8f, and Figure 9) are similar to those at Barrow (Figure 7, Figure 8a, 8b, and 8c). Major differences between surface and space-based observations include: much less ice and mixed phase cloud in the lowest 1 km from space-based observations; greater liquid cloud, and mixed phase cloud above 2 km in the vertical distributions and annual mean of vertical distributions from space-based observations (Figure not shown); comparable monthly mean total cloud amount, higher ice cloud monthly means, lower liquid cloud monthly means, and higher mixed phase cloud monthly means from surface observations relative to space-based observations. In additions, both satellite and surface observations reveal a key difference to the annual cycles of clouds at Eureka versus Barrow. While both sites have a similar annual cycle of ice cloud occurrence with a relative decrease in summer (Figure 8a, and 8d), there are less frequent liquid-containing clouds at Eureka with the annual maximum of these generally shifted to the autumn. These relative annual cycles explain the key differences in total cloud occurrence fraction over the annual cycle and are explained by generally colder and drier conditions in Eureka relative to Barrow (e.g., Shupe 2011).”.*

17. Section 3.2: more information is needed: what is the temporal resolution of the combined product? If monthly means, then this is a combination of the monthly means from surface and satellite? Or are these constructed for coincident observations only? Then how are the two products reconciled in term of surface time average vs satellite spatial average? Line 27: “a complete picture of the “ monthly “cloud fraction vertical distribution”?”

*The blended product is in monthly means. Line 27 has been revised as the reviewer suggested.*

18. Section 3.3: what is the take-home message for this section?

*The following text has been added in the revised manuscript, “These comparisons indicate that liquid water content monthly means from space-based and surface observations show similar annual evolution with noticeable magnitude differences. The ice water content monthly means from space and surface observations share little similarities in annual evolution or magnitude. Further investigation of these differences is warranted in order to combine these products for a complete vertical distribution of cloud water content”.*

19. Conclusions: the first “primary conclusion” is the direct consequence of the known limitations in the CloudSat (surface clutter/low sensitivity) and the CALIPSO (attenuation) instruments. References to other studies should be given. For the second “primary conclusion”, I would be inclined to conclude that surface observations perform well, regardless of cloud altitude. For the third conclusion, I would encourage the authors to discuss a bit more the implications for the annual cycle of the satellite based deficiencies. Finally, although I agree that the blended product is more accurate than surface only observations, I think that the real advantage is if one is to calculate heating rates and/or TOA/surface fluxes, this is where this product might make a difference. This should be discussed.

*All the suggestions are well received, and correspondent discussions have been added in the revised manuscript as suggested by the reviewer. In each of the primary conclusions an additional sentence or more has been added to better capture implications and context.*

20. Finally, two papers come to mind to address the very last sentence of the paper, where combined satellite products were used to evaluate cloud impacts in the Arctic in Kay et al (2008) and Kay and Gettelman (2009). The authors might want to mention these results.

*We agree. Kay et al. (2008) and Kay and Gettelman (2009) used combined satellite products. The last sentence of the manuscript suggests that we need combine surface-based and satellite products, in addition to combined satellite products. However, we appreciate the suggestions, and both references have been included in the revised manuscript.*

Typos

1. Abstract, line 24: remove “annual cycle” after “vertical distribution”
2. Line 24, page 3: “negligible surface above 0.96 km” does not make sense, is “clutter” missing?
3. Line 26, page 6: please add “to” before “penetrate” and “thick” after “optically”
4. Line 34, page 6: replace “the” before “CloudSat” with “that”.
5. Line 13, page 7: add “with” before “2B-CLDCLASS-lidar”
6. Page 8, line 21: remove “This” after “Whether”

7. Page 8. Line 23: “the” instead of “he” before “whole Arctic”

Kay, J. E., and A. Gettelman (2009), Cloud influence on and response to seasonal Arctic sea ice loss, *J. Geophys. Res.*, 114, D18204, doi:10.1029/2009JD011773

Kay, J. E., T. L’Ecuyer, A. Gettelman, G. Stephens, and C. O’Dell (2008), The contribution of cloud and radiation anomalies to the 2007 Arctic sea ice extent minimum, *Geophys. Res. Lett.*, 35, L08503, doi:10.1029/2008GL033451

*All the typos have been corrected. Both references have been added in the revised manuscript.*

The response to the reviewers' comments is in italic.

Abhay Devasthale, SMHI, Norrköping, Sweden 2017-02-06

**Review of Liu et al. doi:10.5194/acp-2016-1132, 2017**

While I am a core satellite believer, I do understand and appreciate the importance of in-situ measurements, especially in anchoring space based observations. And there is perhaps no other region in the world where we desperately need more in-situ observations than in the Arctic. Combining these two (space based and in-situ) observing systems is even better. So I really appreciate the work done by the authors in this regard. I have few issues mentioned below that I regard minor in nature, but need to be explained/elaborated. I also had an opportunity to go through the comments posted by the other reviewer and I broadly agree with her/him and I hope the authors will address them as well.

*We appreciate Dr. Devasthale's valuable comments. The manuscript becomes better with revisions in response to reviewer's comments and suggestions. We have responded to other reviewer's comments point by point, and made correspondent revisions in the revised manuscript.*

1) The authors discuss a great deal about how they compute vertical cloud fraction, but very little (or almost nothing if I haven't missed anything obvious) about the spatial (and temporal) collocation of space based and in-situ measurements. The impact of uncertainties arising from these issues is not be underestimated, especially when you compare and combined products with different spatial resolutions (even at monthly mean scale). Let's say that you (or CALIPSO team) use 15 CALIOP single shots (1/3 km each, 5x3) to generate 5 km product. What happens when this 5 km product is not centered over Barrow or Eureka and you are inconsistently selecting single shots? Have the authors evaluated few individual cases manually to check what to expect when they merge 1/3, 1 and 5 km data with reference to the station in question?

*We totally agree with the reviewer's comments, and thank for his insight. These issues, e.g. cloud frequency from surface observations v.s. spatial coverage from space-based observations, different spatial resolutions, viewing angles, vertical resolution among satellite products, all contribute to the shown differences in this manuscript. By using long-term observations, e.g. over 4 years at Eureka and over 2 years at Barrow (all data we have right now), we believe the temporal and spatial average would mitigate these issues. When longer term data from both surface-based and space-based are available, it is worth to revisit this, and see how the differences would change.*

*Inspired by the reviewer's comments, we add a paragraph in the "Conclusion" as the following, "Cloud frequency from surface is calculated in the temporal domain, while the cloud fraction from space-based observations is calculated in the spatial domain although near the surface sites. Differences in spatial resolution, viewing angles, vertical resolution, instrument sensitivity to clouds and retrieval algorithms may all contribute to the differences in the cloud vertical distributions from different instruments. Long-term averages of products may mitigate the impacts of some of these factors. Causes of the remaining differences are worth further investigation."*

2) It would be helpful if the authors also provide some physical explanation of the seasonal highs and lows in cloud fractions seen in the results. For example, in the case of Barrow, why is cloud



fraction peaking in Feb, Apr and Oct months? Why is there a minimum in Jun and Jul? This is different from Eureka. Why? Perhaps Shupe et al (2011; 2015) already discuss this, but I think the reader still needs at least a brief description of it to make full sense of the differences you observe from these two observing systems.

*A short description of the difference between Barrow and Eureka has been added to the end of Section 3.1.2. This explanation also links to a more detailed discussion of the matter in Shupe (2011). The discussion is “In additions, both satellite and surface observations reveal a key difference to the annual cycles of clouds at Eureka versus Barrow. While both sites have a similar annual cycle of ice cloud occurrence with a relative decrease in summer (Figure 8a, and 8d), there are less frequent liquid-containing clouds at Eureka with the annual maximum of these generally shifted to the autumn. These relative annual cycles explain the key differences in total cloud occurrence fraction over the annual cycle and are explained by generally colder and drier conditions in Eureka relative to Barrow (e.g., Shupe 2011)”.*

3) In the case of Barrow station, I am bit surprised at the differences in CF between 2B-GEORPFO and 2B-GEORPOF-Lidar in Aug (Fig. 2). When you add CALIOP there seems to be increase in clouds in the free troposphere from 1 to 5 km. Instinctively, I would have thought that, in the free troposphere, CALIOP would add those subvisual or super thin clouds that are missed by CPR, located in the upper troposphere lower stratosphere. Nearly 30-40% more clouds are added by GEOPROF-Lidar compared to GEOPROF in the lower and middle troposphere and it seems that even surface measurements missed these clouds. Even more confusing is the fact that CALIPSO 5 km doesn't show these clouds in Aug. So what is happening here? Part of this discrepancy can be due to the attenuation of CALIOP signal and part of it due to high amount thin clouds in the middle and lower troposphere (Devasthale et al. 2011). But it is difficult to say without further investigations.

*I agree with the reviewer's comment. The GEOPROF-Lidar has higher values than the sum of those from 2B-GEOPROF and CALIPSO 5 km in August at Barrow. The reviewer gave some possible causes, and we appreciated that and have included such discussion in the revised manuscript. However, it is still unclear why the 2B-GEOPROF-lidar has higher values than the sum of those from 2B-GEOPROF and CALIPSO 5 km. Though finding the causes is beyond the scope of this study, it is worth further investigation in future work. The following discussion has been added in the revised manuscript.*

*“It is worth pointing out that the 2B-GEOPROF-LIDAR shows higher cloud amount values from 1 km to 5 km in the troposphere than the sum of cloud amounts from 2B-GEOPROF and CALIPSO 5 km. The differences can be partially attributed to the attenuation of CALIOP signal and high amount thin clouds in the middle and lower troposphere (Devasthale et al. 2011). Though attribution investigation is beyond the scope of this study, it is worth further investigation in future studies”.*

4) The authors say that the blended cloud vertical distribution provides a complete picture. But how do we quantitatively know this? After all, we need a third independent reference to make that conclusion.

*We totally agree. A 3-D cloud distribution product would be ideal with known uncertainties. However, such a product does not exist, and probably will not be available in the near future. So, in my humble opinion, we need to work hard on getting the uncertainties of the existing products, and hopefully merging them for better quality. That is the motivation of this study.*

# Cloud vertical distribution from combined surface and space radar/lidar observations at two Arctic atmospheric observatories

Yinghui Liu<sup>1</sup>, Matthew D. Shupe<sup>2</sup>, Zhien Wang<sup>3</sup>, and Gerald Mace<sup>4</sup>

<sup>1</sup>Cooperative Institute of Meteorological Satellite Studies, University of Wisconsin at Madison, Madison, WI, USA

5 <sup>2</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado and NOAA Earth System Research Laboratory, Boulder, CO

<sup>3</sup>University of Wyoming, Laramie, Wyoming

<sup>4</sup>University of Utah, Salt Lake City, UT

10 *Correspondence to:* Yinghui Liu (yinghuil@ssec.wisc.edu)

**Abstract.** Detailed and accurate vertical distributions of cloud properties (such as cloud fraction, cloud phase, and cloud water content) and their changes are essential to accurately calculate the surface radiative flux and to depict the mean climate state. Surface- and space-based active sensors including radar and lidar are ideal to provide this information because of their superior capability to detect clouds and retrieve cloud microphysical properties. In this study, we compare the annual cycles of cloud property vertical distributions from space-based active sensors and surface-based active sensors at two Arctic atmospheric observatories, Barrow and Eureka. Based on the comparisons, we identify the sensors' respective strengths and limitations, and develop a blended cloud property vertical distribution by combining both sets of observations. Results show that surface-based observations offer a more complete cloud property vertical distribution from the surface up to 11 km above mean sea level (AMSL) with limitations in the middle and high altitudes; the annual mean total cloud fraction from space-based observations shows 25-40% fewer clouds below 0.5 km than that from surface-based observations, and space-based observations also show much less ice cloud and mixed phase cloud, and slightly greater liquid cloud, from the surface to 1 km. In general, space-based observations show comparable cloud fraction between 1 km and 2 km AMSL, and greater cloud fraction above 2 km AMSL than those from surface-based observations. The blended product combines the strength of both products to provide a more reliable annual cycle of cloud property vertical distribution from the surface to 11 km AMSL. This information can be valuable for deriving an accurate surface radiative budget in the Arctic and for cloud parameterization evaluation in weather and climate models.

## 1 Introduction

The Arctic has changed dramatically in recent decades, and causes of these changes and their feedbacks to the global climate system are under intense investigation. The Arctic is warming at a higher rate than that of the global average, a phenomenon known as Arctic amplification (Solomon et al. 2007, Serreze and Francis 2006); Arctic sea ice extent has been decreasing dramatically (Serreze et al. 2015), and this trend is expected to continue (Holland and Bitz 2003, Overland and Wang 2013).

Yinghui Liu 3/21/2017 4:50 PM

Deleted: ions

Yinghui Liu 3/31/2017 12:51 PM

Deleted: due to

Matt Shupe 3/29/2017 6:09 PM

Deleted: d

Yinghui Liu 3/31/2017 12:52 PM

Deleted: satellite

Yinghui Liu 3/31/2017 12:52 PM

Deleted: ion stations

Yinghui Liu 3/31/2017 12:52 PM

Deleted: ,

Yinghui Liu 3/31/2017 12:54 PM

Deleted: We use

Matt Shupe 3/29/2017 6:09 PM

Deleted: d

Yinghui Liu 3/31/2017 12:54 PM

Deleted: this data

Yinghui Liu 3/31/2017 12:54 PM

Deleted: to

Yinghui Liu 3/31/2017 12:54 PM

Deleted: to

Matt Shupe 3/29/2017 6:09 PM

Deleted: detailed

Matt Shupe 3/29/2017 6:09 PM

Deleted: see

Matt Shupe 3/29/2017 6:09 PM

Deleted: ; s

Yinghui Liu 3/31/2017 12:56 PM

Deleted: S

Yinghui Liu 3/31/2017 12:55 PM

Deleted:

Matt Shupe 3/29/2017 6:10 PM

Deleted: that

Yinghui Liu 3/21/2017 12:46 PM

Deleted: annual cycle

Yinghui Liu 3/31/2017 12:57 PM

Deleted: the

Yinghui Liu 3/31/2017 12:58 PM

Deleted: a

Yinghui Liu 3/31/2017 12:58 PM

Deleted: that

Changes in the Arctic have likely led to changes in the weather and climate in the midlatitudes through teleconnections in the large-scale circulation (Francis and Vavrus 2012). By studying the factors influencing the Arctic climate system and its changes, we will improve understanding of the Arctic climate and its relationship to the global climate system. The largest uncertainty in predicting the Arctic climate arises from our lack of understanding of the role clouds play in the Arctic climate system (Solomon et al. 2007, Boucher et al. 2013). A complete, accurate description of three dimensional cloud properties is critical to determine the radiation flux both at the surface and at the top of atmosphere (TOA), as well as the radiative heating rate in the atmosphere. Examining and understanding changes in these vertical distributions are key to studying the recent Arctic changes.

Cloud products from space-based combined radar/lidar observations have the potential to provide comprehensive information on the vertical distribution of cloud properties. These observations have been used to describe global cloud spatial distributions and their temporal changes (Li et al. 2015, Naud et al. 2015). However, space-based low cloud observations are limited by radar ground clutter and strong attenuation of lidar signals, especially by liquid and mixed phase clouds (Marchand et al. 2008; Blanchard et al. 2014). Radar reflectivity from CloudSat has been used to generate high vertical resolution longwave and shortwave radiative flux profiles and corresponding heating rates (L'Ecuyer et al. 2008); assessing the product's accuracy shows that CloudSat's weakness in detecting low clouds introduces the largest uncertainty. This product has been improved by the inclusion of complementary cloud and aerosol information mainly from space-based lidar observations (Henderson et al. 2013). Complementing the space-based observations, surface observations have superior performance near the surface (Shupe et al. 2011, Shupe 2011, Zhao and Wang 2010) and in resolving the diurnal cycle at a specific location, with a relatively weaker performance in the middle and upper levels.

Efforts have been made to investigate the differences in cloud fraction/frequency from surface-based and space-based radar-lidar combined observations and their impact on the radiative fluxes at multiple surface stations. Using such observations, Protat et al. (2014) studied the cloud occurrence frequency around Darwin, Australia and found that space-based observations underestimated the cloud occurrence frequency below 2 km above mean sea level (AMSL) (all heights in the text hereinafter are in km AMSL), while surface observations do not detect most of the cirrus clouds above 10 km. Blanchard et al. (2014) investigated the difference in cloud fraction and vertical distribution at Eureka, Canada in the Arctic from surface and space-based combined radar-lidar observations from 2006 to 2010. Among many valuable findings, they found that space-based radar-lidar measurements can depict a complete picture of the cloud vertical profile down to 2 km. Mioche et al. (2015) compared vertical profiles of cloud occurrences from surface lidar and space-based lidar, radar, and combined lidar and radar over the Ny-Ålesund station during March and April 2007, and showed similar results above 2 km as those in Blanchard et al. (2014). [Strength and limitations of these observations are also discussed in other papers, e.g. Kay et al. \(2007\), Kay and Gettleman \(2009\), and Huang et al. \(2012\).](#)

This study focuses on further examining and comparing the performance of space-based and surface based radar-lidar observations and retrievals to capture the vertical distribution of cloud properties, including cloud fraction, cloud phase, and cloud water content, at two Arctic atmospheric observatories, Barrow, Alaska and Eureka, Canada. Since cloud phase has

Matt Shupe 3/29/2017 6:10 PM

Deleted: -

Matt Shupe 3/29/2017 6:10 PM

Deleted: combined

Yinghui Liu 3/31/2017 1:02 PM

Deleted: and with poor spatial coverage

Yinghui Liu 3/31/2017 1:06 PM

Deleted: ion stations

been shown to have a particularly strong impact on Arctic cloud radiative effects on the surface (Shupe and Intrieri 2004), it is particularly important to understand how differences in viewing geometry impact observations of different cloud phases.

Differences between space-based and surface-based cloud (ice cloud, liquid cloud, and mixed phase cloud) amounts, and cloud ice and liquid water contents are shown in term of monthly means. Based on the comparison performed here, this

study also proposes blended products of cloud property vertical distributions from surface and space-based cloud observations at those two Arctic sites to serve as a best estimate cloud product for model and reanalysis evaluation.

Space-based radar and lidar in this paper refer to existing instruments, i.e. Cloud Profiling Radar (CPR) onboard the CloudSat and the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard the Cloud-Aerosol lidar and Infrared Pathfinder Satellite Observation (CALIPSO). However, the conclusions will likely be valid for the space-based radar and lidar instruments in the foreseeable future, i.e. the Atmospheric backscatter LIDar (ATLID), and the CPR onboard the EarthCARE mission (Hélière et al. 2007).

## 2 Data and Method

From the possible Arctic atmospheric observation sites, we have selected Barrow (71°19' N, 156°37' W) and Eureka (80°80' N, 85°57' W) because of the availability of daily cloud vertical profiles from surface observations from 2006 to 2010 when space-based observations are available. The combined radar-lidar cloud fraction best estimation, cloud fraction vertical profiles, cloud phase vertical profiles, and cloud water content vertical profiles, from surface observations at these two sites are described in detail in Shupe et al. (2011), Shupe (2011), and Shupe et al. (2015). These products are based on coincident measurements from the Ka-band cloud radar, depolarization lidars including the micropulse lidar (MPL) at Barrow and the high spectral-resolution lidar (HSRL) at Eureka, microwave radiometer, and radiosondes, which are combined to determine cloud phase (Shupe 2007) and microphysical properties at 1-min temporal and 100-m vertical resolutions.

Observations from *CloudSat* and *CALIPSO* provide an unprecedented opportunity for a spatially extensive picture of cloud cover in the Arctic (Stephens et al. 2002; Winker et al. 2003). The Vertical Feature Mask (VFM) version 3.01 from CALIPSO's CALIOP provides cloud vertical distribution in up to 10 vertical layers at 5 km and 1 km horizontal resolutions, and up to 5 vertical layers at 1/3 km horizontal resolution (Vaughan et al. 2009). The vertical resolution is 30 m below 8.2 km, and 60 m between 8.2 and 20.2 km. A Selective Iterated Boundary Location (SIBYL) scheme is applied to detect all features within a given scene. Strongly scatter features, e.g. stratus clouds, can be identified in a single laser pulse, with the 1/3 km horizontal resolution, and these features are then removed in order to detect any surrounding aerosol layers. Weakly scattering features, e.g. thin cirrus clouds, are detected with the average of several laser pulses, e.g. 5 km horizontal resolution, for higher signal-to-noise ratio. (Vaughan et al. 2005). Compared to the 1 km resolution data, the 5 km resolution product can identify weaker cloud features (Vaughan et al. 2009). Combination of the cloud layer products at 5 km and 1/3 km provides a complete vertical distribution of clouds from CALIPSO (Vaughan et al. 2009, Vaughan et al. 2005). The newly available VFM version 4.10 reports the spatial and optical properties all cloud layers detected at 5-km averaging

Matt Shupe 3/29/2017 6:11 PM

Deleted: of a year

Yinghui Liu 3/31/2017 1:51 PM

Deleted: that are coincident with

Yinghui Liu 3/22/2017 11:00 AM

Deleted: the Cloud-Aerosol lidar and Infrared Pathfinder Satellite Observation (

Yinghui Liu 3/22/2017 11:00 AM

Deleted: )

Yinghui Liu 3/22/2017 11:12 AM

Deleted: Cloud-Aerosol Lidar with Orthogonal Polarization (

Yinghui Liu 3/22/2017 11:12 AM

Deleted: )

Yinghui Liu 3/22/2017 12:19 PM

Formatted: Font:Not Italic

Matt Shupe 3/29/2017 6:12 PM

Deleted: n

Matt Shupe 3/29/2017 6:12 PM

Deleted: n

Matt Shupe 3/29/2017 6:12 PM

Deleted: .

Yinghui Liu 3/31/2017 2:02 PM

Deleted: using an iterative multi-resolution averaging scheme

Yinghui Liu 3/31/2017 2:08 PM

Formatted: Font:Not Italic

resolution, and combination of VFM at 5 km and 1/3 km is no longer needed for a complete cloud vertical distribution. In this study, the CALIPSO products version 3.01 from June 2006 to December 2010 were obtained from the Atmospheric Science Data Center at NASA Langley Research Center.

The CPR onboard CloudSat also provides echo mask, in variable "CPR\_Cloud\_mask" at 125 vertical range bins, with bin size of 240 m, in a product known as the Level 2 geometrical profiling product (2B-GEOPROF) (Marchand et al. 2008). The latest CloudSat cloud mask (R04) has negligible surface contamination from about 0.96 km above the surface. Due to the surface clutter, only strong cloud or precipitation signals can be detected in the lowest approximately 0.7 km, while weaker cloud signals are missed. In this study, a range bin is defined as cloud when the CPR\_Cloud\_mask is equal to or larger than 20, which includes weak echo, good echo, and strong echo. Very weak echo, and echo with likely surface clutter are not included. This threshold is the same as that used in the Radar-Lidar Geometrical Profile Product (2B-GEOPROF-lidar) (Mace et al. 2009, Mace et al. 2014), and a false positive detection of 5% is estimated with this threshold in the 2B-GEOPROF-lidar (Mace et al. 2009). The 2B-GEOPROF-lidar merges the CloudSat GEOPROF (Marchand et al., 2008) and the CALIPSO VFM (Vaughan et al., 2009). 2B-GEOPROF-LIDAR contains parameters for up to five hydrometeor layers, including the cloud base and top heights above mean sea level of each hydrometeor layer in one radar footprint along with the longitude and latitude.

A level 2 combined product, 2B-CLDCLASS-lidar, combines CPR and CALIOP measurements for cloud phase determination into eight basic cloud types (Sassen and Wang, 2012). Ice, water/liquid, and mixed phase clouds are identified for up to 10 layers. 2B-CLDCLASS-lidar collocates CALIOP L1 measurements to CPR footprints, then determines cloud vertical structures (Wang et al. 2008) and cloud phase. The microphysical property differences between water and ice particles, including size, location, falling speed and number concentrations, result in large differences in their radiative properties, and in turn to large differences in the CALIPSO lidar and CloudSat CPR signals. Cloud phase is effectively determined using the different sensitivities of CloudSat radar and CALIPSO lidar to ice crystals and water droplets, together with the cloud top and base temperatures.

Based on the measured CPR radar reflectivity factor, another level 2 product, the CloudSat Radar-Only Cloud Water Content Product (2B-CWC-RO), estimates cloud liquid and ice water content, and effective radius. Effective radius, and water content are retrieved with the assumption that the radar profile is due to a single phase of water, either liquid or ice. Using a simple scheme based on a model temperature profile, this product combines separate liquid and ice profiles into a mixture of ice and liquid phases over the portion of the vertical profile within the proper temperature range. The temperature profile is obtained from European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis data that have been collocated in space and time to the CloudSat radar profile and interpolated to the CloudSat vertical resolution. It should be noted that the retrieval is not designed to determine mixed-phase cloud properties directly.

In this study, vertical profiles of cloud fraction from CALIPSO at 1/3 km, 1 km and 5 km horizontal resolution, 2B-GEOPROF, and 2B-GEOPROF-LIDAR, vertical profiles of cloud phase (ice, liquid, and mixed phase) from 2B-CLDCLASS-lidar, and vertical profiles of cloud effective radius and water content from 2B-CWC-RO are calculated and

- Yinghui Liu 3/31/2017 2:13 PM  
Formatted: Font:Not Italic
- Yinghui Liu 3/31/2017 2:13 PM  
Deleted: T
- Yinghui Liu 3/22/2017 11:10 AM  
Deleted: Cloud Profiling Radar (
- Yinghui Liu 3/22/2017 11:10 AM  
Deleted: )
- Matt Shupe 3/29/2017 6:13 PM  
Deleted: s
- Matt Shupe 3/29/2017 6:13 PM  
Deleted: with
- Matt Shupe 3/29/2017 6:13 PM  
Deleted: and
- Yinghui Liu 3/23/2017 11:47 AM  
Formatted: Font:Not Italic
- Yinghui Liu 3/23/2017 11:47 AM  
Formatted: Font:Not Italic
- Yinghui Liu 3/23/2017 11:48 AM  
Deleted: Radar-Lidar Geometrical Profile Product (
- Yinghui Liu 3/23/2017 11:48 AM  
Deleted: ; Mace et al., 2014)
- Matt Shupe 3/29/2017 6:14 PM  
Deleted: y
- Matt Shupe 3/29/2017 6:14 PM  
Deleted: a
- Yinghui Liu 3/31/2017 3:45 PM  
Deleted: n ECMWF
- Yinghui Liu 3/31/2017 3:44 PM  
Deleted:
- Yinghui Liu 3/31/2017 3:45 PM  
Deleted: ECMWF
- Yinghui Liu 3/31/2017 3:46 PM  
Deleted: and

5 examined. Vertical profiles of all these products within 50 km of the two Arctic atmospheric observation sites, Barrow and Eureka, are extracted and archived. Cloud fraction vertical distribution at a resolution of 30 m is calculated as follows. The mean cloud fraction at each vertical level is calculated as the ratio of number of profiles with cloud detected at this vertical level to the total profile numbers. The cloud vertical distribution from CALIPSO at 1/3 km and 5 km are calculated first, then combined as the mean of the cloud fractions from CALIPSO 1/3 km and 5 km at each vertical level. This combined product, referred as CALIPSO 5 km, provides a complete vertical distribution of clouds from CALIPSO, and is shown in section 3. To compare, the vertical profiles of cloud fraction from CALIPSO at 1/3 km and 1 km are also combined, which is referred as CALIPSO 1km, and shown in section 3. For cloud microphysical property vertical distribution, the mean cloud phase frequency at each vertical level is calculated as the ratio of numbers of profiles with each phase to the total profile numbers. Mean cloud water content for ice (liquid) phase at each vertical level is calculated as the mean values of water content from all available ice (liquid) cloud retrievals at that level. For deriving these statistics, ice in any type of cloud (ice, and mixed phase) is included, while liquid in any type of cloud (liquid and mixed phase) is included. After this step, the vertical resolution of all products is 30 m. Total cloud (ice cloud, liquid cloud, mixed phase cloud) amounts are also calculated, as the ratio of number of profiles with cloud (ice cloud, liquid cloud, mixed phase cloud) detected in any layer to the total number of profiles.

15 Surface-based radar, lidar, and radar-lidar combined products are available from June 2006 to December 2010. Details of the collection and processing of the data can be found in Shupe (2011) and Shupe et al. (2011, 2015). Surface observations of good quality are available at Eureka for most of this time period and at Barrow from middle February 2008 to December 2010. In the text hereinafter, observations at Barrow and Eureka from 2006 to 2010 means observations at Barrow from 20 June 2006 to December 2010, and observations at Eureka from middle February 2008 to December 2010. To be consistent with the surface data, the space-based results are considered over the same time periods as surface observations available at each site. Monthly means are calculated for both surface observations and for the space-based sensors. All heights are above the mean sea level. All surface profiles in a month are accumulated for calculation of monthly means. The monthly mean sample number of the satellite sensors is a function of latitude in the Arctic, with the fewest at 60° N, gradually increasing to a maximum around 80° N (Liu 2015). Both factors are reflected in the large number of samples at Eureka, with over 6000 total samples per month from June 2006 to December 2010 at Eureka, and around 1500 total samples at Barrow per month from middle February 2008 to December 2010. The vertical resolution of the calculated space-based monthly means is interpolated to 100 m to be consistent with and compared to those from surface observations.

Yinghui Liu 3/31/2017 3:47 PM

Deleted: The cloud fraction vertical distribution from CALIPSO at 1/3 km and 5 km are calculated first, then combined as the mean of the cloud fractions from CALIPSO 1/3 km and 5 km at each vertical level. This combined product, referred as CALIPSO 5 km, provides a complete vertical distribution of clouds from CALIPSO, and is shown in section 3. To compare, the vertical profiles of cloud fraction from CALIPSO at 1/3 km and 1 km are also combined, which is referred as CALIPSO 1km, and shown in section 3. For cloud microphysical property vertical distribution, the mean cloud phase frequency at each vertical level is calculated as the ratio of numbers of profiles with each phase to the total profile numbers. Mean cloud water content for ice (liquid) phase at each vertical level is calculated as the mean values of water content from all available ice (liquid) cloud retrievals at that level. For deriving these statistics, ice in any type of cloud (ice, and mixed phase) is included, while liquid in any type of cloud (liquid and mixed phase) is included. After this step, the vertical resolution of all products is 30 m. Total cloud (ice cloud, liquid cloud, mixed phase cloud) amounts are also calculated, as the ratio of number of profiles with cloud (ice cloud, liquid cloud, mixed phase cloud) detected in any layer to the total number of profiles. [1]

Matt Shupe 3/29/2017 6:16 PM

Deleted: at each vertical level as the sum of mean cloud fraction from those from CALIPSO 1/3 km and 5 km at each vertical level. This combined product, referred as CALIPSO 5 km, provides a complete vertical distribution of clouds from CALIPSO, and is shown in section 3. To compare, the vertical profiles of cloud fraction from CALIPSO at 1/3 km and 1 km are also combined, which is referred as CALIPSO 1km, and shown in section 3. For cloud microphysical property vertical distribution, the mean cloud phase frequency at each vertical level is calculated as the ratio of numbers of profiles with each phase to the total profile numbers. Mean cloud water content for ice (liquid) phase at each vertical level is calculated as the mean values of water content from all available ice (liquid) cloud retrievals at that level. For deriving these statistics, ice in any type of cloud (ice, and mixed phase) is included, while liquid in any type of cloud (liquid and mixed phase) is included. After this step, the vertical resolution of all products is 30 m. Total cloud (ice cloud, liquid cloud, mixed phase cloud) amounts are also calculated, as the ratio of number of profiles with cloud (ice cloud, liquid cloud, mixed phase cloud) detected in any layer to the total number of profiles. [2]

Yinghui Liu 3/23/2017 10:55 AM

Deleted: The combined product is referred as CALIPSO 5 km. To compare, the vertical profiles of cloud fraction from CALIPSO at 1/3 km and 1 km are also combined, which is referred as CALIPSO 1km, and shown in section 3. For cloud microphysical property vertical distribution, the mean cloud phase frequency at each vertical level is calculated as the ratio of numbers of profiles with each phase to the total profile numbers. Mean cloud water content for ice (liquid) phase at each vertical level is calculated as the mean values of water content from all available ice (liquid) cloud retrievals at that level. For deriving these statistics, ice in any type of cloud (ice, and mixed phase) is included, while liquid in any type of cloud (liquid and mixed phase) is included. After this step, the vertical resolution of all products is 30 m. Total cloud (ice cloud, liquid cloud, mixed phase cloud) amounts are also calculated, as the ratio of number of profiles with cloud (ice cloud, liquid cloud, mixed phase cloud) detected in any layer to the total number of profiles. [3]

Matt Shupe 3/29/2017 6:17 PM

Deleted: ...and mixed phase) is included. [4]

Matt Shupe 3/29/2017 6:17 PM

Deleted: ...and Shupe et al. (2011, 2015). [5]

### 3 Results

#### 3.1 Cloud fraction vertical distribution

##### 3.1.1 Barrow

Cloud fraction vertical distributions from surface observations at Barrow (Figure 1a) reveal that cloud fractions are greater than 30% at each layer below 0.5 km throughout the year except in March and June. In the low level (surface to 2 km), the cloud fraction vertical distributions show maximum values of between 55% and 85% in October and November. In the middle level (2 km to 6 km), most of the cloud fractions are less than 30%, except local maxima greater than 30% in April and November. Minimal cloud fractions of less than 15% occur above 4 km in January, June and September. In the high level (6 km to 12 km), most cloud fractions are less than 20%, except those between 6 km and 8 km in April, August, and October.

The space-based observations show similar patterns but different values from those of surface observations at Barrow (Figure 2a, 2b, 2c). CloudSat 2B-GEOPROF (Figure 2b) shows little cloud below 0.5 km because of the surface clutter issue, limited cloud distribution between 0.5 km and 1 km, and similar patterns as surface observations above 1 km. CALIPSO 5 km (Figure 2a) shows considerably higher cloud fractions than CALIPSO 1 km (Figure not shown) throughout, and both products show some cloud fraction distribution below 0.5 km. 2B-GEOPROF-LIDAR (Figure 2c) cloud vertical distribution merges information from both CloudSat and CALIPSO, thus providing a more complete vertical distribution than either of those two alone. It is worth pointing out that the 2B-GEOPROF-LIDAR shows higher cloud amount values from 1 km to 5 km in the troposphere than the sum of cloud amounts from 2B-GEOPROF and CALIPSO 5 km. The differences can be partially attributed to the attenuation of CALIOP signal and high amount thin clouds in the middle and lower troposphere (Devasthale et al. 2011). Though attribution investigation is beyond the scope of this study, it is worth further investigation in future studies. Based on the 2B-GEOPROF-LIDAR cloud vertical distribution, the cloud fraction below 0.5 km is less than 30% most of the year, except in May, and November when the local maximum is greater than 30%. In the low level, cloud fraction increases with height, reaches a maximum between 1 km and 1.5 km, and starts to decrease in general. The annual minimum cloud fraction at this level of less than 20% appears in June and July. In the middle level, cloud fractions are mostly between 20% and 40%. The maximum cloud fraction appears in April, August, and December with values greater than 35%. The minimum appears in March and June with values less than 16%. In the high level, cloud fraction of 20% or greater appears most of the time except for November, March and June.

Comparison of cloud vertical distributions from space-based observations and surface observations at Barrow show overall least cloud fraction from CALIPSO 1 km, then CALIPSO 5 km, and 2B-GEOPROF, and overall most cloud fraction from 2B-GEOPROF-LIDAR above 1 km, and all space-based cloud fractions are less than that from surface observations in the lowest 1 km (Figure 2, Figure 3). Compared to cloud fraction vertical distribution from surface observations, CALIPSO 1 km shows less cloud fraction in every month from surface to 6-11 km depending on month (Figure not shown); CALIPSO 5 km shows less cloud fraction from surface to 5 km in every month, and greater cloud fraction above 6 km in most months;

Yinghui Liu 3/31/2017 4:44 PM  
Formatted: Font:Not Italic

Above 1 km 2B-GEOPROF has differences from the surface observations of +20 to -10%. In most months 2B-GEOPROF-LIDAR tends to have greater cloud fractions above 1 km; all space-based cloud fractions show lower cloud fractions below 1 km, with the least from 2B-GEOPROF, then CALIPSO 1 km, CALIPSO 5 km and 2B-GEOPROF-LIDAR. The near surface cloud distributions from 2B-GEOPROF-LIDAR originate from CALIPSO observations and also show much less cloud fraction distributions below 0.5 km, with differences as high as -67% in October. The difference becomes smaller between 0.6 km and 1.2 km. Above 1.2 km, 2B-GEOPROF-LIDAR shows generally greater cloud fractions (up to 27% in September at 5 km) than those from surface observations.

Comparison of the annual mean cloud vertical distributions from space-based observations and surface observations shows that all space-based observations have lower cloud fractions in the lowest 1 km, while 2B-GEOPROF-LIDAR and CALIPSO 5 km have higher cloud fractions at some heights above 1 km (Figure 4a). More specifically, compared to surface observations, below 0.5 km the space-based observations see 25-40% fewer clouds than are observed from the surface; between 1 km and 6 km 2B-GEOPROF and 2B-GEOPROF-LIDAR show slightly greater cloud fraction, while CALIPSO 1 km and 5 km show less cloud fraction; above 6 km, CALIPSO 5 km and 2B-GEOPROF-LIDAR show slightly greater cloud fraction, while CALIPSO 1 km and 2B-GEOPROF show less cloud fraction. For 2B-GEOPROF-LIDAR, the greater cloud fractions above 1 km are due to the combined detection capabilities from CALIPSO 5 km and 2B-GEOPROF. The low cloud fraction from space observations below 1 km can be attributed to surface clutter issue from 2B-GEOPROF near the surface, and the inability of CALIPSO to penetrate optically thick clouds. Surface reported lower cloud fractions above 1 km might be due to the inability of surface lidar to penetrate lower-level optically thick liquid and mixed-phase clouds, along with the difficulty to detect optically thin clouds composed of small ice particles in the middle and upper levels by surface radar.

The annual cycle of monthly mean total cloud amount at Barrow shows relatively low values from January to March, and relatively high values (75% and higher) from April to December (Figure 5a). Monthly means from space observations and surface observations share similarities except 2B-GEOPROF shows much lower fractions in all months, e.g. around 30% in June compared to above 75% from surface observations. The 2B-GEOPROF-LIDAR has the most similar annual cycle to surface observations, with lower monthly means from CALIPSO 5 km, followed by CALIPSO 1 km, and with 2B-GEOPROF showing the lowest values and the largest negative differences from May to September. This is in agreement with results presented in Zygmuntowska et al. (2012) considering that CloudSat does not detect the cloud below approximately 0.5 km. The larger differences from May to September might be attributed to the relatively higher frequency of clouds below 960 m in that time period (Figure 6), which CloudSat does not detect well.

Vertical distributions of ice cloud, liquid cloud, and mixed phase cloud at Barrow from 2006 to 2010 from surface observations are shown in Figure 7 (Shupe 2007, 2011). The main features include the following: Ice clouds are prevalent from the surface up to 9-11 km throughout the year except from the surface to 4.5 km in June, July, and August. The maximum ice cloud fractions occur at low levels from October to April, and at middle levels in April, November, and December with a range between 10% and 30%. At high levels, ice cloud fraction between 10% and 20% appears from June to August. Mixed phase clouds generally occur on average 8-20% at low levels, and on average 2-8% at middle levels. The

Matt Shupe 3/29/2017 6:19 PM

Deleted: by surface radar

Matt Shupe 3/29/2017 6:20 PM

Deleted: A

Matt Shupe 3/29/2017 6:20 PM

Deleted: those

Yinghui Liu 3/21/2017 12:54 PM

Deleted: e

Matt Shupe 3/29/2017 6:20 PM

Deleted: near the surface

Matt Shupe 3/29/2017 6:20 PM

Deleted: Cloud v

Yinghui Liu 3/23/2017 10:35 AM

Deleted: Shupe (2011) presents cloud phase vertical distributions at Barrow and Eureka based on combined surface lidar, radar, microwave radiometer, and radiosonde observations (Shupe 2007). The main features of cloud vertical distributions of ice cloud, liquid cloud, and mixed phase cloud at Barrow from 2006 to 2010 (Figure 7) include the following: Ice clouds are prevalent from the surface up to 9-11 km throughout the year except from the surface to 4.5 km in June, July, and August.



maximum mixed phase cloud fractions, up to 57%, appear between the surface and 1 km from September to November. Liquid clouds appear between the surface and 0.8 km in the warm season mainly from May to September, with a maximum liquid cloud fraction of greater than 40% in the lowest 0.4 km in August.

Cloud phase vertical distributions at Barrow derived with 2B-CLDCLASS-lidar agree in general with the patterns observed above 1 km from surface observations (Figure 7). At Barrow, ice clouds are common throughout the year from 1 km up to 11 km except from the surface to 4.5 km from June to August, when the ice cloud fraction are mostly less than 7%. Liquid cloud fraction of greater than 10% appears mainly from the surface to 0.8 km in May, August, September, and November. Mixed phase clouds appear between 1 km and 3.5 km throughout the year. A maximum of up to 55% appears at 1 km in October. Another local maximum between 15% and 30% extends from 1 km to 6 km in August, which is not shown in the surface observations. There is little mixed phase cloud distribution below 1 km.

One major difference between the vertical distributions of ice, liquid, and mixed phase clouds from space-based and surface observations is that the space-based observations show much less ice cloud and mixed phase cloud, and slightly greater liquid cloud from the surface to 1 km (Figure 7). Above 1 km, the two perspectives show similar annual average profiles, with the space observations seeing slightly higher mixed-phase cloud fractions from 3-5km, slightly higher liquid cloud fraction from 0.5-3 km, higher ice cloud fraction at 10 km, and lower ice cloud fractions at 2-6 km (Figure not shown), although month to month variability can be larger (Figure 7).

Annual cycle of monthly mean ice clouds from surface show greater values throughout the year except January (Figure 8a), similar to the mixed phase cloud amount comparison (Figure 8c). Liquid cloud monthly means from 2B-CLDCLASS-lidar show greater values than those from surface observations in all months except January, June, and July (Figure 8b). Some of the low-level differences may be the result of space-based measurements having difficulties detecting mixed phase clouds with low ice concentration, thus classifying these as liquid.

### 3.1.2 Eureka

All cloud distributions at Eureka show different annual cycles from those at Barrow. Cloud vertical distributions from space-based observations at Eureka are relatively smoother than those from space-based observations at Barrow partly due to greater sample numbers at Eureka. However, general findings about the differences between space-based and surface observations are similar.

Total cloud fraction vertical distribution at Eureka (Figure 1b) from surface observations shows the largest values (up to 55%) between the surface and 0.5 km, except from June to August when low-level values are less than 25% and profile maximum values are above 1 km. The maximum cloud fraction at low levels at Eureka is considerably lower than that at Barrow. At middle levels, the cloud fractions are mainly 10-30% with a local maximum greater than 30% from September to November. At high levels, most of the cloud fractions are less than 20%.

For the vertical distributions of total cloud fraction from space (Figure 2d, 2e, 2f), CALIPSO 5 km (Figure 2d) and 1 km (Figure not shown) show similar patterns with greater values in CALIPSO 5 km. Both show limited cloud below 0.5 km. A

Matt Shupe 3/29/2017 6:21 PM

Deleted: 9a

Matt Shupe 3/29/2017 6:21 PM

Deleted: 9c

Matt Shupe 3/29/2017 6:21 PM

Deleted: 9b

Matt Shupe 3/29/2017 6:22 PM

Deleted: because

Matt Shupe 3/29/2017 6:22 PM

Deleted: e

Matt Shupe 3/29/2017 6:22 PM

Deleted: to

local maximum between 4 km and 6 km appears from October to February in CALIPSO 5 km. 2B-GEOPROF (Figure 2e) shows little cloud below 1 km, and detailed cloud information above 1 km, with maximum fractions between 1 km and 4 km from September to December. 2B-GEOPROF-LIDAR (Figure 2f) merges information from CALIPSO and CloudSat, and presents a comparable cloud vertical distribution to that from surface observations, except near the surface. At low levels, the 2B-GEOPROF-LIDAR cloud fractions are less than 40%, with maximum between 30% and 40% from September to November. At middle levels, a local maximum cloud fraction of between 30% and 35% appears between 2 km and 4 km from September to November; a local minimum cloud fraction of less than 15% appears in March. At high levels, cloud fraction is above 20% from July to November between 6 km and 7.5 km.

Though the total cloud fraction vertical distributions and their annual means at Eureka and Barrow are different (Figure 1, Figure 4), comparison of the space-based cloud vertical distributions and their annual means and those from the surface at Eureka (Figure 3d, 3e, 3f, and Figure 4b) shows qualitatively the same differences as those at Barrow (Figure 3a, 3b, 3c, and Figure 4a). Whether the differences found in cloud detection capabilities from CALIPSO 1 km, CALIPSO 5 km, 2B-GEOPROF, and 2B-GEOPROF-LIDAR regarding total cloud fraction vertical distributions based on comparisons with surface observations at Barrow and Eureka can be generalized to the whole Arctic might be worth further investigation.

The annual cycle of monthly mean cloud amount at Eureka from surface observations shows relatively low values of between 56% and 67% from February to August, and high values of between 67% and 81% from September to February (Figure 5b). Monthly means from space-based observations show general increasing cloud amount from March to September, and then start to decrease gradually. 2B-GEOPROF-LIDAR shows comparable monthly means as CALIPSO 5 km, and both are greater than those from CALIPSO 1 km and 2B-GEOPROF, with the least typically from 2B-GEOPROF.

All space-based monthly means are noticeably smaller from January to March than those from surface observations, and these negative differences might be due to the relatively higher frequency of clouds below 960 m only. Monthly means from 2B-GEOPROF-LIDAR and CALIPSO 5 km are greater from June to August compared to surface observations, which is possibly due to the higher frequency of clouds above 960 m only, which surface observations might miss (Figure 6b).

For surface observations at Eureka, ice clouds are the prevalent cloud type from the surface to up to 11 km throughout the year except in June, July, and August when there are few ice clouds from the surface to 3 km (Figure 9). The maximum ice cloud fraction of up to 40% appears at low levels from November to March. At middle levels, ice cloud fractions are mostly between 15% and 25%, with the exception of lower fractions from June to August. At high levels, ice cloud fractions are mostly below 10% except from July to October. Mixed phase clouds are common at low levels except in July and August, and at middle levels from June to September. A maximum mixed phase cloud fraction between 20% and 30% appears between the surface and 2 km from September to October. Liquid phase clouds are mainly less than 5% throughout the year except in lowest 0.5 km in September and October.

Vertical distributions of ice cloud, liquid cloud, and mixed phase cloud at Eureka from space-based observations show similar patterns above 1 km as those from surface observations (Figure 9). The major differences between surface and space-based observations in the cloud vertical distributions at Eureka (Figure 8d, 8e, 8f, and Figure 9) are similar to those at

Yinghui Liu 3/21/2017 12:55 PM  
Deleted: This

Matt Shupe 3/29/2017 6:22 PM  
Deleted: A

Matt Shupe 3/29/2017 6:22 PM  
Deleted: Cloud v

Yinghui Liu 3/23/2017 2:29 PM  
Deleted: for

Barrow (Figure 7, Figure 8a, 8b, and 8c). Major differences between surface and space-based observations include: much less ice and mixed phase cloud in the lowest 1 km from space-based observations; greater liquid cloud, and mixed phase cloud above 2 km in the vertical distributions and annual mean of vertical distributions from space-based observations (Figure not shown); comparable monthly mean total cloud amount, higher ice cloud monthly means, lower liquid cloud monthly means, and higher mixed phase cloud monthly means from surface observations relative to space-based observations. In additions, both satellite and surface observations reveal a key difference to the annual cycles of clouds at Eureka versus Barrow. While both sites have a similar annual cycle of ice cloud occurrence with a relative decrease in summer (Figure 8a, and 8d), there are less frequent liquid-containing clouds at Eureka with the annual maximum of these generally shifted to the autumn. These relative annual cycles explain the key differences in total cloud occurrence fraction over the annual cycle and are explained by generally colder and drier conditions in Eureka relative to Barrow (e.g., Shupe 2011).

### 3.2 Blended cloud vertical distribution at Barrow and Eureka

While the cloud fraction vertical distributions at Barrow and Eureka show different patterns, the cloud vertical distribution differences between space-based and surface observations are similar for both stations as detailed in Section 3.1. Surface observations show detailed and higher values in the lowest 1 km; space observations provide little cloud information in the lowest 0.5 km, limited information between 0.5 km and 1 km, and comparable or higher values between 1 km and 2 km. In the middle and upper levels, space observations generally show higher values.

Low-level cloud is ubiquitous in the Arctic. For a complete picture of cloud vertical distribution in the Arctic, clouds in the lowest 1 km above AMSL need be included, and such information is better depicted in the surface observations. Here we generate a blended monthly mean cloud fraction vertical distribution for total cloud, ice cloud, liquid cloud, and mixed phase cloud from both surface and space-based observations in monthly means. The monthly mean cloud fraction at every level in the blended product is given as the larger monthly mean cloud fraction of the surface and space-based observations. With this approach, the blended products provide a complete cloud fraction vertical distribution in term of monthly means by drawing strength from surface and space-based products.

Figure 10 presents the blended total cloud fraction vertical distributions from 2B-GEOPROF-LIDAR and surface observations at Barrow and Eureka from 2006 to 2010. The blended product provides a complete picture of the monthly cloud fraction vertical distribution, by blending surface and space-based observations. There is no apparent discontinuity in the cloud fraction vertical distribution near the surface at Barrow or Eureka. Figure 11 shows cloud vertical distributions of ice cloud, liquid cloud, and mixed phase cloud from 2B-CLDCLASS-lidar and surface observations at Barrow and Eureka from 2006 to 2010. The blended cloud phase vertical distributions from space-based observations show similar patterns as those from surface observations with more complete distributions in the middle and high levels. The blended product is smoother for Eureka than for Barrow. The cloud fraction vertical distributions are smooth for all cloud phases.

Matt Shupe 3/29/2017 6:25 PM  
Deleted: than

Yinghui Liu 3/30/2017 7:13 PM  
Deleted: Below 1 km, the cloud fraction vertical distribution is from surface observations; from 1 km to 2 km, the cloud fraction vertical distribution is calculated as a linear combination of surface observation and space-based observation, with the weights of the surface observations changing from 1.0 to 0.0 linearly as a function of altitude, and the weights of the space-based observations changing from 0.0 to 1.0 linearly as a function of altitude; above 2 km, the cloud fraction vertical distribution is from space-based observations.

Yinghui Liu 3/30/2017 7:23 PM  
Deleted: optimally combining

Yinghui Liu 3/30/2017 7:23 PM  
Deleted: between 1 km and 2 km

Matt Shupe 3/29/2017 6:26 PM  
Deleted: total cloud,

Matt Shupe 3/29/2017 6:26 PM  
Deleted: respectively

Yinghui Liu 3/30/2017 7:25 PM  
Deleted: transitions between main surface observations below 1km and main space-based observations above 2 km

### 3.3 Cloud water content

[In addition to fractional occurrence of clouds by phase it is also instructive to examine space and surface-based retrievals of cloud water contents.](#) The ice water content and liquid water content vertical distributions from 2B-CWC-RO and surface observations at Barrow are presented in Figure 12. There is limited information below 1 km from space-based observations.

5 Based on the space-based observations, the ice water content is less than  $40 \text{ mg m}^{-3}$  throughout the year except higher values of up to  $100 \text{ mg m}^{-3}$  from May to August, and in December from 2 km to 6 km; the liquid water content [hass](#), high values of between  $150 \text{ mg m}^{-3}$  and  $300 \text{ mg m}^{-3}$  from June to August from 1 km to 3.5 km, and in February, September and October between 1 km and 2 km. Surface observations show low ice water content of  $20 \text{ mg m}^{-3}$  and less above 4 km, and higher values below 4 km, with maximum values of  $60\text{-}100 \text{ mg m}^{-3}$  from October to February in the lowest 2 km, and in June and

10 July between 1 km and 3 km. [Surface-based](#) liquid water content shows high values of  $150\text{-}250 \text{ mg m}^{-3}$  from May to August from the surface to 5 km, and in September and October from surface to 2 km. The similarity between the ice water content from surface and space-based observations includes that both distributions tend to have higher values in June and July, and from December to February, but at different heights. For liquid water content, both surface and space-based observations show high values from June to August in the lowest 3.5 km, and in September and October below 2 km.

15 In Eureka, the ice water content from space-based observations is less than  $40 \text{ mg m}^{-3}$  throughout the year except values of around  $60 \text{ mg m}^{-3}$  from August to October from 2 km to 5 km, and in April from 2 km to 6 km as shown in Figure 13. The ice water content from surface observations is also below  $40 \text{ mg m}^{-3}$  throughout the year except values of between 60 and  $80 \text{ mg m}^{-3}$  from June to October from the surface to 3 km. Liquid water content from both surface and space-based observations shows low values of  $75 \text{ mg m}^{-3}$  and less from October to April, and high values from June to August below 3 km, with much

20 higher values from space-based observations.

[These comparisons indicate that liquid water content monthly means from space-based and surface observations show similar annual evolution with noticeable magnitude differences. The ice water content monthly means from space and surface observations share little similarities in annual evolution or magnitude. Further investigation of these differences is warranted in order to combine these products for a complete vertical distribution of cloud water content.](#)

### 25 4 Conclusions

This study compares the annual cycles of cloud vertical distributions of total cloud, ice cloud, liquid cloud, and mixed phase cloud [occurrence fractions](#) from combined surface active lidar/radar observations and from multiple space-based active lidar/radar products at two Arctic atmospheric observation stations, Barrow and Eureka. The primary conclusions are as follows:

- 30 • All space-based active radar/lidar cloud observations have limitations in the lowest 1 km AMSL; the surface measurements have superior performance near the surface, and [thereby](#) complement the space-based observations. Surface observations show that the highest total cloud fractions of all cloud, ice cloud, liquid cloud, and mixed phase

Matt Shupe 3/29/2017 6:28 PM  
Deleted: have

Matt Shupe 3/29/2017 6:29 PM  
Deleted: ;

Matt Shupe 3/29/2017 6:29 PM  
Deleted: s

cloud appear between surface and 1 km. All space-based observations show lower total cloud fractions below 1 km, with the least from 2B-GEOPROF, then CALIPSO 1 km, CALIPSO 5 km, and 2B-GEOPROF-LIDAR. The annual mean total cloud fractions from space-based observations show 25-40% fewer clouds below 0.5 km than those from surface-based observations. Compared to surface-based observations, space-based observations show much less ice cloud and mixed phase cloud, and slightly greater liquid cloud from the surface to 1 km. These results are generally consistent with conclusions from previous studies (Protat et al. 2014, Blanchard et al. 2014, Mioche et al. 2015).

• Surface observations perform well in describing the cloud vertical distribution at these observation sites. Above 1 km, space-based observations show similar patterns as surface observations, but different magnitudes for total cloud, ice cloud, liquid cloud, and mixed phase cloud. For satellite-based total cloud fraction, CALIPSO 1 km shows the lowest values, with higher values from CALIPSO 5 km especially above 6 km, and the highest values from 2B-GEOPROF mainly in the middle level. 2B-GEOPROF-LIDAR, which merges CALIPSO and CloudSat, provides the closest vertical distribution to that from surface observations. While the surface observations generally show cloud fractions that are comparable to, or higher than, the satellite-based fractions at most heights, the space observations show greater ice cloud fraction above 9 km, greater liquid cloud fraction in general, and greater mixed phase cloud fraction above 1km.

• For the annual cycle of the total cloud fraction, monthly means from space-based observations are generally lower than those from surface observations. Each perspective has its limitations, with the surface observations missing some high-level cloud and the space-based sensors missing many low-level clouds. Both estimates are likely lower than the true cloud fraction, if those missed clouds are not all overlapping with other clouds. Because low clouds are more prevalent at these locations, the surface-based estimate is likely closer to the true total cloud fraction. Annual cycles of monthly mean cloud occurrence by phase show less ice and mixed phase cloud, and greater liquid cloud from space-based observations. This result suggests that active sensor satellite-based estimates of cloud fraction across the Arctic are likely lower than the true cloud fraction, particularly at low levels and at times of year when low clouds are frequent.

• A blended cloud fraction vertical distribution using the larger value of surface and space-based observations can provide a more complete description of cloud vertical distribution of total cloud, and ice, liquid, and mixed phase clouds from the surface to 11 km. Such a blended product would be important when considering net atmospheric heating rates above these sites. Such approach can be also useful in the tropics for a complete depiction of cloud fraction vertical distribution.

Existing space-based cloud distributions in the lowest 1 km do not capture all clouds, especially ice and mixed phase clouds. How these missed clouds in the lowest 1 km affects the radiation flux calculations at the surface and at the top of the atmosphere is a topic of future work and may impact past studies that examine Arctic surface radiative fluxes as suggested by L'Ecuyer et al. (2008). The blended cloud property vertical distribution can be used as an input to a Monte Carlo radiative transfer model for a more accurate surface radiation flux calculation at these sites. A blended cloud property vertical distribution can also be used to evaluate cloud parameterizations in both weather and climate models (Klaus et al. 2016), to

Matt Shupe 3/29/2017 6:30 PM

**Deleted:** see

Matt Shupe 3/29/2017 6:31 PM

**Deleted:** that

Matt Shupe 3/29/2017 6:31 PM

**Deleted:** This is

Matt Shupe 3/29/2017 6:32 PM

**Deleted:** followed by

Matt Shupe 3/29/2017 6:32 PM

**Deleted:** magnitude

Matt Shupe 3/29/2017 6:32 PM

**Deleted:** as

Matt Shupe 3/29/2017 6:33 PM

**Deleted:** T

Matt Shupe 3/29/2017 6:33 PM

**Deleted:** some

Yinghui Liu 3/23/2017 3:14 PM

**Deleted:**

Matt Shupe 3/29/2017 6:36 PM

**Deleted:** The lack of capability of space-based observations in detecting the low-level cloud would lead to uncertainties in the radiation flux estimations at the surface and at the top of atmosphere, as suggested in L'Ecuyer et al. (2008)

Yinghui Liu 3/30/2017 7:26 PM

**Deleted:** with a weighted combination

Matt Shupe 3/29/2017 6:35 PM

**Deleted:** rather

Matt Shupe 3/29/2017 6:35 PM

**Deleted:** Improvement in radiation flux estimations at the surface and at the top of atmosphere can be achieved with incorporation of such blended products

Yinghui Liu 3/23/2017 3:44 PM

**Deleted:**

Matt Shupe 3/29/2017 6:37 PM

**Deleted:** for improvement

study, Arctic atmosphere-sea ice-ocean interactions (Kay et al. 2008, Kay and Gettleman 2009, Taylor et al. 2015, Liu et al. 2012a), and in other Arctic cloud studies (Devasthale et al. 2011, Liu et al. 2012b, Liu and Key 2016).

Low-level clouds are frequent in the Arctic and important for the surface radiation balance. While space-based cloud observations from active radar/lidar sensors have been critical for improving our understanding of Arctic clouds and their interactions with other climate components in the Arctic, challenges remain in depicting Arctic low-level clouds from space. Surface observations of clouds at existing Arctic atmospheric observatories and a few field campaigns have provided valuable information on Arctic clouds, especially for studying low-level clouds (Tjernström et al. 2014, Uttal et al. 2002). However, such observations are limited in spatial extent and may not represent pan-Arctic cloudiness. Thus it is critical to combine key information from both space-based- and surface cloud measurements to provide the most comprehensive characterization of Arctic clouds possible and to facilitate further understanding of the Arctic climate system.

Cloud frequency from surface is calculated in the temporal domain, while the cloud fraction from space-based observations is calculated in the spatial domain although near the surface sites. Differences in spatial resolution, viewing angles, vertical resolution, instrument sensitivity to clouds, and retrieval algorithms may all contribute to the differences in the cloud vertical distributions from different instruments. Long-term averages of products may mitigate the impacts of some of these factors. Causes of the remaining differences are worth further investigation.

### Acknowledgements

Shupe acknowledges support from the US Department of Energy (DOE) Atmospheric System Research Program (DE-SC0011918) and the National Science Foundation (ARC-0632187). The authors thank Norm Wood, and Ralph Kuehn for their valuable comments on the paper. Ground-based observations from Barrow were obtained from the DOE Atmospheric Radiation Measurement Program. Ground-based observations at Eureka were obtained from the NOAA Earth System Research Laboratory and the Canadian Network for the Detection of Arctic Change (CANDAC). The CALIPSO products from June 2006 to December 2010 were obtained from the Atmospheric Science Data Center at NASA Langley Research Center. The 2B-GEOPROF, 2B-GEOPROF-LIDAR, 2B-CLDCLASS-lidar, and 2B-CWC-RO products from June 2006 to December 2010 were obtained from the CloudSat Data Processing Center at the Colorado State University.

### References

Blanchard, Y., Pelon, J., Eloranta, E. W., Moran, K. P., Delanoë, J., and Sèze, G.: A synergistic analysis of cloud cover and vertical distribution from A-Train and ground-based sensors over the high Arctic station EUREKA from 2006 to 2010, *Journal of Applied Meteorology and Climatology*, 53, 2553-2570, 2014.

Boucher, O., and Coauthors: Clouds and aerosols. *Climate Change 2013: The Physical Science Basis*, T. F. Stocker et al., Eds., Cambridge University Press, 571–657, doi:10.1017/CBO9781107415324.016, 2013.

Matt Shupe 3/29/2017 6:37 PM

Deleted: the

Yinghui Liu 3/23/2017 3:26 PM

Deleted: 9

Yinghui Liu 3/23/2017 4:18 PM

Deleted: 2

Matt Shupe 3/29/2017 6:37 PM

Deleted: improvement of

- [Devasthale, A., Tjernstrom, M., Karlsson, K.-G., Thomas, M. A., Jones, C., Sedlar, J., and Omar, A. H.: The vertical distribution of thin features over the Arctic analysed from CALIPSO observations, \*Tellus Series B-Chemical and Physical Meteorology\*, 63, 77-85, 10.1111/j.1600-0889.2010.00516.x, 2011.](#)
- 5 Devasthale, A., Tjernström, M., Caian, M., Thomas, M. A., Kahn, B. H., and Fetzer, E. J.: Influence of the Arctic Oscillation on the vertical distribution of clouds as observed by the A-Train constellation of satellites, *Atmospheric Chemistry and Physics*, 12, 10535-10544, 2012.
- Francis, J. A., and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes, *Geophysical Research Letters*, 39, 2012.
- Fu-Lung, C., and Zhanqing, L.: Estimating the vertical variation of cloud droplet effective radius using multispectral near-infrared satellite measurements, *Journal of Geophysical Research*, 107, AAC7-1-12, 2002.
- 10 Henderson, D. S., L'Ecuyer, T., Stephens, G., Partain, P., and Sekiguchi, M.: A multisensor perspective on the radiative impacts of clouds and aerosols, *Journal of Applied Meteorology and Climatology*, 52, 853-871, 2013.
- Holland, M. M., and Bitz, C. M.: Polar amplification of climate change in coupled models, *Climate Dynamics*, 21, 221-232, 10.1007/s00382-003-0332-6, 2003.
- 15 [Huang, Y., Siems, S. T., Manton, M. J., Hande, L. B., and Haynes, J. M.: The structure of low-altitude clouds over the Southern Ocean as seen by CloudSat, \*Journal of Climate\*, 25, 2535-2546, 2012.](#)
- [Hélière, A., Lefebvre, A., Wehr, T., Bézy, J.-L., and Durand, Y.: The EarthCARE mission: mission concept and lidar instrument pre-development, 4975-4978.](#)
- 20 [Kay, J. E., L'Ecuyer, T., Gettelman, A., Stephens, G., and O'Dell, C.: The contribution of cloud and radiation anomalies to the 2007 Arctic sea ice extent minimum, \*Geophysical Research Letters\*, 35, 10.1029/2008gl033451, 2008.](#)
- Kay, J. E., and Gettelman, A.: Cloud influence on and response to seasonal Arctic sea ice loss, *Journal of Geophysical Research-Atmospheres*, 114, 10.1029/2009jd011773, 2009.
- Klaus, D., Dethloff, K., Dorn, W., Rinke, A., and Wu, D. L.: New insight of Arctic cloud parameterization from regional climate model simulations, satellite-based, and drifting station data, *Geophysical Research Letters*, 2016.
- 25 L'Ecuyer, T. S., Wood, N. B., Haladay, T., Stephens, G. L., and Stackhouse, P. W.: Impact of clouds on atmospheric heating based on the R04 CloudSat fluxes and heating rates data set, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Li, J., Huang, J., Stamnes, K., Wang, T., Lv, Q., and Jin, H.: A global survey of cloud overlap based on CALIPSO and CloudSat measurements, *Atmospheric Chemistry and Physics*, 15, 519-536, 2015.
- Liu, Y., Key, J., Liu, Z., Wang, X., and Vavrus, S.: A cloudier Arctic expected with diminishing sea ice, *Geophysical Research Letters*, 39, 10.1029/2012GL051251, 2012a.
- 30 Liu, Y., Key, J. R., Ackerman, S. A., Mace, G. G., and Zhang, Q.: Arctic cloud macrophysical characteristics from CloudSat and CALIPSO, *Remote Sensing of Environment*, 124, 159-173, 10.1016/j.rse.2012.05.006, 2012b.

- Liu, Y.: Estimating errors in cloud amount and cloud optical thickness due to limited spatial sampling using a satellite imager as a proxy for nadir-view sensors, *Journal of Geophysical Research: Atmospheres*, 120, 6980-6991, 2015.
- [Liu, Y., and Key, J. R.: Assessment of Arctic cloud cover anomalies in atmospheric reanalysis products using satellite data, \*Journal of Climate\*, 29, 6065-6083, 2016.](#)
- 5 [Mace, G. G., Zhang, Q., Vaughan, M., Marchand, R., Stephens, G., Trepte, C., and Winker, D.: A description of hydrometeor layer occurrence statistics derived from the first year of merged Cloudsat and CALIPSO data, \*Journal of Geophysical Research: Atmospheres\*, 114, 2009.](#)
- Mace, G. G., and Zhang, Q.: The CloudSat radar-lidar geometrical profile product (RL-GeoProf): Updates, improvements, and selected results, *Journal of Geophysical Research-Atmospheres*, 119, 9441-9462, 10.1002/2013jd021374, 2014.
- 10 Marchand, R., Mace, G. G., Ackerman, T., and Stephens, G.: Hydrometeor detection using Cloudsat - An earth-orbiting 94-GHz cloud radar, *Journal of Atmospheric and Oceanic Technology*, 25, 519-533, 10.1175/2007jtecha1006.1, 2008.
- Mioche, G., Jourdan, O., Ceccaldi, M., and Delanoë, J.: Variability of mixed-phase clouds in the Arctic with a focus on the Svalbard region: a study based on spaceborne active remote sensing, *Atmospheric Chemistry and Physics*, 15, 2445-2461,
- 15 2015.
- Naud, C. M., Posselt, D. J., and van den Heever, S. C.: A CloudSat-CALIPSO View of Cloud and Precipitation Properties across Cold Fronts over the Global Oceans, *Journal of Climate*, 28, 6743-6762, 2015.
- Overland, J. E., and Wang, M.: When will the summer Arctic be nearly sea ice free?, *Geophysical Research Letters*, 40, 2097-2101, 2013.
- 20 Protat, A., Young, S. A., McFarlane, S. A., L'Ecuyer, T., Mace, G. G., Comstock, J. M., Long, C. N., Berry, E., and Delanoë, J.: Reconciling Ground-Based and Space-Based Estimates of the Frequency of Occurrence and Radiative Effect of Clouds around Darwin, Australia, *Journal of Applied Meteorology and Climatology*, 53, 456-478, 10.1175/jamc-d-13-072.1, 2014a.
- Protat, A., Young, S. A., McFarlane, S. A., L'Ecuyer, T., Mace, G. G., Comstock, J. M., Long, C. N., Berry, E., and Delanoë, J.: Reconciling ground-based and space-based estimates of the frequency of occurrence and radiative effect of
- 25 clouds around Darwin, Australia, *Journal of Applied Meteorology and Climatology*, 53, 456-478, 2014b.
- Rossow, W. B., Zhang, Y. C., and Wang, J. H.: A statistical model of cloud vertical structure based on reconciling cloud layer amounts inferred from satellites and radiosonde humidity profiles, *Journal of Climate*, 18, 3587-3605, 10.1175/jcli3479.1, 2005.
- Sassen, K., and Wang, Z.: The clouds of the middle troposphere: composition, radiative impact, and global distribution, *Surveys in geophysics*, 33, 677-691, 2012.
- 30 Serreze, M. C., and Francis, J. A.: The arctic amplification debate, *Climatic Change*, 76, 241-264, 10.1007/s10584-005-9017-y, 2006.
- Serreze, M. C., and Stroeve, J.: Arctic sea ice trends, variability and implications for seasonal ice forecasting, *Phil. Trans. R. Soc. A*, 373, 20140159, 2015.



Shupe, M. D., and Intrieri, J. M.: Cloud radiative forcing of the Arctic surface: The influence of cloud properties, surface albedo, and solar zenith angle, *Journal of Climate*, 17, 616-628, 2004.

Shupe, M. D.: A ground-based multisensor cloud phase classifier, *Geophysical Research Letters*, 34, [10.1029/2007JD008737](https://doi.org/10.1029/2007JD008737), 2007.

5 Shupe, M. D.: Clouds at Arctic atmospheric observatories. Part II: Thermodynamic phase characteristics, *Journal of Applied Meteorology and Climatology*, 50, 645-661, [10.1175/JAMC-D-15-0054.1](https://doi.org/10.1175/JAMC-D-15-0054.1), 2011.

Shupe, M. D., Walden, V. P., Eloranta, E., Uttal, T., Campbell, J. R., Starkweather, S. M., and Shiobara, M.: Clouds at Arctic Atmospheric Observatories. Part I: Occurrence and Macrophysical Properties, *Journal of Applied Meteorology and Climatology*, 50, 626-644, [10.1175/2010jamc2467.1](https://doi.org/10.1175/2010jamc2467.1), 2011.

10 Shupe, M. D., Turner, D. D., Zwink, A., Thieman, M. M., Mlawer, E. J., and Shippert, T.: Deriving Arctic cloud microphysics at Barrow, Alaska: algorithms, results, and radiative closure, *Journal of Applied Meteorology and Climatology*, 54, 1675-1689, 2015.

Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., Miller, H. L., and Intergovernmental Panel on Climate, C.: Climate change 2007: Synthesis Report. Contribution of Working Group I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Summary for Policymakers, *Climate change 2007: Synthesis Report. Contribution of Working Group I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Summary for Policymakers*, 22 pp.-22 pp., 2007.

15 Assessment Report of the Intergovernmental Panel on Climate Change. Summary for Policymakers, 22 pp.-22 pp., 2007.

Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z. E., Illingworth, A. J., O'Connor, E. J., Rossow, W. B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., Mitrescu, C., and CloudSat Sci, T.: The CloudSat mission and the A-train - A new dimension of space-based observations of clouds and precipitation, *Bulletin of the American Meteorological Society*, 83, 1771-1790, [10.1175/bams-83-12-1771](https://doi.org/10.1175/bams-83-12-1771), 2002.

20 Taylor, P. C., Kato, S., Xu, K.-M., and Cai, M. C. J. D.: Covariance between Arctic sea ice and clouds within atmospheric state regimes at the satellite footprint level, 120, 12656-12678, 2015.

Tjernström, M., Leck, C., Birch, C. E., Brooks, I. M., Shupe, M. D., Persson, P. O. G., Wheeler, C. R., Sedlar, J., Mauritsen, T., and Paatero, J.: Meteorological conditions in the central Arctic summer during the Arctic Summer Cloud Ocean Study (ASCOS), *Atmospheric Chemistry and Physics*, 12, 6863-6889, 2012.

25 Tjernström, M., Leck, C., Birch, C. E., Bottenheim, J. W., Brooks, B. J., Brooks, I. M., Bäcklin, L., Chang, R.-W., de Leeuw, G., and Di Liberto, L.: The Arctic Summer Cloud Ocean Study (ASCOS): overview and experimental design, *Atmospheric Chemistry and Physics*, 14, 2823-2869, 2014.

Uttal, T., Curry, J. A., McPhee, M. G., and Perovich, D. K.: Surface heat budget of the Arctic Ocean, *Bulletin of the American Meteorological Society*, 83, 255, 2002.

30 Vaughan, M. A., Powell, K. A., Kuehn, R. E., Young, S. A., Winker, D. M., Hostetler, C. A., Hunt, W. H., Liu, Z., McGill, M. J., and Getzewich, B. J.: Fully Automated Detection of Cloud and Aerosol Layers in the CALIPSO Lidar Measurements, *Journal of Atmospheric and Oceanic Technology*, 26, 2034-2050, [10.1175/2009jtecha1228.1](https://doi.org/10.1175/2009jtecha1228.1), 2009.

Matt Shupe 3/29/2017 6:38 PM

Deleted: c

Matt Shupe 3/29/2017 6:38 PM

Deleted: s

Matt Shupe 3/29/2017 6:38 PM

Deleted: a

Vaughan, M. A., Winker, D. M., and Powell, K. A.: CALIOP algorithm theoretical basis document, part 2: Feature detection and layer properties algorithms, Rep. PC-SCI, 202, 87, 2005.

Wang, Z., Stephens, G., Deshler, T., Trepte, C., Parish, T., Vane, D., Winker, D., Liu, D., and Adhikari, L.: Association of Antarctic polar stratospheric cloud formation on tropospheric cloud systems, *Geophysical Research Letters*, 35, 2008.

- 5 Winker, D. M., Pelon, J., and McCormick, M. P.: The CALIPSO mission: Spaceborne lidar for observation of aerosols and clouds, in: *Proceedings of the Society of Photo-Optical Instrumentation Engineers (Spie), Conference on Lidar Remote Sensing for Industry and Environment Monitoring III*, Hangzhou, Peoples R China, 2002, WOS:000182448300001, 1-11, 2003.

Zhao, M., and Wang, Z.: Comparison of Arctic clouds between European Center for Medium-Range Weather Forecasts

- 10 simulations and Atmospheric Radiation Measurement Climate Research Facility long-term observations at the North Slope of Alaska Barrow site, *Journal of Geophysical Research: Atmospheres*, 115, 2010.

Zygmuntowska, M., Mauritsen, T., Quaas, J., and Kaleschke, L.: Arctic Clouds and Surface Radiation—a critical comparison of satellite retrievals and the ERA-Interim reanalysis, *Atmospheric Chemistry and Physics*, 12, 6667-6677, 2012.








Figure 1: Cloud fraction vertical distribution from surface observations at (a) Barrow, and (b) Eureka for 2006-2010 (after Shupe et al. 2011).

5






5 **Figure 2: Cloud fraction vertical distribution 2006-2010 from (a) CALIPSO 5 km, (b) 2B-GEOPROF, and (c) 2B-GEOPROF-lidar at Barrow; (d) CALIPSO 5 km, (e) 2B-GEOPROF, and (f) 2B-GEOPROF-lidar at Eureka.**




Deleted:

Yinghui Liu 3/22/2017 3:55 PM




Deleted:

Yinghui Liu 3/22/2017 3:55 PM




Deleted:

Yinghui Liu 3/22/2017 4:06 PM




Deleted:

Yinghui Liu 3/22/2017 4:06 PM



Deleted:

Yinghui Liu 3/22/2017 4:07 PM






Figure 3: Cloud fraction vertical distribution difference for 2006-2010 of (a) CALIPSO 5km, (b) 2B-GEOPROF, and (c) 2B-GEOPROF-lidar and surface at Barrow; and of (d) CALIPSO 5 km, (e) 2B-GEOPROF, and (f) 2B-GEOPROF-lidar and surface at Eureka.

5








Figure 4: Mean vertical distributions of cloud fraction from surface, and the difference of 2B-GEOPROF-lidar, CALIPSO 1 km, and CALIPSO 5 km, and 2B-GEOPROF minus surface observations at (a) Barrow, and (b) Eureka for 2006-2010.



Deleted:  
Yinghui Liu 3/22/2017 4:09 PM



Deleted:



Yinghui Liu 3/22/2017 4:12 PM

Deleted:

Yinghui Liu 3/22/2017 4:12 PM

Formatted: Normal

Yinghui Liu 3/22/2017 4:12 PM

Formatted: English (UK)

5 Figure 5: Monthly mean cloud fraction from surface, 2B-GEOPROF-lidar, CALIPSO 1km, CALIPSO 5 km, and 2B-GEOPROF at (a) Barrow, and (b) Eureka (bottom) for 2006-2010.







Figure 6: Mean cloud fraction above 960 m only, cloud below 960 m only, and cloud below and above 960 m from surface observations at Barrow (top) and Eureka (bottom) for 2006-2010.






5 **Figure 7:** Vertical Distributions of ice phase cloud (1<sup>st</sup> row), liquid phase cloud (2<sup>nd</sup> row), and mixed-phase cloud (3<sup>rd</sup> row) from 2B-CLDCLASS-lidar (left column), from surface (middle column), and difference of 2B-CLDCLASS-lidar and surface at Barrow for 2006-2010.




Deleted: ... [6]

Yinghui Liu 3/22/2017 4:15 PM




Deleted:

Yinghui Liu 3/22/2017 4:16 PM




Deleted:

Yinghui Liu 3/22/2017 4:16 PM




Deleted:


Yinghui Liu 3/22/2017 4:16 PM




Deleted:



**Figure 8: Monthly mean cloud fraction from surface, and 2B-CLDCLASS-lidar 2006-2010 (a) ice cloud, (b) liquid cloud, and (c) mixed phase cloud at Barrow; 2010 (d) ice cloud, (e) liquid cloud, and (f) mixed phase cloud at Eureka.**




5 **Figure 9:** Vertical Distributions of ice phase cloud (1<sup>st</sup> row), liquid phase cloud (2<sup>nd</sup> row), and mixed-phase cloud (3<sup>rd</sup> row) from 2B-CLDCLASS-lidar (left column), from surface (middle column), and difference of 2B-CLDCLASS-lidar and surface at Eureka for 2006-2010.




Deleted:

Yinghui Liu 3/22/2017 4:17 PM




Deleted:

Yinghui Liu 3/22/2017 4:18 PM




Deleted:

Yinghui Liu 3/22/2017 4:18 PM




Deleted:

Yinghui Liu 3/22/2017 4:18 PM



Deleted:

Yinghui Liu 3/22/2017 4:19 PM







Figure 10: Blended cloud fraction/frequency vertical distribution at Barrow and Eureka with combined surface and space observations from 2B-GEOPROF-lidar for 2006-2010.

5






Figure 11: Blended vertical distributions of (a) ice phase cloud, (b) liquid phase cloud, and (d) mixed-phase cloud at Barrow, and (d) ice phase cloud, (e) liquid phase cloud, and (f) mixed-phase cloud at Eureka from 2B-CLDCLASS-lidar and surface observations for 2006-2010.

5

Yinghui Liu 3/22/2017 4:21 PM


Deleted: ... [7]

Yinghui Liu 3/22/2017 4:22 PM

Deleted: ... [8]

Yinghui Liu 3/22/2017 4:23 PM

Deleted:



**Figure 12: Vertical Distributions of cloud water content for ice cloud from (a) 2B-CWC-RO, and (b) surface, for liquid cloud from (c) 2B-CWC-RO, and (d) surface at Barrow for 2006-2010.**




Figure 13: Same as Figure 12, but for Eureka.