



Source apportionment of NMVOCs in the Kathmandu Valley during the SusKat-ABC international field campaign using positive matrix factorization

Chinmoy Sarkar¹, Vinayak Sinha¹, Baerbel Sinha¹, Arnico K. Panday², Maheswar Rupakheti^{3,4}, and Mark G. Lawrence³

¹Department of Earth and Environmental Sciences, Indian Institute of Science Education and Research (IISER) Mohali, Sector 81, S. A. S. Nagar, Manauli PO, Punjab, 140306, India

²International Centre for Integrated Mountain Development (ICIMOD), Khumaltar, Lalitpur, Nepal

³Institute for Advanced Sustainability Studies (IASS), Berliner Str. 130, 14467 Potsdam, Germany

⁴Himalayan Sustainability Institute (HIMSI), Kathmandu, Nepal

Correspondence to: V. Sinha (vsinha@iisermohali.ac.in)

Abstract. A positive matrix factorization model (US EPA PMF version 5.0) was applied for the source apportionment of the dataset of 37 NMVOCs measured over a period of 19 December 2012 – 30 January 2013 during the SusKat-ABC international air pollution measurement campaign using a Proton Transfer Reaction Time of Flight Mass Spectrometer in the Kathmandu Valley. In all, 5 eight source categories were identified with the PMF model using the new “constrained model operation” mode. Unresolved industrial emissions and traffic source factors were the major contributors to the total measured NMVOC mass loading (17.9 % and 16.8 %, respectively) followed by mixed industrial emissions (14.0 %), while the remainder of the source was split approximately evenly between residential biofuel use and waste disposal (10.9 %), solvent evaporation (10.8 %), biomass 10 co-fired brick kilns (10.4 %), biogenic emissions (10.0 %) and mixed daytime factor (9.2 %). Conditional probability function (CPF) analyses were performed to identify the physical locations associated with different sources. Source contributions to individual NMVOCs showed biomass co-fired brick kilns significantly contribute to the elevated concentrations of several health relevant NMVOCs such as benzene. Despite the highly polluted conditions, biogenic emissions had largest contribution 15 (24.2 %) to the total daytime ozone production potential, even in winter, followed by solvent evaporation (20.2 %), traffic (15.0 %) and unresolved industrial emissions (14.3 %). Secondary organic aerosol (SOA) production had approximately equal contributions from biomass co-fired brick kilns (28.9 %) and traffic (28.2 %). Comparison of PMF results based on the in-situ data versus REAS v2.1 and EDGAR v4.2 emission inventories showed that both the inventories underestimate the 20 contribution of traffic and do not take the contribution of brick kilns into account. In addition, the REAS inventory overestimates the contribution of residential biofuel use and underestimates the contribution of solvent use and industrial sources in the Kathmandu Valley. The quantitative source



apportionment of major NMVOC sources in the Kathmandu Valley based on this study will aid in
improving hitherto largely un-validated bottom up NMVOC emission inventories, enabling more
25 focused mitigation measures and improved parameterizations in chemical-transport models.

1 Introduction

Non-methane volatile organic compounds (NMVOCs) are important atmospheric constituents and
are emitted from both natural and anthropogenic sources (Hewitt, 1999). They are important as pre-
cursors of surface ozone and secondary organic aerosols (SOA) and affect atmospheric oxidation
30 capacity, climate and human health (IPCC, 2013). Thus, identification of NMVOC sources is nec-
essary for devising appropriate mitigation strategies to improve air quality and reduce undesired
impacts of secondary pollutants such as tropospheric ozone and secondary organic aerosol.

Source apportionment of NMVOCs can be achieved by applying source–receptor models to mea-
sured ambient datasets. Ambient NMVOC mixing ratios depend on the emission profiles of the
35 sources contributing to the ambient mixture, their relative source strengths, transport, mixing and
removal processes in the atmosphere. Source receptor models perform statistical analyses on the
dataset to identify and quantify the contribution of different sources to the measured NMVOC con-
centrations (Watson et al., 2001). Positive matrix factorization (PMF) is currently among the most
widely applied receptor models for the source apportionment of NMVOCs, in particular for datasets
40 with high temporal resolution (Anderson et al., 2002; Miller et al., 2002; Kim et al., 2005; Buzcu and
Fraser., 2006; Brown et al., 2007; Vlasenko et al., 2009; Slowik et al., 2010; Yuan et al., 2012; Crippa
et al., 2013; Kaltsonoudis et al., 2016). In comparison to other receptor models based on principal
component analysis/absolute principal component scores (PCA/APCS) (Guo et al., 2004, 2006),
chemical mass balance (CMB) (Na and Pyo Kim., 2007; Morino et al., 2011) and UNMIX (Jorquera
45 and Rappenglück., 2004; Olson et al., 2007), PMF provides more robust results as it does not per-
mit negative source contributions. Moreover, a priori knowledge about the number and signature of
NMVOC source profiles are not required, which is particularly useful and apt for NMVOC source
apportionment studies in a new or understudied atmospheric chemical environment. The recently de-
veloped PMF version 5.0 also allows further refining the solution and reducing rotational ambiguity
50 of the solutions using pre-existing knowledge of emission ratios from known point sources. Source
apportionment of non-methane hydrocarbons (NMHCs) and oxygenated VOCs (OVOCs) using PMF
source–receptor models has been carried out in several previous studies (Shim et al., 2007; Leuchner
and Rappenglück, 2010; Gaimoz et al., 2011; Bon et al., 2011; Chen et al., 2014).

NMVOC emission inventories are frequently associated with large uncertainties (Zhang et al.,
55 2009). This is particularly true for metropolitan cities in the developing world. Emission inventories
can be evaluated using the results obtained from source receptor models such as the PMF model. This
evaluation is important to improve the accuracy of the existing emission inventories and therefore to



develop effective air pollution control strategies. In this study, we report the application of the PMF model for source apportionment of NMVOCs using the NMVOC data measured in the Kathmandu Valley, Nepal, which has been reported and analyzed in detail in Sarkar et al. (2016).

Kathmandu is considered to be amongst the most polluted cities in Asia (Panday et al., 2009). According to the existing Nepalese emission inventory (International Centre for Integrated Mountain Development's (ICIMOD) database) and the REAS v2.1 (Kurokawa et al., 2013) emission inventories residential biofuel use is considered to be the most important anthropogenic source of NMVOCs in the Kathmandu Valley. It is considered to contribute $\sim 67\%$ (REAS) to $\sim 83\%$ (Nepalese inventory), towards the total NMVOC mass loadings. In contrast, EDGAR v4. (Olivier et al., 1994) attributes 66% of the emissions in the Kathmandu Valley to solvent use and a recent emission inventory study conducted by the International Centre for Integrated Mountain Development (ICIMOD) which relied on measurement of particulate matter suggested that traffic is the dominant source (69%) of air pollution in a part of the Kathmandu Valley within the Ring Road (i.e. the Kathmandu Metropolitan City (KMC) and Lalitpur Sub-metropolitan City) and some nearby sub-urban rural areas outside the Ring Road (Pradhan et al., 2012).

The objective of the current study is to identify and quantify the contributions of different emission sources to the ambient wintertime NMVOC concentrations in the Kathmandu Valley using a positive matrix factorization (US EPA PMF 5.0; Brown et al. (2015)) receptor model. NMVOC measurements were carried out at Bode, a suburban site in the Kathmandu Valley over a period of 19 December 2012 – 30 January 2013 during the SusKat-ABC field campaign. The NMVOC measurements, new findings and qualitative analyses of sources have been presented and discussed in Sarkar et al. (2016). The NMVOC measurements suggested significant contribution of varied emission sources such as traffic (associated with high toluene, xylenes and trimethylbenzenes), biomass co-fired brick kilns (associated with high acetonitrile and benzene), industries and wintertime biogenic sources (as characterized by high daytime isoprene). Based on the NMVOCs emission profiles, two distinct periods were identified in the dataset: the first period (19 December 2012 – 3 January 2013) was associated with high daytime isoprene concentrations whereas the second period (4 – 18 January 2013) was associated with sudden increase in acetonitrile and benzene concentrations which was attributed to the start in operations of biomass co-fired brick kilns in the Kathmandu Valley (Sarkar et al., 2016). For quantitative source apportionment, hourly mean measured concentrations of all 37 NMVOCs measured during the instrumental deployment (19 December 2012 – 30 January 2013), were used for the PMF analysis. Sensitivity tests were conducted for the PMF 5.0 model version to evaluate how the new rotational tool called “constrained model operation feature” improves the representation of source profiles in the PMF model output. To identify the physical locations for the identified sources, an important prerequisite for targeted mitigation, conditional probability function (CPF) analyses were also performed. The results obtained from the PMF analyses were compared with three emission inventories – the existing Nepalese inventory, REAS v2.1 (Regional Emission



95 inventory in ASia) and the EDGAR v4.2 (Emissions Database for Global Atmospheric Research)
emission inventory. Additionally, the contributions of each source category to individual NMVOC
mass concentrations, ozone formation potential and formation of secondary organic aerosol (SOA)
were also analyzed.

2 Materials and Methods

100 2.1 Positive Matrix Factorization (PMF)

The United States Environmental Protection Agency's (US EPA) Positive Matrix Factorization (PMF)
receptor model version 5.0 was used for source apportionment of NMVOCs in the Kathmandu Val-
ley. The model is based on the multi-linear engine (ME-2) approach and has been described in detail
by Paatero (1997, 1999). From a data matrix of a number of NMVOCs in a given number of sam-
105 ples, the PMF model helps to determine the total number of possible NMVOC source factors, the
chemical fingerprint (source profile) for each factor, the contribution of each factor to each sample,
and the residuals of the dataset using the following equation (Paatero and Tapper, 1994),

$$X_{ij} = \sum_{k=1}^p g_{ik} f_{kj} + e_{ij} \quad (1)$$

Where, X_{ij} is the NMVOC data matrix with i number of samples and j number of measured
110 NMVOCs which are resolved by the PMF to provide p number of possible source factors with the
source profile f of each source and mass g contributed by each factor to each individual sample,
leaving the residuals e for each sample. To obtain the solution of equation (1), sum of the squared
residuals (e^2) and variation of data points (σ^2) are inversely weighted in PMF as expressed by the
following equation (Paatero and Tapper, 1994),

$$115 Q = \sum_{i=1}^n \sum_{j=1}^m \left(\frac{e_{ij}}{\sigma_{ij}} \right)^2 = \sum_{i=1}^n \sum_{j=1}^m \left(\frac{X_{ij} - \sum_{k=1}^p g_{ik} f_{kj}}{\sigma_{ij}} \right)^2 \quad (2)$$

Where, Q is the object function and a critical parameter for PMF, n is the number of samples, and
 m is the number of considered species. The original data should always be reproduced by the PMF
model within the uncertainty considering the non-negativity constraint for both the predicted source
profile and the predicted source contributions. The explained variability (EV) as given below demon-
120 strates the relative contribution of each factor to the individual compound and can be expressed as
(Gaimoz et al., 2011),

$$EV_{kj} = \frac{\sum_{i=1}^n |g_{ik} f_{kj}| / \sigma_{ij}}{\sum_{i=1}^n (\sum_{k=1}^p |g_{ik} f_{kj}| + |e_{ij}|) / \sigma_{ij}} \quad (3)$$

The explained variability is most useful to policy makers. If the observed mass loading of a com-
pound that is known to be harmful to human health is high, the explained variability will indicate
125 which sources are responsible for most of its emissions and what fraction of the total observed mass
is contributed by each source. Therefore, this allows planning mitigation strategies.



To ascertain the magnitude of random errors that can be caused due to the use of random seeds followed by the selection of the run with the lowest Q due to the existence of infinite solutions with different g_{ik} , f_{kj} and e_{ij} matrices but identical $Q = \sum_{i=1}^n \sum_{j=1}^m (e_{ij}/\sigma_{ij})^2$, bootstrap runs were performed. In the bootstrap runs, the timeseries is partitioned into smaller segments of a user specified length and the PMF is run on each of these smaller segments, for the same number of factors as the original model run. The model output of each bootstrap run is mapped onto the original solution using a cross correlation matrix of the factor contributions g_{ik} of a given bootstrap run with the factor contributions g_{ik} of the same time segment of the original solution using a threshold of the Pearson's correlation coefficient (R) > 0.6 . The bootstrap factor is assigned to the factor with which it is most strongly positively correlated, as long as the value of R is greater than 0.6. If it cannot be attributed to any factor of the original solution it will be termed unmapped. The presence of a high fraction unmapped factor ($> 20\%$) is a clear indication of large random errors and should be investigated carefully. In our analysis, no unmapped factors were present.

For each factor, the factor profile of all bootstrap runs combined is compared with the profile of the original model output. The model will provide a box and whisker plot for the mass loading ($\mu\text{g m}^{-3}$) and percentage of each compound attributed to the factor profile of each of the factors during the bootstrap runs. It will also ascertain for each compound whether or not the original solution for that factor falls into the interquartile range of the bootstrap results and provide this information in a table format.

When all sources are equally strong throughout the entire period, this bootstrap model provides a robust estimate of the total random error. However, if one of the sources is completely absent for a significant fraction of the total hours (like the brick kiln source throughout the first 13 days of the SusKat-ABC campaign), the bootstrap model may overestimate the random error substantially. For such a source, mass loading of all the compounds that contribute strongly to the factor profile of the source will typically be outside the interquartile range. For the same set of compounds, similar behavior could also be seen for the factor profile of several other factors. In such a situation, the error estimate of the bootstrap runs should only be considered as the upper limit of the potential random error.

In addition to the random error, the PMF model has rotational ambiguity. There can be multiple solutions with a different factor profile for all factors for which the model will find a different local minimum of the residual matrix while determining the factor contribution matrix. This fact that different solution for $g_{ik}f_{kj}$ with the same sum of the scaled residuals $Q = \sum_{i=1}^n \sum_{j=1}^m (e_{ij}/\sigma_{ij})^2$ exist is called the rotational ambiguity of the model. The PMF 5.0 has a new feature named as "the constrained model operation" in which the rotational ambiguity of the model can be constrained using external knowledge of the source composition (f_{kj}) or contribution (g_{ik}) matrix. For instance, if a source was inactive for a particular period, then the contribution due to that factor during that time period could be pulled to zero in the model to provide more robust output. Alternatively, the



emission ratios obtained from a particular source through samples collected at the source can also
165 be used to constrain the model. Constraining the PMF model using such external knowledge gives
rise to a penalty in Q (the object function) and a maximum penalty of 5 % is recommended as
a reasonable threshold (Paatero and Hopke, 2009). A detailed discussion of the use of constraints
to a receptor model has been provided in previous studies (Paatero et al., 2002; Rizzo and Scheff,
2007; Paatero and Hopke, 2009; Norris et al., 2009).

170 2.2 Implementation of PMF

PMF was applied to the hourly averaged dataset of 37 ions measured using a Proton Transfer Re-
action Time of Flight Mass Spectrometer (PTR-TOF-MS). All relevant analytical details pertaining
to the site description, meteorology, sampling and quality assurance of the NMVOC dataset has al-
ready been described in detail in the companion paper to this special issue (Sarkar et al., 2016).

175 Briefly, NMVOC measurements during this study were performed in the winter season from 19 De-
cember 2012 until 30 January 2013 at Bode (27.689° N, 85.395° E, 1345 m a.m.s.l.) in Bhaktapur
district, which is a suburban site located in the westerly outflow of the Kathmandu Metropolitan
City. The land use in the vicinity of the measurement site consisted of the following cities - Kath-
mandu Metropolitan City (~ 10 km to the west), Lalitpur Sub-Metropolitan City (~ 12 km south-
180 west of the site) and Bhaktapur Municipality (~ 5 km south-east of the site). The site is located in
the Madhyapur-Thimi Municipality. In addition, the region north of the site had a small forested
area (Nilbarahi Jungle; ~ 0.5 km² area) and a reserve forest (Gokarna Reserve Forest; ~ 1.8 km²
area) at approximately 1.5 km and 7 km from the measurement site, respectively. The Tribhuvan
International Airport is located at about 4 km to the west of the Bode site.

185 All the available data during this aforementioned study (Sarkar et al., 2016) were used for the
PMF analysis and the missing values were replaced by a missing value indicator (-999). To ensure
that differential uncertainties do not drive the object function Q and give undue weightage to cali-
brated organic ions while constructing source profiles, we followed the procedure used by Leuchner
and Rappenglück (2010) for source apportionment of NMVOCs in the Houston Ship Channel area,
190 assigning a constant uncertainty of 20 % for all the ions. The attribution of ions to parent compounds
and corresponding detection limits were as described in Sarkar et al. (2016). Due to its erratic time-
series profile, HCN ($m/z = 28.007$) was classified as a weak species in the PMF input while all
other ions were classified as strong species. All the input data was converted from mixing ratios ppb
to mass concentrations ($\mu\text{g m}^{-3}$) using the relevant temperature, pressure and molecular weight. The
195 total measured NMVOC concentration was calculated by adding the mass concentrations of all mea-
sured NMVOCs and was classified as a weak species in the PMF input. All the measured ions had
a signal to noise (S/N) ratio greater than 2. Table S1 of the supplementary information shows the
signal to noise (S/N) ratios for all input NMVOC species used in the PMF along with other statistical
parameters of the dataset.



Table 1. Diagnostic for the results of the positive matrix factorization (PMF) model run

n (samples)	1006
m (species)	37
k (factors)	8
Q (theoretical)	4480.37
Q (model)	4562.89
Mean ratio NMVOC(estimated)/NMVOC(observed)	0.999

200 PMF model runs ranging from 5 to 12-factor numbers were carried out to ascertain the best solution for this study, consistent with the chemical environment of the Kathmandu Valley. Based on the $Q/Q_{theoretical}$ ratio, the physical plausibility of the factors and the rotational ambiguity of the solution, an 8-factor solution was deemed to be the best for this dataset. When less than 7-factors were employed, several source profiles appeared to be mixed, indicating inadequate resolution of sources.

205 The solution incorporating 7-factors caused strong overlap of mixed industrial emissions with the unresolved industrial emissions factor. This solution was considered inappropriate, as the daytime biogenic emissions could not be separated from the nighttime combustion source of isoprene in the 7-factor solution. Even when the model was nudged towards separating the biogenic emissions and the anthropogenic combustion sources of isoprene using the constraint mode, this separation could
210 only be accomplished with a large penalty on Q in the 7-factor solution. The 9-factor solution had too much rotational ambiguity and assigned brick kiln emissions to two largely co-linear factors, both of which had an incomplete source profile with respect to aromatic compounds and were essentially created to better account for minor variations in the emission ratios associated with brick kiln emissions during the firing up period and the continuous operation later in the campaign.

215 The diagnostics for the 8-factor solution are summarized in Table 1. The eight factors were - 1) traffic, 2) residential biofuel use and waste disposal, 3) mixed industrial emissions, 4) biomass co-fired brick kilns, 5) unresolved industrial emissions, 6) solvent evaporation, 7) mixed daytime source and 8) biogenic emissions. To identify the uncertainty associated with the PMF solution, bootstrap runs were performed 100 times taking 96 hours as the segment length. There were no unmapped
220 factors in the bootstrap runs.

Figure 1 shows the correlation between the estimated total measured NMVOC concentrations calculated using the contributions from all factors (vertical axis) with measured total measured NMVOC concentrations (horizontal axis). An excellent correlation ($r^2 = 0.99$) indicates that PMF model can explain almost all variance in the total measured NMVOC concentrations.

225 The constrained model mode was used to further improve the 8-factor solution. The original model output showed positive correlations between factors such as the biomass co-fired brick kilns and mixed industrial emissions ($r^2 = 0.27$) as well as the residential biofuel use and waste disposal factor with traffic factor ($r^2 = 0.42$). Since this is a new feature and has only recently been used by




Figure 1. Correlation between estimated and observed NMVOC concentrations

Brown et al. (2015) for ambient air data, a detailed description of the implementation procedure and
 230 an analysis of how the constraints affected the model output is provided here. Several constraints
 were used to obtain a more robust PMF solution.

First, the upper limit for the emission ratio of the individual aromatic compounds to isoprene
 as reported by Misztal et al. (2015) were used to constrain the factor profile of primary biogenic
 emissions. As a small fraction of the biogenic isoprene gets attributed to other daytime factor (mixed
 235 daytime) by the PMF model, the same constraint were used on mixed daytime factor and the solvent
 evaporation factor as well.

Second, it was assumed that aromatic compounds and acetonitrile are not photochemically pro-
 duced. Acetic acid is associated with both mixed daytime and solvent evaporation, and so the ratios
 of aromatic compounds and acetonitrile to acetic acid were nudged towards 0.0001 for these two
 240 factors.

Third, to improve the representation of brick kiln emissions, and the residential biofuel use and
 waste disposal in the model, the respective factors, which were clearly identified in the original
 model solution, were nudged using the emission ratios of aromatic compounds to benzene from
 grab samples of domestic waste burning (garbage burning grab sample) and fixed chimney bull's
 245 trench brick kiln emissions (FCBTBK grab sample) collected directly at the point source. This was
 required, because in the original model output, the residential biofuel use and waste disposal factor
 correlated with the traffic factor ($r^2 = 0.42$) while the brick kilns emission factor correlated with
 the mixed industrial emissions factor ($r^2 = 0.27$). This indicates that there was substantial rotational
 ambiguity for these two factor pairs.

250 Nudging was performed by exerting a soft pull allowing for a maximum 0.2% change in Q for
 each constraint. A soft pull allows the change in the Q value up to a certain limit by pulling the
 values to a target value for an expression of elements (the emission ratio). If no minima can be found
 for which the change in $Q = \sum_{i=1}^n \sum_{j=1}^m (e_{ij}/\sigma_{ij})^2$ is less than 0.2% in the $g_{ik}f_{kj}$ matrix after f_{kj}
 has been constrained, no change was made and the original solution was retained. If the condition



Table 2. Inter NMVOC emission ratios used for biogenic, solvent evaporation and mixed daytime factors to nudge the PMF model and the corresponding emission ratios before and after nudging

ERs/Isoprene	ERs used to nudge	BG		SE		MD	
		before nudging	after nudging	before nudging	after nudging	before nudging	after nudging
Acetonitrile	0.002	0.06	0.00	0.00	0.004	2.78	1.75
Benzene	0.002	0.29	0.00	0.52	0.00	0.15	0.00
Toluene	0.012	0.10	0.01	0.39	0.00	4.82	0.00
Styrene	0.002	0.02	0.00	0.06	0.00	0.00	0.002
Xylenes	0.002	0.00	0.0002	0.35	0.41	4.65	0.00
Trimethylbenzenes	0.002	0.06	0.01	0.09	0.00	1.85	0.20
Naphthalene	0.002	0.31	0.30	0.36	0.60	0.00	0.002
ERs/Acetic acid	ERs used to nudge	BG		SE		MD	
		before nudging	after nudging	before nudging	after nudging	before nudging	after nudging
Acetonitrile	0.0001	0.57	0.00	0.00	0.0001	0.07	0.09
Benzene	0.002	1.48	0.00	0.04	0.00	0.01	0.00
Toluene	0.0001	1.01	0.004	0.05	0.00	0.12	0.00
Styrene	0.0001	0.15	0.00	0.01	0.00	0.00	0.0001
Xylenes	0.0001	0.00	0.0001	0.04	0.01	0.12	0.00
Trimethylbenzenes	0.0001	0.59	0.004	0.01	0.00	0.05	0.01
Naphthalene	0.0001	3.08	0.15	0.04	0.01	0.00	0.0001

BG = Biogenic; SE = Solvent evaporation; MD = Mixed daytime

255 can be met without changing Q by more than the threshold, the revised factor profiles will be used
as the base upon which the next constraint in the list of constraints will be executed.

Implementing the constraints mentioned above, significantly improved the representation of bio-
genetic emissions, mixed daytime and solvent evaporation factors. Figure S1 of the supplementary
information shows a comparison of the box and whisker plots of the biogenic emissions, mixed
260 daytime and solvent evaporation factors before and after nudging and demonstrates the significant
improvement after applying constraints.

After nudging, the contribution of the biogenic factor correlated better with solar radiation (r^2
= 0.48) while the mixed daytime factor correlated better with ambient temperature ($r^2 = 0.42$).
The factor profile of the solvent evaporation correlates better with the rise in solar radiation and
265 temperature after sunrise (07:00 - 09:00 LT; $r^2 = 0.53$). Table 2 represents the emission ratios used
to nudge the biogenic, mixed daytime and solvent evaporation factors and provides the corresponding
emission ratios (ERs) before and after nudging.



It can be seen that most constraints on the aromatic to isoprene ratio could be executed without exceeding the penalty on Q . In the biogenic factor, only the naphthalene/isoprene ratio could not be
270 constrained. The solvent evaporation and mixed daytime factors contain only a small fraction of the
total daytime isoprene (8 % and 7 %, respectively). Given the very small overall isoprene mass in
these two factor profiles, few additional ratios did not meet the constraining criteria in these factor
profiles (namely acetonitrile/isoprene and trimethylbenzenes/isoprene ratio in the mixed daytime
factor and the xylenes/isoprene and naphthalene/isoprene ratio in the solvent evaporation factor).
275 Some of these compounds (such as naphthalene) could not be constrained in the same factors while
constraining the ERs with respect to acetic acid.

The fact that the constrained run was incapable of removing naphthalene from the source profiles
of the biogenic and the solvent evaporation source and the fact that the diel profiles of both these
factors show a weak secondary peak between 17:00 - 22:00 LT, seems to indicate that an additional
280 weak combustion source with a high naphthalene emission ratio is possibly poorly represented by the
current 8-factor solution. Cooking on 3-stone fires is known to emit large amounts of benzene and
naphthalene (Stockwell et al., 2015) and the temporal profile of such a cooking source could overlap
with that of the garbage fires. It can be noted that 3-stone fires is still a common way to cook for
construction workers and brick kiln workers staying in temporary camps in the Kathmandu Valley.
285 This would make it challenging for the model to separate these two sources. We will henceforth refer
to the garbage burning factor as the residential biofuel use and waste disposal factor.

Figure S2a of the supplementary information shows the G-space plots for two factors, namely
biomass co-fired brick kilns and mixed industrial emissions. A stronger correlation ($r^2 = 0.42$) ex-
isted in the original solution prior to nudging with ERs of FCBTBK grab samples, which reduced to
290 $r^2 = 0.18$. Similarly, after nudging with ERs of the garbage burning grab sample the correlations be-
tween residential biofuel use and waste disposal was reduced from 0.27 to 0.18, as shown in Figure
S2b. Thus, the new solution fills the solution space better.

Table 3 summarizes the aromatics/benzene emission ratios derived from the PMF (before and
after nudging) and its comparison with the emission ratios obtained from grab samples for biomass
295 co-fired brick kilns and residential biofuel use and waste disposal sources. These emission ratios are
also compared with the ERs reported for 3-stone firewood stoves in Stockwell et al. (2015) and the
mixed garbage burning and open cooking fire sources reported for Nepal in Stockwell et al. (2016).

For the residential biofuel use and waste disposal source, the original model run already had
emission ratios very similar to the garbage burning grab samples of the garbage burning fire. The
300 constrained run improved the agreement further for styrene, trimethylbenzenes and naphthalene.
Constraining this factor with the ERs of 3-stone firewood stoves from Stockwell et al. (2015) instead
of our garbage burning grab samples resulted in a larger penalty on Q and did not improve the
representation of the biogenic, mixed daytime and solvent evaporation factors.



Table 3. Comparison of aromatics/benzene ERs (emission ratios) obtained from PMF (before and after nudging), respective grab samples, the 3-stone firewood source reported in Stockwell et al. (2015) and the mixed garbage burning and open cooking fire sources reported in Stockwell et al. (2016)

ERs/Benzene	FCBTBK grab samples	BK PMF (before nudging)	BK PMF (after nudging)	garbage burning grab samples	RB+WD PMF (before nudging)	RB+WD PMF (after nudging)	3-stone firewood ¹	Mixed garbage ²	Open hardwood cooking ²
Toluene	0.80	0.28	0.35	0.34	0.33	0.34	0.11	0.37	0.27
Styrene	0.08	0.05	0.06	0.16	0.22	0.18	0.09	0.19	0.11
Xylenes	0.58	0.16	0.22	0.25	0.28	0.25	0.10	0.18	0.12
Trimethylbenzenes	0.31	0.06	0.09	0.08	0.16	0.12	0.03	0.02	0.03
Naphthalene	0.09	0.14	0.15	0.09	0.16	0.11	0.40	-	-

1. Stockwell et al. (2015); 2. Stockwell et al. (2016); BK = Biomass co-fired brick kilns; RB+WD = Residential biofuel use and waste disposal

For brick kilns, the emission ratios of the constrained model output runs diverged from the emission ratios of the FCBTBK grab samples. However, the temporal profile of the activity, especially the closure of the brick kilns during the first part of the campaign is better captured by the constrained run and the correlation with mixed industrial emission sources reduced significantly. The FCBTBK grab samples were collected on 6 December 2014, two years after the SusKat study, so differences from the emission profiles observed during the SusKat-ABC campaign are a possibility. Alternatively, the differences could also stem from the inherently variable nature of this source. In particular, naphthalene and benzene were higher in the source profiles of the SusKat-ABC campaign compared to their relative abundances in the FCBTBK grab samples. At the time the FCBTBK grab samples were collected (on 6 December 2014), brick kilns were co-fired using coal, wood dust and sugarcane extracts. It is possible that in January, during peak winter season, a different type of biomass, one associated with higher benzene and naphthalene emissions (e.g. wood) was used in these biomass co-fired brick kilns, resulting in the slight disagreement between the PMF source profile and FCBTBK grab sample signature for this factor. Table S2 of the supplementary information shows the percentage contribution of PMF derived factors obtained from constrained runs with 5, 6, 7, 8 and 9-Factors.

2.3 Collection of grab samples

Grab samples from garbage fires (termed garbage burning) were collected near the measurement site (~ 200 m in the northern direction, upwind of Bode; 27.690° N, 85.395° E) on 7 December 2014 between 15:00 - 15:03 LT. A “brick kiln” grab sample was collected on 6 December 2014 from a fixed chimney bull’s trench brick kiln (FCBTK) co-fired using coal, wood dust and sugarcane



325 extracts. Figure S3 of the supplementary information shows pictures of the grab sample collection
and the instrumental setup for the analysis. The whole air samples were collected in 2 litre glass
flasks that had been validated for the stability of NMVOCs (Chandra et al., in preparation) and were
analyzed within 38 hours of the collection (on 9 December 2014 between 03:42 – 04:05 LT). The
whole air samples (WAS) were diluted (dilution factor of 9.93) using zero air for the quantification of
330 NMVOCs present in the grab samples using a PTR-MS (Sinha et al., 2014). The average background
signals (zero air) were subtracted from each m/z channel and stable data of at least 10 cycles (\sim
10 minutes) were considered for the calculation of mixing ratios as per the protocol described by
Sinha et al. (2014).

2.4 Conditional probability function (CPF) analyses

335 For identifying the physical locations associated with different local sources, conditional probability
function (CPF) analyses were performed. CPF is a well-established method to identify source loca-
tions of local sources based on the measured wind (Fleming et al., 2012). In CPF, the probability of
a particular source contribution from a specific wind direction bin exceeding a certain threshold is
employed which is calculated as follows:

$$340 \quad CPF = \frac{m_{\Delta\theta}}{n_{\Delta\theta}} \quad (4)$$

Where $m_{\Delta\theta}$ represents the number of data points in the wind direction bin $\Delta\theta$ which exceeded the
threshold criterion and $n_{\Delta\theta}$ represents the total number of data points from the same wind direction
bin. For this study, $\Delta\theta$ was chosen as 30° and data for wind speed $> 0.5 \text{ m}^{-1}$ were used.

3 Results and Discussion

345 3.1 Identification of PMF Factors

Figure 2 represents the factor profiles of all the eight factors resolved by the PMF model in which
grey bars indicate the mass concentrations and red lines with markers show the percentage of
a species in the respective factor.

Identification and attribution of these factors is discussed in detail in the following sections.

350 3.1.1 Factor 1 - Traffic

More than 60 % of the total toluene, sum of C8-aromatics, sum of C9-aromatics and ~ 37 % of the
total assorted hydrocarbons ($m/z = 97.102$ and 83.085) were explained by Factor 1. Toluene and
C8-aromatics contributed most (~ 16 % and ~ 13 %, respectively) to the total measured NMVOC
mass of Factor 1. In addition four other compounds also contributed ≥ 5 % to the total mass of this
355 factor (propyne (~ 11 %), acetone (~ 9 %), propene (~ 6 %) and sum of C9-aromatics (~ 5 %)).
The other 31 NMVOCs contributed ~ 40 % of the total measured NMVOC mass to this factor but




Figure 2. Factor profiles of the eight sources obtained by PMF analysis




Figure 3. Timeseries and diel box and whisker plot for Factor 1 (Traffic)

their individual contributions were $\leq 5\%$ each. The diel profile of Factor 1 (Figure 3) showed characteristic evening peak at 17:00 LT with an average concentration of $\sim 40 \mu\text{g m}^{-3}$. This evening peak showed large variability and plume-like characteristics as the average and median diverged frequently. Occasionally, the mass contribution of this factor amounted to $\sim 100 \mu\text{g m}^{-3}$. The high variability during the evening peak hour indicates that the source strength is not equal for all wind directions, but varies with fetch region.

Table 4 shows that the aromatics/benzene emission ratios for this factor are in good agreement with the emission ratios reported by previous studies for vehicular emissions in tunnel experiments and in metropolitan sites/megacities. In view of the diel profile and observed chemical signatures, Factor 1 was attributed to traffic. It can be noted that in winter, rush-hour in the city starts at 16:00 LT, while westerly winds still bring urban air to the measurement site. The morning rush hour in the city takes place in calmer winds which leads to a less sharp peak. It is interesting to note that $\sim 37\%$ of the total styrene was present in this factor and $\sim 31\%$ of the total isoprene was also explained by this factor. Few previous studies have reported traffic related sources of isoprene in urban areas (Borbon et al., 2001; Hellèn et al., 2012) and also estimated isoprene as one of the top 10 contributors to OH reactivity from traffic (Nakashima et al., 2010). Our results indicate that traffic can be a significant source of nighttime isoprene in the Kathmandu Valley.

3.1.2 Factor 2 - Residential biofuel use and waste disposal

Factor 2, too, showed regular evening hour peaks and a bimodal profile (Figure 4). However, the evening peak of average concentrations as high as $\sim 40 \mu\text{g m}^{-3}$ occurred after the traffic peak (at 19:00 LT) and had less variability, indicating that this source is an area source that is spatially spread throughout the Kathmandu Valley. The diel box and whisker plot also has a relatively weak morning peak (at 08:00 LT) with average concentrations of $\sim 18 \mu\text{g m}^{-3}$. Figure 2 shows, that this factor explains 30% of the total styrene, furan, 2-furaldehyde and acrolein.

Most of the measured NMVOC mass in this factor was contributed by acetic acid, propyne, methanol, benzene, propene and acetone + propanal ($\sim 14\%$, $\sim 12\%$, $\sim 10\%$, $\sim 9\%$, $\sim 7\%$ and



Table 4. Emission ratios of NMVOCs/benzene for aromatic hydrocarbons derived from the PMF model for factor attributed to traffic and comparison of ERs with previous studies for traffic source profiles

ERs/Benzene	Kathmandu PMF	Tunnel study, Stockholm ¹	Tunnel study, Hong Kong ²	Tunnel study, Taipei ³	Mexico City ⁴	Los Angeles ⁵
Toluene	3.41	3.89	2.27	2.38	3.47	2.45
C8-aromatics	2.89	2.81	0.87	1.86	3.55	1.38
C9-aromatics	1.20	-	0.77	1.36	2.31	0.48
Styrene	0.30	-	-	0.39	0.17	-
Naphthalene	0.19	-	0.10	-	-	-

1. Kristensson et al. (2004); 2. Ho et al. (2009); 3. Hwa et al. (2002); 4. Bon et al. (2011); 5. Borbon et al. (2013)




Figure 4. Timeseries and diel box and whisker plot for Factor 2 (Residential biofuel use and waste disposal)

~ 6 % respectively). The other 31 measured NMVOCs contributed ~ 42 % to this factor, but their individual contributions were ≤ 5 % each (Figure 2). It was observed that garbage/trash burning activities were more intense during evening hours in winter in the Kathmandu Valley. Table 5 shows a comparison of the aromatics/benzene emission ratios obtained from the PMF, with previously reported aromatics/benzene ratios for waste and trash burning, and with the emission ratios of garbage burning grab samples that were collected in the Kathmandu Valley near the point source (a household waste fire). It can be seen that the aromatics/benzene emission ratios of the PMF output are in excellent agreement with the values obtained for garbage burning grab samples collected in the Kathmandu Valley.

There is some agreement with the emission ratios reported in previous studies, though all of these previous studies found higher emission ratios for styrene. This could indicate that the composition of household waste in the Kathmandu Valley is different (less polystyrene, plastic and more biomass) or that the source profile is mixed with that of a second source, with similar spatial and temporal characteristics. Residential biofuel use is expected to have a similar temporal profile and did not appear as a separate factor in the PMF solution. Therefore, Factor 2 was attributed to residential biofuel use and waste disposal sources collectively.



Table 5. Emission ratios of NMVOCs/benzene for acetonitrile and aromatic hydrocarbons derived from the PMF model for the factor attributed to Residential biofuel use and burning household waste and comparison with previously reported studies and the garbage burning grab samples collected at the point source

ERs/Benzene	Kathmandu PMF	Kathmandu garbage burning grab samples	Mixed garbage burning ¹	Household waste burning ²	Open hardwood cooking ¹	Trash burning ³	Scrap tires burning ²
Acetonitrile	0.23	0.77	-	-	-	0.06	-
Toluene	0.34	0.34	0.37	0.38	0.27	0.41	0.63
C8-aromatics	0.25	0.25	0.19	0.22	0.11	0.10	0.43
C9-aromatics	0.12	0.08	0.18	-	0.12	0.03	0.03
Styrene	0.18	0.16	0.02	0.54	0.03	0.86	0.30
Naphthalene	0.11	0.09	-	0.01	-	0.10	0.30

1. Stockwell et al. (2016) ; 2. Lemieux et al. (2004) ; 3. Stockwell et al. (2015)




Figure 5. Timeseries and diel box and whisker plot for Factor 3 (Mixed industrial emissions)

3.1.3 Factor 3 - Mixed industrial emissions

400 This factor explained 66 % of the total ethanol, which is used as an industrial solvent. Moreover, ~ 20 – 25 % of the total propyne, propene, acetonitrile, dimethyl sulfide (DMS) and furan were also present in this factor. All these compounds have industrial sources (Karl et al., 2003; Kim et al., 2008) as they are widely used as solvents/reactants in various industrial processes and can be emitted during combustion processes. Therefore, Factor 3 was attributed to mixed industrial emissions. Most of the

405 measured NMVOC mass in this factor was contributed by propyne (~ 16 %), acetaldehyde (~ 15 %), ethanol (~ 10 %), propene (~ 9 %), methanol (~ 9 %), benzene (~ 8 %) and acetone + propanal (~ 5 %). The emissions reflect both release of chemicals used in the industrial units as well as emissions associated with combustion of a variety of fuels including biofuels. The other 30 NMVOCs jointly contributed only ~ 28 % of the total measured NMVOC mass and their individual contribution were

410 ≤ 5 % each. The emission strength of industrial sources is typically constant throughout the day and hence the observed mass concentrations are driven by boundary layer dynamics. The diel box and




Figure 6. Timeseries and diel box and whisker plot for Factor 4 (Biomass co-fired brick kilns)

whisker plot (Figure 5) shows a gradual increase in the mass concentrations throughout the night. The highest mass concentration are observed just after sunrise, when the inversion in the mountain Valley is most shallow. This shallow early morning boundary layer is caused by the cold pooling of
415 air at night, which results in an accumulation of cold air at the Valley bottom. The rising sun first warms the upper part of the Valley's atmosphere, while the Valley bottom is still in the shade of the surrounding mountains. Once direct sunlight reaches the Valley bottom, warming and thermally driven convection breaks the shallow boundary layer and wind speeds increase, increasing turbulent mixing under a growing boundary layer. The daytime mass concentrations of the mixed industrial
420 emissions are hence an inverse of the temperature and wind speed profile (Figure 5).

3.1.4 Factor 4 - Biomass co-fired brick kilns

The diel box and whisker plot of factor 4 (Figure 6) shows a profile that is similar to the profile of mixed industrial emissions, indicating that this factor should be attributed to a source that operates 24/7, as its mass loadings, too, represent an inverse of the temperature and wind speed profile. The
425 timeseries of Factor 4 showed sudden increase from 4 January 2013 at exactly the time when brick kilns in the Kathmandu Valley became operational (Sarkar et al., 2016).

Benzene (~ 23 %) contributed most to the total measured NMVOC mass of Factor 4. In addition acetaldehyde (~ 10 %), propyne (~ 8 %), toluene (~ 8 %), acetone (~ 7 %), acetic acid (~ 5 %) and xylenes (~ 5 %) also contributed significantly to the total measured NMVOC mass. The other 30
430 NMVOCs contributed ~ 34 % to the total measured NMVOC mass of this factor, but their individual contribution were ≤ 5 % each. Overall, factor 4 explained ~ 37 % of the total benzene and ~ 24 % of the total acetonitrile mass loading.

It is reported that brick kilns in the Kathmandu Valley burn large quantity of biomass, wood and crop residues along with coal (Stone et al., 2010; Sarkar et al., 2016) that can lead to significant
435 emission of aromatics and acetonitrile (Akagi et al., 2011; Yokelson et al., 2013; Sarkar et al., 2013). Therefore Factor 4 was attributed to the biomass co-fired brick kilns and the conditional probability function analysis (section 3.2) is consistent with this assignment.




Figure 7. Timeseries and diel box and whisker plot for Factor 5 (Unresolved industrial emissions)




Figure 8. Timeseries and diel box and whisker plot for Factor 6 (Solvent evaporation)

3.1.5 Factor 5 - Unresolved industrial emissions

Factor 5 explained $\sim 48\%$ of the total 1,3-butadiene, $\sim 35\%$ of the total methanol, $\sim 30\%$ of the total acetonitrile and 27% of the total propanenitrile and 24% of the total nitromethane. 1,3-butadiene is used in the production of several polymers and acetonitrile and propene can be side products in this process. Propanenitrile is used to start acrylic polymerization reactions in industrial processes. The largest use of methanol worldwide is as feedstock for the plastic industry and nitromethane is used in the synthesis of several important pharmaceutical drugs. It can be noted that several pharmaceutical industries are located in the Thimi area which is only ~ 2 km away from the measurement site. Nitromethane is also emitted from combustion of diesel fired generators (Inomata et al., 2013, 2014; Sekimoto et al., 2013) which are used as a back-up power source by both small and large industrial units in the Kathmandu Valley. It is, therefore, likely that miscellaneous nearby industries contributed significantly to the unresolved factor. The diel profile of Factor 5 (Figure 7) showed morning and evening peaks (at 09:00 - 10:00 LT and 17:00 LT, respectively), which is not typical for industrial emissions, but this factor always had a high background with average mass loadings of $\sim 20 \mu\text{g m}^{-3}$ throughout. The timeseries and diel profile (Figure 7) of this factor did not reveal characteristics that could be related uniquely to a known emission source.

Figure 7 displayed elevated daytime mass concentrations and an evening peak for this factor that occurs slightly before the traffic peak in the early evening during the first part of the SusKat-ABC



campaign (until 25 December). Towards the end of the campaign (from 10 January onwards), the same factor had diurnal variations that showed some similarity to profiles of both the solvent evaporation (morning peak) and mixed industrial emissions (slow rise throughout evening and nighttime) factors. Between 25 December and 10 January, diurnal patterns are weak and peaks in the unresolved factor seem to coincide with peaks in the solvent evaporation factor. This comparison of the diel profiles is shown in Figure S4 of the supplementary information. Since this factor seems to contain contributions of multiple sources and potentially the photooxidation products of their emissions, this factor was termed as the unresolved industrial emissions factor.

Most of the total measured NMVOC mass of Factor 5 was due to oxygenated NMVOCs like methanol (~ 14 %), acetic acid (~ 11 %), acetaldehyde (~ 9 %), acetone (~ 9 %) and formic acid (~ 9 %) but benzene, propyne and propene also contributed > 5 % (~ 9 %, ~ 6 % and ~ 6 %, respectively) to the total measured NMVOC mass of this factor. The other 29 NMVOCs together contributed only ~ 27 % to this factor and their individual contributions were less than 5 %.

3.1.6 Factor 6 - Solvent evaporation

Factor 6 explains approximately 25–40 % of the compounds containing the aldehyde functional group. It explained ~ 39 % of the total acetaldehyde, ~ 27 % of the total formaldehyde and ~ 23 % of 2-furaldehyde. Moreover, ~ 28 % of the total acetic acid and ~ 23 % of the total methylglyoxal were explained by this factor. Acetaldehyde and acetic acid contributed ~ 40 % and ~ 27 % respectively to the total measured NMVOC mass of Factor 6 while formic acid, formaldehyde, acetone and ethanol together contributed ~ 15 % (~ 5 %, ~ 4 % and ~ 3 %, respectively) to the total measured NMVOC mass of this factor. The other 31 species contributed only ~ 18 %. The diel profile (Figure 8) of this factor correlates best with the increase in rates of temperature (dT/dt , $R^2 = 0.41$) and solar radiation (dSR/dt , $R^2 = 0.38$) during the daytime hours (between 06:00 – 17:00 LT; as can be seen in Table S3 of the supplementary information). Factor 6 showed a sharp peak directly after sunrise between 08:00 – 10:00 LT. This time coincides with the maximum increase in both temperature and solar radiation. Average mass loadings of ~ 45 $\mu\text{g m}^{-3}$ were observed during this period. The sharp peaks observed in this factor during morning hours could be explained by the Kathmandu Valley meteorology. After sunrise when air temperatures start to rise, the boundary layer continues to be shallow until direct sunlight reaches the Valley bottom. The accumulation of compounds in a shallow boundary layer contributes to high ambient concentrations. The dilution due to the rising boundary layer and daytime westerly winds in the Valley reduces the concentrations subsequently. Therefore, this factor is attributed as solvent evaporation.

3.1.7 Factor 7 - Mixed daytime

Formic acid and acetic acid contributed most to the total measured NMVOC mass of Factor 7 (~ 25 % and ~ 13 %, respectively) while propyne, methanol and acetone together contributed ~ 26 %




Figure 9. Timeseries and diel box and whisker plot for Factor 7 (Mixed daytime)

(~ 10 %, ~ 8 % and ~ 8 %, respectively). The other 32 species collectively contributed ~ 36 % to this factor but their individual contributions were ≤ 5 %. Like factor 6, this factor, too, has a predominance of oxygenated compounds (that could be due to photooxidation) with a minor contribution from NMVOCs such as acetonitrile and propyne which can be emitted from primary emission sources such as biomass burning and industrial emissions (Hao et al., 1996; Andreae and Merlet , 2001; Akagi et al., 2011). The diel profile of this factor (Figure 9) is similar to that of the ambient temperature and solar radiation with an average mass concentration of $\sim 20 \mu\text{g m}^{-3}$ between 12:00 - 14:00 LT.

Approximately 41 % of the total formamide, ~ 37 % of the total acetamide and ~ 40 % of the total isocyanic acid are explained by this factor. Both formamide and acetamide can be produced by hydroxyl radical initiated photooxidation of primary amines (such as methyl amine) and in turn can photochemically form isocyanic acid through hydroxyl radical mediated oxidation (Roberts et al., 2014; Ge et al., 2011; Sarkar et al., 2016). In addition 34 % of the formic acid and 23 % of the formaldehyde mass was explained by this factor. The timeseries (Figure 9) of this factor showed higher baseline concentrations during second part of the measurement period when primary emissions were higher due to both biomass burning and biomass co-fired brick kiln emissions as described in Sarkar et al. (2016). During this period, influenced strongly by biomass burning sources, specific NMVOCs such as isocyanic acid, formamide and acetamide showed enhancement in their background concentrations. This is likely due to the higher emissions of precursor alkyl amines and other N-containing compounds from the incomplete combustion of biomass (Stockwell et al., 2015) which can form formamide and acetamide via photooxidation. Due to the contribution from both photooxidation and primary emissions, this factor was attributed as the mixed daytime factor.

3.1.8 Factor 8 - Biogenic emissions

Factor 8 explains more of the total isoprene mass than any of the other factors (~ 33 %) and shows a distinct daytime peak with the highest mass loadings of $\sim 32 \mu\text{g m}^{-3}$ observed between 11:00 - 12:00 LT (Figure 10). The diel profile (Figure 10) of this factor correlates best with solar radiation




Figure 10. Timeseries and diel box and whisker plot for Factor 8 (Biogenic emissions)

($R^2 = 0.33$; as can be seen in Table S3 of the supplementary information) during the daytime hours (between 06:00 - 17:00 LT). Average nighttime concentrations of this factor were always less than $10 \mu\text{g m}^{-3}$. The timeseries profile showed very high daytime mass loadings up to $\sim 80 \mu\text{g m}^{-3}$ for the first part of the campaign (19 December 2012 – 2 January 2013) and lower mass loadings as the campaign progressed. This is also consistent with the observation of deciduous trees in the Kathmandu Valley shedding their leaves during peak winter (Sarkar et al., 2016). Therefore, the factor was attributed to biogenic emissions.

Most of the total measured NMVOC mass in this factor was associated with oxygenated NMVOCs namely acetaldehyde, acetic acid, acetone and formic acid which contributed $\sim 21\%$, $\sim 15\%$, $\sim 11\%$ and $\sim 10\%$, respectively to Factor 8. Isoprene contributed $\sim 8\%$ to the total NMVOC mass. The other 32 NMVOCs together contributed $\sim 35\%$.

To summarize, based on the characteristics observed in the factor profiles, factor timeseries and diel plots, Factor 1 was attributed to traffic (TR), Factor 2 was attributed to residential biofuel use and waste disposal (RB+WD), Factor 3 was attributed to mixed industrial emissions (MI), Factor 4 was attributed to biomass co-fired brick kilns (BK), Factor 5 to unresolved industrial emissions (UI), Factor 6 was attributed to solvent evaporation (SE), Factor 7 was attributed to mixed daytime source (MD) and Factor 8 was attributed to biogenic NMVOC emissions (BG). Table S3 of the supplementary information shows the calculated correlation coefficients between the PMF resolved source factors and the independent meteorological parameters.

It can be seen from Table S3 of the supplementary information that during daytime, the solvent evaporation (SE) factor correlated best with the rate of change in solar radiation and the rate of change in ambient temperature ($r = 0.62$ and 0.64 , respectively). This supports the assignment of the solvent evaporation factor as evaporation depends on temperature. However, the change of the saturation vapor pressure for a temperature change from 5°C to 20°C for the dominant compounds (acetaldehyde and acetic acid) is small (less than a factor of 1.3; Betterton and Hoffmann (1988); Johnson et al. (1996)) and, therefore, does not account for the observed magnitude of increase (by a factor of ~ 5) from 06:00 - 09:00 LT. Instead, the temperature dependence of the solubility of



these compounds in an aqueous solution (factor 5-7) would explain a change of this magnitude. The
545 solvent evaporation factor strongly anti-correlated with RH during the nighttime and correlated well
with the unresolved industrial (UI) factor ($r = 0.55$) during daytime. It is, therefore, possible that the
sources of the solvent evaporation and unresolved industrial emission may be identical or at least
spatially co-located. We hypothesize, that compounds, that firstly display a significant solubility in
aqueous solution and secondly a strong temperature dependence of the solubility are attributed to this
550 separate factor. At night, soluble compounds partition into the aqueous phase of the fog aerosol and
hence their mixing ratios will not build up in the nocturnal boundary layer to the same extent as those
of less soluble compounds despite continuous emissions from industrial units. Those compounds
with a high temperature dependence on solubility like acids and aldehydes will rapidly shift to the
gas phase from their nocturnal fog water reservoir when temperatures increase in the morning and
555 their solubility decreases, which manifests itself in a disproportionate (considering only evaporation)
increase of their mixing ratios at that time.

The mixed daytime factor (MD) correlated with solar radiation, ambient temperature and wind
speed ($r = 0.58$, 0.74 and 0.57 , respectively). The biogenic factor (BG) had the best correlation
with solar radiation ($r = 0.57$) during daytime, consistent with its attribution to biogenic emissions.
560 During daytime, the mixed industrial emissions and biomass co-fired brick kiln emissions had very
low mass concentration due to the boundary layer dilution and ventilation effect of high westerly
winds in the Kathmandu Valley (Sarkar et al., 2016). The ambient RH was also lower during the
daytime. Therefore, both the mixed industrial emissions and brick kilns emission showed positive
correlations with ambient RH ($r = 0.65$ and 0.74 , respectively). During nighttime, no significant
565 correlation was observed between the PMF resolved factors except the correlation of the biogenic
factor with the residential biofuel use and waste disposal (RB+WD) factor ($r = 0.58$) which indicates
that the high emissions of oxygenated NMVOCs and isoprene from RB+WD sources could result in
a minor mis-attribution of the combustion derived emissions to the biogenic factor.

3.2 Conditional probability functions (CPF) to determine source directionality

570 Figure 11 shows the Conditional Probability Function (CPF) plots that were used to examine the spa-
tial profile of the eight different PMF source factors. For the CPF plots, only data with wind speed
 $> 0.5 \text{ ms}^{-1}$ were considered. Six factors namely traffic, residential biofuel use and waste disposal,
mixed industrial emissions, unresolved industrial emissions, solvent evaporation and biomass co-
fired brick kilns could be associated clearly with anthropogenic activities and are, therefore, likely to
575 be impacted by spatially fixed sources, while one factor (mixed daytime) was related to photochem-
istry. One factor, biogenic emissions, is natural but can also be attributed to spatially fixed sources
such as forests.

The CPF plot for the traffic factor showed maximum conditional probability (0.4 - 0.7) from
the W-NW direction where the Kathmandu city center and the busiest traffic intersections were




Figure 11. Conditional probability functions (CPF) plots for all source factors resolved by PMF showing wind directional dependency of different source categories

580 located. The conditional probability for the SW and NE wind direction ranged from 0.2-0.4. Two cities, namely Lalitpur (Patan) and Bhaktapur, respectively, are located upwind of the site in these directions. The lowest conditional probability was observed for the SE wind direction.

The residential biofuel use and waste disposal factor showed a high conditional probability of emissions exceeding the mean for air masses reaching the site from most wind directions (0.5 - 0.7 for NW-N, ~ 0.4 for N-NE and S-SW and 0.2 for E-S), indicating that this source is spatially distributed throughout the Kathmandu Valley. Only for the wind sector from SW-NW the conditional probability of this source is low. The reason for this low conditional probability is that every day in the afternoon, winds from the western mountain passes reach the receptor site. The same wind direction is extremely rare after sunset and during the early morning hours, when residential biofuel use and waste disposal mostly occur. Consequently, the conditional probability plot shows low conditional probabilities for this wind sector.

The mixed industrial emissions factor showed the highest conditional probability of air masses with above average mass loadings reaching the receptor site from the NE to SE wind sector ($p = 0.4-0.6$), where Bhaktapur industrial area is located within a distance of 3-4 km upwind of the receptor site. Conditional probabilities of 0.2-0.4 were observed for the NW wind direction where several industries are located.

For brick kilns the highest conditional probability was observed for air masses reaching the receptor site from the NE-SE ($p \sim 0.4$), which had several active brick kilns near the Bhaktapur Industrial Estate, which was ~ 4 km upwind of the receptor site.

600 It is interesting to note that the unresolved industrial emissions factor shows a clear directional dependence ($p = 0.5-0.7$ for the NE-SW wind sector) indicating that this factor, too, can be attributed



to spatially fixed sources in Bhaktapur Industrial Estate and Patan Industrial Estate. Polymer production, manufacturing industries for adhesives, paints and/or pharmaceuticals upwind of the site likely contributed towards the measured NMVOC mass of the unresolved industrial factor.

605 The solvent evaporation factor, too, shows high conditional probabilities for the SE-SW wind direction (Patan Industrial Estate) and low conditional probabilities for the NW-NE wind direction. The conditional probability function shows significant overlap with that of the unresolved industrial emissions factor. It therefore highlights the plausibility that solvent/chemical evaporation or emissions from industrial units are the primary source for this factor although the temperature changes
610 after sunrise drives the partitioning into the gas phase.

Within the bin of calm wind speeds ($< 0.5 \text{ ms}^{-1}$) the maximum conditional probabilities were observed for mixed industrial emissions, unresolved industrial emissions and brick kilns (0.25, 0.18 and 0.18, respectively) which indicates that emissions from these sources tended to accumulate in a shallow boundary layer during stagnant conditions in the Kathmandu Valley. Therefore, using taller
615 chimney stacks, at least for combustion sources, to prevent accumulation of emissions in a shallow boundary layer could potentially improve the air quality of the Valley during foggy nights.

The mixed daytime factor shows no obvious directional dependence for the conditional probability of recording values above the average at the receptor site ($p > 0.3$ for all directions). Slightly higher conditional probabilities ($p \sim 0.6$) are recorded for air masses reaching the receptor site from the
620 N-NE and S-SW wind direction.

The biogenic factor showed high conditional probabilities for air masses reaching the receptor site from the SW to N direction ($p = 0.5$ to 1) where few forested areas such as Nilbarahi jungle and Gokarna forest were located. Also forested areas in mountain slopes in the SW and NW direction and the midday fetch region being frequently from this sector explains the directional dependency
625 of the biogenic factor.

The CPF analysis of the PMF model output clearly indicates that spatially fixed sources are responsible for a significant fraction of the overall measured NMVOC mass loadings and opens up the possibility to identify and mitigate emissions or at least the build-up of pollutants in a shallow inversion.

630 3.3 Source contribution to the total measured NMVOC mass loading and comparison with emission inventories

Figure 12 shows a pie chart summarizing contributions of individual sources to the total measured NMVOC mass loading. Total measured NMVOC mass loading was calculated by summing up the concentrations of individual measured NMVOCs (in $\mu\text{g m}^{-3}$). The distribution shows that biogenic
635 sources and the mixed daytime factor contributed only 10% and 9.2%, respectively, to the total measured NMVOC mass loading while all the anthropogenic sources collectively contributed $\sim 80\%$ to the total measured NMVOC mass loading.




Figure 12. Contributions of various sources to the total NMVOC mass loading observed at Bode, a semi-urban site in the Kathmandu Valley

According to two widely used emission inventories, namely REAS v2.1 (Regional Emission inventory in ASia) and EDGAR v4.2 (Emissions Database for Global Atmospheric Research) (Kurokawa et al., 2013; Olivier et al., 1994) and the existing Nepalese inventory obtained from the International Centre for Integrated Mountain Development's (ICIMOD) database, residential biofuel use is considered to be the pre-dominant source of anthropogenic NMVOC emissions in Nepal. When the analysis is spatially restricted to the Kathmandu Valley for those inventories that provide gridded emissions (as shown in Figure 13), differences between EDGAR v4.2 and REAS v2.1 appear.

The EDGAR v4.2 inventory (for the year 2008) attributes only 10.6% of the total anthropogenic NMVOC emissions in the Kathmandu Valley (85.2-85.5 Longitude and 27.6-27.8 Latitude) to be due to residential biofuel use and an additional 8.9% to solid waste disposal. These numbers are in reasonable agreement with our PMF output, which attributes 13.5% instead of 19.5% of the total measured NMVOC mass to these two sources combined.

The REAS v2.1 inventory (for the year 2008) estimates that 67.2% of the total wintertime (December and January) anthropogenic NMVOC emissions in the Kathmandu Valley (85.25-85.5 Longitude and 27.5-27.75 Latitude) originate from residential and commercial biofuel use — a significant overestimation when the numbers are compared to our PMF output and the EDGAR v4.1 inventory. The national Nepali emission inventory, too apportions a large share of the total national annual NMVOCs emissions to residential and commercial biofuel use (83.1%). It, therefore, appears, that while apportioning the emissions spatially, the REAS v2.1 emission inventory does not fully account for the socio-economic differences between rural and urban areas. The EDGAR v4.2 emission inventory, on the other hand, seems to apportion most of the national consumption of LPG cooking gas to the highly urbanised Kathmandu valley and correspondingly scales down the emission from biofuel use within the Kathmandu valley. In absolute terms the annual NMVOC emissions



attributed to domestic fuel usage within the Kathmandu valley by EDGAR v4.2 are a factor of 3.6 lower compared to the annual NMVOC emissions attributed to this sector by REAS v2.1.

The EDGAR inventory considers solvent use (66 %) and mixed industrial emissions to represent the second most important source of NMVOCs. Solvent use and other industrial emissions (8.5 %) combined account for 74.5 %. Collectively they are considered to contribute $\sim 10\%$ to the total anthropogenic NMVOC mass in the EDGAR v4.2 inventory, while our PMF results attribute 52.8 % of the measured NMVOCs to solvent use and industrial emissions combined. It should be noted, that solvent use and other factors related to industrial emissions (mixed industrial and unresolved industrial) must be combined while comparing our PMF output with emission inventories. Both the mixed industrial emission factor and the unresolved industrial emission factor contain a significant NMVOC mass fraction from industrial solvent use, but also combustion related emissions from industrial units. Unfortunately, industrial solvent use and industrial combustion emissions from co-located units cannot be cleanly segregated using the PMF model, which relies on spatio-temporal patterns while building factor profiles. Overall, our PMF output agrees with the EDGAR v4.2 inventory, that industries are the dominant source of NMVOCs in the Kathmandu valley. According to the REAS v2.1 inventory, solvent use is considered to be the second most dominant contributor (29.8 %) to wintertime NMVOC emissions in the Kathmandu valley. Solvents and other industrial emissions (0.9 %) combined account for 30.7 % of the total wintertime NMVOC emissions in the REAS v2.1 emission inventory. Since, most of the national consumption of solvents and a significant share of Nepal's industrial production is concentrated in the Kathmandu valley, the discrepancies between the REAS v2.1 emission inventory and our results indicate, that the REAS v2.1 emission inventory does not sufficiently account for the special status of the Kathmandu valley while spatially apportioning emissions. The emissions that EDGAR v4.2 attributed to solid waste disposal, industries, the transport sector, and solvent use within the Kathmandu valley are a factor of 17.4, 14.0, 7.4 and 3.3 times higher compared to what the REAS v2.1 inventory attributes to the same sectors for the same geographical area.

The annual Nepalese inventory (for the year 2000) considers solvent and paint use to be the second largest contributor to the anthropogenic NMVOC emissions in Nepal, while industries are considered to make an insignificant overall contribution (0.7 %). These numbers cannot be compared to our results in a meaningful manner, as the national emissions in particular for sectors such as domestic fuel usage and agricultural waste burning may be dominated by the rural hinterland, while our PMF results apply to the largest urban agglomeration in Nepal.

Traffic was considered to contribute only between $\sim 1.3\%$ (in the REAS v2.1 inventory) to a maximum of $\sim 2.6\%$ (in EDGAR v4.2 inventory) of the total anthropogenic NMVOC emissions in the Kathmandu valley. This stands in stark contrast to the results of our PMF analyses, which indicate traffic contributes ca. 20 %, solvent evaporation and industrial solvent/chemical usage accounts for ca. 36 % (unresolved industrial emissions + solvent evaporation) and other industrial emissions




Figure 13. Comparison of the PMF derived contribution of anthropogenic sources with NMVOCs source contribution according to the existing Nepalese, REAS and EDGAR emission inventory

(mixed industrial emissions + brick kilns) account for ca. 30 of the total measured anthropogenic NMVOC mass loading in the Kathmandu valley. According to the recent study of the vehicle fleet in Kathmandu valley Shrestha et al. (2013), transport sector NMVOC emissions in the Kathmandu valley for the year 2010 amounted to 7654 t y^{-1} , a number that is 10 times higher than the number currently in the EDGAR v4.2 inventory and 72 times higher than the number currently in the REAS v2.1 inventory. If the emission estimate of (Shrestha et al., 2013) was incorporated into EDGAR v4.2 inventory without any further changes, the percentage share of transport sector emissions to the total NMVOC emissions would increase to 38.7%, while the contribution of domestic fuel usage and waste disposal would drop to 12.7% (PMF 13.5%) and the contribution of industrial emissions and solvent use would drop to 48.6% (PMF 52.8%). Our PMF results, however, seem to suggest, that 2012 transport sector emissions have decreased by $\sim 50\%$ compared to the 2010 emissions presented in (Shrestha et al., 2013), possibly due to a reduction of the number older vehicles in the fleet.

Inefficient biomass co-fired brick kilns are a unique industrial source in the Kathmandu Valley, and contributed significantly ($\sim 15\%$) to the total measured anthropogenic NMVOC mass loading. The existing Nepalese inventory considers contributions of brick kilns only to the emission of particulate matter (PM_{10} and $\text{PM}_{2.5}$), while the two other emission inventories do not include emissions from brick kilns in the Kathmandu Valley at all. If transport sector NMVOC emissions of $\sim 3800 \text{ t y}^{-1}$ and an additional $\sim 2400 \text{ t y}^{-1}$ NMVOC emissions from brick kilns, were included in the EDGAR v4.2 emission inventory, the EGAR emission inventory and our PMF output would agree perfectly




Figure 14. Contribution of PMF derived source factors to acetonitrile and aromatic NMVOCs

(within $\pm 0.2\%$) on the relative contribution of all sources, without changing the contribution from any of the other sources.

720 Only two sources, domestic fuel usage (on account of the changed heating demand) and agricultural waste burning are expected to have significant seasonality. Jointly, they account for less than 10% of the total NMVOC emissions. Since cooking needs persist throughout the year and the decrease in agricultural waste burning outside harvest season may be partially offset by leaf-litter burning (a source currently not in the model), it is likely that the failure to account for seasonal effects imparts an uncertainty of less than 1% on the overall result of our analysis.

725 The REAS v2.1 emission inventory for the Kathmandu valley, on the other hand, seems to require large corrections. While our analysis of the REAS inventory was restricted to December and January, annual averages of individual sources differ by less than $\pm 10\%$ from the winter values. Therefore, the difference in the time window selected for the analysis cannot explain the observed discrepancies to the EDGAR emission inventory.

3.4 Source contribution to individual NMVOCs

Figure 14 represents the pie charts showing contribution of the eight source factors to individual NMVOCs such as acetonitrile, benzene, styrene, toluene, sum of C8-aromatics (xylenes and ethyl-



Table 6. Emission ratios of NMVOCs/benzene for acetonitrile and aromatic hydrocarbons derived from the PMF model for different sources and comparison with the ratios for different source categories reported in previous studies

ERs/Benzene	RB+WD	BK	MI	UI	Garbage burning grab samples	Waste burning ¹	Wood burning ²	Charcoal burning ²
Acetonitrile	0.23	0.14	0.25	0.36	0.77	0.06	-	-
Toluene	0.34	0.35	0.18	0.30	0.34	0.41	0.05	0.50
C8-aromatics	0.18	0.06	0.08	0.00	0.25	0.10	-	0.46
C9-aromatics	0.25	0.22	0.06	0.12	0.08	0.03	-	-
Styrene	0.12	0.09	0.09	0.04	0.16	0.86	-	-
Naphthalene	0.11	0.15	0.20	0.05	0.09	0.10	-	-

1. Stockwell et al. (2015); 2. Tsai et al. (2003); RB+WD = Residential biofuel use and waste disposal; BK = Biomass co-fired brick kilns; MI = Mixed industrial emissions; UI = Unresolved industrial emissions

benzene) and sum of C9-aromatics (trimethylbenzenes and propylbenzene). Maximum contribution
735 to the acetonitrile mass concentration was observed from the unresolved industrial emission sources
(~ 30 %) followed by the biomass co-fired brick kilns emission (~ 24 %) and mixed industrial emis-
sion (~ 20 %) factors. Residential biofuel use and waste disposal features only fourth (~ 18 %). The
same sources also contribute most to benzene emissions, indicating that fuel usage, rather than its
application as solvent/chemical reagents in industrial processes is responsible for most of the in-
740 dustrial acetonitrile emissions. It also indicates that industrial rather than residential biofuel usage
contributes more towards outdoor NMVOC air pollution. Most of the benzene (which is a human
carcinogen) can be attributed to biomass co-fired brick kilns (~ 37 %), mixed industrial (~ 17 %)
and unresolved industrial (~ 18 %) sources. Residential biofuel use again featured only fourth as
far as the contribution towards mixing ratios of this compound in the outdoor environment is con-
745 cerned. Table 6 shows a comparison of NMVOCs/benzene emission ratios for four PMF derived
sources (residential biofuel use and waste disposal, biomass co-fired brick kilns, mixed industrial
and unresolved industrial sources) to the emission ratios obtained from the grab samples collected
for garbage burning in the Kathmandu Valley and the previously reported emission ratios for waste
burning, wood burning and charcoal burning sources.

750 Residential biofuel use and waste disposal contributed ~ 28 % of the total styrene which were
emitted significantly from waste burning. However, traffic was found to be equally important as
a styrene source (~ 37 %) in the Kathmandu Valley. Recently, styrene has been detected from traffic
and was found to have high emission ratios with respect to benzene after cold startup of engines and
in LPG fuel (Alves et al., 2015). Biomass co-fired brick kilns and mixed industrial emissions also
755 contribute significantly (~ 21 % and ~ 14 %, respectively) towards styrene mass loadings. Traffic
was found to be the most important source of higher aromatics including toluene, C8-aromatics, and




Figure 15. Contribution of PMF derived sources to formamide, acetamide, isocyanic acid and formic acid

C9-aromatics (> 60 %). Biomass co-fired brick kilns were the second largest contributors towards their mass loadings, while residential biofuel usage and waste disposal ranked third.

Figure 15 shows the pie charts summarizing contributions of PMF derived sources to two newly
760 quantified compounds in the Kathmandu Valley, namely formamide and acetamide along with iso-
cyanic acid and formic acid. All these compounds showed maximum contribution from the mixed
daytime factor (~ 34 % to ~ 41 %) due to the photo-oxidation source. As discussed previously in
Sarkar et al. (2016) and in section 3.1.7, both formamide and acetamide are formed primarily as
a result of photooxidation of amine compounds and N-containing compounds. These can be emitted
765 from the various inefficient combustion processes in the Kathmandu Valley. Photooxidation of these
amides further forms isocyanic acid (reaction schematic is shown in Figure S5 of the supplemen-
tary information). Apart from the mixed daytime source, unresolved industrial emissions factor also
contributed significantly to all these compounds (~ 22 % to ~ 23 %) as they are used as reactants
(e.g. formic acid is used as reactant to produce formamide in industries) or produced during different
770 industrial processes (such as formamide is produced in pharmaceuticals and plastic industries). Sol-
vent evaporation factor contributed ~ 19 % to formamide while biogenic factor contributed ~ 14 %
to formic acid. Contributions from all the other sources to these NMVOCs were < 10 %.

Figure 16 represents the pie charts showing contribution of the eight sources derived from PMF
to 1,3-butadiene and oxygenated compounds namely methanol, acetone, acetaldehyde, ethanol and
775 acetic acid. It can be seen from Figure 16 that emissions of all these compounds in the Kath-
mandu Valley were dominated by different industrial activities. The total unresolved industrial emis-




Figure 16. Contribution of PMF derived sources to 1,3-butadiene and oxygenated NMVOCs such as methanol, acetone, acetaldehyde, ethanol and acetic acid

sions factor dominated the contribution to 1,3-butadiene (~ 48 %), methanol (~ 35 %) and acetone (~ 22 %). Residential biofuel use and waste disposal also contributed significantly to 1,3-butadiene (~ 21 %) and methanol (~ 16 %). Traffic was found to have significant contribution to acetone (~ 21 %). It is known that acetaldehyde, ethanol and acetic acid are used as solvents in different industries and it was found that industrial sources obtained from PMF (mixed industrial + unresolved industrial + solvent evaporation) together contributed ~ 72 % of the total acetaldehyde, 100 % of the total ethanol and ~ 47 % of the total acetic acid. Biogenic sources also had significant contribution to acetaldehyde and acetic acid (~ 17 % and ~ 14 %, respectively) whereas residential biofuel use and waste disposal contributed to ~ 15 % of the total acetic acid.

Figure 17 represents a timeseries of daily mean relative contribution of the PMF derived sources during SusKat-ABC campaign. As discussed in Sarkar et al. (2016), the whole campaign can be divided into three different periods based on the measurements – first period (from the start of the campaign until 3 January 2013) was associated with high daytime isoprene emissions due to strong biogenic emissions, the second period (4 – 18 January 2013) was marked by enhancements in acetonitrile and benzene concentrations due to the kick start of the biomass co-fired brick kilns in the




Figure 17. Daily mean relative contribution of PMF derived eight sources during SusKat-ABC campaign

Kathmandu Valley and in the third period (19 January until the end of the campaign), more oxygenated NMVOCs were observed which was believed to be due to the stable operation of the brick kilns and more contribution from the industrial sources. PMF derived results also supports these
795 observation as can be seen in Figure 17. It can be seen that from the start of the campaign until 3 January 2013 contribution of PMF derived biogenic sources were $> 20\%$ for most of the time while contribution from the brick kilns emission factor was negligible ($\leq 5\%$). From 4 January until 18 January 2013, the contribution of brick kilns increased significantly ($\sim 20\%$ to $\sim 40\%$) as almost all brick kilns in the Kathmandu Valley became operational. After 18 January until the end of the
800 campaign, the contribution of brick kilns become lower due to its stable operation.

During the first period, contribution of traffic was found to be higher ($\sim 20\%$ to $\sim 30\%$) compared to the rest of the campaign. The higher contribution of the mixed daytime source during the second and third part of the campaign was due to the early morning and daytime photooxidation of the precursor compounds which were emitted as a result of biomass co-fired brick kilns and other
805 biomass burning emissions during these periods. The mixed industrial emissions factor contributed almost equally throughout the campaign (contributing $\sim 10\%$ to $\sim 15\%$) but the solvent evaporation and the unresolved industrial emissions factor contributed more during the second and third part of the campaign (increase of $\sim 10\%$).

3.5 Source contribution to daytime ozone production potential and SOA formation

810 Figure 18a represents the source contribution to daytime O_3 production potential while Figure 18b represents the contribution of different classes of compounds measured in the Kathmandu Valley to the daytime O_3 production potential as discussed in Sarkar et al. (2016). The daytime O_3 production potential for individual sources was calculated by summing up the O_3 production potential




Figure 18. Daytime O_3 production potential obtained a) from the source contribution using PMF and b) from the measurements performed in the Kathmandu Valley

for the individual compounds which was calculated according to the method described by Sinha
815 et al. (2012). The distribution of the daytime O_3 production potential obtained from the measure-
ments (Figure 18b) shows that $\sim 78\%$ of the total daytime O_3 production potential was due to
the contribution from isoprene and oxygenated NMVOCs which indicated dominance of biogenic
emissions and photochemistry in the Kathmandu Valley even in the winter. But the distribution of
different sources obtained from PMF to daytime O_3 production potential shows that the biogenic
820 factor together with the photochemistry factor (mixed daytime) contributed only $\sim 30\%$ of the total
 O_3 production potential. The remaining $\sim 70\%$ was contributed by anthropogenic sources. While
solvent evaporation contributed most ($\sim 20\%$) to the total daytime O_3 production potential, traffic
and unresolved industrial emission stood second and third, respectively, in terms of anthropogenic
ozone precursor emissions. Residential biofuel use and waste disposal, and biomass co-fired brick
825 kilns while potentially important from a human health perspective, contributed only a minor fraction
of the total anthropogenically emitted ozone precursors.

The consequence of including only a subset of NMVOCs is an underestimation of the OH re-
activity and hence ozone production potential, which scales directly with the OH reactivity. Based
on measured methane and 63 non methane hydrocarbon measurements in the city of Lahore which
830 is much larger and by all indications more polluted than Kathmandu Barletta et al. (2016) the au-
thors reported a maximum contribution of about 14% due to all alkanes including methane to the
total measured OH reactivity. Despite high concentration abundances in urban atmospheric envi-
ronments, the rate constants of these species are typically 100 times lower than compounds like
isoprene, and hence their contribution to the total OH reactivity is much lower. For example, even
835 3 ppm methane (observed only in plumes) would contribute only $\sim 0.5\text{ s}^{-1}$ to the total OH re-
activity and hence make an insignificant contribution to the ozone production potential. Hence, our
analyses of the ozone production potential may underestimate the total ozone production potential
by 15–25%, if we can extrapolate the observations from another South Asian city like Lahore.

Secondary organic aerosol (SOA) production was calculated using the concentrations and the
840 known SOA yields for benzene, toluene, styrene, xylene, trimethylbenzenes, naphthalene and iso-




Figure 19. Contribution of PMF derived eight sources to the SOA formation in the Kathmandu Valley

prene (Ng et al., 2007; Chan et al., 2009; Yuan et al., 2013; Kroll et al., 2006). As the biomass co-fired brick kilns and the traffic factor contains most of the reactive aromatic compounds, they appeared to be the dominant contributors to SOA production (as shown in Figure 19) in the Kathmandu Valley.

845 4 Conclusions

The PMF model results reveal several new results regarding the source apportionment of NMVOCs in the Kathmandu Valley. Speciation of NMVOCs in the emission inventory for Nepal only includes compound classes (e.g. alkanes, alkenes etc.) and not specific compounds. Also, the existing emission inventories (e.g. REAS v2.1, EDGAR v4.2; Kurokawa et al. (2013); Olivier et al. (1994) and
850 Nepalese inventory (ICIMOD)) are highly uncertain as there has been no validation using in-situ measurements of these mostly bottom up inventories which rely on fuel and source emission factors measured in other technologically different regions of the world (primarily the US and Europe). By using the specific NMVOC emission tracer data measured in the Kathmandu Valley and constraining the PMF with measured source profiles of complex sources (e.g. biomass co-fired brick kilns, residential solid biofuel use and waste disposal), it is shown that the contribution from sources such as
855 residential solid biofuel use and waste disposal is overestimated in the REAS v2.1 emission inventory. At the same time, the emissions from industrial sources are underestimated. Both REAS v2.1 and EDGAR v4.2 underestimate the contribution of traffic and do not include brick kiln emissions. The presence of elevated concentrations of several health relevant NMVOCs (e.g. benzene) could be
860 attributed to the biomass co-fired brick kiln sources.

This study has provided quantitative information regarding the contributions of the major NMVOC sources in the Kathmandu Valley. This will enable focused mitigation efforts by policy makers and practitioners to improve the air quality of the Kathmandu Valley by reducing emissions of both toxic NMVOCs and formation of secondary pollutants. The results will also enable significant improve-



865 ments in existing NMVOC emission inventories so that chemical-transport models can be parameter-
ized more accurately over the South Asian region and the air quality-climate predictions by models
can become more reliable.

Authors' contributions

870 Sections of this study were submitted in part fulfilment of the PhD work of C.S. carried out under
the supervision of V.S. at IISER Mohali. The VOC dataset QA/QC and analyses were performed
by C.S. and V.S. whereas B.S. designed and set up the PMF model and ensured QA/QC of PMF
output which was performed by C.S. A.P. helped with interpretation of PMF results and suggested
grab sampling experiments at an early stage. C.S., V.S. and B.S. wrote the paper and all co-authors
discussed the results and commented on the paper.

875 *Acknowledgements.* Chinmoy Sarkar and Vinayak Sinha acknowledge the support extended by the Founding
Director of IISER Mohali, Professor N. Sathyamurthy to enable participation of the IISER Mohali team in the
SusKat-ABC campaign. Chinmoy Sarkar acknowledges the Ministry of Human Resources and Development
(MHRD), India and IASS Potsdam, Germany for funding with a service contract. IASS Potsdam funded the
deployment of the PTR-TOF-MS by the IISER Mohali team in Kathmandu and local logistical support was
880 provided by Khadak S. Mahata, Dipesh Rupakheti, Bhogendra Kathayat at the Bode site.

This study was partially supported by core funds of ICIMOD contributed by the governments of Afghanistan,
Australia, Austria, Bangladesh, Bhutan, China, India, Myanmar, Nepal, Norway, Pakistan, Switzerland, and the
United Kingdom.

All the data reported in this article can be obtained from the corresponding author by sending an email to
885 vsinha@iisermohali.ac.in.



References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039-4072, doi:10.5194/acp-11-4039-2011, 2011.
- 890 Alves, C. A., Lopes, D. J., Calvo, A. I., Evtyugina, M., Rocha, S., and Nunes, T.: Emissions from Light-Duty Diesel and Gasoline in-use Vehicles Measured on Chassis Dynamometer Test Cycles, *Aerosol and Air Quality Research*, doi:10.4209/aaqr.2014.01.0006, 2015.
- Anderson, M. J., Daly, E. P., Miller, S. L., and Milford, J. B.: Source apportionment of exposures to volatile organic compounds: II. Application of receptor models to TEAM study data, *Atmospheric Environment*, 36, 895 3643-3658, doi:10.1016/S1352-2310(02)00280-7, 2002.
- Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochemical Cycles*, 15, 955-966, doi:10.1029/2000gb001382, 2001.
- Barletta, B., Simpson, I. J., Blake, N. J., Meinardi, S., Emmons, L. K., Aburizaiza, O. S., Siddique, A., Zeb, J., Yu, L. E., Khwaja, H. A., Farrukh, M. A. and Blake, D. R.: Characterization of carbon monoxide, methane and nonmethane hydrocarbons in emerging cities of Saudi Arabia and Pakistan and in Singapore, *J. Atmos.* 900 *Chem.*, 11, 2399-2421, doi:10.1007/s10874-016-9343-7, 2016.
- Betterton, E. A., and Hoffmann, M. R.: Henry's law constants of some environmentally important aldehydes, *Environmental Science & Technology*, 22, 1415-1418, doi:10.1021/es00177a004, 1988.
- Bon, D. M., Ulbrich, I. M., de Gouw, J. A., Warneke, C., Kuster, W. C., Alexander, M. L., Baker, A., Beyersdorf, A. J., Blake, D., Fall, R., Jimenez, J. L., Herndon, S. C., Huey, L. G., Knighton, W. B., Ortega, J., 905 Springston, S., and Vargas, O.: Measurements of volatile organic compounds at a suburban ground site (T1) in Mexico City during the MILAGRO 2006 campaign: measurement comparison, emission ratios, and source attribution, *Atmos. Chem. Phys.*, 11, 2399-2421, doi:10.5194/acp-11-2399-2011, 2011.
- Borbon, A., Fontaine, H., Veillerot, M., Locoge, N., Galloo, J. C., and Guillermo, R.: An investigation into 910 the traffic-related fraction of isoprene at an urban location, *Atmospheric Environment*, 35, 3749-3760, doi:10.1016/S1352-2310(01)00170-4, 2001.
- Borbon, A., Gilman, J. B., Kuster, W. C., Grand, N., Chevaillier, S., Colomb, A., Dolgorouky, C., Gros, V., Lopez, M., Sarda-Esteve, R., Holloway, J., Stutz, J., Petetin, H., McKeen, S., Beekmann, M., Warneke, C., Parrish, D. D., and de Gouw, J. A.: Emission ratios of anthropogenic volatile organic compounds in northern 915 mid-latitude megacities: Observations versus emission inventories in Los Angeles and Paris, *Journal of Geophysical Research: Atmospheres*, 118, 2041-2057, doi:10.1002/jgrd.50059, 2013.
- Brown, S. G., Frankel, A., and Hafner, H. R.: Source apportionment of VOCs in the Los Angeles area using positive matrix factorization, *Atmospheric Environment*, 41, 227-237, doi:10.1016/j.atmosenv.2006.08.021, 2007.
- 920 Brown, S. G., Eberly, S., Paatero, P., and Norris, G. A.: Methods for estimating uncertainty in PMF solutions: Examples with ambient air and water quality data and guidance on reporting PMF results, *Science of The Total Environment*, 518-519, 626-635, doi:10.1016/j.scitotenv.2015.01.022, 2015.
- Buzzu, B., and Fraser, M. P.: Source identification and apportionment of volatile organic compounds in Houston, TX, *Atmospheric Environment*, 40, 2385-2400, doi:10.1016/j.atmosenv.2005.12.020, 2006.



- 925 Chan, A. W. H., Kautzman, K. E., Chhabra, P. S., Surratt, J. D., Chan, M. N., Crouse, J. D., Kürten, A., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from photooxidation of naphthalene and alkylnaphthalenes: implications for oxidation of intermediate volatility organic compounds (IVOCs), *Atmos. Chem. Phys.*, 9, 3049-3060, doi:10.5194/acp-9-3049-2009, 2009.
- Chen, W. T., Shao, M., Lu, S. H., Wang, M., Zeng, L. M., Yuan, B., and Liu, Y.: Understanding primary and secondary sources of ambient carbonyl compounds in Beijing using the PMF model, *Atmos. Chem. Phys.*, 14, 3047-3062, 10.5194/acp-14-3047-2014, 2014.
- 930 Crippa, M., Canonaco, F., Slowik, J. G., El Haddad, I., DeCarlo, P. F., Mohr, C., Heringa, M. F., Chirico, R., Marchand, N., Temime-Roussel, B., Abidi, E., Poulain, L., Wiedensohler, A., Baltensperger, U., and Prevot, A. S. H.: Primary and secondary organic aerosol origin by combined gas-particle phase source apportionment, *Atmos. Chem. Phys.*, 13, 8411-8426, 10.5194/acp-13-8411-2013, 2013.
- Fleming, Z. L., Monks, P. S., and Manning, A. J.: Review: Untangling the influence of air-mass history in interpreting observed atmospheric composition, *Atmospheric Research*, 104-105, 1-39, doi:10.1016/j.atmosres.2011.09.009, 2012.
- Gaimoz, C., Sauvage, S., Gros, V., Herrmann, F., Williams, J., Locoge, N., Perrussel, O., Bonsang, B., d'Ár-gouges, O., Sarda-Estéve, R., and Sciare, J.: Volatile organic compounds sources in Paris in spring 2007. Part II: source apportionment using positive matrix factorisation, *Environmental Chemistry*, 8, 91-103, doi:dx.doi.org/10.1071/EN10067, 2011.
- 940 Ge, X., Wexler, A. S., and Clegg, S. L.: Atmospheric amines - Part I. A review, *Atmospheric Environment*, 45, 524-546, doi:10.1016/j.atmosenv.2010.10.012, 2011.
- 945 Guo, H., Wang, T., and Louie, P. K. K.: Source apportionment of ambient non-methane hydrocarbons in Hong Kong: Application of a principal component analysis/absolute principal component scores (PCA/APCS) receptor model, *Environmental Pollution*, 129, 489-498, doi:10.1016/j.envpol.2003.11.006, 2004.
- Guo, H., Wang, T., Blake, D. R., Simpson, I. J., Kwok, Y. H., and Li, Y. S.: Regional and local contributions to ambient non-methane volatile organic compounds at a polluted rural/coastal site in Pearl River Delta, China, *Atmospheric Environment*, 40, 2345-2359, doi:10.1016/j.atmosenv.2005.12.011, 2006.
- 950 Hao, W. M., Ward, D. E., Olbu, G., and Baker, S. P.: Emissions of CO₂, CO, and hydrocarbons from fires in diverse African savanna ecosystems, *Journal of Geophysical Research: Atmospheres*, 101, 23577-23584, doi:10.1029/95JD02198, 1996.
- Hellén, H., Tykkä, T., and Hakola, H.: Importance of monoterpenes and isoprene in urban air in northern Europe, *Atmospheric Environment*, 59, 59-66, doi:10.1016/j.atmosenv.2012.04.049, 2012.
- 955 Hewitt, C. N.: *Reactive Hydrocarbons in the Atmosphere*, Academic Press, ISBN: 978-0-12-346240-4, San Diego, 1999.
- Ho, K. F., Lee, S. C., Ho, W. K., Blake, D. R., Cheng, Y., Li, Y. S., Ho, S. S. H., Fung, K., Louie, P. K. K., and Park, D.: Vehicular emission of volatile organic compounds (VOCs) from a tunnel study in Hong Kong, *Atmos. Chem. Phys.*, 9, 7491-7504, doi:10.5194/acp-9-7491-2009, 2009.
- 960 Hwa, M. Y., Hsieh, C. C., Wu, T. C., and Chang, L. F. W.: Real-world vehicle emissions and VOCs profile in the Taipei tunnel located at Taiwan Taipei area, *Atmospheric Environment*, 36, 1993-2002, doi:10.1016/S1352-2310(02)00148-6, 2002.



- Inomata, S., Tanimoto, H., Fujitani, Y., Sekimoto, K., Sato, K., Fushimi, A., Yamada, H., Hori, S., Kuzumazawa, Y., Shimono, A., and Hikida, T.: On-line measurements of gaseous nitro-organic compounds in diesel vehicle exhaust by proton-transfer-reaction mass spectrometry, *Atmospheric Environment*, 73, 195-203, doi:10.1016/j.atmosenv.2013.03.035, 2013.
- Inomata, S., Fujitani, Y., Fushimi, A., Tanimoto, H., Sekimoto, K., and Yamada, H.: Field measurement of nitromethane from automotive emissions at a busy intersection using proton-transfer-reaction mass spectrometry, *Atmospheric Environment*, 96, 301-309, doi:10.1016/j.atmosenv.2014.07.058, 2014.
- IPCC: Impacts, Adaptation and Vulnerability : Working Group II Contribution to the Intergovernmental Panel on Climate Change : Fifth Assessment Report (AR5): Summary for Policymakers. , Intergovernmental Panel on Climate Change. Working Group Impacts, 2013.
- Johnson, B., Berton, E., and Craig, D.: Henry's law coefficients of formic and acetic acids, *Journal of Atmospheric Chemistry*, 24, 113-119, doi:10.1007/bf00162406, 1996.
- Jorquera, H., and Rappenglück, B.: Receptor modeling of ambient VOC at Santiago, Chile, *Atmospheric Environment*, 38, 4243-4263, doi:10.1016/j.atmosenv.2004.04.030, 2004.
- Kaltonoudis, C., Kostenidou, E., Florou, K., Psichoudaki, M., and Pandis, S. N.: Temporal variability and sources of VOCs in urban areas of Eastern Mediterranean, *Atmos. Chem. Phys.*, 16, 14825-14842, doi:10.5194/acp-16-14825-2016, 2016.
- Karl, T., Jobson, T., Kuster, W. C., Williams, E., Stutz, J., Shetter, R., Hall, S. R., Goldan, P., Fehsenfeld, F., and Lindinger, W.: Use of proton-transfer-reaction mass spectrometry to characterize volatile organic compound sources at the La Porte super site during the Texas Air Quality Study 2000, *Journal of Geophysical Research: Atmospheres*, 108, 4508, doi:10.1029/2002jd003333, 2003.
- Kim, E., Brown, S. G., Hafner, H. R., and Hopke, P. K.: Characterization of non-methane volatile organic compounds sources in Houston during 2001 using positive matrix factorization, *Atmospheric Environment*, 39, 5934-5946, doi:10.1016/j.atmosenv.2005.06.045, 2005.
- Kim, K. H., Hong, Y. J., Pal, R., Jeon, E. C., Koo, Y. S., and Sunwoo, Y.: Investigation of carbonyl compounds in air from various industrial emission sources, *Chemosphere*, 70, 807-820, doi:10.1016/j.chemosphere.2007.07.025, 2008.
- Kristensson, A., Johansson, C., Westerholm, R., Swietlicki, E., Gidhagen, L., Wideqvist, U., and Vesely, V.: Real-world traffic emission factors of gases and particles measured in a road tunnel in Stockholm, Sweden, *Atmospheric Environment*, 38, 657-673, doi:10.1016/j.atmosenv.2003.10.030, 2004.
- Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol Formation from Isoprene Photooxidation, *Environmental Science & Technology*, 40, 1869-1877, doi:10.1021/es0524301, 2006.
- Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058, doi:10.5194/acp-13-11019-2013, 2013.
- Lemieux, P. M., Lutes, C. C., and Santoianni, D. A.: Emissions of organic air toxics from open burning: a comprehensive review, *Progress in Energy and Combustion Science*, 30, 1-32, doi:10.1016/j.pecs.2003.08.001, 2004.



- Leuchner, M., and Rappenglück, B.: VOC source-receptor relationships in Houston during TexAQS-II, *Atmospheric Environment*, 44, 4056-4067, doi:10.1016/j.atmosenv.2009.02.029, 2010.
- 1005 Miller, S. L., Anderson, M. J., Daly, E. P., and Milford, J. B.: Source apportionment of exposures to volatile organic compounds. I. Evaluation of receptor models using simulated exposure data, *Atmospheric Environment*, 36, 3629-3641, doi:10.1016/S1352-2310(02)00279-0, 2002.
- Misztal, P. K., Hewitt, C. N., Wildt, J., Blande, J. D., Eller, A. S. D., Fares, S., Gentner, D. R., Gilman, J. B., Graus, M., Greenberg, J., Guenther, A. B., Hansel, A., Harley, P., Huang, M., Jardine, K., Karl, T., Kaser, L., Keutsch, F. N., Kiendler-Scharr, A., Kleist, E., Lerner, B. M., Li, T., Mak, J., Nölscher, A. C., Schnitzhofer, R., Sinha, V., Thornton, B., Warneke, C., Wegener, F., Werner, C., Williams, J., Worton, D. R., Yassaa, N., and Goldstein, A. H.: Atmospheric benzenoid emissions from plants rival those from fossil fuels, *Scientific Reports*, 5, 12064, doi:10.1038/srep12064, 2015.
- 1010 Misztal, P. K., Hewitt, C. N., Wildt, J., Blande, J. D., Eller, A. S. D., Fares, S., Gentner, D. R., Gilman, J. B., Graus, M., Greenberg, J., Guenther, A. B., Hansel, A., Harley, P., Huang, M., Jardine, K., Karl, T., Kaser, L., Keutsch, F. N., Kiendler-Scharr, A., Kleist, E., Lerner, B. M., Li, T., Mak, J., Nölscher, A. C., Schnitzhofer, R., Sinha, V., Thornton, B., Warneke, C., Wegener, F., Werner, C., Williams, J., Worton, D. R., Yassaa, N., and Goldstein, A. H.: Atmospheric benzenoid emissions from plants rival those from fossil fuels, *Scientific Reports*, 5, 12064, doi:10.1038/srep12064, 2015.
- 1015 Morino, Y., Ohara, T., Yokouchi, Y., and Ooki, A.: Comprehensive source apportionment of volatile organic compounds using observational data, two receptor models, and an emission inventory in Tokyo metropolitan area, *Journal of Geophysical Research: Atmospheres*, 116, doi:10.1029/2010jd014762, 2011.
- Na, K., and Pyo Kim, Y.: Chemical mass balance receptor model applied to ambient C2-C9 VOC concentration in Seoul, Korea: Effect of chemical reaction losses, *Atmospheric Environment*, 41, 6715-6728, doi:10.1016/j.atmosenv.2007.04.054, 2007.
- 1020 Nakashima, Y., Kamei, N., Kobayashi, S., and Kajii, Y.: Total OH reactivity and VOC analyses for gasoline vehicular exhaust with a chassis dynamometer, *Atmospheric Environment*, 44, 468-475, doi:10.1016/j.atmosenv.2009.11.006, 2010.
- Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem. Phys.*, 7, 3909-3922, doi:10.5194/acp-7-3909-2007, 2007.
- 1025 Norris, G., Vedantham, R., Wade, K., Zahn, P., Brown, S., Paatero, P., Eberly, S., and Foley, C.: Guidance Document for PMF Applications with the Multilinear Engine, EPA 600/R-09/032, 2009.
- Olivier, J. G. J., Bouwman, A. F., van der Maas, C. W. M., and Berdowski, J. J. M.: Emission Database for Global Atmospheric Research (EDGAR), in: *Non-CO₂ Greenhouse Gases: Why and How to Control?*, edited by: van Ham, J., Janssen, L. J. H. M., and Swart, R. J., Springer Netherlands, 93-106, 1994.
- 1030 Olson, D. A., Norris, G. A., Seila, R. L., Landis, M. S., and Vette, A. F.: Chemical characterization of volatile organic compounds near the World Trade Center: Ambient concentrations and source apportionment, *Atmospheric Environment*, 41, 5673-5683, doi:10.1016/j.atmosenv.2007.02.047, 2007.
- 1035 Paatero, P., and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, 111-126, doi:10.1002/env.3170050203, 1994.
- Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometrics and Intelligent Laboratory Systems*, 37, 23-35, doi:10.1016/S0169-7439(96)00044-5, 1997.
- Paatero, P.: The Multilinear Engine-A Table-Driven, Least Squares Program for Solving Multilinear Problems, Including the n-Way Parallel Factor Analysis Model, *Journal of Computational and Graphical Statistics*, 8, 854-888, doi:10.1080/10618600.1999.10474853, 1999.
- 1040



- Paatero, P., Hopke, P. K., Song, X. H., and Ramadan, Z.: Understanding and controlling rotations in factor analytic models, *Chemometrics and Intelligent Laboratory Systems*, 60, 253-264, doi:10.1016/S0169-7439(01)00200-3, 2002.
- 1045 Paatero, P., and Hopke, P. K.: Rotational tools for factor analytic models, *Journal of Chemometrics*, 23, 91-100, doi:10.1002/cem.1197, 2009.
- Panday, A. K., Prinn, R. G., and Schär, C.: Diurnal cycle of air pollution in the Kathmandu Valley, Nepal: 2. Modeling results, *Journal of Geophysical Research: Atmospheres*, 114, D21308, doi:10.1029/2008jd009808, 2009.
- 1050 Pradhan, B. B., Dangol, P. M., Bhaunju, P. M., and Pradhan, S.: Rapid Urban Assessment of Air Quality for Kathmandu, Nepal: Summary, ICIMOD, 2012.
- Rizzo, M. J., and Scheff, P. A.: Utilizing the Chemical Mass Balance and Positive Matrix Factorization models to determine influential species and examine possible rotations in receptor modeling results, *Atmospheric Environment*, 41, 6986-6998, doi:10.1016/j.atmosenv.2007.05.008, 2007.
- 1055 Roberts, J. M., Veres, P., VandenBoer, T. C., Warneke, C., Graus, M., Williams, E. J., Lefer, B. L., Brock, C. A., Bahreini, R., Öztürk, F., Middlebrook, A. M., Wagner, N. L., Dubè, W. P. A., and de Gouw, J. A.: New insights into atmospheric sources and sinks of isocyanic acid, HNCO, from recent urban and regional observations, *Journal of Geophysical Research: Atmospheres*, 119, 1060-1072, doi:10.1002/2013JD019931, 2014.
- 1060 Sarkar, C., Kumar, V., and Sinha, V.: Massive emissions of carcinogenic benzenoids from paddy residue burning in North India, *Current Science*, 104, 1703 - 1709, 2013.
- Sarkar, C., Sinha, V., Kumar, V., Rupakheti, M., Panday, A., Mahata, K. S., Rupakheti, D., Kathayat, B., and Lawrence, M. G.: Overview of VOC emissions and chemistry from PTR-TOF-MS measurements during the SusKat-ABC campaign: high acetaldehyde, isoprene and isocyanic acid in wintertime air of the Kathmandu Valley, *Atmos. Chem. Phys.*, 16, 3979-4003, doi: 10.5194/acp-16-3979-2016, 2016.
- 1065 Sekimoto, K., Inomata, S., Tanimoto, H., Fushimi, A., Fujitani, Y., Sato, K., and Yamada, H.: Characterization of nitromethane emission from automotive exhaust, *Atmospheric Environment*, 81, 523-531, doi:10.1016/j.atmosenv.2013.09.031, 2013.
- Shim, C., Wang, Y., Singh, H. B., Blake, D. R., and Guenther, A. B.: Source characteristics of oxygenated volatile organic compounds and hydrogen cyanide, *Journal of Geophysical Research: Atmospheres*, 112, doi:10.1029/2006jd007543, 2007.
- 1070 Shrestha, S. R., Oanh, N. T. K., Xu, Q., Rupakheti, M., and Lawrence, M.: Analysis of the vehicle fleet in the Kathmandu Valley for estimation of environment and climate co-benefits of technology intrusions, *Atmospheric Environment*, 81, 579-590, doi:10.1016/j.atmosenv.2013.09.050, 2013.
- 1075 Sinha, V., Williams, J., Diesch, J. M., Drewnick, F., Martinez, M., Harder, H., Regelin, E., Kubistin, D., Bozem, H., Hosaynali-Beygi, Z., Fischer, H., Andrés-Hernández, M. D., Kartal, D., Adame, J. A., and Lelieveld, J.: Constraints on instantaneous ozone production rates and regimes during DOMINO derived using in-situ OH reactivity measurements, *Atmos. Chem. Phys.*, 12, 7269-7283, doi:10.5194/acp-12-7269-2012, 2012.



- 1080 Sinha, V., Kumar, V., and Sarkar, C.: Chemical composition of pre-monsoon air in the Indo-Gangetic Plain measured using a new air quality facility and PTR-MS: high surface ozone and strong influence of biomass burning, *Atmos. Chem. Phys.*, 14, 5921-5941, doi:10.5194/acp-14-5921-2014, 2014.
- Slowik, J. G., Vlasenko, A., McGuire, M., Evans, G. J., and Abbatt, J. P. D.: Simultaneous factor analysis of organic particle and gas mass spectra: AMS and PTR-MS measurements at an urban site, *Atmos. Chem. Phys.*, 10, 1969-1988, doi:10.5194/acp-10-1969-2010, 2010.
- 1085 Stockwell, C. E., Veres, P. R., Williams, J., and Yokelson, R. J.: Characterization of biomass burning emissions from cooking fires, peat, crop residue, and other fuels with high-resolution proton-transfer-reaction time-of-flight mass spectrometry, *Atmos. Chem. Phys.*, 15, 845-865, doi:10.5194/acp-15-845-2015, 2015.
- Stockwell, C. E., Christian, T. J., Goetz, J. D., Jayarathne, T., Bhave, P. V., Praveen, P. S., Adhikari, S., Maharjan, R., DeCarlo, P. F., Stone, E. A., Saikawa, E., Blake, D. R., Simpson, I. J., Yokelson, R. J., and Panday, A. K.: Nepal Ambient Monitoring and Source Testing Experiment (NAMaSTE): emissions of trace gases and light-absorbing carbon from wood and dung cooking fires, garbage and crop residue burning, brick kilns, and other sources, *Atmos. Chem. Phys.*, 16, 11043-11081, doi:10.5194/acp-16-11043-2016, 2016.
- 1090 Stone, E. A., Schauer, J. J., Pradhan, B. B., Dangol, P. M., Habib, G., Venkataraman, C., and Ramanathan, V.: Characterization of emissions from South Asian biofuels and application to source apportionment of carbonaceous aerosol in the Himalayas, *Journal of Geophysical Research: Atmospheres*, 115, D06301, doi:10.1029/2009jd011881, 2010.
- Tsai, S. M., Zhang, J., Smith, K. R., Ma, Y., Rasmussen, R. A., and Khalil, M. A. K.: Characterization of Non-Methane Hydrocarbons emitted from various Cookstoves used in China, *Environmental Science & Technology*, 37, 2869-2877, doi:10.1021/es026232a, 2003.
- 1100 Vlasenko, A., Slowik, J. G., Bottenheim, J. W., Brickell, P. C., Chang, R. Y. W., Macdonald, A. M., Shantz, N. C., Sjostedt, S. J., Wiebe, H. A., Leitch, W. R., and Abbatt, J. P. D.: Measurements of VOCs by proton transfer reaction mass spectrometry at a rural Ontario site: Sources and correlation to aerosol composition, *Journal of Geophysical Research: Atmospheres*, 114, D21305, doi:10.1029/2009JD012025, 2009.
- 1105 Watson, J. G., Chow, J. C., and Fujita, E. M.: Review of volatile organic compound source apportionment by chemical mass balance, *Atmospheric Environment*, 35, 1567-1584, doi:10.1016/S1352-2310(00)00461-1, 2001.
- Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., de Gouw, J., Akagi, S. K., Urbanski, S. P., Veres, P., Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, T. J., Hosseini, S., Miller, J. W., Cocker Iii, D. R., Jung, H., and Weise, D. R.: Coupling field and laboratory measurements to estimate the emission factors of identified and unidentified trace gases for prescribed fires, *Atmos. Chem. Phys.*, 13, 89-116, doi:10.5194/acp-13-89-2013, 2013.
- 1110 Yuan, B., Shao, M., de Gouw, J., Parrish, D. D., Lu, S., Wang, M., Zeng, L., Zhang, Q., Song, Y., Zhang, J., and Hu, M.: Volatile organic compounds (VOCs) in urban air: How chemistry affects the interpretation of positive matrix factorization (PMF) analysis, *Journal of Geophysical Research: Atmospheres*, 117, D24302, doi:10.1029/2012jd018236, 2012.
- Yuan, B., Hu, W. W., Shao, M., Wang, M., Chen, W. T., Lu, S. H., Zeng, L. M., and Hu, M.: VOC emissions, evolutions and contributions to SOA formation at a receptor site in Eastern China, *Atmos. Chem. Phys.*, 13, 8815-8832, doi:10.5194/acp-13-8815-2013, 2013.



- 1120 Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, Atmos. Chem. Phys., 9, 5131-5153, doi:10.5194/acp-9-5131-2009, 2009.