

## **Response to reviewers' comments**

Journal: ACP

Title: Nitropolyaromatic hydrocarbons - gas-particle partitioning, mass size distribution, and formation along transport in marine and continental background air

Author(s): Gerhard Lammel et al.

MS No.: acp-2016-1145

MS Type: Research article

We would like to thank the reviewers for their thoughtful reading, comments and questions, which considerably helped to improve this manuscript. We have addressed all comments below and have indicated the corresponding **modifications** in the revised version of the manuscript which is **attached to this response (changes highlighted)**.

### **Colour coding below:**

- (1) **comments from Referees,**
- (2) **author's response,**
- (3) **author's changes**

### **Anonymous Referee #1**

This manuscript provides a study of nitro-PAHs in 2 background locations. The study includes results about their atmospheric concentrations, gas/particle partitioning measurements and modeling, particle size distribution and substance patterns together with an evaluation of the urban influence of the samples using a Lagrangian particle dispersion model. Overall, the manuscript results are scientifically relevant and well presented. The modeling approach used to track the air mass history and to understand the nitro-PAHs sources and processes is very interesting and inovative. Thus, I recommend the publication of this paper in ACP after some minor revisions listed below.

Line 46: OK to cite review papers but you should also cite some key papers.  
Early key papers will be added. Sentence will be re-ordered.

Line 47: mutagenicity yes. Toxicity, no: What about metals, PAHs, other POPs. ...?  
True. Will be corrected

Line 55: recent paper about that: (Keyte et al., 2016) -  
This paper is cited (line 55). No change will be made.

Line 59-60: OK for 2-NFlt. Instead of 3-NFlt, better to give as an example for primary source, 1-nitropyrene.

Both 3NFLT and 1NPYR are dominated by the primary sources, i.e. diesel and gasoline (Ciccioli et al., 1989; IARC, 1989) vehicles for 3NFLT, and diesel and gasoline vehicles (Ciccioli et al., 1989; IARC, 1989), coal-fired power plants, Al smelters, carbon blacks (IARC, 1989), and coal stoves (Tang et al., 2005) for 1NPYR. But, based on smog-chamber studies; also heterogeneous oxidation by OH radicals was indicated as a (secondary) source for 1NPYR (Jariyasopit et al., 2014b, mentioned in the text, lines 397-398). No change will be made.

### **References:**

Ciccioli P., Cecinato A, Brancaleoni E, Draisci R, Liberti A: Evaluation of nitrated polycyclic aromatic hydrocarbons in anthropogenic emission and air samples: A possible means of detecting reactions of carbonaceous particles in the atmosphere. Aerosol Sci. Technol. 1989, 10, 296-310.

IARC: Diesel and gasoline engine exhausts and some nitroarenes. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Vol. 46, International Agency for Research on Cancer, Lyon. 1989, URL: <http://monographs.iarc.fr/ENG/Monographs/vol46/volume46.pdf>.

Tang N, Hattori T, Taga R, Igarashi K, Yang XY, Tamura K, Kakimoto H, Mishukov VF, Toriba A, Kizu R, Hayakawa K: Polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons in urban air particulates and their relationship to emission sources in the Pan–Japan Sea countries. Atmos. Environ. 2005, 39, 5817-5826.

Line 68: 2007 instead of 2006

Will be corrected

Line 72: an altitude (remote) site in the French Alps has been also investigated in (Albinet et al., 2008).

Will be corrected i.e., added to list of studies in remote environments

Line 174: PM10 determined by gravimetry using dedicated filters? Using automatic measurement instrumentation? Precise method and references.

Will be specified

Lines 261-266: These results refer to the particulate phase. What about the gaseous phase? You should discuss both phases or the total. Heterogenous photolysis is true for PAHs too. You could also have an enhanced formation of nitro-PAHs during night-time. Is the pattern the same for all PAHs and all nitro-PAHs or did you notice some differences for specific compounds? The time trend of some compounds should be shown somewhere as an example.

This is a misunderstanding: These results refer to the total, gas + particulate phases.

Will be specified

Line 286, Table 3: what about the altitude (remote) site in the Alps? The concentrations seemed very low.

Thanks. Results at high altitude site in the Alps will be added to Table 3 and discussed.

Line 289:  $\alpha_s$  should be defined before. This refers to the particulate phase, right? -

Particulate mass fraction  $\theta$ . The parameter will be introduced

Lines 293-300: this paragraph should be moved in the “Mass size distributions” section.

Thanks, yes, we agree. Will be moved.

Line 344, Figure 3: There is a problem with the x-axis, it should be the opposite or the size distribution shown is wrong. PAH and nitro-PAHs are mainly associated to the finest aerosol particles. You should also show the standard deviation of the particle size distribution for each size range. As it is shown on this figure, you should remove these data from the Table 1.

Thanks, yes, the x-axes were inverted erroneously. Plots will be corrected. Standard deviation of the mean will be shown together with (absolute) mass size distributions.

Lines 374-377: To evaluate the influence of primary or secondary sources of nitro-PAHs, it is better to focus on the 2-NFlt/1-NPyr ratio.

Indeed, this ratio had been used in the literature to this end, and is applied in this study (lines 431-439). However, nowadays, this appears inappropriate, as heterogeneous oxidation by OH radicals was indicated to be a possible secondary source for 1NPYR (Jariyasopit et al., 2014b, mentioned in the text, lines 397-398).

Lines 378-388: What about the day/night variations? Did you notice anything specific showing a predominance of day-time or night-time processes at the continental site? This is then useful to support the findings of the following paragraph (Lines 389-396).

The day/night variation of the ratio 2NFLT/1NPYR, the concentration of a secondarily formed over a primary emitted NPAH, reflects the combination of the diurnal variations of 1NPYR emissions, 2NFLT formation rate, and photochemical sinks of both substances, all unknown in quantitative terms. The day/night variation was 2NFLT/1NPYR  $\approx 12/\approx 16$  for day/night, which, therefore, is considered insignificant. No immediate conclusion could be drawn.

#### References:

Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., 2007. Polycyclic aromatic hydrocarbons (PAHs), nitrated PAHs and oxygenated PAHs in ambient air of the Marseilles area (South of France): Concentrations and sources. *Sci. Total Environ.* 384, 280-292.

Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., Jaffrezo, J.L., 2008. Nitrated and oxygenated derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys. Part 1: Concentrations, sources and gas/particle partitioning. *Atmos. Environ.* 42, 43-54.

Keyte, I.J., Albinet, A., Harrison, R.M., 2016. On-road traffic emissions of polycyclic aromatic hydrocarbons and their oxy- and nitro- derivative compounds measured in road tunnel environments. *Sci. Total Environ.*

#### **Anonymous Referee #2**

The Lammel et al. manuscript reports on measurements, partitioning calculations, and back-trajectory modeling of PAHs and nitro-PAHs at two European sites. The sites were chosen to study the transport of PAHs and nitro-PAHs from urban locations. Samples were collected using high-volume samplers, and particles were collected onto quartz fiber filters and gases using polyurethane foam plugs. Samples were extracted using dichloromethane and extracts were analyzed by gas chromatography-mass spectrometry. Partitioning calculations were performed using a recent poly-parameter linear free energy relationship. Particle trajectories were modeling using HYSPLIT and FLEXPART. There are a lot of high quality data and analyses presented in the manuscript, and the work should prove to be a valuable contribution to the field. There are no concerns with the methodology employed. The primary concern is with the quality of the representation-the technical writing and organization. There are a lot of grammatical errors and undefined symbols and acronyms. It is hard to follow the results as they are presented, particularly with regard to the discussions and conclusions being presented.

I recommend significant attention to the quality and clarity of the presentation, and perhaps another round of reviews following revision. Several specific suggestions and recommendations are provided below regarding grammar/symbols/abbreviations through the abstract and introduction. The intent here is to provide examples of the types of changes that need to be made throughout the manuscript; all such needed changes are not noted beyond the introduction.

Abstract: Why are the fractional doses not reported for the background site? Or is that being used to characterize the fraction at the marine site? This is not clear in the abstract as written.

The reason is that advection to the continental background site was influenced by a too homogeneous source distribution in central Europe (during the campaign). This explanation was given in lines 245-247 of the main text (no change).

On line 25, it is suggested to replace “were received” with “were calculated”.

Will be corrected

Line 22-23: For easier reading, recommend to move “with 2-nitrofluranthene . . . most abundant” out of the parenthesis and to the end of the sentence, and then start a new sentence with “While the concentration. . .”

Will be corrected

Line 31: NPYR is not defined or otherwise discussed

True. Acronym will be introduced

Line 36-37: Seems out of place. Recommend to remove or better integrate with rest of paragraph.

True. Will be integrated into rest of paragraph.

Line 49-50: Add comma after “more than one third”. On line 50, ambient what? Aerosols?

Yes. Will be specified.

Line 55: Insert “a” between “only” and “few”

Will be corrected

Line 60: Suggest to change “indicative for” to “indicative of” or “indicators for”

Yes, thanks. Will be corrected

Line 63: SVOCs is not defined

True. Acronym will be introduced

Line 69: Suggest removing “However . . . studied”. It has already been established (lines 66-67) that most NPAH observations are in urban areas. Could then use something like “Though there are a few studies in unpolluted environments: studies were conducted by. . .”.

Will be phrased as suggested.

Also suggest to change all the “were” to “have been” (lines 70-75) as measurements may be ongoing or planned.

Will be phrased as suggested.

Line 66-67: Suggest to change “suspect to” to “expect to”

True. Will be phrased as suggested.

Line 79-80: Suggest to change “lack of data. . .obvious” to “However, there is limited NPAH data from remote atmospheric environments. . .”

Will be phrased as suggested.

Line 80: Revise “aim of study to study”

Thanks. ‘...to study’ will be replaced by ‘...to characterise’

Line 112-113: What was the collection period of for the QFF samples at K-pusztá?

The collection period at the continental site will be specified in the sentence (previously in lines 111-113).

Line 116: Ni is undefined

Yes, thanks. Will be specified.

The comments/suggestions below address the clarity of the manuscript and analyses/conclusions presented. There are some technical questions here, but I think that more of the questions/comments arise from organizational structure and imprecise wording.

Conceptually, the urban fractional dose is clear; but how exactly were they calculated? I think much more time needs to be spent on the discussion of the calculated doses, sensitivities, and uncertainties to support the first paragraph of the conclusions section.

The calculation of the fractional dose was explained in the Supplementary material, S2.1.2. This text (S2.1.2) will be shifted to the main text (new Section 2.5), including all equations used for calculation of the dose. Uncertainties will be addressed. The new text will read:

#### “2.5 Quantification of urban influence on samples

The potential urban influence for individual samples collected at the marine site was based on the fraction of released Lagrangian particles which travelled through an urban boundary layer. A backward run from the sampling site was performed with Lagrangian particles (i.e. air parcels) being released during the entire sampling period. Three urban areas were considered, i.e. Izmir ( $\approx 300$  km direct distance,  $38.2\text{-}38.8^\circ\text{N}/26.2\text{-}27.3^\circ\text{E}$ ), Athens ( $\approx 300$  km,  $37.8\text{-}38.1^\circ\text{N}/23.5\text{-}23.8^\circ\text{E}$ ) and Istanbul ( $\approx 500$  km away,  $40.8\text{-}41.1^\circ\text{N}/28.6\text{-}29.5^\circ\text{E}$ ).

The urban fractional dose,  $D_{u\ i}$ , an air mass collected in sample  $i$  had received for a given simulation period  $\Delta t$  can be derived as:

$$(4) \quad D_{u\ i} = \sum_t N_{\text{blua}}(t) \times \Delta t_{\text{Rblua}} / (N_{\text{tot}}(t) \times \Delta t_i)$$

with  $N_{\text{blua}}(t)$  = number of virtual particles within the urban boundary layer during the specific time step, model output time resolution  $\Delta t_{\text{Rblua}} = 0.5$  h, and  $N_{\text{tot}}(t)$  = number of virtual particles present during the specific time step. Under the given flow conditions in the region, a 2-day time horizon is considered here. Hence, the simulation period is given as:

$$(5) \quad \Delta t_i = \Delta t_{\text{sample}} + 48\text{h}$$

with  $\Delta t_{\text{sample}}$  being the sampling time.  $D_{u\ i}$  takes values between 0 and 1, corresponding to none or all, respectively, of the entire sample air having crossed the urban boundary layer. The  $D_u$  time series with allocation to 3 urban areas is shown in the SM, Fig. S3.

The comparison of urban influence in samples of various sample volume,  $V$ , requires normalisation to  $V$ , a relative dose (equ. (5), with  $n$  = total number of samples collected). Values of  $D_{ru\ i}$  may exceed 1.

$$(6) \quad D_{ru\ i} = [\sum_n V_n / (nV_i)] \times D_{u\ i}$$

The urban fractional dose,  $D_{u\ i}$ , accuracy is limited by the meteorological input data (here  $0.125^\circ \times 0.125^\circ$  resolution, hourly) and boundary layer depth calculation. In the FLEXPART model, the latter is done according to Vogelezang and Holtslag, 1996.”

Line 209-213: The discussion of the two OM phases is confusing as written. Is one phase low molecular mass/water soluble OM and the other phase high molecular mass/organic soluble OM?

This text will be rephrased. The new text will read: “OM is assumed to be constituted of two separate phases, low to mid molecular mass, both organic soluble and water soluble OM. For these, ppLFER equations for dimethyl sulfoxide-air (representing the low molecular mass range) and for polyurethane ether-air (representing the high molecular mass OM) are used, respectively (Shahpoury et al., 2016).”

Line 229: What is meant by “the substance patterns are similar”? Diurnal patterns? Monomer ratios? Monomer fractions? (also Line 254)

Composition of NPAH mixture. Will be specified.

Line 241: Assume “this” data set refers to the marine data set? Language throughout could be more specific and precise (see also comment above).

Yes. Will be specified

Line 256: What is the evidence for the secondary emissions of 4rPAHs? Are the cited studies part of the same campaign? Or is this an inference based on a prior study? Or using measured monomers or ratios using a method in the cited study?

Yes, the cited studies focussed on the atmosphere-surface exchange during the campaigns. This will be clarified to avoid confusion. The revised sentence (lines 256-259) will read: “The investigation of the diffusive air-surface exchange processes during the measurements presented here showed that 4rPAHs were in fact influenced by secondary emissions, namely throughout day and night from the soil at the continental site (by average 16.3 and 9.3  $\text{pg m}^{-2} \text{h}^{-1}$  for FLT and PYR, respectively; Degrendele et al., 2016) or occasionally from surface seawater at the marine site (during at least 1 day-time interval out of in total 3 of this data subset; Lammel et al., 2016).”

Line 262-267: Is the day/night ratio of PAHs to n-PAHs sufficiently explained by photochemistry? Or may there be measurement-based explanations as well? For example, do the citations referenced support that the photolysis rates of n-PAHs are faster than those of PAHs? If so, this should be explicitly stated. Is the much higher day/night ratio of PAHs surprising given reported oxidation timescales of hours? What exactly is meant by “the same could be reflected” in line 267? Emission rates are governing ratios (based on line 271)? Or that n-PAHs are photochemically degraded faster than precursors (based on line 275)?

Are the lowest reported remote concentrations due to changing urban emissions? Changing photochemistry? Or improvements of in analytical techniques? Some combination? Further discussion on this point is warranted.

Further elaboration on NPAH sources and sinks is impossible, as not enough is known about NPAH atmospheric fate. In fact, this was a major motivation for this study. While the atmospheric lifetimes of 4rPAHs are limited by homogeneous reactions, no such statement is possible for NPAHs. Their lifetimes are eventually limited by heterogeneous photolysis or heterogeneous thermal reactions, but cannot be quantified, as these rate coefficients depend on PM composition and have almost not studied at all (with the exception of wood smoke particles; Fan et al., 1996). Therefore, it is unknown, whether NPAHs are photochemically degraded faster than their precursors. The day/night ratio of PAHs could not be explained by the sinks (stronger during day-time), but is in agreement with the finding of secondary emissions (stronger during day-time) at the site (discussed in lines 262-263).

Also, NPAH field studies are scarce, such that little data exist for comparison (Table 3) and no conclusions can be drawn from existing data with regard to temporal trends.

In order to put day/night ratios into better context, the sentences discussed here (lines 262-271) will be rephrased; “NPAH lifetimes may be limited by heterogeneous photolysis, but available kinetic data are scarce and limited to few aerosol types” (specification in line 265-266). Furthermore, limited knowledge of NPAH atmospheric fate will be highlighted by adding “...but their atmospheric lifetimes are still unknown.” To a sentence in the introduction (namely: “These substances have been suggested as tracers for air pollution on the time scales of hours to days” in lines 61-62); The sentence “.More studies into NPAH atmospheric fate, both field observations and kinetic data, are needed in order to assess and quantify spatial and temporal trends, the long-range transport potential and persistence..” (will be added to the conclusions, after line 457).

Line 289-290: Particle phase distributions? What does the symbol theta represent?

Thanks. Will be introduced at first mentioning (line 190)

Was the pp-LFER also used for the PAHs? Discussion focused on n-PAHs.

In this study gas-particle partitioning of parent PAHs was not studied. However, the ppLFER model used had been applied previously on parent PAHs, which showed good prediction power across three urban and rural sites in Europe and the Mediterranean (Shahpoury et al., 2016).

How sensitive are the partitioning results to the assumptions in the pp-LFER? What are the uncertainty bounds on the phase predictions? Are they significant enough to influence any conclusions presented regarding phase partitioning?

More discussion on the shallow slopes observed for the marine n-PAHs would be valuable. The model does not seem to be capturing any of the observed variance.

In lack of experimental data, the ppLFER model for most NPAHs is based on estimated solute-specific Abraham solvation parameters for the various types of molecular interactions. The estimation is based on a group contribution method (ACD, 2015). Experimentally based descriptors used for one NPAH, i.e. 9-nitroanthracene, lead to better predictions (RMSE = 0.33, log  $K_p$  units) than the estimated descriptors (RMSE = 0.75). The sensitivity of assumptions on PM phase composition, made in the model had been tested and it was found that the mixing ratios of the WSOSOM and organic polymers' phases, as well as the soot surface area do not contribute significantly to the discrepancy ( $\ll 1$ , log  $K_p$  units).

The predicted variation of  $K_p$  is significantly less than the observed for four NPAHs at the marine and one at the continental site (discussed in lines 335-349). The reason is unknown. Apart from PM properties it might be related to sampling or sample handling, although a number of such issues can be excluded, as same sampler, same temperature range, same sampling protocols applied across sites with both satisfactory and deficient agreement between predicted and observed  $K_p$ .

Satisfactory and deficient agreement between predicted and observed  $K_p$  across sites will be discussed accordingly.

Sentences to be added: "While the sensitivity of assumptions regarding PM phase composition, made in the model do not contribute significantly to the deviations ( $\ll 1$ , log  $K_p$  units), a significant part can be attributed to the usage of estimated solute-specific Abraham solvation parameters (taken from ACD, 2015), in lack of experimentally based descriptors. E.g., for an urban site (Tomaz et al., 2016) it was found that experimentally based descriptors used for 9NPAH lead to better predictions than the estimated descriptors i.e., RMSEs differed by 0.43 log units (...) The reason is unknown. Moreover, sampling or sample handling artefacts cannot be excluded, even so same temperature range, sampler and sampling protocols applied across sites with both satisfactory and deficient agreement between predicted and observed  $K_p$ ."

Is there a distinction between observed yield and potential yield? Or are these terms being used interchangeably?

All observed yields are conservatively considered to be potential yields. This will be better specified by replacing " $c_{PAH}$  (total concentrations) and reflects an upper estimate, as other PAH photochemical sinks, such as formation of oxy-PAHs, are neglected" by "... $c_{PAH}$  (total concentrations). This yield is called 'potential' as it reflects an upper estimate, as other PAH photochemical sinks, such as formation of oxy-PAHs, are neglected".

Line 436-437: What is meant by "hardly limited by"?

Will be rephrased: "The mass size distribution is determined by the particle size upon emission (primary NPAHs) and condensation and redistribution in the aerosol along transport, hence, does not include the short-lived coarse mass fraction. This indicates a high long-range transport potential."

Line 446-447: “Levels” of what?  
NPAH levels. Will be specified.

Further correction:

In Table 1 (in the line: ‘marine gas’) and Table 2 (in the column ‘marine θ’) minor mistakes (which do not impact on the conclusions) were found and will be corrected.



1           **Nitro-polycyclic aromatic hydrocarbons - gas-particle partitioning, mass size**  
2           **distribution, and formation along transport in marine and continental background air**

3  
4           Gerhard Lammel <sup>1,2\*</sup>, Marie D. Mulder <sup>1</sup>, Pourya Shahpoury <sup>2</sup>, Petr Kukučka <sup>1</sup>, Hana Lišková  
5                           <sup>1</sup>, Petra Příbylová <sup>1</sup>, Roman Prokeš <sup>1</sup>, Gerhard Wotawa <sup>3</sup>

6  
7           <sup>1</sup> Masaryk University, Research Centre for Toxic Compounds in the Environment, Brno,  
8                           Czech Republic

9           <sup>2</sup> Max Planck Institute for Chemistry, Multiphase Chemistry Department, Mainz, Germany

10           <sup>3</sup> Zentralanstalt fuer Meteorologie und Geodynamik, Wien, Austria

11  
12           \*lammel@recetox.muni.cz

13  
14           **Abstract**

15 Nitro-polycyclic aromatic hydrocarbons (NPAH) are ubiquitous in polluted air but little is  
16 known about their abundance in background air. NPAHs were studied at one marine and one  
17 continental background site i.e., a coastal site in the southern Aegean Sea (summer 2012) and  
18 a site in the central Great Hungarian Plain (summer 2013), together with the parent  
19 compounds, PAHs. A Lagrangian particle dispersion model was used to track air mass history.  
20 Based on Lagrangian particle statistics, the urban influence on samples was quantified for the  
21 first time as a fractional dose to which the collected volume of air had been exposed to.

22 At the remote marine site, the 3-4 ring **Nnitro**-PAH (sum of 11 targeted species) concentration  
23 was 23.7 pg m<sup>-3</sup> while the concentration of 4-ring PAHs (6 species) was 426 pg m<sup>-3</sup>. **2-**  
24 **nitrofluoranthene (2NFLT) and 3-nitrophenanthrene were the most abundant NPAHs.** Urban  
25 fractional doses in the range <0.002–5.4% were **calculated**. At the continental site, the  
26  $\Sigma_{11}$  3-4rNPAH and  $\Sigma_6$  4rPAH were 58 and 663 pg m<sup>-3</sup>, respectively, with 9-nitroanthracene  
27 and 2NFLT being highest concentrated amongst the targeted NPAHs. The NPAH levels  
28 observed in the marine background are the lowest ever reported and remarkably lower, by  
29 more than one order of magnitude, than one decade before. Day-night variation of NPAHs at

30 the continental site reflected shorter lifetime during the day, possibly because of photolysis of  
31 some NPAHs. The yields of formation of 2NFLT and 2-nitropyrene (2NPYR) in marine air  
32 seem to be close to the yields for OH-initiated photochemistry observed in laboratory  
33 experiments under high NO<sub>x</sub> conditions. Good agreement is found for prediction of NPAH  
34 gas-particle partitioning using a multi-phase poly-parameter linear free energy relationship.  
35 Sorption to soot is found less significant for gas-particle partitioning of NPAHs than for  
36 PAHs.

37 The NPAH levels determined in the southeastern outflow of Europe confirm intercontinental  
38 transport potential.

39

40 **Keywords:** long-range transport potential, semi-volatile organic compounds, PAH  
41 photochemistry,

42

### 43 1. Introduction

44 PAHs may undergo chemical transformations in the gaseous and in the particulate phase  
45 (Finlayson-Pitts and Pitts, 2000; Keyte et al., 2013). Nitro-PAHs (NPAHs), ~~earlier predicted~~  
46 ~~based on smog-chamber experiments (Atkinson and Arey, 1994), and later~~ observed in urban  
47 and rural areas (Nielsen et al., 1984; Feilberg et al., 2001; Finlayson-Pitts and Pitts, 2000;  
48 Keyte et al., 2013) and predicted based on smog-chamber experiments (Atkinson and Arey,  
49 1994), seem to be most significant derivatives: Mutagenicity ~~and toxicity~~ of atmospheric  
50 aerosols in general is mostly related to NPAHs (Grosjean et al., 1983; Garner et al., 1986;  
51 Finlayson-Pitts and Pitts, 2000; Claxton et al., 2004; Hayakawa, 2016). A large part, more  
52 than one third, of the mutagen potential of ambient aerosols may be attributable to NPAHs  
53 (Schuetzle, 1983).

54 Secondary formation of NPAH from PAHs is thought to occur on short time scales (hours).  
55 This has been observed for PAHs collected on filters (Ringuet et al., 2012a; Zimmermann et  
56 al., 2013; Jaryasopit et al., 2014a, 2014b), and also in in urban plumes (Bamford and Baker,  
57 2003; Arey et al., 1989; Reisen and Arey, 2005). Although many NPAHs are emitted from  
58 road traffic, only a few are abundant in this source type (Arey, 1998; Keyte et al., 2013 and  
59 2016; Inomata et al., 2015; Alves et al., 2016). The occurrence of various isomers of  
60 nitrofluoranthene (NFLT) and nitropyrene (NPYR) can be used to study PAH sources, PAH  
61 chemical transformations and the role of the photo-oxidants hydroxyl radical (OH) and nitrate

62 radical (NO<sub>3</sub>) (Ciccioli et al., 1996; Finlayson-Pitts and Pitts 2000). E.g., 3- and 2-  
63 nitrofluoranthene (3-, 2NFLT) are indicative ~~of~~ primary and secondary sources,  
64 respectively. These substances have been suggested as tracers for air pollution on the time  
65 scales of hours to days (Ciccioli et al., 1996; Finlayson-Pitts and Pitts 2000; Keyte et al.  
66 2013), but their atmospheric lifetimes are still unknown.

67 Like their precursors, NPAHs are semivolatile organic compounds (SVOCs), partitioning  
68 between the phases of the atmospheric aerosol. Similar to other SVOCs, the NPAHs' phase  
69 distribution was found to depend on temperature (summer and winter campaigns in the Alps;  
70 Albinet et al., 2008b) and results from both absorptive as well as adsorptive contributions  
71 (Tomaz et al., 2016). NPAHs have primarily been observed in polluted areas (e.g. Pitts et al.,  
72 1985; Ramdahl et al., 1986; Garner et al., 1986; Albinet et al., 2007 and 2008a; Ringuet et  
73 al., 2012a and 2012b; Zimmermann et al., 2012; Barrado et al., 2013; Li et al., 2016).  
74 ~~However, NPAHs have hardly been studied in the unpolluted environment. Though there are a~~  
75 ~~few studies were conducted in the rural environments~~ i.e., in Germany (Ciccioli et al., 1996),  
76 in the French Alps (100-1000 pg m<sup>-3</sup> range for the sum of 10 NPAHs; Albinet et al., 2008a)  
77 and in northern China (Li et al., 2016). Very few measurements ~~have been~~ were performed in  
78 the remote atmospheric environment i.e., in the Mediterranean (Tsapakis and Stephanou,  
79 2007), high altitude sites in the Himalayas (single data; Ciccioli et al. 1996) and French Alps  
80 (Albinet et al., 2008a), and in the Arctic (with so-called Arctic haze; Masclat et al. 1988;  
81 Halsall et al. 2001). With regard to the long-range transport potential, the state of the  
82 knowledge is that at least some NPAHs are ~~expected~~ ~~suspect~~ to go into intercontinental  
83 transport (Lafontaine et al., 2015) and might be ubiquitous in the global atmosphere (Ciccioli  
84 et al., 1996).

85 However, ~~there is limited~~ ~~lack of~~ NPAH data from remote atmospheric environments is  
86 obvious and little is known about their long-range transport potential. The aim of this study  
87 was to ~~characterise~~ ~~study~~ the long-range transport potential of NPAHs by measurements at  
88 remote sites of Europe, addressing the continental background and the outflow of the  
89 continent.

90

## 91 **2. Methodology**

### 92 **2.1 Sampling**

93 High-volume air sampling was conducted at a marine background site, Finokalia  
94 (35.3°N/25.7°E, 250 m a.s.l.), in the context of a coordinated field experiment 2-13 July 2012

95 (Lammel et al., 2015) and at a continental background site in central Europe, K-pusztá  
96 (46°58'N/19°33'E, 125 m a.s.l.; Degrendele et al., 2016), 5-16 August 2013. The Finokalia site  
97 is located on a cliff at the northern coast of Crete, some 70 km east of major significant  
98 anthropogenic emissions (Iraklion, a city of 100000 inhabitants with airport and industries;  
99 Mihalopoulos et al., 1997; Kouvarakis et al., 2000). The K-pusztá site is located on a clearing,  
100 characterised by uncultivated grassland, in a mostly coniferous forest in the Hungarian  
101 (Pannonian) Great Plain, ca. 70 km and 270 km southeast of Budapest and Vienna,  
102 respectively ( $\approx 2$  mn inhabitants each). The background site character of both observatories  
103 had been demonstrated (Borbély-Kiss et al., 1988; Kouvarakis et al., 2000; Vrekoussis et al.,  
104 2005). Meteorological and trace gas measurements are covered by both observatories, which  
105 are stations of the EMEP network (EMEP, 2015).

106 High volume air samples were collected using a HV-100P (Baghirra, Prague, Czech  
107 Republic), equipped with a multi-stage cascade impactor (Andersen Instruments Inc.,  
108 Fultonville, New York, USA, series 230, model 235) with five impactor stages, corresponding  
109 to 10–7.2, 7.2–3, 3–1.5, 1.5–0.95 and 0.95–0.49  $\mu\text{m}$  of aerodynamic particle size,  $D_p$  (spaced  
110 roughly equal  $\Delta \log D$ ), a backup filter collecting particles  $< 0.49 \mu\text{m}$  and, downstream, two  
111 polyurethane foam plugs (PUFs, Molitan, Břeclav, Czech Republic, density  $0.030 \text{ g cm}^{-3}$ ,  
112 placed in a glass cartridge), together 10 cm high. Particles were sampled on slotted QFF  
113 substrates (TE-230-QZ, Tisch Environmental Inc., Cleves, USA,  $14.3 \times 13.7 \text{ cm}$ ) and glass  
114 fibre filters (Whatman,  $20.3 \times 25.4 \text{ cm}$ ). The filters had been cleaned prior to use by heating  
115 108 ( $330^\circ\text{C}$ ). PUFs were cleaned (8 hour-extraction in acetone and 8 hours in  
116 dichloromethane (DCM)), wrapped in two layers of aluminum foil, placed into zip-lock  
117 polyethylene bags and kept in the freezer prior to deployment. The sampler was operated at  
118 constant flow rate of  $68 \text{ m}^3 \text{ h}^{-1}$ . Day/night sampling (changing at sunset and sunrise) of  
119 gaseous samples (PUF) was performed at both sites ( $V = 600\text{-}1000 \text{ m}^3$ ), while at the marine  
120 site the impactor filter (QFF) samples were collected over 24 h (5) or 48 h (3).

121 PUFs were cleaned (8 hour-extraction in acetone and 8 hours in dichloromethane (DCM)),  
122 wrapped in two layers of aluminum foil, placed into zip-lock polyethylene bags and kept in  
123 the freezer prior to deployment. The sampler was operated at constant flow rate of  $68 \text{ m}^3 \text{ h}^{-1}$ .  
124 Day/night sampling of gaseous samples (PUF) was performed at both sites ([12 h](#),  $V \approx 700 \text{ m}^3$ ),  
125 while at the marine site the impactor filter (QFF) samples were collected over 12 h ( $n = 1$ ), 24  
126 h (4) or 48 h (3).

127 Particle number concentration,  $N$ , was determined by an optical particle counter (Grimm  
128 model 107, Ainring, 31 channels between 0.25 and 32  $\mu\text{m}$  of aerodynamic particle diameter,  
129  $D$ ). Aerosol surface concentration,  $S$  ( $\text{cm}^{-1}$ ), was derived as  $S = \pi \sum_i N_i D_i^2$  assuming  
130 sphericity. Hereby, true  $S$  will be underestimated, in particular if particles of irregular form  
131 were abundant (e.g. Jaenicke, 1988). Comparisons with absolute methods (e.g. Pandis, et al.  
132 1991) suggest that the discrepancy may reach up to a factor of 2-3. The mass median diameter  
133 ( $D_m$ ,  $\mu\text{m}$ ), was derived as  $\log D_m = \frac{\sum_i m_i \log D_i}{\sum_i m_i}$  with  $m_i$  denoting the mass in size class  
134  $i$ ,  $D_i$  being the geometric mean diameter collected on stage  $i$  of the cascade impactor.

135

## 136 2.2 Chemical analysis

137 All air samples were extracted with DCM using an automatic warm Soxhlet extractor (Büchi  
138 B-811, Switzerland). Deuterated PAHs (D8-naphthalene, D10-phenanthrene, D12-perylene;  
139 Wellington Laboratories, Canada) were used as surrogate standards for both PAHs and  
140 NPAHs. Deuterated PAHs proved to be suitable surrogate standards for NPAHs. These were  
141 spiked on each PUF prior to extraction. The extract was split in two parts, 1/9 for PAHs and  
142 Nitro-PAHs analysis, 9/10 for PBDEs, PCBs and OCPs. The PAHs and Nitro-PAHs aliquot  
143 was a subject to open column chromatography clean-up. Glass column (1 cm i.d.) was filled  
144 with 5 g activated silica (150°C for 12 h), sample was loaded and eluted with 10 mL *n*-  
145 hexane, followed by 40 mL DCM. The cleaned sample was evaporated under a stream of  
146 nitrogen in a TurboVap II apparatus (Biotage, Sweden), transferred into a conical GC vial and  
147 spiked with recovery standard, terphenyl, the volume was reduced to 100  $\mu\text{L}$ .

148 GC-MS analysis of 4-ring PAHs (fluoranthene (FLT), pyrene (PYR), benzo(b)fluorene  
149 (BBN), benzo(a)anthracene (BAA), triphenylene (TPH) and chrysene (CHR)) and 2-4 ring  
150 NPAHs (1- and 2-nitronaphthalin (1-, 2NNAP), 3- and 5-nitroacenaphthene (3-, 5NACE), 2-  
151 nitrofluorene (2NFLN), 9-nitroanthracen (9NANT), 3- and 9-nitrophenanthren (3-, 9NPHE),  
152 2- and 3-nitrofluoranthene (2-, 3NFLT), 1- and 2-nitropyrene (1-, 2NPYR), 7-  
153 nitrobenz(a)anthracene (7NBAA), 6-nitrochrysene (6NCHR) was performed using a gas  
154 chromatograph atmospheric pressure chemical ionization tandem mass spectrometer (GC-  
155 APCI-MS/MS) instrument, Agilent 7890A GC (Agilent, USA), equipped with a 60m  $\times$   
156 0.25mm  $\times$  0.25 $\mu\text{m}$  DB-5MSUI column (Agilent, J&W, USA), coupled to Waters Xevo TQ-S  
157 (Waters, UK). Injection was 1  $\mu\text{L}$  splitless at 280°C, with He as carrier gas at constant flow  
158 1.5 mL  $\text{min}^{-1}$ . The GC oven temperature program was as follows: 90°C (1 min), 40°C/min  
159 to 150°C, 5°C/min to 250°C (5 min) and 10°C/min to 320°C (5 min). APCI was used in

160 charge transfer conditions. The isomers 2- and 3NFLT were not separated by the GC method,  
161 but co-eluted and are reported as sum.

162 Recovery of native analytes varied 72-102% for PAHs and deuterated PAHs, 70-110% for  
163 NPAHs (details see supplementary material (SM), Table S1a). The results were not recovery  
164 corrected. The mean of field blank values was subtracted from the sample values. Values  
165 below the mean + 3 standard deviations of the field blank values were considered to be  
166 <LOQ. Field blank values of some analytes in air samples were below the instrument limit of  
167 quantification (ILOQ), which corresponded to 0.004-0.069  $\text{pg m}^{-3}$  for NPAHs (except for  
168 INNAP for which it ranged 0.60-0.87  $\text{pg m}^{-3}$ ) and 0.010-0.126  $\text{pg m}^{-3}$  for 4-ring PAHs  
169 (except for FLT and PYR for which it ranged 0.17-0.59  $\text{pg m}^{-3}$ ) (Table S1).

170 Higher LOQs were determined for some of the NPAHs and for all 4-ring PAHs in gaseous air  
171 samples (PUFs), namely 0.006-0.009 ng (corresponding to 3.5-8.0  $\text{pg m}^{-3}$ ) for 3NACE and  
172 2NPYR, 0.028-0.097 (corresponding to 16-86  $\text{pg m}^{-3}$ ) for 2NNAP, 2NFLT and 1NPYR, and  
173 0.10-0.27 ng (corresponding to  $\approx$ 60-240  $\text{pg m}^{-3}$ ) for 4-ring PAHs (except for FLT and PYR for  
174 which it was 1.71 and 1.05 ng, respectively, corresponding to  $\approx$ 600-1500  $\text{pg m}^{-3}$ ). In  
175 particulate phase samples, where separate field blanks for the 2 different QFFs were  
176 determined (on the impactor stages on one hand side and the backup filter on the other hand  
177 side), higher LOQs were determined for some of the NPAHs and for all 4-ring PAHs, namely  
178 0.008-0.089 ng (corresponding to 4.6-79  $\text{pg m}^{-3}$ ) for 2NNAP, 2NFLT, 1NPYR and 2NPYR,  
179 0.26-0.31 ng (corresponding to 150-274  $\text{pg m}^{-3}$ ) for 9NANT, and 0.05-0.22 ng (corresponding  
180 to  $\approx$ 30-200  $\text{pg m}^{-3}$ ) for 4-ring PAHs (except for FLT and PYR for which it was 0.79 and 0.36  
181 ng, respectively, corresponding to  $\approx$ 200-700  $\text{pg m}^{-3}$ ).

182 The breakthrough in PUF samples was estimated (Pankow, 1989; ACD, 2015; Melymuk et  
183 al., 2016), and as a consequence, 2-3 ring PAHs and 2-ring NPAHs results were excluded  
184 from this study as their sampling may have been incomplete. We, therefore, report  $\Sigma_6$ 4rPAH  
185 and  $\Sigma_{11}$ 3-4rNPAH.

186 Particulate matter mass ( $\text{PM}_{10}$ ) was determined by gravimetry (microbalance, filters  
187 accommodated to stable temperature and humidity, 3 replicate weighings), and organic matter  
188 (OM) and elemental carbon (EC) contents of PM by a thermal-optical method (Sunset Lab.,  
189 USA; EUSAAR protocol).

190

## 191 2.3 Gas-particle partitioning

192 Gas-particle partitioning was studied by applying a multiphase ppLFER model, which was  
193 recently introduced (Shahpoury et al., 2016). In brief, partitioning of semivolatile compounds  
194 in air can be described (Yamasaki et al., 1982), by

195

$$196 \quad (1) \quad K_p = c_{ip} / (c_{ig} \times c_{PM})$$

197

198 where  $K_p$  ( $\text{m}^3 \text{air} (\text{g PM})^{-1}$ ) is the temperature dependent partitioning coefficient,  $c_{PM}$  ( $\text{g m}^{-3}$ ) is  
199 the concentration of particulate matter in air,  $c_{ip}$  and  $c_{ig}$  are the analyte (i) concentrations ( $\text{ng}$   
200  $\text{m}^{-3}$ ) in the particulate and gas phase, respectively.  $K_p$  can be predicted using models based on  
201 single- and poly-parameter linear free energy relationships (spLFER, ppLFER). spLFER's  
202 relate the partitioning coefficient to one physic-chemical property i.e., assume one process to  
203 determine the sorption process, while ppLFER's in principle account for all types of  
204 molecular interactions between solute and matrix (Goss and Schwarzenbach, 2001). The  
205 observed particulate mass fraction, data,  $\theta = c_p / (c_g + c_p)$  (Table 2), were tested with both a  
206 spLFER and a ppLFER model. The spLFER chosen is the widely used  $K_{oa}$  model of Finizio et  
207 al., 1997 (results presented in the Supplementary material (SM), S2.3). The ppLFER is a  
208 multi-phase model recently presented (Shahpoury et al., 2016) and applied for NPAHs  
209 (Tomaz et al., 2016). It is based on linear solvation energy relationships (Abraham, 1993;  
210 Goss, 2005):

211

$$212 \quad (2) \quad \log K_p = eE + sS + aA + bB + lL + c$$

$$213 \quad (3) \quad \log K_p = sS + aA + bB + vV + lL + c$$

214

215 where capital letters E, S, A, B, L, and V are solute-specific Abraham solvation parameters for  
216 excess molar refraction (describes interactions between  $\pi$ - and lone (n-) electron pairs),  
217 polarizability/ dipolarity, solute H-bond acidity, solute H-bond basicity, logarithm of solute  
218 hexadecane-air partitioning coefficient (unitless), and McGowan molar volume ( $\text{cm}^3$   
219  $\text{mol}^{-1}$ )/100, respectively (Endo and Goss, 2014). The corresponding parameters e, s, a, b, l,  
220 and v reflect matrix-specific solute-independent contribution to  $K_p$ . In lack of experimental  
221 data, the solute descriptors for NPAHs were taken from M.H. Abraham (personal  
222 communication). The multi-phase ppLFER considers adsorption onto soot,  $(\text{NH}_4)_2\text{SO}_4$ , and  
223  $\text{NH}_4\text{Cl}$ , and absorption into particulate organic matter (OM). OM is assumed to be constituted  
224 of two separate phases, low to ~~mid~~high molecular mass, both organic soluble and water

225 | soluble OM. ~~For these, represented by a ppLFER equations~~ for dimethyl sulfoxide-air  
226 | ~~(representing the low molecular mass range) on one hand side, and high molecular mass OM,~~  
227 | ~~represented by a ppLFER equation and~~ for polyurethane ether-air ~~(representing the high~~  
228 | ~~molecular mass OM) are used, respectively~~ (Shahpoury et al., 2016).

229 | A conventional single-parameter LFER ( $K_{oa}$ ) model is applied, too.

230

## 231 | 2.4 Air mass history analysis

232 | The HYSPLIT (Draxler and Rolph, 2003) and FLEXPART (Stohl et al., 1998, 2005) models  
233 | were used to identify air mass histories over 10 and 2 days, respectively. The possible  
234 | influence of polluted air on samples was quantified using a novel method of applying  
235 | Lagrangian particle statistics (FLEXPART, see SM, S2.2). To this end, for the entire sampling  
236 | period, one particle per second was released. The model output is generated at  $0.062^\circ$  ( $\approx 7$   
237 | km), every 30 minutes and expressed as 'residence time' i.e., a measure of the time particles  
238 | resided in grid cells. ECMWF meteorological data ( $0.125^\circ \times 0.125^\circ$  resolution, hourly) were  
239 | used as input.

240

## 241 | 2.5 Quantification of urban influence on samples

242 | The potential urban influence for individual samples collected at the marine site was based on  
243 | the fraction of released Lagrangian particles which travelled through an urban boundary layer.  
244 | A backward run from the sampling site was performed with Lagrangian particles (i.e. air  
245 | parcels) being released during the entire sampling period. Three urban areas were considered,  
246 | i.e. Izmir ( $\approx 300$  km direct distance,  $38.2$ - $38.8^\circ$ N/ $26.2$ - $27.3^\circ$ E), Athens ( $\approx 300$  km,  $37.8$ -  
247 |  $38.1^\circ$ N/ $23.5$ - $23.8^\circ$ E) and Istanbul ( $\approx 500$  km away,  $40.8$ - $41.1^\circ$ N/ $28.6$ - $29.5^\circ$ E).

248 | The urban fractional dose,  $D_{u,i}$ , an air mass collected in sample  $i$  had received for a given  
249 | simulation period  $\Delta t$  can be derived as:

250

$$251 | \underline{(4) \quad D_{u,i} = \sum_t N_{blua}(t) \times \Delta t_{Rblua} / (N_{tot}(t) \times \Delta t_i)}$$

252

253 | with  $N_{blua}(t)$  = number of virtual particles within the urban boundary layer during the specific  
254 | time step, model output time resolution  $\Delta t_{Rblua} = 0.5$  h, and  $N_{tot}(t)$  = number of virtual

Formatiert: Zeilenabstand: 1.5 Zeilen



255 particles present during the specific time step. Under the given flow conditions in the region, a  
256 2-day time horizon is considered here. Hence, the simulation period is given as:

$$258 \quad (5) \quad \Delta t_i = \Delta t_{\text{sample}} + 48\text{h}$$

259  
260 with  $\Delta t_{\text{sample}}$  being the sampling time.  $D_{u,i}$  takes values between 0 and 1, corresponding to  
261 none or all, respectively, of the entire sample air having crossed the urban boundary layer. The  
262  $D_u$  time series with allocation to 3 urban areas is shown in the SM, Fig. S3.

263 The comparison of urban influence in samples of various sample volume,  $V$ , requires  
264 normalisation to  $V$ , a relative dose (equ. (5), with  $n$  = total number of samples collected).  
265 Values of  $D_{ru,i}$  may exceed 1.

$$267 \quad (6) \quad D_{ru,i} = [\sum_n V_n / (nV_i)] \times D_{u,i}$$

268  
269 The urban fractional dose,  $D_{u,i}$ , accuracy is limited by the meteorological input data (here  
270  $0.125^\circ \times 0.125^\circ$  resolution, hourly) and boundary layer depth calculation. In the FLEXPART  
271 model, the latter is done according to Voegelezang and Holtslag, 1996.

### 273 3. Results and discussion

274 The NPAH levels are distinctly lower at the marine than at the continental site,  $\sum_{11 \text{ 3-4r}} \text{NPAH} =$   
275  $22.5$  and  $58.5 \text{ pg m}^{-3}$ , respectively (Table 1). The NPAHs showing the highest concentrations  
276 were 2NFLT and 3NPHE at the marine (Fig. 1b) and 9NANT and 2NFLT at the continental  
277 site (Fig. 1d, Table 2). The substance patterns (composition of NPAH mixture) at both sites  
278 are similar, though ( $R^2 = 0.76$ ,  $P > 0.99$ , t-test). At the marine site, advection was northerly,  
279 with air masses originating (time horizon 10 days) in eastern and central Europe and, towards  
280 the end of the campaign, in the western Mediterranean. The site was placed into the  
281 southeastern outflow of Europe.  $\text{NO}_x$  ( $0.2$ - $0.6 \text{ ppbv}$ ), EC ( $0.2$ - $0.8 \text{ } \mu\text{g m}^{-3}$ ) and  $\text{PM}_{10}$  ( $18.3$ - $39.3$   
282  $\text{ } \mu\text{g m}^{-3}$ ) reflect background conditions. Air mass history analysis suggests that the somewhat  
283 elevated concentration in the first sample collected at the marine site (Fig. 1a) is related to  
284 long-range transport influenced by passage over the urban areas of Izmir and Istanbul (urban

285 fractional dose  $D_u = 5.0\%$ , in contrast to the mean which was  $1.6\%$ ; Fig. S3). Overall, urban  
286 fractional dose in the range  $<0.002\text{--}5.4\%$  was received at the marine site. Across all samples  
287 at the marine site,  $D_u$  is found to be significantly correlated with the pollutant sum  
288 concentrations  $\Sigma_{6-4r}$  PAH and  $\Sigma_{11\ 3-4r}$  NPAH ( $R^2 = 0.61$  and  $0.69$ , respectively, both  $P > 0.99$ ).

289 From the marine sites data set, subsets of each two samples are formed, representing  
290 minimum (i.e., almost no influence from industrialised area 48 hours prior to arrival  
291 (hereforth called ‘marine background’, urban fractional dose  $D_u = 0.4\%$ ) and maximum  
292 observed influence (hereforth called ‘background with urban influence’,  $D_u = 3.1\%$ ; SM Table  
293 S2, Figure S3). The results for these subsets are listed in Tables 1-3. Such classification was  
294 not deemed meaningful for the samples collected at the continental site, as the relevant source  
295 distribution in central Europe was too homogeneous during this episode. Advection was  
296 mostly from northwest and partly from easterly directions, with air mass origin (time horizon  
297 of 10 days) mostly in central Europe and, to a lesser extent in eastern Europe and the western  
298 Balkans. The  $\text{NO}_2$  ( $1.2\text{--}2.6$  ppbv), total carbon ( $3\text{--}6$   $\mu\text{g m}^{-3}$ ) and  $\text{PM}_{10}$  ( $10.7\text{--}46.3$   $\mu\text{g m}^{-3}$ )  
299 levels during the campaign reflect continental background conditions.

300 The 4-ring PAH concentrations in samples from the continental site on the one hand, and in  
301 background air with urban influence collected at the marine site (urban areas  $300\text{--}500$  km  
302 away) on the other hand, are similar (Table 2). Also, the substance patterns are more similar  
303 than when relating all samples at the marine site i.e.,  $R^2 = 0.88$  ( $P > 0.999$ , t-test) instead of  $R^2$   
304  $= 0.76$ . The investigation of the diffusive air-surface exchange processes during the  
305 measurements presented here showed thatAt both sites, 4rPAHs were in fact influenced by  
306 secondary emissions, namely throughout day and night from the soil at the continental site (by  
307 average  $16.3$  and  $9.3$   $\text{pg m}^{-2} \text{h}^{-1}$  for FLT and PYR, respectively; Degrendele et al., 2016) or  
308 occasionally from surface seawater at the marine site (during at least 1 day-time interval out of  
309 in total 3 of this data subset; Lammel et al., 2016). In the data set from the continental site, we  
310 study day/night (D/N) effects (subsets listed in Tables 1-3, too): PAH concentrations ( $c_{\text{tot}}$ )  
311 were  $\approx 60\%$  higher during the day than during the night, while  $c_{\text{tot}}$  of NPAH-concentrations  
312 were by average  $\approx 5\%$  lower during the day (Table 2). NPAHs are subject to photolysis, while  
313 PAHs are not. At the site, the PAH concentrations were driven by re-volatilisation from soil,  
314 determined by temperature variation (Degrendele et al., 2016). For NPAHs (partly primary  
315 emitted) this indicates that the higher emissions during the day (due to re-volatilisation and  
316 road traffic) were compensated by shorter lifetime. NPAH lifetimes may be limited by  
317 (because of heterogeneous photolysis, but available kinetic data are scarce and limited to few

318 aerosol types (Fan et al., 1996; Feilberg and Nielsen, 2000, 2001; García-Berríos et al.,  
319 2017). ~~Also, The same could be reflected in~~ different NPAH/PAH ratios (the potential NPAH  
320 yields), which were 5.6% and 8.9% at the marine and continental sites, respectively, reflect  
321 the combination of emission sources and photochemical sinks. The NPAH/PAH ratios at the  
322 two sites were influenced by similar substance patterns upon emission, similar irradiation  
323 (summer, no or almost no clouds) and deposition velocities ( $\theta$  in the range 0.05-0.20 for  $\Sigma_{11} 3-$   
324  $4rNPAH$  and  $\Sigma_6 4rPAH$ , no precipitation), but different re-volatilisation fluxes and different  
325 characteristic transport times elapsed. Distance to major urban source areas was 300- >1000  
326 km at the marine and 100-500 km at the continental site. The NPAH/PAH ratios being lower  
327 at the more distant receptor site, the marine site, may suggest that photochemical degradation  
328 of NPAHs along transport was on average faster than degradation of the precursors. ~~xxx~~

329  
330 The NPAH levels observed in marine background air are the lowest ever reported.  
331 Remarkably, the concentrations are much lower, by more than one order of magnitude, than  
332 one decade before at the same site during the same season (Tsapakis and Stephanou, 2007).  
333 The concentrations observed now are a factor of 4-10 lower than in a forest site in Amazonia  
334 two decades before (which might have been influenced by biomass burning emissions), ~~and~~  
335 ~~also~~ a factor of 3 lower (for 2NPYR) than observed at an extremely remote site in the  
336 Himalayas two decades before (Ciccioli et al., 1996), ~~and comparable to a high altitude site in~~  
337 ~~the Alps (with the exception of 2NPYR which was observed one order of magnitude higher~~  
338 ~~there in winter; Albinet et al., 2008a; Table 3)~~. The NPAH levels observed at the marine site  
339 with influence of pollution and at the continental site are comparable, but also at the lower end  
340 of the range spanned by previous observations at rural and remote sites (Table 3).

341

### 342 **Gas-particle partitioning**

343 The time-weighted mean NPAH phase distributions ( $\Sigma_{11} 3-4rNPAH$ ) differ, corresponding to  $\theta$   
344 = 0.05 and 0.17 at the marine and continental sites, respectively, – despite similar  
345 temperatures (Table 1). In contrast and despite of similar temperature ranges, the 4-ring  
346 PAHs' ( $\Sigma_6 4rPAH$ ) particulate mass fraction was higher at the marine than at the continental  
347 site ( $\theta = 0.42$  and  $0.20$ , respectively). ~~The NPAH mass size distribution had its maximum in~~  
348 ~~the <0.49  $\mu m$  size range at both sites. The 4-ring PAHs mass size distribution had 2 maxima,~~  
349 ~~<0.49  $\mu m$  and between 0.95 and 1.5  $\mu m$ , at the marine site, but one at <0.49  $\mu m$  at the~~

350 ~~continental site (Table 1). This is probably related to the presence of aged aerosol at the~~  
351 ~~marine site vs. a larger contribution of fresh aerosols at the continental site. This is,~~  
352 ~~furthermore, supported by the analysis of air mass origins that shows significant influence of~~  
353 ~~urban areas for only few samples at the marine and for all samples at the continental site (SM~~  
354 ~~S2).~~

355 Both 4-ring PAHs and 3-4 ring NPAHs were more associated with PM in polluted air than in  
356 clean air. This trend is weak for PAHs with  $\theta = 0.02$  for  $\Sigma_6 4rPAH$  in marine background but  
357  $0.07$  in background with urban influence (and  $\theta = 0.09$  and  $0.20$  for CHR; Table 2), but is  
358 obviously strong for NPAHs, namely  $\theta = 0.19$  for 2NPYR in marine background but  $0.69$  in  
359 background with urban influence,  $\approx 0.93$  in polluted continental air, and  $\theta = 0.01$  for  $\Sigma_{11} 3-$   
360  $4rNPAH$  in marine background but  $0.22$  in background with urban influence (Table 2). The  
361 urban influenced air at the marine site is also reflected in a much higher OC (a factor of 3  
362 higher than the all-campaign mean) and elevated EC, (less prominent,  $\approx 50\%$  above mean).  
363 This confirms the understanding that gas-particle partitioning of both PAHs (Lohmann and  
364 Lammel, 2004; Shahpoury et al., 2016) and NPAHs (Tomaz et al., 2016) is mostly determined  
365 by absorption in POM and adsorption to soot. When comparing polluted air at the continental  
366 site and background with urban influence at the marine site, a strong shift of  $\Sigma_6 4rPAH$   
367 towards the particulate phase,  $\theta \approx 0.21$  vs.  $0.07$ , respectively, is found, while for  $\Sigma_{11} 3-4rNPAH$   
368  $\theta$  are similar i.e.,  $\approx 0.16$  vs.  $0.22$ , respectively. This phase partitioning trend of the 4rPAHs  
369 could be explained by sorption to EC, which is a factor of  $\approx 2$  higher, but not by OC (only  
370  $\approx 20\%$  higher). In conclusion, these observations consistently indicate that sorption to soot is  
371 less significant for gas-particle partitioning of NPAHs than for PAHs.

372 While NPAHs were significantly phase-shifted ( $\theta = 0.24$  during day-time but  $\theta = 0.58$  during  
373 night-time), this was not the case for 4rPAHs ( $\theta = 0.18$  during day-time and  $\theta = 0.23$  during  
374 night-time). This is in line with the perception that the temperature sensitivity of phase change  
375 is stronger for the substance class with stronger molecular interactions in the condensed phase,  
376 NPAHs. E.g., the enthalpies of phase change between air and OC of FLT and NFLT are  $-98$   
377 and  $-75 \text{ kJ mol}^{-1}$ , respectively (OC represented by DMSO; ACD, 2015).

378 Good agreement is found for the prediction of NPAH partitioning using the multi-phase (3-  
379 phase) ppLFER with most values predicted within one order of magnitude of the observed  
380 values (Fig. 2; quantification of deviations in S2.3.1). While the sensitivity of assumptions  
381 regarding PM phase composition, made in the model do not contribute significantly to the

382 deviations ( $\ll 1$ , log  $K_p$  units), a significant part can be attributed to the usage of estimated  
383 solute-specific Abraham solvation parameters (taken from ACD, 2015), in lack of  
384 experimentally based descriptors. E.g., for an urban site (Tomaz et al., 2016) it was found that  
385 experimentally based descriptors used for 9NPAH lead to better predictions than the estimated  
386 descriptors i.e., RMSEs differed by 0.43 log units. The agreement of the ppLFER prediction is  
387 better than assuming absorption (into OM) to be the only relevant process ( $K_{oa}$  model; see  
388 S2.3.2, Fig. S5). The same was found when studying gas-particle partitioning of NPAHs in  
389 urban air (Tomaz et al., 2016). This supports the perception that gas-particle partitioning of  
390 NPAHs is governed by various molecular interactions with OM, with its polarity being well  
391 represented by DMSO, better than by octanol. Earlier, it had been found for eight 3-4rNPAHs  
392 at urban and rural sites (Li et al., 2016) that the dual model, assuming adsorption (to soot) and  
393 absorption (into OM) predicts better than single adsorption (to the total aerosol surface i.e.,  
394 Junge-Pankow) or single absorption ( $K_{oa}$ ) models do.

395 The interactions with the aerosol matrix of 9NPHE (continental site) and 5NACE, 2NFLN,  
396 2NFLT and 1NPYR (marine site) are less well represented than other NPAHs by the model as  
397 suggested by low slopes of their  $\log K_p \text{ experimental} / \log K_p \text{ predicted}$  relationships. The reason is  
398 unknown. Moreover, sampling or sample handling artefacts cannot be excluded, even so same  
399 temperature range, sampler and sampling protocols applied across sites with both satisfactory  
400 and deficient agreement between predicted and observed  $K_p$ . Further conclusions are not  
401 supported by the limited amount of data and uncertainties on both the model (estimated  
402 ppLFER parameters) and experimental (concentrations close to LOQ) sides.

#### 403 404 **Mass size distribution**

405 The NPAH mass size distribution had its maximum in the  $<0.49 \mu\text{m}$  size range at both sites.  
406 The 4-ring PAHs mass size distribution had 2 maxima,  $<0.49 \mu\text{m}$  and between 0.95 and 1.5  
407  $\mu\text{m}$ , at the marine site, but one at  $<0.49 \mu\text{m}$  at the continental site (Table 1). This is probably  
408 related to the presence of aged aerosol at the marine site vs. a larger contribution of fresh  
409 aerosols at the continental site. This is, furthermore, supported by the analysis of air mass  
410 origins that shows significant influence of urban areas for only few samples at the marine and  
411 for all samples at the continental site (SM S2).

412 Sums of NPAHs' and PAHs' mass size distributions are found unimodal with the maximum  
413 in particles  $<0.49 \mu\text{m}$ , except PAHs at the marine site, which shows a second maximum  
414 between 1.5 and 3.0  $\mu\text{m}$  (Fig. 3). At the marine site, 50 and 69% of 1NPYR and 2NFLT,

415 respectively, were found associated with particles  $<0.45\mu\text{m}$  and 68 and 86%, respectively,  
416 with particles  $<0.95\mu\text{m}$ , and even more, 83% and 100%, respectively, with particles  $<0.45\mu\text{m}$   
417 at the continental site.

418  $\Sigma_6 4\text{rPAH}$  mass size distributions are shifted to larger particles in background with urban  
419 influence as compared to marine background air (both collected at the marine site) i.e., MMD  
420 = 0.19 and 0.28, respectively. However, such a trend is not apparent for NPAHs (Table 2). The  
421 size shift of PAHs is not corresponding to the  $\text{PM}_{10}$  mass size distribution: The MMD of  $\text{PM}_{10}$   
422 for all samples collected at the marine site was  $0.58\mu\text{m}$ , while it was  $1.13$  and  $0.62\mu\text{m}$  in the  
423 marine background and background with urban influence data subsets, respectively. The  $\text{PM}_{10}$   
424 as well as the OC mass size distributions were bimodal with maxima corresponding to  $<0.49$   
425  $\mu\text{m}$  and  $3.0\text{-}7.2\mu\text{m}$  particles (MMDs listed in Table 2), while the EC mass size distribution  
426 was unimodal, with the maximum concentration in the finest fraction. At the continental site,  
427 the  $\Sigma_{11} 3\text{-}4\text{rNPAH}$  mass size distribution was bimodal with maxima corresponding to  $<0.49$   
428  $\mu\text{m}$  and  $7.2\text{-}10\mu\text{m}$  particles, while the  $\Sigma_6 4\text{rPAH}$  mass size distribution was unimodal, with the  
429 maximum concentration in the finest fraction (for all samples as well as for day and night data  
430 subsets; Table 1).

431 The formation of a second maximum, at larger particles than emitted, reflects the  
432 redistribution of semivolatile organics in an aged aerosol, hence, is expected at receptor sites  
433 such as the marine site. This was also observed in polluted air, at rural and suburban sites, but  
434 not at traffic sites or in winter at a rural site, when primary emissions dominated (unimodal;  
435 Albinet et al., 2008b; Ringuet et al., 2012b).

436

#### 437 **Substance patterns and NPAH formation during long-range atmospheric transport**

438 Among the targeted NPAHs and apart from NNAPs, which were highest concentrated,  
439 2NFLT and 3NPHE prevailed at the marine site (accounting together for  $\approx 60\%$  of the NPAH  
440 mass, excluding the NNAPs), while at the continental site 9NANT and 2NFLT prevailed  
441 (accounting for  $\approx 65\%$  together) (Fig. 1, summarised in Fig. S4). The analytical method did not  
442 separate the isomers 2NFLT and 3NFLT, but at receptor sites, far from diesel emissions it  
443 appears justified to assume  $c_{2\text{NFLT}} \gg c_{3\text{NFLT}}$  (Finlayson-Pitts and Pitts, 2000; Zimmermann et  
444 al., 2012). The ratio 1NPYR/2NPYR is higher,  $\approx 1$ , at the continental site than at the marine  
445 site ( $\approx 0.25$ ), which reflects the significance of primary sources for polluted air (Atkinson and  
446 Arey, 1994; Finlayson-Pitts and Pitts, 2000; Zimmermann et al., 2012). This ratio was found

447 similarly high or even higher at urban sites (Ringuet et al., 2012c; Tomaz et al., 2016).  
448 Similarly, the ratio 2NFLT/1NPYR, the concentration of a secondarily formed over a primary  
449 emitted NPAH, has been used as indicator for fresh emissions (if  $< 5$ ) vs. photochemically  
450 aged air mass (Keyte et al., 2013). These values were  $\gg 5$  in 21 out of 22 and 7 out of 8  
451 samples at the marine and continental sites, respectively. The only sample collected at the  
452 continental site with elevated primary NPAH (2NFLT/1NPYR = 4.3) was possibly influenced  
453 by emissions from Budapest, which was passed by the advected air within the last hours  
454 before arrival. The only sample collected at the marine site with elevated primary NPAH  
455 (2NFLT/1NPYR = 5.9) was indeed directly influenced by emissions into the boundary layer  
456 above the Izmir and Istanbul metropolitan areas (urban fractional dose  $D_u = 5.0\%$  for samples  
457 no. 1 and 2 in Fig. S3). In conclusion, these results from receptor / background sites confirm  
458 the existing knowledge about primary emitted and secondarily formed NPAHs.

459 The ratio of two secondarily formed NPAHs, 2NFLT/2NPYR, indicative for day- vs. night-  
460 time formation paths (Atkinson and Arey, 1994; Ciccioli et al., 1996), is found  $\approx 2$  at the  
461 marine and  $\approx 8$  at the continental site (normalised to the precursor ratio i.e.,  
462 2NFLT/2NPYR/(FLT/PYR); Table 4). Such low values point to day-time (OH initiated)  
463 formation, while night-time ( $\text{NO}_3$  initiated) formation was negligible, practically excluded at  
464 the marine site. This is in line with the perception that  $\text{NO}_3$  must have been very low in this  
465 remote environment. ( $\text{NO}_x$  levels at the marine site were in the range 0.2-0.6 ppbv). A similar  
466 conclusion had been drawn in a semi-rural environment (Feilberg et al., 2001).

467 For 2NFLT and 2NPYR (secondary sources only) and for 1NPYR, which has mostly primary  
468 sources (Finlayson-Pitts and Pitts, 2000; Ringuet et al., 2012a; Jariyasopit et al., 2014a,  
469 2014b) we infer the potential yields (Table 4). Here, yield is defined as  $c_{\text{NPAH}}/c_{\text{PAH}}$  (total  
470 concentrations). This yield is called 'potential' as it ~~and~~ reflects an upper estimate, as other  
471 PAH photochemical sinks, such as formation of oxy-PAHs, are neglected. The yield of  
472 2NFLT in polluted air exceeds the one in background air only slightly, while the yield of  
473 2NPYR in polluted air exceeds the one in background air much more (a factor of 3 higher).

474 As expected, the highest potential yield of 1NPYR is found in polluted air (both sites),  
475 reflecting the dominance of primary emissions of 1NPYR. Similarly, higher yields of  
476 secondary NPAHs are found for marine background air compared to background air with  
477 urban influence (marine site), reflecting the longer reaction times elapsed since PAH  
478 emission. The yield for 2NFLT,  $c_{2\text{NFLT}}/c_{\text{FLT}}$ ,  $\approx 2$ -4% at both sites ranges higher than the one  
479 for 2NPYR,  $c_{2\text{NPYR}}/c_{\text{PYR}}$ , which is found  $\approx 0.5$ -2%. Note that because of the co-elution of

480 2NFLT and 3NFLT, and neglect of 3NFLT, the so derived values of  $c_{2NFLT}/c_{FLT}$  represent  
481 actually upper estimates. Apart from sites which were immediately influenced by PAH sources  
482 (road traffic, power plant, biomass burning), only very few studies reported NPAH together  
483 with precursor data in both phases of ambient air.  $c_{2NPYR}/c_{PYR} = 1.0\%$ , similar to our finding at  
484 remote sites, but a very high  $c_{2NFLT}/c_{FLT} = 12.9\%$  were reported from a suburban site in France  
485 in summer during day-time (corresponding values for night-time were 2.0 and 9.4%,  
486 respectively; Ringuet et al., 2012c). 2NFLT was not separated from 3NFLT (similar to our  
487 data set). A suburban site will be influenced by direct 3NFLT emissions, such that  $c_{2NFLT}/c_{FLT}$   
488 is an upper estimate. Much lower ratios,  $c_{2NFLT}/c_{FLT} = 0.20\%$  and  $c_{2NPYR}/c_{PYR} = 0.08\%$  were  
489 reported as the median values for 90 sites of various categories, rural and urban, in northern  
490 China in summer (Lin et al., 2015). These yields are somewhat higher for the subset of the  
491 rural sites. The potential yields found at the marine site in our study are close to the yields for  
492 OH-initiated photochemistry observed in laboratory experiments under high  $NO_x$  conditions  
493 i.e., 3% for  $c_{2NFLT}/c_{FLT}$  and 0.5% for  $c_{2NPYR}/c_{PYR}$  (Atkinson and Arey, 1994).

494

#### 495 **4. Conclusions**

496 For the first time pollution contained in individual background air samples was quantified, by  
497 means of a fractional dose. The fractional dose indicated how much the collected volume of  
498 air had been exposed to an urban boundary layer within a given time horizon. This is found  
499 suitable to discriminate among samples and discuss results, clearly beyond qualitative  
500 reasoning on back trajectories alone. The concept could be applied to any type of  
501 georeferenced origin and might be useful to track the influence of land use of various kind, or  
502 ship and aircraft routes.

503 Our measurements confirmed occurrence of mutagenic NPAHs, earlier reported from polluted  
504 atmospheric environments of America, Europe and Asia, also for the European background  
505 atmosphere and the outflow of Europe. These substances obviously go into intercontinental  
506 transport and might be indeed ubiquitous. The ~~long-range-transport-potential-is-hardly-limited~~  
507 ~~by-the~~ mass size distribution, ~~which~~ is determined by the particle size upon emission (primary  
508 NPAHs) and condensation and redistribution in the aerosol along transport, hence, does not  
509 include the short-lived coarse mass fraction. This indicates a high long-range transport  
510 potential. However, the observation of 3.8 and 0.92  $\text{pg m}^{-3}$  of 2NFLT and 2NPYR,  
511 respectively, measured at the southeastern outflow of Europe (the lowest ever reported



512 concentrations; Table 3), may indicate that their abundance in the remote global environment  
513 could be less than anticipated. Earlier, this was based on a single measurement of 2NPYR, 3  
514  $\text{pg m}^{-3}$ , at an extremely remote site in central Asia two decades before (Ciccioli et al., 1996).  
515 Moreover, this air, classified as marine background, was not completely clean, but had been  
516 exposed to a non-zero fractional urban pollution dose (0.4% of the total, time horizon of 2  
517 days). More measurements at remote sites should verify [NPAH](#) levels globally. PAHs have  
518 been abated significantly in Europe during the last decades (EEA, 2014), which should also be  
519 reflected in long-term trends of their derivatives. However, a temporal trend for the Aegean or  
520 the southeastern outflow of Europe in summer cannot be inferred based on the current and the  
521 earlier (2002; Tsapakis and Stephanou, 2007) campaign data. NPAHs should be included in  
522 monitoring programmes to better assess the exposure of human health hazards of atmospheric  
523 pollution, even in remote areas.

524 Understanding of NPAH formation in ambient air is still rudimentary. Although our  
525 observations of a potential NPAH yield are in agreement with laboratory studies of OH-  
526 initiated photochemistry, the kinetics of NPAHs, both formation from parent PAHs and  
527 photolysis remains to be quantitatively studied under relevant conditions for the background  
528 atmosphere i.e., low  $\text{NO}_x$  and on various aerosol matrices including sea salt, respectively.  
529 [More studies into NPAH atmospheric fate, both field observations and kinetic data, are](#)  
530 [needed in order to assess and quantify spatial and temporal trends, the long-range transport](#)  
531 [potential and persistence.](#)

### 533 **Acknowledgements**

534 We thank Christos I. Efstathiou (Masaryk University), András Hoffer, Gyula Kiss (MTA-PE  
535 Air Chemistry Research Group, Veszprém), Jiří Kohoutek (MU), Giorgos Kouvarakis  
536 (University of Crete, Iraklion) and Lajos Szöke (Hungarian Meteorological Service) for on-  
537 site support, Giorgos Kouvarakis and Krisztina Labancz (Hungarian Meteorological Service)  
538 for meteorological and trace gas data, Michael H. Abraham (University College London) for  
539 providing ppLFR solute descriptors, Ignacio Pisso (NILU, Kjeller, Norway) for model post-  
540 processing scripts and Céline Degrendele (MU) and Manolis Tsapakis (Hellenic Centre for  
541 Marine Research, Gournes) for discussion. This research was supported by the Czech Science  
542 Foundation (n° P503 16-11537S), the Czech Ministry of Education, Youth and Sports (n°  
543 LO1214 and LM2015051), and the European Union FP7 (n° 262254, ACTRIS).

545 **Compliance with Ethical Standards** No potential conflicts of interest (financial or non-  
546 financial) exist.

### 547 **References**

548 Abraham, M.H.: Scales of solute hydrogen-bonding: their construction and application to physicochemical and  
549 biochemical processes. *Chem. Soc. Rev.*, 22, 73–83, 1993.

550 ACD: ACD/Labs Absolv Software, Advanced Chemistry Development, Toronto, Canada, 2015.

551 Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Simultaneous analysis of oxygenated and  
552 nitrated polycyclic aromatic hydrocarbons on standard reference material 1649a (urban dust) and on natural  
553 ambient air samples by gas chromatography-mass spectrometry with negative ion chemical ionisation. *J.*  
554 *Chrom. A*, 1121, 106-113, 2006.

555 Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Polycyclic aromatic hydrocarbons (PAHs),  
556 nitrated PAHs and oxygenated PAHs in ambient air of the Marseilles area (South of France): Concentrations  
557 and sources. *Sci. Total Environ.*, 384, 280-292, 2007.

558 Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., and Jaffrezo, J.L.: Nitrated and oxygenated  
559 derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys part 1:  
560 concentrations, sources and gas/particle partitioning. *Atmos. Environ.*, 42, 43-54, 2008a.

561 Albinet, A., Leoz-Garziandia, E., Budzinski, H., Villenave, E., and Jaffrezo, J.L.: Nitrated and oxygenated  
562 derivatives of polycyclic aromatic hydrocarbons in the ambient air of two French alpine valleys part 2: Particle  
563 size distribution. *Atmos. Environ.*, 42, 55-64, 2008b.

564 Alves, C.A., Vicente, A.M.P., Gomes, J., Nunes, T., Duarte, M., and Bandowe, B.A.M.: Polycyclic aromatic  
565 hydrocarbons (PAHs) and their derivatives (oxygenated-PAHs, nitrated-PAHs and azaarenes) in size-  
566 fractionated particles emitted in an urban road tunnel. *Atmos. Res.*, 180, 128-137, 2016.

567 Arey, J.: Atmospheric reactions of of PAHs including formation of nitro-arenes. In: *The handbook of*  
568 *Environmental Chemistry, Vol. 3I: PAHs and Related Compounds* (Neilson AH, ed.), Springer, Berlin, pp.  
569 347-385, 1998.

570 Arey, J., Atkinson, R., Zielinska, B., and McElroy, P.A.: Diurnal concentrations of volatile polycyclic aromatic  
571 hydrocarbons and nitroarenes during a photochemical air pollution episode in Glendora, California. *Environ.*  
572 *Sci. Technol.*, 23, 321-327, 1989.

573 Atkinson, R., and Arey, J.: Atmospheric chemistry of polycyclic aromatic hydrocarbons: Formation of  
574 atmospheric mutagens. *Environ. Health Persp.*, 102, 117-126. 1994.

575 Bamford, H.A., and Baker, J.E.: Nitro-polycyclic aromatic hydrocarbon concentrations and sources in urban and  
576 suburban atmospheres of the Mid-Atlantic region. *Atmos. Environ.*, 37, 2077-2091, 2003.

577 Barrado, A.I., García, S., Sevillano, M.L., Rodríguez, J.A., and Barrado, E.: Vapor-phase concentrations of  
578 PAHs and their derivatives determined in a large city: Correlations with their atmospheric aerosol. *Chemosph.*,  
579 93, 1678-1684, 2013.

580 Borbély-Kiss, I., Haszpra, L., Koltay, E., László, S., Mészáros, A., Mészáros, E., and Szabó, S.: Elemental  
581 concentrations and regional signatures in atmospheric aerosols over Hungary. *Physica Scripta*, 37, 299-304,  
582 1988.

583 Ciccioli, P., Cecinato, A., Brancaleoni, E., Frattoni, M., Zacchei, P., Miguel, A., and de Castro Vasconcellos, P.:  
584 Formation and transport of 2-nitrofluoranthene and 2-nitropyrene of photochemical origin in the troposphere.  
585 *J. Geophys. Res.*, 101, 19567-19582, 1996.

586 Claxton, L.D., Matthews, P.P., and Warren, S.H.: The genotoxicity of ambient outdoor air, a review: Salmonella  
587 mutagenicity. *Mutation Res. Rev.* 567, 347-399, 2004.

588 Degrendele, C., Audy, O., Hofman, J., Kučerik, J., Kukučka, P., Mulder, M.D., Přibylková, P., Prokeš, R., Šánka,  
589 M., Schaumann, G.E., and Lammel, G.: Diurnal variations of air-soil exchange of semi-volatile organic  
590 compounds (PAHs, PCBs, OCPs and PBDEs) in a central European receptor area. *Environ. Sci. Technol.*, 50,  
591 4278–4288, 2016.

592 Draxler, R.R., and Rolph, G.D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model  
593 access via NOAA ARL READY. NOAA Air Resources Laboratory, Silver Springs, USA. 2003, Available from:  
594 <http://www.arl.noaa.gov/ready/hysplit4.html>.

595 EEA: European emission inventory report 1990-2012 under the UNECE Convention on Long-range Transboundary  
596 Air Pollution (LRTAP), European Environment Agency Technical Report No. 12/2014, Copenhagen, 2014, 132  
597 pp.

598 EMEP: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components. Co-operative  
599 Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe, EMEP  
600 Report No. 1/2015, Oslo, 150+78 pp. Available from:  
601 [http://emep.int/publ/reports/2015/EMEP\\_Status\\_Report\\_1\\_2015.pdf](http://emep.int/publ/reports/2015/EMEP_Status_Report_1_2015.pdf), 2015

602 Endo, S., and Goss, K.U.: Applications of poly-parameter linear free energy relationships in environmental  
603 chemistry. *Environ. Sci. Technol.*, 48, 12477–12491, 2014.

604 Fan, Z.H., Kamens, R.M., Hu, J.X., Zhang, J.B., and McDow, S.: Photostability of nitro-polycyclic aromatic  
605 hydrocarbons on combustion soot particles in sunlight. *Environ. Sci. Technol.*, 30, 1358–1364, 1996.

606 Feilberg, A., and Nielsen, T.: Effect of aerosol chemical composition on the photodegradation of nitro-PAHs.  
607 Environ. Sci. Technol., 34, 789-797, 2000.

608 Feilberg, A., and Nielsen, T.: Photodegradation of nitro-PAHs in viscous organic media used as models of organic  
609 aerosols. Environ. Sci. Technol., 35, 108-113, 2001.

610 Feilberg, A., Poulsen, M.W.B., Nielsen, T., and Skov, H.: Occurrence and sources of particulate nitro polycyclic  
611 aromatic hydrocarbon in ambient air in Denmark. Atmos. Environ., 35, 353-366, 2001.

612 Finizio, A., Mackay, D., Bidleman, T., and Harner, T.: Octanol-air partition coefficient as a predictor of  
613 partitioning of semi-volatile organic chemicals to aerosols. Atmos. Environ., 31, 2289-2296, 1997.

614 Finlayson-Pitts, B.J., and Pitts, J.N.: Chemistry of the Upper and Lower Atmosphere: Theory, Experiments,  
615 Application, San Diego (Academic Press), USA, 2000

616 García-Berrios, Z.I., Arce, R., Burgos-Martínez, M., and Burgos-Polanco, N.D.: Phototransformations of  
617 environmental contaminants in models of the aerosol: 2 and 4-nitropyrene. J. Photochem. Photobiol. A, 332,  
618 131-140, 2017.

619 Garner, R.C., Stanton, C.A., Martin, C.N., Chow, F.L., Thomas, W., Hübner, D., and Herrmann, R.: Bacterial  
620 mutagenicity and chemical analysis of polycyclic aromatic hydrocarbons and some nitro derivatives in  
621 environmental samples collected in West Germany. Environ Mutagen, 8, 109-117, 1986.

622 Goss, K.U.: Predicting the equilibrium partitioning of organic compounds using just one linear solvation energy  
623 relationship (LSER). Fluid Phase Equilib., 233, 19-22, 2005.

624 Goss, K.U., and Schwarzenbach, R.P.: Linear free energy relationships used to evaluate equilibrium partitioning  
625 of organic compounds. Environ. Sci. Technol., 35, 1-9, 2001.

626 Grosjean, D., Fung, K., and Harrison, J.: Interactions of polycyclic aromatic hydrocarbons with atmospheric  
627 pollutants. Environ. Sci. Technol., 17, 673-679, 1983.

628 Halsall, C.J., Sweetman, A.J., Barrie, L.A., and Jones, K.C.: Modelling the behaviour of PAHs during  
629 atmospheric transport from the United Kingdom to the Arctic. Atmos. Environ. 35, 255-267.

630 Hayakawa, K. (2016) Environmental behaviors and toxicities of polycyclic aromatic hydrocarbons and  
631 nitropolycyclic aromatic hydrocarbons. Chem. Pharm. Bull., 64, 83-94, 2001.

632 Inomata, S., Fushimi, A., Sato, K., Fujitani, Y., and Yamada, H.: 4-Nitrophenol, 1-nitropyrene, and 9-  
633 nitroanthracene emissions in exhaust particles from diesel vehicles with different exhaust gas treatments.  
634 Atmos. Environ., 110, 93-102, 2015.

635 Jaenicke, R.: Aerosol physics and chemistry. Landolt-Börnstein Neue Serie b, 4, 391-457, 1988.

636 Jariyasopit, N., Zimmermann, K., Schrlau, J., Arey, J., Atkinson, R., Yu, T.W., Dashwood, R.H., Tao, S., and  
637 Massey Simonich, S.L.: Novel NPAH formation from heterogeneous reactions of PAHs with NO<sub>2</sub>, NO<sub>3</sub>/N<sub>2</sub>O<sub>5</sub>,  
638 OH radicals, and OH radicals: Prediction, laboratory studies and mutagenicity. Environ. Sci. Technol., 48,  
639 412-419, 2014a.

640 Jariyasopit, N., Zimmermann, K., Schrlau, J., Arey, J., Atkinson, R., Yu, T.W., Dashwood, R.H., Tao, S., and  
641 Massey Simonich, S.L.: Heterogeneous reactions of particulate matter-bound PAHs and NPAHs with  
642 NO<sub>3</sub>/N<sub>2</sub>O<sub>5</sub>, OH radicals, and O<sub>3</sub> under simulated long-range atmospheric transport conditions: Reactivity and  
643 mutagenicity. Environ. Sci. Technol., 48, 10155-10164, 2014b.

644 Keyte, I.J., Harrison, R.M., and Lammel, G.: Chemical reactivity and long-range transport potential of polycyclic  
645 aromatic hydrocarbons – a review. Chem. Soc. Rev., 42, 9333-9391, 2013.

646 Keyte, I.J., Albinet, A., and Harrison, R.M.: On-road traffic emissions of polycyclic aromatic hydrocarbons and  
647 their oxy- and nitro- derivative compounds measured in road tunnel environments. Sci. Total Environ., 566-  
648 567, 1131-1142, 2016.

649 Kouvarakis, G., Tsigaridis, K., Kanakidou, M., and Mihalopoulos, N.: Temporal variations of surface regional  
650 background ozone over Crete Island in the Southeast Mediterranean. J. Geophys. Res., 105, 4399-4407, 2000.

651 Lafontaine, S., Schrlau, J., Butler, J., Jia, Y.L., Harper, B., Harris, S., Bramer, L.M., Waters, K.M., Harding, A.,  
652 and Massey Simonich, S.L.: Relative influence of trans-Pacific and regional atmospheric transport of PAHs in  
653 the Pacific Northwest, U.S. Environ. Sci. Technol., 49, 13807-13816, 2015.

654 Lammel, G., Audy, O., Besis, A., Efstathiou, C., Eleftheriadis, K., Kohoutek, J., Kukučka, P., Mulder, M.D.,  
655 Přibylková, P., Prokeš, R., Rusina, T., Samara, C., Sofuoglu, A., Sofuoglu, S.C., Taşdemir, Y., Vassilatou, V.,  
656 Voutsas, D., and Vrana, B.: Air and seawater pollution and air-sea gas exchange of persistent toxic substances  
657 in the Aegean Sea: spatial trends of PAHs, PCBs, OCPs and PBDEs. Environ. Sci. Pollut. Res., 22, 11301-  
658 11313, 2015.

659 Lammel, G., Meixner, F.X., Vrana, B., Efstathiou, C., Kohoutek, J., Kukučka, P., Mulder, M.D., Přibylková, P.,  
660 Prokeš, R., Rusina, T.S., Song, G.Z., and Tsapakis, M.: Bi-directional air-sea exchange and accumulation of  
661 POPs (PAHs, PCBs, OCPs and PBDEs) in the nocturnal marine boundary layer. Atmos. Chem. Phys., 16,  
662 6381-6393, 2016.

663 Li, W., Shen, G.F., Yuan, C.Y., Wang, C., Shen, H.Z., Jiang, H., Zhang, Y.Y., Chen, Y.C., Su, S., Lin, N., and  
664 Tao, S.: The gas/particle partitioning of nitro- and oxy-polycyclic aromatic hydrocarbons in the atmosphere of  
665 northern China. *Atmos. Res.*, 172-173, 66-73, 2016.

666 Lin, Y., Qiu, X.H., Ma, Y.Q., Ma, J., Zheng, M., and Shao, M.: Concentrations and spatial distribution of  
667 polycyclic aromatic hydrocarbons (PAHs) and nitrated PAHs (NPAHs) in the atmosphere of North China, and  
668 the transformation from PAHs to NPAHs. *Environ. Pollut.*, 196, 164-170, 2015.

669 Lohmann, R., and Lammel, G.: Adsorptive and absorptive contributions to the gas particle partitioning of  
670 polycyclic aromatic hydrocarbons: State of knowledge and recommended parameterisation for modelling,  
671 *Environ. Sci. Technol.*, 38, 3793-3803, 2004.

672 Masclat, P., Pistikopoulos, P., Beyne, S., and Mouvier, G.: Long-range transport and gas/particle distribution of  
673 polycyclic aromatic hydrocarbons at a remote site in the Mediterranean sea, *Atmos. Environ.*, 22, 639-650,  
674 1988.

675 Melymuk, L., Bohlin-Nizzetto, P., Prokeš, R., Kukučka, P., and Klánová, J.: Sampling artifacts in active air  
676 sampling of semivolatile organic contaminants: Comparing theoretical and measured artifacts and evaluating  
677 implications for monitoring networks. *Environ. Pollut.*, 217, 97-106, 2016.

678 Mihalopoulos, N., Stephanou, E., Pilitsidis, S., Kanakidou, M., and Bousquet, P.: Atmospheric aerosol  
679 composition above the Eastern Mediterranean region. *Tellus B*, 49, 314-326, 1997.

680 Nielsen, T., Seitz, B., and Ramdahl, T.: Occurrence of nitro-PAH in the atmosphere in a rural area. *Atmos.*  
681 *Environ.*, 18, 2159-2165, 1984.

682 Pandis, S.N., Baltensperger, U., Wolfenbarger, J.K., and Seinfeld, J.H.: Inversion of aerosol data from the  
683 epiphaniometer. *J. Aerosol Sci.*, 22, 417-428, 1991.

684 Pankow, J.F.: Overview of the gas phase retention volume behaviour of organic compounds on polyurethane  
685 foam. *Atmos. Environ.*, 23, 1107-1111, 1989.

686 Pitts, J.N., Arey, J., Zielinska, B., Winer, A.M., and Atkinson, R.: Determination of 2-nitro-fluoranthene and 2-  
687 nitropyrene in ambient particulate organic matter: Evidence for atmospheric reactions. *Atmos. Environ.*, 19,  
688 1601-1608, 1985.

689 Ramdahl, T., Zielinska, B., Arey, J., Atkinson, R., Winer, A.M., and Pitts, J.N.: Ubiquitous occurrence of 2-  
690 nitro-fluoranthene and 2-nitropyrene in air. *Nature*, 321, 425-427, 1986.

691 Reisen, F., and Arey, J.: Atmospheric reactions influence seasonal PAH and nitro-PAH concentrations in the Los  
692 Angeles Basin. *Environ. Sci. Technol.*, 39, 64-73, 2005.

693 Ringuet, J., Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Reactivity of polycyclic aromatic  
694 compounds (PAHs, NPAHs and OPAHs) adsorbed on natural aerosol particles exposed to atmospheric  
695 oxidants. *Atmos. Environ.*, 61, 15-22, 2012.

696 Ringuet, J., Leoz-Garziandia, E., Budzinski, H., Villenave, E., and Albinet, A.: Particle size distribution of  
697 nitrated and oxygenated polycyclic aromatic hydrocarbons (NPAHs and OPAHs) on traffic and suburban sites  
698 of a European megacity: Paris (France). *Atmos. Chem. Phys.*, 12, 8877-8887, 2012b.

699 Ringuet, J., Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Diurnal/nocturnal concentrations  
700 and sources of particulate-bound PAHs, OPAHs and NPAHs at traffic and suburban sites in the region of Paris  
701 (France). *Sci. Total Environ.*, 437, 297-305, 2012c.

702 Schauer, C., Niessner, R., and Pöschl, U.: Analysis of nitrated polycyclic aromatic hydrocarbons by liquid  
703 chromatography with fluorescence and mass spectrometry detection: air particulate matter, soot, and reaction  
704 product studies. *Anal. Bioanal. Chem.*, 378, 725-736, 2004.

705 Schuetzle, D.: Sampling of vehicle emissions for chemical analysis and biological testing. *Environ. Health*  
706 *Perspect.*, 47, 65-80, 1983.

707 Shahpoury, P., Lammel, G., Albinet, A., Sofuoğlu, A., Dumanoglu, Y., Sofuoğlu, S.C., Wagner, Z., and Ždimal,  
708 J.: Model evaluation for gas-particle partitioning of polycyclic aromatic hydrocarbons in urban and non-urban  
709 sites in Europe – Comparison between single- and poly-parameter linear free energy relationships based on a  
710 multi-phase aerosol scenario. *Environ. Sci. Technol.*, 50, 12312-12319, 2016.

711 Stohl, A., Hitznerberger, M., and Wotawa, G.: Validation of the Lagrangian particle dispersion model  
712 FLEXPART against large scale tracer experiments. *Atmos. Environ.*, 32, 4245-4264, 1998.

713 Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion  
714 model FLEXPART version 6.2. *Atmos. Chem. Phys.*, 5, 2461-2474, 2005.

715 Tomaz, S., Shahpoury, P., Jaffrezou, J.L., Lammel, G., Perraudin, E., Villenave, E., and Albinet, A.: One year  
716 study of polycyclic aromatic compounds at an urban site in Grenoble (France): seasonal variations, gas/particle  
717 partitioning and cancer risk estimation. *Sci. Total Environ.*, 565, 1071-1083, 2016.

718 Tsapakis, M., and Stephanou, E.G.: Diurnal cycle of PAHs, nitro-PAHs and oxy-PAHs in a high oxidant capacity  
719 marine background atmosphere. *Environ. Sci. Technol.*, 41, 8011-8017, 2007.

720 Vincenti, M., Maurino, V., Minero, C., and Pelizzetti, E.: Determination of nitro-substituted polycyclic aromatic  
721 hydrocarbons in the Antarctic airborne particulate. *Intl. J. Environ. Anal. Chem.*, 79, 257-272, 2001.

722 | [Vogelezang, D. H. P., and Holtslag, A. A. M.: Evaluation and model impacts of alternative boundary-layer height](#)  
723 | [formulations. Bound.-Layer Met., 81, 245–269, 1996.](#)  
724 | Vrekoussis, M., Liakakou, E., Kocak, M., Kubilay, N., Oikonomou, K., Sciare, J., and Mihalopoulos, N. (2005)  
725 | Seasonal variability of optical properties of aerosols in the Eastern Mediterranean, Atmos. Environ. 39, 7083-  
726 | 7094.  
727 | Yamasaki, H., Kuwata, K., and Miyamoto, H.: Effects of ambient temperature on aspects of airborne polycyclic  
728 | aromatic hydrocarbons. Environ. Sci. Technol., 16, 189–194, 1982.  
729 | Zimmermann, K., Atkinson, R., Arey, J., Kojima, Y., and Inazu, K.: Isomer distributions of molecular weight 247  
730 | and 273 nitro-PAHs in ambient samples, NIST diesel SRM, and from radical-initiated chamber reactions.  
731 | Atmos. Environ., 55, 431-439, 2012.  
732 | Zimmermann, K., Jariyasopit, N., Massey Simonich, S.L., Tao, S., Atkinson, R., and Arey, J.: Formation of  
733 | NPAHs from the heterogeneous reaction of ambient particle-bound PAHs with N<sub>2</sub>O<sub>5</sub>/NO<sub>3</sub>/NO<sub>2</sub>. Environ. Sci.  
734 | Technol., 47, 8434-8442, 2013.  
735

736 Table 1. Overview time weighted mean concentrations in the particulate and gas phases, and  
 737 ambient temperature. Data subsets (B = background, P = polluted, D = day mean, N = night  
 738 mean) and mass size distribution (<0.45/0.45-0.95/0.95-1.5/1.5-3.0/3.0-7.2/7.2-10  $\mu\text{m}$   
 739 aerodynamic equivalent diameter) in brackets.

Site	Phase	$\Sigma_{11} 3\text{-}4\text{rNPAH}$ ( $\text{pg m}^{-3}$ )	$\Sigma_6 4\text{rPAH}$ ( $\text{pg m}^{-3}$ )
Marine	particulate (n = 8)	4.1 (3.5/0.6/0.2/0.03/0.03/0.00) (B: 0.2/P: 8.7)	43 (28/8.1/1.2/6.2/4.3/1.7) (B: 7.9/P: 42.4)
	gas (n = 21)	18.4 (B: 13.2/P:31.1)	403 (B:321/P:580)
	T( $^{\circ}\text{C}$ )	25.6 (B: 27.1/P: 22.0)	
Continental	particulate (n = 22)	24.3 (20.5/2.9/0.7/0.04/0.06/0.15) (D:13.9/N:34.6)	129 (87/28/12/0.6/0.0/0.0) (D:146/N:116)
	gas (n = 22)	34.2 (D:42.9/N:25.5)	517 (D:649/N:384)
	T( $^{\circ}\text{C}$ )	23.1 (D:28.8/N:17.5)	

740

741

742 Table 3. Comparison of total (g + p) concentrations in air,  $c_{tot}$  (pg/m<sup>3</sup>), with other  
 743 measurements at remote and rural sites

	1NPYR (pg/m <sup>3</sup> )	2NFLT (pg/m <sup>3</sup> )	2NPYR (pg/m <sup>3</sup> )	References
background CEu summer 2013	1.1	15 <sup>a</sup>	1.3	this work (continental)
E Mediterranean summer 2012	0.74	8.6 <sup>a</sup>	2.5	this work (marine)
E Mediterranean clean summer 2012	0.21	3.8 <sup>a</sup>	0.92	this work (marine background <sup>b</sup> )
E Mediterranean clean summer 2002	-	29	21	Tsapakis and Stephanou, 2007
Ross Sea coast, Antarctica	<0.02 <sup>c</sup>	<0.03 <sup>c</sup>		Vincenti et al., 2001
Himalayas, Nepal 1991	-	-	3	Ciccioli et al., 1996
Forest Amazonia 1993	2	15	8	
Rural Northern Germany 1991	-	-	3	
Rural Denmark winter-spring 1982	9±5 <sup>c</sup>	-	-	Nielsen et al., 1984
Semi-rural Denmark all year 1998-99	40	97	6.3	Feilberg et al., 2001
Remote Alps 2002	2.2	-	-	Schauer et al., 2004
Rural Alps 2002	6.6	-	-	
Rural Alps <sup>d</sup> winter 2002-03	21	96 <sup>a</sup>	81	Albinet et al., 2008a
Rural Alps <sup>d</sup> summer 2003	4.2	28 <sup>a</sup>	5.7	
<u>Remote Alps<sup>e</sup> winter 2002-03</u>	<u>2.4</u>	<u>1.3<sup>a</sup></u>	<u>14.8</u>	
<u>Remote Alps<sup>e</sup> summer 2003</u>	<u>0.6</u>	<u>1.8<sup>a</sup></u>	<u>0.7</u>	
Rural Southern France 2004	0.6	2.6 <sup>a</sup>	1.6	Albinet et al., 2007

744 <sup>a</sup> co-eluted with 3NFLT, assuming  $c_{3NFLT} = 0$

745 <sup>b</sup> samples No. 9, 10, 19 and 22 in Fig. S3

746 <sup>c</sup> particulate phase concentration only

747 <sup>d</sup> Val de Maurienne sites (Albinet et al., 2008a)

748 <sup>e</sup> Plan de l'Aiguille site (Albinet et al., 2008a)

749

750 Table 4: Selected 4-ring PAHs and primary and secondary 3-4 ring NPAH total (g + p) time-  
 751 weighted mean concentrations  $\pm\sigma$  ( $\mu\text{g m}^{-3}$ ). Potential yields,  $C_{\text{NPAH}}/C_{\text{PAH}}$ , in brackets.  $\sigma$  given  
 752 for  $n > 2$ .

Site		Marine			Continental
Data subset		all (n = 8 <sup>a</sup> )	marine background (n = 2 <sup>b</sup> )	background with urban influence (n = 2 <sup>c</sup> )	all (n = 22)
Primary	FLT	213±161	161	259	342±215
	PYR	146±130	103	188	226±131
Primary and secondary (potential yield)	2NFLN <sup>d</sup>	0.038±0.12	<0.18	0.15	0.034±0.044
	1NPYR	0.62±1.1 (0.4±0.2%)	0.21 (0.2%)	1.4 (0.7%)	1.1±0.6 (0.6±0.3%)
Secondary (yield)	2NFLT <sup>e</sup>	7.7±8.5 (3.6±2.0%)	1.68 (1.0%)	11.0 (4.2%)	15±10 (6.5±7.5%)
	2NPYR	2.2±2.6 (1.5±0.7%)	0.92 (0.9%)	3.3 (1.8%)	1.3±1.7 (0.74±1.09%)

753 <sup>a</sup> 8 filter and 21 PUF samples

754 <sup>b</sup> 2 filter and 6 PUF samples i.e., No. 9-10 and 19-22 in Fig. S3 (urban fractional dose  $D_u =$   
 755 0.4%)

756 <sup>c</sup> 2 filter and 5 PUF sample i.e., No. 1-2 and 15-18 in Fig. S3 ( $D_u = 3.1\%$ )


757 <sup>d</sup> no yield given as  $C_{\text{FLN}}$  not quantified

758 <sup>e</sup> co-eluted with 3NFLT, assuming  $C_{3\text{NFLT}} = 0$

759




760 Fig. 1: Time series of absolute (a, c;  $\text{pg m}^{-3}$ ) and relative (b, d) total (gas + particulate) NPAH  
 761 concentrations at the (a, b) marine (24 h means shown<sup>a</sup>) and (c, d) continental site (day / night  
 762 means)




767 <sup>a</sup> gas-phase (PUF) sampled day / night, particulate phase (filter) sampled 1-4 subsequent days /  
 768 nights, 4 during the period 7.-12.7.12  
 769

770 Fig. 2: Predicted versus experimental  $\log K_p$  ( $\text{m}^3 \text{air g}^{-1} \text{PM}$ ) for NPAHs using the multi-phase  
771 ppLFER model at the (a) marine and (b) continental site

772 a.




773  
774 b.



775  
776  
777

778 Fig. 3. Time-weighted mean  $\Sigma_6$  4rPAH and  $\Sigma_{11}$  3-4rNPAH mass size distributions ( $\text{pg m}^{-3}$ ) at  
 779 the marine and continental sites. ~~Upper cut off of impactor stage given in  $\mu\text{m}$  of aerodynamic~~  
 780 ~~particle size.~~ The error bars show the standard deviation from the campaign mean.



781

782

Table 2. Total (g + p) time-weighted concentrations,  $c_{tot}$  ( $\text{pg m}^{-3}$ ), particulate mass fraction,  $\theta = c_p / c_{tot}$ , and mass median diameter (MMD,  $\mu\text{m}$ ), of of 2-4-ring NPAHs and 4-ring PAHs at the marine (as ‘mean (background mean/ urban influence mean)’,  $n = 8(2^a/2^b)$ ) and continental (as ‘mean (day mean/ night mean)’,  $n = 22(11/11)$ ) sites, together with temperature and supporting aerosol parameters ( $\text{PM}_{10}$  and carbonaceous mass fractions). LOQ = limit of quantification, nd = no data.

	Marine			Continental		
	$c_{tot}$ ( $\text{pg m}^{-3}$ )	$\theta$	MMD ( $\mu\text{m}$ )	$c_{tot}$ ( $\text{pg m}^{-3}$ )	$\theta$	MMD ( $\mu\text{m}$ )
FLT	226 (161/259)	0.07 (0.03/0.07)	0.58 (0.43/0.52)	342 (432/251)	0.11 (0.11/0.12)	0.062 (0.101/0.034)
PYR	158 (103/188)	0.04 (0.01/0.05)	0.21 (0.022/0.22)	226 (276/176)	0.18 (0.18/0.19)	0.075 (0.105/0.055)
BBN	4.1 (2.0/5.5)	0.01 (nd/0.05)	0.022 (nd/0.022)	15 (16/13)	0.61 (0.58/0.65)	0.079 (0.127/0.053)
BAA	2.8 (<LOQ/3.4)	0.28 (nd/0.29)	0.022 (nd/0.022)	16 (14/18)	0.91 (0.90/0.92)	0.070 (0.090/0.060)
TPH	12 (8.5/14)	0.02 (nd/0.05)	0.022 (nd/0.022)	23 (26/21)	0.51 (0.41/0.63)	0.070 (0.090/0.057)
CHR	23 (10/29)	0.22 (0.09/0.20)	0.15 (0.022/0.15)	41 (44/38)	0.75 (0.71/0.80)	0.074 (0.105/0.055)
$\Sigma_6$ 4rPAH	426 (284/499)	0.07 (0.02/0.07)	0.31 (0.19/0.28)	663 (808/517)	0.21 (0.19/0.25)	0.071 (0.10/0.051)
3NACE	0.21 (0.17/0.39)	0.05 (0.00/0.14)	0.022 (nd/0.022)	1.0(1.0/1.0)	0.05 (0.01/0.11)	0.022 (nd/0.022)
5NACE	1.8 (1.5/2.0)	0.07 (0.00/0.00)	0.022 (nd/nd)	6.8 (7.6/6.0)	0.03 (0.01/0.05)	0.022 (0.022/0.022)
2NFLN	0.01 (<LOQ/0.15)	0.02 (nd/0.00)	1.19 (nd/nd)	0.035 (0.035/0.034)	0.00 (0.00/0.00)	nd
9NPHE	0.73 (0.84/0.55)	0.00 (0.00/0.00)	nd	0.21 (0.28/0.13)	0.36 (0.43/0.20)	0.022 (0.022/nd)

3NPHE	4.8 (3.4/5.0)	0.00 (nd/nd)	nd	7.4 (10.0/4.8)	0.24 (0.15/0.44)	0.109 (0.067/0.116)
9NANT	4.2 (1.1/8.2)	0.00 (0.00/0.00)	nd	22 (22/22)	0.23 (0.14/0.33)	0.022 (0.022/0.022)
2NFLT <sup>c</sup>	8.6 (3.8/11)	0.32 (nd/0.45)	0.040 (nd/0.080)	15 (13/18)	0.78 (0.54/0.95)	0.054 (0.035/0.050)
1NPYR	0.75 (0.21/1.4)	0.33 (0.00/0.58)	0.061 (nd/0.14)	1.1 (1.1/1.2)	0.82 (0.76/0.88)	0.030 (0.031/0.029)
2NPYR	2.5 (0.92/3.3)	0.53 (0.19/0.69)	0.058 (0.060/0.055)	1.3 (0.73/2.0)	0.93 (0.83/0.97)	0.070 (0.040/0.061)
7NBAA	<LOQ	nd	nd	2.5 (0.77/4.2)	0.91 (0.56/0.97)	0.074 (0.038/0.057)
6NCHR	0.02 (<LOQ/0.07)	1.00 (nd/1.00)	2.12 (nd/2.12)	0.01 (<LOQ/0.02)	1.00 (nd/1.00)	0.022 (nd/0.022)
Σ <sub>113</sub> - 4rNPAH	23.7 (11.8/32.0)	0.22 (0.01/0.22)	0.34(0.33/0.34)	58 (56/59)	0.16 (0.13/0.17)	0.039 (0.036/0.040)
PM <sub>10</sub> (µg/m <sup>3</sup> )	34.9 (21.0/55.5)		0.58 (1.13/0.62)	22.1 (19.5/24.5)		nd
EC (µg/m <sup>3</sup> )	0.11 (0.09/0.17)		0.03(0.05/0.03)	0.31 (0.28/0.33)		0.21(0.19/0.22)
OC (µg/m <sup>3</sup> )	1.9 (1.5/3.0)		0.17(0.16/0.15)	3.6 (3.3/3.9)		0.16(0.13/0.18)
T (°C)	25.6 (27.0/22.2)			23.1 (28.8/17.5)		

<sup>a</sup> 2 filter and 4 PUF samples i.e., No. 9, 10, 19 and 22 in Fig. S3

<sup>b</sup> 1 filter and 1 PUF sample i.e., No. 1 and 2 in Fig. S3

<sup>c</sup> co-eluted with 3NFLT, assuming  $c_{3NFLT} = 0$