

Submitted on 28 Sep 2016

Anonymous Referee #2

Anonymous during peer-review: Yes No

Anonymous in acknowledgements of published article: Yes No

Recommendation to the Editor

1) Scientific Significance

Does the manuscript represent a substantial contribution to scientific progress within the scope of this journal (substantial new concepts, ideas, methods, or data)?

Excellent Good Fair Poor

2) Scientific Quality

Are the scientific approach and applied methods valid? Are the results discussed in an appropriate and balanced way (consideration of related work, including appropriate references)?

Excellent Good Fair Poor

3) Presentation Quality

Are the scientific results and conclusions presented in a clear, concise, and well structured way (number and quality of figures/tables, appropriate use of English language)?

Excellent Good Fair Poor

For final publication, the manuscript should be

accepted as is

accepted subject to technical corrections

accepted subject to minor revisions

reconsidered after major revisions

I would be willing to review the revised paper, if the Editor considers it necessary

I would NOT be willing to review the revised paper
rejected

Please note that this rating only refers to this version of the manuscript!

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

Referee Comment

General comment:

The authors have been greatly improving the manuscript according to my comments.

Before final publication I have a few minor issues I would recommend to take care of.

These minor comments are listed below.

Detailed scientific comments:

Abstract

Q1:Page 1, Line 13:

Comment: Which size distribution? Number? Please specify!

A1: We have changed 'size distribution' to 'number distribution' on Page1 Line13.

Q2:Page 2, Line 6-7:

Comment: It is not the visibility that leads to health problems. Please correct the sentence!

A2: We have clarified ‘Visibility degradation is mainly caused by the increase of particle number or mass concentration and can lead to a variety of health problems (such as cardiovascular disease, respiratory system diseases, etc.)’ to ‘Visibility degradation is mainly caused by the increase of particle number or mass concentration. The increase of particulate pollution can lead to a variety of health problems (such as cardiovascular disease, respiratory system diseases, etc.)’ on Page2 Line6-8.

Q3:Page 2, Line 12-14:

Comment: It is not the chemical composition itself, rather more the resulting refractive index that affect the optical properties. Please say that!

A3: We have clarified ‘The particle number size distribution, chemical composition and relative refractive index determined by the chemical composition are the important parameters that affect the optical properties of the particles’ to ‘The particle number size distribution, chemical composition and relative refractive index are the important parameters that affect the optical properties of the particles’ on Page2 Line12-14.

Q4:Page 2, Line 18:

... which has ...

A4: We have changed’... which have...’ to ‘... which has...’ on page2 Line18.

Q5:Page 4, Line 5:

Remove Q18!

A5: We have removed ‘Q18’ on Page4 Line6.

2 Experiment and Methods

Q6:Figure 1:

Comment: I would rather more describe this as “data coverage” in the figure description.

A6: We have changed ‘Fig. 1 Missing data from instruments during the observation period’ to ‘Fig. 1 Data coverage from instruments during the observation period’ on Page19 Line13.

Q7:Page 5, Line 9:

... were fired ...

A7: We have changed ‘were fires’ to ‘were fired’ on Page5 Line10.

Q8:Page 5, Line 11:

... analysis ...

A8: We have changed ‘analyzing’ to ‘analysis’ on Page5 Line12.

Q9:Page 5, Line 22:

... EC and OC concentrations ...

A9: We have changed ‘EC and OC concentration’ to ‘EC and OC concentrations’ on Page5 Line23.

Page 6, Line 15:

Q10: Remove Q23!

A10: We have removed 'Q23' on Page6 Line16.

Q11: Page 7, Line 11:

... EC determines the ...

A11: We have changed 'EC can describe the...' to 'EC determines the ...'

Q12: Page 8, Line 11-13:

Comment: Check the sentence "550nm is the ...". It does not make sense!

A12: We have changed '550nm is the...' to 'Wavelength of 550 nm is the...' on Page8 Line13.

3 Results and Discussion

Q13: Figure 4:

... in different size segments ...

A13: We have changed 'Fig. 4 Time series of κ in different sizes during the observation period' to 'Fig. 4 Time series of κ in different size segments during the observation period' on Page20 Line4.

Q14: Page 9, Line 26:

... lower compared to ...

A14: We have changed '...lower than when $\lambda = 940$ nm' to 'lower compared to $\lambda = 940$ nm' on Page9 Line27.

Q15: Figure 8:

Comment: Figure 9 is actually discussed before Figure 8. Please restructure!

A15: We have exchanged Figure 9 and Figure 8 according to review's comment on page.

Q16: Page 11, Line 14-15 and ongoing:

Comment: Check this sentence! You now talk about Figure 10! You mix up the Figure numbers in the following.

A16: We are so sorry that mix up the numbers. We have changed 'Fig. 9(a)' to 'Fig. 10(a)' on Page11 Line15 and ongoing.

Q17: Figure 10b:

Comment: I still do not understand Figure 10b!

A17: Fig. 10(b) represented the variation in Q with respect to RH. From Fig.10a, we find that the extinction coefficient of particles in the $0.01\text{-}0.2\mu\text{m}$ size range increased the fastest with the increased RH, followed by the extinction coefficients of particles in the $0.2\text{-}0.5\mu\text{m}$ and $1.0\text{-}2.0\mu\text{m}$ size ranges.

While the particles in the $0.5\text{-}1.0\mu\text{m}$ range had the largest κ , which means that the variability in diameter cannot explain the lack of obvious increase in the extinction coefficients in the $0.5\text{-}1.0\mu\text{m}$ size range.

According to the Mie theory, the impact of RH on particles was reflected in two aspects: the variability

in diameter and the efficiency factor (Q). Thus, we give the Fig. 10(b), and find that Q increased significantly in the 0.01-0.2 μ m, 0.2-0.5 μ m, and 1.0-2.0 μ m size ranges with the increase in RH and that Q declined slightly in the 0.5-1.0 μ m and 2.0-10.0 μ m size ranges at high RH values. Because λ =550 nm, the increase in the scale parameter in the 0.01-0.2 μ m, 0.2-0.5 μ m, and 1.0-2.0 μ m size ranges favors the increase in Q, whereas the increase of the scale parameter in the 0.5-1.0 μ m size range leads to a decrease in Q. So, we draw the conclusion that variation in the scale parameter leads to variation in Q, which is the main reason that growth multiples of the extinction coefficients vary at different RH levels.

4 Conclusions

Q18:Page 13, Line 11:

... of the extinction coefficients ...

A18: We have changed ‘...of the extinctions...’ to ‘of the extinction coefficients’ on Page13 Line11

Submitted on 01 Oct 2016

Anonymous Referee #1

Anonymous during peer-review: Yes No

Anonymous in acknowledgements of published article: Yes No

Recommendation to the Editor

1) Scientific Significance

Does the manuscript represent a substantial contribution to scientific progress within the scope of this journal (substantial new concepts, ideas, methods, or data)?

Excellent **Good** Fair Poor

2) Scientific Quality

Are the scientific approach and applied methods valid? Are the results discussed in an appropriate and balanced way (consideration of related work, including appropriate references)?

Excellent Good **Fair** Poor

3) Presentation Quality

Are the scientific results and conclusions presented in a clear, concise, and well structured way (number and quality of figures/tables, appropriate use of English language)?

Excellent Good Fair **Poor**

For final publication, the manuscript should be

accepted as is

accepted subject to technical corrections

accepted subject to minor revisions

reconsidered after major revisions

I would be willing to review the revised paper, if the Editor considers it necessary

I would NOT be willing to review the revised paper

rejected

Please note that this rating only refers to this version of the manuscript!

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

I think the manuscript has been improved from the previous one. The results the manuscript presented are interesting and of importance, which fill the data pool of extinction properties globally. However, I still have some comments regarding to the current version:

Q1:

1) The presentation or writing of the manuscript should be improved much more. There are too many long, wired, improper statements in the manuscript, for example: Page 3, line 4-6; Page 3, line 22-25; Page 4, line 5; Page 5, line 22-30; Page 7, line 14-16; Page 11, line 14-15, ect.... Please consider going through the whole manuscript and maybe asking someone who has more experience in English writing to check it.

A1: We amended the paper again, especially the sentences mentioned above.

Q2:

2) The author presented 11 figures, eight of which were about time series. Since the article is not focusing on time series analysis, I suggest the author replace some of them as scatter plot. For instance, the author compared PM2.5 with visibility and PM2.5 with the total particle number concentration. However, it is not obvious to see how well their negative or positive correlations in the time series of Fig. 2 or Fig. 3. Hence, scatter plot might be more useful. From the time series of Fig. 4 to Fig. 8, no critical findings or conclusions were driven by the author and the information presented in these time series figures are obviously not clear. I suggest the author consider moving them into supplementary material or giving more discussion.

A2: We also considered what reviewer said. While the time series figures in our study are basic data, and are used to be basis of the following calculations. For instance, Fig. 2 and Fig. 3 mainly give the observed results of some variates. Fig. 4 and Fig. 5 are the key parameters in our model, and their time series are the basis of the calculation of extinction coefficients. Fig.6 to Fig.9 are the calculated results that calculated by our model.

For the correlations that reviewer mentioned, the correlation among the visibility, PM2.5 and the total particle number concentration are not the main content. We therefore only give the correlation coefficient.

Q3:

3) I am not satisfied with the author's response to Question 6 from Referee 1 (see acp-2016-119-AC1-supplement-2). The internally mixed and externally mixed results are not shown in Fig. 1, which is not consistent with its figure caption and figure legend or where are the green and red curves?

A3: We are so sorry that referee is not satisfied with our reply to Question 6. The Fig.1 we give including the internally mixed and externally mixed results, and found them to be consistent with those of the core-shell model. Maybe the coincidence of lines caused the misunderstanding. Thus, we give them respectively. The values of the extinction coefficient calculated by the core-shell model are

between those of the internally mixed model and externally mixed model. The relationships between the model-derived extinction coefficients and measurement-derived extinction coefficients using the three mixed models are given below ($\lambda=550\text{nm}$). The result of the internally /externally mixed model is consistent with the current values. Considering the article length and structure, we discussed only the core-shell model in this paper.

Submitted on 01 Oct 2016

Anonymous Referee #1

Anonymous during peer-review: Yes No

Anonymous in acknowledgements of published article: Yes No

Recommendation to the Editor

1) Scientific Significance

Does the manuscript represent a substantial contribution to scientific progress within the scope of this journal (substantial new concepts, ideas, methods, or data)?

Excellent **Good** Fair Poor

2) Scientific Quality

Are the scientific approach and applied methods valid? Are the results discussed in an appropriate and balanced way (consideration of related work, including appropriate references)?

Excellent Good **Fair** Poor

3) Presentation Quality

Are the scientific results and conclusions presented in a clear, concise, and well structured way (number and quality of figures/tables, appropriate use of English language)?

Excellent Good Fair **Poor**

For final publication, the manuscript should be

accepted as is

accepted subject to technical corrections

accepted subject to minor revisions

reconsidered after major revisions

I would be willing to review the revised paper, if the Editor considers it necessary

I would NOT be willing to review the revised paper

rejected

Please note that this rating only refers to this version of the manuscript!

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

I think the manuscript has been improved from the previous one. The results the manuscript presented are interesting and of importance, which fill the data pool of extinction properties globally. However, I still have some comments regarding to the current version:

Q1:

1) The presentation or writing of the manuscript should be improved much more. There are too many long, wired, improper statements in the manuscript, for example: Page 3, line 4-6; Page 3, line 22-25;

Page 4, line 5; Page 5, line 22-30; Page 7, line 14-16; Page 11, line 14-15, ect.... Please consider going through the whole manuscript and maybe asking someone who has more experience in English writing to check it.

A1: We amended the paper again, especially the sentences mentioned above.

Q2:

2) The author presented 11 figures, eight of which were about time series. Since the article is not focusing on time series analysis, I suggest the author replace some of them as scatter plot. For instance, the author compared PM2.5 with visibility and PM2.5 with the total particle number concentration. However, it is not obvious to see how well their negative or positive correlations in the time series of Fig. 2 or Fig. 3. Hence, scatter plot might be more useful. From the time series of Fig. 4 to Fig. 8, no critical findings or conclusions were driven by the author and the information presented in these time series figures are obviously not clear. I suggest the author consider moving them into supplementary material or giving more discussion.

A2: We also considered what reviewer said. While the time series figures in our study are basic data, and are used to be basis of the following calculations. For instance, Fig. 2 and Fig. 3 mainly give the observed results of some variates. Fig. 4 and Fig. 5 are the key parameters in our model, and their time series are the basis of the calculation of extinction coefficients. Fig.6 to Fig.9 are the calculated results that calculated by our model.

For the correlations that reviewer mentioned, the correlation among the visibility, PM2.5 and the total particle number concentration are not the main content. We therefore only give the correlation coefficient.

Q3:

3) I am not satisfied with the author's response to Question 6 from Referee 1 (see acp-2016-119-AC1-supplement-2). The internally mixed and externally mixed results are not shown in Fig. 1, which is not consistent with its figure caption and figure legend or where are the green and red curves?

A3: We are so sorry that referee is not satisfied with our reply to Question 6. The Fig.1 we give including the internally mixed and externally mixed results, and found them to be consistent with those of the core-shell model. Maybe the coincidence of lines caused the misunderstanding. Thus, we give them respectively. The values of the extinction coefficient calculated by the core-shell model are between those of the internally mixed model and externally mixed model. The relationships between the model-derived extinction coefficients and measurement-derived extinction coefficients using the three mixed models are given below ($\lambda=550\text{nm}$). The result of the internally /externally mixed model is consistent with the current values. Considering the article length and structure, we discussed only the core-shell model in this paper.




Fig.1 Time series of measurement-derived extinction coefficients and model-derived extinction coefficients (core-shell/internally/externally model)




Fig.2 Correlation coefficient of model-derived extinction coefficients (core-shell/internally/externally model) and measurement-derived extinction coefficients

● Analysis of extinction properties as a function of relative humidity using a κ -EC-Mie model in Nanjing

Zefeng. Zhang¹, Yan. Shen¹, Yanwei. Li^{1,2}, Bin. Zhu¹, Xingna. Yu¹

¹Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Key

5 Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science & Technology, Nanjing, 210044, China

²Fenghua Meteorological Bureau of Zhejiang Province, Fenghua 315500, China

Correspondence to: Y. L Li (liyw@nuist.edu.cn)

10 **Abstract** The relationship between relative humidity (RH) and extinction properties is of widespread concern. In this study, a hygroscopic parameter (κ) and the volume fraction of elemental carbon (EC) were used to characterize the chemical characteristics of particles, and a core-shell model was built based on these characteristics. **The number distribution**, chemical compositions and RH were measured in Nanjing from 15/10/2013 to 13/11/2013. The model-derived extinction coefficients of particles were 15 fitted with the program of coated spheres form Bohren & Huffman (BHCOAT), and the values correlated well with the measurement-derived extinction coefficients ($r^2=0.81$), which suggested that the core-shell model produced reasonable results. The results show that more than 81% of the extinction coefficient in Nanjing was due to particles in the 0.2-1.0 μm size range. Under dry conditions, the higher mass fraction of particles in the 0.2-1.0 μm size range caused the higher extinction coefficient. An 20 increase in RH led to a significant increase in the extinction coefficient, although the increases differed among the different size segments. The corresponding functions are given in this study. For $\lambda=550$ nm, the extinction coefficient contributions of the 0.01-0.2 μm , 0.2-0.5 μm , and 1.0-2.0 μm size ranges increased significantly with the increase in RH, whereas the extinction coefficient contributions of the 0.5-1.0 μm and 2.0-10.0 μm size ranges decreased slightly.

25 1 Introduction

The degradation of visibility is likely the most readily perceived impact of aerosol pollution and has been used as a visual indicator of ambient air quality (Watson, 2002). Visibility throughout the world has generally decreased in recent decades, especially in Asia. In China, horizontal visibility has significantly decreased since 1980 (Che et al., 2007; Qian and Giorgi, 2000; Qian et al., 2007; Streets et al., 2008; Fu

et al., 2013; Yang et al., 2015; Liu et al., 2015). For example, in Guangzhou, one of the largest cities in the Pearl River Delta (PRD), low visibility occurs $150 \text{ days} \cdot \text{year}^{-1}$ (Deng et al., 2008). In the Beijing-Tianjin-Hebei region, the annual average number of continuous haze events has increased, accounting for more than half of the total haze days in a year (Zhang et al., 2015). In the Yangtze River Delta (YRD) region, visibility has decreased at the rate of $2.41 \text{ km} \cdot \text{decade}^{-1}$ (Gao et al., 2011). **Visibility degradation is mainly caused by the increase of particle number or mass concentration. The increase of particulate pollution can lead to a variety of health problems (such as cardiovascular disease, respiratory system diseases, etc.) and can further lead to an increase of traffic accidents, which has a serious impact on human health and activities (Tie et al., 2009; Wu et al., 2005; Chang et al., 2009).** As a result, visibility problems have received unprecedented attention in recent years.

Under dry conditions, the extinction of particles is the main factor affecting visibility (Covert et al., 1972; Deng et al., 2008; Watson, 2002). **The particle number size distribution, chemical composition and relative refractive index are the important parameters that affect the optical properties of the particles** (Day et al., 2000; Ma et al., 2012; Cheng et al., 2008a; Wen and Yeh, 2010). However, many of the aerosol components are hygroscopic and take up water as a function of the relative humidity (RH) (Clarke et al., 2004; Covert et al., 1972). When the RH is high, even at subsaturated conditions, the hygroscopic growth of the particles can lead to an increase in size and a decrease in the refractive index, **which has significant effects** on the extinction properties (Cheng et al., 2008b; Covert et al., 1972; Stock et al., 2011). Furthermore, the physicochemical properties of aerosols can lead to variable hygroscopic growth, and the extinction associated with different particles (with differences in size and chemistry) is significantly different under the same RH. Overall, visibility will decrease when the RH increases (Charlson, 1969; Covert et al., 1972; Stock et al., 2011; Day and Malm, 2001). Some studies show that extinction can increase by more than 100% when the RH exceeds 70%-80% (Mcmurry, 2000; Zhang and Mcmurry, 1992; Tang, 1996). Therefore, the study of the effect of RH on the extinction coefficient is very important.

Interest in the relationship between aerosol composition, RH and visibility dates back to at least to the studies of Wright (1940) on the atmospheric opacity over Valentia, Ireland (Wright, 1940). Currently, we can calculate the extinction coefficient accurately based on the Mie theory (Bohren and Huffman, 2008) as long as we have information of the overall aerosol population. However, atmospheric particles consist of a complicated mixture of various chemical compositions, and it is very difficult to obtain

complete data on the physicochemical properties of all particles. In field observations, a fitting formula is often used to assess the contribution of RH, though this method cannot reflect the differences in physicochemical properties of particles based on the observed aerosol, and fitting curves are different at different sites(Chen et al., 2010; Yu et al., 2016). Another method used an empirical formula which

5 based on chemical composition and different RH to calculate the extinction coefficient. However, this empirical formula may not be suitable for other locations. Therefore, we have established a model that

has few variables and for which each variable can be obtained based on conventional observations. The three-component model is an important hypothesis(Cheng et al., 2006). The first component is elemental carbon (EC), which is the light-absorbing component. The real and imaginary parts of EC are extremely

10 high, and a typical value is 1.8- 0.54*i* (Lee and Tien, 1981; Redemann et al., 2000). Water is the second component, which only scatters the incident radiation with the lowest refractive index of 1.33-0.0*i* (Levoni et al., 1997). Aside from EC and water, the rest of the aerosol components primarily only scatter light, and their refractive indices were very similar, with a real part of approximately 1.53 and an imaginary part of nearly zero. This component is called the non-light-absorbing component (Lin et al.,

15 Tang, 1996; Wex et al., 2002). Wex's study showed that the simplification is reasonable (2002).

She found that, under dry conditions, there was no statistically significant effect on the deviation between the measured and calculated scattering coefficients when varying the mass fractions of the nearly pure light-scattering compositions within their general concentration levels. Thus, as long as we know the volume fraction of EC, we can better describe the extinction properties of the particles.

20 Particles show hygroscopic growth as the RH increases. Petters and Kreidenweis (2007) proposed a simple hygroscopic parameter, κ , that can be used to calculate the hygroscopic growth factor (GF) at different levels of RH. κ can be considered a function of the volume fraction of the hygroscopic components which are non-light absorbing and the volume fraction of the non-hygroscopic components which are light-absorbing and can be assumed to be EC here. In this way hygroscopic and optical

25 properties can be understood to have a strong linkage. Supposing that non-light-absorbing material is uniformly mixed with water after hygroscopic growth, we can determine the changes in volume of both the real part and imaginary parts of the particles, then we can calculate the extinction coefficient of particles (Chen et al., 2012).Therefore, we can calculate the extinction properties of particles accurately at different levels of RH based on the Mie theory according to the three-component model, as long as we

30 obtain the volume fraction of EC of the observed aerosols, the hygroscopic parameter (κ) of the

observed aerosols and the hypothesized mixed mode of the observed aerosols.

In this study, the volume fraction of EC and the hygroscopic parameter (κ) were obtained using film sampling. Film sampling was used to analyze the chemical composition of particles that were most common and oldest, with abundant observational data. In this study, the volume fraction of EC was

5 obtained from the film sampling conducted using an Anderson instrument in Nanjing and κ was calculated according to the ZSR rule(Petters and Kreidenweis, 2007; Stokes and Robinson, 1966). The extinction coefficient calculated by the above method had a good relationship with the visibility, which verified the reasonableness of our method. Based on this result, we further explored the growth curve of the extinction coefficient in dependence of RH and the variety of extinction contributions provided by 10 different particle sizes ranges.

2 Experiment and methods

2.1 Measurement location and sampling

The sampling site was on the roof of a twelve-story building at the Nanjing University of Information Science and Technology in the Pukou District of Nanjing (32.207 N; 118.717 E), 40 m above ground

15 level. Visibility and meteorological parameters were obtained from the detection base near the sampling site at a distance of less than 1.5 km. The sampling period was from 15/10/2013 to 13/11/2013. The periods of instrumental observation are shown in Fig. 1, and the missing data were due to power failure. Moreover, because the system measurement error is high at high RH values, we excluded the data with an RH >90% and visibility <1 km.

20 Fig. 1 Data coverage from instruments during the observation period

2.2 Instruments and data analysis

2.2.1 Instruments

A wide-range particle spectrometer (WPS; MSP Corporation model 1000XP) is a recently introduced commercial instrument with the unique ability to measure the size distributions of aerosols with

25 diameters from 0.01 to 10 μ m (Liu, 2010). The WPS combines the principles of differential mobility analysis (DMA), condensation particle counting (CPC) and laser light scattering (LPS). DMA and CPC are used to measure particles' number concentration in the size range of 10-500nm, and LPS is used to

measure particles' number concentration in the range of 0.35-10 μ m. One complete scan of the entire size range with a 3s scanning period for each channel takes approximately 5 min, and a detailed description was provided previously(An et al., 2015; Kang et al., 2013).

Anderson is a nine-stage impact sampler that is produced by the Thermo Electron Corporation (USA). It was used to collect aerosol samples. The sampling flow rate is 28.3 L \cdot min $^{-1}$. The size distributions are provided in section as follows: \leq 0.43 μ m, 0.43-0.65 μ m, 0.65-1.1 μ m, 1.1-2.1 μ m, 2.1-3.3 μ m, 3.3-4.7 μ m, 4.7-5.8 μ m, 5.8-9.0 μ m and $>$ 9.0 μ m. We used cellulose filters for ionic species and quartz filters for EC

and OC(organic carbon). **Before use, quartz filters were fired for 5h at 800 °C to lower the blank levels for EC and OC.** All of these filters were kept in a refrigerator for cryopreservation. **Every sample was collected continuously for 23h and then kept in a refrigerator before analysis (Zou et al., 2014).**

Water-soluble ions were measured with a chromatograph (850 professional IC). Na $^+$, NH $_4^+$, Ca $^{2+}$, Mg $^{2+}$, K $^+$, F $^-$, Cl $^-$, NO $_2^-$, NO $_3^-$ and SO 4^{2-} were analyzed in this study. Chromatography includes the use of a column oven, a conductivity detector, an 858 auto-injector and a MagIC net chromatography workstation (Metrohm, Switzerland). The column oven consists of a Metrosep C4150/4.0 separation column and Metrosep A Supp 5150/4.0 separation column. The eluent was set at 3.2mmol \cdot L $^{-1}$ Na₂CO₃+1.0mmol \cdot L $^{-1}$ NaHCO₃ for anions and 1.7mmol \cdot L $^{-1}$ HNO₃+0.7mmol \cdot L $^{-1}$ pyridine carboxylic acid for cations. The column temperature was maintained at 30 °C. The flow-rate was 1.0mL \cdot min $^{-1}$, and the inject volume was 20 μ L. The detection limits for Na $^+$, NH $_4^+$, K $^+$, Mg $^{2+}$, Ca $^{2+}$, F $^-$, Cl $^-$, NO $_2^-$, NO $_3^-$ and SO 4^{2-} were 0.001, 0.005, 0.001, 0.001, 0.001, 0.01, 0.01, 0.01, 0.01, 0.01mg \cdot L $^{-1}$ respectively (An et al., 2015).

The EC and OC concentrations were determined with a thermal/optical carbon analyzer (Model 2001A, DRI). The samples were heated to 140, 280, 480 and 580 °C in pure He to determine OC1, OC2, OC3 and OC4, respectively. Then the samples were heated to 580, 740 and 840 °C in 2%O₂/98%He to determine EC1, EC2 and EC3, respectively. During the heating process, some volatilized organic compounds were converted to carbon dioxide (CO₂) through an oxidizer (heated manganese dioxide, MnO₂). CO₂ was reduced to methane (CH₄) through a methanator. Finally, the CH₄ equivalents were quantified with a flame ionization detector (FID). The charring effect can transform part of organic carbon into pyrolysis carbon under anaerobic heating. Hence, the correction for pyrolysis was made by continuously monitoring the filter through a 633nm He-Ne laser in order not to underestimate OC or include some pyrolyzed OC in the EC fraction.

By monitoring the change of reflected light in the heating process, the initial reflected light is an diacritical point of OC and EC (Miao et al., 2015; Zou et al., 2014).

PM_{2.5} was detected with a β -ray particulate continuous monitor (Thermo Fisher) with the working

principle of measuring the particles' mass concentration through the β -ray attenuation. Visibility data were collected with a CJY-1 visibility meter (CAMA Measurement & Control Equipments Co., Ltd). The visibility meter was used to measure the scattering coefficient of the particles and its light source wavelength was 940nm. The accuracy was $\pm 10\%$, and the data update rate was 1min. A detailed 5 description of these two instruments was provided previously (Yu et al., 2016).

2.2.2 Calculation of the hygroscopic parameter (κ)

κ can be calculated according to many methods (Liu et al., 2014; Miao et al., 2015; Petters and Kreidenweis, 2007). In this study, κ was calculated with the ZSR rule(Petters and Kreidenweis, 2007)according to the chemical composition of the particles. For an inorganic component, we 10 considered a system containing H^+ , NH_4^+ , HSO_4^- , SO_4^{2-} , and NO_3^{2-} . We used the ion pairing method from Gysel et al. (2007), and his method is more precise than the ADDEM model ($\Delta g/g < 2\%$) (Topping et al., 2005). For each species, the molecular weight, κ and density are described in detail in Table 1 (Gysel et 15 al., 2007; Kreidenweis et al., 2008; Petters and Kreidenweis, 2008; Topping et al., 2005). Moreover, we considered the effect of water-soluble organic components (WSOC) on hygroscopic growth and assumed $\kappa_{\text{org}}=0.1$ (Jimenez et al., 2009; King et al., 2010).

We obtained the mass of each pure species according to the pairing method. Supposing a dry particle's density of $1.7 \text{ g} \cdot \text{cm}^{-3}$ (Wehner et al., 2008), we calculated the volume of the dry particle. Aside from the WSOC and the four types of inorganic components in Table 1, we assumed that other components do not contribute to the hygroscopic properties of the aerosols. According to the ZSR rule, κ is given by Eq. (1):

$$20 \quad \kappa = \sum_{i=1}^N \kappa_i \frac{V_{i,\text{dry}}}{V_{\text{tol,dry}}} \quad (1)$$

where N is the number of pure materials, κ_i is the hygroscopic parameter of the i^{th} pure material, $V_{i,\text{dry}}$ is the volume of the i^{th} pure material in the dry condition, and $V_{\text{tol,dry}}$ is the total volume of the dry particle.

25 Table 1 Properties of each pure material component

2.2.3 Calculation of the hygroscopic growth factor (GF)

The hygroscopic growth behavior of particles can be described by the theory of Köhler (1936). The theory of Köhler considered the Kelvin effect and Raoult effect and established a relationship among the

saturation ratio S (at sub-saturation, S is equivalent to RH), diameter and solute properties. Introducing the hygroscopic parameter κ (Petters and Kreidenweis, 2007), the hygroscopic growth factor (GF) is determined as follows in Eq. (2):

$$GF = \left(1 + \frac{\kappa \cdot S}{\exp\left(\frac{4\sigma_{s/a} M_w}{RT\rho_w}\right) - S}\right)^{\frac{1}{3}}$$

5 (2)

where M_w is the molecular weight of water, R is the ideal gas constant, ρ_w is the density of water, T is the temperature with a value of 20 °C, and $\sigma_{s/a}$ is assumed to be the surface tension coefficient between water and air (when $T=20^{\circ}\text{C}$, $\sigma_{s/a}=0.0728 \text{ N m}^{-1}$).

2.3 Methods

10 2.3.1 Model-derived extinction coefficients

Under dry conditions, the volume fraction of EC determines the volume fraction of the light-absorbing component and κ can describe the hygroscopicity of the particle. We believe that these two parameters can be used to calculate the extinction coefficient of a single particle accurately at different RH levels.

The physicochemical properties can be different for the same size of particles. It is unrealistic to

15 describe the physicochemical properties of the aerosols individually. Therefore, the internally mixed model, externally mixed model, and core-shell model are often used to describe the chemical composition of aerosol particles in practical studies (Lesins et al., 2002; Cheng et al., 2006; Hao et al., 2010). The calculated results of the extinction coefficient by the core-shell model are usually between those of the internally mixed model and externally mixed model(Hao et al., 2010). Therefore, the 20 core-shell model was used in this study.

In this study, the core-shell model operates under the following assumptions: 1) particles of the same size have the same physicochemical properties, and particles are spherical; 2) under dry conditions, particles are composed of a light-absorbing component (EC, $1.8-0.54i$) and a non-light-absorbing component ($1.53-0i$), and the EC is a spherical "core" that is always at the center of the particle; 3) GF is a function of

25 κ and the hygroscopic uptake of EC is minor, and the non-light-absorbing material is uniformly mixed with water after hygroscopic exposure. Considering that the methods of film sampling and WPS differ significantly in time resolution, we made the following assumptions: 1) the chemical compositions of particles were unchanged for a given diameter segment of Anderson; 2) the chemical composition of

particles remained unchanged over the course of a day.

According to the hypothesis of the core-shell model, we can calculate the model-derived extinction coefficients of the particles using Eq. (3). N was the number of size segments of the WPS (0.01-10 μm), with a value was 67. In this study, particles in the range of 0.5-10 μm were measured by LPS. The term n

5 represents the number concentration of size segment N_i (i from 1 to 67), and r_i is the median radius corresponding to N_i . Q_{ext} is an efficiency factor calculated with the BHCOAT program, Q_{ext} is defined as the extinction cross section of particle divided by the geometric cross section of particle. The input/output parameters of Q_{ext} and the formulas are listed in Table 2. In Table 2, X is a scale parameter.

10 D_0 is the diameter of a single particle under dry conditions. λ is the incident light wavelength($\lambda=550$ nm,

15 940 nm). The wavelength of light source of the visibility meter was 940nm, and the calculated value of $\lambda=940$ nm was used to contrast with the observed value of the visibility meter. **Wavelength of 550 nm is the most sensitive wavelength for the human eye, and its calculated value was consistent with the value that the most sensitive for human eye.** GF is the hygroscopic growth factor, which was calculated using Eq. (2). If $\text{RH}=0$, then $\text{GF}=1$. The complex refractive index was calculated with the volume weighting

15 method after the hygroscopic growth of the particle (Lesins et al., 2002).

$$b_{\text{ext, model-derived}} = \sum_{i=1}^N Q_{\text{ext}} \times \pi(r \times GF)_i^2 \times n(r_i) \quad (3)$$

Table 2 Input/output parameters of the efficiency factor (Q)

2.3.2 Measurement-derived extinction coefficients

The meteorological optical range is determined as (Zhang, 2007):

$$20 \quad \text{MOR} = \frac{1}{\sigma} \ln \frac{|c|}{\epsilon} = \frac{1}{\sigma} \ln \frac{1}{0.05} = \frac{3.0}{\sigma} \quad (4)$$

where σ is the extinction coefficient of the particles, ϵ is the visual threshold with a value of 0.05(MOR is equal to the visibility when $\epsilon=0.05$), and c is the target characteristic coefficient. When the target is black, $c=1$.

Hence, the measured extinction coefficient can be calculated from the visibility as:

$$25 \quad b_{\text{ext, measurement-derived}} = \frac{1}{\text{visibility}} \ln \frac{1}{0.05} = \frac{3.0}{\text{visibility}} \quad (5)$$

3 Results and discussion

3.1 Aerosol properties and visibility during the measurement period

Time series of RH, visibility, measurement extinction coefficient, and $PM_{2.5}$ during the observation period is shown in Fig. 2. The measurement extinction coefficient was calculated as 3.0/visibility

5 (Seinfeld and Pandis, 2012). The picture shows that the visibility has a strong negative correlation with $PM_{2.5}$ and RH ($r=-0.7$ and -0.62 , respectively). A time series of number size distribution for dry particles

is given by Fig. 3, We find that the periods with a high number concentration had a good consistency

with the periods of a high $PM_{2.5}$ mass concentration ($r=0.7$) . Fig. 4 shows the time series of κ for

different particle size segments. κ was calculated according to the ZSR rule, which is described in detail

10 in Section 2.2.2. Fig. 5 shows the time series of the volume fraction of EC in different size segments, and

the volume fraction of EC was calculated using data from the Anderson instrument. Fig. 4 and Fig. 5

show that κ and the volume fraction of EC changed over time, but the variation between size segments is

higher compared to the variation over time within one size segment especially interactive for κ . The

reason for this difference may be that the particle size was closely related to the sources.

15 Fig. 2 Time series of RH, visibility, extinction coefficient, and $PM_{2.5}$ during the observation period

Fig. 3 Time series of particle number size distribution (dry particles) during the observation period

Fig. 4 Time series of κ in different size segments during the observation period

Fig. 5 Time series of the volume fraction of EC for different size segments during the observation period

20 3.2 Comparative analysis of the model-derived and measurement-derived extinction coefficients by the core-shell model

Figure 6 shows the relative values of the model and measurement values of the extinction coefficient from the core-shell model. When $\lambda=940$ nm, the calculated and measured values of extinction coefficient were in good agreement ($r^2=0.81$), which indicated that using the hygroscopic parameter (κ) and volume

25 fraction of EC to characterize the chemical characteristics of particles was reasonable. When $\lambda=550$ nm,

the correlation coefficient of the calculated and measured values ($r^2=0.714$) was slightly lower compared

to $\lambda= 940$ nm, mainly due to the differences in the wavelength of the light source. Comparing the

extinction values of 550 nm and 940 nm, we found that the model-derived extinction coefficient at 550

nm was higher, mainly due to the differences in scale parameters, which led to a Q that was larger when

$\lambda=550$ nm. Because 550 nm is the most sensitive wavelength for the human eye, the following section focuses on the measurements and calculations at $\lambda=550$ nm for discussion.

Fig. 6 Relationships among the calculated and measured values based on the core-shell model ($\lambda=550$

nm, 940 nm)

5 **3.3 Contributing fraction of the extinction coefficient for different size segments under dry conditions**

In the core-shell model, we defined GF=1 and then used Eq. (3) to calculate the extinction coefficients of particles under dry conditions. We can calculate the extinction coefficients of particles in different size segments with different median radii (r). In this study, particle size was divided into five segments:

10 0.01-0.2 μ m, 0.2-0.5 μ m, 0.5-1.0 μ m, 1.0-2.0 μ m, and 2.0-10.0 μ m. Fig. 7(a) shows the time series of

different size segments to the dry aerosol extinction coefficient, and Fig. 7(b) shows the relative contributing fraction of different size segments to the dry aerosol extinction coefficient. Fig. 7(b) shows

that the relative contributing fraction of different size segments to the dry aerosol extinction coefficient were significantly different. On average, the 0.2-0.5 μ m and 0.5-1.0 μ m ranges together contributed more

15 than 81% of the extinction coefficients, much higher than their total PM₁₀ mass fraction (45%). This result suggests that, an increase in the proportion of particles in the 0.2-1.0 μ m size range in PM₁₀ will

result in an even greater increase in the extinction capacity relative to the unit mass of the particles. This result is consistent with the results of Kang et al. (2013). **To verify this point, we present Fig.8**, which

reflects the extinction capacity relative to the unit mass in different size segments under dry/wet conditions. The y-axis is the ratio of the extinction coefficient to the mass concentration for different size segments. From the picture, we can find that extinction capacity relative to the unit mass in the

20 0.2-2 μ m range was much stronger than that of the other segments. This result explains why the particles in the 0.2-2 μ m range are the most important for the reduction of the visibility, especially

those in the 0.5-1 μ m range.

25 Fig. 7 Time series (a) and the relative contributing fraction(b) of different size segments to the dry aerosol extinction coefficient

Fig. 8 Extinction capacity relative to unit mass in different size segments under dry/wet condition

Fig. 9 Time series (a) and the relative contributing fraction(b) of different size segments to the wet aerosol extinction coefficient

3.4 Effects of relative humidity on the extinction coefficient

For ambient RH, we can calculate the extinction coefficients of particles in different size segments using Eq. (3). Fig. 9(a) shows the time series of different size segments to the wet aerosol extinction coefficient, and Fig. 9(b) shows the relative contributing fraction of different size segments to the wet

5 aerosol extinction coefficient. Comparing Fig. 7 and Fig. 9, we found that the extinction coefficients of different size segments to the wet condition were larger than for particles under dry conditions. Simultaneously, the relative contributing fraction of different size segments to the aerosol extinction coefficient underwent significant changes. Generally speaking, when particles were in the 0.01-0.2 μm , 0.2-0.5 μm and 1.0-2.0 μm size ranges, the relative contribution fraction of the extinction coefficients all increased, especially for fine particles (Table 3). When particles were in the 0.5-1.0 μm and 2.0-10.0 μm size ranges, the relative contribution fraction of the extinction coefficients decreased.

10 Table 3 Contribution fraction of the model-derived extinction coefficients at dry/wet condition and mass fraction in PM_{10} at dry condition

The growth multiples of the extinction coefficients in different size segment (as shown in fig.10a) was

15 calculated through wet aerosol extinction coefficients in size segment (as shown in fig. 9a) divided by the dry aerosol extinction coefficients in size segment (as shown in fig. 7a). The y-axis represents growth multiples of the extinction coefficients compared to dry conditions. The x-axis represents the variability of RH. There are five fitting curves in Fig. 10(a), representing different size segments, and the correlation coefficient (r^2) of each fitting curve was larger than 0.9. This result suggests that, on different

20 days, the changes in the enhancement of extinction with the RH in the same size segment were consistent. In addition, the extinction coefficient of particles in the 0.01-0.2 μm size range increased the fastest with the increased RH, followed by the extinction coefficients of particles in the 0.2-0.5 μm and 1.0-2.0 μm size ranges. The extinction coefficients of particles in the 0.5-1.0 μm and 2.0-10.0 μm size ranges did not obviously increase with the increased RH.

25 The impact of RH on particles was reflected in two aspects: the variability in diameter and the efficiency factor (Q). The growth of particles was determined by the hygroscopic parameter (κ). As κ increased, GF also increased. Fig. 4 shows the time series of κ for different particle sizes during the observations. The particles in the 0.5-1.0 μm range had the largest κ , which means that the variability in diameter cannot explain the lack of obvious increase in the extinction coefficients in the 0.5-1.0 μm size range. To obtain

Q following the influence of RH, we performed the following calculation. Firstly, we assumed that the RH had no effect on Q, which means that Q was equivalent to the value under dry conditions. Secondly, we calculated the extinction coefficient of particles in different size segments using Eq. (3) (indicated by the letter *b*). Lastly, [Fig. 9\(a\)](#) was divided by *b* to produce Fig. 10(b), which represented the variation in Q with respect to RH. Fig. 10(b) shows that Q increased significantly in the 0.01-0.2 μm , 0.2-0.5 μm , and 1.0-2.0 μm size ranges with the increase in RH and that Q declined slightly in the 0.5-1.0 μm and 2.0-10.0 μm size ranges at high RH values. The fitting curve and the calculated values are significantly different. For particles in the 0.01-0.2 μm , 0.2-0.5 μm , 0.5-1.0 μm and 1.0-2.0 μm size ranges, the correlation coefficients are all high. Because $\lambda=550$ nm, the increase in the scale parameter in the 0.01-0.2 μm , 0.2-0.5 μm , and 1.0-2.0 μm size ranges favors the increase in Q, whereas the increase of the scale parameter in the 0.5-1.0 μm size range leads to a decrease in Q. For particles in the 2.0-10.0 μm range, the correlation coefficient is very low. This finding suggested that the effect of the variation in the scale parameter on Q was significantly different on different days. In summary, variation in the scale parameter leads to variation in Q, which is the main reason that growth multiples of the extinction coefficients vary at different RH levels.

Because the average particle size distribution and chemical composition in each size segment are known, we can calculate the average contribution fraction of the extinction coefficients in each size segment with the increase in RH. The calculation results are shown in Fig. 11, which illustrates that the extinction coefficient was primarily related to particles in the 0.2-0.5 μm and 0.5-1.0 μm size ranges. Generally speaking, an increase in RH will lead to an increase in the extinction coefficient, but the rate of increase in the extinction coefficient was significantly different in each size segment. With an increase in RH, the fractions of the extinction coefficients contributed by the 0.01-0.2 μm , 0.2-0.5 μm , and 1.0-2.0 μm size ranges increased considerably, whereas the fractions of the extinction coefficient contributed by the 0.5-1.0 μm and 2.0-10.0 μm size ranges decreased slightly.

Fig. 10 Growth multiples of the extinction coefficients (a) and the change in the efficiency factor (b) for different size segments at ambient relative humidity

Fig. 11 Relationship between the contribution fraction of the extinction coefficient in different size segments and relative humidity (RH)

4 Conclusions

In this study, a hygroscopic parameter (κ) and the volume fraction of elemental carbon (EC) were used to characterize the chemical characteristics of particles and a core-shell model was built based on these characteristics. In the core-shell model, the real part and the imaginary part of the refractive index, the scale parameters were both functions of RH. The extinction coefficients of particles fitted with the BHCOAT program correlated well with the measured values ($r^2=0.81$) that were derived from the visibility, which suggested that using κ and the volume fraction of EC to characterize the chemical characteristics of particles was reasonable.

In the core-shell model, when $\lambda=550$ nm, the contribution fractions of the extinction coefficient of different size segments were significantly different. Under the dry condition, more than 81% of the extinction coefficients in Nanjing were contributed by particles in the 0.2-1.0 μm size range, a much higher percentage than their PM_{10} mass fraction (45%). This finding suggested that, for PM_{10} , an increase in the mass proportion of particles in 0.2-1.0 μm size range results in an even greater increase in the extinction capacity.

With the increase in RH, the extinction capacity of particles will grow significantly. In this study, the formula for the increase in extinction coefficients in different size segments is given. At given RH, the growth rate of extinction coefficients differs significantly among different size segments. The growth rates are related to κ , but the variation in the scale parameter leads to variations in Q , which is the main reason that the growth multiples of the extinction coefficient differ at different RH values. With the increase in RH, the extinction coefficient contribution fractions increase for particles in the 0.01-0.2 μm , 0.2-0.5 μm and 1.0-2.0 μm size ranges but decrease for particles in the 0.5-1.0 μm and 2.0-10.0 μm size ranges.

Acknowledgement:

This work was supported jointly by the National Natural Science Foundation of China (Grant No. 25 41275152 and 41005071), and the National Key Research and Development Program of China (2016YFA0602003).

References

An, J., Wang, H., Shen, L., Zhu, B., Zou, J., Gao, J., and Kang, H.: Characteristics of new particle

formation events in Nanjing, China: Effect of water-soluble ions, *Atmos. Environ.*, 108, 32-40, doi: 10.1016/j.atmosenv.2015.01.038, 2015.

Bohren, C. F., and Huffman, D. R.: *Absorption and Scattering of Light by Small Particles*, John Wiley & Sons, New York, 2008.

5 Chang, D., Song, Y., and Liu, B.: Visibility trends in six megacities in China 1973–2007, *Atmos. Res.*, 94, 161-167, 2009.

Charlson, R. J.: Atmospheric visibility related to aerosol mass concentration: review, *Environ. Sci. Technol.*, 3, 913-918, 1969.

Che, H., Zhang, X., Li, Y., Zhou, Z., and Qu, J. J.: Horizontal visibility trends in China 1981–2005, *Geophys. Res. Lett.*, 34, L24706, doi: 10.1029/2007GL031450, 2007.

Chen, J., Zhao, C. S., Ma, N., Liu, P. F., Göbel, T., Hallbauer, E., Deng, Z. Z., Ran, L., Xu, W. Y., and Liang, Z.: A parameterization of low visibilities for hazy days in the North China Plain, *Atmos. Chem. Phys.*, 12, 4935-4950, 2012.

Chen, Y., Zhao, D., Chai, F., Liang, G., Xue, Z., Wang, B., Liang, Y., Chen, Y., and Zhang, M.: 15 Correlation between the atmospheric visibility and aerosol fine particles concentrations in Guangzhou and Beijing, *China Environ. Sci.*, 7, 967-971, 2010.

Cheng, Y. F., Eichler, H., Wiedensohler, A., Heintzenberg, J., Zhang, Y. H., Hu, M., Herrmann, H., Zeng, L. M., Liu, S., and Gnauk, T.: Mixing state of elemental carbon and non - light - absorbing aerosol components derived from in situ particle optical properties at Xinken in

20 Pearl River Delta of China, *J. Geophys. Res. Atmos.* (1984–2012), 111, 2006.

Cheng, Y. F., Heintzenberg, J., Wehner, B., Wu, Z. J., Su, H., Hu, M., and Mao, J. T.: Traffic restrictions in Beijing during the Sino-African Summit 2006: aerosol size distribution and visibility compared to long-term in situ observations, *Atmos. Chem. Phys.*, 8, 7583-7594, 2008a.

Cheng, Y. F., Wiedensohler, A., Eichler, H., Su, H., Gnauk, T., Brüggemann, E., Herrmann, H., 25 Heintzenberg, J., Slanina, J., and Tuch, T.: Aerosol optical properties and related chemical apportionment at Xinken in Pearl River Delta of China, *Atmos. Environ.*, 42, 6351-6372, 2008b.

Clarke, A. D., Shinozuka, Y., Kapustin, V. N., Howell, S., Huebert, B., Doherty, S., Anderson, T., Covert, D., Anderson, J., and Hua, X.: Size distributions and mixtures of dust and black 30 carbon aerosol in Asian outflow: Physiochemistry and optical properties, *J. Geophys. Res.*

Atmos., 109, 839-856, 2004.

Covert, D. S., Charlson, R. J., and Ahlquist, N. C.: A study of the relationship of chemical composition and humidity to light scattering by aerosols, *J. Appl. Meteorol.*, 11, 968-976, 1972.

Day, D. E., Malm, W. C., and Kreidenweis, S. M.: Aerosol light scattering measurements as a function 5 of relative humidity, *J. Air Waste Manage. Assoc.*, 50, 710-716, 2000.

Day, D. E., and Malm, W. C.: Aerosol light scattering measurements as a function of relative humidity: a comparison between measurements made at three different sites, *Atmos. Environ.*, 35, 5169-5176, 2001.

Deng, X., Tie, X., Wu, D., Zhou, X., Bi, X., Tan, H., Li, F., and Jiang, C.: Long-term trend of visibility 10 and its characterizations in the Pearl River Delta (PRD) region, China, *Atmos. Environ.*, 42, 1424-1435, doi: 10.1016/j.atmosenv.2007.11.025, 2008.

Fu, C., Wu, J., Gao, Y., Zhao, D., and Han, Z.: Consecutive extreme visibility events in China during 1960–2009, *Atmos. Environ.*, 68, 1-7, doi: 10.1016/j.atmosenv.2012.11.035, 2013.

Gao, L., Jia, G., Zhang, R., Che, H., Fu, C., Wang, T., Zhang, M., Jiang, H., and Yan, P.: Visual range 15 trends in the Yangtze River Delta Region of China, 1981–2005, *J. Air Waste Manage. Assoc.*, 61, 843-849, doi: 10.3155/1047-3289.61.8.843, 2011.

Gysel, M., Crosier, J., Topping, D. O., Whitehead, J. D., Bower, K. N., Cubison, M. J., Williams, P. I., Flynn, M. J., McFiggans, G. B., and Coe, H.: Closure study between chemical composition and hygroscopic growth of aerosol particles during TORCH2, *Atmos. Chem. Phys.*, 7, 6131-6144, doi: 10.5194/acp-7-6131-2007, 2007.

Hao, L., Yang, W., Wu, T., Shi, G., Zhao, J., and Tan, Y.: Analysis of radiative properties of BC-sulfate 20 aerosols, *Plateau Meteorol.*, 1238-1245, 2010.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, J. E., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J.

R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525-1529, doi: 10.1126/science.1180353, 2009.

5 Kang, H., Zhu, B., Su, J., Wang, H., Zhang, Q., and Wang, F.: Analysis of a long-lasting haze episode in Nanjing, China, *Atmos. Res.*, 120, 78-87, 2013.

King, S. M., Rosenoern, T., Shilling, J. E., Chen, Q., Wang, Z., Biskos, G., McKinney, K. A., Pöschl, U., and Martin, S. T.: Cloud droplet activation of mixed organic-sulfate particles produced by the photooxidation of isoprene, *Atmos. Chem. Phys.*, 10, 3953-3964, 2010.

10 Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Trans. Faraday Soc.*, 32, 1152-1161, 1936.

Kreidenweis, S. M., Petters, M. D., and DeMott, P. J.: Single-parameter estimates of aerosol water content, *Environ. Res. Lett.*, 3, 035002, 2008.

Lee, S. C., and Tien, C. L.: Optical constants of soot in hydrocarbon flames, *Symposium (International) on Combustion*, 18, 1159-1166, 1981.

15 Lesins, G., Chylek, P., and Lohmann, U.: A study of internal and external mixing scenarios and its effect on aerosol optical properties and direct radiative forcing, *J. Geophys. Res. Atmos.* (1984–2012), 107, AAC 5-1-AAC 5-12, 2002.

Levoni, C., Cervino, M., Guzzi, R., and Torricella, F.: Atmospheric aerosol optical properties: a database of radiative characteristics for different components and classes, *Appl. Optics*, 36, 20 8031-8041, 1997.

Lin, Z. J., Tao, J., Chai, F. H., Fan, S. J., Yue, J. H., Zhu, L. H., Ho, K. F., and Zhang, R. J.: Impact of relative humidity and particles number size distribution on aerosol light extinction in the urban area of Guangzhou, *Atmos. Chem. Phys.*, 13, 1115-1128, 2013.

25 Liu, B. Y. H.: A wide-range particle spectrometer for aerosol measurement from 0.010 μm to 10 μm , *Aerosol. Air Qual. Res.*, 10, 125-139, doi: 10.4209/aaqr.2009.10.0062, 2010.

Liu, H. J., Zhao, C. S., Nekat, B., Ma, N., Wiedensohler, A., van Pinxteren, D., Spindler, G., Müller, K., and Herrmann, H.: Aerosol hygroscopicity derived from size-segregated chemical composition and its parameterization in the North China Plain, *Atmos. Chem. Phys.*, 14, 2525-2539, 2014.

30 Liu, X., Hui, Y., Yin, Z. Y., Wang, Z., Xie, X., and Fang, J.: Deteriorating haze situation and the severe

haze episode during December 18–25 of 2013 in Xi'an, China, the worst event on record, Theoretical & Applied Climatology, 1-15, 2015.

Ma, N., Zhao, C. S., Müller, T., Cheng, Y. F., Liu, P. F., Deng, Z. Z., Xu, W. Y., Ran, L., Nekat, B., and Pinxteren, D. v.: A new method to determine the mixing state of light absorbing carbonaceous using the measured aerosol optical properties and number size distributions, *Atmos. Chem. Phys.*, 12, 2381-2397, 2012.

McMurry, P. H.: A review of atmospheric aerosol measurements, *Atmos. Environ.*, 34, 1959-1999, 2000.

Miao, Q., Zhang, Z., Li , Y., Duan, Q., Qin, X., and Xu , B.: Size distributions of carbonaceous aerosols and their potential sources at Mt.Huang during Summer., *Environ. Sci. Chinese*, 35, 1938-1946, 2015.

Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, 2007.

Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity – Part 2: Including solubility, *Atmos. Chem. Phys.*, 8, 6273-6279, doi: 10.5194/acp-8-6273-2008, 2008.

Qian, Y., and Giorgi, F.: Regional climatic effects of anthropogenic aerosols? The case of southwestern China, *Geophys. Res. Lett.*, 27, 3521-3524, doi: 10.1029/2000GL011942, 2000.

Qian, Y., Wang, W., Leung, L. R., and Kaiser, D. P.: Variability of solar radiation under cloud-free skies in China: The role of aerosols, *Geophys. Res. Lett.*, 34, L12804, doi: 10.1029/2006GL028800, 2007.

Redemann, J., Turco, R. P., Liou, K. N., Russell, P. B., Bergstrom, R. W., Schmid, B., Livingston, J. M., Hobbs, P. V., Hartley, W. S., and Ismail, S.: Retrieving the vertical structure of the effective aerosol complex index of refraction from a combination of aerosol in situ and remote sensing measurements during TARFOX, *J. Geophys. Res. Atmos.* (1984–2012), 105, 9949-9970, 2000.

Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons, New York, 2012.

Stock, M., Cheng, Y. F., Birmili, W., Massling, A., Wehner, B., Müller, T., Leinert, S., Kalivitis, N., Mihalopoulos, N., and Wiedensohler, A.: Hygroscopic properties of atmospheric aerosol

particles over the Eastern Mediterranean: implications for regional direct radiative forcing under clean and polluted conditions, *Atmos. Chem. Phys.*, 11, 4251-4271, 2011.

Stokes, R. H., and Robinson, R. A.: Interactions in Aqueous Nonelectrolyte Solutions. I. Solute, *Journal of Physical Chemistry*, 70, 1966.

5 Streets, D. G., Yu, C., Wu, Y., Chin, M., Zhao, Z., Hayasaka, T., and Shi, G.: Aerosol trends over China, 1980 – 2000 ☆, *Atmos. Res.*, 88, 174–182, 2008.

Tang, I. N.: Chemical and size effects of hygroscopic aerosols on light scattering coefficients, *J. Geophys. Res.*, 101, 19,245-219,250, 1996.

Tie, X., Wu, D., and Brasseur, G.: Lung cancer mortality and exposure to atmospheric aerosol particles 10 in Guangzhou, China, *Atmos. Environ.*, 43, 2375-2377, 2009.

Topping, D. O., McFiggans, G. B., and Coe, H.: A curved multi-component aerosol hygroscopicity model framework: Part 1 – Inorganic compounds, *Atmos. Chem. Phys.*, 5, 1205-1222, doi: 10.5194/acp-5-1205-2005, 2005.

Watson, J. G.: Visibility: Science and regulation, *J. Air Waste Manage. Assoc.*, 52, 628-713, doi: 15 10.1080/10473289.2002.10470813, 2002.

Wehner, B., Birmili, W., Ditas, F., Wu, Z., Hu, M., Liu, X., Mao, J., Sugimoto, N., and Wiedensohler, A.: Relationships between submicrometer particulate air pollution and air mass history in Beijing, China, 2004–2006, *Atmos. Chem. Phys.*, 8, 6155-6168, 2008.

Wen, C.-C., and Yeh, H.-H.: Comparative influences of airborne pollutants and meteorological 20 parameters on atmospheric visibility and turbidity, *Atmos. Res.*, 96, 496-509, 2010.

Wex, H., Neusüß, C., Wendisch, M., Stratmann, F., Koziar, C., Keil, A., Wiedensohler, A., and Ebert, M.: Particle scattering, backscattering, and absorption coefficients: An in situ closure and sensitivity study, *J. Geophys. Res. Atmos.* (1984–2012), 107, LAC 4-1-LAC 4-18, 2002.

Wright, H. L.: Atmospheric opacity at valentia, *Q. J. R. Meteorol. Soc.*, 66, 209–213, 1940.

25 Wu, D., Tie, X., Li, C., Ying, Z., Lau, A. K.-H., Huang, J., Deng, X., and Bi, X.: An extremely low visibility event over the Guangzhou region: A case study, *Atmos. Environ.*, 39, 6568-6577, 2005.

Yang, Y. Q., Wang, J. Z., Gong, S. L., Zhang, X. Y., Wang, H., Wang, Y. Q., Wang, J., Li, D., and Guo, J. P.: PLAM - a meteorological pollution index for air quality and its applications in fog-haze 30 forecasts in North China, *Atmospheric Chemistry & Physics*, 16, 9077-9106, 2015.


Yu, X., Ma, J., An, J., Yuan, L., Zhu, B., Liu, D., Wang, J., Yang, Y., and Cui, H.: Impacts of meteorological condition and aerosol chemical compositions on visibility impairment in Nanjing, China, *Journal of Cleaner Production*, 131, 112-120, <http://dx.doi.org/10.1016/j.jclepro.2016.05.067>, 2016.

5 Zhang, W.: Principle and method of atmospheric detection, Meteorological Press, 2007.

Zhang, X., and McMurry, P. H.: Evaporative losses of fine particulate nitrates during sampling, *Atmos. Environ.*, 26, 3305–3312, 1992.

Zhang, Y., Zhang, P., Wang, J., Qu, E., Liu, Q., and Li, G.: Climatic characteristics of persistent haze events over Jingjiji during 1981-2013, *Meteorol. Mon.*, 311-318, 2015.

10 Zou, J., An, J., Wang, H., Shao, P., Duan , Q., Xue , G., and Pang , B.: Distribution characteristics of pollution gases and water soluble ion in aerosol during the Asian youth games of Nanjing, China, *Environ. Sci.*, 35, 4044-4051, 2014.



15 Fig. 1 Data coverage from instruments during the observation period




Fig. 2 Time series of RH, visibility, extinction coefficient, and PM_{2.5} during the observation period




Fig. 3 Time series of number size distribution (dry particles) during the observation period




Fig. 4 Time series of κ in different size segments during the observation period




Fig. 5 Time series of the volume fraction of EC for different size segments during the observation period




Fig. 6 Relationships among the calculated and measured values based on the core-shell model ($\lambda=550$ nm, 940 nm)





Fig. 7 Time series (a) and the relative contributing fraction(b) of different size segments to the dry aerosol extinction coefficient



5 Fig. 8 Extinction capacity relative to unit mass in different size segments under dry/wet condition





Fig. 9 Time series (a) and the relative contributing fraction(b) of different size segments to the wet aerosol extinction coefficient



10 Fig. 10 Growth multiples of the extinction coefficients (a) and the change of efficiency factor (b) in different size segments at ambient relative humidity




Fig. 11 Relationship between the contribution fraction of extinction coefficients in different size segments and relative humidity (RH)

5 Table 1 Properties of each pure material

	Molecular weight	Density (g cm ⁻³)	κ
NH_4NO_3	80.04	1.72	0.68
H_2SO_4	98.08	1.83	1.13
NH_4HSO_4	115.11	1.78	0.56
$(\text{NH}_4)_2\text{SO}_4$	132.14	1.77	0.53
WSOC		1.40	0.10

Table 2 Input/output parameters of efficiency factor (Q)

input parameters	output parameters
$X_{cor} = \frac{\pi \cdot \sqrt[3]{\frac{V_{EC}}{V_{tol}}} \times D_0}{\lambda}$	Q_{ext}
$X_{man} = \frac{\pi \cdot D_0 \cdot GF}{\lambda}$	ω_0
$m_{cor} = (1.8, 0.54)$	
$m_{man} = \left(\frac{1.53 + 1.33(GF-1)^3}{(GF-1)^3 + 1}, 0 \right)$	

Table 3 Contribution fraction of the model-derived extinction coefficients at dry/wet condition and mass fraction in

10 PM₁₀ at dry condition

	0.01-0.2 μm	0.2-0.5 μm	0.5-1.0 μm	1.0-2.0 μm	2.0-10.0 μm
Contribution fraction of the model-derived extinction coefficients at dry condition ^a	3.4%	28.3%	52.6%	7.6%	8.0%
Contribution fraction of the model-derived extinction coefficients at wet condition ^a	4.6%	33.3%	47.6%	7.8%	6.9%
Mass fraction in PM ₁₀ at dry condition ^a	8.5%	17.4%	27.6%	13.2%	33.3%