

Referee comments are in plain text.

Our responses are indented, in italicized text.

Anonymous Referee #1

In this paper, Chen et al. use kinetic analysis of OH p-HBA formation in illuminated snow samples to infer OH photochemical formation rates and steady-state concentrations in (assumed) pristine snow taken from Arctic regions. This work is important and timely: in a rapidly changing and increasingly human-impacted Arctic environment, understanding the contributions to local atmospheric reactivity and oxidation state due to chemistry associated with "pristine" snow and firm air will give a yardstick against which to measure any changes due to human activities.

Although the concentrations of OH determined here are certainly in line with expectations, I do have some concerns about some of the methods and assumptions used to derive these. I outline these below.

First, as noted by the authors on page 15 of the MS (in the "Implications and Uncertainties" section), the experiments were all carried out by first melting the snow sample, then adding the benzoic acid, then re-freezing, carrying out the illumination, then melting again for analysis. What we know about snow reactivity (Bartels-Rausch et al. ACP (2014); Dominé et al., JPCA (2013)) indicates that morphology may indeed play a significant role in chemistry, so it could well be important here as well. As well, Kahan et al. (ACP (2010), ES&T (2010)) have shown that OH reactivity on ice surfaces may be orders of magnitude different from that within the matrix. These considerations suggest that care should be taken in trying to infer OH concentrations within a disordered layer.

We agree with the reviewer that sample morphology and solute location could be important factors in ice photochemistry, including in our OH experiments. We tried to use a volatile probe (benzene) to determine OH in the field at Summit without melting the snow, but benzene was too volatile to stick appreciably to the snow grains. Thus we needed to use benzoate, which required melting and refreezing the snow.

We have expanded upon the solute partitioning and solute mixing uncertainties in the revised manuscript (section 3.5). As part of this we have added an additional sentence in to indicate that our results should be considered a "first-order estimate of hydroxyl radical kinetics in natural snow".

There do not appear to be significant amounts of HOOH at the air-ice interface (QLL) in natural snow (or in our reconstituted samples), so the QLL enhancement seen by Kahan and others is likely of little relevance to OH. We have revised the paragraph on uncertainties to more clearly state that HOOH in natural snow is probably in the bulk ice matrix, while HOOH in our reconstituted samples is probably in liquid-like regions. (As an aside, the QLL enhancements seen by Kahan in the cited references are a factor of 4 – 5, and not the orders of magnitude stated by the reviewer.)

Second, also as noted in the same section by the authors, the re-freezing not only alters the "natural" morphology, but could also alter the partitioning of the HOOH and organics within the sample. I think it is fair to say that we do not yet have any quantitative understanding about how different solutes distribute themselves as an aqueous solution freezes.

We agree and have included further discussion of HOOH location in our revised paragraph. We have some new experimental evidence that the solutes in samples such as ours will be

pushed into internal liquid-like regions and have cited this in-process manuscript in Cryosphere in the revised text of section 3.5.

Third, I do worry a bit that the bulk of the MS discusses laboratory kinetic results which the authors conclude to have been significantly impacted by an unknown contaminant. What are the implications of this for the general results?

The laboratory results have two main purposes. One is to compare OH kinetics in solution and ice, which can't be done in the (freezing) field. The second (initially unintended) purpose is to evaluate the approach of bringing field samples into the laboratory for study, which is commonly used since it's difficult to study snow photochemical processes in the field. Our laboratory results in the current manuscript show that such an approach is potentially perilous because of contamination. We were quite careful in our laboratory study of OH to avoid contamination, but our results show that it was a major problem. So while the laboratory results are not useful for understanding OH in field snow, they are a quantitative example of the contamination issues that can arise from the common practice of studying field samples in the laboratory.

On a more technical note, I am not convinced that the slopes and intercepts obtained from 4-point fits (such as those displayed in Figure 2) are as well constrained as the authors imply. Perhaps some discussion is warranted along these lines.

While we have used standard statistical propagation of errors, we agree with the reviewer that the resulting errors do not generally express all of the uncertainties inherent in the measurements. We addressed this in two ways in the original manuscript: (1) the discussion of the contamination of the samples studied in the laboratory (which dwarfs the experimentally determined uncertainties) and (2) reporting our field results to only one significant figure (Table 1). To this we have added our new statement of uncertainty in section 3.5, the statement that our results are a first-order estimate of OH kinetics. Together we think these components give a representative picture of the overall uncertainties.

Also on a technical note, I find the presentation in Section 2.7, concerning how the data were treated, to be quite confusing.

We have modified Section 2.7 in several places in order to clarify data treatment and the experimental methods.

Anonymous Referee #2

The manuscript describes a detailed kinetic study on the photochemistry of OH in ice. The manuscript is very clear and shows that experiments were very carefully done and evaluated. The data are sound and conclusions well justified. This and relevance of the hydroxyl radical in atmospheric chemistry of polar areas clearly grants publication after a minor addition. In particular, I was impressed by the comparison of OH and O₂ nicely discussing the freeze-concentration effect and how it differs for different reaction systems.

General comment on freeze-concentration effect and Liquid-Like-Regions: The term freeze-concentration effect was, to the best of my knowledge, first used to describe increasing reaction rates with the shrinking volume of the liquid fraction in a binary ice- solution system. As soon as the system was completely frozen, i.e. below the eutectic, reactivity ceased. (Takenaka, N., Ueda, A., Daimon, T., Bandow, H., Dohmaru, T., and Maeda, Y. "Acceleration Mechanism of Chemical Reaction by Freezing: the Reaction of Nitrous Acid with Dissolved Oxygen" *The Journal of Physical Chemistry* 100, no. 32 (1996): 13874–13884. doi:10.1021/jp9525806; Takenaka, N., Ueda, A., and Maeda, Y. "Acceleration of the Rate of Nitrite Oxidation by Freezing in Aqueous-Solution" *Nature* 358, no. 6389 (1992): 736–738. doi:10.1038/358736a0). I recommend to stick to this terminology which implies that one either has a freeze-concentration effect AND a liquid fraction in a frozen (binary) system or a (potentially high) reactivity in a completely frozen system that might take place in LLR/in a qll/in a liquid-like brine/or on the surface. More general, I'd appreciate a discussion on the phase behaviour of your samples. As the composition is not known, this is agreeable difficult. Nevertheless, maybe reactivity is similar in ice and liquid, because the ice is actually a mixture of ice and a reactant solution. For example, if H₂O₂ would be the origin of OH, one might expect liquid well down to -50 C (Foley, W. T. and Giguère, P. A. "Hydrogen Peroxide and Its Analogues: II. Phase Equilibrium in the System Hydrogen Peroxide-Water" *Canadian Journal of Chemistry-Revue Canadienne De Chimie* 29, no. 2 (1951): 123–132.) May I therefore suggest to include a short discussion on the possibility of the presence of liquid as reaction medium in your samples, both in the introduction and when comparing the results.

We consider a sample to be completely frozen when it has reached thermodynamic equilibrium with the surrounding sub-freezing temperature. At temperatures above the eutectic, the frozen sample should contain (mostly) pure water ice, concentrated liquid-like regions (LLRs) containing the solutes, and gas bubbles. (Below the eutectic temperature all of the solutes should be in a solid, frozen phase, although we and others have found evidence for reactive LLRs even below the eutectic (Bower and Anastasio, 2013; Cho et al., 2002).

The freeze-concentration factor, as defined by Takenaka (Takenaka and Bandow, 2007), is the ratio of solute concentration in the LLRs ("micropockets") compared to in the initial solution. In this same work the authors describe the LLR composition and kinetics using freezing-point depression. So our use of the terms "freeze-concentration effect" and "freeze-concentration factor", and our kinetic treatment of our data, are consistent with this work by Takenaka. We have modified the introduction to include this reference and mention that the freeze-concentration effect can also enhance the rate of thermal reactions in ice.

Consistent with these ideas, our recent imaging work on laboratory samples show three phases: pure ice, gas bubbles, and liquid-like regions containing the solutes (Hullar and Anastasio, In review). HOOH is the major precursor for OH, and is a major solute in the samples, but other solutes (including NO₃⁻, SO₄²⁻, NH₄⁺, and Ca²⁺) together account for approximately half of the total solutes (Anastasio et al., 2007; Dibb et al., 2010). Thus,

while a HOOH-water system doesn't capture the full complexity of our refrozen snow samples, the solutes are likely primarily in concentrated LLRs.

Specific comments:

P1 - 12ff: Laboratory studies show that . . . Does the contrast to other oxidant refer to both OH kinetics and concentration? Or only to the concentration? Could you reword.

We have modified the sentence to clarify our meaning.

P 3 - 15: While it has not been measured experimentally, the freeze-concentration effect might also alter . . . This is certainly a very valid hypothesis. I would suggest to underline it further by stating a few examples where reactivity was observed to change, referring to Klan and/or Donaldson earlier work (Bartels-Rausch, T., Jacobi, H.-W., Kahan, T. F., Thomas, J. L., Thomson, E. S., Abbatt, J. P. D., Ammann, M., Blackford, J. R., Bluhm, H., Boxe, C., Dominé, F., Frey, M. M., Gladich, I., Guzman, M. I., Heger, D., Huthwelker, T., Klán, P., Kuhs, W. F., Kuo, M. H., Maus, S., Moussa, S. G., McNeill, V. F., Newberg, J. T., Pettersson, J. B. C., Roeselova, M., and Sodeau, J. R. "A Review of Air–Ice Chemical and Physical Interactions (AICI): Liquids, Quasi-Liquids, and Solids in Snow" *Atmospheric Chemistry and Physics* 14, no. 3 (2014): 1587–1633. doi:10.5194/acp-14-1587-2014)

In this portion of the introduction we are referring specifically to the possibility that the concentration of hydroxyl radical might be enhanced in ice compared to in solution. Since there are no published data that address this topic (e.g., the Bartels-Rausch paper does not), we haven't added a citation. We do talk more broadly about oxidants in general earlier in the paragraph and we have added a new reference from Grannas' group (Fede and Grannas, 2015) to this portion of the text.

P 8 - 10: This is surprising as the apparent rate constant of OH towards organics spans several orders of magnitude (Schwarzenbach, R. P., Gschwend, P. M., and Imboden, D. M. "Environmental Organic Chemistry" (2005): doi:10.1002/0471649643). Would this imply that reactivity of DOC is dominated by one class of organics? Could you comment (or further support) that statement in the manuscript.

We were surprised by this result as well, although it turns out that the surface water community has known this for some time. As we discuss in detail in our previous work (Arakaki et al., 2013), while there are a wide range of bimolecular OH rate constants for organic compounds, when expressed in terms of carbon concentration (i.e., in units of $L \text{ mol-C}^{-1} \text{ s}^{-1}$, rather than in terms of the molar concentration of compound), the average OH rate constant is fairly robust over a very wide range of carbon numbers (from approximately 2 to 30). Since the organic carbon composition in natural samples is very complex, likely composed of 1000s of compounds, we suspect that the relatively robust result in natural samples is because of averaging across so many species. On the other hand, the relative standard deviation of the average rate constant is 50% for atmospheric waters, so there is still significant spread between different samples. We have added sentence about this to section 2.8 in the revised manuscript; for more information we encourage readers to see the Arakaki et al. reference.

P 11 - 6: "if the blank samples. . .contamination Milli-Q. . .". This is a reasonable assumption. Could you further support it by stating the type of Milli-Q used, does it use UV and does it maybe not filter the organics efficiently? Then one might expect high peroxide concentrations.

We defined the type of Milli-Q system in section 2.1. We have added a note that the system does not treat the water with UV radiation in section 2.1 and on page 11 in the revised manuscript.

References cited:

Anastasio, C., Galbavy, E. S., Hutterli, M. A., Burkhart, J. F., and Friel, D. K.: Photoformation Of Hydroxyl Radical On Snow Grains At Summit, Greenland, *Atmospheric Environment*, 41, 5110-5121, 2007.

Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S., Kimura, T., Tsuchi, A., and Miyagi, Y.: A General Scavenging Rate Constant for Reaction of Hydroxyl Radical with Organic Carbon in Atmospheric Waters, *Environmental Science & Technology*, 47, 8196-8203, 2013.

Bower, J. P. and Anastasio, C.: Using Singlet Molecular Oxygen To Probe The Solute And Temperature Dependence Of Liquid-Like Regions In/On Ice, *J Phys Chem A*, 117, 6612-6621, 2013.

Cho, H., Shepson, P. B., Barrie, L. A., Cowin, J. P., and Zaveri, R.: NMR Investigation Of The Quasi-Brine Layer In Ice/Brine Mixtures, *Journal of Physical Chemistry B*, 106, 11226-11232, 2002.

Dibb, J. E., Ziemba, L. D., Luxford, J., and Beckman, P.: Bromide and other ions in the snow, firn air, and atmospheric boundary layer at Summit during GSHOX, *Atmospheric Chemistry and Physics*, 10, 9931-9942, 2010.

Fede, A. and Grannas, A. M.: Photochemical Production of Singlet Oxygen from Dissolved Organic Matter in Ice, *Environmental Science & Technology*, 49, 12808-12815, 2015.

Hullar, T. and Anastasio, C.: Direct visualization of solute locations in laboratory ice samples, *The Cryosphere*, doi: 10.5194/tc-2015-197, In review. <http://dx.doi.org/10.5194/tc-2015-5197>, In review.

Takenaka, N. and Bandow, H.: Chemical Kinetics Of Reactions In The Unfrozen Solution Of Ice, *Journal of Physical Chemistry A*, 111, 8780-8786, 2007.

Hydroxyl Radical in/on Illuminated Polar Snow: Formation Rates, Lifetimes, and Steady-State Concentrations

Zeyuan Chen¹, Liang Chu¹, Edward S. Galbavy^{1,2}, Keren Ram¹, Cort Anastasio¹

¹Department of Land, Air, and Water Resources, University of California, Davis, CA, 95616, USA.

5 ²Now at NOVA Engineering and Environmental, Panama City Beach, FL, 32408, USA.

Correspondence to: Cort Anastasio (canastasio@ucdavis.edu)

Keywords: ice photochemistry, snowpack, oxidants, organic carbon, reactive halogens

Abstract. While hydroxyl radical ($\cdot\text{OH}$) in the snowpack is likely a dominant oxidant for organic
10 species and bromide, little is known about the kinetics or steady-state concentrations of $\cdot\text{OH}$ on/in snow
and ice. Here we measure the formation rate, lifetime, and concentration of $\cdot\text{OH}$ for illuminated polar
snow samples studied in the laboratory and in the field. Laboratory studies show that $\cdot\text{OH}$ kinetics and
steady-state concentrations are essentially the same for a given sample studied as ice and liquid; this is;
15 in contrast to other photooxidants, which show a concentration enhancement in ice relative to solution
as a result of kinetic differences in the two phases. The average production rate of $\cdot\text{OH}$ in samples
studied at Summit, Greenland is 5 times lower than the average measured in the laboratory, while the
average $\cdot\text{OH}$ lifetime determined in the field is 5 times higher than in the laboratory. These differences
indicate the polar snows we studied in the laboratory are affected by contamination, despite significant
20 efforts to prevent this; our results suggest similar contamination might be a widespread problem in
laboratory studies of ice chemistry. Steady-state concentrations of $\cdot\text{OH}$ in clean snow studied in the field
at Summit, Greenland range from $(0.8 \text{ to } 3) \times 10^{-15} \text{ M}$, comparable to values reported for mid-latitude
cloud and fog drops, rain, and deliquesced marine particles, even though impurity concentrations in the
snow samples are much lower. Partitioning of firm-air $\cdot\text{OH}$ to the snow grains will approximately double
the steady-state concentration of snow-grain hydroxyl radical, leading to an average $[\cdot\text{OH}]$ in near-
25 surface, summer Summit snow of approximately $4 \times 10^{-15} \text{ M}$. At this concentration, the $\cdot\text{OH}$ -mediated

lifetimes of organics and bromide in Summit snow grains are approximately 3 days and 7 hours, respectively, suggesting that hydroxyl radical is a major oxidant for both species.

1 Introduction

Hydroxyl radicals ($\cdot\text{OH}$) are ubiquitous in the atmosphere and react readily with most organic compounds; as such, they play an important role in chemical processing and in controlling the oxidizing capacity of the troposphere (Grannas et al., 2007b; Thompson and Stewart, 1991). Although $\cdot\text{OH}$ has been studied in atmospheric and terrestrial waters (including cloud and fog drops and surface waters) (Anastasio and McGregor, 2001; Arakaki et al., 2013; Ashton et al., 1995; Herrmann et al., 2010), little is known of $\cdot\text{OH}$ in snowpacks (Beyersdorf et al., 2007; Galbavy et al., 2007).

10 Snowpacks have two general components: the solid snow grains (and their associated impurities) and the interstitial (firn) air (Bartels-Rausch et al., 2014). From hydrocarbon decay measurements, $\cdot\text{OH}$ concentrations in the near-surface firn air at Summit, Greenland peak at 1.5×10^6 molecules cm^{-3} in the spring (close to the ambient air value) and 3.2×10^6 molecules cm^{-3} in July (approximately 20 – 30% lower than ambient) (Beyersdorf et al., 2007). There are no measurements of $\cdot\text{OH}$ concentrations in/on
15 snow grains, but some is known about the production rate of $\cdot\text{OH}$ (P_{OH}). Based on laboratory determinations of the quantum yields for $\cdot\text{OH}$ formation from photolysis of hydrogen peroxide (HOOH), nitrate (NO_3^-) and nitrite (NO_2^-), estimated values of P_{OH} in polar surface snow during summer are on the order of 100 - 300 nM hr^{-1} , with nearly all from photolysis of HOOH (Chu and Anastasio, 2005, 2003; France et al., 2007). The only measurements of $\cdot\text{OH}$ formation rates in/on snow
20 grains are at Summit, with summer values typically in the same range as estimated previously and HOOH accounting for 97% or more of photoformed $\cdot\text{OH}$ (Anastasio et al., 2007). While the sink for $\cdot\text{OH}$ has not been measured in snow grains, based on snow composition it appears that organic compounds are the dominant $\cdot\text{OH}$ sinks (Anastasio et al., 2007; Grannas et al., 2004). There are also no measurements of the $\cdot\text{OH}$ steady-state concentration in snow grains, which makes it difficult to estimate
25 the importance of $\cdot\text{OH}$ as an oxidant for organics and bromide.

As discussed in these previous studies, $\cdot\text{OH}$ reactions in/on snow are likely an important sink for snowpack trace species, which will influence the lifetimes, toxicities, and transformations of these

contaminants. For example, $\cdot\text{OH}$ reacts with snow-grain organic matter to form volatile organic compounds (VOCs) such as formaldehyde, which can be released to the atmosphere (Anastasio et al., 2007; Anastasio and Jordan, 2004; Domine and Shepson, 2002; Grannas et al., 2004; Jacobi et al., 2006). Oxidation by $\cdot\text{OH}$ can also convert snowpack halides (especially Br^-) into reactive volatile halogens, such as Br_2 , which can alter ozone and hydrocarbon chemistry in both the snow interstitial air and the atmospheric boundary layer (Abbatt et al., 2010; Anastasio et al., 2007; Chu and Anastasio, 2005; Grannas et al., 2007b; Pratt et al., 2013; Thomas et al., 2011; Wren et al., 2013). Additionally, $\cdot\text{OH}$ might alter ice core records of past atmospheres by reacting with trace species in snow (Anastasio and Jordan, 2004).

Recent work has shown that the steady-state concentrations of some oxidants can be much higher in/on illuminated ice compared to in the same sample studied as solution (Bower and Anastasio, 2013a; Bower and Anastasio, 2013b; [Fede and Grannas, 2015](#)). For example, the singlet oxygen concentration in ice can be higher by a factor of approximately 10,000 (Bower and Anastasio, 2013a) while the ice concentration of a triplet excited state can ~~increase-be higher~~ by a factor of roughly 100 (Chen, In preparation). These enhancements occur because of the freeze-concentration effect, where solutes are excluded to liquid-like regions (LLRs) in/on the ice, resulting in much higher effective concentrations in these small domains (Bower and Anastasio, 2013a; Bower and Anastasio, 2013b; Cho et al., 2002; Grannas et al., 2007a). [This same freeze-concentration effect is responsible for the enhancement in rates seen for some second-order thermal reactions in ice \(Takenaka and Bandow, 2007\).](#)

While it has not been measured experimentally, the freeze-concentration effect might also alter $\cdot\text{OH}$ concentrations within snow and ice (compared to the corresponding solution), which would alter ~~its~~ the impacts of this important radical. With this in mind, our goals in this work are to: (1) measure the steady-state concentration of $\cdot\text{OH}$, and the $\cdot\text{OH}$ kinetics (i.e., its rate of formation and lifetime), in polar snow samples, (2) compare $\cdot\text{OH}$ measurements in samples studied in the laboratory and in the field, and (3) examine how $\cdot\text{OH}$ kinetics and concentrations vary between solution and ice. To achieve these goals we used a benzoate probe technique to characterize photoformed $\cdot\text{OH}$ in polar snow samples studied in the lab (as solution and ice) and in the field (studied as ice).

2 Experimental Methods

2.1 Materials

Hydrogen peroxide and acetonitrile (Optima) were from Fisher, sodium benzoate (99%) and 2-nitrobenzaldehyde (2NB; 98%) were from Sigma-Aldrich, and *p*-hydroxybenzoate (*p*-HBA; 98%) was from TCI America. All chemicals were used as received. Purified water (“Milli-Q water”) was obtained from a Milli-Q Plus system ($\geq 18.2 \text{ M}\Omega \text{ cm}$) with an upstream Barnstead B-Pure cartridge to remove organics; the Milli-Q system does not use UV radiation to mineralize organic contaminants.

2.2 Snow sample collection

Surface snow samples (approximately 0 – 3 cm depth) were collected from undisturbed areas within the clean air sectors at Summit, Greenland (72.6°N, 38.5°W, 3200 m elevation) and Dome C, Antarctica (75.1°S, 123.4°E, 3270 m elevation). For the 2005 Summit samples, snow was removed with polytetrafluoroethylene (PTFE) instruments, set on a 3 ft \times 3 ft Teflon sheet, mixed, and placed in glass Schott bottles (100 or 250 mL). In 2006 and 2007, samples were collected directly into the Schott bottles. See Supplemental Table S1 for sampling conditions. Samples to be studied in the laboratory were held in cold, dark storage at Summit and Dome C and shipped frozen to UC Davis, where they were stored in a freezer ($-20 \text{ }^\circ\text{C}$) for 2 to 26 months prior to being studied. Samples studied in the field at Summit were used within a week of collection. The field blank for Summit was prepared using Milli-Q from the field that was frozen in Schott bottles and shipped back to UC Davis with the samples. The Dome C field blank was Milli-Q water shipped from our laboratory, frozen at Dome C, and then shipped back with the samples.

2.3 Laboratory sample preparation

Each laboratory sample was studied twice: once to determine the formation rate of $\cdot\text{OH}$ (P_{OH}) and once to determine the apparent rate constant for $\cdot\text{OH}$ destruction (k'_{OH}). For each type of test, the snow sample was first melted overnight in the refrigerator. To determine P_{OH} , 200 μM benzoate (BA) was added to a portion of the melted sample to scavenge essentially all of the $\cdot\text{OH}$ formed during illumination. For measuring k'_{OH} , we took four aliquots of the sample, added 100 μM HOOH to each (to

increase the rate of $\cdot\text{OH}$ formation and better determine the $\cdot\text{OH}$ sink), and added a different BA concentration (typically between 20 and 200 μM) to each. Samples studied as ice were frozen in a covered, custom-built, Peltier-cooled freeze chamber at $-10\text{ }^\circ\text{C}$ (Bower and Anastasio, 2013a). Laboratory blanks were prepared in the same way as the samples but with fresh Milli-Q instead of melted snow. Based on recent imaging work (Hullar and Anastasio, In review), we expect that our laboratory-prepared ice samples contain solutes in concentrated liquid-like regions, often in contact with internal air bubbles, in a (mostly) pure ice matrix.

2.4 Laboratory container types and cleaning treatments

We tested three types of sample containers in the lab to explore which would minimize contamination:

(1) 1-mL white PTFE Teflon beakers (15 mm H, 8 mm ID, Fisher Scientific), (2) 4-mL rectangular quartz cells (1-cm path length) with air-tight screw caps and Teflon septa (FUV, Spectrocell), and (3) 400- μL quartz tubes (30 mm L, 5 mm ID, 1 mm wall thickness) custom-made from GE 021 quartz and sealed with white silicone caps. We also explored two methods of extra cleaning for the containers after completing our normal cleaning procedures: (1) add Milli-Q to the container and illuminate for 24 hours with 254 nm radiation in an RPR-100 photoreactor equipped with 16 mercury lamps (25 W) (Southern New England Ultraviolet Company), and (2) in the same way but with 100 μM HOOH added to the Milli-Q prior to illumination as a photochemical source of $\cdot\text{OH}$.

2.5 Laboratory sample illumination and analysis

For laboratory determinations of $\cdot\text{OH}$ kinetics, we apportioned samples (with added BA or with BA and 100 μM HOOH; Section 2.3) into cleaned (HOOH + UV) quartz tubes with silicon caps. Samples were held in a custom designed, Peltier-cooled chamber illuminated with simulated sunlight (Ram and Anastasio, 2009). The solar simulator simulates total global solar radiation at a solar zenith angle of 48.2° . Prior to illumination, samples were allowed to thermally equilibrate in the illumination system chamber in the dark (30 min for liquid samples and 60 - 80 min for ice samples). During illumination $\cdot\text{OH}$ reacts with BA to form the stable product *p*-HBA. To determine the rate of *p*-HBA formation, 50 μL of sample (10 μL for rinse and 40 μL for injection) was removed at known times and *p*-HBA was

measured using HPLC with UV-Vis detection (Anastasio et al., 2007). For ice samples, the entire tube was removed for a given time point. We used the same procedures for laboratory and field samples. Dark control samples were run in parallel in the illumination system chamber, using the same sample and container conditions except that the tube and cap were covered with aluminum foil. Lastly, we measured the photon flux (I_λ) in all samples by measuring the direct photodegradation of 4 μM of 2NB ($j_{2\text{NB}}$) in the same type of container and same temperature as that in the sample measurement (Chu and Anastasio, 2003).

2.6 Field studies

For experiments at Summit, the snow sample in the sealed Schott bottle was left in a heated building ($\sim 10^\circ\text{C}$) in the dark to melt during the day. That evening we divided the sample into several portions, added a different concentration of BA (2 – 11 μM) to each, and created ice pellets by freezing 1-mL aliquots of sample on a PTFE sheet within a trench dug below the Summit Science Lab. (Note that we had planned to add 10 times more BA to the samples, but mistakenly made a lower concentration BA stock solution that was not discovered until after the field campaign ended.) After freezing we kept the pellets outdoors (below freezing) in 30-mL amber glass jars wrapped in aluminum foil until the moment of exposure. To start illumination, we placed the pellets on the snow surface, and quickly picked up an ice pellet as the time zero point (Anastasio et al., 2007). Samples were then collected at known times, put into sealed jars, melted in the dark, and analyzed for *p*-HBA and BA using HPLC. For dark samples we used the same pellets, but placed under a tub next to the sunlit samples to eliminate illumination. We also simultaneously illuminated blank controls, which were ice pellets made with Milli-Q water (and BA) instead of snow.

2.7 $\cdot\text{OH}$ kinetic analysis

For the laboratory work we studied each sample twice (section 2.3): once to determine the rate of $\cdot\text{OH}$ formation (in a sample portion with added BA) and once to determine the rate constant for $\cdot\text{OH}$ loss (with added HOOH and BA). In laboratory samples used to determine the formation rate of $\cdot\text{OH}$ we add (i.e., with 200 μM BA added), we expect so that essentially all of $\cdot\text{OH}$ will react with BA. Under this

condition the experimentally measured rate of $\cdot\text{OH}$ formation ($P_{\text{OH,exp}}$) is determined as

$$P_{\text{OH,exp}} = P_{p\text{-HBA}}/Y_{p\text{-HBA}} \quad (1)$$

where $P_{p\text{-HBA}}$ is the rate of $p\text{-HBA}$ formation, determined from the slope of a linear regression of [$p\text{-HBA}$] versus illumination time, and $Y_{p\text{-HBA}}$ is the yield of $p\text{-HBA}$ from the $\cdot\text{OH}$ reaction with BA (0.19 ± 0.0068 and 0.081 ± 0.014 for solution and ice, respectively (Anastasio and McGregor, 2001; Chu and Anastasio, 2003)).

In laboratory samples with 100 μM added HOOH (and different amounts of BA), we determined the apparent rate constant for $\cdot\text{OH}$ destruction, k'_{OH} , which is the inverse of the $\cdot\text{OH}$ lifetime (τ_{OH}). This pseudo-first-order rate constant reflects the concentrations and reactivities of all of the $\cdot\text{OH}$ sinks in each sample and is equal to the product of the second-order rate constant for each sink i and its concentration, $[i]$, summed over all of the sinks, i.e., $k'_{\text{OH}} = \Sigma k_{\text{OH}+i}[i]$. The value of k'_{OH} for each sample is determined from a linear regression of $1/P_{p\text{-HBA}}$ versus $1/[\text{BA}]$ (i.e., the “inverse” plot) (Anastasio and McGregor, 2001; Zhou and Mopper, 1990):

$$k'_{\text{OH}} = k_{\text{BA}+\cdot\text{OH}} \times (\text{slope} / y - \text{intercept}) \quad (2)$$

where $k_{\text{BA}+\cdot\text{OH}}$ is the second-order rate constant for reaction of BA and $\cdot\text{OH}$ (in $\text{M}^{-1}\text{s}^{-1}$), estimated from $k_{\text{BA}+\cdot\text{OH}} = \exp(26.6 - (1194.8/T))$ for both solution and ice samples (Ashton et al., 1995). We subtracted the k'_{OH} contribution from the 100 μM HOOH added to each sample; this value is $4.5 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ based on a 2nd-order rate constant (Dorfman, 1973) for $\cdot\text{OH}$ and HOOH of $4.5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$. Lastly, we determined the steady-state concentration of $\cdot\text{OH}$ by combining the measured production rate (from the sample without added HOOH) and the measured k'_{OH} ,

$$[\cdot\text{OH}]_{\text{exp}} = \frac{P_{\text{OH,exp}}}{k'_{\text{OH}}} \quad (3)$$

The experimentally determined values of $P_{\text{OH,exp}}$ and $[\cdot\text{OH}]_{\text{exp}}$ (i.e., under the illumination conditions in our laboratory) were then normalized to give the values expected under midday, summer solstice sunlight conditions at Summit:

$$P_{\text{OH,Sum}} = P_{\text{OH,exp}} \times \frac{j_{2\text{NB,Sum}}}{j_{2\text{NB,exp}}} \quad (4)$$

$$[\cdot\text{OH}]_{\text{Sum}} = [\cdot\text{OH}]_{\text{exp}} \times \frac{j_{2\text{NB,Sum}}}{j_{2\text{NB,exp}}} \quad (5)$$

where $j_{2\text{NB,Sum}}$ is the rate constant for loss of the actinometer 2NB under midday, summer solstice sunlight at Summit (0.020 s^{-1}) (Galbavy et al., 2010) and $j_{2\text{NB,exp}}$ is the value measured on the day of a given laboratory experiment.

~~In the For_ field, each samples was only studied once, with different pellets from a given sample containing different concentrations of BA (section 2.6). For each BA concentration in the sample we determined the rate of p-HBA formation from a linear regression of [p-HBA] versus illumination time. We then combined these rates in an “inverse” plot of of $1/P_{p\text{-HBA}}$ versus $1/[\text{BA}]$ and, used it to we~~
determined $P_{\text{OH,Sum}}$, k'_{OH} (Eq. 2), and $[\cdot\text{OH}]_{\text{Sum}}$ ~~using the “inverse” plot from the ice pellets studied with a range of BA concentrations for the sample:~~

$$P_{\text{OH,Sum}} = (y - \text{intercept} \times Y_{p\text{-HBA}})^{-1} \quad (6)$$

$$[\cdot\text{OH}]_{\text{Sum}} = (k_{\text{BA}+\cdot\text{OH}} \times \text{slope} \times Y_{p\text{-HBA}})^{-1} \quad (7)$$

In contrast to our past work (Anastasio et al., 2007), [BA] in the ice pellets was stable and therefore was not needed to normalize p-HBA concentrations. Because the rates of formation and steady-state concentrations for the field samples were measured using ambient Summit sunlight, they were not normalized to $j_{2\text{NB,Sum}}$.

2.8 Calculated organic carbon concentration

Organic compounds are likely the dominant sink for $\cdot\text{OH}$ in/on snow grains. As shown by Arakaki et al., the second-order rate constant for $\cdot\text{OH}$ reaction with organic carbon (in units of $\text{L} (\text{mol-C})^{-1} \text{ s}^{-1}$) is very similar in different atmospheric waters and even in surface waters (Arakaki et al., 2013). These similarities indicate that the apparent $\cdot\text{OH}$ scavenging rate constant in environmental waters is primarily controlled by the organic carbon concentration and is relatively insensitive to differences in the complex mixtures of organic compounds in different samples. This result might be because natural samples

likely contain thousands of organic compounds, leading to a relatively robust average scavenging rate constant between different samples. Based on this finding Thus we estimated the concentration of dissolved organic carbon in each sample, [DOC] (mol-C L⁻¹), using:

$$[\text{DOC}] \approx \frac{k'_{\text{OH}}}{k_{\text{C,OH}}} \quad (8)$$

5 where k'_{OH} is the measured pseudo-first-order $\cdot\text{OH}$ scavenging rate constant (s⁻¹) and $k_{\text{C,OH}}$ is the general bimolecular rate constant between $\cdot\text{OH}$ and organic carbon, determined as $(3.8 \pm 1.9) \times 10^8$ L (mol-C)⁻¹ s⁻¹ in atmospheric waters (Arakaki et al., 2013).

3 Results and Discussion

3.1 Minimizing background contamination for $\cdot\text{OH}$ sinks

10 To examine whether our containers or handling might add $\cdot\text{OH}$ -scavenging contaminants to our samples, we first performed a series of tests on laboratory Milli-Q blanks in three different types of containers (quartz cells, Teflon beakers, and quartz tubes) with three types of additional cleaning methods (no additional cleaning, 254-nm UV treatment, HOOH + UV treatment). These additional methods were added after finishing with our standard cleaning procedures, which are described below.

15 After the additional cleaning, fresh Milli-Q and 100 μM HOOH was added to each container, BA was added at a range of concentrations to different sample aliquots, and the samples were illuminated in order to determine k'_{OH} .

The first container type we examined was a 1-cm rectangular, air-tight quartz cell, where the standard cleaning is copious Milli-Q rinsing before and after a rinsing with a 50:50 MeOH:1 M H₂SO₄

20 solution. As shown in Figure 1, without any additional cleaning, k'_{OH} at 293 K is 1.1×10^5 s⁻¹, which is quite high and corresponds to an estimated organic carbon concentration of 300 $\mu\text{mol-C L}^{-1}$ based on Eq. (8). UV irradiation decreases this background by approximately a factor of two, while the combination of HOOH + UV treatment decreases the background by a factor of 10 (Figure 1). However, because the expensive quartz cells had a tendency to crack upon sample freezing, we next

25 tested Teflon beakers as a container.

Ice and solution Milli-Q blanks in Teflon beakers have k'_{OH} values of $(1 - 1.7) \times 10^5 \text{ s}^{-1}$ using only our standard cleaning condition, which is Alconox wash, Milli-Q rinse, ethanol rinse, Milli-Q rinse, and then an overnight Milli-Q soak. In contrast, additional HOOH + UV treatment can produce $\cdot\text{OH}$ destruction apparent rate constants as low as seen in the quartz cells ($8 \times 10^3 \text{ s}^{-1}$). However, HOOH + UV treated Milli-Q blanks studied as ice were approximately 5 – 10 times higher, suggesting that contaminant gases can adsorb onto the beaker samples and that a closed container is needed.

Therefore, we tested homemade quartz tubes with silicon caps as our third container type. The standard cleaning procedure for these tubes is the same as the Teflon beakers. Figure 1 shows that with additional UV + HOOH treatment the quartz tubes can achieve $\cdot\text{OH}$ rate constants as low as the other containers, corresponding to organic carbon concentrations as low as $20 \mu\text{mol-C L}^{-1}$. Since the quartz tubes with HOOH + UV treatment gave low amounts of background $\cdot\text{OH}$ scavengers, were inexpensive, and only occasionally broke upon freezing, we conducted all laboratory measurements of P_{OH} and k'_{OH} in snow samples using these containers.

3.2 $P_{\text{OH,Sum}}$, k'_{OH} , and $[\cdot\text{OH}]_{\text{sum}}$ in laboratory illuminated samples

Figure 2 shows the results of laboratory experiments conducted on one of the snow samples (Summit 0526). Figure 2a illustrates that the rate of *p*-HBA formation increases with increasing [BA] due to [BA] intercepting a larger fraction of the photoformed $\cdot\text{OH}$ compared to the natural $\cdot\text{OH}$ scavengers in the sample. The competition between [BA] and other scavengers can be seen more clearly in Figure 2b where $P_{p\text{-HBA}}$ increases as the [BA] concentration increases and plateaus at BA concentrations of $50 \mu\text{M}$ and higher, which indicates BA is scavenging most OH at and above this concentration. Thus essentially all $\cdot\text{OH}$ will be scavenged by BA in our samples studied in the laboratory to measure $P_{\text{OH,exp}}$, where we used $200 \mu\text{M}$ of BA. Lastly, Figure 2c shows an example of the “inverse” plot used to find k'_{OH} (Eq. 2) in the laboratory samples, which is then combined with $P_{\text{OH,Sum}}$ (Eq. 4) to determine $[\cdot\text{OH}]$ (Eqs. 3 and 5).

Figure 3 shows values of $P_{\text{OH,Sum}}$ measured at 263 and 274 K for three different Summit samples, one Dome C sample, field blanks from Summit and Dome C, and a laboratory blank. Values of $P_{\text{OH,Sum}}$ are similar at Summit and Dome C (although there is only one Dome C sample) for ice samples (263 K)

and range from approximately 330 – 840 nM hr⁻¹, which is somewhat higher than past summer results measured at Summit (130 to 610 nM hr⁻¹) (Anastasio et al., 2007) and higher than the calculated rate of •OH formation on snow grains from HOOH photolysis (approximately 250 nM hr⁻¹) (Chu and Anastasio, 2005). Solution values (274 K) of $P_{OH,Sum}$ are similar to their corresponding ice values but generally slightly higher; this is likely due in part to the temperature dependence of the •OH quantum yield from HOOH photolysis, which is 10% higher at 274 K than at 263 K (Chu and Anastasio, 2005). Blank rates range from 80 to 160 nM hr⁻¹ and represent 10 to 40% of the corresponding sample values.

Figure 4 depicts the apparent rate constant for •OH destruction (k'_{OH}) and its inverse – the lifetime of •OH (τ_{OH}) – for the same set of samples shown in Figure 3. As was seen for P_{OH} , rate constants for •OH destruction (k'_{OH}) are similar between solution and ice for a given sample and are in the same range for the Summit and Dome C samples. On average, k'_{OH} in solution at 274 K is 20% higher than the ice value at 263 K. This small difference is probably due to the temperature dependence of the •OH bimolecular rate constants with organic scavengers. For example, $k_{BA^+,OH}$ at 274 K is 20% higher than the value at 263 K; many of the natural scavengers probably have similar temperature dependence.

k'_{OH} values for the samples without blank correction are in the range of $(3.0 - 7.8) \times 10^4 \text{ s}^{-1}$, while blank values range from $(0.83 - 2.8) \times 10^4 \text{ s}^{-1}$ (Figure 4). The average of the blank values is roughly 10, 30, and 60% of the corresponding average sample values at 293, 274, and 263 K, respectively. This suggests that the sample values without blank correction are upper bounds of the true values. We have not corrected the sample values for the blanks because the blank levels might or might not reflect contamination in the samples. For example, if the blank values are the result of ~~low levels of~~ contamination in the Milli-Q water (which was not treated with UV radiation) then this will not affect the samples. On the other hand, if the blank values result from low levels of organics in the illumination containers, then the samples would be expected to have the same background level.

Using results from Figure 3 and 4, we can combine $P_{OH,Sum}$ and k'_{OH} to calculate $[•OH]_{Sum}$ (Eq. 7). As shown in Figure 5, steady-state •OH concentrations normalized to midday, Summit summer solstice sunlight are $(2 - 5) \times 10^{-15} \text{ M}$ and are similar between ice and solution. These snow concentrations of •OH are comparable to average values in mid-latitude cloud and fog drops, marine particles, and rain, which are generally $(0.5 - 7) \times 10^{-15} \text{ M}$ (Anastasio and McGregor, 2001; Arakaki et al., 2013). This is

somewhat surprising since the polar snow samples are much cleaner than the mid-latitude drops and particles, which typically have orders of magnitude higher concentrations of contaminants; for example, organic carbon concentrations in Davis, CA fog waters (Anastasio and McGregor, 2001) are approximately 100 times higher than levels in polar snow (see below). In fact, $P_{\text{OH,Sum}}$ and k'_{OH} are both
 5 much smaller in the snow samples (expressed relative to the melted snow volume) than in atmospheric hydrometeors. However, since $[\cdot\text{OH}]$ is equal to the ratio of these parameters (Eq. 7), its value is very similar across mid-latitude drops and particles (Arakaki et al., 2013; Arakaki et al., 1999) as well as for the snow samples here.

This correlation between $\cdot\text{OH}$ sources and sinks is also responsible for the very similar
 10 concentrations of $\cdot\text{OH}$ between the solution and ice results for a given sample (Figure 5). The $\cdot\text{OH}$ concentration is determined by the balance between its rate of formation and its pseudo-first-order rate constant for loss:

$$[\cdot\text{OH}] = \frac{P_{\text{OH}}}{k'_{\text{OH}}} \quad (9)$$

For the melted snow samples this is approximately equal to

$$15 \quad [\cdot\text{OH}]_{\text{LIQ}} \approx \frac{j(\text{HOOH} \rightarrow \cdot\text{OH})[\text{HOOH}]_{\text{LIQ}}}{\sum k_{i+\cdot\text{OH}}[i]_{\text{LIQ}}} \quad (10)$$

where $j(\text{HOOH} \rightarrow \cdot\text{OH})$ is the rate constant for $\cdot\text{OH}$ formation from HOOH photolysis and i represents natural (likely organic) $\cdot\text{OH}$ scavengers. Upon freezing the snow solution, solutes should be mostly excluded to liquid-like regions containing high concentrations of solutes (Bartels-Rausch et al., 2014); for example, at 263 K the total solute concentration in LLRs is predicted to be 5.4 M (Cho et al., 2002),
 20 which represents a freeze-concentration factor of approximately 50,000 for a snow sample with an initial (melted) total solute concentration of 100 μM . As a result of this enhancement in solute concentrations, both P_{OH} and k'_{OH} in the LLRs should be approximately 50,000 times higher than in the melted sample. The resulting expression for $[\cdot\text{OH}]$ in the LLRs is

$$[\cdot\text{OH}]_{\text{LLR}} \approx \frac{j(\text{HOOH} \rightarrow \cdot\text{OH})[\text{HOOH}]_{\text{LLR}}}{\sum k_{i+\cdot\text{OH}}[i]_{\text{LLR}}} \quad (11)$$

25 Since the LLR concentration of a given solute is larger than the initial solution concentration by

approximately the freeze-concentration factor (F), we can re-express Eq. (11) as

$$[\cdot\text{OH}]_{\text{LLR}} \approx \frac{j(\text{HOOH} \rightarrow \cdot\text{OH})[\text{HOOH}]_{\text{LIQ}}F}{\sum k_{i+\cdot\text{OH}}[i]_{\text{LIQ}}F} \quad (12)$$

The similarities in the ice and solution results in Figure 5 indicate that the freeze-concentration factor is essentially the same for both the $\cdot\text{OH}$ sources and sinks. Thus, the freeze-concentration factor approximately cancels in Eq. (12) and the steady-state concentration for $\cdot\text{OH}$ in LLRs of ice is essentially the same as in solution. In contrast, in the case of $^1\text{O}_2^*$, the source of singlet oxygen is enhanced by the freeze-concentration factor but the sink (liquid H_2O) is essentially the same in LLRs and solution (Bower and Anastasio, 2013a; Bower and Anastasio, 2013b). The result is that $[^1\text{O}_2^*]$ in ice LLRs is enhanced by the ~~freeze-freeze~~-concentration factor, which is on the order of 10^4 or higher for typical polar snows (Bower and Anastasio, 2013a).

We can also use equations 9 – 12 to gain a better understanding of kinetics in the liquid-like regions. While Figures 3 and 4 show that P_{OH} and k'_{OH} are essentially the same for a sample studied as solution or ice, this similarity is somewhat misleading. As shown in the numerator of equation 12, the rate of $\cdot\text{OH}$ production in the LLRs should be F times higher than the solution rate. However, since we melt the sample to analyze it, we cannot determine the concentration of photoproducted p -HBA in the LLR volume, but only in the melted sample volume, which is higher than the LLR volume by a factor of F . The same should be true for the apparent rate constant for $\cdot\text{OH}$ loss: as shown in the denominator of equation 12, k'_{OH} in the LLR should be F times higher than the solution value, but we do not see this since we can only do our analysis in the melted sample. Thus, on a whole (melted) sample volume basis, P_{OH} and k'_{OH} are the same in solution and ice, but in the native LLR volume, both the rate of $\cdot\text{OH}$ formation and apparent rate constant for $\cdot\text{OH}$ loss are much higher than in solution. On the other hand, $[\cdot\text{OH}]$ is essentially the same in both ice and solution, regardless of whether we consider LLR or total melted sample volume, since it is the ratio of P_{OH} to k'_{OH} .

3.3 P_{OH} , k'_{OH} , and $[\cdot\text{OH}]$ in the field

In addition to the Summit and Dome C samples that we studied in the laboratory, we also studied a number of snow samples in the field at Summit. Of the 38 field samples (Figure 6), most were studied

only at one BA concentration, but four (experiments 207, 225, 254 and 263) used 2 to 4 BA concentrations and thus can be used to derive P_{OH} , k'_{OH} , and $[^{\bullet}OH]$. As can be seen from Table 1, P_{OH} values (7 to 200 nM hr⁻¹) for these four samples studied in the field are, on average, 5 times lower than values for samples studied in the laboratory as ice (350 to 600 nM hr⁻¹). While these are different sets of samples, this large difference indicates that the samples studied in the laboratory were contaminated, either from shipping, storage, or handling during the experiments. We can also estimate $^{\bullet}OH$ production rates for the 34 field samples that were studied at only one BA concentration (i.e., the green diamonds in Figure 6). Using the average of the slopes from the four experiments with multiple BA concentrations in Figure 6 (644 hr $\mu M^{-1} \mu M$), we can extrapolate each green diamond in Figure 6 to its corresponding y-intercept, and then use Eq. 6 to estimate P_{OH} . The resulting $^{\bullet}OH$ formation rates range from 9 to 370 nM hr⁻¹, with an average ($\pm \sigma$) value of 100 ± 90 nM hr⁻¹ ($n = 34$); these values are comparable to P_{OH} values for the four field samples studied with multiple BA concentrations (Table 1).

Similar to the field-lab relationship in P_{OH} , k'_{OH} measured in the field, $(0.2 - 1) \times 10^4$ s⁻¹, is, on average, 5 times lower than values measured in the laboratory ice samples, $(3.0 - 6.2) \times 10^4$ s⁻¹. These large differences in P_{OH} and k'_{OH} between samples studied in the lab and field indicate that transport to Davis, storage, and/or handling in the laboratory added significant amounts of contaminants that are both sources and sinks for $^{\bullet}OH$. This is also apparent from laboratory measurements of k'_{OH} for the blanks, which are only somewhat smaller than the sample results (Figure 4). In contrast to the P_{OH} and k'_{OH} results, values of $[^{\bullet}OH]$ measured in field are quite similar to those determined in the laboratory, with average values of 2×10^{-15} M and 3×10^{-15} M, respectively. However, $[^{\bullet}OH]$ is only close between the two sets of samples because the contamination in both P_{OH} and k'_{OH} essentially cancels in the calculation of the steady-state $^{\bullet}OH$ concentration (Eq. 9).

3.4 Organic carbon in samples

We next use our measured values of k'_{OH} to estimate concentrations of organic carbon in our samples. As shown in Table S4, organic carbon concentrations in snow samples studied in the laboratory range from 90 to 190 $\mu mol-C/L$ while the field and lab blanks (Milli-Q) range from 20 to 100 $\mu mol-C/L$. In contrast, estimated organic carbon concentrations in the field samples are 5 to 30 $\mu mol-C/L$ (Table 1);

these much lower DOC values for the field samples are consistent with the idea that there was significant contamination in the samples studied in the lab. Compared to previously reported OC values, the snow organic carbon concentrations estimated from our field k'_{OH} values are at the lower end of previous results for snow from Summit and Alert and are similar to snow and sea ice values from Barrow (Table 2). In comparison, concentrations in ice core samples tend to be significantly lower, indicating significant mineralization or volatilization of the organic carbon after snow deposition.

3.5 Implications and Uncertainties

In addition to photolysis of HOOH and other chromophores in/on snow grains, transport of firm air $\cdot OH$ is another source of hydroxyl radical to snow grains. Based on firm air concentrations of $\cdot OH$ measured at Summit (Beyersdorf et al., 2007), we previously estimated that the rate of mass transport of firm air $\cdot OH$ to the snow grains is roughly equal to the rate of $\cdot OH$ photoformation in/on the snow grains (Chu and Anastasio, 2005). Thus, accounting for gas-to-grain partitioning of $\cdot OH$ will approximately double our measured values of $P_{OH,Sum}$ and $[\cdot OH]_{Sum}$: based on our field results, the resulting snow-grain steady-state concentrations of $\cdot OH$ at 263 K are $(2 - 6) \times 10^{-15}$ M, with an average of 4×10^{-15} M. Based on 2nd-order rate constants of $\cdot OH$ with dissolved organic compounds (typically 1×10^9 M⁻¹ s⁻¹) (Ross, 1988) and bromide (1.06×10^{10} M⁻¹ s⁻¹) (Zehavi and Rabani, 1972), we estimate OC and Br⁻ lifetimes as 3 days and 7 hours, respectively. This suggests that snow-grain $\cdot OH$ is significant in the transformation of both snowpack organics and bromide, resulting in emissions of both volatile organic compounds and reactive halogen gases (Anastasio et al., 2007; Anastasio and Robles, 2007).

There are several important uncertainties in our results. Perhaps the most important is the uncertain impact of melting the snow samples and refreezing them as ice pellets, ~~(which is required in order to add the $\cdot OH$ probe).~~ This likely alters the partitioning of solutes between the different ice reservoirs, which could change the mixing and collocation of solutes, e.g., possibly moving hydrogen peroxide that is trapped in the bulk ice matrix into liquid-like layers. ~~Given that~~ The most important change in partitioning might be for HOOH, which ~~is the dominant source of snow-grain $\cdot OH$ (accounting for more than 97% of $\cdot OH$ formed)~~ (Anastasio et al., 2007; Chu and Anastasio, 2005) and is thought to be present in the bulk ice of surface snow (Jacobi et al., 2004; Neftel et al., 1984). ~~Our sample preparation~~

probably moves HOOH's movement from the bulk ice matrix to liquid-like regions (Hullar and Anastasio, In review) LLRs; this ~~during sample preparation~~ might increase the rate constant for HOOH photolysis (and thus the steady-state concentration of $\cdot\text{OH}$) by roughly a factor of 2 or 3 ~~based on measurements~~ since of HOOH photolysis in LLRs (Chu and Anastasio, 2005) appears to be more rapid than ~~flash-frozen ices in bulk ice~~ (Beine et al., 2012). ~~The change in solute locations~~ Melting and ~~refreezing the snow~~ might also affect the mixing state of solutes, possibly ~~moving combining species that were~~ initially separate ~~species together~~ or possibly having the opposite effect. The potential impact of this movement is highly uncertain. Given these potential alterations in solute location and mixing state associated with the melting and refreezing required for the benzoate probe method, our field measurements should be, at present, considered a first-order estimate of hydroxyl radical kinetics in natural snow. While our results are the first experimental determinations of the $\cdot\text{OH}$ sink and steady-state concentration in snow samples, we hope that future developments include a snow-grain $\cdot\text{OH}$ techniques that does not disturb snow morphology or solute location.

4 Conclusion

15 We have made the first complete measurements of $\cdot\text{OH}$ kinetics and steady-state concentrations in illuminated snow samples. For a given sample, we find that $\cdot\text{OH}$ concentrations are essentially the same whether the sample is studied as ice or liquid; the same is true for OH production rates and OH lifetimes. This lack of enhancement on ice is different from what we have found for singlet molecular oxygen and triplet excited states, both of which are enhanced by orders of magnitude in/on ice
20 compared to in solution.

Production rates of $\cdot\text{OH}$ in samples studied at Summit during summer range between approximately 10 – 200 nM hr⁻¹, similar to previously measured and modeled rates. The lifetime of $\cdot\text{OH}$ in these samples is on the order of 70 to 500 μs , corresponding to organic carbon concentrations of 30 to 5 $\mu\text{mol-C L}^{-1}$. $\cdot\text{OH}$ concentrations in near-surface snow at Summit are $(2 - 6) \times 10^{-15}$ M, with
25 approximately equal contributions from photolysis of snow-grain impurities (mostly HOOH) and partitioning of $\cdot\text{OH}$ from the firm air. Compared to these samples studied in the field, samples studied in the laboratory show higher $\cdot\text{OH}$ production rates and lower $\cdot\text{OH}$ lifetimes, both as a result of

contaminants, despite significant efforts to minimize contamination. Similar contamination might be important for many laboratory studies of snow chemistry and should be characterized when possible. Based on our field measurements, hydroxyl radical in/on deposited snow grains and on atmospheric snow and ice likely plays an important role in the transformation of organic compounds and bromide. 5 These $\cdot\text{OH}$ -mediated reactions will release volatile organic compounds and reactive bromine species to the atmospheric boundary layer, leading to ozone depletion, mercury oxidation, and alterations to HO_x chemistry.

Author contributions. L. Chu and C. Anastasio designed and carried out the laboratory sample 10 experiments. E. S. Galbavy, K. Ram, and C. Anastasio designed and carried out the field sample experiments. Z. Chen and C. Anastasio wrote the manuscript with contributions from L. Chu.

Acknowledgement. We dedicate this work in memory of Harry Beine, our friend and colleague, who travelled the world to explore the chemistry of Earth's air and snow. The authors thank the National 15 Science Foundation and the Greenland Home Rule Ministry of Environment and Nature for granting us permission to do research at Summit, Greenland. We also thank Barry Lefer, Bernhard Stauffer, Karen Guldbaek Schmidt, and Manuel Hutterli for snow samples; VECO Polar for superb logistical support; and Ted Hullar for editorial help. Funding for this work was provided by the National Science Foundation (grants 0455055 and 1214121).

20 **References**

Abbatt, J., Oldridge, N., Symington, A., Chukalovskiy, V., McWhinney, R. D., Sjostedt, S., and Cox, R. A.: Release of Gas-Phase Halogens by Photolytic Generation of OH in Frozen Halide-Nitrate Solutions: An Active Halogen Formation Mechanism?, *Journal of Physical Chemistry A*, 114, 6527-6533, 2010.

25 Anastasio, C., Galbavy, E. S., Hutterli, M. A., Burkhart, J. F., and Friel, D. K.: Photoformation Of Hydroxyl Radical On Snow Grains At Summit, Greenland, *Atmospheric Environment*, 41, 5110-5121, 2007.

- Anastasio, C. and Jordan, A. L.: Photoformation Of Hydroxyl Radical And Hydrogen Peroxide In Aerosol Particles From Alert, Nunavut: Implications For Aerosol And Snowpack Chemistry In The Arctic, *Atmospheric Environment*, 38, 1153-1166, 2004.
- 5 Anastasio, C. and McGregor, K. G.: Chemistry Of Fog Waters In California's Central Valley: 1. In Situ Photoformation Of Hydroxyl Radical And Singlet Molecular Oxygen, *Atmospheric Environment*, 35, 1079-1089, 2001.
- Anastasio, C. and Robles, T.: Light Absorption By Soluble Chemical Species In Arctic And Antarctic Snow, *Journal of Geophysical Research-Atmospheres*, 112, 14, 2007.
- 10 Arakaki, T., Anastasio, C., Kuroki, Y., Nakajima, H., Okada, K., Kotani, Y., Handa, D., Azechi, S., Kimura, T., Tshako, A., and Miyagi, Y.: A General Scavenging Rate Constant for Reaction of Hydroxyl Radical with Organic Carbon in Atmospheric Waters, *Environmental Science & Technology*, 47, 8196-8203, 2013.
- Arakaki, T., Miyake, T., Shibata, M., and Sakugawa, H.: Photochemical Formation And Scavenging Of Hydroxyl Radical In Rain And Dew Waters, *Nippon Kagaku Kaishi*, 1999. 335-340, 1999.
- 15 Ashton, L., Buxton, G. V., and Stuart, C. R.: Temperature-Dependence Of The Rate Of Reaction Of OH With Some Aromatic-Compounds In Aqueous-Solution - Evidence For The Formation Of A Pi-Complex Intermediate, *Journal of the Chemical Society-Faraday Transactions*, 91, 1631-1633, 1995.
- 20 Bartels-Rausch, T., Jacobi, H. W., Kahan, T. F., Thomas, J. L., Thomson, E. S., Abbatt, J. P. D., Ammann, M., Blackford, J. R., Bluhm, H., Boxe, C., Domine, F., Frey, M. M., Gladich, I., Guzman, M. I., Heger, D., Huthwelker, T., Klan, P., Kuhs, W. F., Kuo, M. H., Maus, S., Moussa, S. G., McNeill, V. F., Newberg, J. T., Pettersson, J. B. C., Roeselova, M., and Sodeau, J. R.: A Review Of Air-Ice Chemical And Physical Interactions (AICI): Liquids, Quasi-Liquids, And Solids In Snow, *Atmospheric Chemistry and Physics*, 14, 1587-1633, 2014.
- 25 Beine, H., Anastasio, C., Domine, F., Douglas, T., Barret, M., France, J., King, M., Hall, S., and Ullmann, K.: Soluble Chromophores In Marine Snow, Seawater, Sea Ice And Frost Flowers Near Barrow, Alaska, *Journal of Geophysical Research-Atmospheres*, 117, 12, 2012.
- Beyersdorf, A. J., Blake, N. J., Swanson, A. L., Meinardi, S., Dibb, J. E., Sjostedt, S., Huey, G., Lefer, B., Rowland, F. S., and Blake, D. R.: Hydroxyl Concentration Estimates In The Sunlit Snowpack At Summit, Greenland, *Atmospheric Environment*, 41, 5101-5109, 2007.
- 30 Bower, J. P. and Anastasio, C.: Measuring A 10,000-Fold Enhancement Of Singlet Molecular Oxygen ($^1\text{O}_2^*$) Concentration On Illuminated Ice Relative To The Corresponding Liquid Solution, *Atmospheric Environment*, 75, 188-195, 2013a.

- Bower, J. P. and Anastasio, C.: Using Singlet Molecular Oxygen To Probe The Solute And Temperature Dependence Of Liquid-Like Regions In/On Ice, *J Phys Chem A*, 117, 6612-6621, 2013b.
- Chen, Z., Anastasio, C.: Concentrations Of A Triplet Excited State Are Enhanced In Illuminated Ice, In preparation. In preparation.
- 5 Cho, H., Shepson, P. B., Barrie, L. A., Cowin, J. P., and Zaveri, R.: NMR Investigation Of The Quasi-Brine Layer In Ice/Brine Mixtures, *Journal of Physical Chemistry B*, 106, 11226-11232, 2002.
- Chu, L. and Anastasio, C.: Formation Of Hydroxyl Radical From The Photolysis Of Frozen Hydrogen Peroxide, *Journal of Physical Chemistry A*, 109, 6264-6271, 2005.
- Chu, L. and Anastasio, C.: Quantum Yields Of Hydroxyl Radical And Nitrogen Dioxide From The
10 Photolysis Of Nitrate On Ice, *Journal of Physical Chemistry A*, 107, 9594-9602, 2003.
- Domine, F. and Shepson, P. B.: Air-Snow Interactions And Atmospheric Chemistry, *Science*, 297, 1506-1510, 2002.
- Dorfman, L., Adams, GE: Reactivities Of The Hydroxyl Radicals In Aqueous Solutions, NDRL/NIST Solution Kinetics Database, Gaithersburg, MD, U.S. Department of Commerce, Standard Reference
15 Data Program, 1973. 1973.
- Fede, A. and Grannas, A. M.: Photochemical Production of Singlet Oxygen from Dissolved Organic Matter in Ice, *Environmental Science & Technology*, 49, 12808-12815, 2015.
- France, J. L., King, M. D., and Lee-Taylor, J.: Hydroxyl (OH) Radical Production Rates In Snowpacks From Photolysis Of Hydrogen Peroxide (H₂O₂) And Nitrate (NO₃⁻), *Atmospheric Environment*, 41,
20 5502-5509, 2007.
- Galbavy, E. S., Anastasio, C., Lefer, B. L., and Hall, S. R.: Light Penetration In The Snowpack At Summit, Greenland: Part I - Nitrite And Hydrogen Peroxide Photolysis, *Atmospheric Environment*, 41, 5077-5090, 2007.
- Galbavy, E. S., Ram, K., and Anastasio, C.: 2-Nitrobenzaldehyde As A Chemical Actinometer For
25 Solution And Ice Photochemistry, *Journal of Photochemistry and Photobiology a-Chemistry*, 209, 186-192, 2010.
- Grannas, A. M., Bausch, A. R., and Mahanna, K. M.: Enhanced Aqueous Photochemical Reaction Rates After Freezing, *Journal of Physical Chemistry A*, 111, 11043-11049, 2007a.
- Grannas, A. M., Jones, A. E., Dibb, J., Ammann, M., Anastasio, C., Beine, H. J., Bergin, M.,
30 Bottenheim, J., Boxe, C. S., Carver, G., Chen, G., Crawford, J. H., Domine, F., Frey, M. M., Guzman, M. I., Heard, D. E., Helmig, D., Hoffmann, M. R., Honrath, R. E., Huey, L. G., Hutterli, M., Jacobi, H.

- W., Klan, P., Lefer, B., McConnell, J., Plane, J., Sander, R., Savarino, J., Shepson, P. B., Simpson, W. R., Sodeau, J. R., von Glasow, R., Weller, R., Wolff, E. W., and Zhu, T.: An Overview Of Snow Photochemistry: Evidence, Mechanisms And Impacts, *Atmospheric Chemistry and Physics*, 7, 4329-4373, 2007b.
- 5 Grannas, A. M., Shepson, P. B., and Filley, T. R.: Photochemistry And Nature Of Organic Matter In Arctic And Antarctic Snow, *Global Biogeochemical Cycles*, 18, 10, 2004.
- Herrmann, H., Hoffmann, D., Schaefer, T., Brauer, P., and Tilgner, A.: Tropospheric Aqueous-Phase Free-Radical Chemistry: Radical Sources, Spectra, Reaction Kinetics and Prediction Tools, *Chemphyschem*, 11, 3796-3822, 2010.
- 10 Hullar, T. and Anastasio, C.: Direct visualization of solute locations in laboratory ice samples, *The Cryosphere*, doi: 10.5194/tc-2015-197, In review. <http://dx.doi.org/10.5194/tc-2015-5197>, In review.
- Jacobi, H. W., Annor, T., and Quansah, E.: Investigation Of The Photochemical Decomposition Of Nitrate, Hydrogen Peroxide, And Formaldehyde In Artificial Snow, *Journal of Photochemistry and Photobiology a-Chemistry*, 179, 330-338, 2006.
- 15 Jacobi, H. W., Bales, R. C., Honrath, R. E., Peterson, M. C., Dibb, J. E., Swanson, A. L., and Albert, M. R.: Reactive trace gases measured in the interstitial air of surface snow at Summit, Greenland, *Atmospheric Environment*, 38, 1687-1697, 2004.
- Lafreniere, M. J. and Sharp, M. J.: The Concentration And Fluorescence Of Dissolved Organic Carbon (DOC) In Glacial And Nonglacial Catchments: Interpreting Hydrological Flow Routing And DOC Sources, *Arctic Antarctic and Alpine Research*, 36, 156-165, 2004.
- 20 Legrand, M., Preunkert, S., Schock, M., Cerqueira, M., Kasper-Giebl, A., Afonso, J., Pio, C., Gelencser, A., and Dombrowski-Etchevers, I.: Major 20th Century Changes Of Carbonaceous Aerosol Components (EC, Winoc, DOC, HULIS, Carboxylic Acids, And Cellulose) Derived From Alpine Ice Cores, *Journal of Geophysical Research-Atmospheres*, 112, 12, 2007.
- 25 Neftel, A., Jacob, P., and Klockow, D.: Measurement of hydrogen peroxide in polar ice samples, *Nature*, 311, 43-45, 1984.
- Pratt, K. A., Custard, K. D., Shepson, P. B., Douglas, T. A., Pohler, D., General, S., Zielcke, J., Simpson, W. R., Platt, U., Tanner, D. J., Huey, L. G., Carlsen, M., and Stirm, B. H.: Photochemical Production Of Molecular Bromine In Arctic Surface Snowpacks, *Nature Geoscience*, 6, 351-356, 2013.
- 30 Preunkert, S., Legrand, M., Stricker, P., Bulat, S., Alekhina, I., Petit, J. R., Hoffmann, H., May, B., and Jourdain, B.: Quantification of Dissolved Organic Carbon at Very Low Levels in Natural Ice Samples by a UV-Induced Oxidation Method, *Environmental Science & Technology*, 45, 673-678, 2011.

Ram, K. and Anastasio, C.: Photochemistry Of Phenanthrene, Pyrene, And Fluoranthene In Ice And Snow, Atmospheric Environment, 43, 2252-2259, 2009.

Ross, A. B., Mallard, W. G., Helman, W. P., Buxton, G. V., Huie, R. E., & Neta, P. : NDRL-NIST solution kinetics database, ver. 3.0, Notre Dame Radiation Laboratory, Notre Dame, IN and National
5 Institute of Standards and Technology, Gaithersburg, MD 1998, J. Phys. Chem. Ref. Data, 17, 1988.

Takenaka, N. and Bandow, H.: Chemical Kinetics Of Reactions In The Unfrozen Solution Of Ice, Journal of Physical Chemistry A, 111, 8780-8786, 2007.

Thomas, J. L., Stutz, J., Lefer, B., Huey, L. G., Toyota, K., Dibb, J. E., and von Glasow, R.: Modeling Chemistry In And Above Snow At Summit, Greenland - Part 1: Model Description And Results,
10 Atmospheric Chemistry and Physics, 11, 4899-4914, 2011.

Thompson, A. M. and Stewart, R. W.: Effect Of Chemical-Kinetics Uncertainties On Calculated Constituents In A Tropospheric Photochemical Model, Journal of Geophysical Research-Atmospheres, 96, 13089-13108, 1991.

Wren, S. N., Donaldson, D. J., and Abbatt, J. P. D.: Photochemical Chlorine And Bromine Activation
15 From Artificial Saline Snow, Atmospheric Chemistry and Physics, 13, 9789-9800, 2013.

Zehavi, D. and Rabani, J.: Oxidation Of Aqueous Bromide Ions By Hydroxyl Radicals - Pulse Radiolytic Investigation, Journal of Physical Chemistry, 76, 312-319, 1972.

Zhou, X. L. and Mopper, K.: Determination Of Photochemically Produced Hydroxyl Radicals In Seawater And Fresh-Water, Marine Chemistry, 30, 71-88, 1990.

20

|

Tables and Figure Captions

Figure 1. Effects of container type and cleaning on the apparent rate constant for $\cdot\text{OH}$ destruction in Milli-Q blanks. Open bars represent containers with no treatment, slashed bars have UV treatment (container with Milli-Q illuminated for ≥ 8 hours with 254 nm radiation), and solid bars are containers treated with 100 μM added HOOH followed by ≥ 8 hours of UV treatment. Bars within a given treatment are listed in chronological order, from the initial to the final experiment. Error bars represent ± 1 standard error, based on propagated errors of the linear regression of $1/p\text{-HBA}$ vs. $1/[\text{BA}]$ and the 2nd-order rate constant for $\cdot\text{OH}$ with BA.

Figure 2. Example of kinetic results for Summit sample 0526 studied in the laboratory. Panel a) shows the increase in $p\text{-HBA}$ concentration during illumination in 4 aliquots of the sample containing different initial BA concentrations ($[\text{BA}]_0$) illuminated for 30 min with simulated sunlight. The diamonds (blue), squares (red), triangles (green), and circles (gray) correspond to BA concentrations of 20, 30, 50, and 200 μM , respectively. The lines represent linear regression fits to data at each BA concentration (with y-intercept fixed at zero). Panel b) shows the measured production rate of $p\text{-HBA}$ for each BA concentration and the accompanying regression fit. Error bars represent ± 1 standard error based on the linear regression slopes in panel a. Panel c) is the “inverse” plot and linear regression fit, whose slope and y-intercept are used to calculate P_{OH} , k'_{OH} and $[\cdot\text{OH}]$ (see Sect. 2.7 for more details). Error bars based on ± 1 standard error calculated in panel b.

Figure 3. Rate of $\cdot\text{OH}$ formation in Summit and Dome C snow samples studied as solution (274 K, red) and as ice (263 K, blue), after normalization to Summit summer solstice sunlight. Error bars represent ± 1 standard error, based on propagated errors in the rate of $p\text{-HBA}$ formation ($P_{p\text{-HBA}}$) and $Y_{p\text{-HBA}}$. Sample values are not adjusted for $\cdot\text{OH}$ formation in the blanks.

Figure 4. Laboratory-determined apparent rate constant for $\cdot\text{OH}$ destruction (left axis) and the corresponding $\cdot\text{OH}$ lifetime (right axis) in Summit and Dome C snow samples and Milli-Q field and lab

blanks (with 100 μM added HOOH) at 293 K (green), 274 K (red), and 263 K (blue). Error bars represent ± 1 standard error, based on propagated standard errors (see text for details).

Figure 5. Laboratory-determined steady-state $\cdot\text{OH}$ concentrations in Summit and Dome C snow samples at 274 K (red) and 263 K (blue). Error bars represent ± 1 standard error, based on propagated errors in the slope and y-intercept of the “inverse” plots.

Figure 6. Measurements of $\cdot\text{OH}$ in Summit snow samples studied in the field, including four samples studied with multiple BA concentrations (with experiment number listed). Green hollow diamonds represent the 34 other snow samples that were studied using just one BA concentration.

Table 1. Summary of snow samples studied in the field at Summit, Greenland.

Date	Expt #	Illumination Time ^a	T Range (K)	P_{OH} (nM/hr)	k'_{OH} (s ⁻¹)	τ_{OH} (μ s)	[[•] OH] (10^{-15} M)	[DOC] ^b (μ M)
5/30/05	207	10:38 - 16:01	252-254	7	2×10^3	500	0.8	5
6/24/05	225	11:44 - 15:54	259-261	100	9×10^3	100	3	20
7/28/05	254	10:38 - 15:45	266-268	200	1×10^4	70	3	30
8/3/05	263	10:07 - 15:34	260-265	50	1×10^4	100	1	30

^a Local time^b Estimated concentration of dissolved organic carbon in each sample (in units of μ mol-C / L), calculated based on the measured k'_{OH} value using Eq. (8).

5

Table 2. Average dissolved organic carbon concentrations in snow and ice from past reports.

Type	Location	Time Frame	Average [DOC] ($\mu\text{mol-C/L}$)	References
Snow	Summit, Greenland ^a	Spring (2001)	48	(Grannas et al., 2004)
		Summer (2001)	47	
		Fall (2001)	33	
		Winter (2002)	36	
	Alert, Nunavut, Canada ^a	May (2002)	58	(Grannas et al., 2004)
		February (2002)	17	
	Barrow, Alaska	Feb – Apr (2009)	20	(Beine et al., 2012)
	Taylor Valley, Antarctica ^a		< 8	(Lyons et al., 2007)
	Alberta, Canada ^b	October (1999)	23, 34	(Lafreniere and Sharp, 2004)
Ice Cores	Col du Dome	1850 – 1976	9.6 – 25.5 ^c	(Legrand et al., 2007)
	Vostok, Antarctica	9970 BP	0.43	(Preunkert et al., 2011)
	D47, Antarctica	9970 BP	0.14	(Preunkert et al., 2011)
	South Pole	1010 BP	0.62	(Preunkert et al., 2011)
	Summit, Greenland	Winter (1020)	0.8	(Preunkert et al., 2011)
		Summer (1020)	3	
	Mt. Blanc, French Alps	Winter (1925-1936)	3.8	(Preunkert et al., 2011)
		Summer (1925-1936)	8.2	
Sea Ice	Barrow, Alaska		26	(Beine et al., 2012)

^a Total (dissolved and particulate) organic carbon concentration.

^b Snowmelt from a forest (23 $\mu\text{mol-C/L}$) and meadow (34 $\mu\text{mol-C/L}$) sample.

^c Range of [DOC] values, not the average.




Figure 1






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6