

1

2 Pan-Eurasian Experiment (PEEX):

3 Towards a holistic understanding of the feedbacks and interactions in the land - atmosphere -
4 ocean- society continuum in the Northern Eurasian region

5

6 Hanna K. Lappalainen^{1,2}, Veli-Matti Kerminen¹, Tuukka Petäjä¹, Theo Kurten³, Aleksander
7 Baklanov^{4,5}, Anatoly Shvidenko⁶, Jaana Bäck⁷, Timo Vihma², Pavel Alekseychik¹, Meinrat O.
8 Andreae⁸, Stephen R. Arnold⁹, Mikhail Arshinov¹⁰, Eija Asmi², Boris Belan¹⁰, Leonid
9 Bobylev¹¹, Sergey Chalov¹², Yafang Cheng¹³, Natalia Chubarova¹², Gerrit de Leeuw^{1,2}, Aijun
10 Ding¹³, Sergey Dobrolyubov¹², Sergei Dubtsov¹⁴, Egor Dyukarev¹⁵, Nikolai Elansky¹⁶,
11 Kostas.Eleftheriadis¹⁷, Igor Esau¹⁸, Nikolay Filatov¹⁹, Mikhail Flint²⁰, Congbin Fu¹³, Olga
12 Glezer²¹, Aleksander Gliko²², Martin Heimann²³, Albert A. M. Holtslag²⁴, Urmas Hörrak²⁵,
13 Juha Janhunen²⁶, Sirkku Juhola²⁷, Leena Järvi¹, Heikki Järvinen¹, Anna Kanukhina²⁸, Pavel
14 Konstantinov¹², Vladimir Kotlyakov²⁹, Antti-Jussi Kieloaho¹, Alexander S. Komarov³⁰, Joni
15 Kujansuu¹, Ilmo Kukkonen³¹, Ella-Maria Duplissy¹, Ari Laaksonen², Tuomas Laurila², Heikki
16 Lihavainen², Alexander Lisitzin²⁰, Aleksander Mahura⁵, Alexander Makshtas³², Evgeny
17 Mareev³³, Stephany Mazon¹, Dmitry Matishov^{34,†}, Vladimir Melnikov^{35,36}, Eugene
18 Mikhailov³⁷, Dmitri Moiseev¹, Robert Nigmatulin²⁰, Steffen M. Noe³⁸, Anne Ojala⁷, Mari
19 Pihlatie¹, Olga Popovicheva³⁹, Jukka Pumpanen⁴⁰, Tatjana Regerand¹⁹, Irina Repina¹⁶, Aleksei
20 Shcherbinin²⁷, Vladimir Shevchenko²⁰, Mikko Sipilä¹, Andrey Skorokhod¹⁶, Dominick V.
21 Spracklen⁹, Hang Su¹³, Dmitry A. Subetto¹⁹, Junying Sun⁴¹, Arkady Yu. Terzhevik¹⁹, Yuri
22 Timofeyev³⁷, Yuliya Troitskaya³³, Veli-Pekka Tynkkynen⁴², Viacheslav I. Kharuk⁴³, Nina
23 Zaytseva²², Jiahua Zhang⁴⁴, Yrjö Viisanen², Timo Vesala¹, Pertti Hari⁷, Hans Christen
24 Hansson⁴⁵, Gennady G. Matvienko¹⁰, Nikolai S. Kasimov¹², Huadong Guo⁴⁴, Valery Bondur⁴⁶,
25 Sergej Zilitinkevich^{1,2,12,33} and Markku Kulmala¹

26 1 Department of Physics, University of Helsinki, 00014 Helsinki, Finland

27 2 Finnish Meteorological Institute, Research and Development, 00101 Helsinki, Finland

28 3 Department of Chemistry, University of Helsinki, 00014 Helsinki, Finland

29 4 World Meteorological Organization, 1211 Genève, Switzerland

30 5 Danish Meteorological Institute, Research and Development Department, 2100,

31 Copenhagen

32 6 International Institute for Applied Systems Analysis, 2361 Laxenburg, Austria

33 7 Department of Forest Sciences, University of Helsinki, 00014 Helsinki, Finland

34 8 Biogeochemistry and Multiphase Chemistry Departments, Max Planck Institute for
35 Chemistry, 55020 Mainz, Germany

36 9 Institute for Climate and Atmospheric Science, School of Earth and Environment,
37 University of Leeds, Leeds, LS2 9JT, UK

38 10 Institute of Atmospheric Optics, Russian Academy of Sciences, Tomsk 634021, Russia

39 11 Nansen International Environmental and Remote Sensing Center, St. Petersburg, Russia

40 12 Lomonosov Moscow State University, Faculty of Geography, Moscow 119899, Russia

41 13 Institute for Climate and Global Change Research & School of Atmospheric Sciences,
42 Nanjing University, 210023 Nanjing, China

43 14 Institute of Chemical Kinetics & Combustion, Russian Academy of Sciences, 630090
 44 Novosibirsk, Russia
 45 15 Institute of Monitoring of Climatic & Ecological Systems SB RAS, 634055 Tomsk,
 46 Russia
 47 16 A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences, Russia
 48 17 National Centre of Scientific Research "DEMOKRITOS", Greece
 49 18 Nansen Environmental and Remote Sensing Center/Bjerknes Centre for Climate
 50 Research, 5006 Bergen, Norway
 51 19 Northern Water Problems Institute, Karelian Research Center, Russian Academy of
 52 Sciences, 185003 Petrozavodsk, Russia
 53 20 P.P. Shirshov, Institute of Oceanology, Russian Academy of Sciences, Russian Academy
 54 of Sciences, 117997 Moscow, Russia
 55 21 Institute of Geography, Russian Academy of Sciences, Moscow, Russia
 56 22 Department of Earth Sciences of the Russian Academy of Sciences, Russian Academy of
 57 Sciences, 119991, Moscow, Russia
 58 23 Max-Planck-Institute for Biogeochemistry, 07745 Jena, Germany
 59 24 Wageningen University, 6708 Wageningen, Nederland
 60 25 Institute of Physics, University of Tartu, 18 Ülikooli St., 50090 Tartu, Estonia.
 61 26 University of Helsinki, Department of World Cultures, 00014 Helsinki, Finland
 62 27 Department of Environmental Sciences, University of Helsinki, 00014 Helsinki, Finland
 63 28 Russian State Hydrometeorological University, 195196 Saint Petersburg, Russia
 64 29 Institute of Geography, Russian Academy of Sciences, Moscow, Russia
 65 30 Institute of Physico-chemical & Biological Problems in Soil Science, Russian Academy
 66 of Sciences, 142290 Institutskaya, Russia
 67 31 University of Helsinki, Geophysics and Astronomy, 00014 Helsinki, Finland
 68 32 Arctic & Antarctic Research Institute, Russian Academy of Sciences, St. Petersburg
 69 199397, Russia
 70 33 Department of Radiophysics, Nizhny Novgorod State University, Nizhny Novgorod,
 71 Russia
 72 34 Southern Center of Russian Academy of Sciences, Rostov on Don, Russia, [†] deceased,
 73 20.August.2015
 74 35 Tyumen Scientific Center, Siberian Branch, Russian Academy of Science, Russia
 75 36 Tyumen State University, 625003 Tyumen, Russia
 76 37 Saint Petersburg State University, 7/9 Universitetskaya nab., St. Petersburg, 199034
 77 Russia
 78 38 Institute of Agricultural and Environmental Sciences, Estonian University of Life
 79 Sciences, 51014 Tartu, Estonia
 80 39 Skobeltsyn Institute of Nuclear Physics, Moscow State University, Department
 81 Microelectronics, Russia
 82 40 University of Eastern Finland, Department of Environmental Science, P.O.Box 1627
 83 FI-70211 Kuopio, Finland
 84 41 Graduate University of Chinese Academy of Sciences, 100049 Beijing, China
 85 42 Aleksanteri Institute and Department of Social Research, 00014 University of Helsinki,
 86 Finland
 87 43 Sukachev Forest Institute, Russian Academy of Sciences, Krasnoyarsk 660036, Russia
 88 44 Institute of Remote Sensing and Digital Earth, Chinese Academy of Sciences, Beijing,
 89 100094, China
 90 45 Environmental Science and Analytical Chemistry, Stockholm University, Sweden
 91 46 AEROCOSMOS Research Institute for Aerospace Monitoring, 105064, Moscow, Russia

92 *Correspondence to:* Hanna K. Lappalainen (hanna.k.lappalainen(at)helsinki.fi)

93 Abstract. The Northern Eurasian regions and Arctic Ocean will very likely undergo substantial
94 changes during the next decades. The arctic-boreal natural environments play a crucial role in
95 the global climate via albedo change, carbon sources and sinks, as well as atmospheric aerosol
96 production from biogenic volatile organic compounds. Furthermore, it is expected that global
97 trade activities, demographic movement, and use of natural resources will be increasing in the
98 Arctic regions. There is a need for a novel research approach, which not only identifies and
99 tackles the relevant multi-disciplinary research questions, but is also able to make a holistic
100 system analysis of the expected feedbacks. In this paper, we introduce the research agenda of
101 the Pan-Eurasian Experiment (PEEX), a multi-scale, multi-disciplinary and international
102 program started in 2012 (<https://www.atm.helsinki.fi/peex/>). PEEX is setting a research
103 approach where large-scale research topics are investigated from a system perspective and
104 which aims to fill the key gaps in our understanding of the feedbacks and interactions between
105 the land-atmosphere-aquatic-society continuum in the Northern Eurasian region. We introduce
106 here the state of the art of the key topics in the PEEX research agenda and give the future
107 prospects of the research, which we see relevant in this context.

108 Contents

109	1. Introduction.....	4
110	2.System perspective approach.....	5
111	2.1. Land system – state of the art and future research needs	8
112	2.1.1. Changing land ecosystem processes.....	8
113	2.1.2 Ecosystem structural changes and resilience.....	11
114	2.1.3. Risk areas of permafrost thawing	14
115	2.2. Atmospheric system - state of the art and future research needs	16
116	2.2.1. Atmospheric composition and chemistry	16
117	2.2.2. Urban air quality.....	22
118	2.2.3. Atmospheric circulation and weather	24
119	2.3. Aquatic system - state of the art and future research needs	27
120	2.3.1. The Arctic Ocean in the climate system.....	27
121	2.3.2. Arctic marine ecosystem	30
122	2.3.3. Lakes, wetlands and large-scale river systems	32
123	2.4. Social system – state of the art and future research needs	36
124	2.4.1. Land use and natural resources.....	36
125	2.4.2. Natural hazards	39

126	2.4.3. Social transformations	45
127	3. From process studies towards system understanding and quantification of feedbacks of	
128	Arctic-boreal regions	48
129	3.1. Hydrological cycle	50
130	3.2. Carbon cycle.....	51
131	3.3. Nitrogen cycle	53
132	3.4. Phosphorus cycle.....	55
133	3.5. Sulfur cycle	56
134	4. From system understanding to mitigation and adaptation strategies and decision making	
135	58

136

137

138 1. Introduction

139 The global environment is changing rapidly due to anthropogenic influences. As a result,
140 we are already facing several “Grand Challenges” in the 21st century (e.g., Smith 2010; Bony
141 et al., 2015, IPCC; Randers 2012). Two of these challenges, climate change and air quality,
142 are strongly influenced by human activities and their impacts on changing atmospheric
143 composition, more specifically on the concentrations of greenhouse gases, reactive trace gases,
144 and aerosol particles. In the future, the arctic-boreal natural environment will play a crucial
145 role in the global climate via albedo changes, carbon sources and sinks, as well as aerosol
146 production from biogenic volatile organic compounds (Arneeth et al., 2010; 2014; Ballantyne
147 et al., 2012; Carslaw et al., 2010; Kulmala et al., 2014).

148 In order to advance our understanding on interlinked grand challenges further, we need
149 a research approach that helps us to construct a holistic scientific understanding of the
150 feedbacks and interactions within the continuum of land-atmosphere-aquatic-systems and
151 society across different spatial and temporal scales. Therefore, we have established the Pan-
152 Eurasian Experiment (PEEX) program (<https://www.atm.helsinki.fi/peex/>), which is a multi-
153 disciplinary, multi-scale research initiative focusing on understanding biosphere-ocean-
154 cryosphere-climate-society interactions and feedbacks (Lappalainen et al., 2014, Kulmala et
155 al., 2015). PEEX fills some of the most critical scientific gaps needed for a holistic
156 understanding of the feedback mechanisms characteristic of the Northern Eurasian

geographical domain. Boreal forests and peat lands characterize the vast land areas of Northern Eurasia, with a major part of them situated in the Russian territory. In addition to natural environments, the PEEEX research program is also interested in different human-influenced environments: from urban to countryside, from megacities to non-populated remote areas, from areas of dispersed settlements and sparsely-built environments to heavily-industrialized regions. Thus, the research approach covers the Arctic and boreal regions situated in Northern Eurasia, and also the marine environments of the Arctic Ocean. PEEEX operates in an integrative way using tools from natural and social sciences such as in-situ and satellite observations, laboratory experiments, multi-scale models, and statistical data analyses, together with socio-economic analyses. The PEEEX research agenda covers spatial scales from regional to global and temporal scales and from seconds to decades (Kulmala et al., 2011b). The scientific results will be used for developing new climate scenarios on global and regional scales, for constructing reliable early-warning systems, and for the mitigation and adaptation planning of the Northern societies in the most efficient way. PEEEX aims to contribute to climate policy concerning topics important to the Northern Eurasian environment,, helping societies in building a sustainable future.

2.System perspective approach

Earth (System) Sciences (ESS) has emerged as one of the most rapidly developing scientific fields. The recent growth of ESS has been facilitated by the importance of understanding the fundamental scientific processes of climate change and air quality, as well as the increasing societal impact of this research area. The development has mainly taken place among natural sciences, while the collaboration between natural and social sciences to tackle climate change issues has started to emerge relatively slowly. A multi- and cross-disciplinary approach is thus needed to advance the solution-oriented understanding of grand challenges and to apply new knowledge for reliable climate scenario development, mitigation and

adaptation, as well as early warning system development. In addition to enhanced collaboration between different branches of science, there is a need for a next generation of multidisciplinary scientists able to connect the scientific issues with an understanding of the societal dimensions related to the grand challenges.

Climate change can be considered as the main driving force for system changes and their feedback dynamics, especially in the Arctic-boreal regions. It has already been estimated that the future warming in Northern high latitudes regions will be, on average, larger than that experienced at lower latitudes (IPCC, 2013, 2014). The climate change driven processes taking place in the Arctic provide a good example on how important it is to quantify feedback dynamics and at the same time study the specific research topics from the land-atmosphere-hydrosphere-cryosphere-societal system perspective. For example, the surface radiation balance regulates the melting and freezing of the pack ice, which in turn is a key climate regulator. Model simulations of Arctic clouds are particularly deficient, impeding correctly simulated radiative fluxes, which are vital for the estimation of the snow/ice-albedo feedback (Vavrus et al., 2009). Important, yet poorly-quantified players in the Arctic atmospheric system and climate change are the short-lived climate forcers (SLCF), such as black carbon and ozone. The climatic impacts of SLCFs are tightly connected with cryospheric changes of the land system, and associated with human activities. Models display diverse and often poor skill in simulating SLCF abundances both at the surface and vertically through the troposphere at high latitudes (Eckhardt et al., 2015; Emmons et al., 2015; Monks et al., 2015).

PEEX is setting a research approach where the large-scale research questions are studied from a system perspective, and which is also filling the key gaps in our understanding of the feedbacks and interactions between the land, atmosphere, aquatic, and societal systems in the Northern Eurasian region (Kulmala et al., 2015). We have structured the research agenda so that we have highlighted three thematic research areas per system (Fig. 1). The identification

of these key thematic research areas has been based on a bottom-up approach by researchers coming from Europe, Russia, and China, participating PEEEX meetings and conferences starting from 2012. These researchers first introduced a wide spectrum of specific research topics relevant to the Northern Eurasian region, which were then evaluated and classified. This bottom up process led to the so-called “system-based” structure with altogether twelve thematic research areas. This approach will piece by piece lead into a holistic system understanding, quantifying the dominant feedbacks and interactions between the systems, and providing an understanding of the dynamics of Arctic-boreal biogeochemical cycles (e.g., water, carbon, nitrogen, phosphorus, sulfur). In our approach, climate change is the key driver in the dynamics of the land, atmosphere, aquatic and societal systems (Kulmala et al., 2015). The large-scale thematic areas of each system and many of the research highlight topics introduced by the PEEEX research agenda are fundamentally related to climate-change-driven shifting GHG and SLCF formation processes and their primary and secondary feedbacks between socioeconomic and biogeochemical systems. When studying the Arctic-boreal feedback loops in a wider context, the PEEEX agenda addresses China as the most crucial source area of atmospheric pollution, having a significant impact on the chemical composition of the atmosphere over Northern Eurasia (Monks et al, 2015). One must keep in mind that solving air quality – climate interactions is also the key to practical solutions on local air quality problems in China.

In this paper, we introduce the state of the art of the selected thematic research areas and summarize the future research needs at large scale. This introduction serves a White Paper of the PEEEX research community. The thematic research areas relevant to the Land System are related to “changing ecosystem processes” (2.1.1), “ecosystem structural changes and resilience” (2.1.2), and “risk areas of permafrost thawing” (2.3.1). In the Land System research agenda, we address the following key issues: changing boreal forests biomass, Arctic greening,

and permafrost processes. The main research areas of the Atmospheric System research are “the specific characterization of the atmospheric composition and chemistry” (2.2.1), “urban air quality” (2.3.2.), and “the atmospheric circulation and weather” (2.2.3). In terms of atmospheric systems, we address oxidants, trace gases, greenhouse gases, and aerosols as atmospheric key components. We highlight that future advances in predicting urban air quality and improving weather forecasting are strongly based the the atmospheric boundary layer dynamics research (Holtslag et al., 2013).

The thematic research areas relevant to the Aquatic System are “the Arctic Ocean in the climate system” (2.3.1), “the Arctic maritime ecosystems” (2.3.2), and “the lakes, wetland and rivers systems” (2.3.3). Under these research areas, we focus on topics like Arctic sea ice changes, marine gross primary production, and Arctic pelagic food webs under environmental changes. Lakes and large-scale river systems have multiple roles and aspects of the physical environments, starting from water chemistry and algal blooms, and ending up with carbon and methane dynamics.

The thematic areas of the Societal System have a number of dimensions, but in the first phase the primary interest lies on studying the consequences of “the increasing use of natural resources” (2.4.1), on the growing number of “natural hazards” (2.4.2), and on “the social transformations” (2.4.3) in the Northern Eurasian region. We see topics like the future Siberian forest area together with fuel balance, forest fires effecting the carbon and nitrogen balance, and societal dimensions related to infrastructure degradation as the most important future research areas. In Section 3, we investigate the connections and interlinks between those four systems.

2.1. Land system – state of the art and future research needs

2.1.1. Changing land ecosystem processes

In the future, many Arctic-boreal processes will respond sensitively to climate change, affecting ecosystem productivity and functions. These changes may lead to unprecedented consequences, e.g., in the magnitude of the ecosystem carbon sinks, production of aerosol precursor gases, and surface albedo. We need first to develop methods for identifying the land regions and processes that are especially sensitive to climate change. Only after that are we able to analyze their responses.

Boreal forests are one of the largest terrestrial biomes, and account for around one third of the Earth's forested area (Global Forest Watch, 2002; <http://www.globalforestwatch.org/>). Nearly 70 % of all boreal forests are located in the Siberian region. The forest biomass, soils, and peatlands in the boreal forest zone together constitute one of the world's largest carbon reservoirs (Bolin et al., 2000; Kasischke, 2000; Schepaschenko et al., 2013). Due to their large forest surface areas and huge stocks of carbon (~320 gigatonnes of carbon; GtC), the boreal and Arctic ecosystems are significant players in the global carbon budget. Furthermore, permafrost, a dominant feature of Siberian landscapes, stores around 1700 GtC (Tarnocai et al., 2009). Boreal forests form the main vegetation zone in the catchment areas of large river systems, so they are an important part of the global water-energy-carbon feedbacks.

The forest biomass forms a climate feedback via the anticipated changes in nutrient availability and temperatures, affecting carbon sequestered into both the aboveground biomass and soil compartment. The Siberian forests are currently assumed to be a carbon sink, although with a large uncertainty range of 0-1 PgC yr⁻¹ (Gurney et al., 2002). However, these ecosystems are vulnerable to global climate change in many ways, and the effects on ecosystem properties and functioning are complicated. While higher ambient CO₂ concentrations and longer growing seasons may increase plant growth and productivity, as well as the storage of carbon to soil organic matter (e.g., Ciais et al., 2005; Menzel et al., 2006), warming affects respiration and ecosystem water relations in the opposite way (Bauerle et al.,

2012; Parmentier et al., 2011). Expected acceleration of fire regimes might also substantially impact the carbon balance in Arctic and boreal regions (Shvidenko and Schepaschenko, 2013).

One example of the potentially large feedbacks is the critical role that permafrost plays in supporting the larch forest ecotone in northern Siberia. The boreal forests in the high latitudes of Siberia are a vast, rather homogenous ecosystem dominated by larch. The total area of larch forests is around 260 million ha, or almost one-third of all forests in Russia. Larch forests survive in the semi-arid climate because of the unique symbiotic relationship they have with permafrost. The permafrost provides enough water to support larch domination, and the larch in turn blocks radiation, protecting the permafrost from intensive thawing during the summer season. The anticipated thawing of permafrost could decouple this relationship, and may cause a strong positive feedback, intensifying the warming substantially.

The ambient temperature, radiation intensity, vegetation type, and foliar area are the main constraints for the emission of biogenic volatile organic compounds (BVOCs) (Laothawornkitkul et al., 2009). This makes BVOC emissions sensitive to both climate and land use changes, via, e.g., increased ecosystem productivity or the expansion of forests into tundra regions. Although the inhibitory effect of CO₂ on the process level may be important, Arctic greening may strongly enhance the production of BVOCs in northern ecosystems (Arneeth et al., 2007; Sun et al., 2013). Open tundra may also act as a significant source for BVOCs, especially if the snow cover period changes (Aaltonen et al., 2012; Faubert et al., 2012). This would lead to negative climate feedbacks involving either aerosol-cloud or aerosol-carbon cycle interactions (Kulmala et al., 2013; 2014; Paasonen et al., 2013). Linear trends in the annual maximum Normalized Difference Vegetation Index (NDVI) over 15 years in the Northern areas of the Yamalo-Nenets, Okrug region in Russia, provide supporting evidence of the increasing biological activity and greening, and the potential for enhanced BVOC emissions (Fig.2).

In summary, even small proportional changes in ecosystem carbon uptake can switch terrestrial ecosystems from a net carbon sink to a carbon source, with consequent impacts on atmospheric CO₂ concentrations and global temperatures (e.g., Bala et al., 2013; Bodman et al., 2013, Mukhortova et al., 2015). This process has already been observed, particularly in disturbed forests of Northern Asia (Shvidenko and Schepaschenko 2014). Currently, we do not fully understand all the factors influencing carbon storage, or the links between biogeochemical cycles of carbon, water, and nutrients in a changing climate. However, the changes in these processes may be large, and their impacts may either amplify or decrease climate change, especially in the high northern latitudes.

2.1.2 Ecosystem structural changes and resilience

The ecosystem structural changes are tightly connected to adaptation needs, and to the development of effective mitigation and adaptation strategies. Predictions concerning the shifting of vegetation zones are important for estimating the impacts of the region on future global GHG, BVOC, and aerosol budgets. Furthermore, natural and anthropogenic stresses, such as land use changes and biotic and abiotic disturbances, are shaping ecosystems in the Arctic and boreal regions and have many important feedbacks to climate (see e.g., the review by Gauthier et al., 2015). In a warmer climate, northern ecosystems may become susceptible to insect outbreaks, drought, devastating forest fires, and other natural disasters. In addition, human impacts may cause sudden or gradual changes in ecosystem functioning. The ecosystem resilience is dependent on both the rate and magnitude of these changes. Recent studies have concluded that current estimates very likely overestimate the resilience of global forests and particularly boreal forests (Allen et al., 2015). In some cases, the changes may lead to system imbalance and to the crossing of a tipping point, after which the effects are irreversible. One of the most relevant research topics for the land system are to determine the

333 structural changes and tipping points of the ecosystem changes in the Northern Pan Eurasian
334 region.

335 Part of the expected ecosystem structural changes is related to the lengthening of the
336 growing season that is taking place the Arctic-boreal regions due to climate change. This
337 phenomenon, called “Arctic Greening”, is due to increased plant biomass growth and
338 advancing tree lines, turning previously open tundra into shrubland or forest (Myneni et al.,
339 1997; Xu et al., 2013). However, "browning" as a proxy of decreased productivity has also
340 been observed during recent decades in many boreal regions (Lloyd and Bunn 2007), including
341 vast territories of Central Siberia, together with a general downward trend in basal area
342 increment after the mid-20th century (Berner et al., 2013) and the overall decline in green from
343 2011 to 2014 in Arctic regions (Phoenix and Bjerke 2016). Current predictions on the extent
344 and magnitude of these processes vary significantly (Tchebakova et al., 2009; Hickler et al.,
345 2012; Shvidenko et al., 2013). It has been estimated that the northward shift of bioclimatic
346 zones in Siberia will be as large as 600 km by the end of this century (Tchebakova et al., 2009).
347 By taking into account that the natural migration rate of boreal tree species cannot exceed 200-
348 500 m per year, such a forecast implies major vegetation changes in huge areas. In addition,
349 we need to have a deeper understanding of the future role of the browning process and to re-
350 analyze the previous model predictions of arctic greening: to what extent are they wrong, and
351 why (Phoenix and Bjerke 2016)? This has important biophysical consequences and climatic
352 feedbacks. Changes in vegetation cover can, e.g., lead to albedo changes and therefore higher
353 net absorption of radiation in regions covered by forests compared to open vegetation (Jeong
354 et al., 2011). This modifies the local heat and vapor fluxes, and affects boundary layer
355 conditions as well as both local and larger-scale climate (Sellers et al., 1997).

356 Northern peatlands contain a significant part of the global soil organic matter reservoirs
357 (45% of the world's soil carbon; Post et al., 1982), and comprise one of the world’s largest

GHG sources (in particular CH₄) (IPCC 2013). The hydrological conditions are a major factor in determining the functioning of peatlands as carbon source or sink, and the carbon balance of the vast northern peatlands is extremely sensitive to human influence, be it through either management or climate change. For example, thawing of permafrost peatlands in tundra regions might change tundra ecosystems from a stable state into a dynamically changing and alternating land-water mosaic, with dramatic impacts on their GHG production (Heikkinen et al., 2004; Repo et al., 2009). Today, peatland management activities range from drainage and peat harvesting to establishing crop plantations and forests. A complete understanding of the climatic effects of peatland management remains a challenging question (Maljanen et al., 2010).

Northern ecosystems are frequently suffering from increased stresses and deterioration. There is seldom a single and clear cause for forest dieback, but rather the ecosystems are suffering from multiple stresses simultaneously (e.g., Kurz et al., 2008 a,b; Allen et al., 2010). This implies that a single stress factor may not be very dramatic for the resilience of the system, but when occurring simultaneously in combination with others, the system may cross a threshold (i.e., tipping point), and this may have dramatic consequences. Such perturbations and disturbances can include long-term pollutant exposures, but also stochastic events such as fires, flooding, windstorms, or insect population outbreaks, and human activities such as deforestation or the introduction of exotic plant or animal species. Disturbances of sufficient magnitude or duration can profoundly affect an ecosystem, and may force an ecosystem to reach a threshold beyond which a different regime of processes and structures predominates. Climate warming, precipitation changes during growth periods, and permafrost changes will substantially increase water stress, and consequently increase the risk of mortality for trees. This process has already clearly intensified over the entire circumpolar boreal belt (Allen et

al., 2010). As a consequence, ecosystems may turn into carbon sources rather than sinks (Parmentier et al., 2011).

In the future, boreal forest diebacks may occur due to mass infections of invasive pathogens or herbivores, such as the autumnal moth (*Epirrita autumnata*) or mountain bark beetle (*Dendroctonus ponderosae*), that have previously been climatically controlled by harsh winter conditions. The growth and life cycles of herbivores or their habitat conditions may change in such a way that the outbreak frequencies and intensities of previously relatively harmless herbivore populations increase (Hunter et al., 2014). At the same time as climate is changing, boreal vegetation is also exposed to increased anthropogenic influences by pollutant deposition and land use changes (Dentener et al., 2006; Bobbink et al., 2010; Savva and Berninger, 2010). Large industrial complexes may lead to local forest diebacks, as has been observed in the Kola region (e.g., Nöjd and Kauppi, 1995; Tikkanen, 1995; Kukkola et al., 1997) and in some regions of Siberia (Baklanov et al., 2013). Societal transformations may lead to abandonment of agricultural land or deterioration of previously managed forests.

2.1.3. Risk areas of permafrost thawing

The major part of the Northern Eurasian geographical region is covered by continuous permafrost. The fate of permafrost soils in high latitudes is important for global climate with regard to all greenhouse gases. Thawing of permafrost will also substantially alter the hydrological regimes, particularly in Northern Asia, which will lead to increasing water stress in forests and explosive enlargement of fire extent and severity as well as post fire successions (Shvidenko et al., 2013). These scenarios underline the urgent need for systematic permafrost monitoring, together with GHG measurements in various ecosystems. The treatment of permafrost conditions in climate models is still not fully developed (Bala et al., 2013). The major question is, how fast will the permafrost thaw proceed and how will it affect ecosystem

processes and ecosystem-atmosphere feedbacks, including hydrology and greenhouse gas cycling.

Understanding of the feedbacks between carbon and water cycling, ecosystem functioning, and atmospheric composition related to permafrost thawing is one of the important topics of the land system study (Heimann and Reichstein, 2008; Schuur et al., 2009; Arneeth et al., 2010). In high-latitude ecosystems with large, immobile carbon pools in peat and soil, the future net CO₂ and CH₄ exchange will depend on the extent of near-surface permafrost thawing, local thermal and hydrological regimes, and interactions with the nitrogen cycle (Tarnocai et al., 2009). The extra heat produced during microbial decomposition could accelerate the rate of change in active-layer depth, potentially triggering a sudden and rapid loss of carbon stored in carbon-rich Siberian pleistocene loess (yedoma) soils (Khvorostyanov et al., 2008).

The connection between the climate and the thermal conditions in the subsurface layers (soil and bedrock) is an important aspect. The warming of the atmosphere will inevitably result in the warming of the permafrost layer, and is easily observed in deep borehole temperature data. However, the changes depend on the soil and rock type as well as on the pore-filling fluids. As long as the pore-fill is still ice, the climatic changes are reflected mainly in the thickness of the active layer, and in slow diffusive temperature changes of the permafrost layer itself. In areas where the ground is dominated by low ground temperatures and thick layers of porous soil types (e.g., sand, silt, peat), the latent heat of the pore filling ice will efficiently ‘buffer’ and retard the final thawing. This is one of the reasons why relatively old permafrost exists at shallow depths in high-porosity soils. On the other hand, quite different conditions prevail in low-porosity areas, e.g., in crystalline rock areas.

The permafrost dynamics affect methane fluxes in many ways. Hot spots such as mud ponds emitting large amounts of CH₄ may form when permafrost mires thaw. In contrast, lakes

have occasionally disappeared as a result of the intensification of soil water percolation (Smith et al., 2005). The rapid loss of summer ice, together with increasing temperature and melting ice deposits, results in coastal erosion, physical destruction of surface in hilly areas, activation of old carbon and elevated CO₂ and CH₄ emissions from sea bottom sediments (Vonk et al., 2012). High methane emissions have been observed from the East Siberian Arctic shelf (Shakhova et al., 2010).

2.2. Atmospheric system - state of the art and future research needs

2.2.1. Atmospheric composition and chemistry

Atmospheric composition plays a central role in the Northern Eurasian climate system. In addition to greenhouse gases and their biogeochemical cycling discussed in more detail in section 3.2, key compounds in this regard are ozone and other oxidants, carbon monoxide, numerous organic compounds, as well as different types of aerosols and their precursors (SO₂ will be discussed in chapter 3). At the moment, there is a serious gap in our knowledge on tropospheric composition and chemistry over Russia and China, with particularly few observation programs being active over Siberia (Crutzen et al., 1998; Ramonet et al., 2002; Paris et al., 2008; Kozlova et al., 2008; Uttal et al., 2015, Paris et al., 2010 a,b; Sasakawa et al., 2010; Chi et al., 2013; Saeki et al., 2013; Ding et al., 2013a, 2013b; Berchet et al., 2015; Heimann et al., 2014).

There is thus an urgent need for harmonized, coordinated and comprehensive greenhouse gas, trace gas and aerosol in-situ observations over Northern Eurasia and China (long-term transport aspect) comparable to European and circumpolar data observations. In Fig. 3 we illustrate the geographical coverage of the ground stations that will be part of the coordinated, coherent, and hierarchic observation network in the Northern Eurasian region and in China.

2.2.1.1. Main pollutants

Little is known about whether and how the regional ozone budget in northern Pan-Eurasia differs from that in the rest of the northern hemisphere (Ding et al., 2008; Berchet et al., 2013). Arctic tropospheric ozone is significantly influenced by long-range import of ozone and precursors from mid-latitude sources, as well as by boreal wildfires (Ding et al., 2009; Wespes et al., 2012; Paris et al., 2010b; Vivchar et al., 2009). The role of biomass burning emissions in the ozone budget in high latitudes remains controversial. While most studies suggest significant ozone production in boreal smoke plumes (e.g., Paris et al., 2010b; Parrington et al., 2013; Jolleys et al., 2015), some observations from individual plumes suggest that O₃ production in boreal wildfire plumes may be weaker, or even turn into net destruction, compared to fire plumes at lower latitudes (Liang et al., 2011; Jaffe and Wigder, 2012). Recent modeling work has suggested that boreal fires produce a substantial large-scale enhancement in summertime ozone at high latitudes, which appears to be highly sensitive to differences in partitioning of reactive nitrogen among models (Arnold et al., 2015). The boreal biosphere, on the other hand, provides a large sink for tropospheric ozone (Paris et al., 2010b; Parrington et al., 2013). Given their importance for air quality and global greenhouse gas budget, more atmospheric measurements of O₃, its precursors and other pollutants over Siberia are needed (see Elansky, 2012). This is particularly the case in light of increasing local Arctic sources of ozone precursors (NO_x, VOCs) from, e.g., shipping and fossil fuel resource extraction (Roiger et al., 2015). Such datasets would be particularly useful for the evaluation of atmospheric chemistry models and satellite products.

The changes in the abundance of anthropogenic aerosols and their precursors in Northern Eurasia have been extensive during the last decades (Granier et al., 2011), and this has almost certainly contributed to the very different regional warming patterns over these areas (e.g., Shindell and Faluvegi, 2009). The main anthropogenic aerosols in this context are primary carbonaceous particles, consisting of organic and black carbon, as well as secondary sulfate

particles produced during the atmospheric transport of sulfur dioxide. These species, as well as nitrate, have also been found to dominate the aerosol composition at the ZOTTO site in central Siberia (Mikhailov et al., 2015a, 2015b; Ryshkevich et al., 2015). These aerosols cause large perturbations to the regional radiation budget downwind of major source areas in the Northern Eurasian region, and the resulting changes in cloud properties and atmospheric circulation patterns may be important even far away from these sources (Koch and Del Genio, 2010; Persad et al., 2012). In the snow-covered parts of Eurasia, long-range transported aerosols containing black carbon and deposited onto snow tend to enhance the spring and early-summer melting of the snow, with concomitant warming over this region (Flanner et al., 2009; Goldenson et al., 2012; Meinander et al., 2015; Atlaskina et al., 2015).

The most important natural aerosol type over large parts of Eurasia is secondary organic aerosol originating from atmospheric oxidation of biogenic volatile organic compounds (BVOC) emitted by boreal forests and possibly other ecosystems. Studies conducted in the Scandinavian part of the boreal zone indicate that new particle formation associated with BVOC emissions is the dominant source of aerosol particles and cloud condensation nuclei during summer time (Mäkelä et al., 1997; Kulmala et al., 2001; Tunved et al., 2006; Asmi et al., 2011; Hirsikko et al., 2011). The production of secondary organic aerosols associated with BVOC emissions has been estimated to induce large direct and indirect radiative effects over the boreal forest zone (Spracklen et al., 2008; Tunved et al., 2008; Lihavainen et al., 2009, 2015; Scott et al., 2014). The few continuous measurement data sets from Siberia suggest similarities in the frequency and seasonal pattern of new particle formation events between Siberia and Nordic stations (Dal Maso et al., 2007; Arshinov et al., 2012; Asmi et al., 2015). Measurements conducted at the ZOTTO site in central Siberia have shown that biogenic secondary organic aerosols reach high concentrations in summer and dominate the aerosol composition during this season (Mikhailov et al., 2015a,b; Ryshkevich et al., 2015). At this

site, however, new particle formation events are seen much less frequently than at the Nordic stations (Heintzenberg et al., 2011). At present, relatively little is known about the overall contribution of biogenic emissions to aerosol number or mass concentrations, or to the cloud condensation nuclei budget, in Northern Eurasia.

Other important natural aerosol types in Northern Eurasia are sea spray, mineral dust, and primary biogenic aerosol particles. Sea spray aerosol makes an important contribution to the atmospheric aerosol over the Arctic Ocean and its coastal areas (Zábori et al., 2012, 2013), and influences cloud properties over these regions (Tjernström et al., 2013). The climatic effects of sea spray are expected to change in the future as a result of changes in the sea ice cover and ocean temperatures (Struthers et al., 2011). Mineral dust particles affect regional climate and air quality over large regions in Asia, especially during periods of high winds and moderate precipitation. Mineral dust and primary biological aerosol particles (PBAP) particles are also effective ice nuclei (Hoose and Möhler, 2012), and have the potential to influence the radiative and other properties of mixed-phase cold clouds in the arctic-boreal regions. Over Northern Eurasia, PBAP typically contribute more than 20% of PM_{2.5} organic aerosol mass concentrations (Heald and Spracklen, 2009) and 25% of supermicron aerosol number concentrations (Spracklen and Heald, 2014). Ice nucleation, in general, is one of the key microphysical processes in the atmosphere that remain ill understood. However, a novel theoretical approach (Laaksonen, 2015; Laaksonen and Malila, 2016) has been shown to be superior to older theories in the case of water nucleation on solid surfaces, and it may open a completely new avenue in the studies of atmospheric ice formation.

Satellites provide information about spatial distributions of the column-integrated concentrations of aerosols (e.g., de Leeuw and Kokhanovsky, 2011; Andreae, 2009) and various trace gases including ozone and its precursors (Burrows et al., 2011). These atmospheric constituents are generally retrieved using passive instruments, which have good

sensitivity near the surface. However, retrieving information on the near-surface concentrations of pollutants requires assumptions on their vertical distributions. For instance, the retrieval of tropospheric ozone from satellite observations requires corrections for the high concentrations in the upper troposphere and lower stratosphere. For aerosols, which can only be retrieved in clear sky conditions, the situation may be complicated when disconnected layers are present with different types of aerosols. A solution may be the retrieval of aerosol vertical variation or the height of the aerosol layer using, e.g., active instruments (lidars), or retrieval using spectrally-resolved observations in the Oxygen-A band (e.g., Hollstein and Fisher, 2013), or instruments providing multiple viewing algorithms such as MISR (Nelson et al., 2013) or AATSR (Virtanen et al., 2014). Another complication for aerosols may be the vertical variation of the physical and chemical properties, which renders it difficult to obtain closure between column and ground-based in situ measurements (Zieger et al., 2015 and references cited therein). Nevertheless, good progress has been made in aerosol retrieval, and column-integrated aerosol measurements (AOD) from satellites and ground-based observations compare favorably (e.g., de Leeuw et al., 2015; Kolmonen et al., 2015). Measurements of trace gases from space using wavelengths in the thermal infrared suffer from low sensitivity in the lower troposphere (Pommier et al., 2010). All these factors may render the comparison against local ground-based in-situ observations difficult, although a possible way out could be the use of chemical transport models constrained by the satellite column measurements (e.g., de Laat et al., 2009; Stavrou, 2012; 2014), possibly together with sub-orbital airborne measurements of relevant species. Satellite-measured AOD has been successfully applied to obtain information on ground based aerosol mass concentrations (PM_{2.5}) (Xu et al., 2015; van Donkelaar et al., 2015). In addition, the use of multiple satellite instruments, with different characteristics, is proposed to obtain more accurate information on the transport of aerosols and trace gases and their vertical distribution (e.g., Naeger et al.,

2015). Recently, a technique has been demonstrated that makes it possible to derive CCN concentrations at cloud base using remote sensing of cloud properties (Rosenfeld et al., 2016).

2.2.1.2. Large-scale pollutant transport and sources

Of particular interest is the pollutant transport to Arctic areas, where they can influence the radiation budget and climate in various ways (Stohl, 2006; Warneke et al., 2009; Meinander et al., 2013; Eckhardt et al., 2015). Model simulations suggest that European emissions dominate Arctic pollutant burdens near the surface, with sources from North America and Asia more important in the mid and upper troposphere (Monks et al., 2015). The impact and influence of China and its polluted megacities on Arctic and boreal areas is a topic of key importance, given recent and rapid Chinese industrialization. Inter-continental pollution transport has also become of increased concern due to its potential influence on regional air quality. The pollutant export from North America and Asia has been characterized by intensive field campaigns (Fehsenfeld et al., 2006; Singh et al., 2006), but long-term research approaches are lacking.

Emissions from forest fires (van der Werf et al., 2006; Sofiev et al., 2013) and from agricultural fires in southern Siberia, Kazakhstan, and Ukraine (Korontzi et al., 2006) in spring and summer are large sources of trace gases such as carbon monoxide (Nédélec et al., 2005; Konovalov et al., 2014), as well as aerosol particles (Konovalov et al., 2015). Aerosols emitted by forest fires are of particular interest, since the strength of this source type depends on both climate change and human behavior (Pechony and Shindell, 2010), and since particles emitted by these fires have potentially large radiative effects over Eurasia (Randerson et al., 2006). We need comprehensive top-down emissions estimates, using inverse modeling constrained by satellite observations, in order to provide quantitative information on the source strength of aerosols and trace gases emitted by open fires.

Air pollution in monsoon Asia has two main characteristics. First, the total pollutant emission rate from fossil fuel combustion sources is very high, leading to a high concentration

of primary and secondary pollutants in Asia, especially in eastern China and northern India. Observations show that Asia is the only region where the concentrations of key pollutants, such as nitrogen oxides (Richter et al., 2005; Mijling et al., 2013) and their end-product ozone (Ding et al., 2008; Wang et al., 2009; Verstraeten et al., 2015), are still increasing. Second, in addition to the anthropogenic fossil fuel combustion pollutants, monsoon Asia is also influenced by intensive pollution from seasonal biomass burning and dust storms. For example, intensive forest burning activities often take place in south Asia during spring and in Siberia during summer, whereas intensive anthropogenic burning of agricultural straw takes place in the north and east China plains. Dust storms frequently occur in the Taklimakan and Gobi deserts in northwest China, and this dust is often transported over eastern China, southern China, the Pacific Ocean and even the entire globe (Nie et al., 2014). After mixing with other anthropogenic pollutants, biomass burning and mineral dust aerosols have been found to cause complex interactions in the climate system (Ding et al., 2013; Nie et al., 2014).

2.2.2. Urban air quality

The northern Eurasian urban environments are characterized by cities with strong anthropogenic emissions from local industry, traffic, and housing in Russia and China, and by megacity regions with alarming air quality levels like those of Moscow and Beijing. Bad air quality has serious health effects and damages ecosystems. In Beijing, for example, concentrations of atmospheric fine particles have been found to be more than 10 times higher than the safe level recommended by the World Health Organization (WHO) (Zheng et al., 2015). Furthermore, atmospheric pollutants and oxidants play a central role in climate change dynamics via their direct and indirect effects on global albedo and radiative transfer. A deeper understanding of the unpredicted chemical reactions between pollutants and identification of the most relevant feedbacks between air quality and climate at northern high latitudes and in

China is the most urgent task helping us to find practical solutions for more healthy air (Kulmala, 2015).

In Siberian cities, the air quality is strongly linked to climatic conditions typical for Siberia. Stable atmospheric stratification and temperature inversions are predominant weather patterns for more than half of the year. This contributes to the accumulation of different pollutants in the lowest layers of the atmosphere, thus increasing their impact on ecosystems and humans. In addition to the severe climatic conditions, human impacts on the environment in industrial areas and large cities continue to increase. In winter time, shallow and stably-stratified planetary boundary layers (PBL) typical for northern Scandinavia and Siberia are especially sensitive to even weak impacts and, therefore, deserve particular attention, especially in the conditions of environmental and climate change (Zilitinkevich and Esau, 2009; Esau et al., 2012; Davy and Esau, 2014; Wolf et al., 2014 ; Wolf and Esau, 2014). Unstably stratified PBLs interact with the free atmosphere mainly through turbulent ventilation at the PBL upper boundary (Zilitinkevich, 2012). This mechanism, still insufficiently understood and poorly modeled, controls the development of convective clouds, as well as dispersion and deposition of aerosols and gases, which are essential features of heat waves and other extreme weather events.

The worst air pollution episodes are usually associated with stagnant weather conditions with a shallow PBL, which promotes the accumulation of intensively emitted pollutants near the surface. The lower PBL is also influenced by the heavy pollution itself through its direct or indirect effects on solar radiation and hence the surface sensible heat flux (e.g., Ding et al., 2013b). The boundary layer - air pollution feedback will decrease the height of the PBL and result in an even more polluted PBL (Ding et al., 2013b; Wang et al., 2014, Petäjä et al., 2016). Therefore, considering the complex land surface types (city clusters surrounded by agricultural areas) and pollution sources, improving our understanding of the associated feedbacks is very

important for forecasting extreme air pollution episodes and for long-term policymaking. In order to understand this topic, more vertical measurements using aircraft, balloons, and remote sensing techniques, as well as advanced numerical models including all relevant processes and their couplings, are needed.

Planetary boundary layers are subject to diurnal variations, absorb surface emissions, control microclimate, air pollution, extreme colds, and heat waves, and are sensitive to human impacts. Very stable stratification in the atmosphere above the PBL prevents the compounds produced by the surface fluxes or surface emissions from efficiently penetrating from the PBL into the free atmosphere. This means that the PBL height and turbulent fluxes through the PBL upper boundary control local features of climate and extreme weather events, such as the heat waves associated with convection, or the strongly stable stratification events triggering the air pollution (Zilitinkevich et al., 2015). This concept (equally relevant to the hydrosphere) illustrates the importance of modeling and monitoring the atmospheric PBL height, which varies from dozens to thousands of meters (Zilitinkevich, 1991; Zilitinkevich et al., 2007; Zilitinkevich and Esau, 2009). To carry out a comprehensive inventory of the PBL height over Northern Eurasia is urgently needed.

2.2.3. Atmospheric circulation and weather

The ongoing environmental change and its amplification in Northern Eurasia pose special challenges to the prediction of weather-related hazards, and also to long-term impacts. A key question is how the atmospheric dynamics (synoptic scale weather, boundary layer characteristics) will change in Arctic and boreal regions. The recent changes in the Arctic sea-ice have been much more rapid than models and scientists anticipated about ten years ago. The role of the Arctic Ocean in the climate system and sea-ice changes have affected mid-latitude weather and climate, with central and eastern Eurasia among the regions with strongest effects (Vihma, 2014; Overland et al., 2015) (see section 2.3.1).

2.2.3.1. Atmospheric dynamics

The reliability of weather forecasts, and the extension of the time-range of useful forecasts is needed for minimizing economic and human losses from extreme weather and extreme weather related natural hazards. In Europe, this range is currently on average about 8–9 days (Bauer et al., 2015), which allows reliable early warnings to be issued for weather related hazards, such as windstorms and extreme precipitation events with flash floods. The time-range of useful forecasts has typically increased by a day per decade over the past three decades (Uppala et al., 2005). In the Northern Eurasian region, improved predictions can be used, for instance, for better prediction of thermal comfort conditions in Northern cities (Konstantinov et.al, 2014). A strong urban heat island effect has already been observed in urban areas of the Arctic with complex spatial and temporal structures (Konstantinov et al., 2015).

Understanding of planetary boundary layer (PBL) processes is particularly important for improving the weather predictions. The representation of boundary layer clouds, and their further coupling to convection in stable conditions is not currently well understood. Quantification of the behavior of the PBL over the Northern Eurasian region is needed in analyses of spatial and temporal distribution of the surface fluxes, in predictions of microclimate and extreme weather events, and in modeling clouds and air quality.

The development of diagnostic and modeling methods for aero-electric structures is important for a study of both convective and electric processes in the lower troposphere (Shatalina et al., 2005; 2007). Convection in the PBL leads to the formation of aero-electric structures, manifested in ground-based measurements as short-period electric-field pulsations with periods from several to several hundreds of seconds (Anisimov et al., 1999; 2002). The sizes of such structures are determined by the characteristic variation scales of aerodynamic and electrodynamics parameters of the atmosphere, including the PBL and surface-layer

height, as well as by the inhomogeneities in the ground (water) surface. Formed as a result of convective processes and the capture of positive and negative charged particles (both ions and aerosols) by convective elements (cells), aero-electric structures move with the airflow along the Earth's surface. The further evolution of convective cells results, in particular, in cloud formation.

2.2.3.2. Global Electric Circuit

The global electric circuit (GEC) is an important factor connecting the solar activity and upper atmospheric processes with the Earth's environment, including the biosphere and climate (Dolezalek et al. 1974, Singh et al. 2004). Thunderstorm activity maintains this circuit, whose appearance is dependent on atmospheric conductance variations over a wide altitude range. The anthropogenic impact on the GEC through aviation, forest fires and electromagnetic pollution has been noted with great concern, and the importance of lightning activity in climate processes has been recognized. The GEC forms because of two reasons: the continuous operation of ionization sources, which provides an exponential growth of the conductivity in the lower atmosphere, and the continuous operation of thunderstorm generators, providing a high rate of electrical energy generation and dissipation in the troposphere. Therefore, the GEC is influenced by both geophysical and meteorological factors, and can serve as a convenient framework for the analysis of possible inter-connections between atmospheric electrical phenomena and climate processes. Further exploration of the GEC as part of the climate system studies, specifically its effect on the balance between the Earth's ionosphere and global circuit, requires accurate modeling of the GEC stationary state and its dynamics (Mareev 2010). Special attention should be paid to the observations and modeling of generators (thunderstorms, electrified shower clouds, mesoscale convective systems) in the global circuit.

2.3. Aquatic system - state of the art and future research needs

2.3.1. The Arctic Ocean in the climate system

The essential processes related to the interaction between the Arctic ocean and other components of the Earth system include the air-sea exchange of momentum, heat, and matter (e.g., moisture, aerosol, trace gases, CO₂, and CH₄), and the dynamics and thermodynamics of sea ice. The most dramatic change in the Arctic Ocean has been the rapid decline of the sea ice cover. Since the early the 1980s, the Arctic sea ice extent has decreased by roughly 50 % in summer and autumn (Cavalieri and Parkinson 2012), while the winter sea ice thickness in the central Arctic has decreased by approximately 50 % (Kwok and Rothrock 2009). Arctic sea ice changes have serious teleconnections. Despite the warming climate, wintertime cold spells in East Asia have become more frequent, stronger and longer lasting in this century compared with the 1990s (Kim et al., 2014). It also seems that the strong decline of the Arctic sea ice has favored atmospheric pressure patterns that generate cold-air outbreaks from the Arctic to East Asia (Mori et al., 2014; Kug et al., 2015; Overland et al., 2015). The reasons for and the future evolution of the sea ice decline, as well as its effects on the ocean, atmosphere and surrounding continents are among the current study topics of the Arctic climate system. Other major issues include the role of the ocean in the Arctic amplification of climate change, greenhouse gas exchange between the ocean, sea ice, and atmosphere, and aerosol budgets in the marine Arctic (Smedsrud et al., 2013). The key question here is related to the changes of sea ice extent and thickness, and to the terrestrial snow cover change.

Many of the processes considered to be responsible for the Arctic amplification of climate warming are related to the ocean and sea ice (Döscher et al., 2014). Among these, the snow/ice albedo feedback has received the most attention (e.g., Flanner et al., 2011). This feedback is strongest when sea ice is replaced by open water, but it starts to play a significant role already in spring when the snowmelt on top of sea ice begins. This is because of the large albedo difference between dry snow (albedo about 0.85) and wet, melting, bare ice (albedo

about 0.40). More work is needed to understand quantitatively the reduction of snow/ice albedo during the melting season, including the effects of melt ponds and pollutants in the snow. Other amplification mechanisms related to the ocean include increased heat transports from lower latitudes to the Arctic (Polyakov et al., 2010; Döscher et al., 2014) and fall-winter energy loss from the ocean (Screen and Simmonds, 2010). Furthermore, the melting of sea ice strongly affects evaporation, and hence the water vapor and cloud radiative feedbacks (Sedlar et al., 2011), and the PBL thickness, which controls the sensitivity of the air temperature to heat input into the PBL (Esau et al., 2012, Davy and Esau 2016). The relative importance of the mechanisms affecting the Arctic amplification of climate warming are not yet well known (See also Pithan and Mauritsen, 2014; Cohen et al., 2014).

The rapid decline of the Arctic sea ice cover has tremendous effects on navigation and exploration of natural resources. To be able to predict the future evolution of the sea ice cover, the first priority is to understand better the reasons, including the role of black carbon (see Bond et al., 2013), behind the past and ongoing sea ice evolution. Several processes have contributed to the decline of Arctic sea ice cover, but the role of these processes needs better quantification (Smedsrud et al., 2013; Vihma et al., 2014). Further studies are needed on the impacts of changes in cloud cover and radiative forcing (Kay et al., 2008), atmospheric heat transport (Kapsch et al., 2013) and oceanic heat transport (Döscher et al., 2014). In addition, as the ice thickness has decreased, the sea ice cover becomes increasingly sensitive to the ice-albedo feedback (Perovich et al., 2008). Other issues calling for more attention include the reasons for the earlier onset of the spring melt (Maksimovich and Vihma, 2012), changes in the phase of precipitation (Screen and Simmonds, 2011), and large-scale interaction between the sea ice extent, sea surface temperature distribution, and atmospheric dynamics (cyclogenesis, cyclolysis, and cyclone tracks) as discussed, e.g., by Outten et al., (2013).

In addition to thermodynamic processes, another factor affecting the sea ice cover in the Arctic is the drift of sea ice. The momentum flux from the atmosphere to the ice is the main driver of sea-ice drift, which is poorly represented in climate models (Rampal et al., 2011). This currently hinders a realistic representation of sea-ice drift patterns in large-scale climate models. Furthermore, the progressively thinning ice pack is becoming increasingly sensitive to wind forcing (Vihma et al., 2012). In the future, research has to address the main processes that determine the momentum transfer from the atmosphere to the sea ice, including the effects of atmospheric stratification and sea ice roughness.

To understand better the links between the Arctic Ocean and terrestrial Eurasia, there is a particular need to study the effects of Arctic sea ice decline on Eurasian weather and climate (Section 2.2.3). Another poorly studied problem related to the Arctic Ocean is the role of sea ice as a source of aerosol precursors, and in the gas exchange between the ocean and atmosphere (Parmentier et al., 2013). Preliminary results of field studies at the drifting stations North Pole 35 and 36 (Makshtas et al., 2011) showed that the shrinking sea ice cover could be the reason for increasing CO₂ uptake from the atmosphere over the annual cycle, and for the growth of the seasonal amplitude of CO₂ concentrations in the Arctic.

Climate models project that air temperatures and precipitation will increase over the Arctic Ocean, and that this may have important effects on the structure of sea ice. Increased snow load on a thinner ice may in the future cause flooding of seawater on ice in the Arctic, which results in the formation of snow ice. Increased snow melt and rain, on the other hand, results in increased percolation of water to the snow-ice interface, where it re-freezes, forming super-imposed ice (Cheng et al., 2008). Snow ice and super-imposed ice have granular structures, and differ thermodynamically and mechanically from the sea ice that currently prevails in the Arctic.

781 The changes in the Arctic Ocean have opened some, albeit limited, possibilities for
782 seasonal prediction. These are mostly related to the large heat capacity of the ocean: if there is
783 little sea ice in the late summer and early autumn, this tends to cause large heat and moisture
784 fluxes to the atmosphere, favoring warm, cloudy weather in late autumn and early winter (Liu
785 et al., 2012; Stroeve et al., 2012). On the other hand, the reduction of the sea ice thickness may
786 decrease the possibilities for seasonal forecasting of ice conditions in the most favorable
787 navigation season in late summer - early autumn. This is because a thin ice is very sensitive to
788 unpredictable anomalies in the atmospheric forcing. For example, in August 2012 a single
789 storm caused a reduction of the sea ice extent by approximately 1 million km². The reduced
790 sea ice extent in the winter months has significant impacts on convective clouds. Observations
791 revealed a gradually increasing frequency of the convective cloud fields over Norwegian and
792 Barents Seas (Chernokulsky and Mokhov, 2012; Esau and Chernokulsky, 2015). The
793 unusually strong atmospheric convection and weaker virtual potential temperature inversions
794 create favorable conditions for the extreme Arctic cold outbreaks and meso-scale cyclones
795 known as Polar Lows (Kolstad et al., 2009).

796 It is vital to enhance routine observations, data assimilation techniques and prediction
797 models in order to properly monitor the physical state of the environment. Longer-term
798 impacts of the reduced ice cover are largely unknown, because the scientific community has
799 had only little time to create new knowledge on essential climate variables across the domain
800 (see section 2.3.1). To improve preparedness, new observational evidence is therefore needed
801 to reduce uncertainties in the system dynamics both on short and longer time-scales.

802 2.3.2. Arctic marine ecosystem

803 The ice cover of the Arctic Ocean is undergoing fast changes, including a decline of
804 summer ice extent and ice thickness (see 2.3.1). This results in a significant increase of the
805 ice-free sea surface in the vegetation season, and an increase in the duration of the growing

season itself. The key topic of future research is the joint effect of Arctic warming, ocean freshening, pollution load, and acidification on the Arctic marine ecosystem, primary production, and carbon cycle.

New ice-free areas of the Arctic Ocean could result in a pronounced growth of the annual gross primary production (GPP), increased phytoplankton biomass and a loss of ice-rich algae communities associated with the low ice sheet surface (Bluhm et al., 2011). Progressive increase of oil and natural gas drilling and transportation over the shelf areas will be escalating the environmental changes of the Arctic marine ecosystems. Furthermore, there is a risk of irreversible changes in marine Arctic productivity and key biogeochemical cycles, and the potential for CO₂ absorption by marine ecosystem. Processes involving the Arctic may also affect adjacent boreal areas.

We do not know how the climatically-induced increase in GPP and phytoplankton biomass will influence the productivity of higher trophic levels of the Arctic ecosystem. In typical Arctic ecosystems, the most important consumers are large-sized herbivorous copepods, which have life cycles synchronized with the temperature as well as the seasonal algae dynamics (Kosobokova, 2012). Another important consumer community are the small-sized herbivorous copepods, which are important especially in shelf ecosystems. An increase in the phytoplankton production in fall, together with an increase in the sea temperature, may influence the populations of small-sized copepods, and increase their role in mass and energy flow in the ecosystems. Our current understanding of the role of small copepods in the Arctic ecosystems is limited (Arashkevich et al., 2010). An increase in surface water temperature may “open the Arctic doors” for new species, and change the Arctic pelagic food webs, energy flows, and biodiversity.

Increases in the Arctic sea temperature may lead to populations from neighboring regions penetrating the Arctic ecosystem, changing the structure and functioning of native

ecosystems. For example, a 1.5 °C water temperature increase in the Bering Sea during the mid-1970s allowed the Alaskan Pollock to penetrate the Arctic ecosystem, and occupy a place as a keystone species for several years, supporting one of the world's largest regional fish harvests (Shuntov et al., 2007). The Bering Sea ecosystem is very rich compared to the Arctic ecosystems. Currently, we are not aware of food sources sufficient for supporting massive invader populations even in case of climate-induced changes in ecosystems. However, the appearance of aggressive new species even in low numbers may dramatically impact the sensitive Arctic ecosystems and have effects on the future regulation of international fisheries in the Arctic.

We have only recently begun to understand the processes that regulate freshwater-marine ecosystem interactions in estuarine zones (Flint, 2010). The mechanisms determining the impact of riverine waters over the Arctic shelves and the central deep-basin, and their dependence on specific climatic forces, are still poorly understood. In order to determine the impact of riverine waters, it is important to locate new flagship-stations or permanent observation points in the estuaries of large Siberian rivers. The changing riverine discharge to the Arctic shelves may amplify the impact of climate warming on the Arctic marine ecosystems. Degradation of permafrost, soil erosion, changes in snow cover and summer precipitation may all lead to changes in flood timing, and also to an increase in the amount of fresh water and materials of terrestrial origin, including organic matter and nutrients, annually delivered to the Arctic shelves, and further to the Arctic basin (Gustafsson et al., 2011). Human-driven land use changes to drainage basins and associated river systems have the potential to increase the speed of delivery of pollutants to the Arctic sea.

2.3.3. Lakes, wetlands and large-scale river systems

In the last decade, the combined effects of air pollution and climate warming on freshwater systems have received increasing attention (Skjelkvåle and Wright, 1998; Schindler et

al., 2001, Alcamo et al., 2002; Sanderson et al., 2006; Feuchtmayr et al., 2009; Sereda et al., 2011). It is important to understand the future role of Arctic boreal lakes, wetlands, and large river systems, including thermokarst lakes and running waters of all size, in biogeochemical cycles, and how these changes affect livelihoods, agriculture, forestry, and industry. The water chemistry of lakes without any direct pollution sources in the catchment area can be expected to reflect regional characteristics of water chemistry, as well as global anthropogenic processes, such as climate change and long-range air pollution (Müller et al., 1998; Moiseenko et al., 2001; Battarbee et al., 2005). The current ground-based stream flow-gauging network over the Northern Eurasian region does not provide adequate spatial coverage for many scientific and water management applications, including the verification of the land-surface runoff contribution to the recipients of intra-continental runoff. Special field laboratories, with joint observation and modeling capabilities in hydrometeorology, sedimentology, and geochemistry are needed to understand the spreading of tracers and pollutants as part of current and future global environmental fluxes.

The gradient in water chemistry from the tundra to the steppe zones in Siberia can provide insight into the potential effects of climate change on water chemistry. In the last century, long-range trans-boundary air pollution led to changes in the geochemical cycles of sulfur, nitrogen, metals, and other compounds in many parts of the world (Schlesinger, 1997; Vitousek et al., 1997a,b; Kvaeven et al., 2001; Skjelkvåle et al., 2001). Environmental pollution problems include also the waterborne spreading of nutrients and pesticides from local agricultural areas, heavy metals often originating from mining areas, and other elements and chemicals, such as persistent organic pollutants from urban and industrial areas. Shifts in downstream loads cause changes in river and delta dynamics. One example of important study area is the Selenga river basin, which is located in the center of Eurasia, extends from Northern Mongolia into southern Siberia (Russia), and has its outlet at Lake Baikal. The Selenga river

basin and Lake Baikal are located in the upstream part of the Yenisei River system, which discharges into the Arctic Ocean. Lake Baikal has the largest lake volume in the world at about 23000 km³ (comprising 20 % of all unfrozen freshwater in the world), hosts a unique ecosystem (Granina, 1997), and is an important regional water resource (Garmaev and Khristoforov, 2010; Brunello et al., 2006). There are numerous industries and agricultural activities within the Selenga river basin, which affect the water quality of the lake and its tributaries. Mining is well-developed in the region (e.g., Karpoff and Roscoe, 2005; Byambaa and Todo, 2011), and heavy metals accumulate in biota and in sediments of the Selenga River delta and Lake Baikal (Boyle et al., 1998; Rudneva et al., 2005; Khazheeva et al., 2006).

In addition to water chemistry, the role of aquatic systems as a net sink or source for atmospheric CO₂ is presently under debate. When precipitation or other processes transport large volumes of organic matter from land into nearby lakes and streams, the carbon of this matter effectively disappears from the carbon budget of the terrestrial ecosystem (Huotari et al., 2011). The enhanced decomposition of soil organic matter may significantly affect the transport of terrestrial carbon to rivers, estuaries, and the coastal ocean. The contribution of this process to the global and regional carbon budgets is unknown. Thus, the biological processes taking place in the terrestrial ecosystem (e.g., photosynthesis, respiration, and decomposition) and in the aquatic ecosystem are interlinked. The observed higher temperature response of aquatic ecosystems compared to terrestrial ecosystems indicates that a substantial part of the carbon respired or emitted from the aquatic system must be of terrestrial origin (Yvon-Durocher et al., 2012). Long-term measurements carried out during all seasons in the littoral zone of Lake Baikal showed that maximum CO₂ sink and emission rates are observed in August and December (during the pre-ice period), respectively, and the total CO₂ flux from the atmosphere into the littoral zone of Lake Baikal was estimated to be 3–5 g CO₂ m⁻² (Domysheva et al., 2013).

906 The Siberian lakes situated in tundra and forest-tundra zones are in general poorly
907 studied. In their natural state, their productivity is low, but their ecosystems are highly sensitive
908 to external influences. Profuse blooming of cyanobacteria is usually associated with urban and
909 industrial effluents and nutrient run-off. An assessment is needed of the impact of climate
910 change in the northern Eurasian region on eutrophication, accompanied by blooms of
911 cyanobacteria. Besides, the northern Eurasian region is characterized by thaw lakes, which
912 comprise 90% of the lakes in the Russian permafrost zone (Romanovskii et al., 2002). These
913 lakes, which are formed in melting permafrost, have long been known to emit CH₄. The latest
914 observations of the lakes in the permafrost zone of northern Siberia indicate that they are
915 releasing much more CH₄ into the atmosphere than previously thought. Rather than being
916 emitted in a constant flow, 95 % of CH₄ comes from random bubbling in disperse locations.
917 In coming decades, this could become a more significant factor in global climate change
918 (Walter et al., 2006).

919 One direct consequence of climate change is the explosive reproduction of toxic
920 cyanobacteria (*Nodularia*, *Microcystis*, *Anabaena*, *Aphanizomenon*, *Planktothrix*) and
921 diatoms (*Pseudo-nitzschia*) (Moore et al., 2008; Paerl and Huisman, 2009). These blooms
922 occur in ponds, lakes, reservoirs, and bays of the sea. Cyanobacteria and diatoms excrete
923 especially dangerous carcinogens and neurotoxins into the water. The toxicity of some
924 cyanotoxins exceeds the toxicity of currently banned warfare agents. Antidotes to these toxins
925 do not exist at the moment.

926 Water conservation has received an increasing attention in China, and multiple new
927 projects have been initiated recently. Especially the construction of water transfer, reservoir,
928 and irrigation schemes have received much attention, because the central and western regions
929 of China are suffering from water shortages. These projects are expected to improve water
930 usage and security, especially for agricultural activities, and to provide sufficient water

resources for local societies (Mu, 2014). In China, the river systems are dominated by rivers flowing from the Tibetan plateau to the Pacific Ocean. The Yangtze is the longest river in China, and flows from the Tibetan plateau to Shanghai. The Yellow river is the second longest in China, and it is characterized by seasonal flooding which causes great economic and societal losses. The Amur River forms the northern border with Russia. The Haihe River flows through Beijing to Tianjin, and is under heavy stress from the highly populated and industrialized capital metropolitan region. Only one river from China flows to the Arctic Ocean: the Ertix River, which flows to the north through Kazakhstan, across Siberian Russia, finally joining the Ob River, which flows to the Arctic Ocean.

2.4. Social system – state of the art and future research needs

2.4.1. Land use and natural resources

The fundamental large-scale task is to estimate how human actions such as land use changes, energy production, the use of natural resources, changes in energy efficiency, and the use of renewable energy sources will influence the environments and societies of the Northern Eurasian region. For example, the industrial development of Siberia should be considered as one of most important drivers of future land use and land cover changes in Russia. Siberia is a treasure chest of natural resources of Russia, containing 85 % of its prospected gas reserves, 75 % of its coal reserves, and 65 % of its oil reserves. Siberia has more than 75 % of Russia's lignite, 95 % of its lead, approximately 90 % of its molybdenum, platinum, and platinoids, 80 % of its diamonds, 75 % of its gold, and 70 % of its nickel and copper (Korynty, 2009).

During the 20th century, a considerable transformation of landscapes in the tundra and taiga zones in northern Eurasia has occurred as a result of various industrial, socio-economic and demographic processes, leading to the industrial development of previously untouched territories (Bergen et al., 2013). This has led to a decrease in the rural population and, mostly after the 1990s, to decrease in agricultural activities. There has also been a significant

reduction in agricultural land use, and its partial replacement by zonal forest ecosystems (Lyuri et al., 2010). According to recent estimates, the total area of abandoned agricultural land in Russia in the 1990s to 2010s is at about 57 million ha, of which 18 million ha have been restored by forests and 6 million ha of this are located in Asian Russia (Schepaschenko et al., 2015). As a result, these areas have become active accumulators of atmospheric CO₂ (Kalinina et al., 2009). These new forests (“substituting resources”) could form the basis for sustainable development in these regions, in case relevant management programs for the forests re-established on abandoned lands are going to be implemented.

The dynamics of land cover, particularly forests, have been documented since 1961 when the results of the first complete inventory of Russian forests were published. According to official statistics, the area of forests in Asian Russia increased by around 80 million ha during 1961-2009, mostly before the middle of the 1990s. This large increase is explained by improved quality of forest inventories in remote territories, natural reforestation, mostly during the Soviet era as a result of forest fire suppression, and encroaching forest vegetation in previously non-forested land. Based on official statistics, the area of cultivated agricultural land in the region decreased by around 10 million ha between 1990 and 2009. After the year 2000, the forested area in Siberia decreased, mostly due to fire and the impacts of industrial transformations in high latitudes (Shvidenko and Schepaschenko, 2014). A critical decrease in the forest area has also been observed in the most populated areas with intensive forest harvesting particularly in the southern part of Siberia and the Far East. For example, in the Krasnoyarsk Krai, the total area of forests decreased by 5 %, while that of mature coniferous forests decreased by 25 %. Overall, the typical processes in these regions are a dramatic decline in the quality of forests, unsustainable use of forest resources, and insufficient governance and forest management in the region, including frequent occurrence of illegal logging, natural, and human-induced disturbances (Shvidenko et al., 2013).

982 Future land use and land cover changes will crucially depend on how successfully the
983 strategy of sustainable development of northern territories is developed and implemented. An
984 effective system for the adaptation of boreal forests to global change needs to be developed
985 and implemented in the region. An “ecologization” of the current practices of industrial
986 development of previously untouched territories would allow a substantial decrease in the
987 physical destruction of landscapes, and halt the decline of surrounding ecosystems due to air
988 pollution and water and soil contamination (Kotilainen et al., 2008).

989 The expected changes in the climate and environment will have multiple and
990 complicated impacts on ecosystems, with consequent land cover changes. The alteration of
991 fire regimes and the thawing of permafrost will intensify the process of “green desertification”
992 in large areas. Climate warming will have multiple effects on soil-vegetation-snow
993 interactions. For example, in a warmer climate, mosses and other vegetation grow faster,
994 providing a better thermal insulation of the permafrost in summer, and better feeding
995 conditions for reindeer. However, snow can also more easily accumulate on thicker vegetation,
996 thus protecting the deeper soil from cooling during the winter (Tishkov, 2012).

997 Both Russia’s north and east possess abundant mineral resources (Korytnyi, 2009). The
998 resource orientation of northern and eastern Russia’s economy, which has not changed for
999 centuries, increased in the post-Soviet period, and has been influenced primarily by the product
1000 market. It is also expected that the natural resource development sector will continue to
1001 dominate the economy in the majority of these territories for the next decades.

1002 A crucial factor in greenhouse gas emission dynamics is the fuel balance. In Russia,
1003 features of the fuel balance have led to an increased pollution. On average, specific emissions
1004 in the northern and eastern cities of Russia, where coal accounts for most of the power
1005 generation, are three times higher than in cities where power is generated mainly from gas or
1006 fuel oil (Bondur, 2011a). The geographical location, undeveloped infrastructure, harsh

climate, and coal burning are the main reasons for increased levels of anthropogenic pollution in these areas (Bondur and Vorobev, 2015; Bondur 2014). In small towns, low-capacity boiler rooms are the main source of emissions. Usually, the lack of financial resources leads to the use of low-quality coal and obsolete boilers. In the steppe zone of Asian Russia, Mongolia, Kazakhstan, and Buryatia, the main source of emissions is the burning of harvest residues.

The dynamics of GHG emissions in Russia are largely determined by the economic conditions of production. The economic crisis in 1990-1998 slowed down environmental degradation to some extent: emissions generally decreased by 40 %. However, the underlying environmental problems not only remained unresolved, but significantly deepened, and turned into systemic problems. The most polluting industries were more resistant to the decline in production. Technological degradation took place, cleaning systems were eliminated, and production shifted to part-time, leading to inefficient capacity utilization. Significant amounts of pollution continued to be emitted from the domestic sector. Emissions decreased in most regions of the country, and in 83% of the cities, but much more slowly than production. As a result, the specific emissions (per product cost at comparable prices) had grown by the end of the 1990s in all categories of cities, except cities with more than 1 million inhabitants (Bityukova et al., 2010). All this can cause negative impacts on ecosystems. For example, there are about 2 million ha of technogenic deserts around Norilsk. Norilsk is probably the biggest smelter in the world, and produces more than 2 million tons of pollutants per year (Groisman et al., 2013).

2.4.2. Natural hazards

2.4.2.1. Extreme weather and fire occurrence

The frequency and intensity of weather extremes have increased substantially during the last decades in Europe, Russia, and China. Further acceleration is expected in the future (IPCC 2013). The evolving impacts, risks, and costs of weather extremes on population, environment, transport, and industry have so far not been properly assessed in the Northern latitudes of

Eurasia. New knowledge is needed for improving the forecasting of extreme weather events, for understanding the effect of wildfires on radiative forcing and atmospheric composition in the region, for estimating the impacts of weather extremes on major biogeochemical cycles, and for understanding the effects of disturbances in forests on the emissions of BVOC and VONs (volatile organic nitrogen) (Bondur, 2011b, 2015; Bondur, Ginsburg, 2016). How do changes in the physical, chemical and biological state of the different ecosystems and the inland, water, and coastal areas affect the economies and societies in the region, and vice versa?

The number of large hydrometeorological events in Russia that cause substantial economic and social losses has increased by more a factor of two from 2001 to 2013 (State Report, 2014). The main hazards are related to atmospheric processes on various temporal and spatial scales, including strong winds, floods and landslides caused by heavy precipitation, and fires caused by drought and extreme temperatures. High temperatures and long droughts can substantially decrease the productivity and cause high dieback in dark coniferous forests. Hurricanes occur fairly often in the forest zone. For example, a hurricane destroyed about 78000 ha of forest in the Irkutsk region in July 2004 (Vaschuk and Shvidenko, 2006). However, there are no reliable statistics on many types of natural hazards.

In order to build scenarios of the future frequency and properties of weather-related hazards, one should first analyze the atmospheric mechanisms behind the circulation structures responsible for these hazards: the cyclones related to strong winds and heavy precipitation and the anticyclones related to drought and fires episodes. Studying the cyclone/anticyclone tracks, frequency and intensity can provide a statistical basis for understanding the geographical distribution and properties of the major atmospheric hazards and extremes (e.g., Shmakin and Popova, 2006). For future climate projections, atmospheric hazards and extremes should be

1057 interpreted from the viewpoint of cyclone/anticyclone statistics, and possible changes in the
1058 cyclone/anticyclone geography and frequency should be analyzed.

1059 Fires are the most important natural disturbances in the boreal forests. Fires strongly
1060 determine the structure, composition and functioning of the forest. Each year, about 0.5–1.5
1061 % of the boreal forest burns. Since boreal forests cover 15 % of the Earth's land surface, this
1062 is a significant area (Kasischke, 2000; Conard et al., 2002; Bondur, 2011b, 2015). Climate
1063 change already substantially impacts fire regimes in northern Eurasia. More frequent and
1064 severe catastrophic (mega-) fires have become a typical feature of the fire regimes. Such fires
1065 envelope areas of up to a hundred thousand hectares within large geographical regions, lead to
1066 the degradation of forest ecosystems, decrease the biodiversity, may spread to usually
1067 unburned wetlands, cause large economic losses, deteriorate life conditions and health of local
1068 populations, and lead to “green desertification”, which is an irreversible transformation of the
1069 forest cover for long periods (Shvidenko and Schepaschenko, 2013, Bondur, 2011b, 2015).
1070 Megafires also lead to specific weather conditions over the affected areas that are comparable
1071 in size to large-scale pressure systems (~30 million ha and more). The annually burned area in
1072 the Russian territory was estimated to be $8.2 \pm 0.8 \cdot 10^6$ ha during 1998-2010, and about two
1073 thirds of this area consisted of boreal forests. For this period, the fire carbon balance (total
1074 amount of carbon in the burnt fuel) was estimated to be 121 ± 28 Tg C year⁻¹ (Shvidenko et al.,
1075 2011). Current model projections suggest that the number of fires will double by the end of
1076 this century. The extent of catastrophic fires escaping from the control and fire intensity are
1077 projected to increase. Due to increased severity of fire and deeper soil, carbon emissions from
1078 fires are predicted to increase by a factor of 2 to 4 (Gromtsev, 2002; Malevsky-Malevich et
1079 al., 2008; Flanningan et al., 2009; Shvidenko et al., 2011). During and after fires, significant
1080 changes take place in the forest ecosystems, including the soil. These changes include: (i) a
1081 significant amount of biomass is combusted, and large amounts of carbon and nitrogen are

released to the atmosphere in the form of carbon dioxide and other gases or particles (Harden et al., 2000; Andreae and Merlet, 2001; Kaiser et al., 2012; Konovalov et al., 2014; Kulmala L. et al., 2014); (ii) fire alters the microbial community structure in the soil, as well as the structure of the vegetation (Dooley and Treseder, 2012; Sun et al., 2015); (iii) fires determine the structure of the vegetation, succession dynamics and the fragmentation of forest cover, tree species composition, and the productivity of boreal forests (Gewehr et al., 2014) and (iv) fire is one of the crucial drivers controlling the dynamics of the carbon stock of boreal forests (Jonsson and Wardle, 2010; Köster et al., 2014).

Disturbances resulting from fire, pest outbreaks and diseases also have substantial effects on the emissions of BVOCs and volatile organic nitrogen compounds (Isidorov, 2001), and consequently on atmospheric aerosol formation. The acceleration of fire regimes will also affect the amount of black carbon in the atmosphere, and thus has an effect on the albedo of the cryosphere.

2.4.2.2. Permafrost degradation and infrastructures

The degradation of permafrost will cause serious damage both to infrastructure and to ecosystems and water systems in the Northern Eurasian region. This includes, for example, damage to pipelines and buildings, deformation of roads and railroads in Russia, Mongolia and China, variations in the ion distribution in soil water in young and ancient landslides, cryogenic landslides, spatial and temporal changes of grass and willow vegetation, saline water accumulation in local depressions of the permafrost table, and formation of highly saline lenses of ground water called ‘salt traps’.

Due to the large extent of permafrost-covered areas in northern Eurasia (for ecosystem effects, see sections 2.1.1 and 2.1.2), there are numerous infrastructural issues related to possible changes in the thickness and temperature of the frozen part of the subsurface, and thus in the mechanical soil properties. Climate change -induced changes in the cryosphere are probably among the most dramatic issues affecting the infrastructure in northern Eurasia, as

1108 this infrastructure is literally standing on permafrost. Moreover, an interesting coupling may
1109 be related to the decreasing ice-cover of the Arctic Ocean, which results in increased humidity
1110 and precipitation on the continent, and thus a further thickening and longer duration of the
1111 annual snow cover. Snow is a good thermal insulator, and influences the average ground
1112 surface temperature, thus playing a potentially important role in speeding up the thawing of
1113 permafrost.

1114 The increased risk of damage to local infrastructure, such as buildings and roads, can
1115 cause significant social problems, and exerts pressure on the local economies. Thawing
1116 permafrost is structurally weak, and places a variety of infrastructure at risk. For example, the
1117 failure of buildings, roads, pipelines, or railways can have dramatic environmental
1118 consequences, as seen in the 1994 breakdown of the pipeline to the Vozei oilfield in northern
1119 Russia, which resulted in a spill of 160,000 tons of oil - the world's largest terrestrial oil spill
1120 (United Nations Environment Program, 2013). Maintenance and repair costs related to
1121 permafrost thaw and degradation of infrastructure in northern Eurasia have recently increased,
1122 and will most probably increase further in the future. This is an especially prominent problem
1123 in discontinuous permafrost regions, where even small changes in the permafrost temperature
1124 can cause significant damage to infrastructure. Most settlements in permafrost zones are
1125 located on the coast, where strong erosion places structures and roads at risk. After damage to
1126 the infrastructure, local residents and indigenous communities are often forced to relocate.
1127 This can cause changes in, or even disappearances of, local societies, cultures, and traditions
1128 (United Nations Environment Program, 2013).

1129 2.4.2.3. Changing sea environments and the risk of accidents in coastal regions

1130 In northern Eurasia, from the eastern part of the Barents Sea to the Bering Sea, the
1131 permafrost is located directly on the seacoast. In many of these coastal permafrost areas, sea
1132 level rise and continuing permafrost degradation leads to significant coastal erosion, and to
1133 the possibility of a collapse of coastal constructions such as lighthouses, ports, houses, etc. In

1134 this region, the sea level rise is coupled to the permafrost degradation in a complex way, and
1135 should be focused on in future studies.

1136 Understanding and measuring artificial radionuclides in marine ecosystems is needed
1137 for improving emergency preparedness capabilities, and for developing risk assessments of
1138 potential nuclear accidents. The awareness of the general public and associated stakeholders
1139 across the region should also be raised concerning the challenges and risks associated with
1140 nuclear technologies, environmental radioactivity, and emergency preparedness. The current
1141 state of radioactive contamination in terrestrial and marine ecosystems in the European Arctic
1142 region will be studied by examining environmental samples collected from Finnish Lapland,
1143 Finnmark, and Troms in Norway, the Kola Peninsula, and the Barents Sea. The results will
1144 provide updated information on the present levels, occurrence and fate of radioactive
1145 substances in the Arctic environments and food chains. The results will also allow us to
1146 estimate where the radioactive substances originate from, and what risks they may pose in case
1147 of accidents.

1148 Annual expeditions for sample collection are needed for the development of models to
1149 predict the distribution of radionuclides in the northern marine environment, and for the
1150 assessment of the current state of radioactive contamination in marine ecosystems in the
1151 European Arctic region. In view of recent developments and increased interests in the
1152 European Arctic region for oil and gas extraction, special attention needs to be given to the
1153 analysis of norms (naturally occurring radioactive materials) in order to understand current
1154 levels. The future focus should be put on atmospheric modeling, and on the assessment of
1155 radionuclide distributions in the case of accidents leading to the release of radioactive
1156 substances to the environment in the European Arctic region. This includes the assessment of
1157 nuclear accident scenarios for dispersion modeling.

2.4.3. Social transformations

Climate and weather strongly affect the living conditions, mostly in the Eastern part of the Northern Eurasian societies, influencing people's health, incidence of diseases and adaptive capacity. The vulnerability of societies, including their adaptive capacity, varies greatly depending on both their physical environment, and on their demographic structure and economic activities. There is a need to analyze the scientific background and robustness of the adaptation and mitigation strategies (AMS) of the region's societies, and their resilience capacity, with special emphasis on the forest sector and agriculture. The future research needs are in understanding in what ways populated areas are vulnerable to climate change; how their vulnerability can be reduced and their adaptive capacities improved; what responses should be identified to mitigate and adapt to climate changes.

Health issues are also important in multidisciplinary studies of northern Eurasia, as the living conditions of both humans and livestock are changing dramatically. Short-lived climate forcers (SLCF), such as black carbon, ozone, and aerosol particles, are important players in both air quality and Arctic climate change and their impacts are not yet quantified. Black carbon has a special role when designing future emission control strategies, since it is the only major aerosol component whose reduction is likely to be beneficial to both climate and human health. These changes can be expressed through complex parameters combining the direct effects of, e.g., temperature and wind speed, with indirect effects of several climatic and non-climatic factors such as the atmospheric pressure variability, or the frequency of unfavorable weather events, such as heat waves or strong winds. During the last decades, living conditions in northern Eurasia have generally improved, but with a significant regional and seasonal variation (Zolotokrylin et al., 2012).

Both northern and eastern Eurasia have small and diminishing populations, mainly due to the migration outflow started in the 1990s due to severe and unfavorable living conditions combined with changing state policies with respect to the development of the northern

territories. This reversed the previous long-standing pattern of migration inflow. The combination of outflow and natural population decrease (with some regional exceptions in several ethnic republics and autonomous regions (*okrugs*) with oil and gas industry) led to a steady population decline in most regions in northern and eastern Russia from 1990s. In the post-soviet period, the population of eastern Russia decreased by 2.7 million, while the population of Russia's Arctic zone decreased by nearly by one third (500 000 people), in contrast to the majority of the world's Arctic territories (Glezer, 2007a, 2007b). The population change in northeastern Russia was particularly remarkable: the Chukotka autonomous okrug lost 68 % of its population, the Magadan Oblast lost 59 %, and the Kamchatka Krai lost 33 %.

Geographical and ethnic factors influence the demography and settlement pattern in the region. Geographical factors include environmental conditions and the mixture of urban and rural territories. Areas with a large proportion of indigenous people employed in traditional nature management were exposed to relative small post-soviet transformations in the 1990's and 2000's. In contrast, the largest transformations occurred in areas with a larger proportion of Russian people and developed mining industries. The differences in the transformations between settlements with predominantly indigenous and predominantly Russian populations are evident. For example, in the Chukotka Autonomous Okrug, the former remained mostly intact, with only small decreases in population, while the latter disappeared entirely or were significantly depopulated (Litvinenko, 2012; 2013).

When assessing the impacts of climate change and other environmental changes on human societies, it should be taken into account that the urban environments in northern Eurasian cities and towns situated in the less favored regions are currently incapable of mitigating unfavorable impacts. The impact of climate parameters, such as temperature (including seasonal, weekly, and daily cycles, and extreme values), strong winds, snowfall, snowstorms, and precipitation should be investigated. Both the frequency and the duration of

weather events should be considered. These climate parameters influence human health, incidence of diseases, adaptation potential, and economic development in general. Furthermore, it is important to explore the interactions between the environmental change and post-soviet transformations of natural resource utilization in northern Eurasia in order to assess the complexity of their socio-ecological consequences at regional and local levels (Litvinenko, 2012; Tynkkynen, 2010). The population dynamics of the northern Russian regions in 1990-2012, and the linkage between intra-regional differences in population dynamics, spatial transformations of natural resources utilization, and ethnic composition of the populations should be clarified. It would be desirable to develop an “early warning system” for the timely mitigation of the negative socio-ecological effects of both environmental changes, and changes in the availability of natural resources as well as accident like leakages in gas and oil pipelines. Such systems would be useful for federal, regional, and local authorities, as well as for local communities.

It should also be taken into account that the majority of the world’s ethnic groups are small and engaged in culturally specialized methods of subsistence, so any change in their immediate environment may lead to their traditional way of life becoming unsustainable. These changes may be due to rising sea levels, warming seawater, melting ice cover, thawing permafrost, flooding rivers, changing rain patterns, or moving vegetational zones. These are direct effects of climate change and environmental deterioration on ethnodiversity. However, even more threatening are the indirect effects. The immediate environment of small ethnic groups is often vulnerable to the adverse impact of majority populations representing governments and nations. The effects of climate change may lead to a rapid and massive transfer of majority populations to areas previously inhabited by small ethnic groups.

1232 3. From process studies towards system understanding and quantification of feedbacks of
1233 Arctic-boreal regions

1234 The system understanding helps us to understand the behavior of feedbacks between the
1235 land, atmosphere, aquatic, and societal/economic systems. To be able to provide a system
1236 understanding, we need to understand the individual processes, and based on process
1237 understanding we are then able to quantify different biogeochemical cycles. Via
1238 biogeochemical cycles, the energy and matter flows are linked to a wider system context,
1239 which enables us to analyze the feedback phenomena. Feedbacks are essential components of
1240 our climate system, as they either increase or decrease the changes in climate-related
1241 parameters in the presence of external forcings (IPCC, 2013).

1242 The effects of climate change on biogeochemical cycles are still inadequately
1243 understood, and many feedback mechanisms are difficult to quantify (Arneth et al., 2010;
1244 Kulmala et al., 2014). They are related to, for example, the coupling of carbon and nitrogen
1245 cycles, permafrost processes and ozone phytotoxicity (Arneth et al., 2010), or to the emissions
1246 and atmospheric chemistry of biogenic volatile organic compounds (Grote and Niinemets,
1247 2008; Mauldin et al., 2012), subsequent aerosol formation processes (Kulmala et al., 2004;
1248 Tunved et al., 2006; Kulmala et al., 2011a; Hirsikko et al., 2011) and aerosol-cloud interactions
1249 (McComiskey and Feingold, 2012; Penner et al., 2012; Rosenfeld et al., 2014).

1250 The northern Eurasian Arctic-boreal geographical region covers a wide range of
1251 interactions and feedback processes between humans and natural systems. Humans are acting
1252 both as the source of climate and environmental changes, and as recipient of their impacts. The
1253 PEEX research agenda is addressing the most relevant research topics related to the process
1254 dynamics in the land, atmospheric, aquatic, and society systems relevant to Northern regions.
1255 PEEX also aims to quantify the range of emissions and fluxes from different types of
1256 ecosystems and environments and links to ecosystem productivity (see also Su et al., 2011;
1257 Kulmala and Petäjä, 2011; Bäck et al., 2010). This new knowledge helps us to obtain a holistic

view on the changes in biogeochemical cycles and feedbacks in the future Arctic-boreal system (Fig 4). PEEX will also take into consideration that there may exist previously unknown sources and processes (Su et al., 2011; Kulmala and Petäjä, 2011; Bäck et al., 2010).

Holistic representations of feedback loops potentially relevant to Arctic-boreal systems have been given by Charlson et al., (1987), Quinn and Bates (2011), and by Kulmala et al., (2004; 2014). The “CLAW” hypothesis (“CLAW” acronym refers to Charlson, Lovelock, Andreae and Warren) connects the ocean biochemistry and climate via a negative feedback loop involving cloud condensation nuclei production due to dimethylsulfoniopropionate (DMSP) and DMS biosynthesis by marine phytoplankton (e.g., Quinn and Bates, 2011; Ducklow et al., 2001; O’Dowd et al., 2004; de Leeuw et al., 2011; Malin et al., 1993; O’Dowd and de Leeuw, 2007). The COBACC (CONTinental Biosphere-Aerosol-Cloud-Climate) hypothesis suggests two partly overlapping feedbacks that connect the atmospheric carbon dioxide concentration, ambient temperature, gross primary production, biogenic secondary organic aerosol formation, clouds, and radiative transfer (Kulmala et al., 2004; 2014; also see section 2.1.1.). The quantification of these feedback loops under changing climate is crucial for reliable Earth system modelling and predictions.

In the context of the COBACC feedback loop, the key large-scale research questions are the changing cryospheric conditions and consequent changes in ecosystem feedbacks affecting the Arctic-boreal climate system and weather. Furthermore, we should estimate the net effects of various feedback effects (CLAW, COBACC) on land cover changes, photosynthetic activity, GHG exchanges, BVOC emissions, aerosol and cloud formation, and radiative forcing at regional and global scales. In our analysis, we should also take in account the urbanization processes and social transformations (see section 2.4.3), which are changing the regional climates. In this task, we should also study the key gaps of the biogeochemical cycles.

3.1. Hydrological cycle

Climate change may profoundly affect most of the components of the hydrological cycle, giving rise to positive or negative feedbacks (Fig 5). While variations in the hydrological cycle often take place at regional or local scales, they can also give rise to large-scale or even global changes. Knowledge of the hydrological cycle in general and particularly related to permafrost is crucial for predicting the resilience and transformation of forest ecosystems coupled with permafrost (Osawa et al., 2009).

In addition to permafrost processes, another important issue in high latitudes is precipitation. Precipitation is a critical component of the hydrological cycle, having a great spatial and temporal variability. The lack of understanding of some precipitation-related processes, combined with the lack of global measurements of sufficient detail and accuracy, limits the quantification of different components of the hydrological cycle like precipitation, evapotranspiration, CCN formation etc. This is especially true in high-latitude regions, where observations and measurements are particularly sparse, and processes poorly understood.

Recent retrievals of multiple satellite products for each component of the terrestrial water cycle provide an opportunity to estimate the water budget globally (Sahoo et al., 2011) (Fig. 5). Global precipitation is retrieved at very high spatial and temporal resolution by combining microwave and infrared satellite measurements (Sorooshian et al., 2000; Kummerow et al., 2001; Joyce et al., 2004; Huffman et al., 2007). Large-scale estimates of global precipitation have been derived by applying energy balance, process, and empirical models to satellite-derived surface radiation, meteorology, and vegetation characteristics (e.g., Mu et al., 2007; Su et al., 2007; Fisher et al., 2008; Sheffield et al., 2010). The water storage change component can be obtained from satellite data, and the water level in lakes and large-scale river systems can be estimated from satellite altimetry with special algorithms developed for terrestrial waters (Berry et al., 2005; Velicogna et al., 2012; Troitskaya et al., 2012; 2013).

3.2. Carbon cycle

It is not clear how future climate will modify incoming terrestrial net primary production (NPP) and outgoing (e.g., heterotrophic soil respiration (HSR)) carbon fluxes to and from terrestrial ecosystems. It is likely that the transformation of Russian forests is a tipping element for the climate system by end of the century over huge areas, even though uncertainties in such forecasts are significant (Gauthier et al., 2015). The role of boreal and Arctic lakes and catchment areas in carbon storage dynamics is poorly quantified (Fig. 6).

The terrestrial biosphere is a key regulator of atmospheric chemistry and climate via its carbon uptake capacity (Arneeth et al., 2010; Heimann and Reichstein, 2008). The Eurasian area holds a large pool of organic carbon both within the above- and below-ground living biota, in the soil, and in frozen ground, stored during the Holocene and the last ice age. The area also contains vast stores of fossil carbon. According to estimates of carbon fluxes and stocks in Russia made as part of a full carbon account by the land-ecosystem approach (Shvidenko et al., 2010a; Schepaschenko et al., 2011; Dolman et al., 2012), terrestrial ecosystems in Russia served as a net carbon sink of 0.5-0.7 Pg(C) per year during the last decade. Forests provided above 90 % of this sink. The spatial distribution of the carbon budget shows considerable variation, and substantial areas, particularly in permafrost regions and in disturbed forests, display both sink and source behavior. The already clearly observable greening of the Arctic is going to have large consequences on the carbon sink in the upcoming decades (Myneni et al., 1997; Zhou et al., 2001), although future predictions are uncertain. The Net Ecosystem Carbon Budget (NECB) or Net Biome Production (NBP) are usually a sensitive balance between carbon uptake through forest growth, ecosystem heterotrophic respiration, and carbon release during and after disturbances such as fire, insect outbreaks, or weather events such as exceptionally warm autumns (Piao et al., 2008; Vesala et al., 2010). This balance is delicate, and for example in the Canadian boreal forest the estimated net carbon

balance is close to carbon neutral due to fires, insects, and harvesting cancelling the carbon uptake from forest net primary production (Kurz and Apps, 1995; Kurz et al., 2008).). Long-term measurements of the concentrations of CO₂ and other carbon gases at selected sites, especially using tall towers such as the ZOTTO tower, will be essential for constraining the large-scale carbon fluxes in the PEEX region (Heimann et al., 2014).

Plant growth and carbon allocation in boreal forest ecosystems depend critically on the supply of recycled nutrients within the forest ecosystem. In the nitrogen-limited boreal and Arctic ecosystems, the biologically available nitrogen (NH₄ and NO₃) is in short supply, although the flux of assimilated carbon below ground may stimulate the decomposition of nitrogen-containing soil organic matter (SOM), and the nitrogen uptake of trees (Drake et al., 2011; Phillips et al., 2011). The changes in easily decomposable carbon could enhance the decomposition of old SOM (Kuzyakov, 2010; Karhu et al., 2014), and thus increase the turnover rates of nitrogen in the rhizosphere, with possible growth-enhancing feedbacks on vegetation (Phillips et al., 2011).

Arctic warming is promoting terrestrial permafrost thaw and shifting hydrologic flow paths, leading to fluvial mobilization of ancient carbon stores (Karthé et al., 2014). Observed permafrost thaw acts as a significant and preferentially degradable source of bioavailable carbon in Arctic freshwaters, which is likely to increase as permafrost thaw intensifies, causing positive climate feedbacks in response to on-going climate change (Mann et al., 2015). Significant differences in fluvial carbon input between headwaters and downstream reaches of large Arctic catchments (Yenisey and Lena) have been identified, but the problem is very poorly explained yet. At the same time, the fluvial export by the largest rivers is considered to be an order of magnitude less than coastal erosion in the East Siberian Arctic Shelf (Semiletov et al., 2011). The Lena's particulate organic carbon export is estimated to be two orders of

magnitude less than the annual input of eroded terrestrial carbon onto the shelf of the Laptev and East Siberian seas.

Although inland waters are especially important as lateral transporters of carbon, their direct carbon exchange with the atmosphere, so-called outgassing, has been recognized to be a significant component in the global carbon budget (Bastviken *et al.*, 2011; Regnier *et al.*, 2013). In the boreal pristine regions, forested catchment lakes can vent *ca.* 10 % of the terrestrial NEE (Net Ecosystem Exchange), thus weakening the terrestrial carbon sink (Huotari *et al.*, 2011). There is a negative relationship between the lake size and gas saturation, and especially small lakes are relatively large sources of CO₂ and CH₄ (e.g., Kortelainen *et al.*, 2006; Vesala, 2012). However, on a landscape level, large lakes can still dominate the GHG fluxes. Small lakes also store relatively larger amounts of carbon in their sediments than larger lakes. The role of lakes as long-term sinks of carbon, and simultaneously as clear emitters of carbon-containing gases, is strongly affected by the physics of the water column. In lakes with very stable water columns and anoxic hypolimnion sediments, carbon storage is especially efficient, but at the same time, these types of lakes emit CH₄. In general, the closure of landscape-level carbon balances is virtually impossible without studying the lateral carbon transfer processes (Pumpanen *et al.*, 2014), and the role of lacustrine ecosystems as GHG sources/sinks. Besides lakes, these studies should include rivers and streams, which could be even more important than lakes as transport routes of terrestrial carbon and as emitters of GHGs (Huotari *et al.*, 2013). In addition, the role of VOC emissions as a part of the carbon budget needs to be quantified.

3.3. Nitrogen cycle

Nitrogen is the most abundant element in the atmosphere. However, most of the atmospheric nitrogen is in the form of inert N₂, which is unavailable most for plants and microbes, and can only be assimilated into terrestrial ecosystems through biological N₂

fixation (Canfield et al., 2010). Only cryptogamic covers and certain organisms living in symbiosis with plants are capable of nitrogen fixation, making nitrogen the main growth-limiting nutrient in terrestrial ecosystems (Elbert et al., 2012; Lenhart et al., 2015). Human perturbations to the natural nitrogen cycle have, however, significantly increased the availability of nitrogen in the environment (Fig. 7). These perturbations mainly stem from the use of fertilizers in order to increase crop production to meet the demands of the growing population (European Nitrogen Assessment, 2010), although atmospheric nitrogen deposition may also play a significant role in some areas. The increased use of fertilizer nitrogen, and consequent perturbations in nitrogen cycling, also cause severe environmental problems such as eutrophication of terrestrial and aquatic ecosystems, atmospheric pollution, and ground water deterioration (European Nitrogen Assessment, 2010).

Emission of reactive nitrogen (NO, NO₂, HONO, ammonia, amines) from soils (Su *et al.*, 2011; Korhonen *et al.*, 2013), fossil fuel burning, and other sources links the nitrogen cycle to atmospheric chemistry and secondary aerosol formation in the atmosphere. There are indications that emissions of N₂O from the melting permafrost regions in the Arctic may significantly influence the global N₂O budget and hence contribute to the positive radiative forcing by greenhouse gases (Repo et al., 2009; Elberling et al., 2011).

In natural terrestrial ecosystems, nitrogen availability limits ecosystem productivity, linking the carbon and nitrogen cycles closely together (Gruber and Galloway, 2008). The increasing temperatures due to climatic warming accelerate nitrogen mineralization in soils, leading to increased nitrogen availability and transport of reactive nitrogen from terrestrial to aquatic ecosystems. This perturbed and accelerated nitrogen cycling may lead to large net increases in the carbon sequestration of ecosystems (Magnani et al., 2007). The large surface area of boreal and Arctic ecosystems implies that even small changes in nitrogen cycling or feedbacks to the carbon cycle may be important on the global scale (Erisman et al., 2011). For

instance, increased atmospheric nitrogen deposition has led to higher carbon sequestration in boreal forests (Magnani et al., 2007). However, the feedback mechanisms from increased perturbations of the nitrogen cycle may change the dynamics of the emissions of other greenhouse gases, hence complicating the overall effects. For instance, the stimulated carbon uptake of forests due to increased atmospheric nitrogen deposition can largely be offset by the simultaneously increased soil N₂O emissions (Zaehle et al., 2011). In the Arctic, the melting permafrost may lead to high emissions of N₂O (Repo et al., 2009; Elberling et al., 2010), which may significantly influence the global N₂O budget.

Understanding the processes within the nitrogen cycle, the interactions of reactive nitrogen with the carbon and phosphorus cycles, atmospheric chemistry and aerosols, as well as their links and feedback mechanisms is therefore essential in order to fully understand how the biosphere affects the atmosphere and the global climate (Kulmala and Petäjä, 2011).

3.4. Phosphorus cycle

Phosphorus (P) is, together with nitrogen (N), one of the limiting nutrients for terrestrial ecosystem productivity and growth, while in marine ecosystems, phosphorus is the main limiting nutrient for productivity (Whitehead and Crossmann, 2012). The role of P in nutrient limitation in natural terrestrial ecosystems has not been recognized as widely as that of N (Vitousek et al., 2010).

In the global phosphorus biogeochemical cycle, the main reservoirs are in continental soils, where phosphorus in mineral form is bound to soil parent material, and in ocean sediments (Fig.8). Sedimentary phosphorus originates from riverine transported material eroded from continental soils. The atmosphere plays a minor role in the phosphorus cycle, and the phosphorus cycle does not have a significant atmospheric reservoir. Atmospheric phosphorus mainly originates from aeolian dust, sea spray, and combustion (Wang et al.,

2014). Gaseous forms of phosphorus are scarce, and their importance for atmospheric processes is unknown (Glindemann et al., 2005).

Southwestern Siberian soils have lately been reported to contain high concentrations of plant-available phosphorus (Achat et al., 2013), which may enhance the carbon sequestration of the ecosystems if they are not too limited by nitrogen. In soils, phosphorus is found mainly in mineral form and bound to the soil parent material such as apatite minerals. The amount of phosphorus in the parent material is a defining factor for phosphorus limitation, and the weathering rate determines the amount of phosphorus available for ecosystems. In ecosystems, most of the available phosphorus is in organic forms (Achat et al., 2013; Vitousek et al., 2010). In ecosystems growing on phosphorus-depleted soils, the productivity is more likely to be nitrogen-limited in early successional stages, and gradually shift towards phosphorus limitation as the age of the site increases (Vitousek et al., 2010). In freshwater ecosystems, excess phosphorus leads to eutrophication, which has ecological consequences, such as the loss of biodiversity due to changes in physicochemical properties and in species composition (Conley et al., 2009). Due to the scarcity of studies focusing on ecosystem phosphorus cycling, the effects of climate change on physicochemical soil properties and phosphorus availability, and the interactions of the phosphorus cycle with the cycles of carbon and nitrogen, are largely unknown.

3.5. Sulfur cycle

Sulfur is released naturally through volcanic activity, as well as through weathering of the Earth's crust. The largest natural atmospheric sulfur source is the emission of dimethyl sulfide (DMS) from oceanic phytoplankton (Andreae, 1990). DMS is converted to sulfur dioxide (SO₂), sulfuric acid (H₂SO₄) and methyl sulfonic acid (MSA) via gas-phase oxidation. However, human activities have a major effect on the global sulfur cycle via vast emissions of SO₂ from fossil fuel burning and smelting activities. The main sink of SO₂ is oxidation to

sulfuric acid in both gas and liquid phases, and subsequent removal from the atmosphere via precipitation and dry deposition.

Global anthropogenic SO₂ emissions are predicted to decrease significantly by the year 2100 (IPCC, special report on emissions scenarios, SRES, 2000). Emissions in Europe and North America started to decrease already in the 1970s, but this decrease is still overwhelmed on a global scale by increasing emissions in eastern Asia and other strongly developing regions of the world (Smith et al., 2011). The current global anthropogenic SO₂ emissions are about 120 Tg per year, with Europe, the former Soviet Union and China together responsible for approximately 50 % (Smith et al., 2011). Global natural emissions of sulfur, including DMS, are significantly smaller (a few tens of Tg per year; Smith et al., 2001). Anthropogenic emissions dominate especially over the continents. The main sources of SO₂ are coal and petroleum combustion, metal smelting, and shipping, with minor contributions from biomass burning and other activities.

SO₂ emissions in Eurasia have a large spatial variability. Smelters in the Russian Arctic areas emit vast amounts of SO₂, significantly affecting the regional environment. In 2007 Blacksmith Institute experts estimated, that the smelter complexes in Norilsk, with annual emissions of 2 Tg are alone responsible for more than 1.5 % of global SO₂ emissions. However, the emissions from the smelters in the Kola Peninsula, while still remarkably high, have decreased significantly during the past decades (Paatero et al., 2008), thus altering the impact of human activities on the regional climate and environment. In general, existing anthropogenic activities are slowly becoming more sulfur-effective and less polluting. However, the emergence of new sulfur-emitting activities and infrastructures partially counteracts this development.

The behavior of future changes in SO₂ emissions in the PEEX research area is uncertain. In northern Eurasia, natural resources like fossil fuels, metals, minerals, and wood are vast,

and their utilization is becoming more and more attractive due increasing demand. This will most likely lead to an increase in human activities (e.g., mining, oil drilling, shipping) in this area (e.g., Smith, 2010, and references therein). For example, sulfur emissions in China increased rapidly until 2006, and then decreased by 9.2 % to 30.8 Tg in 2010 due to the wide application of flue-gas desulfurization (FGD) equipment in power plants (Lu and Zhang 2011). Sulfur emissions in Europe have decreased significantly during the last decades (Jones and Harrison 2011).

Most of the natural and anthropogenic SO_2 is removed from the atmosphere by liquid-phase oxidation to H_2SO_4 , and subsequent precipitation. In areas with high sulfur loadings, acid rain leads to acidification of soils and waters (Fig. 9). The main final sink of sulfur is the oceans. A fraction of SO_2 is oxidized to H_2SO_4 in the gas phase in a reaction chain initiated by the reaction of SO_2 with the hydroxyl radical, OH. Especially in forested areas of Eurasia, reactions of SO_2 with a second important oxidant type, the stabilized Criegee intermediates originating from biogenic VOC emissions, also produce significant amounts of H_2SO_4 (Mauldin et al., 2012). Gas-phase sulfuric acid plays a key role in the Earth's atmosphere by triggering secondary aerosol formation, thus connecting anthropogenic SO_2 emissions to global climate via aerosol-cloud interactions. Particles containing sulfuric acid, or sulfate, are also connected with air quality problems and human health deterioration. Understanding the spatial and temporal evolution of SO_2 emissions in northern Eurasia, along with atmospheric sulfur chemistry, is crucial for understanding and quantifying the impacts of anthropogenic activities and SO_2 emissions on air quality, acidification, as well as on regional and global climate.

4. From system understanding to mitigation and adaptation strategies and decision making

Climate change and weather extremes are already affecting the living conditions of Northern Eurasian societies. The vulnerability of the Northern environments and societies, including their adaptive capacity and buffering thresholds, varies greatly depending on their current and future physical environment as well as their demographic structure and economic activities. The PEEX program as a whole is built on four pillars, namely (i) research, (ii) research infrastructure, (iii) impact on society, and (iv) knowledge transfer and capacity building. The scientific outcome of the first two pillars will be addressing the future state of the physical environment and its interactions and feedbacks with the demographic structure and economic activities in the Arctic boreal system. Periodic PEEX assessments will be delivered for constructing mitigation and adaptation strategies of the Northern societies and for use of regional and governmental decision-making. The PEEX approach is applicable to China, when taking into account the specific geographical, climatological, and social characteristics of that region.

The integrative approach of the PEEX first two pillars provides both analytical and operational answers to our research questions, which can be utilized in solving interlinked grand challenges using pillars iii) and iv). These will also contribute to the Earth System sciences (ESS) questions as a whole (see ESS questions: Schellnhuber et al., 2004). The implementation of the PEEX research agenda starts with process studies in the frame of three main topics determined for the land, atmosphere, aquatic, and social systems of the Northern Eurasian region. The research approach is designed to answer the analytical questions on the major dynamical patterns and feedback loops relevant to Earth system science in the Northern context. The PEEX program has defined altogether 12 large-scale research questions for the 12 main topics in the Northern Eurasian domain (Kulmala et al., 2016). At the same time, PEEX adheres to several operational ESS questions, including “what level of complexity and resolution have to be achieved in Earth System modelling?”, “what are the best techniques for

analyzing and predicting irregular events?”, “what might be the most effective global strategy for generating, processing, and integrating relevant Earth system datasets?”, and “what are the most appropriate methodologies for integrating natural science and social science knowledge?” (Schellnhuber et al., 2004).

In terms of the level of complexity and resolution in Earth System modelling, PEEEX builds on a multi-scale modelling and observation approach originally introduced by Kulmala et al., (2009). PEEEX will construct its own multi-scale modelling platform (Lappalainen et al., 2014). In terms of generating, processing and integrating relevant Earth system datasets, a detailed conceptual design of the PEEEX research infrastructure (RI) will include a concept design of a coherent in-situ observation network, coordinated use of remote sensing observations and standardized and harmonized data procedures, as well as a data system. One of the first tasks of PEEEX-RI is to fill in the observational gap in atmospheric in-situ and ground base remote sensing data in the Northern Eurasia, especially in Siberia. This approach is based on the coordination of existing observation activities (Alekseychik et al., 2016), but also making plans for a new infrastructure needed. PEEEX-RI development will be largely based on the SMEAR (Station for Measuring Ecosystem-Atmosphere Relations) concept (Kulmala et al., 2016), which has been developed by the University of Helsinki Division of Atmospheric Sciences together with Division of Forest Ecology starting from 1995 (Hari and Kulmala, 2005; Hari et al., 2016). The SMEAR concept provides a state-of-the-art foundation for establishing a PEEEX observation system to be integrated into the global GEOSS data system. Furthermore, detailed design of greenhouse gas, aerosol, cloud, and trace gas measurements, and observation of biological activity will find synergies with the major European land-atmosphere observation infrastructures, such as ICOS (a research infrastructure to determine the greenhouse gas balance of Europe and adjacent regions), ACTRIS (aerosols,

clouds, and trace gases research infrastructure), GAW (Global Atmospheric Watch), and AnaEE (the experimentation in terrestrial ecosystem research).

PEEX is interested in developing methodologies for integrating natural science and social science knowledge as part of the operational Earth sustainable system questions (Schellnhuber et al., 2004). The first-priority task in this case is to establish an integrated geographical information background (Ribeiro et al., 2009; Hunt and Sanchez-Rodriguez, 2009; Shvidenko et al., 2010; Skryzhevskaya et al., 2015). A common information background would be the first step serving the development of a common language of integrated studies. For example, we need spatially and temporally explicit descriptions of terrestrial ecosystems, landscapes, atmosphere, and hydrosphere. A common information background would be a unified base for the PEEX modelling platform and for the development of integrated modelling clusters, which could combine ecological, economic, and social dimensions. It could provide a historical background for the future trajectories of land cover, state and resilience of ecosystems, stability of landscapes, and dynamics of environmental indicators of environment. The already existing Integrated Land Information System could be utilized here for combining all historical knowledge about the region and all scientific results obtained by past, current, and future studies across the region (e.g., Schepaschenko et al., 2011; Shvidenko and Schepaschenko, 2014).

In addition to data services, PEEX is developing procedures for integrating and linking natural science and social science knowledge and data. As one example, we need to analyze data on emission sources together with population health risk factors, environment pollution, food security, drinking water quality, changes in the spreading areas of infectious diseases, and changes in the general epidemiological situation (Bityukova and Kasimov, 2012; Malkhazova et al., 2013). Via novel multidisciplinary data interfaces and data procedures, we are able to connect satellite observations with inverse modeling, provide fast updates to

emission inventories, estimate the emission for the climate models, and, in the end, provide climate and air quality scenarios and the storylines of the future development of the arctic-boreal region (Fig. 10).

In terms of strategic questions of the ESS, such as “what is the optimal mix of adaptation and mitigation measures to respond global change?” or “what is the structure of an effective and efficient system of global and development of institutions?”, PEEEX is an active player in creating direct contacts with the stakeholders, so that its scientific information and services will receive an optimal impact on decision making. Furthermore, the PEEEX approach endorses the Earth System Manifesto. (https://www.atm.helsinki.fi/peex/images/manifesti_peex_ru_hub2.pdf), which addresses three strategic tasks: (i) construction of novel observation systems, (ii) finding consensus addressing necessary mitigation and adaptation actions in different parts of the world, and (iii) operational prerequisites for technological development to moderate the global change towards the sustainable Earth System. In this framework, PEEEX will work closely with influential organizations, such as the Intergovernmental Panel for Climate Change (IPCC) delivering PEEEX assessment of Arctic-boreal region, the Future Earth acting as an Arctic-Boreal Hub, and the Digital Earth via demonstrating novel methods for integrating in-situ data with satellite observations.

Acknowledgements

The preparatory work of the PEEEX approach, started in 2012, has been mainly based on the in-kind contribution of several European, Russian, and Chinese research institutes. The work presented here would not have been possible without the collaboration of the PEEEX meetings participants. In 2012-2015 we have organized five PEEEX meetings in Helsinki (2012, 2015), Moscow (2013), Hyytiälä (2013), and Saint Petersburg (2014), and the first PEEEX

Science conference in Helsinki, Finland in February 2015. PEEEX has also been active in a frame of international coordination activities and has as such been listed as GEOSS - Gold region project, IGBP-iLEAPS Arctic and boreal regional node, Digital Earth, Arctic Council – SAON Task, one of the main collaborators of the International Eurasian Academy of Sciences (IEAS) and Future Earth Arctic-boreal hub.

In addition, we would like acknowledge the support and funding from the following bodies: Academy of Finland Centre of Excellence (grant no. 272041), “International Working Groups, Markku Kulmala” Grant by Finnish Cultural Foundation; ICOS 271878, ICOS-Finland No. 281255, ICOS-ERIC No. 281250, No. 259537, 218094, 255576, 286685, 280700 and 259537 funded by Academy of Finland; Beautiful Beijing project funded by TEKES, In GOS DEFROST and CRAICC (no 26060) and Nordforsk CRAICC-PEEX (amendment to contact 26060) funded by Nordforsk. "European-Russian Centre for cooperation in the Arctic and Sub-Arctic environmental and climate research" (EuRuCAS, grant 295068); Erasmus+ CBHE project ECOIMPACT 561975 funded by EU 7th Framework Program.

We thank many Russian projects, which have been contributed PEEEX and have been granted by national funding organizations: Russian Mega-Grant No. 11.G34.31.0048 (University of Nizhny Novgorod), Russian Ministry of Education and Science Grants (unique project identifiers RFMEFI58614X0004 and RFMEFI58314X0003, ISR “AEROCOSMOS”, 2014-2016); Russian Science Foundation projects No. 15-17-20009 (University of Nizhny Novgorod) and No. 15-17-30009 (Faculty of Geography, Moscow University, 2015-2018), RFBR Grant № 14-05-91759 (ISR “AEROCOSMOS”, 2014-2016), Russian Science Foundation projects No. 14-47-00049 (A.M. Obukhov Institute of Atmospheric Physics RAS, 2014-2016), No. 11.37.220.2016 (St.Petersburg State University) and No. 14-27-00083 (the Geochemical foundations of PEEEX), No.15-5554020 (SINP Moscow University, 2015-2016) and No. № 14-17-00096 (Saint-Petersburg State University, 2014-2016). We also

acknowledge Presidium of the Russian Academy of Sciences (Program No. 4); the Branch of Geology, Geophysics and Mining Sciences of RAS (Program No. 5); interdisciplinary integration projects of the Siberian Branch of the Russian Academy of Science No. 35, No. 70 and No. 131; Russian Foundation for Basic Research (Grants No. 14-05-00526, No. 14-05-00590, No. 14-05-93108) and Russian Science Foundation project No. № 14-17-00096 (Saint-Petersburg State University, 2014-2016). AARI thanks CNTP 1.5.3.2 and 1.5.3.4 of Roshydromet. Institute of Geography thanks Russian Academy of Sciences and Russian Science Foundation (project № 14-27-00133).

University of Tartu together with University of Life Sciences, Estonia, acknowledge the European Commission through European Regional Fund (the “Internationalization of Science Programme” project INSMEARIN (10.1-6/13/1028) and the “Estonian Research Infrastructures Roadmap” project Estonian Environmental Observatory, 3.2.0304.11-0395). The University of Tartu, Estonia, acknowledges the institutional research funding IUT20-11 of the Estonian Ministry of Education and Research for it’s support for the development of SMEAR Estonia at Järvelja. Nansen Center, Norway, acknowledges the International Belmont Forum project “Anthropogenic Heat Islands in the Arctic - Windows to the Future of the Regional Climates, Ecosystems and Societies” No.: 247468/E10.

References

- Aaltonen, H., Pumpanen, J., Hakola, H., Vesala, T., Rasmus, S., and Back, J.: Snowpack concentrations and estimated fluxes of volatile organic compounds in a boreal forest, *Biogeosciences*, 9, 2033-2044, 10.5194/bg-9-2033-2012, 2012.
- Achat, D.L., Bakker, M.R., Augusto, L., Derrien, D., Gallegos, N., Lashchinskiy, N., Milin, S., Nikitich, P., Raudina, T., Rusalimova, O., Zeller, B., and Barsukov, P.: Phosphorus status

1653 of soils from contrasting forested ecosystems in southwestern Siberia: effects of
 1654 microbial and physicochemical properties, *Biogeoscience*, 10, 733-752, 2013.

1655 Alcamo, J., Mayerhofer, P., Guardans, R., van Harmelen, T., van Minnen, J., Onigkeit, J.,
 1656 Posch, M., and de Vries, B.: An integrated assessment of regional air pollution and climate
 1657 change in Europe: findings of the AIR-CLIM Project, *Environ. Sci. Policy*, 5, 257-272, 2002.

1658 Alekseychik, P., Lappalainen, H.K., Petäjä, T., Zaitseva, N., Arshinov, M., Shevchenko,
 1659 V., Makshtas, A., Dubtsov, S., Mikhailov, E., Lapshina, E., Kirpotin, S., Kurbatova, Y.,
 1660 and Kulmala, M.: PEEX potential ground station network: an overview, *J. Geography*
 1661 *Environment, Sustainability*, in press, 2016.

1662 Allen, C.D., Makalady A.K., Chechjuini, H., Bachelet, D., McDowell, N., Vennetier, M.,
 1663 Kitzberger, T., Rigling, A., Breshears, D., Hogg, E.D. (Ted), Gonzalez, P., Fensham, R.,
 1664 Zhang Z., Castro, J., Demidova, N., Lim, J.-M., Allard, G., Running, S.W., Semerci, A., and
 1665 Cobb, N.: A global overview of drought and heat-induced tree mortality reveals emerging
 1666 climate change risk for forests, *Forest Ecology & Management*, 259, 660-684, 2010.

1667 Allen, C.D., Breshears, D.D., and McDowel, N.G.: On underestimation of global
 1668 vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene,
 1669 *Ecosphere* 6(8), 129, <http://dx.doi.org/10.1890/ES15-00203.1>, 2015.

1670 Andreae, M. O.: Ocean-atmosphere interactions in the global biogeochemical sulfur cycle,
 1671 *Mar. Chem.*, 30, 1-29, 1990.

1672 Andreae, M. O.: Correlation between cloud condensation nuclei concentration and aerosol
 1673 optical thickness in remote and polluted regions, *Atmos. Chem. Phys.*, 9, 543–556, 2009.

1674 Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning,
 1675 *Global Biogeochem. Cy.*, 15, 955–966, doi:10.1029/2000GB001382, 2001.

1676

1677 Anisimov, S.V., Mareev, E. A., and Bakastov, S. S.: On the generation and evolution of
1678 aeroelectric structures in the surface layer, *J. Geophys. Res.*, 104, 14359–14368, 1999.

1679 Anisimov, S.V., Mareev, E. A., Shikhova, N. M., and Dmitriev, E. M.: Universal spectra of
1680 electric field pulsations in the atmosphere, *Geophys. Res. Lett.*, 29, 2217–2220, 2002.

1681 Arashkevich, E.G., Flint, M.V., Nikishina, A.B., Pasternak, A.F., Timonin, A.G., Vasilieva,
1682 J.V., Mosharov, S.A., and Soloviev, K.A.: The role of zooplankton in transformation of
1683 organic matter in the Ob Estuary, on the shelf and in the deep regions of the Kara Sea,
1684 *Oceanology*, 50, 766-779, 2010.

1685 Arneth, A., Niinemets, Ü., Pressley, S., Bäck, J., Hari, P., Karl, T., Noe, S., Prentice, IC.,
1686 Serça, D., Hickler, T., Wolf, A., and Smith, B.: Process-based estimates of terrestrial
1687 ecosystem isoprene emissions: incorporating the effects of a direct CO₂-isoprene interaction,
1688 *Atmos. Chem. Phys.*, 7, 31-53, 2007.

1689 Arneth, A., Harrison, S.P., Tsigaridis, K., Menon, S., Bartlein, P.J., Feichter, H., Korhola, A.,
1690 Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., Vesala, T., and Zaehle, S.: Terrestrial
1691 biogeochemical feedbacks in the climate system: from past to future, *Nature Geoscience*, 3,
1692 525-532, 2010.

1693 Arneth, A., Svenningsson, B., Swietlicki, E., Tarozzi, L., Decesari, S., Facchini, M.C.,
1694 Birmili, W., Sonntag, A., Wiedensohler, A., Boulon, J., Sellegri, K., Laj, P., Gysel, M.,
1695 Bukowiecki, N., Weingartner, E., Wehrle, G., Laaksonen, A., Hamed, A., Joutsensaari, J.,
1696 Petäjä, T., Kerminen, V.-M., and Kulmala M.: EUCAARI ion spectrometer measurements at
1697 12 European sites — analysis of new particle formation events, *Atmos. Chem. Phys.*, 10,
1698 7907-7927, 2010.

1699 Arneth, A., Olin, S., Makkonen, R., Paasonen, P., Holst, T., Kajos, M., Kulmala, M.,
 1700 Maximov, T., Miller, P.A., and Achurgers, G.: Future biogeochemical forcing in Eastern
 1701 Siberia: cooling or warming?, *Atmos. Chem. Phys. Discuss.*, 14, 19149-19179,
 1702 doi:10.5194/acpd-14-19149-2014, 2014.

1703 Arnold, S. R., Emmons, L. K., Monks, S. A., Law, K. S., Ridley, D. A., Turquety, S.,
 1704 Tilmes, S., Thomas, J. L., Bouarar, I., Flemming, J., Huijnen, V., Mao, J., Duncan, B. N.,
 1705 Steenrod, S., Yoshida, Y., Langner, J., and Long, Y. Biomass burning influence on high-
 1706 latitude tropospheric ozone and reactive nitrogen in summer 2008: a multi-model analysis
 1707 based on POLMIP simulations, *Atmos. Chem. Phys.*, 15, 6047-6068, 2015.

1708 Arshinov, M., Yu., M. Belan, B.D., Kozlov, A.V., Antokhin, P.N., Davydov, D.K., and
 1709 Arshinova V.G.: Continuous measurements of aerosol size distribution at two Siberian
 1710 stations: new particle formation bursts, in: European Aerosol Conference, Granada, Spain, 2-
 1711 7 September 2012, A-WG01S1P25: <http://www.eac2012.com/EAC2012Book/files/341.pdf>,
 1712 2012.

1713 Asmi, E., Kivekäs, N., Kerminen, V.-M., Komppula, M., Hyvärinen, A.-P., Hatakka, J.,
 1714 Viisanen, Y., and Lihavainen, H.: Secondary new particle formation in Northern Finland
 1715 Pallas site between the years 2000 and 2010, *Atmos. Chem. Phys.*, 11, 12959-12972,
 1716 doi:10.5194/acp-11-12959-2011, 2011.

1717 Asmi, E., Kondratyev, V., Brus, D., Laurila, T., Lihavainen, H., Backman, J., Vakkari, V.,
 1718 Aurela, M., Hatakka, J., Viisanen, Y., Uttal, T., Ivakhov, V., and Makshtas, A.: Aerosol size
 1719 distribution seasonal characteristics measured in Tiksi, Russian Arctic, *Atmos. Chem. Phys.*
 1720 *Discuss.*, 15, 18109-18149, doi:10.5194/acpd-15-18109-2015, 2015.

1721 Atlaskina, K., Berninger, F., and de Leeuw, G.: Satellite observations of changes in snow-
 1722 covered land surface albedo during spring in the Northern Hemisphere, *The Cryosphere*, 9,
 1723 1879-1893, doi:10.5194/tc-9-1879-2015, 2015.

1724 Bala, G., Krishna, S., Narayanappa, D., Cao, L., Caldeira, K., and Nemani, R.: An estimate
 1725 of equilibrium sensitivity of global terrestrial carbon cycle using NCAR CCSM4, *Clim Dyn*,
 1726 40, 1671–1686, 2013.

1727 Ballantyne, A.P., Alden, C.B., Miller, J.B., Tans, P.P., and White, J. W. C.: Increase in
 1728 observed net carbon dioxide uptake by land and oceans during the past 50 years, *Nature*, 488,
 1729 70–72, doi:10.1038/nature11299, 2012.

1730 Baklanov, A.A., Penenko, V.A., Mahura, A.G., Vinogradova, A.A., Elansky, N.F., Tsvetova,
 1731 E.A., Rigina, O., Yu., Maksimenkov, L.O., Nuterman, R.B., Pogarskii, F.A., and A. Zakey,
 1732 A.: Aspects of atmospheric pollution in Siberia. In Gutman G. and Groisman P. (eds),
 1733 *Regional Environmental Changes in Siberia and Their Global Consequences*, Springer, 2013,
 1734 pp. 303-346, 2013.

1735 Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., and Enrich-Prest, A.: Freshwater
 1736 methane emissions offset the continental carbon sink, *Science*, 331, 50, doi:
 1737 10.1126/science.1196808, 2011.

1738 Battarbee, R. W., Patrick, S., Kernan, M., Psenner, R., Thies, H., Grimalt, J., Rosseland, B.
 1739 O., Wathne, B., Catalan, J., Mosello, R., Lami, A., Livingstone, D., Stuchlik, E.,
 1740 Straskrabova, V., and Raddum, G.: High mountain lakes and atmospherically transported
 1741 pollutants. In: Huber, U. M., Bugmann, H. K. M., and Reasoner, M. A. (Eds), *Global Change*
 1742 *and Mountain Regions*, Springer, 2005.

1743 Bauer, P., Thorpe, A., and Brunet, G.: The quiet revolution of numerical weather prediction,
 1744 Nature, 525, 47–55, doi:10.1038/nature14956, 2015.

1745 Bauerle, W.L., Bauerlea, W.L., Orenc, R., Wayc, D.A., Qianc, S.S., Stoyf, P.C., Thorntong,
 1746 P.E., Bowdena, J.D., Hoffmanh, F.M., and Reynoldsi, R.F.: Photoperiodic regulation of the
 1747 seasonal pattern of photosynthetic capacity and the implications for carbon cycling, PNAS,
 1748 109, www.pnas.org/cgi/doi/10.1073/pnas.1119131109, 2012.

1749 Berchet, A., Paris, J.-D., Ancellet, G., Law, K. S., Stohl, A., Nédélec, P., Arshinov, M., Yu,
 1750 M., Belan, B. D., and Ciais, P.: Tropospheric ozone over Siberia in spring 2010: remote
 1751 influences and stratospheric intrusion, Tellus B, 65, 19688,
 1752 <http://dx.doi.org/10.3402/tellusb.v65i0.19688>, 2013.

1753 Berchet, A., Pison, I., Chevallier, F., Paris, J.-D., Bousquet, P., Bonne, J.-L.,
 1754 Arshinov, M. Y., Belan, B. D., Cressot, C., Davydov, D. K., Dlugokencky, E. J.,
 1755 Fofonov, A. V., Galanin, A., Lavrič, J., Machida, T., Parker, R., Sasakawa, M., Spahni, R.,
 1756 Stocker, B. D., and Winderlich, J.: Natural and anthropogenic methane fluxes in Eurasia: a
 1757 mesoscale quantification by generalized atmospheric inversion, Biogeosciences, 12, 5393-
 1758 5414, doi:10.5194/bg-12-5393-2015, 2015.

1759 Bergen, K.M., Hitztaler, S.K., Kharuk, V.I., Krankina, O.N., Loboda, T.V., Zhao, T.,
 1760 Shugart, H. H., and Sun, G.: Human dimensions of environmental change in Siberia,
 1761 Regional Environmental Changes in Siberia and Their Global Consequences, Groisman, P.
 1762 Y., and Gutman, G., (Eds),

1763 Berner, L.T., Beck, P.S.A., Bunn A.G., and Goetz, S.J.: Plant response to climate change
 1764 along the forest-tundra ecotone in northeastern Siberia, Global Change Biology, 19 (11),
 1765 3449–3462, 2013.

1766 Berry, P. A. M., Garlick, J. D., Freeman, J. A., and Mathers, E. L.: Global inland water
 1767 monitoring from multi-mission altimetry, *GRL*, 32, L16401, doi:10.1029/2005GL022814,
 1768 2005.

1769 Bluhm, B.A., Gebruk, A.V., Gradinger, R., Hopcroft, R.R., Huettmann, F., Kosobokova,
 1770 K.N., Sirenko, B.I., and Weslawski, J.M.: Arctic Marine Biodiversity: An Update of Species
 1771 Richness and Examples of Biodiversity Change, *Oceanography*, 24, 232-248, 2011.

1772 Bityukova, V.R. and Kasimov, N.S.: Atmospheric pollution of Russia's cities: assessment of
 1773 emissions and immissions based on statistical data, *Geofizika*, 29, 53-67, 2012.

1774 Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M.,
 1775 Bustamante, M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.W., Fenn,
 1776 M., Gilliam, F., Nordin, A., Pardo, L., and de Vries, W.: Global assessment of nitrogen
 1777 deposition effects on terrestrial plant diversity: a synthesis, *Ecol. Appl.*, 20, 30-59, 2010.

1778 Bodman, R.W., Rayner, P.J., and Karoly D.J.: Uncertainty in temperature projections
 1779 reduced using carbon cycle and climate observations, *Nature Climate Change*, 3, 725-729,
 1780 2013.

1781 Bolin, B., Ciais, P., Cramer, W., Jarvis, P., Kheshgi, H., Nobre, C., Semenov, S., and
 1782 Steffen, W.: Global perspective, In: Watson, R.T, Noble, I.R., Bolin, B., Ravindranath, N.H.,
 1783 Verardo D.J., and Dokken, D.J. (Eds.), *Land use, land-use change, and forestry. A special*
 1784 *report of the IPCC*, Cambridge University Press, Cambridge, 23-51, 2000.

1785 Bond, T.C., Doherty, S.J., Fahey, D.W., Foster, P.M., Berntsen, T., DeAngelo, B.J., Flanner,
 1786 M.G., Ghan, S., Kärcher, B., Kock, D., Kinne, S., Kondo, Y., Quinn, P.K., Sarofilm, M.C.,
 1787 Schulz, M., Venkataram, C., Zhang, H., Zhang, S., Belloiin, N., Guttukunda, S.K., Hopke,
 1788 P.K., Jacobson, M.Z., Kaiser, J.W., Klimont Z., Lohmann, U., Schwarz, J.P., Shindell, D.,

1789 Storelmo, T., Warren, S.G., Zender, C.S.: Bounding the role of black carbon in the climate
 1790 system: A scientific assessment, JGR, 118, 5380-5552, DOI: 10.1002/jgrd.50171., 2013.

1791 Bondur V.G.: Aerospace Methods and Technologies for Monitoring Oil and Gas Areas and
 1792 Facilities // Izvestiya, Atmospheric and Oceanic Physics, 47 (9), 1007-1018, 2011a.

1793 Bondur V.G.: Satellite Monitoring of Wildfires during the Anomalous Heat Wave of 2010 in
 1794 Russia // Izvestiya, Atmospheric and Oceanic Physics, 47(9), 1039-1048, 2011b.

1795 Bondur V.G.: Modern Approaches to Processing Large Hyperspectral and Multispectral
 1796 Aerospace Data Flows. Izvestiya, Atmospheric and Oceanic Physics, 50(9), 840-852, DOI:
 1797 10.1134/S0001433814090060, 2014.

1798 Bondur V.G. and Vorobev V.E.: Satellite monitoring of impact Arctic regions, Izvestiya,
 1799 Atmospheric and Oceanic Physics, 51(9), 949–968, 2015.

1800 Bondur V.G. and Ginsburg A.S.: Emissions of carbon-containing gases and aerosols resulted
 1801 from wildfires in Russia according to space-borne monitoring data, Doklady Akademii
 1802 nauk., 466(4), 473-477, 2016.

1803 Bondur, V.G.: Space-borne monitoring of trace gas and aerosol emissions during wildfires in
 1804 Russia, Issledovanie Zemli iz Kosmosa, 6, 3-19, 2015.

1805 Boyle, J. F., Mackay, A.W., Rose, N. L., Flower, R. J., and Appleby, P. G.: Sediment heavy
 1806 metal records in Lake Baikal: natural and anthropogenic sources, J. Palaeolimnol, 20, 135–
 1807 150, 1998.

1808 Brunello, A.J., Molotov, V. C., Dugherkhuu, B., Goldman, C., Khamaganova, E. ,
 1809 Strijhova, T., and Sigman, R.: Lake Baikal. Experiences and Lessons Learned Brief, Tahoe-
 1810 Baikal Institute, South Lake Tahoe, CA, USA, 2006.

1811 Burrows, J.P., Platt, U., and Borrel, P., (Editors): The Remote Sensing of Tropospheric
 1812 Composition from Space, 536 pp., Springer-Verlag Berlin Heidelberg ISBN: 978-3-642-
 1813 14790-6 doi: 10.1007/978-3-642-14791-3., 359-313, 2011.

1814 Byambaa, B. and Todo, Y.: Technological Impact of Placer Gold Mine on Water Quality:
 1815 Case of Tuul River Valley in the Zaamar Goldfield, Mongolia, World Academy of Science,
 1816 Eng. Tech., 54, 2011.

1817 Bäck, J., Aaltonen, H., Hellen, H., Kajos, M. K., Patokoski, J., Taipale, R., Pumpanen, J.,
 1818 and Heinonsalo, J.: Variable emissions of microbial volatile organic compounds (MVOCs)
 1819 from root-associated fungi isolated from Scots pine, Atmos. Environ., 44, 3651-3659,
 1820 doi:10.1016/j.atmosenv.2010.06.042, 2010.

1821 Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S.,
 1822 and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth
 1823 system, Atmos. Chem. Phys., 10, 1701-1737, doi:10.5194/acp-10-1701-2010, 2010.

1824 Cavalieri, D. J. and Parkinson, C. L.: Arctic sea ice variability and trends, 1979-2010,
 1825 Cryosphere. 6, 881–889, 2012.

1826 Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton,
 1827 atmospheric sulphur, cloud albedo and climate, Nature, 326, 655-661, 1987.

1828 Cheng, B., Zhang, Z., Vihma, T., Johansson, M., Bian, L., Li, Z., and Wu, H.: Model
 1829 experiments on snow and ice thermodynamics in the Arctic Ocean with CHINARE2003
 1830 data, J. Geophys. Res., 113, C09020, doi:10.1029/2007JC004654, 2008.

1831 Chernokulsky A.V. and Mokhov I.I.: Climatology of total cloudiness in the Arctic: An
 1832 intercomparison of observations and reanalyses, Adv. in Meteorol., 2012, Article ID 542093,
 1833 15 pages, doi: 10.1155/2012/542093, 2012.

1834 Chi, X., Winderlich, J., Mayer, J.-C., Panov, A. V., Heimann, M., Birmili, W., Heintzenberg,
 1835 J., Cheng, Y., and Andreae, M. O.: Long-term measurements of aerosols and carbon
 1836 monoxide at the ZOTTO tall tower to characterize polluted and pristine air in the Siberian
 1837 Taiga: *Atmos. Chem. Phys.*, 13, 12,271-12,298, doi:10.5194/acp-13-12271-2013, 2013.

1838 Ciais, P., Janssens, I., Shvidenko, A., Wirth, C., Malhi, Y., Grace, J., Schulze, E.D.,
 1839 Heimann, M., Phillips, O., and Dolman, A.J.: The potential for rising CO₂ to account for the
 1840 observed uptake of carbon by tropical, temperate, and boreal forests biome. In Griffiths H. &
 1841 Jarvis P.J. (Eds), *The Carbon Budget of Forest Biomes*, Garland Science, BIOS Scientific
 1842 Publishers, 109-149, 2005.

1843 Cohen, J., Screen, J.A., Furtado, J.C., Barlow, M., Whittleston, D., Coumou, D., Francis, J.,
 1844 Dethloff, K., Entekhabi, D., Overland, J., and Jones, J.: Recent Arctic amplification
 1845 and extreme mid-latitude weather, *Nature Geoscience*, 7, 627–637, doi:10.1038/ngeo2234,
 1846 2014.

1847 Conard, S.G., Sukhinin, A.I., Stocks B.J., Cahoon, D.R., Davidenko, E.P., and Ivanova,
 1848 G.A.: Determining effects of area burned and fire severity on carbon cycling and emissions
 1849 in Siberia, *Climatic Change*, 55, 197-211, 2002.

1850 Conley, D.J., Paerl, H.Q., Howarth, R.W., Boesch, D.F., Seitzinger, S.P., Havens, K.E.,
 1851 Christiane Lancelot, C., and Likens, G.E.: Controlling eutrophication: nitrogen and
 1852 phosphorus, *Science*, 323, 1014-1015, 2009.

1853 Crutzen, P.J., Elansky, N.F., Hahn, M., Golitsyn, G.S., Benninkmeijer, C.A.M., Scharffe,
 1854 D.H., Belikov, I.B., Maiss, M., Bergamaschi, P., Röckmann, T., Grisenko A.M., and
 1855 Sevostyanov, V.M.: Trace gas measurements between Moscow and Vladivostok using the
 1856 Trans-Siberian Railroad, *J. Atmos. Chem.*, 29, 179-194, 1998.

1857 Dal Maso, M., Sogacheva, L., Aalto, P.P., Riipinen, I., Komppula, M., Tunved, P.,
 1858 Korhonen, L., Suur-Uski, V., Hirsikko, A., Kurtén, T., Kerminen, V.-M., Lihavainen, H.,
 1859 Viisanen, Y., Hansson, H.-C., and Kulmala, M.: Aerosol size distribution measurements at
 1860 four Nordic field stations: identification, analysis and trajectory analysis of new particle
 1861 formation bursts, *Tellus*, 59B, 350-361, 2007.

1862 Davy, R. and Esau, I.: Global climate models' bias in surface temperature trends and
 1863 variability, *Environmental Research Letters*, 9, 114024, 2014, doi: 10.1088/1748-
 1864 9326/9/11/114024

1865 Davy, R. and Esau, I.: Differences in the efficacy of climate forcings explained by variations
 1866 in atmospheric boundary layer depth, *Nature Communications*, 7, 11690,
 1867 2016,doi:10.1038/ncomms11690

1868 de Leeuw, G., Andreas, E.L., Anguelova, M.D., Fairall, C.W., Lewis, E.R., O'Dowd, C.,
 1869 Schulz, M., and Schwartz, S.E.: Production flux of sea spray aerosol, *Rev. Geophys.*, 49,
 1870 RG2001, doi:10.1029/2010RG000349, 2011.

1871 de Leeuw, G., Holzer-Popp, T., Bevan, S., Davies, W., Descloîtres, J., Grainger, R.G.,
 1872 Griesfeller, J., Heckel, A., Kinne, S., Klüser, L., Kolmonen, P., Litvinov, P., Martynenko,
 1873 D., North, P.J.R., Ovigneur, B., Pascal, N., Poulsen, C., Ramon, D., Schulz, M., Siddans, R.,
 1874 Sogacheva, L., Tanré, D., Thomas, G.E., Virtanen, T.H., von Hoyningen Huene, W.,
 1875 Vountas, M., and Pinnock, S.: Evaluation of seven European aerosol optical depth retrieval
 1876 algorithms for climate analysis, *Remote Sensing of Environment*, 162, 295–315, DOI
 1877 information: 10.1016/j.rse.2013.04.023, 2015.

1878 Dentener, F., Drevet, J., Lamarque, J.F., Bey, I., Eickhout, B., Fiore, A.M., Hauglustaine, D.,
 1879 Horowitz, L.W., Krol, M., Kulshrestha, U.C., Lawrence, M., Galy-Lacaux, C., Rast, S.,

1880 Shindell, D., Stevenson, D., Noije, T.V., Atherton, C., Bell, N., Bergman, D., Butler, T.,
 1881 Cofala, J., Collins, B., Doherty, R., Ellingsen, K., Galloway, J., Gauss, M., Montanaro, V.,
 1882 Müller, J.F., Pitari, G., Rodriguez, J., Sanderson, M., Solmon, F., Strahan, S., Schultz, M.,
 1883 Sudo, K., Szopa, S., and Wild, O.: Nitrogen and sulfur deposition on regional and global
 1884 scales: a multimodel evaluation, *Global Biogeochemical Cycles*, 20, 21, doi:
 1885 10.1029/2005GB002672, 2006.

1886 Ding, A.J., Wang, T., Thouret, V., Cammas, J.-P., and Nedelec, P.: Tropospheric ozone
 1887 climatology over Beijing: Analysis of aircraft data from the MOZAIC program, *Atmos.*
 1888 *Chem. Phys.*, 8, 1-13, 2008.

1889 Ding, A.J., Wang, T., Xue, L.K., Gao, J., Stohl, A., Lei, H.C., Jin, D.Z. Ren, Y., Wang, X.Z.,
 1890 Wei, X.L., Qi, Y.B., Liu, J., and Zhang, X.Q.: Transport of north China air pollution by
 1891 mid-latitude cyclones: Case study of aircraft measurements in summer 2007, *J. Geophys.*
 1892 *Res.*, 114, D08304, 2009.

1893 Ding, J., Zhong, J., Yang, Y., Li, B., Shen, G., Su, Y., Wang, C., Li, W., Shen, H., Wang, B.,
 1894 Wang, R., Huang, Y., Zhang, Y., Cao, H., Zhun, Y., Simonich, S. L. M., and Tao, S.:
 1895 Occurrence and exposure to polycyclic aromatic hydrocarbons and their derivatives in a rural
 1896 Chinese home through biomass fuelled cooking, *Environ. Poll.*, 169, 160-166, 2012.

1897 Ding, A., Fu, C.B., Yang, X.Q., Sun, J.N., Peteja, T., Kerminen, V.-M., Wang, T., Xie.,
 1898 Y.N., Herrmann, E., Zheng, L.F., W. Nie, W., Liu, Q., Wei, X.L., and Kulmala, M.: Intense
 1899 atmospheric pollution modifies weather: a case of mixed biomass burning with fossil fuel
 1900 combustion pollution in the eastern China, *Atmospheric Chemistry and Physics*, 13, 10545-
 1901 10554, 2013a.

1902 Ding, A. J., Fu, C. B., Yang, X. Q., Sun, J. N., Zheng, L.F., Xie, Y.N., Hermann, E., Nie, W.,
 1903 Petäjä, T., Kerminen, V.M., and Kulmala, M.: Ozone and fine particle in the western
 1904 Yangtze River Delta: an overview of 1 yr data at the SORPES station, *Atmos. Chem. Phys.*,
 1905 13, 11, 5813-5830, doi:10.5194/acp-13-5813-2013, 2013b.

1906 Dolezalek, H., Reiter, R., and Landsberg, H.E. (editors): *Electrical Processes in*
 1907 *Atmospheres. Proceedings of the Fifth International Conference on Atmospheric Electricity*
 1908 *held at Garmisch-Partenkirchen (Germany), 2–7 September 1974. Dr. Dietrich Steinkopff*
 1909 *Verlag, Darmstadt, 1977. ISBN: 978-3-642-85296-1 (Print) 978-3-642-85294-7 (Online)*

1910 Domysheva, V., Panchenko, M., Pestunov, D., and Sakirko M.: Air-Water Carbon Dioxide
 1911 Exchange in the Littoral Zone of Lake Baikal (Ice-Free Period), *International Journal of*
 1912 *Geosciences*, 4, 1339-1345, doi:[10.4236/ijg.2013.410130](https://doi.org/10.4236/ijg.2013.410130), 2013.

1913 Ducklow, H., Steinber, D.K., and Buesseler, K.O.: Upper ocean carbon export and the
 1914 biological pump, *Oceanography*, 14, 50-58, 2001.

1915 Dooley, S. and Treseder, K.: The effect of fire on microbial biomass: a meta-analysis of field
 1916 studies, *Biogeochemistry*, 109, 49, 2012.

1917 Dolman, A. J., Shvidenko, A., Schepaschenko, D., Ciais, P., Tchebakova, N., Chen, T., van
 1918 der Molen, M. K., Beileli Marchesini, L., Maximov, T. C., Maksyutov, S., and Schulze, E.-
 1919 D.: An estimate of the terrestrial carbon budget of Russia using inventory-based, eddy
 1920 covariance and inversion methods, *Biogeosciences*, 9, 5323-5340, doi:10.5194/bg-9-5323-
 1921 2012, 2012.

1922 Drake, J.E., Gallet-Budynek A., Hofmockel, K.S., Bernhardt, E., Billings, S., Jackson, R. B.,
 1923 Johnsen, K.S., Lichter, J., McCarthy, H.R., McCormack L., Moore, D., Oren, R., Palmroth,
 1924 S., Phillips, R.P., Phippen, J.S., Pritchard, S., Treseder, K.K., Schlesinger W.H., DeLucia E.,

1925 and Finzi, A.C.: Increases in the flux of carbon belowground stimulate nitrogen uptake and
 1926 sustain the long-term enhancement of forest productivity under elevated CO₂, *Ecol. Lett.*, 14,
 1927 349–357, doi: 10.1111/j.1461-0248.2011.01593.x, 2011.

1928 Döscher, R., Vihma, T., and Maksimovich, E.: Recent advances in understanding the Arctic
 1929 climate system state and change from a sea ice perspective: a review, *Atmos. Chem. Phys.*,
 1930 14, 13571-13600, doi:10.5194/acp-14-13571-2014, 2014.

1931 Eckhardt, S., Quennehen, B., Olivié, D.J.L., Berntsen, T.K., Cherian, R. Christensen, J.H.,
 1932 Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog,
 1933 Ø. , Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K.S., Lund, M.T., Mahmood,
 1934 R., Massling, A., Myriokefalitakis, S., Nielsen, I.E., Nøjgaard, J.K., Quaas, J., Quinn, P.K.,
 1935 Raut, J.-C., Rumbold, S.T., Schulz, M., Sharma, S., Skeie, R.B., Skov, H., Uttal, T., von
 1936 Salzen, K., and Stohl A.: Current model capabilities for simulating black carbon and sulfate
 1937 concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive
 1938 measurement data set, *Atmos. Chem. Phys.*, 15, 9413-9433, 2015.

1939 Elansky, N.F., Belikov, I.B., Lavrova, O.V., Skorokhod, A.I., Shumsky, R.A.,
 1940 Brenninkmeijer, C.A.M., and Tarasova, O.A.: Chapter 8. Train-Based Platform for
 1941 Observations of the Atmosphere Composition (TROICA Project) P. 175-196, In “Air
 1942 Pollution-Monitoring, Modelling and Health”. Ed. Mukesh Khare ISBN: 978-953-51-0424-
 1943 7, Tech., 386, doi 10.5772/1801, 2012.

1944

1945 Elberling, B., Christiansen, H. H., and Hansen, B. U.: High nitrous oxide production from
 1946 thawing permafrost, *Nature Geoscience*, 3, 332-335, 2010.

1947 Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Budel, B., Andreae, M. O., and Poschl,
 1948 U.: Contribution of cryptogamic covers to the global cycles of carbon and nitrogen, *Nature*
 1949 *Geosci*, 5, 459-462, 2012.

1950 Emmons, L. K., Arnold, S. R., Monks, S. A., Huijnen, V., Tilmes, S., Law, K. S., Thomas, J.
 1951 L., Raut, J.-C., Bouarar, I., Turquety, S., Long, Y., Duncan, B., Steenrod, S., Strode, S.,
 1952 Flemming, J., Mao, J., Langner, J., Thompson, A. M., Tarasick, D., Apel, E. C., Blake, D.
 1953 R., Cohen, R. C., Dibb, J., Diskin, G. S., Fried, A., Hall, S. R., Huey, L. G., Weinheimer, A.
 1954 J., Wisthaler, A., Mikoviny, T., Nowak, J., Peischl, J., Roberts, J. M., Ryerson, T., Warneke,
 1955 C., and Helmig, D. : The POLARCAT Model Intercomparison Project (POLMIP): Overview
 1956 and evaluation with observations, *Atmos. Chem. Phys.*, 15, 6721-6744, 2015.

1957 Engvall -Stjernberg A.-C., Skorokhod, A., Paris, J.-D., Elansky, N., Nédélec, P., and Stohl
 1958 A.: Low concentrations of near-surface ozone in Siberia, *Tellus B*, 64, doi:
 1959 10.3402/tellusb.v64i0.11607, 2012.

1960 Erisman, J.W., Grinsven, H.V., Grizzetti, B., Bouraoui, F., and Powlson, D.: The European
 1961 nitrogen problem in a global perspective, in: Sutton, M.A., Howard, C.M., Erisman, J.W.,
 1962 Billen, G., Bleeker, A., Grennfelt, P., and Hansen, J. (Eds.) :The European Nitrogen
 1963 Assessment, Cambridge University Press, Cambridge, UK, 2011.

1964 Esau, I.; Davy, R., and Outten, S.:Complementary explanation of temperature response in the
 1965 lower atmosphere, *Environmental Research Letters*, 7, 044026, 2012, doi: 10.1088/1748-
 1966 9326/7/4/044026

1967 Esau, I.N. and Chernokulsky A.V.,: Convective Cloud Fields in the Atlantic Sector of the
 1968 Arctic: Satellite and Ground-Based Observations, *Izvestiya, Atmospheric and Oceanic*
 1969 *Physics*, 51, 1007–1020, 2015.

1970 Esau, I.; Miles, V. V.; Davy, R.; Miles, M. W., and Kurchatova, A. Trends in normalized
 1971 difference vegetation index (NDVI) associated with urban development in northern West
 1972 Siberia, *Atmospheric Chemistry and Physics*, 16, 9563–9577, doi: 10.5194/acp-16-9563-
 1973 2016 , 2016.

1974 European Nitrogen Assessment, Sutton, M. A., Howard, C. M., Willem Erisman, J., Billen,
 1975 G., Bleeker, A., Grennfelt, P., van Grinsven, H., and Grizzetti, B. (Eds.): Cambridge
 1976 University Press, 2010.

1977 Faubert, P., Tiiva, P., Michelsen, A., Rinnan, Å., Ro-Poulsen, H., and Rinnan, R.: The shift
 1978 in plant species composition in a subarctic mountain birch forest floor due to climate change
 1979 would modify the biogenic volatile organic compound emission profile, *Plant and Soil*, 352,
 1980 199–215, doi:10.1007/s11104-011-0989-2, 2012.

1981 Fehsenfeld, F. C., Fehsenfeld, F. C., Ancellet, G., Bates, T. S., Goldstein, A. H., Hardesty,
 1982 M., Honrath, R., Law, K. S., Lewis, A. C., Leaitch, R., McKeen, S., Meagher, J., Parrish, D.
 1983 D., P. Pszenny, A. A., Russell, P. B., Schlager, H., Seinfeld, J., Talbot, R., and Zbinden, R.:
 1984 International Consortium for Atmospheric Research on Transport and Transformation
 1985 (ICARTT): North America to Europe-Overview of the 2004 summer field study, *J. Geophys.*
 1986 *Res.*, 111, D23S01, doi:10.1029/2006JD007829, 2006.

1987 Feuchtmayr, H., Moran, R., Hatton, K., Cannor, L., Yeyes, T., Harley, J., and Arkinson, D.:
 1988 Global warming and eutrophication: effects on water chemistry and autotrophic communities
 1989 in experimental hypertrophic shallow lake mesocosms, *J. Appl. Ecol.*, 46, 713-723, 2009.

1990 Fisher, J. B., Tu, K. P., and Baldocchi, D. D.: Global estimates of the land-atmosphere water
 1991 flux based on monthly AVHRR and ISLSCP-II data, validated at 16 FLUXNET sites,
 1992 *Remote Sensing of Environment*, 112, 901-919, 2008.

- 1993 Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan,
1994 V., and Rasch, P. J.: Springtime warming and reduced snow cover from carbonaceous
1995 particles, *Atmos. Chem. Phys.*, 9, 2481-2497, 2009.
- 1996 Flanner, M. G., Shell, K. M., Barlage, M., Perovich, D. K., and Tschudi, M. A.: Radiative
1997 forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and
1998 2008, *Nature Geoscience*, 4, 151-155, 2011.
- 1999 Flannigan, M.D., Stocks, B.J., Turetsky, M. R., and Wotton, B. M.: Impact of climate change
2000 on fire activity and fire management in the circumboreal forest, *Global Change Biology*, 15,
2001 549-560, 2009.
- 2002 Flint, M.V. (Ed.): *Ecosystem of the Kara Sea*, *Oceanology*, 50(5), 637-820, 2010.
- 2003 Garmaev, E.Zh. and Khristoforov, A.V.: *Water resources of Rivers in Lake Baikal Basin:*
2004 *Principles of their use and protection* Novosibirsk (in Russia), GEO, 2010.
- 2005 Gauthier, S., Bernier P., Kuuluvainen T., A. Z. Shvidenko, A.Z., and Schepaschenko,
2006 D.G.: *Boreal forest health and global change*, *Science*, 349, 819- 822, 2015.
- 2007 Gewehr, S., Drobyshev, I., Berninger, F., and Yves Bergeron, Y.: Soil characteristics
2008 mediate the distribution and response of boreal trees to climatic variability, *Canadian Journal*
2009 *of Forest Research*, 44, 487-498, 10.1139/cjfr-2013-0481, 2014.
- 2010 Glezer, O.B.: *Population and Its Settlement Pattern*, in: *Space, Population, and Economics of*
2011 *Yugra. Socioeconomic Transformation of the Khanty-Mansi Autonomous Okrug*,
2012 *Artobolevsky, S.S., Glezer, O.B. (Eds.), Ekonomist*, 169-191, (in Russian), 2007a.
- 2013 Glezer, O.B.: *The Development of the North Abroad: Experience and Lessons*,
2014 *Environmental Planning and Management*, 4, 62-76, (in Russian), 2007b.

2015 Glindemann, D., Edwards, M., Liu, J., and Kusch, P.: Phosphine in soils, sludges, biogases
2016 and atmospheric implications - a review, *Ecological Engineering*, 24, 457-463, 2005.

2017 Goldenson, N., Doherty, K. S., Bitz, C. M., Holland, M. M., Light, B., and Conley, A. J.:
2018 Arctic climate response to forcing from light-absorbing particles in snow and sea ice in
2019 CESM, *Atmos. Chem. Phys.*, 12, 7903-7920, 2012.

2020 Granier, C., Bessagnet, B., Bond, T., and D'Angiola A.: Evolution of anthropogenic and
2021 biomass burning emissions of air pollutants at global and regional scales during the 1980-
2022 2010 periods, *Climatic Change*, 109, 163-190, 2011.

2023 Granina, L.Z.: The chemical budget of Lake Baikal—A review, *Limnol. Oceanogr.*, 42, 373-
2024 378, 1997.

2025 Groisman, P. Ya., Gutman, G., Shvidenko, A.Z., Bergen, K., Baklanov, A.A., and
2026 Stackhouse, Jr., P.W.: Introduction—Regional features of Siberia. Groisman, P. Ya., and
2027 Gutman, G. (Eds), *Regional Environmental Changes in Siberia and Their Global*
2028 *Consequences*, Springer, 1-17, 2013.

2029 Gromtsev, A.: Natural Disturbance Dynamics in Boreal Forests of European Russia: a
2030 Review, *Silva Fenn.*, 36, 41, 2002.

2031 Grote, R. and Niinemets, Ü.: Modeling volatile isoprenoid emissions – a story with split
2032 ends, *Plant Biol.*, 10, 8–28, 2008.

2033 Gruber, N., and Galloway, J. N.: An Earth-system perspective of the global nitrogen cycle,
2034 *Nature*, 451, 293–296, 2008.

2035 Guan, T., Yao, M., Wang, J., Fang, Y., Hu, S., Wang, Y., Dutta, A., Yang, J., Wu, Y., Hu,
 2036 M., and Zhu, T.: Airborne endotoxin in fine particulate matter in Beijing, *Atmos. Environ.*,
 2037 97, 35-42, 2014.

2038 Gurney, K.R., Rachel M., Law R.M., Denning A.S., Rayner P.J., Baker D., Bousquet Ph.,
 2039 Bruhwiler L., Chen Y.-H., Ciais Ph., Fan S., Fung I.Y., Gloor M., Heimann M., Higuchi K.,
 2040 John J., Maki T., Maksyutov Sh., Masarie K., Peylin Ph., Prather M., Pak B.C., Randerson
 2041 J., Sarmiento J., Taguchi S., Takahashi T., and Yuen Ch.-W.: Towards robust regional
 2042 estimates of CO₂ sources and sinks using atmospheric transport models, *Nature (Gr. Brit.)*,
 2043 415, 626-630, 2002.

2044 Gustafsson, Ö., van Dongen, B. E., Vonk, J. E., Dudarev, O. V., and Semiletov, I. P.:
 2045 Widespread release of old carbon across the Siberian Arctic echoed by its large rivers,
 2046 *Biogeosciences*, 8, 1737-1743, doi:10.5194/bg-8-1737-2011, 2011.

2047 Heald, C. L. and Spracklen, D.V.: Atmospheric budget of primary biological aerosol particles
 2048 from fungal spores, *Geophys. Res. Lett.*, 36, L09806, doi:10.1029/2009GL037493, 2009.
 2049

2050 Holtslag, A.A.M., G. Svensson, G., Baas, P., S. Basu, S., Beare, B., Beljaars, A.C.M.,
 2051 Bosveld, F.C., Cuxart, J., Lindvall, J., Steeneveld, G.J., Tjernström, M., and. Van De Wiel,
 2052 B. J. H : Stable Atmospheric Boundary Layers and Diurnal Cycles: Challenges for Weather
 2053 and Climate Models. *Bull. Amer. Meteor. Soc.*, 94, 1691-1706, doi:
 2054 <http://dx.doi.org/10.1175/BAMS-D-11-00187.1>, 2013.

2055 Harden, J. W., Trumbore, S. E., Stocks, B. J., Hirsch, A., Gower, S. T., O'Neill, K. P., and
 2056 Kasischke, E. S.: The role of fire in the boreal carbon budget, *Global Change Biology*, 6,
 2057 174-184, doi: 10.1046/j.1365-2486.2000.06019.x, 2000.

2058 Hari, P., Petäjä, T., Bäck, J., Kerminen, V-M., Lappalainen, H. K., Vihma, T., Laurila, T.,
 2059 Viisanen, Y., Vesala, T., and Kulmala, M.: Conceptual design of a measurement network of
 2060 the global change Atmos. Chem. Phys. Discuss., 15, 21063–21093, 2016.

2061 Hari, P. and Kulmala, M.: Stations for Measuring Ecosystem – Atmosphere Relations
 2062 (SMEAR II), Boreal Environment Research, 10, 315-322, 2005.

2063 Heikkinen, J. E. P., Virtanen, T., Huttunen, J. T., Elsakov, V., and Martikainen, P. J.: Carbon
 2064 balance in East European tundra, Glob. Biogeochem. Cycles, 18, GB, 1023, 2004.

2065 Heald, C. L. and Spracklen, D.V.: Atmospheric budget of primary biological aerosol
 2066 particles from fungal spores, Geophys. Res. Lett., 36, L09806, doi:10.1029/2009GL037493,
 2067 2009.

2068 Heimann, M. and Reichstein, M.: Terrestrial ecosystem carbon dynamics and climate
 2069 feedbacks, Nature, 451, 289-292 , doi:10.1038/nature06591, 2008.

2070 Heimann, M., Schulze, E.-D., Winderlich, J., Andreae, M.O., Chi, X., Gerbig, C., Kolle, O.,
 2071 Kubler, K., Lavric, J., Mikhailov, E., Panov, A., Park, S., Rodenbeck, C., and Skorochood A.:
 2072 The Zotino Tall Tower Observatory (ZOTTO): Quantifying Large Scale Biogeochemical
 2073 Changes in Central Siberia, Nova Acta Leopoldina, 117, 51-64, 2014.

2074 Heintzenberg, J., Birmili, W., Otto, R., Andreae, M. O., Mayer, J.-C., Chi, X.-G., and Panov,
 2075 A.: Aerosol particle number size distributions and particulate light absorption at the ZOTTO
 2076 tall tower (Siberia), 2006-2009, Atmos. Chem. Phys., 11, 8703-8719, 0.5194/acp-11-8703-
 2077 2011, 2011.

2078 Hemispheric Transport of Air Pollution Working Group (HTAP): Hemispheric transport of
 2079 air pollution: Part A: ozone and particulate matter, in: Air pollution Studies 17, Dentener, F.,
 2080 Keating, T., and Akimoto, H. (Eds.), United Nations, Geneva, 2010.

2081 Hickler, T., Vohland, K., Feehan, J., Miller, P.A., Smith, B., Costa, L., Giesecke, T.,
 2082 Fronzek, S., Carter, T.R., Cramer, W., Kühn, I., and Sykes, M.T.: Projecting the future
 2083 distribution of European potential natural vegetation zones with a generalized, tree species-
 2084 based dynamic vegetation model, *Global Ecology and Biogeography*, 21, 50-63, 2012.

2085 Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, K., Manninen, H. E., Ehn, M., Hörrak, U.,
 2086 Kerminen, V.-M., Laakso, L., McMurry, P. H., Mirme, A., Mirme, S., Petäjä, T., Tammet, H.,
 2087 Vakkari, V., Vana, M., and Kulmala, M.: Atmospheric ions and nucleation: a review of
 2088 observations, *Atmospheric Chemistry and Physics*, 11, 767–798, 2011.

2089 Hollstein, A., Fischer, J., Carbajal Henken, C., and Preusker, R.: Bayesian cloud detection for
 2090 MERIS, AATSR, and their combination, *Atmos. Meas. Tech. Discuss.*, 7, 11045-11085,
 2091 doi:10.5194/amtd-7-11045-2014, 2014.

2092 Hoose, C. and Möhler, O.: Heterogeneous ice nucleation of atmospheric aerosols: a review
 2093 of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817-9854, 2012.

2094 Huffman, G. J., Adler, R. F., Bolvin, D. T., Gu, G. J., Nelkin, E. J., Bowman, K. P., Hong,
 2095 Y., Stocker, E.F., and Wolff, D.B.: The TRMM multisatellite precipitation analysis
 2096 (TMPA): Quasi-global, multiyear, combined-sensor precipitation estimates at fine scales, *J.*
 2097 *Hydrometeorology*, 8, 38-55, 2007.

2098 Hunter, M., Kozlov, M.V., Itämies, J., Pullainen, E., Bäck, J., Kyrö E.-M, and Niemelä, P.:
 2099 Current temporal trends in moth abundance are counter to predicted effects of climate
 2100 change in an assemblage of subarctic forest moths, *Global Change Biology*, 1723-37, doi:
 2101 10.1111/gcb.12529, 2014.

2102 Huotari, J., Ojala, A., Peltomaa, E., Nordbo, A., Launiainen, S., Pumpanen, J., Rasilo, T.,
 2103 Hari, P., and Vesala, T.: Long-term direct CO₂ flux measurements over a boreal lake: Five

2104 years of eddy covariance data, *Geophysical Res. Letters*, 38, L18401, doi:
 2105 10.1029/2011GL048753, 2011.

2106 Huotari, J., Haapanala, S., Pumpanen, J., Vesala, T., and Ojala, A.: Efficient gas exchange
 2107 between a boreal river and the atmosphere, *Geophys. Res. Lett.*, 40, 5683-5686, 2013.

2108 Inst. Health Metrics & Evaluation (IHME): Global burden of diseases, injuries, and risk
 2109 factors study 2010. GBD profile: China, [www.healthmetricsandevaluation.org/gbd/country-](http://www.healthmetricsandevaluation.org/gbd/country-profiles)
 2110 [profiles](http://www.healthmetricsandevaluation.org/gbd/country-profiles), 2013.

2111 IPCC 2013. *Climate Change: The Physical Science Basis*. Working Group I Contribution to
 2112 the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker T.F.,
 2113 Qin D., Plattner G.-K. , Tignor M., Allen S.K., Boschung J., Nauels A., Xia Y., Bex V. &
 2114 Midgley P. M. (eds.). Cambridge University Press, Cambridge, United Kingdom and New
 2115 York, NY, US, 2013.

2116 IPCC 2014. *Climate Change: Impacts, Adaptation, and Vulnerability*. Working Group II
 2117 Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate
 2118 Change, 2014.

2119 Isidorov, V.A.: *Organic chemistry of the atmosphere*, Khimizdat, Moscow, Russia (in
 2120 Russian), 2001.

2121 Jaffe, D. A. and Wigder, N. L.: Ozone production from wildfires: a critical review, *Atmos.*
 2122 *Environ.*, 51, 1-10, 2012.

2123 Jeong, J.-H., Kug, J.-S., Kim, B.-M., Min, S.-K., Linderholm, H. W., Ho, C.-H., Rayner, D.,
 2124 Chen, D., and Jun, S.-Y.: Greening in the circumpolar high-latitude may amplify warming in
 2125 the growing season, *Climate Dynamics*, 38, 1421–143, 2011.

2126 Jolleys, M. D., Coe, H., McFiggans, G., Taylor, J. W., O'Shea, S. J., Le Breton, M.,
 2127 Bauguitte, S. J. B., Moller, S., Di Carlo, P., Aruffo, E., Palmer, P. I., Lee, J. D., Percival, C.
 2128 J., and Gallagher, M. W., Properties and evolution of biomass burning organic aerosol from
 2129 Canadian boreal forest fires: *Atmos. Chem. Phys.*, 15, 3077-3095, doi:10.5194/acp-15-3077-
 2130 2015, 2015.

2131 Jones A. M. and Harrison R. M.: Temporal trends in sulphate concentrations at European sites
 2132 and relationships to sulphur dioxide, *Atmos. Environ.* 45, 873-882, 2011.

2133 Jonsson, M. and Wardle, D.A.: Structural equation modeling reveals plant-community
 2134 drivers of carbon storage in boreal forest ecosystems, *Biology Letters*, 6, 116-119, 2010.

2135 Joyce, R. J., Janowiak, J. E., Arkin, P. A., and Xie, P. P.: CMORPH—A method that
 2136 produces global precipitation estimates from passive microwave and infrared data at high
 2137 spatial and temporal resolution, *J. Hydrometeorology*, 5, 487-503, 2004.

2138 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette,
 2139 J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning
 2140 emissions estimated with a global fire assimilation system based on observed fire radiative
 2141 power, *Biogeosciences*, 9, 527–554, doi:10.5194/bg-9-527-2012, 2012.

2142 Kalinina O., Goryachkin, S.V., Karavaeva, N.A., Lyuri, D.I., Najdenko, L., and Giani, L.:
 2143 Self-restoration of post-agrogenic sandy soils in the southern Taiga of Russia: Soil
 2144 development, nutrient status, and carbon dynamics, *Geoderma*, 152: 35-42, 2009.

2145 Kapsch, M.-L., Graversen, R. G., and Tjernström, M.: Springtime atmospheric energy
 2146 transport and the control of Arctic summer sea-ice extent, *Nature Climate Change*, 3, 744–
 2147 748, doi:10.1038/nclimate1884, 2013.

2148 Karhu, K., Auffret, M. D., Dungait, J. A. J., Hopkins D. W., Prosser J. I., Singh B. K.,
 2149 Subke, J-A., Wookey, P. A., Ågren G. I., Sebastia, M.-T., Gouriveau F., Bergkvist G., Meir
 2150 P., Nottingham A. T., Salina, N., and Hartley I. P.: Temperature sensitivity of soil respiration
 2151 rates enhanced by microbial community response, *Nature*, 513, 81–84,
 2152 doi:10.1038/nature13604, 2014.

2153 Karpoff, B.S. and Roscoe W.E.: Report on Placer Gold Properties in the Tuul Valley,
 2154 Zaamar Goldfield, Mongolia, Roscoe Postle Associates INC, Toronto, Ontario, 1- 66, 2005.

2155 Karthe, D., Kasimov, N., Chalov, S., Shinkareva, G., Malsy, M., Menzel, L., Theuring, P.,
 2156 Hartwig, M., Schweitzer, C., Hofmann, J., Priess, J., and Lychagin, M.: Integrating Multi-
 2157 Scale Data for the Assessment of Water Availability and Quality in the Kharaa -
 2158 Orkhon - Selenga River System, *Geography, Environment, Sustainability*, 3, 65-86, 2014.

2159 Kasischke, E.S.: Boreal ecosystems in the global carbon cycle, in: *Fire, Climate Change and*
 2160 *Carbon Cycling in the Boreal Forest*, Kasischke, E.S., Stocks, B.J. (Eds.), *Ecological*
 2161 *Studies*, 138, 19-30, 2000.

2162 Kay, J. E., L'Ecuyer, T., Gettelman, A., Stephens, G., and O'Dell, C.: The contribution of
 2163 cloud and radiation anomalies to the 2007 Arctic sea ice extent minimum, *Geophys. Res.*
 2164 *Lett.*, 35, L08503, doi:10.1029/2008GL033451, 2008.

2165 Keeling, C. D., Chin, J. F. S., and Whorf, T. P.: Increased activity of northern vegetation
 2166 inferred from atmospheric CO₂ measurements, *Nature*, 382, 146 – 149, 1996.

2167 Khvorostyanov, D. V., Ciais, P., Krinner, G., and Ziv, S. A.: Vulnerability of East Siberia's
 2168 frozen carbon stores to future warming, *Geophys. Res. Lett.*, 35, L10703, 2008.

2169 Kim, B.-M., Son, S.-W. , Min, S.-K., Jeong, J.-H. , Kim, S.-J., Zhang, X., Shim, T., and
2170 Yoon, J.-H.: Weakening of the stratospheric polar vortex by Arctic sea-ice loss, Nat.
2171 Commun., 5, 4646, doi:10.1038/ncomms5646, 2014.

2172 Koch, D. and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and
2173 synthesis, Atmos. Chem. Phys., 10, 7685-7696, 2010.

2174 Kolmonen, P., Sogacheva, L., Virtanen, T.H., de Leeuw, G., and Kulmala, M.: The
2175 ADV/ASV AATSR aerosol retrieval algorithm: current status and presentation of a full-
2176 mission AOD data set, International Journal of Digital Earth, 1-17, DOI:
2177 10.1080/17538947.2015.1111450, 2015.

2178 Kolstad, E.W., Bracegirdle, T.J., and Seierstad, I.A.: Marine cold air outbreaks in the North
2179 Atlantic: temporal distribution and associations with large-scale atmospheric circulation, Clim.
2180 Dyn., 33, 187–197, 2009.

2181 Konovalov, I. B., Berezin, E. V., Ciais, P., Broquet, G., Beekmann, M., Hadji-Lazaro, J.,
2182 Clerbaux, C., Andreae, M. O., Kaiser, J. W., and Schulze, E.-D., Constraining CO₂ emissions
2183 from open biomass burning by satellite observations of co-emitted species: a method and its
2184 application to wildfires in Siberia, Atmos. Chem. Phys., 14, 10383–10410, doi:10.5194/acp-
2185 14-10383-2014, 2014.

2186 Konovalov, I. B., Beekmann, M., Berezin, E. V., Petetin, H., Mielonen, T., Kuznetsova, I.
2187 N., and Andreae, M. O., The role of semi-volatile organic compounds in the mesoscale
2188 evolution of biomass burning aerosol: a modelling case study of the 2010 mega-fire event in
2189 Russia: Atmos. Chem. Phys., 15, 13,269–13,297, doi:10.5194/acp-15-13269-2015, 2015.

2190 Konstantinov, P.I., Varentsov, M.I., and Malinina, E.P.: Modeling of thermal comfort
 2191 conditions inside the urban boundary layer during Moscow's 2010 summer heat wave (case-
 2192 study), *Urban Climate*, 10, 563–572, 2014.
 2193

2194 Konstantinov, P.I., Grishchenko, M.Y., and Varentsov, M. I.: Mapping of Arctic Cities
 2195 Urban Heat Island Based on the Composition of Field Meteorological Measurements and
 2196 Satellite Derived Imagery (Example of Apatity, Kola Peninsula), *Issledovanie Zemli iz*
 2197 *kosmosa, Izvestija, Atmospheric and Oceanic Physics*, 51, doi 10.7868/S0205961415030069
 2198 (in press), 2015.

2199 Korhonen, J. F. J., Pihlatie, M., Pumpanen, J., Aaltonen, H., Hari, P., Levula, J., Kieloaho,
 2200 A.-J., Nikinmaa, E., Vesala, T., and Ilvesniemi, H.: Nitrogen balance of a boreal Scots pine
 2201 forest, *Biogeosciences*, 10, 1083-1095, 2013.

2202 Korontzi, S., McCarty, J., Loboda, T., Kumar, S., and Justice, C.: Global distribution of
 2203 agricultural fires in croplands from 3 years of Moderate Resolution Imaging
 2204 Spectroradiometer (MODIS) data, *Global Biogeochem Cycles*, 20, GB2021,
 2205 doi:10.1029/2005GB002529, 2006.

2206 Kortelainen, P., Rantakari, M., Huttunen, J. T., Mattsson, T., Alm, J., Juutinen, S., Larmola,
 2207 T., Silvola, J., and Marikainen, P.: Sediment respiration and lake trophic state are important
 2208 predictors of large CO₂ evasion from small boreal lakes, *Global Change Biology*, 12, 1554–
 2209 1567, 2006.

2210 Korytnyi, L.M.: Urgent tasks of geographical resources management, in: Vaganov, E.A. et
 2211 al. (Eds.) *Resource Economics, Environmental Economics and Climate Change, Materials of*
 2212 *the International Conference*, July, 1-7, Siberian Federal University, 359-366, 2009.

2213 Kotilainen, J., Tysiachniouk, M., Kuliasova, A., Kuliasov, I., and Pchelkina, S.: The potential
 2214 for ecological modernisation in Russia: scenarios from the forest industry, *Environmental*
 2215 *Politics*, 17, 58-77, 2008.

2216 Kosobokova, K.N.: *Zooplankton of the Arctic Basin*, GEOS, Moscow, 2012.

2217 Kozlova, E. A., Manning, A. C., Kisilyakhov, Y., Seifert, T., and Heimann, M.: Seasonal,
 2218 synoptic, and diurnal-scale variability of biogeochemical trace gases and O₂ from a 300-m
 2219 tall tower in central Siberia, *Global Biogeochem. Cycles*, 22, GB4020,
 2220 doi:10.1029/2008GB003209, 2008.

2221 Kug, J.-S., Jeong, J.-H., Jang, Y.-S., Kim, B.-M., Kim, Folland, C. K., Min, S.-K., and Son,
 2222 S.-W.: Two distinct influences of Arctic warming on cold winters over North America and
 2223 East Asia, *Nature Geosci.*, 8, 759-762, doi: 10.1038/ngeo2517, 2015.

2224 Kukkola, E., Huttunen, S., Bäck, J., and Rautio, P.: Scots pine needle injuries at subarctic
 2225 industrial sites, *Trees*, 11, 378-387, 1997.

2226 Kulmala, L., Aaltonen, H., Berninger F., Kieloaho A.-J., Levula J., Bäck J., Hari P., Kolari
 2227 P., Korhonen J.F.J., Kulmala M., Nikinmaa E., Pihlatie M., Vesala T., and Pumpanen, J.:
 2228 Changes in biogeochemistry and carbon fluxes after the clear-cutting and partial burning of
 2229 slash, *Agric. Forest Meteorol.*, 188, 33-44, 2014.

2230 Kulmala, L., Aaltonen, H., Berninger, F., Kieloaho, A.-J., Levula, J., Bäck, J., Hari, P.,
 2231 Kolari, P., Korhonen, J.F.J., Kulmal,a M., Nikinmaa, E., Pihlatie, M., Vesala, T., and
 2232 Pumpanen, J.: Short term changes in biogeochemistry and the fluxes of carbon in a boreal
 2233 Spruce forest two three after a clear cut and partial burning of slash, *Agric. Forest Meteorol.*,
 2234 188, 33-44, 2014.

2235 Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen,
 2236 P., Hämeri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation
 2237 mode particles, *Tellus B*, 53, 479-490, 2001.

2238 Kulmala, M., Suni, T., Lehtinen, K. E. J., Dal Maso, M., Boy, M., Reissell, A., Rannik, U.,
 2239 Aalto, P., Keronen, P., Hakola, H., Back, J. B., Hoffmann, T., Vesala, T., and Hari, P.: A
 2240 new feedback mechanism linking forests, aerosols, and climate, *Atmos. Chem. Phys.*, 4, 557-
 2241 562, doi:10.5194/acp-4-557-2004, 2004.

2242 Kulmala, M., Laakso, L., Lehtinen, K. E. J., Riipinen, I., Dal Maso, M., Anttila, T., Kerminen,
 2243 V. M., Horrak, U., Vana, M., and Tammet, H.: Initial steps of aerosol growth, *Atmospheric
 2244 Chemistry and Physics*, 4, 2553–2560, 2004.

2245 Kulmala, M., Asmi, A., Lappalainen, H. K., Carslaw, K. S., Pöschl, U., Baltensperger, U.,
 2246 Hov, Ø., Brenquier, J.-L., Pandis, S. N., Facchini, M. C., Hansson, H.-C., Wiedensohler, A.,
 2247 and O'Dowd, C. D.: Introduction: European Integrated project on Aerosol Cloud Climate and
 2248 Air Quality interactions (EUCAARI) - integrating aerosol research from nano to global
 2249 scales, *Atmos. Chem. Phys.*, 9, 2825-2841, 2009.

2250 Kulmala, M., Alekseychik, P., Paramonov, M., Laurila, T., Asmi, E., Arneth, A.,
 2251 Zilitinkevich, S., and Kerminen, V.-M.: On measurements of aerosol particles and
 2252 greenhouse gases in Siberia and future research needs, *Boreal Env. Res.*, 16, 337-362, 2011a.

2253 Kulmala, M., Asmi, A., Lappalainen, H. K., Baltensperger, U., Brenguier, J.-L., Facchini, M.
 2254 C., Hansson, H.-C., Hov, Ø., O'Dowd, C. D., Pöschl, U., Wiedensohler, A., Boers, R.,
 2255 Boucher, O., de Leeuw, G., Denier van der Gon, H. A. C., Feichter, J., Krejci, R., Laj, P.,
 2256 Lihavainen, H., Lohmann, U., McFiggans, G., Mentel, T., Pilinis, C., Riipinen, I., Schulz,
 2257 M., Stohl, A., Swietlicki, E., Vignati, E., Alves, C., Amann, M., Ammann, M., Arabas, S.,

2258 Artaxo, P., Baars, H., Beddows, D. C. S., Bergström, R., Beukes, J. P., Bilde, M., Burkhardt,
 2259 J. F., Canonaco, F., Clegg, S. L., Coe, H., Crumeyrolle, S., D'Anna, B., Decesari, S.,
 2260 Gilardoni, S., Fischer, M., Fjaeraa, A. M., Fountoukis, C., George, C., Gomes, L., Halloran,
 2261 P., Hamburger, T., Harrison, R. M., Herrmann, H., Hoffmann, T., Hoose, C., Hu, M.,
 2262 Hyvärinen, A., Hörrak, U., Iinuma, Y., Iversen, T., Josipovic, M., Kanakidou, M., Kiendler-
 2263 Scharr, A., Kirkevåg, A., Kiss, G., Klimont, Z., Kolmonen, P., Komppula, M., Kristjánsson,
 2264 J.-E., Laakso, L., Laaksonen, A., Labonnote, L., Lanz, V. A., Lehtinen, K. E. J., Rizzo, L.
 2265 V., Makkonen, R., Manninen, H. E., McMeeking, G., Merikanto, J., Minikin, A., Mirme, S.,
 2266 Morgan, W. T., Nemitz, E., O'Donnell, D., Panwar, T. S., Pawlowska, H., Petzold, A.,
 2267 Pienaar, J. J., Pio, C., Plass-Duelmer, C., Prévôt, A. S. H., Pryor, S., Reddington, C. L.,
 2268 Roberts, G., Rosenfeld, D., Schwarz, J., Seland, Ø., Sellegri, K., Shen, X. J., Shiraiwa, M.,
 2269 Siebert, H., Sierau, B., Simpson, D., Sun, J. Y., Topping, D., Tunved, P., Vaattovaara, P.,
 2270 Vakkari, V., Veefkind, J. P., Visschedijk, A., Vuollekoski, H., Vuolo, R., Wehner, B., Wildt,
 2271 J., Woodward, S., Worsnop, D. R., van Zadelhoff, G.-J., Zardini, A. A., Zhang, K., van Zyl,
 2272 P. G., Kerminen, V.-M., Carslaw, K. S. and Pandis, S. N.: General overview: European
 2273 Integrated project on Aerosol Cloud Climate and Air Quality interactions (EUCAARI) –
 2274 integrating aerosol research from nano to global scales, *Atmos. Chem. Phys.*, 11, 13061-
 2275 13143, 2011b.

 2276 Kulmala, M. and Petäjä, T.: Soil nitrites influence atmospheric chemistry, *Science*, 333,
 2277 1586-1587, 2011.

 2278 Kulmala, M., Nieminen, T., Chellapermal, R., Makkonen, R., Bäck, J., and Kerminen, V.-
 2279 M.: Climate feedbacks linking the increasing atmospheric CO₂ concentration, BVOC
 2280 emissions, aerosols and clouds in forest ecosystems. In: *Biology, Controls and Models of*
 2281 *Tree Volatile Organic Compound Emissions*, Ü Niinemets and R. K Monson (Eds.),
 2282 Springer, 489-508, 2013.

2283 Kulmala, M., Nieminen, T., Nikandrova, A., Lehtipalo, K., Manninen, H. E., Kajos, M. K.,
 2284 Kolari, P., Lauri, A., Petäjä, T., Krejci, R., Hansson, H.-C., Swietlicki, E., Lindroth, A.,
 2285 Christensen, T. R., Arneth, A., Hari, P., Bäck, J., Vesala, T., and Kerminen, V.-M.: CO₂-
 2286 induced terrestrial climate feedback mechanism: From carbon sink to aerosol source and
 2287 back, *Boreal Env. Res.*, 19, Suppl. B, 122-131, 2014.

2288 Kulmala, M., Lappalainen, H.K., Petäjä, T., Kurten, T., Kerminen, V-M., Viisanen, Y., Hari,
 2289 P., Bondur, V., Kasimov, N., Kotlyakov, V., Matvienko, G., Baklanov, A., Guo, H., Ding,
 2290 A., Hansson, H-C., and Zilitinkevich, S.: Introduction: The Pan-Eurasian Experiment (PEEX)
 2291 – multi-disciplinary, multi-scale and multi-component research and capacity building
 2292 initiative, *Atmos. Chem. Phys.*, 15, 13085-13096, doi:10.5194/acp-15-13085-2015, 2015.

2293 Kulmala, M.: China's choking cocktail, *Nature*, 526, 497, 2015.
 2294

2295 Kulmala, M., Lappalainen, H.K., Petäjä, T., Kerminen, V.M., Viisanen, Y., Matvienko,
 2296 G., Melnikov, V., Baklanov, A., Bondur, V., Kasimov, N., and Zilitinkevich, Z.: Pan-
 2297 Eurasian Experiment (PEEX) Program: Grant Challenges in the Arctic-boreal context,
 2298 *J.Geography, Environment and Sustainability*, 2, 5-18., 2016.

2299 Kummerow, C., Hong, Y., Oleson, W. S., Yang, S., Adler, R. F., Mccollum, J., Ferraro, R.,
 2300 Petty, G., Shin, D.B., and Wilheit, T.T.: The evolution of the Goddard profiling algorithm
 2301 (GPROF) for rainfall estimation from passive microwave sensors, *J. Applied Meteorology*,
 2302 40, 1801-1820, 2001.

2303 Kurz, W.A. and Apps, M.J.: An analysis of future carbon budgets of Canadian boreal forests,
 2304 *Water Air Soil Pollut.*, 82, 321-331, 1995.

2305 Kurz, W. A., Dymond, C. C., Stinson, G., Rampley, G. J., Neilson, E. T., Carroll, A. L.,
2306 Ebata, T., and Safranyik, L.: Mountain pine beetle and forest carbon feedback to climate
2307 change, *Nature*, 452, 987-990, 2008.

2308 Kurz, W. A., Stinson, G., Rampley, G. J., Dymond, C. C., and Neilson, E. T.: Risk of natural
2309 disturbances makes future contribution of Canada's forests to the global carbon cycle highly
2310 uncertain, *PNAS*, 105, 1551–1555, doi10.1073pnas.0708133105, 2008.

2311 Kuzyakov, Y.: Priming effects: Interactions between living and dead organic matter, *Soil*
2312 *Biology and Biochemistry*, 42, 1363–1371, 2010.

2313 Kvaeven, B., Ulstein, M. J., and Skjelkvåle, B. L.: ICP Waters - An international programme
2314 for surface water monitoring, *WaterAir Soil Pollut.*, 130, 775-780, 2001.

2315 Kwok, R. and Rothrock, D. A.: Decline in Arctic sea ice thickness from submarine and
2316 ICESat records: 1958–2008, *Geophys. Res. Lett.*, 36, L15501, doi:10.1029/2009GL039035,
2317 2009.

2318 Köster, K., Berninger, F., Lindén, A., and Pumpanen, J.: Recovery in fungal biomass is
2319 related to decrease in soil organic matter turnover time in a boreal fire chronosequence,
2320 *Geoderma*, 235–236, 74–82, 2014.

2321 Laaksonen, A. A.: Unifying model for adsorption and nucleation of vapors on solid surfaces,
2322 *J. Phys. Chem. A*, 119, 3736-3745, 2015.

2323 Laaksonen, A. and Malila, J.: An adsorption theory of heterogeneous nucleation of water
2324 vapour on nanoparticles, *Atmos. Chem. Phys.* 16, 135-143, doi:10.5194/acp-16-135-2016,
2325 2016.

2326 de Laat, A. T. J., van der A, R.J. and van Weele, M.: Evaluation of tropospheric ozone
 2327 columns derived from assimilated GOME ozone profile observations, *Atmos. Chem. Phys.*, 9,
 2328 8105–8120, 2009.

2329 Lappalainen, H.K., Petäjä, T., Kujansuu, J., Kerminen, V-M., Shvidenko, A., Bäck, J.,
 2330 Vesala, T., Vihma, T., de Leeuw, G., Lauri, A., Ruuskanen, T., Flint, M., Zaitseva, N.,
 2331 Arshinov, M., Spracklen, D., Arnold, S., Juhola, S., Lihavainen, H., Viisanen, Y.,
 2332 Chubarova, N., Filatov, N., Skorokhod, A., Elansky, N., Dyukarev, E., Hari, P., Kotlyakov,
 2333 V., Kasimov, N., Bondur, V., Matvienko, G., Baklanov, A., Guo H., Zilitinkevich, S., and
 2334 Kulmala, M.: Pan-Eurasian Experiment (PEEX) – a research initiative meeting the grand
 2335 challenges of the changing environment of the northern Pan-Eurasian arctic-boreal areas, *J.*
 2336 *Geography Environment Sustainability*, 2, 13-48, 2014.

2337 Laothawornkitkul, J., Taylor, J. E., Paul, N. D., and Hewitt, C. N.: Biogenic volatile organic
 2338 compounds in the Earth system, *New Phytologist*, 183, 27-51, 2009.

2339 Lenhart, K., Weber, B., Elbert, W., Steinkamp, J., Clough, T., Crutzen, P., Pöschl, U., and
 2340 Keppler, F.: Nitrous oxide and methane emissions from cryptogamic covers, *Global Change*
 2341 *Biology*, 21, 3889-3900, 10.1111/gcb.12995, 2015.

2342 Liang, Q., Rodriguez, J. M., Douglass, A. R., Crawford, J. H., Olson, J. R., Apel, E., Bian,
 2343 H., Blake, D. R., Brune, W., Chin, M., Colarco, P. R., da Silva, A., Diskin, G. S., Duncan, B.
 2344 N., Huey, L. G., Knapp, D. J., Montzka, D. D., Nielsen, J. E., Pawson, S., Riemer, D. D.,
 2345 Weinheimer, A. J., and Wisthaler, A., Reactive nitrogen, ozone and ozone production in the
 2346 Arctic troposphere and the impact of stratosphere-troposphere exchange: *Atmos. Chem.*
 2347 *Phys.*, 11, 13181-13199, doi:10.5194/acp-11-13181-2011, 2011.

2348 Lihavainen, H., Kerminen, V.-M., Tunved, P., Aaltonen, V., Arola, A., Hatakka, J.,
 2349 Hyvärinen, A., and Viisanen, Y.: Observational signature of the direct radiative effect by
 2350 natural boreal forest aerosols and its relation to the corresponding first indirect effect, *J.*
 2351 *Geophys. Res.*, 114, D20206, doi:10.1029/2009JD012078, 2009.

2352 Lihavainen, H., Asmi, E., Aaltonen, V., Makkonen U., and Kerminen, V.-M.: Direct
 2353 radiative feedback due to biogenic secondary organic aerosol estimated from boreal forest
 2354 site observations, *Environ. Res. Lett.*, 10, 104005, 2015.

2355 Liston, G., McFadden, J., Sturm, M., and Sr., R. P.: Modelled changes in arctic tundra snow,
 2356 energy and moisture fluxes due to increased shrubs, *Global Change Biology*, 8, 17-32, 2002.

2357 Litvinenko, T.V.: Socioecological Consequences of Transformation of Natural Resources
 2358 Utilization in Russia's Eastern Part in Post-Soviet Period, *Regional Research of Russia*, 2,
 2359 284-295, 2012.

2360 Litvinenko, T.V.: Post-Soviet Transformation of Natural Resources Utilization and its
 2361 Impact on Population Dynamics in Chukotka Autonomous Okrug, *Izvestiya of Russian*
 2362 *Academy of Sciences, Geography*, 2, 30-42, (in Russian), 2013.

2363 Liu, Y.Y., Parinussa, R.M., Dorigo, W.A., De Jeu, R. A. M. , W. Wagner W., van Dijk, A. I.
 2364 J. M., McCabe, M.F., and Evans, J.P.: Developing an improved soil moisture dataset by
 2365 blending passive and active microwave satellite-based retrievals, *Hydrology and Earth*
 2366 *System Sciences*, 15, 425 - 436, 2011.

2367 Lloyd, A. H. and Bunn, A. G.: Responses of the circumpolar boreal forest to 20th century
 2368 climate variability, *Environmental Research Letters*, 2, 045013, doi:10.1088/1748-
 2369 9326/2/4/045013, 2007.

2370 Lu, J. and Cai, M.: Quantifying contributions to polar warming amplification in an idealized
 2371 coupled general circulation model, *Clim Dy*, 34, 669-687, 2010.

2372 Lyuri, D.I., Goryachkin, S.V., Karavaeva, N.A., Denisenko, E.A., and Nefedova, T.G.:
 2373 Dynamics of agricultural lands in Russia in 20th century and post-agrogenic progradation of
 2374 vegetation and soils, GEOS, Moscow, Russia (in Russian), 2010.

2375 Magnani, F., Mencuccini, M., Borghetti, M., Berbigier, P., Berninger, F., Delzon, S., Grelle,
 2376 A., Hari, P., Jarvis, P.G., Kolari, P., Kowalski, A.S., Lankreijer, H., Law, B.E., Lindroth, A.,
 2377 Loustau, D., Manca, G., Moncrieff, J.B., Rayment, M., Tedeschi, V., Valentini, R., and
 2378 Grace, J.: The human footprint in the carbon cycle of temperate and boreal forests, *Nature*,
 2379 447, 848-850, 2007.

2380 Makshtas, A., Nedashkovsky, A., P., and Uttal, T.: The role of the Arctic sea ice in carbon
 2381 dioxide exchange, *AMS Conf. on Polar Meteor. Ocean*, 3 May, 2011.

2382 Maksimovich, E. and Vihma, T.: The effect of surface heat fluxes on interannual variability
 2383 in the spring onset of snow melt in the central Arctic Ocean, *J. Geophys. Res.*, 117,
 2384 C07012, doi:10.1029/2011JC007220, 2012.

2385 Malevsky-Malevich, S.P., Molkentin, E.K., Nadyozhina, E.D., and Shklyarevich, O.B.: An
 2386 assessment of potential change in wildfire activity in the Russian boreal forest zone induced
 2387 by climate warming during the twenty-first century, *Climatic Change*, 86, 463-474, 2008.

2388 Malin, G., Turner, S., Liss, P., Holligan, P., and Harbour, D.: Dimethylsulphide and
 2389 dimethylsulphoniopropionate in the Northeast Atlantic during the summer coccolithophore
 2390 bloom, *Deep Sea Research Part I: Oceanographic Research Papers*, 40, 1487-1508, 1993.

2391 Maljanen, M., Sigurgsson, B., Guðmundsson, J., Óskarsson, H., Huttunen, J., and
 2392 Martikainen, P.: Greenhouse gas balances of managed peatlands in the Nordic countries -
 2393 present knowledge and gaps, *Biogeosciences*, 7, 2711-2738, 2010.

2394 Malkhazova, S. M., Mironova, V. V., Kotova, T. V., Shartova N. V., and Orlov D. S.:
 2395 Natural Focal Diseases in Russia: Monitoring and Mapping, *Geography Environment*
 2396 *Sustainability*, 4, 4-12, 2013.

2397 Mann, P. J., Davydova, A., Zimov, N., Spencer, R., Davydov, S., Bulygina, E., Zimov, S.,
 2398 and Holmes, R.: Controls on the composition and lability of dissolved organic matter in
 2399 Siberia's Kolyma River basin, *J. Geophys. Res.* 117, G01028, 2012.

2400 Mareev, E.: Global electric circuit research: achievements and prospects, *Uspekhi*
 2401 *Fizicheskikh Nauk and P N Lebedev Physics Institute of the Russian Academy of Sciences*,
 2402 *Physics-Uspekhi*, 53, 504- 511, DOI: 10.3367/UFNe.0180.201005h.052, 2010.

2403 Mauldin, R.L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T.,
 2404 Stratmann, F., Kerminen, V.-M., and Kulmala, M.: A new atmospherically relevant oxidant
 2405 of sulphur dioxide, *Nature*, 488, 193-196, 2012.

2406 McComiskey, A., and Feingold, G.: The scale problem in quantifying aerosol indirect
 2407 effects, *Atmos. Chem. Phys.*, 12, 1031-1049, doi:10.5194/acp-12-1031-2012, 2012.

2408 Menzel, A., Sparks, T.H., Estrella, N., Koch, E., Aasa, A., Aha, R., Alm-Kubler, K., Bissolli,
 2409 P., Braslavska, O., Briede, A., Chmielewski, F.M., Crepinsek, Z., Curnel, Y., Dahl, Å.,
 2410 Defila, C., Donnelly, A., Filella, Y., Jatca, K., Måge, F., Mestre, A., Nordli, O., Penuelas,
 2411 J., Pirinen, P., Remisova, V., Scheifinger, H., Striz, M., Susni, A., Van Vliet, A.J.H.,
 2412 Wielgolaski, F-E., Zach, S., and Züst, A.: European phenological response to climate change
 2413 matches the warming pattern, *Global Change Biol.*, 12, 1969-1976, 2006.

2414 Meinander, O., Kazadzis, S., Arola, A., Riihelä, A., Räisänen, P., Kivi, R., Kontu, A.,
 2415 Kouznetsov, R., Sofiev, M., Svensson, J., Suokanerva, H., Aaltonen, V., Manninen, T.,
 2416 Roujean, J.-L., and Hautecoeur, O.: Spectral albedo of seasonal snow during intensive melt
 2417 period at Sodankylä, beyond the Arctic Circle, *Atmos. Chem. Phys.*, 13, 3793-3810,
 2418 doi:10.5194/acp-13-3793-2013, 2013.

2419 Mijling, B., van der R.J., and Zhang, Q.: Regional nitrogen oxides emission trends in East
 2420 Asia observed from space, *Atmos. Chem. Phys.*, 13, 12003-12012, doi:10.5194/acp-13-
 2421 12003-2013, 2013.

2422 Mikhailov, E. F., Mironov, G. N., Pöhlker, C., Chi, X., Krüger, M. L., Shiraiwa, M., Förster,
 2423 J.-D., Pöschl, U., Vlasenko, S. S., Ryshkevich, T. I., Weigand, M., Kilcoyne, A. L. D., and
 2424 Andreae, M. O., Chemical composition, microstructure, and hygroscopic properties of
 2425 aerosol particles at the Zotino Tall Tower Observatory (ZOTTO), Siberia, during a summer
 2426 campaign: *Atmos. Chem. Phys.*, 15, 8847–8869, doi:10.5194/acp-15-8847-2015, 2015a.

2427 Mikhailov, E. F., Mironova, S. Y., Makarova, M. V., Vlasenko, S. S., Ryshkevich, T. I.,
 2428 Panov, A. V., and Andreae, M. O., Studying seasonal variations in carbonaceous aerosol
 2429 particles in the atmosphere over Central Siberia: *Izvestija, Atmos. Ocean. Phys.*, 51, 423-
 2430 430, doi:10.1134/S000143381504009X, 2015b.

2431 Moiseenko, T.I., Kydrjavzeva, L.P., and Sandimirov, S.S.: Principles and methods of water
 2432 quality studies for airborne polluted water bodies: case study of Kola Subarctic, *Water Res.*,
 2433 27, 81-86, 2001.

2434 Monks, S. A., Arnold, S. R., Emmons, L. K., Law, K. S., Turquety, S., Duncan, B. N.,
 2435 Flemming, J., Huijnen, V., Tilmes, S., Langner, J., Mao, J., Long, Y., Thomas, J. L.,
 2436 Steenrod, S. D., Raut, J. C., Wilson, C., Chipperfield, M. P., Diskin, G. S., Weinheimer, A.,

2437 Schlager, H., and Ancellet, G.: Multi-model study of chemical and physical controls on
 2438 transport of anthropogenic and biomass burning pollution to the Arctic, *Atmos. Chem. Phys.*,
 2439 15, 3575-3603, 2015.

2440 Moore, S. K., Trainer, V. L., Mantua, N. J., Parker, M. S., Laws, E. A., Bacher, L. C., and
 2441 Fleming, L. E.: Impacts of climate variability and future climate change on harmful algal
 2442 blooms and human health, *Environmental Health* 7, S4. doi: 10.1186/1476-069x-7-s2-s4,
 2443 2008.

2444 Mori, M., Watanabe, M., Shiogama, H., Inoue, J., and Kimoto, M.: Robust Arctic sea-ice
 2445 influence on the frequent Eurasian cold winters in past decades, *Nat. Geosci.*, 7, 869–873,
 2446 doi:10.1038/NGEO2277, 2014.

2447 Moseley, C. (ed.): *Atlas of the World's languages in danger*. Paris: UNESCO Publishing,
 2448 2010. <http://www.unesco.org/languages-atlas/>).

2449 Mu, Q., Heinsch, F. A., Zhao, M., and Running, S. W.: Development of a global
 2450 evapotranspiration algorithm based on MODIS and global meteorology data, *Remote*
 2451 *Sensing of Environment*, 111, 519-536, 2007.

2452 Mukhrtova, L., Schepaschenko, D., Shvidenko, A., McCallum, I., and Kraxner, F.: Soil
 2453 contribution to carbon budget of Russian forests, *Agricultural and Forest Meteorology*, 200,
 2454 97-118, 2015.

2455 Müller, B., Lotter, A F., and Sturm, A. A.: Influence of catchment quality and altitude on the
 2456 water and sediment composition of 68 small lakes in Central Europe, *Aquat. Sci., Research*
 2457 *Across Boundaries*, 60, 316-337, 1998.

2458 Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G., and Nemani, R. R.: Increased plant
 2459 growth in the northern high latitudes from 1981 to 1991, *Nature*, 386, 698–702, 1997.

2460 Mäkelä, J. M., Aalto, P., Pohja, T., Nissinen, A., Palmroth, S., Markkanen, T., and Kulmala,
 2461 M.: Observations of ultrafine aerosol particle formation and growth in boreal forest,
 2462 Geophysical Research Letters, 24, 1219-1222, 1997.

2463 Naeger, A. R., Gupta, P., Zavodsky, B., and McGrath, K. M.: Monitoring and tracking the
 2464 trans-Pacific transport of aerosols using multi-satellite aerosol optical depth retrievals,
 2465 Atmos. Meas. Tech. Discuss., 8, 10319-10360, doi:10.5194/amtd-8-10319-2015, 2015.

2466 Nédélec, P., Thouret, V., Brioude, J., Sauvage, B., Cammas, J. P., and Stohl, A.: Extreme
 2467 CO concentrations in the upper troposphere over northeast Asia in June 2003 from the in situ
 2468 MOZAIC aircraft data, Geophysical Research Letters, 32, doi:10.1029/2005GL023141,
 2469 2005.

2470 Nelson, D. L., Garay, M. J., Kahn, R. A., and Dunst, B. A.: Stereoscopic Height and Wind
 2471 Retrievals for Aerosol Plumes with the MISR INteractive eXplorer (MINX), Remote Sens.,
 2472 5, 4593– 4628, doi:10.3390/rs5094593, 2013.

2473

2474 Nie, W., Ding, A., Wang, T., Kerminen, V-M., George, C., Xue, L., Wang, W., Zhang, Q.,
 2475 Petäjä, T., Qi, X., Gao, X., Wang, X., Yang, X., Fu, C., and Kulmala, M.: Polluted dust
 2476 promotes new particle formation and growth, Scientific Reports, 4, 6634,
 2477 doi:10.1038/srep06634, 2014.

2478 Nöjd, P. and Kauppi, P.: Growth of Scots pine in a changing environment. In: Tikkanen E,
 2479 Niemela, I. (Eds) Kola Peninsula pollutants and forest ecosystems in Lapland. Final report of
 2480 the Lapland Forest Damage Project. Finland's Ministry of Agriculture and Forestry, The
 2481 Finnish Forest Research Institute, Gummerus Kirjapaino Oy, Jyväskylä, 61-64, 1995.

2482 O'Dowd, C. D., Facchini, M.C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi,
 2483 S., Yoon, Y. J., and Putaud, J.-P.: Biogenically driven organic contribution to marine
 2484 aerosol, *Nature*, 431, 676–680, doi:10.1038/nature02959, 2004.

2485 O'Dowd, C.D. and de Leeuw, G.: Marine Aerosol Production: a review of the current
 2486 knowledge, *Phil. Trans. R. Soc. A* , 365, 1753 - 1774, doi:10.1098/rsta.2007.2043, 2007.

2487 Osava, A., Zyryanova, O.A., Matsuura, Y., Kajamoto, T., and Wein, R.W. (eds): *Permafrost*
 2488 *ecosystems*, Springer, 502 pp., 2009.

2489 Outten, S., Davy, R., and Esau, I.: Eurasian winter cooling: Intercomparison of Reanalyses
 2490 and CMIP5 data sets, *Atmospheric and Oceanic Science Letters*, 6, 324-331,
 2491 doi:10.3878/j.issn.1674-2834.12.0112, 2013.

2492 Overland, J., Francis, J., Hall, R., Hanna, E., Kim, S.-J., and Vihma, T.: The Melting Arctic
 2493 and Mid-latitude Weather Patterns: Are They Connected?, *J. Clim.*, published online,
 2494 <http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-14-00822.1>, 2015.

2495 Overland, J.E. and Wang, M.: Large-scale atmospheric circulation changes are associated
 2496 with the recent loss of Arctic sea ice, *Tellus*, 62, DOI: 10.1111/j.1600-0870.2009.00421.x
 2497 2009, 2010.

2498 Paasonen, P., Asmi, A., Petäjä, T., Kajos, M. K., Äijälä, M., Junninen, H., Holst, T., Abbatt,
 2499 J. P. D., Arneth, A., Birmili, W., Denier van der Gon, H., Hamed, A., Hoffer, A., Laakso, L.,
 2500 Laaksonen, A., Leaitch, W. R., Plass-Dulmer, C., Pryor, S. C., Räisänen, P., Swietlicki, E.,
 2501 Wiedensohler, A., Worsnop, D. R., Kerminen, V.-M., and Kulmala, M.: Evidence for
 2502 negative climate feedback: warming increases aerosol number concentrations, *Nature*
 2503 *Geosci.*, 6, 438–442, doi:10.1038/ngeo1800, 2013.

2504 Paatero, J., Dauvalter, V., Derome, J., Lehto, J., Pasanen, J., Vesala, T., Miettinen, J.,
 2505 Makkonen, U., Kyrö, E-M., Jernström, J., Isaeva, L., and Derome, K.: Effects of Kola air
 2506 pollution on the environment in the western part of the Kola Peninsula and Finnish Lapland -
 2507 Final report, Finnish Meteorological Institute Rep., 6, 2008.

2508 Paerl, H. W. and Huisman, J.: Climate change: a catalyst for global expansion of harmful
 2509 cyanobacterial blooms, *Environmental Microbiology Reports*, doi:10.1111/j.1758-
 2510 2229.2008.0004.x, 2009.

2511 Palmer, T. and Slingo, J.: Uncertainty in weather and climate prediction, *Phil. Trans. R. Soc.*
 2512 *A*, 369, 4751-4767, doi:10.1098/rsta.2011.0161, 2011.

2513 Pan, Y., Birdsey R.A., Fang, J., Houghton, R., Kauppi, P.E., Kurz, W.A., Phillips, O.I.,
 2514 Shvidenko, A., Lewis, S.L., Canadell, J.G., Ciais, P., Jackson, R.B., Pacala, S.W., McGuire,
 2515 A.D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D. A.: Large and Persistent Carbon Sink
 2516 in the World's Forests, *Science*, 333, 988-993, 2011.

2517 Paris, J.-D., Ciais, P., Nedelec, P., Ramonet, M., Golytsin, G., Granberg, I., Athier, G.,
 2518 Boumard, F., Cousin, J.-M., Cayez, G., and Stohl, A.: The YAK-AEROSIB transcontinental
 2519 aircraft campaigns: new insights on the transport of CO₂, CO and O₃ across Siberia, *Tellus*
 2520 *B*, 60, 551-568, 2008.

2521 Paris, J.-D., Ciais, P., Nedelec, P., Stohl, A., Belan, B. D., Arshinov, M. Y., , Carouge, C.,
 2522 Golitsyn, G. S., and Granberg, I. G.: New Insights on the Chemical Composition of the
 2523 Siberian Air Shed from the YAK-AEROSIB Aircraft Campaigns, *B. Am. Meteorol. Soc.*, 91,
 2524 625–641, 2010.a

2525 Paris, J. D., Stohl, A., Ciais, P., Nedelec, P., Belan, B. D., Arshinov, M. Y., and Ramonet,
 2526 M.: Source-receptor relationships for airborne measurements of CO₂, CO and O₃ above
 2527 Siberia: a cluster-based approach: *Atmos. Chem. Phys.*, 10, 1671-1687, 2010b.
 2528
 2529 Parmentier, F. J. W., van der Molen, M.K., van Huissteden, J., Karsanaev, S.A., Kononov,
 2530 S.V., Suzdalov, D.A., Maximov, T.C., and Dolman, A.J.: Longer growing seasons do not
 2531 increase net carbon uptake in the northeastern Siberian tundra, *J. Geophys. Res.*, 116,
 2532 G04013, doi:10.1029/2011JG001653, 2011.
 2533 Parmentier, F.-J. W., Christensen, T.R., Sørensen, L.L., Rysgaard, S., McGuire, A.D., Miller
 2534 P.A., and Walker, D.A.: The impact of lower sea-ice extent on arctic greenhouse-gas
 2535 exchange, *Nature Clim. Change*, 3, 195–202, 2013.
 2536 Pechony, O. and Shindell, D.: Driving forces of global wildfires over the past millenium and
 2537 forthcoming century, *PNAS*, 107, 19167-19170, 2010.
 2538 Penner, J. E., Zhou C., and Xu, L.: Consistent estimates from satellites and models for the
 2539 first aerosol indirect forcing, *Geophys. Res. Lett.*, 39, L13810, doi:10.1020/2012GL051870,
 2540 2012.
 2541 Perovich, D. K., Richter-Menge, J. A., Jones, K. F., and Light, B.: Sunlight, water, and ice:
 2542 Extreme Arctic sea ice melt during the summer of 2007, *Geophys. Res. Lett.*, 35, L11501,
 2543 doi:10.1029/2008GL034007, 2008.
 2544 Persad, G. G., Ming, Y., and Ramaswamy, V.: Tropical troposphere-only response to
 2545 absorbing aerosols, *J. Climate*, 25, 2471-2480, 2012.

2546 Petäjä, T., Järvi, L., Kerminen, V.-M., Ding, A. J., Sun, J. N., Nie, W., Kujansuu, J.,
 2547 Virkkula, A., Yang, X.-Q., Fu, C. B., Zilitinkevich, S., and Kulmala, M.: Enhanced air
 2548 pollution via aerosol-boundary layer feedback in China, *Scientific Reports* 6, 18998,
 2549 doi:10.1038/srep18998, 2016.

2550 Phillips, R.P., Finzi, A.C., and Bernhardt, E.S.: Enhanced root exudation induces microbial
 2551 feedbacks to N cycling in a pine forest under long-term CO₂ fumigation, *Ecology Letters*, 14,
 2552 187-194, 2011.

2553 Piao, S., Ciais, P., Friedlingstein, P., Peylin, P., Reichstein, M., Luyssaert, S., Margolis, H.,
 2554 Fang, J., Barr, A., Chen, A., Grelle, A., Hollinger, D.Y., Laurila, T., Lindroth, A.,
 2555 Richardson, A.D., and Vesala, T.: Net carbon dioxide losses of northern ecosystems in
 2556 response to autumn warming, *Nature*, 451, 49-53, 2008.

2557 Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in
 2558 contemporary climate models, *Nature Geoscience*, 7, 181–184, doi:10.1038/ngeo2071, 2014.

2559 Phoenix, G.K. and J.W. Bjerke, J.W.: Arctic browning: extreme events and trends reversing
 2560 arctic greening, *Global Change Biology*, 22, 2960–2962, 2016.

2561 Polyakov, I. V., Timokhov, L. A., Alexeev, V. A., Bacon, S., Dmitrenko, I. A., Fortier, L.,
 2562 and Toole, J.: Arctic Ocean warming contributes to reduced polar ice cap, *J. Phys.*
 2563 *Oceanogr.*, 40, 2743–2756, 2010.

2564 Pommier, M., Law, K. S., Clerbaux, C., Turquety, S., Hurtmans, D., Hadji-Lazaro, J.,
 2565 Coheur, P.-F., Schlager, H., Ancellet, G., Paris, J.-D., Nédélec, P., Diskin, G. S., Podolske, J.
 2566 R., Holloway, J. S., and Bernath, P.: IASI carbon monoxide validation over the Arctic during
 2567 POLARCAT spring and summer campaigns, *Atmos. Chem. Phys.*, 10, 10655-10678,
 2568 doi:10.5194/acp-10-10655-2010, 2010.

2569 Post, W. M., Emanuel, W. R., Zinke, P. J., and Stangenberger, A. G.: Soil carbon pools and
 2570 world life zones, *Nature*, 298, 156-159, 1982.

2571 Pumpanen, J., Linden, A., Miettinen, H., Kolari, P., Ilvesniemi, H., Mammarella, I., Hari,
 2572 P., Nikinmaa, E., Heinonsalo, J., Bäck, J., Ojala, A., Berninger, F., and Vesala, T.:
 2573 Precipitation and net ecosystem exchange are the most important drivers of DOC flux in
 2574 upland boreal catchments, *J. Geophys. Res. – Biogeosciences* DOI: 10.1002/2014JG002705,
 2575 2014.

2576 Quinn, P.K. and Bates, T.S.: The case against climate regulation via oceanic phytoplankton
 2577 sulphur emissions, *Nature*, 480, 51-56, 2011.

2578 Ramonet, M., Ciais, P., Nepomniachii, I., Sidorov, K., Neubert, R.E.M., Picard, D., Kazan,
 2579 V., Biraud, S.C., Gusti, M., Schulze, E.D., and Lloyd, J.: Three years of aircraft based trace
 2580 gas measurements over Fyodoroskye southern taiga forest, 300 km North-West of Moscow,
 2581 *Tellus B*, 54, 713-734, 2002.

2582 Randers, J.: A Global Forecast for the Next Forty Years - 2052. A Report to the Club of
 2583 Rome Commemorating the 40th Anniversary of The Limits to Growth. Editor Joni Praded.
 2584 Chelsea Green Publishing. 395 pp. 2012.

2585 Rampal, P., Weiss, J., Dubois, C., and Campin, J. M.: IPCC climate models do not capture
 2586 Arctic sea ice drift acceleration: Consequences in terms of projected sea ice thinning and
 2587 decline, *J. Geophys. Res.*, 116, C00D07, 2011.

2588 Randersson, J. T., Liu, H., Flanner, M. G., Chambers, S. D., Jin, Y., Hess, P. G., Pfister, G.,
 2589 Mack, M. C., Treseder, K. K., Welp, L. R., Chapin, F. S., Harden, J. W., Goulden, M. L.,
 2590 Lyons, E., Neff, J. C., Schuur, E. A. G., and Zender, C. S.: The impact of boreal forest fire
 2591 on climate warming, *Science*, 314, 1130-1132, 2006.

2592 Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F.T., Gruber, N., Janssens, I.A., Laruelle,
 2593 G.G., Lauerwald, R., Luyssaert, S., Andersson, A.J., Arndt, S., Arnosti, A., Borges A.V.,
 2594 Dale, A.W., Gallego-Sala, A., Godd  ris, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina,
 2595 T., Joos, F., LaRowe, D.E., Leifeld, J., Meysman, F.J.R., Munhoven, G., Raymond, P.A.,
 2596 Spahni, R., Suntharalingam, P., and Thullner, M.: Anthropogenic perturbation of the carbon
 2597 fluxes from land to ocean, *Nature Geoscience* doi:10.1038/NGEO1830, 2013.

2598 Repo, M., Susiluoto, S., Lind, S., Jokinen, S., Elsakov, V., Biasi, C., Virtanen, T., and
 2599 Martikainen, P. J.: Large N₂O emissions from cryoturbated peat soil in tundra, *Nature*
 2600 *Geoscience*, 2, 189 -192, 2009.

2601 Ribeiro, M., Losenno, C., Dworak, T., Massey, E., Swart, R., Benzie, M., and Laaser, C.:
 2602 Design of guidelines for the elaboration of Regional Climate Change Adaptation Strategies,
 2603 Ecologic Institute, Vienna, 2009.

2604 Richter, A., Burrows, J.P., N  szlig, H., Granier, C., and Niemeier, U.: Increase in
 2605 tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129-132,
 2606 doi:10.1038/nature04092, 2005.

2607 Roiger, A., Thomas, J.L., Schlager, H., Law, K.S., Kim, J., Schafler, A., Weinzierl, B.,
 2608 Dahlkottter, F., Krisch, I., Marelle, L., Minikin, A., Raut, J.C., Reiter, A., Rose, M.,
 2609 Scheibe, M., Stock, P., Baumann, R., Bouarar, I., Clerbaux, C., George, M., Onishi, T., and
 2610 Flemming J.: Quantifying emerging local anthropogenic emissions in the Arctic region: The
 2611 ACCESS aircraft campaign experiment, *Bull. Amer. Meteor. Soc.*, 96, 441–460, 2015.

2612 Romanovsky, V., Burgess, M., Smith, S., Yoshikawa, K., and Brown, J. : Permafrost
 2613 temperature records: Indicators of climate change, *EOS*, 83, 589–594, 2002.

2614 Rosenfeld, D., Andreae, M.O., Asmi, A., Chin, M., de Leeuw, G., Donovan, D., Kahn, R.,
 2615 Kinne, S., Kivekäs, N., Kulmala, M., W. Lau, W., Schmidt, S., Suni, T., Wagner, T., Wild,
 2616 M., and Quaas, J.: Global observations of aerosol-cloud-precipitation-climate interactions,
 2617 Rev. Geophys., 52, 750–808, doi:[10.1002/2013RG000441](https://doi.org/10.1002/2013RG000441), 2014.

2618 Rosenfeld, D., Zheng, Y., Hashimshoni, E., Pöhlker, M. L., Jefferson, A., Pöhlker, C., Yu,
 2619 X., Zhu, Y., Liu, G., Yue, Z., Fischman, B., Li, Z., Giguzin, D., Goren, T., Artaxo, P.,
 2620 Barbosa, H. M. J., Pöschl, U., and Andreae, M. O.: Satellite retrieval of cloud condensation
 2621 nuclei concentrations by using clouds as CCN chambers, Proc. Natl. Acad. Sci., 113, 5828-
 2622 5834, doi:[10.1073/pnas.1514044113](https://doi.org/10.1073/pnas.1514044113), 2016.

2623 Ryshkevich, T. I., Mironov, G. N., Mironova, S. Y., Vlasenko, S. S., Chi, X., Andreae, M.
 2624 O., and Mikhailov, E. F., Comparative analysis of hygroscopic properties of atmospheric
 2625 aerosols at ZOTTO Siberian background station during summer and winter campaigns of
 2626 2011: Izvestiya, Atmospheric and Oceanic Physics, 51, 512-519, 2015.

2627 Saeki, T., Maksyutov, S., Sasakawa, M., Machida, T., Arshinov, M., Tans, P.J., Conway, T.,
 2628 Saito, M., Valsala, V., Oda, T., Andres, R. J., and Belikov D.: Carbon flux estimation for
 2629 Siberia by inverse modeling constrained by aircraft and tower CO₂ measurements, J.
 2630 Geophys. Res., 118, 1100-1122, doi:[10.1002/jgrd.50127](https://doi.org/10.1002/jgrd.50127), 2013.

2631 Sahoo, A. K., Pan, M., Troy, T. J., Vinukollu, R., Sheffield, J., and Wood, E. F.: Reconciling
 2632 the global terrestrial water budget using satellite remote sensing, Remote Sens. Environ.,
 2633 115, 1850-1865, doi:[10.1016/j.rse.2011.03.009](https://doi.org/10.1016/j.rse.2011.03.009), 2011.

2634 Sanchez-Rodriguez, R.: Learning to adapt to climate change in urban areas. A review of
 2635 recent contributions, Current Opinion in Environmental Sustainability, 1, 201–206, 2009.

2636 Sanderson, M.G., Collins, W.J., Johnson, C.E., and Derwent, R.G.: Present and future acid
 2637 deposition to ecosystems: the effect of climate change, *Atmos. Environ.*, 40, 1275-1283,
 2638 2006.

2639 Sasakawa, M., Shimoyama, K., Machida, T., Tsuda, N., Suto, H., Arshinov, M., Davydov,
 2640 D., Fofonov, A., Krasnov, O., Saeki, T., Koyama, Y., and Maksyutov, S.: Continuous
 2641 measurements of methane from a tower network over Siberia, *Tellus B*, 62, 403-416, doi:
 2642 10.1111/j.1600-0889.2010.00494.x, 2010.

2643 Savva, Y. and Berninger, F.: Sulphur deposition causes a large scale growth decline in boreal
 2644 forests in Eurasia, *Global Biogeochemical Cycles*, 24, GB3002,
 2645 doi:10.1029/2009GB003749, 2010.

2646 Sazhin, A.F., Romanova, N.D., and Mosharov, S.A.: Bacterial and primary production in the
 2647 pelagic zone of the Kara Sea, *Oceanology*, 50, 759-765, 2010.

2648 Schepaschenko, D., McCallum, I., Shvidenko, A., Fritz, S., Kraxner, F., and Obersteiner,
 2649 M.: A new hybrid land cover dataset for Russia: a methodology for integrating statistics,
 2650 remote sensing and in situ information, *J. Land Use Science*, 6, 245 -259, 2011.

2651 Schepaschenko, D.G., Shvidenko, A.Z., Lesiv, M.V., Ontikov, P.V., Schepaschenko, M.V.,
 2652 and Kraxner, F.: Area of Russia's forests and its dynamics based on synthesis of remote
 2653 sensing products, *Forest Science*, 3, 163-171, (in Russian), 2015.

2654 Schellnhuber, J., Cruzen, P., Clark, W., Claussen, M., and Held, H. (editors): *Earth System
 2655 Analysis for Sustainability*, Dahlem Workshop on Earth System Analysis for Sustainability
 2656 Berlin, May 25-30, 2003, MIT Press, 454 p, 2004.

2657 Schindler, D.W.: The cumulative effects of climate warming and other human stresses on
 2658 Canadian freshwaters in the new millennium, *Can. J. Fish. Aquat. Sci.*, 58, 18-29, 2001.

2659 Schlesinger, W. H.: Biogeochemistry: an analysis of global change, 2nd ed. Academic Press,
 2660 San Diego, California, 1997.

2661 Schuur, E.A.G., Vogel, J. G., Crummer, K.G., Lee H., Sickman, J.O., and Osterkamp, T.E.:
 2662 The effect of permafrost thaw on old carbon release and net carbon exchange from tundra,
 2663 Nature, 556 - 559, 2009.

2664 Scott, C. E., Rap, A., Spracklen, D.V., Forster, P. M., Carslaw, K.S., Mann, G.W., Pringle,
 2665 K.J., Kivekäs, N., Kulmala, M., Lihavainen, H., and Tunved, P.: The direct and indirect
 2666 radiative effects of biogenic secondary organic aerosol, Atmos. Chem. Phys., 14, 447–470.
 2667 2014.

2668 Screen, J. A. and Simmonds, I.: Declining summer snowfall in the Arctic: causes, impacts
 2669 and feedbacks, Clim. Dyn., 38, 2243-2256, doi:10.1007/s00382-011-1105-2, 2012.

2670 Screen, J. A. and Simmonds, I.: Increasing fall-winter energy loss from the Arctic Ocean and
 2671 its role in Arctic temperature amplification, Geophys. Res. Lett., 37, L16707,
 2672 doi:10.1029/2010GL044136, 2010.

2673 Sedlar, J., Tjernström, M., Mauritsen, T., Shupe, M., Brooks, I., Persson, P. O. G., Birch, C.
 2674 E., Leck, C., Sirevaag, A., and Nicolaus, M.: A transitioning Arctic surface energy budget:
 2675 the impacts of solar zenith angle, surface albedo and cloud radiative forcing, Clim. Dynam.,
 2676 37, 1643-1660, doi:10.1007/s00382-010-0937-5, 2011.

2677 Sellers, P.J., Dickinson, R. E., Randall, D.A., Betts, A. K., Hall, F. G., Berry, J. A., Collatz,
 2678 G. J., Denning, A. S., Mooney, H. A., Nobre, C. A., Sato, N., Field, C. B., and Henderson-
 2679 Sellers, A.: Modeling the exchanges of energy, water, and carbon between continents and the
 2680 atmosphere, Science, 24, 502-509, doi:10.1126/science.275.5299.502, 1997.

2681 Semiletov, I. P., Pipko, I. I., Shakhova, N. E., Dudarev, O. V., Pugach, S. P., Charkin, A. N.,
 2682 Kosmach, D., Gustafsson, O., and McRoy, C. P.: Carbon transport by the Lena River from
 2683 its headwaters to the Arctic Ocean, with emphasis on fluvial input of terrestrial particulate
 2684 organic carbon vs. carbon transport by coastal erosion, *Biogeosciences*, 8, 2407–2426, 2011.

2685 Sereda, J., Bogard, M., Hudson, J., Helps, D., and Dessouki, T.: Climate warming and the
 2686 onset of salinization: rapid changes in the limnology of two northern plains lakes,
 2687 *Limnologica*, 41, 1-9, 2011.

2688 Shakhova, N., Semiletov, I., Salyuk, A., Yusupov, V., Kosmach, D., and Gustafsson, O.:
 2689 Extensive methane venting to the atmosphere from sediments of the East Siberian Arctic
 2690 Shelf, *Science*, 327, 1246-1250, 2010.

2691 Shatalina, M. V., Mareev, E. A., Anisimov, S. V., and Shikhova, N.M.: Modeling of the
 2692 Electric-Field Dynamics in the Atmosphere Using the Test-Structure Method, *Radiophys.*
 2693 *Quantum El.*, 48, 575 - 586, doi:10.1007/s11141-005-0102-x, 2005.

2694 Shatalina, M. V., Mareev, E. A., Anisimov, S. V., and Shikhova, N. M.: Recovery of space
 2695 charge distribution by the method of test structures, in: *Proc. Int. Conf. Atm. Electr, ICAE*
 2696 *07, Beijing, China, 2007.*

2697 Sheffield, J., Wood, E. F., and Munoz-Arriola, F.: Long-term regional estimates of
 2698 evapotranspiration for Mexico based on downscaled ISCCP data, *Journal of*
 2699 *Hydrometeorology*, 11, 253-275, 2010.

2700 Shindell, D. and Faluvegi, G.: climate response to regional radiative forcing during the
 2701 twentieth century, *Nature Geosci.*, 2, 294-300, 2009.

2702 Shmakin, A. B. and Popova, V. V.: Dynamics of climate extremes in northern Eurasia in the
 2703 late 20th century. *Izvestia Akad. Nauk, Atmospheric and Oceanic Physics*, 42, 157-166,
 2704 2006.

2705 Shuntov, V. P., Dulepova, E. G., Temnih, O. C., Volkov, A. F., Naydenko, S. V.,
 2706 Chuchukalo, V. I., and Volkov, I. V.: Condition of biological resources in relation to
 2707 dynamics of macroecosystems in economic zone of Russian Far East Seas, *Dynamics of*
 2708 *the ecosystems and contemporary problems of conservation of potential bioresources of*
 2709 *Russian Seas*, Chapter 2, Vladivostok, Dalnauka, 75-176, 2007.

2710 Shvidenko, A., Schepaschenko, D., McCallum, I., and Nilsson, S.: Can the uncertainty of full
 2711 carbon accounting of forest ecosystems be made acceptable to policy makers?, *Climatic*
 2712 *Change*, 103, 137-157, 2010.

2713 Shvidenko, A., Schepaschenko, D., Vaganov, E. A., Sukhinin, A. I., Maksyutov, Sh. Sh.,
 2714 McCallum, I., and Lakyda, I. P.: Impacts of wildfire in Russia between 1998-2010 on
 2715 ecosystems and the global carbon budget, *Proc. Russian Academy of Sciences (Doklady*
 2716 *Earth Sciences)*, 441, 1678-1682, 2011.

2717 Shvidenko, A., Gustafson, E., McGuire, A.D., Kharuk, V.I., Svhepaschenko, D.G., Shugart,
 2718 H.H., Tchebakova, N.M., Vygodskaya, N.N., Onuchin, A.A., Hayes, D.J., McCallum, I.,
 2719 Maksyutov, S., Mukhortova, L.V., Soja, A.J., Beletti-Marchesini, L., Kurbatova, J.A.,
 2720 Oltchev, A.V., Parfenova, E.I., and Shuman, J.K.: Terrestrial ecosystems and their change,
 2721 in: *Regional Environmental Changes in Siberia and their Global Consequences*, Groisman, P.
 2722 Y., and Gutman, G. (Eds.), Springer, 171-249, 2013.

2723 Shvidenko, A.Z. and Schepaschenko, D.G.: Climate change and wildfires in Russia,
 2724 *Contemporary Problems of Ecology*, 6, 683-692, 2013.

2725 Shvidenko, A., Schepaschenko, D., Kraxner, F., and Obersteiner, M.: Terrestrial ecosystems
 2726 full carbon account as a fuzzy system: An attempt to understand uncertainties, in: 9th Int.
 2727 CO₂ Conf., Beijing, China, 3-7 June, 2013.

2728 Shvidenko, A. and Schepaschenko, D.: Carbon budget of Russian forests, *Siberian Journal of*
 2729 *Forest Science* (in Russian), 1, 69-92, 2014.

2730 Singh, D., Singh, R., and Kamra, A. The Electrical Environment of the Earth's Atmosphere:
 2731 A Review, *Space Science Reviews*, 113, 375–408, 2004.

2732 Singh, H. B., Brune, W. H., Crawford, J. H., Jacob, D. J., and Russell, P. B.: Overview of the
 2733 summer 2004 Intercontinental Chemical Transport Experiment-North America (INTEX-A),
 2734 *J. Geophys. Res.*, 111, D24S01, doi:10.1029/2006JD007905, 2006.

2735

2736 Skjelkvåle, B.L., Stoddard, J.L., and Andersen, T.: Trends in surface water acidification in
 2737 Europe and North America (1989-1998), *Water Air Soil Pollut.*, 130, 787-792, 2001.

2738 Skjelkvåle, B.L., and Wright, R.F.: Mountain lakes; sensitivity to acid deposition and global
 2739 climate change, *Ambio*, 27, 280-286, 1998.

2740 Skryzhevskaya, Y., Tynkkynen, V.-P., and Leppänen, S.: Russia's climate policies and local
 2741 reality, *Polar Geography*, 38, 146-170, 2015.

2742 Smedsrud, L., Esau, I., Ingvaldsen, R., Eldevik, T., Haugan, P., Li, C., Lien, V., Olsen,
 2743 A., Omar, A., Otterå, O., Risebrobakken, B., Sandø, A., Semenov, V., and Sorokina, S.: The
 2744 role of the Barents Sea in the Arctic climate system, *Reviews of Geophysics*, 51, 1-35, 2013.

2745 Smith, L. C.: *The World in 2050: Four forces shaping civilization's northern future*,
 2746 Brockman Inc., 2010.

2747 Smith, L.C., Sheng, Y., MacDonald, G.M., and Hinzman, L.D.: Disappearing arctic lakes,
2748 Science, 308, 1429, 2005.

2749 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.:
2750 Anthropogenic sulfur dioxide emissions: 1850-2005, Atmos. Chem. Phys., 11, 1101-1116,
2751 2011.

2752 Smith, S. J., Pitcher, H., and Wigley, T. M. L.: Global and Regional Anthropogenic Sulfur
2753 Dioxide Emissions, Global Planet Change, 29, 99-119, 2001.

2754 Sofiev, M., Vankevich, R., Ermakova, T., and Hakkarainen, J.: Global mapping of maximum
2755 emission heights and resulting vertical profiles of wildfire emissions, Atmos. Chem. Phys.,
2756 13, 7039-7052, doi:10.5194/acp-13-7039-2013, 2013.

2757 Sorooshian, S., Hsu, K. L., Gao, X., Gupta, H. V., Imam, B., and Braithwaite, D.: Evaluation
2758 of PERSIANN system satellite-based estimates of tropical rainfall, Bulletin of the American
2759 Meteorological Society, 81, 2035-2046, 2000.

2760 Spracklen, D. V., Bonn B., and Carslaw, K. S.: Boreas forests, aerosols and the impacts on
2761 clouds and climate, Philos. Trans. R. Soc., 266A, 1-11, doi:10.1098/rsta.2008.0201, 2008.

2762 Spracklen, D. V. and Heald, C. L.: The contribution of fungal spores and bacteria to regional
2763 and global aerosol number and ice nucleation immersion freezing rates, Atmos. Chem. Phys.,
2764 14, 9051-9059, doi:10.5194/acp-14-9051-2=014, 2014.

2765 State Report: State Report on State and Protection of Environment of the Russian Federation
2766 in 2011, Roshydromet, Moscow, 2011.

2767 Stavrou, T., Müller, J.-F., Peeters, J., Razavi, A., Clarisse, L., Clerbaux, C., Coheur, P.-F.,
2768 Hurtmans, D., De Mazière, M., Vigouroux, C., Deutsch, N. M., Griffith, D. W. T., Jones,

2769 N., and Paton-Walsh, C.: Satellite evidence for a large source of formic acid from boreal and
 2770 tropical forests, *Nature Geosci.*, 5, 26–30, doi:10.1038/ngeo1354, 2012.

2771 Stavrakou, T., Müller, J.-F., Bauwens, M., De Smedt, I., Van Roozendaal, M., Guenther, A.,
 2772 Wild, M., and Xia, X.: Isoprene emissions over Asia 1979–2012: impact of climate and land-
 2773 use changes, *Atmos. Chem. Phys.*, 14, 4587–4605, doi:10.5194/acp-14-4587-2014, 2014.

2774 Stohl, A.: Characteristics of atmospheric transport into the Arctic troposphere, *J. Geophys.*
 2775 *Res.*, 111, D11306, doi:10.1029/2005JD006888, 2006.

2776 Stroeve, J.C., Serreze, M.C., Holland, M.M., Kay, J.E., Maslanik, J., and Barrett, A.P.: The
 2777 Arctic's rapidly shrinking sea ice cover: a research synthesis, *Climatic Change*, 110, 1005-
 2778 1027, doi:10.1007/s10584-011-0101-1, 2012.

2779 Struthers, H., Ekman, A. M. L., Glantz, P., Iversen, T., Kirkevåg, A., Mårtensson, M.,
 2780 Seland, O., and Nilsson, E. D.: The effect of sea ice loss on sea salt aerosol concentrations
 2781 and the radiative balance in the Arctic, *Atmos. Chem. Phys.*, 11, 3459–3477, 2011.

2782 Su, H., Wood, E. F., McCabe, M. F., and Su, Z.: Evaluation of remotely
 2783 sensed evapotranspiration over the CEOP EOP-1 reference sites, *Journal of the*
 2784 *Meteorological Society of Japan*, 85A, 439–459, 2007.

2785 Su, H., Cheng, Y., Oswald, R., Behrendt, T., Trebs, I., Meixner, F.X., Andreae, M.O.,
 2786 Cheng, P., Zhang, Y., and Pöschl, U.: Soil nitrite as a source of atmospheric HONO and OH
 2787 radicals, *Science*, 333, 1616–1618, 2011.

2788 Sun, H., Santalahti, M., Pumpanen, J., Köster, K., Berninger, F., Raffaello, T., Jumpponen,
 2789 A., Asiegbu, F.O., and Heinonsalo, J.: Fungal community shifts in structure and function
 2790 across a boreal forest fire chronosequence, *Applied and Environmental Microbiology*, 81,
 2791 7869–80, 2015.

2792 Sun, Z., Niinemets, Ü., Hüve, K., Rasulov, B., and Noe, S.M.: Elevated atmospheric CO₂
 2793 concentration leads to increased whole-plant isoprene emission in hybrid aspen (*Populus*
 2794 *tremula x Populus tremuloides*), New Phytologist, 198, 788-800, doi: 10.1111/nph.12200,
 2795 2013.

2796 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.:
 2797 Soil organic carbon pools in the northern circumpolar permafrost region, Global
 2798 Biogeochem. Cycles, 23, GB2023, doi:10.1029/2008GB003327, 2009.

2799 Tchebakova, N.M., Parfenova, E.I., and Soja, A.J.: Effect of climate, permafrost and fire on
 2800 vegetation change in Siberia in a changing climate, Env. Res. Lett., 4, 045013, 2009.

2801 Tikkanen, E.: Conclusions. In: Tikkanen, E., Niemelä, I. (Eds.) Kola Peninsula pollutants
 2802 and forest ecosystems in Lapland. Final report of the Lapland Forest Damage Project.
 2803 Finland's Ministry of Agriculture and Forestry, The Finnish Forest Research Institute,
 2804 Gummerus Kirjapaino Oy, Jyväskylä, 71-81, 1995.

2805 Tishkov, A. A.: Biogeographical Consequences of Natural and Anthropogenic Climate
 2806 Changes, Biology Bull. Rev., 2, 132-140, 2012.

2807 Tjernström, M., Leck, C., Birch, C. E., Bottenheim, J. W., Brooks, B. J., Brooks, I. M.,
 2808 Bäcklin, L., Chang, R. Y.-W., de Leeuw, G., Di Liberto, L., de la Rosa, S., Granath, E.,
 2809 Graus, M., Hansel, A., Heintzenberg, J., Held, A., Hind, A., Johnston, P., Knulst, J., Martin,
 2810 M., Matrai, P. A., Mauritsen, T., Müller, M., Norris, S. J., Orellana, M. V., Orsini, D. A.,
 2811 Paatero, J., Persson, P. O. G., Gao, Q., Rauschenberg, C., Ristovski, Z., Sedlar, J., Shupe, M.
 2812 D., Sierau, B., Sirevaag, A., Sjogren, S., Stetzer, O., Swietlicki, E., Szczodrak, M.,
 2813 Vaattovaara, P., Wahlberg, N., Westberg, M., and Wheeler, C. R.: The Arctic Summer Cloud

2814 Ocean Study (ASCOS): overview and experimental design, *Atmos. Chem. Phys.*, 14, 2823-
2815 2869, doi:10.5194/acp-14-2823-2014, 2014.

2816 Troitskaya, Y., Rybushkina, G., Soustova, I., Balandina, G., Lebedev, S., and Kostianoy, A.:
2817 Adaptive retracking of Jason1 altimetry data for inland waters: the example of the Gorky
2818 Reservoir, *Int. J. Remote Sens.*, 33, 7559-7578, 2012.

2819 Troitskaya, Y., Troitskaya, Y., Ezhova, E. V., Sergeev, D. A., Kandaurov, A. A., Baidakov,
2820 G. A., Vdovin, M. I., and Zilitinkevich, S.: Momentum and buoyancy transfer in atmospheric
2821 turbulent boundary layer over wavy water surface. Part 2: Wind–wavespectra, *Nonlin,*
2822 *Processes Geophys.*, 20, 841–856, 2013.

2823 Tunved, P., Hansson, H.-C., Kerminen, V.-M., Ström, J., Dal Maso, M., Lihavainen, H.,
2824 Viisanen, Y., Aalto, P.P., Komppula, M., and Kulmala, M.: High natural aerosol loading
2825 over boreal forests, *Science*, 312, 261-263, 2006.

2826 Tynkkynen, V-P.: From mute to reflective: Changing governmentality in St Petersburg and
2827 the priorities of Russian environmental planning, *Journal of Environmental Planning and*
2828 *Management*, 53, 1-16, 2010.

2829 Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Da Costa Bechtold, V., Fiorino,
2830 M., Gibson, J.K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S.,
2831 Sokka, N., Allan, R.P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Van
2832 De Berg, L., Bidlot, L., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M.,
2833 Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B.J., Isaksen, L., Janssen, P. A.
2834 E. M., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette, J-J., Rayner, N. A., Saunders, R.
2835 W., Simon, P., Sterl, A., Trenberth, K.E., Untch, A., Vasiljevic, D., Viterbo, P., and

2836 Woollen, J.: The ERA-40 re-analysis, Q. J. R. Meteorol. Soc., 131, 2961-3012,
 2837 doi:10.1256/qj.04.176, 2005.

2838 Uttal, T., Starkweather, S., Drummond, J., Vihma, T., Makshtas, A., Darby, L., Burkhart, J.,
 2839 Cox, C., Schmeisser, L., Haiden, T., Maturilli, M., D. Shupe, M., de Boer, G., Saha, A.,
 2840 Grachev, A., Crepinsek, S., Bruhwiler, L., Goodison, B., McArthur, B., Walden, V.,
 2841 Dlugokencky, E., Persson, O., Lesins, G., Laurila, T., Ogren, J., Stone, R., Long, C., Sharma,
 2842 S., Massling, A., Turner, D., Stanitski, D., Asmi, E., Aurela, M., Skov, H., Eleftheriadis, K.,
 2843 Virkkula, A., Platt, A., Førland, E., Iijima, Y., Nielsen, I., Bergin, M., Candlish, L., Zimov,
 2844 N., Zimov, S., O'Neill, N., Fogal P., Kivi, R., Konopleva-Akish, E.A., Verlinde, J., Kustov,
 2845 V.Y., Vasel, B., Ivakhov, V.M., Viisanen, Y., and Intrieri, J.M.: International Arctic Systems
 2846 for Observing the Atmosphere (IASOA): An International Polar Year Legacy Consortium,
 2847 Bulletin of the American Meteorological Society, doi: [http://dx.doi.org/10.1175/BAMS-D-](http://dx.doi.org/10.1175/BAMS-D-14-00145.1)
 2848 [14-00145.1](http://dx.doi.org/10.1175/BAMS-D-14-00145.1), 2015.

2849 Vavrus, S., Waliser, D., Schweiger, A., and Francis, J.: Simulations of 20th and 21st century
 2850 Arctic cloud amount in the global climate models assessed in the IPCC AR4, Climate
 2851 Dynamics, 12, 1099-1115, doi: 10.1007/s00382-008-0475-6, 2009.

2852 Vaschuk, L. N. and Shvidenko, A. Z.: Dynamics of forests of Irkutsk region. Irkutsk, Russia
 2853 (in Russian), 2006.

2854 Velicogna, I., Tong, J., Zhang, T., and Kimball, J. S.: Increasing subsurface water storage in
 2855 discontinuous permafrost areas of the Lena River basin, Eurasia, detected from GRACE,
 2856 Geophys. Res. Lett., 39, L09403, doi:10.1029/2012GL051623, 2012.

2857 Verstraeten, W.W., Neu, J.L., Williams, J.E., Bowman, K.W., Worden, J.R., and Boersma,
 2858 K.F.: Rapid increases in tropospheric ozone production and export from China, *Nature*
 2859 *Geoscience* 8, 690–695, doi:10.1038/ngeo2493, 2015.

2860 Vesala, T., Launianen, S., Kolari, P., Pumpanen, J., Sevanto, S., Hari, P., Nikinmaa, E.,
 2861 Kaski, P., Mannila, H., Ukkonen, E., Piao, S.L., and Ciais, P.: Autumn temperature and
 2862 carbon balance of a boreal Scots pine forest in Southern Finland, *Biogeosciences*, 7, 163-
 2863 176, 2010.

2864 Vesala, T., Eugster, W., and Ojala, A.: Eddy Covariance Measurements over Lakes. In:
 2865 *Eddy Covariance - A practical guide to measurements and data analysis* (M. Aubinet, T.
 2866 Vesala and D. Papale). Springer, the Netherlands, 365-376, 2012.

2867 Vihma, T., Tisler, P., and Uotila, P.: Atmospheric forcing on the drift of Arctic sea ice in
 2868 1989-2009, *Geophys. Res. Lett.*, 39, L02501, doi:10.1029/2011GL050118, 2012.

2869 Vihma, T.: Effects of Arctic sea ice decline on weather and climate: a review, *Surv.*
 2870 *Geophys.*, 35, 1175–1214, DOI 10.1007/s10712-014-9284-0, 2014.

2871 Vihma, T., Pirazzini, R., Fer, I., Renfrew, I. A., Sedlar, J., Tjernström, M., Lüpkes, C.,
 2872 Nygård, T., Notz, D., Weiss, J., Marsan, D., Cheng, B., Birnbaum, G., Gerland, S., Chechin,
 2873 D., and Gascard, J. C.: Advances in understanding and parameterization of small-scale
 2874 physical processes in the marine Arctic climate system: a review, *Atmos. Chem. Phys.*, 14,
 2875 9403–9450, doi:10.5194/acp-14-9403-2014, 2014.

2876 Virtanen, T. H., Kolmonen, P., Rodríguez, E., Sogacheva, L., Sundström, A.-M., and
 2877 de Leeuw, G.: Ash plume top height estimation using AATSR, *Atmos. Meas. Tech.*, 7, 2437-
 2878 2456, doi:10.5194/amt-7-2437-2014, 2014.

2879 Vitousek, P.M., Mooney, H.A., Lubchenco, J., and Melillo, J.M.: Human domination of
 2880 Earth's ecosystems, *Science*, 277, 494-499, 1997a.

2881 Vitousek, P. M., Aber, J.D., Howarth, R.W., Likens, G.E., Matson, P.A., Schindler, D.W.,
 2882 Schlesinger, W.H., and Tilman, D.G.: Human alteration of the global nitrogen cycle: sources
 2883 and consequences, *Ecological Applications*, 7, 737-750, 1997b.

2884 Vitousek, P.M., Porder, S., Houlton, B.Z., and Chadwick, O.A.: Terrestrial phosphorus
 2885 limitation: mechanisms, implications, and nitrogen-phosphorus interactions, *Ecological*
 2886 *Applications*, 20, 5-15, 2010.

2887 Vivchar, A. V., Moiseenko, K. B., Shumskii, R. A., and Skorokhod, A. I.: Identifying
 2888 anthropogenic sources of nitrogen oxide emissions from calculations of Lagrangian
 2889 trajectories and the observational data from a tall tower in Siberia during the spring-summer
 2890 period of 2007: *Izvestiya Atmospheric and Oceanic Physics*, 45, 302-313,
 2891 doi:10.1134/s0001433809030049, 2009.

2892 Vonk, J.E., Sánchez-García, L., van Dongen, B.E., Alling, V., Kosmach, D., Charkin, A.,
 2893 Semiletov, I.P., Dudarev, O.V., Shakhova, N., Roos, P., Eglinton, T.I., Andersson, A., and
 2894 Gustafsson, Ö.: Activation of old carbon by erosion of coastal and subsea permafrost in
 2895 Arctic Siberia. *Nature* 489, 137-140, 2012.

2896 Walter, K. M., Zimov, S. A., Chanton, J. P., Verbyla, D., and Chapin III, F. S.: Methane
 2897 bubbling from Siberian thaw lakes as a positive feedback to climate warming, *Nature*, 443,
 2898 71-75. doi:10.1038/nature05040, 2006.

2899 Wang, T., Wei, X.L., Ding, A., Poon, C.N., Lam, K.S., Li, Y.S., Chan, L.Y., and Anson, M.:
 2900 Increasing surface ozone concentrations in the background atmosphere of Southern China,
 2901 1994-2009, *Atmos. Chem. Phys.*, 9, 6217-6227, 2009.

2902 Wang, R., Balkanski, Y., Boucher, O., Ciais, P., Peñuelas, J., and Tao, S.: Significant
 2903 contribution of combustion-related emissions to the atmospheric phosphorus budget, *Nature*,
 2904 8, 48–54, doi:10.1038/ngeo2324, 2014.

2905 Warneke, C., Bahreini, R., Brioude, J., Brock, C. A., de Gouw, J. A., Fahey, D. W., Froyd,
 2906 K. D., Holloway, J. S., Middlebrook, A., Miller, L., Montzka, S., Murphy, D. M., Peischl, J.,
 2907 Ryerson, T. B., Schwarz, J. P., Spackman, J. R., and Veres, P.: Biomass burning in Siberia
 2908 and Kazakhstan as an important source for haze over the Alaskan Arctic in April 2008,
 2909 *Geophys. Res. Lett.*, 36, L02813, doi:10.1029/2008GL036194, 2009.

2910 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and
 2911 Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to
 2912 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

2913 van Donkelaar, A., Martin, R. V., Brauer, M., and Boys, B. L.: Use of Satellite Observations
 2914 for Long-Term Exposure Assessment of Global Concentrations of Fine Particulate Matter,
 2915 *Environ. Health Perspect.*, 123, 135–143, doi:10.1289/ehp.1408646, 2015.

2916 Wespes, C., Emmons, L., Edwards, D. P., Hannigan, J., Hurtmans, D., Saunois, M., Coheur,
 2917 P.-F., Clerbaux, C., Coffey, M. T., Batchelor, R. L., Lindenmaier, R., Strong, K.,
 2918 Weinheimer, A. J., Nowak, J. B., Ryerson, T. B., Crounse, J. D., and Wennberg, P. O.:
 2919 Analysis of ozone and nitric acid in spring and summer Arctic pollution using aircraft,
 2920 ground-based, satellite observations and MOZART-4 model: source attribution and
 2921 partitioning, *Atmos. Chem. Phys.*, 12, 237–259, doi:10.5194/acp-12-237-2012, 2012.

2922 Whitehead, P.G. and Crossman, J.: Macronutrient cycles and climate change: key science
 2923 areas and an international perspective, *Sci Total Environ.*, 434, 13–17, 2012.

2924 Wolf, T. and Esau, I.: Air quality hazards under present and future climate conditions in
 2925 Bergen, Norway, *Urban climate*, 10, 801-814, doi: 10.1016/j.uclim.2014.10.006, 2014.

2926 Wolf, T., Esau, I., and Reuder, J.: Analysis of the vertical temperature structure in the
 2927 Bergen valley, Norway, and its connection to pollution episodes, *Journal of Geophysical*
 2928 *Research (Atmosphere)*, 119, doi:10.1002/2014JD022085, 2014.

2929 Xu, L., Guo, H., Boyd, C.M., Klein, M., Bougiatioti, A., Cerully, K.M., Hite, J.R.,
 2930 Isaacman-VanWertz, G., Kreisberg, N.M., Knote, C., Olson, K., Koss, A., Goldstein, A.H.,
 2931 Hering, S.V., de Gouw, J., Baumann, K., Lee, S.H., Rodney A.N., Weber, R.J., and Ng
 2932 N.L.: Effects of anthropogenic emissions on aerosol formation from isoprene and
 2933 monoterpenes in the southeastern United States, *Proc Natl Acad Sci USA* 112:37–42, 2015.

2934 Yvon-Durocher, G., Caffrey, J.M., Cescatti, A., Dossena, M., del Giorgio, P., Gasol, J.M.,
 2935 Montoya, J. M., Pumpanen, J., Staehr, P.A., Trimmer, M., Woodward, G., and Allen, A.P.:
 2936 Reconciling the temperature dependence of respiration across timescales and ecosystem
 2937 types, *Nature*, 487, 472-476, doi:10.1038/nature11205, 2012.

2938 Zábori, J., Krejci, R., Ekman, A. M. L., Mårtensson, E. M., Ström, J., de Leeuw, G., and
 2939 Nilsson, E. D.: Wintertime Arctic Ocean sea water properties and primary marine aerosol
 2940 concentrations, *Atmos. Chem. Phys.*, 12, 10405-10421, doi:10.5194/acp-12-10405-2012,
 2941 2012.

2942 Zábori, J., Krejci, R., Ström, J., Vaattovaara, P., Ekman, A. M. L., Salter, M. E., Mårtensson,
 2943 E. M., and Nilsson, E. D.: Comparison between summertime and wintertime Arctic Ocean
 2944 primary marine aerosol properties, *Atmos. Chem. Phys.*, 13, 4783–4799, doi:10.5194/acp-
 2945 13-4783-2013, 2013.

2946 Zaehle, S., Ciais, P., Friend, A. D., and Prieur, V.: Carbon benefits of anthropogenic reactive
 2947 nitrogen offset by nitrous oxide emissions, *Nature Geoscience*, 4, 601-605, 2011.

2948 Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T.,
 2949 Kimoto, T., Chang, D., Pöschl, U., Cheng, Y. F., and He, K. B.: Exploring the severe winter
 2950 haze in Beijing: the impact of synoptic weather, regional transport and heterogeneous
 2951 reactions, *Atmos. Chem. Phys.*, 15, 2969-2983, 10.5194/acp-15-2969-2015, 2015.

2952 Zhou, L., Tucker, C.J., Kaufmann, R.K., Slayback, D., Shabanov, N.V., Fung, I., and
 2953 Myneni, R.B.: Variations in northern vegetation activity inferred from satellite data of
 2954 vegetation index during 1981 to 1999, *J. Geophys. Res.*, 106, D17, 20,069-20,083, 2001.

2955 Zieger, P., Aalto, P. P., Aaltonen, V., Äijälä, M., Backman, J., Hong, J., Komppula, M.,
 2956 Krejci, R., Laborde, M., Lampilahti, J., de Leeuw, G., Pfüller, A., Rosati, B., Tesche, M.,
 2957 Tunved, P., Väänänen, R., and Petäjä, T.: Low hygroscopic scattering enhancement of boreal
 2958 aerosol and the implications for a columnar optical closure study, *Atmos. Chem. Phys.*, 15,
 2959 7247-7267, doi:10.5194/acp-15-7247-2015, 2015.

2960 Zilitinkevich, S.S.: *Turbulent Penetrative Convection*, Avebury Technical, Aldershot, 1991.

2961 Zilitinkevich, S., Esau, I., and Baklanov, A.: Further comments on the equilibrium height of
 2962 neutral and stable planetary boundary layers, *Quart. J. Roy. Met. Soc.*, 133, 265-271, 2007.

2963 Zilitinkevich, S.S. and Esau, I.N.: Planetary boundary layer feedbacks in climate system and
 2964 triggering global warming in the night, in winter and at high latitudes, *Geography*,
 2965 *Environment and Sustainability*, 1, 2, 20-34, 2009, doi: 10.1002/qj.27

2966 Zilitinkevich, S., Kulmala, M., Esau, I., and Baklanov, A.: Megacities – refining models to
 2967 personal environment, *WMO Bulletin*, 64, 20-22, 2015.

Zilitinkevich S.S.: The Height of the Atmospheric Planetary Boundary layer: State of the Art and New Development – Chapter 13 in “National Security and Human Health Implications of Climate Change”, edited by H.J.S. Fernando, Z. Klaić, J.L. McKulley, NATO Science for Peace and Security Series – C: Environmental Security (ISBN 978-94-007-2429-7), Springer, 147-161, 2012.

Zolotokrylin, R., Titkova, T. B., Cherenkova, E. A., and Vinogradova, V. V.: Satellite index for evaluation of climatic extremes in dru areas, Mod Stud Earth Remote Sens Sp., 9, 114-121 (in Russian), 2012.




Figure 1: The thematic research areas relevant to the Northern Eurasian land system include Land-Topic-1 “changing ecosystem processes”, Land-Topic-2 “ecosystem structural changes and resilience” and Land-Topic--3 “risk areas of permafrost thawing”. For the atmospheric

system they are Atmosphere-topic-1 “atmospheric composition and chemistry”, Atmosphere-
 topic -2 “Urban air quality”, are Atmosphere-topic-3, “atmospheric circulation and weather”,
 for the aquatic system they are Aquatic-topic-1 “Arctic Ocean in the climate system” ,
 Aquatic-topic-2 “maritime ecosystems”, Aquatic-topic-3 “Lakes and large river systems” and
 for the social system they are Society-topic-1 “natural resources and anthropogenic activities”,
 Society-topic-2 “natural hazards” and Society-Topic-3“social transformations”.





Figure 2: Linear trends in the annual maximum Normalized Difference Vegetation Index (NDVI) obtained from analysis of the MODIS 0.25 km data product for 2000-2014 over the North-Western Siberia region in Russia. The trends are given in the NDVI changes per 15 years. The yellow colors show the decreasing NDVI, which corresponds to decreasing biological production; the blue colors show the increasing NDVI. More detailed analysis of the trends is given in Esau et al. (2016).

2995

2996

2997



2998 Figure 3: Map showing the existing ACTRIS (Aerosols, clouds and trace gases Research
2999 Infrastructure Network) and ICOS (Integrated Carbon Observations System) stations in
3000 Europe (blue), stations making atmospheric and/or ecosystem measurements in Russia (red),
3001 INTERACT (International Network for Terrestrial Research and Monitoring in the Arctic)
3002 stations in Russia (light blue) and China Flux stations in China (yellow). However, all of these
3003 stations need certain upgrades.

3004




Figure 4: In urban and industrialized regions, the process understanding of biogeochemical cycles includes anthropogenic sources, such as industry and fertilizers, as essential parts of the biogeochemical cycles.




Figure 5: Hydrological cycle schematics.





Figure 6: Carbon cycling in the Arctic will change as the climate warms. Figure after ACIA, 2004. (Impacts of a Warming Arctic: Arctic Climate Impact Assessment (ACIA) Overview Report).

3019

3020


3021



3022

3023 Figure 7: Schematic figure for terrestrial nitrogen cycle.

3024



3025

3026 Figure 8: Schematic figure of the phosphorus cycle.





Figure 9: Schematic figure of the sulfur cycle.



3034 Figure 10. An example of the study approach to be implemented by PEEX for integrating
3035 natural science and social science knowledge and generating climate predictions and narratives
3036 of the Northern regions.