

Reviewer comments in black, our responses in red. Revised paper with tracked changes is appended after our responses.

Response to RC1 from Anonymous Referee #2, 31 March 2016

Specific comments:

1. Page 2 Lines 29-31 (P2L29-31 hereafter): I would have been more comfortable if two different models were used for the TR and DA. As it has been done, the DA experiments are constrained “twice” with the same information (the first time through the background that comes from using in the DA runs the TR model, and the second time by assimilating pseudo-observations derived from the TR itself).

We appreciate the reviewer’s comments regarding the “identical twin” approach used in this study. This study, along with the previous two papers, was designed to lay the groundwork for understanding the interactions of tracers and winds in the data assimilation framework. Given the limitations of the shallow water model as an approximation to the full 3-D atmosphere, we do not expect quantitative results to be applicable to the real world. However, our hope is that these studies will benefit future, more realistic, studies.

I appreciate the statement “results are therefore likely to be overly optimistic”, and the care the authors put in designing the experiments, for instance by providing an initial error in the background resulting from imposing the Day=20 T=6h TR fields to be the initial condition for the DA, but I do wonder how significant the differences between the Day=20 T=0 and Day=20 T=6h TR fields really are. This could perhaps be commented in the paper.

The initial zonal and meridional wind errors as a function of latitude are shown by the black lines on newly numbered Fig. 9 and Fig. 10, respectively. Since these aren’t referenced until much later in the paper, we decided to add a brief discussion of the initial error in Sec 2.3. We also added a time series plot of the global RMSE of the zonal wind and WEP (new Fig. 2) as described further below, which addresses the issue of spin-up.

2. Related to point 1 above: How responsive is the system? This aspect can help the interpretation of the results on two fronts: *a*) to disentangle the impact of essentially using the same constraint twice (in the sense mentioned above) and possibly quantify the actual wind extraction potential of ozone data assimilation (i.e. understand how overly optimistic the results are); and *b*) to know if at day 10 the system is completely spun up. A comment on this point should perhaps be included as well.

We included in the revision a time series plots of the global RMSE of the zonal wind and WEP (new Fig. 2) to show how the system evolves. This figure is for the optimal results with hybrid 4DVar assimilation of height and ozone using ensemble size of 100 members. This shows that the system is fairly well spun up after about 6 days. The last two figures (newly numbered as Fig. 9 and 10) also

provide an indication of the overall responsiveness of the system, since they show the initial and final errors as a function of latitude.

3. Section 2.2 and P5L20: In my understanding, inflation and localization are used in EnKF to address issues (such as filter divergence or long range spurious correlations) that result from under-sampling, i.e. having an ensemble so small in size that is not statistically representative of the state of the system.

Yes, this is our understanding as well.

I'd argue that this does not apply to the case of the "large" ensemble with 1518 members and it would also explain why for that ensemble size the WEP value does not significantly change as function of the localization length (figure 2).

Yes, this is likely the case.

The authors may want to comment on this and explain why and in what sense the large ensemble results that used no localization were worse than those with localization (P5L20).

The ensemble initialization technique we used involved sampling the truth run at regular time intervals. For the large ensembles, this required using a small time interval (36 minutes) and sampling late into the truth run when the dynamical variability of the system was settling down. Therefore, the ensemble members may not have been completely independent. Since the ensemble wasn't "perfect", applying some localization actually improved the system. We added a comment on this.

4. P5L4-6: I appreciate the reference to A15, but perhaps the authors can say something more here on the NNMI and make this manuscript more self-contained. Also in A15, instead of NNMI the authors used the acronym NMI. It would be good perhaps to either use a consistent acronym or acknowledge the use of a different name.

We will include some additional discussion of the normal mode initialization to make the paper more self-contained, and will also change to the acronym "NMI" rather than NNMI.

5. P7L11-12: I guess the wind vector RMSE is computed as in equation 6 of A15, perhaps a reference can be added.

This is nearly the same calculation. The only difference is that in the present paper we applied an area-weighting factor that we didn't apply in A15. We have included a comment to this effect.

6. Fig 7: You may want to specify in the right panel caption how the relative difference is computed.

We added a description to explain how the relative difference is computed.

I find interesting the reduced efficiency of the 50-member ensemble compared with those with 25 and 100 members. Is the reason for such a result known?

The reason for obtaining somewhat reduced efficiency with 50 members is unknown. This may be related to the fact that we tuned the system with one length scale for the assimilation of height and ozone (note that this feature doesn't occur for the height only assimilation). For unknown reasons, using a single localization length may have adversely affected the 50 member experiments more than the other experiments. In principle, we could simultaneously tune separate lengths for the two variables, since it is clear from newly numbered Fig. 3, and also discussion in A15, that the optimal length scale for ozone is smaller than for height. Why this would adversely affect the 50 member case is unclear, but it is a factor that is worth considering for future work.

Response to RC2 from Thomas Milewski, 22 May 2016

Specific notes

1) The choice of using streamfunction and velocity potential as control variables seems to be based on the experiments ran in A15 with the EnKF. What about the impact of choosing this set of control variables versus zonal and meridional winds in the 4Dvar system?

We haven't examined the impact of variable choice in the 4DVar system, but this would be an interesting follow up study. In the analytic algorithm that we used, based on Daley's formulation, the wind/wind and wind/geopotential height correlations are actually derived from streamfunction and velocity potential correlations using the f -plane approximation and isotropic assumption. So while the correlations are currently available for either set of variables, specification of the error standard deviations seems to be more straight-forward when using zonal and meridional wind. For example, we assume globally constant values for these quantities, which should be a reasonable assumption. It is less intuitive how to specify error standard deviations for streamfunction and velocity potential.

2) How did the authors come up with the value of 1518 as the number of dynamical state variables in the T21 system?

There are 253 modes for each variable used in spectral T21 system. This can be calculated by the following algorithm (in IDL notation). For given truncation NN, the total number of modes for a single global field is:

```
NN=21
totalmodes=0
for m=0,NN do begin
  nummodes=NN-abs(m)+1
  totalmodes=totalmodes+nummodes
endfor
```

Since there are 3 variables in the dynamical system, this results in $253 \times 3 = 759$ modes. And, since each mode is characterized by a complex amplitude, we multiply this by 2 to get the degrees of freedom for the dynamical system of 1518.

If I understand correctly, this is roughly the number of degrees of freedom in the dynamical system, but does it include the influence of O3?

Yes, the full system with O3 would have $253 \times 4 \times 2 = 2024$ degrees of freedom.

If not, this might explain why the optimal blending factor was not 1.0 for the large ensemble Z/O3 assimilation (before NNMI).

Yes, this may be a partial explanation of why the optimal blending is not 1.0, since our ensemble size is less than that of the entire system. We actually tried running the system with the size 2024 members. However, results were slightly worse than running with 1518 members. We think this may be due to inadequacies in the ensemble initialization (see also response to RC1, question 3).

3) Optimal localization lengths for Z-only and Z/O3 assimilation are very different, suggesting that O3 error covariance structures and Z error covariance structures are probably different. Ideally, this should require separate localization lengths. Did you try tuning Z and O3 localization lengths separately?

We didn't try tuning the Z and O3 localization lengths separately, but that would be a logical follow-on to this project. Having an additional tuning parameter (2 length scales rather than 1) would undoubtedly improve the results, but would likely not change the overall conclusions. In addition, it would be interesting to explore tuning the system using correlation lengths that vary with latitude.

4) In figures 7, 8 and 9, the hybrid system is only compared to 4DVar, not EnKF. Since EnKF seems to outperform 4DVar for moderate to large ensemble sizes (at least in terms of WEP), it would be instructive to see the improvement that the hybrid system brings with respect to second-best performing system.

In the revision we added the EnKF results to the original Figs. 7, 8, and 9 (newly renumbered as Figs. 8, 9, and 10) along with some discussion.

5) Considering that this paper seems to complete a trilogy on the topic of wind extraction in a hierarchy of data assimilation systems, it would be interesting to have a final paragraph in the "Conclusions" section that is a more extensive review of the behavior of the different DA systems, possibly including the pros and cons of each.

This is an excellent idea. We decided to move much of the current conclusions in to a summary section and then used the conclusions section for a more extensive review.

Technical notes

1) P.6 L.16: Please correct "The WEP value for of Z/O3"

Done

Hybrid ensemble 4DVar assimilation of stratospheric ozone using a global shallow water model

Douglas R. Allen, Karl W. Hoppel, David D. Kuhl

Remote Sensing Division, Naval Research Laboratory, Washington, DC, USA

5 *Correspondence to:* D. R. Allen (douglas.allen@nrl.navy.mil)

Abstract. Wind extraction from stratospheric ozone (O_3) assimilation is examined using a hybrid ensemble 4DVar shallow water model (SWM) system coupled to the tracer advection equation. Stratospheric radiance observations are simulated using global observations of the SWM fluid height (Z), while O_3 observations represent sampling by a typical polar-orbiting satellite. Four ensemble sizes were examined (25, 50, 100, and 1518 members), with the largest ensemble equal to the number of dynamical state variables. The optimal length scale for ensemble localization was found by tuning an ensemble Kalman filter (EnKF). This scale was then used for localizing the ensemble covariances that were blended with conventional covariances in the hybrid 4DVar experiments. Both optimal length scale and optimal blending coefficient increase with ensemble size, with optimal blending coefficients varying from 0.2 to 0.5 for small ensembles to 0.5 to 1.0 for large ensembles. The hybrid system outperforms conventional 4DVar for all ensemble sizes, while for large ensembles the hybrid produces similar results to the offline EnKF. Assimilating O_3 in addition to Z benefits the winds in the hybrid system, with the fractional improvement in global vector wind increasing from ~35% with 25 and 50 members to ~50% with 1518 members. For the smallest ensembles (25 and 50 members), the hybrid 4DVar assimilation improves the zonal wind analysis over conventional 4DVar in the Northern Hemisphere (winter-like) region and also at the equator, where Z observations alone have difficulty constraining winds due to lack of geostrophy. For larger ensembles (100 and 1518 members), the hybrid system results in both zonal and meridional wind error reductions, relative to 4DVar, across the globe.

1 Introduction

The extraction of wind information from stratospheric ozone (O_3) assimilation using a 4D data assimilation (DA) system is an attractive prospect, given the paucity of direct wind observations in the stratosphere. The tracer-wind relationship has been examined with a variety of DA systems including the extended Kalman Filter (EKF, Daley, 1995, 1996), 4D Variational assimilation (4DVar, Riishøjgaard, 1996; Peuch et al., 2000; Andersson et al., 2007; Peubey and McNally, 2009; Semane et al., 2009; Han and McNally, 2010; Dragani and McNally, 2013; Allen et al., 2013, 2014), and Ensemble Kalman Filter (EnKF, Milewski and Bourqui, 2011, Allen et al., 2015). While idealized studies have shown strong potential for wind extraction from tracer assimilation, attempts to assimilate O_3 using realistic numerical weather prediction (NWP) systems have produced mixed results (see Allen et al., 2015 for a discussion). In an effort to understand the problem in more detail,

we previously developed a shallow water model (SWM) test case representing Northern Hemisphere (NH) winter stratosphere conditions. Assimilation experiments using both 4DVar (Allen et al., 2014; hereinafter A14) and EnKF (Allen et al., 2015; hereinafter, A15) showed that tracer assimilation is useful for wind extraction, but also raised issues such as sensitivity to measurement errors, localization, and choice of DA state variables, and the problem of imbalance.

5

Another approach to evaluating O₃-wind interaction in DA is to blend the 4DVar static covariance with flow-dependent ensemble covariance within the 4DVar. This hybrid 4DVar method is becoming increasingly popular at operational NWP centers (Buehner et al., 2010; Bonavita et al., 2012; Clayton et al., 2013; Kuhl et al., 2013; Kleist and Ide, 2015). In this paper, we extend our previous work by examining O₃-wind interactions using a hybrid 4DVar system within the SWM framework. Tuning of the length scale for the ensemble covariance localization as well as the covariance blending parameter are examined, in addition to probing the limits of wind extraction with a large ensemble experiment.

10

The layout of the paper is as follows. Section 2 describes the SWM hybrid 4DVar system and the experimental design. Section 3 describes hybrid results using both small and large ensembles, relative to the size of the state vector. Section 4 presents a discussion of the optimal assimilation experiments. ~~Sections 5 and 6 provide a summary and conclusions, respectively are provided in Section 5.~~

15

2. Model description

2.1 Forecast model, truth run, and observations

The forecast model is the spectral SWM described in A14 and A15. For this paper, the model was run at a lower resolution of triangular truncation T21 (64 longitudes \times 32 latitudes, for a Gaussian grid spacing of $\sim 5.6^\circ$ at the equator) rather than T42 in order to facilitate a large number of tuning experiments and to allow the full background error covariance to be stored in active memory. To accommodate the lower resolution, the horizontal fourth order diffusion coefficient was increased from $5.0 \times 10^{15} \text{ m}^4 \text{ s}^{-1}$ to $8.9 \times 10^{16} \text{ m}^4 \text{ s}^{-1}$, which maintains an e-folding damping for the highest wavenumber of approximately one day. Other settings are the same, including a 10 km global mean height and time step of 120 s.

20

25

The truth run (TR) is similar to that used in A14 and A15. The system is initialized with a zonal jet with maximum wind of 60 ms^{-1} in the NH, which is in geostrophic balance with the fluid height. A time-dependent topographic forcing is applied over the first 20 days. The shape of the forcing is the same as in A14 and A15 (i.e., a zonal wave 1 mountain centered at 45°N), but the mountain height is increased from 1250 m to 1750 m to allow greater dynamical variability in the T21 system. After day 20, the topography is flat for the rest of the TR; since the assimilation begins on day 20, there is no topography

30

during the DA experiments. We use the same forecast model for the TR and the DA (i.e., “identical twin” experiments), making our results a best case scenario. The results are therefore likely to be overly optimistic.

5 Observations of O_3 and fluid height (Z) were generated by sampling the TR with the same frequency as in A15. The O_3 observations mimic Aura Microwave Limb Sounder sampling (one observation every 24.5 s), while Z observations are pseudo-random in space and time, with the same sampling frequency. We increased the error standard deviations to more realistic values of 0.3 parts per million volume (ppmv) instead of 0.408 ppmv for O_3 , and 200 m instead of 50 m for height. The 200 m error for Z corresponds to approximately 1°K, using the scaling explanation in A15. Experiments assimilating either Z only (referred to as “ Z assimilation”) or Z and ozone (referred to as “ Z/O_3 assimilation”) are performed.

10 2.2 Ensemble Kalman filter

The EnKF is described in detail in A15. Briefly, it is a “perturbed observations” EnKF (Houtekamer and Mitchell, 1998; Evensen, 2003), with data assimilated in 20 minute batches. The EnKF analysis equation can be solved using different combinations of control variables. In this study, we use streamfunction, velocity potential, Z , and O_3 (the EnKF- $\psi\chi$ system), which was shown in A15 to have less imbalance than when zonal and meridional wind are used as the horizontal flow variables (also discussed in Kepert, 2009). To avoid filter divergence, we apply a state space covariance inflation factor (Anderson, 2007) to the background ensemble before assimilating observations. The inflation factor is designed to alter the global average ensemble spread in the streamfunction to match the global Root Mean Square Error (RMSE) in the streamfunction. We also apply the elementwise (Schur product) localization (e.g., Houtekamer and Mitchell, 2001) using Eq. (4.10) of Gaspari and Cohn (1999).

20 2.3 Hybrid 4DVar

The SWM 4DVar DA system is described in A14. The 4DVar minimizes a standard cost function using the accelerated representer approach (Xu et al., 2005; Rosmond and Xu, 2006) with a perfect model assumption. The conventional initial background error covariance $\mathbf{B}_0^{\text{con}}$ is calculated using an analytic formulation that employs wind-geopotential correlations based on approximate geostrophic balance on an f -plane, i.e., constant Coriolis parameter with latitude (Daley, 1991; Daley and Barker, 2001). There is no coupling between O_3 and dynamical variables in $\mathbf{B}_0^{\text{con}}$, but coupling [between these variables](#) does develop implicitly over the 4DVar time window. The background error standard deviations are adaptively tuned to match the globally averaged error standard deviations (with respect to the TR), as discussed in A14. The tangent linear model is also run at T21 resolution with the same diffusion coefficient and time step as in the nonlinear model. The 4DVar system runs with a 6-hour update cycle, and the 6-hour analysis at the end of one window is used to initialize the analysis at the start of the subsequent window.

For the horizontal correlation used in $\mathbf{B}_0^{\text{con}}$, we found that the function used in A14 (single order auto regressive (SOAR) function with 1000 km length, sloping up to ~~150250~~ 1500 km in the tropics; see A14, Fig. 1a) was near optimal for the current experiments, even though the observations used in this study are much more sparse than in A14. Single observation experiments revealed that increasing the length scale in the $\mathbf{B}_0^{\text{con}}$ introduces more gravity waves into the SWM system. The imbalance is minimized with smaller length scales, since the f -plane assumption is more accurate. Reformulating the analytic balance for larger correlation lengths, or applying either a digital filter or nonlinear normal mode initialization within the variational solver, may further optimize the system. However, this is beyond the scope of the current paper.

To run the hybrid system (see Fig. 1 for schematic diagram), we first perform a 10-day EnKF simulation. We then run the hybrid 4DVar over the same 10-day period in which the ensemble covariance, $\mathbf{B}_0^{\text{ens}} = \mathbf{X}'\mathbf{X}'^T / (N_{\text{ens}} - 1)$, is calculated at the start of each 6-hour window using the ensemble states \mathbf{X}' . The prime indicates perturbation from the ensemble mean, and N_{ens} is the ensemble size. The ensemble covariance is then blended together with $\mathbf{B}_0^{\text{con}}$ using $\mathbf{B}_0^{\text{hybrid}} = (1 - \alpha)\mathbf{B}_0^{\text{con}} + \alpha\mathbf{S} \circ \mathbf{B}_0^{\text{ens}}$. Here α is a blending coefficient between 0 and 1, \mathbf{S} is the localization function, and the open circle indicates the Schur product. Using the offline EnKF facilitates running the hybrid system with a range of parameters without having to compute the ensemble each time. Tests in which the EnKF is re-centered about the 4D-Var analysis at the beginning of each cycle, as it would be done in an operational setting, produce similar results.

The experimental design is similar to A14 and A15. The DA experiments begin 20 days into the TR (day 20, 0 h), with the initial state defined as the TR state that is offset 6 h from the initial time (i.e., day 20, 6 h). This initial 6 h offset, or mismatch, between the TR and the initial background fields is the source of the initial background error. The initial wind errors range from $\sim 2\text{-}3 \text{ ms}^{-1}$ in the extratropics to $\sim 3\text{-}6 \text{ ms}^{-1}$ in the tropics (see thin black lines on Figs. 9 and 10). -We then perform 10-day assimilation ~~runs~~ experiments and compare the final wind errors with the initial wind errors. To illustrate how the wind errors evolve with time, Fig. 2a shows the global RMSE of the zonal wind over a 10 day DA period (the meridional wind shows a similar trend) for one of the hybrid 4DVar experiments. The wind error starts at $\sim 3.3 \text{ ms}^{-1}$, but drops rapidly over the first several days before leveling out near day 6, suggesting that the system is well spun-up after ~ 6 days of assimilation. Fig. 2b shows tThe Wind Extraction Potential (WEP), which is a normalized diagnostic relating the analyzed RMSE of the vector wind to the initial RMSE of the vector wind (a WEP value of 100% indicates perfect winds, while 0% indicates no improvement). Details of the WEP calculation are provided in A15. One slight difference is that in this study the RMSE of the vector wind included an area-weighting factor that wasn't applied in Eq. (6) of Allen et al. (2015). Fig. 2b shows the WEP starts at zero, but increases rapidly over the first several days, before leveling out at $\sim 83\%$ by day 10 of the assimilation experiment.

There are two main “tuning” parameters that we are considering for this study: the ensemble covariance localization length scale (L) and the hybrid blending coefficient (α). While the localization length used in the hybrid blending does not have to ~~be the same as~~ match what is that used in the EnKF, sensitivity tests showed that using the same length for both provides optimal or near-optimal results. Therefore in the ~~tests-experiments~~ for this paper the same length is always used for both. Note that since inflation is automatically adjusted in a self-consistent manner with the TR, it does not require tuning.

A15 quantified imbalance due to erroneous gravity wave modes that enter the EnKF system via imbalance in the analysis increments. Since the TR is virtually free of gravity waves due to the nature of the topographic forcing, any imbalance is considered to be unwanted noise. The imbalance can be reduced by judicious choice of flow variables and by tuning the localization length. In addition, A15 showed that application of nonlinear normal mode initialization (NMI) as a post-processing diagnostic ~~has been shown to benefit~~ improved the analysis in the EnKF system. In the NMI approach, the SWM state is first decomposed into three different mode types: eastward gravity waves, westward gravity waves, and rotational waves. The state is then adjusted using the Machenhauer (1977) condition, which reduces the time tendencies of the complex amplitudes of the gravity wave modes. We apply five iterations to solve the nonlinear balance equation using a cutoff frequency of 1.0 day⁻¹. This removes much of the imbalance and results in better agreement with the TR. For each of the ~~tests-experiments~~ in this study (4DVar, EnKF, and hybrid), we therefore also compare results with and without NMI post-processing.

3. Results

3.1 Tuning the localization length

In order to examine the sensitivity of the 4DVar to the quality of the ensemble covariance, the offline EnKF is run at different ensemble sizes. Three “small” ensemble experiments are performed with 25, 50, and 100 members, and one “large” ensemble experiment is performed with 1518 members, which equals the number of ~~dynamical state variables~~ degrees of freedom in the T21 SWM system. The large ensemble experiment is used to explore the maximum benefit of ensemble covariance blending in the 4DVar system, while the small ensemble experiments test the performance of limited, or more practical, ensemble sizes. To initialize the small (large) ensembles, we sampled the TR at 6-h (36-min) intervals, starting at day 20. Experiments were also performed with 2024 members, which equals the total number of degrees of freedom in the system, including ozone. However, results were slightly worse than with 1518 members. This was likely caused by inadequate initialization of the large ensemble through sampling of the TR. Since the TR does not have topographic forcing, the system becomes less dynamically active as time progresses, and therefore the initial ensembles for the large runs may not be completely independent.

For the small ensembles, the tuning of the localization length was performed using 10-day EnKF experiments with a range of localization lengths, starting at 500 km and increasing in 500 km increments until the 10-day WEP values showed an obvious maximum. Due to intensive computation time, the large ensemble experiments were not finely tuned, rather localization lengths of 10,000, 15,000 and 20,000 km were tested. Tests with the large ensemble were also performed with no localization, but results were slightly worse. This may be caused by inadequacies of the initialization procedure, as explained above. Figure 2-3 shows the WEP as a function of length for eight combinations of observations (Z or Z/O_3) and ensemble size (25, 50, 100, and 1518). Most experiments show smoothly-varying WEP as a function of length, with a well-defined peak. At 25 members the peak is narrow for Z/O_3 , while for 100 members and Z only, the peak is quite broad. The optimal lengths (i.e., producing maximum WEP) are indicated by vertical red lines; see Table 1 for numerical values. For the large ensembles, the WEP is not very sensitive to localization length for the large ensembles, since the ensemble is sampling nearly all of the background error state.

One main conclusion from these tests is that both optimal length and optimal WEP increase with ensemble size. For small ensemble experiments, optimal lengths are also larger for the Z only assimilation than for Z/O_3 assimilation. At 100 members, the optimal length of 14,000 km for Z assimilation is quite long; this is likely due to the large-scale structure of the Z fields in this experiment, combined with the relatively low resolution T21 system. Application of NMI increases the WEP for all ensemble sizes (see dotted lines in Fig. 2-3 and Table 1), with a larger impact on the Z/O_3 assimilation. This is consistent with A15, which showed that assimilation of O_3 tends to produce more gravity waves than Z only, where there was very little imbalance. For the small ensemble experiments with Z/O_3 , the optimal length scale also increases when NMI is applied.

3.2 Tuning the hybrid blending coefficient

We next tune the blending coefficient in the hybrid 4DVar system by performing 10-day experiments with values of α ranging from 0.0 to 1.0, in 0.1 increments, for each of the eight experiments. Figure 3-4 (top row) presents the WEP values for Z assimilation as a function of α (NMI results are dotted lines). For each ensemble size, the optimal α is indicated (vertical red line) in addition to the range of α values that produce WEP within 0.5% of the maximum (vertical dashed lines). This range provides an indication of the flatness of the peak and the degree of flexibility in choosing α . Plots of these ranges as a function of α are also provided in Fig. 4-5.

For Z assimilation, the conventional 4DVar ($\alpha=0$) has WEP=67.6% (69.1% with NMI). For each ensemble size, WEP initially increases with α , showing that ensemble covariances provide useful flow-of-the-day information in the system. The optimal blending coefficient for Z assimilation increases from 0.1 for 25 members to 1.0 for 1518 members (see also Fig. 4-5). The latter result indicates that for a “perfect” ensemble (i.e., one that samples the entire error space), the hybrid

system benefits from using as much of the ensemble covariance information as possible. This is expected, since the large ensemble ~~samples all of the dynamical state space~~ has the same number of degrees of freedom as the dynamical state space of the SWM. The WEP values for ~~of~~-Z/O₃ assimilation are provided in the bottom row of Fig. 34. The optimal blending coefficient varies from 0.2 for 50 members to 0.70 for 1518 members (see also Fig. 45). While the optimal blending coefficient increases monotonically for Z assimilation, for Z/O₃ assimilation, the coefficient decreases from 25 to 50 members. The exact cause of this behavior is unknown, but it may be due tuning the system with one length scale for both ozone and height-. That the optimal α for 1518 members is not exactly unity suggests that the ensemble is not perfectly sampling the entire error space when O₃ is included in the state. While the WEP for conventional 4DVar ($\alpha = 0$) is 78%, the peak hybrid WEP is ~86% for 100 members and ~89% for 1518 members. As it will be discussed below, the hybrid provides more benefit over 4DVar when O₃ is assimilated along with the Z, suggesting strong O₃-wind correlations.

Figure 5-6 shows the amount of imbalance entering the system for each experiment. Here we define “imbalance” as the global RMS difference in Z fields before and after NNMI post-processing. For each ensemble size, the imbalance varies with α , with a minimum value that decreases with increasing ensemble size. More ensemble members therefore results in less imbalance in the system. The WEP values for Z assimilation indicate only a slight improvement when NNMI is applied (dotted lines in Fig. 34). The bulk of the improvements with ensemble size in the Z assimilation experiments are likely due to more reliable information in the larger ensembles, rather than to reduced imbalance. As with Z assimilation, the imbalance decreases with ensemble size for Z/O₃ assimilation (Fig. 56, bottom row). For 25 members, the imbalance increases monotonically with α , while for 1518 members the imbalance shows a minimum at $\alpha = 0.7$. There is a slightly larger benefit to applying NNMI for Z/O₃ assimilation than for Z assimilation (dotted lines in Fig. 34), particularly for small ensemble sizes.

4. Discussion of results with optimal tuning

We now ~~compare~~examine the results from all three DA systems when using the optimal hybrid tuning parameters. Figure 6-7 shows the optimal hybrid 10-day WEP values for hybrid (blue) and EnKF (black)(blue) for all ensemble sizes and for 4DVar (red) along with the 4DVar (red) and EnKF (black) results. The hybrid system outperforms the 4DVar, with WEP values increasing with ensemble size for both Z and Z/O₃ assimilation. The hybrid outperforms the EnKF for small ensembles, while at 100 and 1518 members the results are similar. As ensemble size increases, it will likely become more difficult for the hybrid to beat the offline EnKF. There are therefore two limiting values of the hybrid system. The case of one ensemble member would be analogous to conventional 4DVar, while for large ensemble size the hybrid results are limited by the EnKF.

To quantify the added value of O_3 relative to the baseline system that assimilates only Z , Fig. 7-8 shows the difference in global RMS vector wind error between the two sets of runs (Z assimilation and Z/O_3 assimilation). Note that larger positive numbers on Fig. 7-8 indicate smaller wind errors when adding O_3 to the system. The absolute difference (Fig. 78, left) shows that in the 4DVar system (red lines), O_3 reduces the wind error by $\sim 0.32 \text{ ms}^{-1}$. In the hybrid system (blue lines), the wind error reduction is larger at 25, 100, and 1518 members, and similar at 50 members. With NMI applied (dotted lines), the O_3 benefit for the hybrid is larger than for 4DVar at all ensemble sizes. In Fig. 7-8 (right) the error reduction is given as a fractional reduction of the error when only Z is assimilated. The reduction is $\sim 30\%$ for 4DVar, but increases to $\sim 36\text{-}49\%$ for the hybrid (and up to $\sim 56\%$ for 100 members with NMI). These results show that the added value of O_3 to the wind field is larger in the hybrid system than in 4DVar. This highlights the benefits of having initial O_3 -wind covariances in the hybrid system that are not available in conventional 4DVar. The EnKF results (black lines) are also included in Fig. 8. Except for the large ensemble experiments, the relative benefit of adding O_3 is smaller in the EnKF system than in the hybrid. This suggests that the O_3 -wind interaction benefits both from the ensemble covariances as well as the variational DA approach.

Lastly, we examine the wind errors as a function of latitude ~~to see where the benefit of the hybrid system is largest~~. Figure 8 9 shows initial zonal wind errors (thin black lines) along with final 4DVar (red), ~~and~~ hybrid (blue), ~~and~~ EnKF (thick black) errors. ~~Both~~ All three systems show strong reductions from the initial errors. For Z assimilation (top row), the 25-member hybrid shows a zonal wind improvement over 4DVar at high NH latitudes. Since the TR is dynamically forced with a wave centered at 45°N , this result is not surprising. With 50 or more members, the hybrid provides additional improvement in the tropics and parts of the Southern Hemisphere (SH). The hybrid Z assimilation also reduces the meridional wind errors (Fig. 910, top row), ranging from modest NH improvements at 25 members to global improvements at 1518 members. The hybrid system shows generally smaller errors than EnKF for 25 and 50 members, but results for these two DA systems are similar at 100 and 1518 members, as also shown in Fig. 7. Application of NMI does not alter the errors very much for Z assimilation.

The zonal wind errors for Z/O_3 assimilation are plotted in Fig. 8-9 (bottom row). For 25 members, the hybrid system shows reduced errors in the NH and tropics, relative to 4DVar, while there are some slight increases in zonal wind errors near 30°S and 60°S . Why errors would increase at some latitudes when adding ensemble information is unclear, but it might be due to spurious correlations that are not localized. Since the optimization of the length is based on globally-averaged WEP, we might expect some regions to have increased errors. An ensemble localization scale that varies with latitude might be useful to consider here, but this is beyond the scope of this paper. As ensemble size increases, the hybrid errors decrease, although even at 100 members hybrid errors are still slightly larger than 4DVar errors at 60°S . At 1518 members, hybrid errors are smaller at all latitudes, and the tropical peak seen in the 4DVar errors is considerably reduced. This suggests that ensemble Z -wind correlations in the tropics are more reliable than the conventional correlations based on analytic balance assumptions. The hybrid system shows generally smaller errors than the EnKF for Z/O_3 assimilation at 25, 50, and 100 members, but results are similar at 1518 members.

For meridional winds (Fig. 910, bottom), the hybrid system with Z/O₃ assimilation has smaller errors than 4DVar in the NH and in the midlatitude SH for small ensemble sizes, while at large ensemble sizes, the hybrid wind errors are smaller than 4DVar errors at all latitudes. The EnKF shows generally larger errors at small ensemble sizes than the hybrid, with a peak at over 2 ms⁻¹ near 20°S for 25 members. At large ensemble sizes the hybrid and EnKF results are similar. Including O₃ in the DA system reduces the overall wind errors, particularly in the tropics. This is consistent with the EnKF results in A15, which showed reductions in tropical errors when O₃ was assimilated. This is important, since Z observations alone have difficulty constraining the tropical winds, even in the hybrid system. But the O₃-wind tropical correlations have information that can reduce the wind errors there. We note as a caveat, however, that we have not yet attempted to include O₃ chemistry in the system, which may limit the tropical O₃ gradients, particularly in the middle and upper stratosphere. However, in the lower stratosphere, where the O₃ photochemical lifetime is long, we might expect hybrid O₃ assimilation to provide a benefit particularly to the tropical winds. Finally, we note that application of NMI slightly reduces zonal and meridional wind errors for Z/O₃ assimilation for all three DA systems. For 25 members, the reduction occurs at nearly all latitudes, while for 1518 members the reduction is confined to the extratropics.

5. ConclusionsSummary

The problem of wind extraction from tracer observations in hybrid 4DVar data assimilation was examined in this study using a shallow water model (SWM) system coupled to an O₃ advection equation. While previous studies (A14 and A15) examined conventional 4DVar and EnKF simulations, this study combines the best of both systems by blending the ensemble covariance with the conventional covariance at the beginning of the variational assimilation window. The results show that O₃ provides added value in a system already constrained by height (in lieu of temperature for the SWM) observations, and that for small ensemble sizes, relative to the degrees of freedom in the state, the hybrid provides better results than either conventional 4DVar or EnKF.

Using a relatively low resolution system, we were able to probe the limits of the benefit of hybrid covariance blending. Both the optimal localization length and the optimal blending parameter generally increased with ensemble size, so that with large ensemble size (spanning the dynamical state space), the optimal blending is essentially 1.0. For small ensembles (25 or 50 members), values of 0.2 to 0.5 produced better results. With large ensemble size, the hybrid system produced wind errors comparable with the offline EnKF, while for small ensemble size, the hybrid results were closer to 4DVar, suggesting the limiting benefits of hybrid blending. Overall, the hybrid outperforms the 4DVar, suggesting value in combining high-rank conventional background error covariance with localized ensemble flow-of-the-day information when attempting wind extraction from tracers.

While wind extraction potential (WEP) was highest for large ensembles, even small ensembles provided information that benefitted the hybrid system. We should note, however, that the ensemble size relative to the state vector is much larger than it would be currently possible for a full 3D NWP system. Therefore we hesitate to extrapolate the results to a full system. Another caveat is that the truth run we used was relatively smooth, due to the large-scale forcing applied. This may favor the 4DVar, since the tangent linear model (TLM) is likely to do quite well in this regime (see, for example, TLM errors in Fig. 3 of A14). Further tests with more complicated flows, such as the case of barotropic instability, would be valuable to examine the benefit of ensembles in highly nonlinear regimes.

The issue of balance also plays a role in DA with the SWM system. For small ensembles, imbalance generally increases as more ensemble information is added. When nonlinear normal mode initialization (NNMI) is applied as a post-processing diagnostics, it benefits the winds in the hybrid system. The SWM, with minimal diffusion and no other physical parameterizations, is much more sensitive to imbalance than a typical operational system. How these results translate to operational systems is unclear, but at minimum they may provide some guidance as to when filtering (digital filter or NNMI) may be useful.

6. Conclusions

~~Several future directions are being considered for this work with the SWM system. We would like to examine the effects of O_3 photochemistry on the wind extraction. It is likely that when the photochemical lifetime is short, the ability to extract wind from O_3 may be reduced, since advection is not the dominant term on the O_3 continuity equation. We also need to test how this limited resolution study scales upward to more realistic systems. In addition, we plan to completely eliminate the tangent linear model and adjoint using different ensemble variational approaches (e.g., Buchner et al., 2010, 2013, 2015; Lorene et al. 2015; Frolov and Bishop, 2016). Finally, we plan to apply what we've learned from these SWM studies to devise experiments with an operational system, building on the work of Allen et al. (2013), who examined O_3 assimilation in a pre-operational version of the Navy Global Environmental Model.~~ The present work culminates a series of three papers (A14, A15, and the current work) examining the impact of tracer assimilation on winds using three modern operational data assimilation techniques (4DVar, EnKF, and hybrid 4DVar, respectively). The overarching goal of the tracer assimilation on winds is to use tracer data to fill the gaps in direct wind observations, particularly in the upper troposphere, stratosphere, and mesosphere. Since trace gas observations are not generally available at sufficient resolution for deriving feature-track winds, they must be combined with model background information to produce an analysis using a 4D data assimilation system. A pilot study using a full 3-D NWP model with a 4DVar system (Allen et al., 2013) showed that wind extraction from ozone assimilation in the stratosphere is possible, but also highlighted limitations due to geophysical conditions, tracer observation

quality and error specifications, and limited observing sampling patterns. These results motivated the authors to pursue a more detailed theoretical study of the problem using the shallow water model (SWM) framework with a variety of data assimilation systems. Below we provide some overall reflections on the SWM experiments and suggestions for future work.

5 In the first SWM study (A14), we examined the relative benefit of assimilation different tracers (ozone, nitrous oxide, and water vapor) in a conventional 4DVar system. Since the conventional 4DVar does not have correlations between tracers and wind in the initial background error covariances, the only way that tracer assimilation can affect the winds is through the adjoint of the TLM, which propagates sensitivities of the cost function with respect to tracer observations backwards in time. This approach is effective, as long as the background error covariances are correctly modeled. The 4DVar system has an advantage over the ensemble methods in that imbalance appears to play only a minor role. The analysis increments also tend to be smoother than when ensembles are used. One important conclusion of this study was that wind extraction will be easier with certain tracer characteristics (e.g., large background gradients and small observation error standard deviations).

10 The second SWM study (A15), which took the EnKF approach, illustrated the benefits of using ensemble correlations to propagate information from ozone observations to the dynamical variables. This proved to be very effective for extracting wind information from ozone, even in a relatively data-rich environment. The issue of spurious gravity waves played a larger role in EnKF than in 4DVar, due to imbalance caused when localizing the covariance. Imbalance was shown to be reduced by judicious choice of variables and increased ensemble size. However, imbalance is still an issue that needs to be studied further in the context of tracer assimilation using ensemble methods.

15 The current study showed that the largest benefit to the winds from ozone assimilation occurs in the hybrid 4DVar, which combines the benefits of variational DA with flow-of-the-day covariances generated from ensembles. While imbalance is still an issue, the blending of ensemble covariance with conventional covariance reduces the generation of gravity waves. The additional tuning required in the hybrid system does somewhat limit the applicability of the method. However, much of the tuning (e.g., localization) will already be performed in development of the 4DVar and EnKF (or other ensemble approach) systems. The main additional requirement is the tuning of the blending coefficient, which can be coarsely done if the sensitivity is small. The overall conclusion is that hybrid 4DVar offers the most promising approach (of the three DA methods we examined) for tracer-wind extraction in NWP.

20 The current study showed that the largest benefit to the winds from ozone assimilation occurs in the hybrid 4DVar, which combines the benefits of variational DA with flow-of-the-day covariances generated from ensembles. While imbalance is still an issue, the blending of ensemble covariance with conventional covariance reduces the generation of gravity waves. The additional tuning required in the hybrid system does somewhat limit the applicability of the method. However, much of the tuning (e.g., localization) will already be performed in development of the 4DVar and EnKF (or other ensemble approach) systems. The main additional requirement is the tuning of the blending coefficient, which can be coarsely done if the sensitivity is small. The overall conclusion is that hybrid 4DVar offers the most promising approach (of the three DA methods we examined) for tracer-wind extraction in NWP.

25 Several future directions are being considered for this work with the SWM system. We would like to examine the effects of O₃ photochemistry on the wind extraction. It is likely that when the photochemical lifetime is short, the ability to extract wind from O₃ may be reduced, since advection is not the dominant term on the O₃ continuity equation. We would also like need to test how this limited resolution study scales upward to more realistic systems. Further work is needed in separate tuning of the localization lengths for different variables, as well as filtering out gravity waves within the hybrid 4DVar

5 ~~system. In addition, we would like to plan to completely explore eliminating~~ the tangent linear model and adjoint using different ensemble variational approaches (e.g., Buehner et al., 2010, 2013, 2015; Lorenc et al. 2015; Frolov and Bishop, 2016). Finally, ~~we plan~~ we plan to apply what we've learned from these SWM studies to devise experiments with an operational hybrid 4DVar system, building on the work of Allen et al. (2013), who examined wind benefits from O₃ assimilation in a pre-operational version of the Navy Global Environmental Model 4DVar system.

Acknowledgments

~~We would like to thank Thomas Milewski, one anonymous reviewer, and the editor for helpful comments on the manuscript.~~

10 ~~This work was funded by the U. S. Office of Naval Research.~~ Douglas R. Allen and Karl W. Hoppel acknowledge support from Office of Naval Research base funding via Task BE-033-02-42. David D. Kuhl and Karl W. Hoppel acknowledge support from Office of Naval Research base funding via Task BE-435-050.

References

- Allen, D. R., Hoppel, K. W., Nedoluha, G. E., Kuhl, D. D., Baker, N. L., Xu, L., and Rosmond, T. E.: Limitations of wind extraction from 4D-Var assimilation of ozone, *Atmos. Chem. Phys.*, 13, 3501-3515, doi:10.5194/acp-13-3501-2013, 2013.
- 15 Allen, D. R., Hoppel, K. W., and Kuhl, D. D.: Wind extraction potential from 4D-Var assimilation of stratospheric O₃, N₂O, and H₂O using a global shallow water model, *Atmos. Chem. Phys.*, 14, 3347-3360, doi:10.5194/acp-14-3347-2014, 2014.
- Allen, D. R., Hoppel, K. W., and Kuhl, D. D.: Wind extraction potential from ensemble Kalman filter assimilation of stratospheric ozone using a global shallow water model, *Atmos. Chem. Phys.*, 15, 5835-5850, doi:10.5194/acp-15-5835-2015, 2015.
- 20 Anderson, J. L.: An adaptive covariance inflation error correction algorithm for ensemble filters, *Tellus*, 59A, 210-224, doi:10.1111/j.1600-0870.2006.00216.x, 2007.
- Andersson, E., E. Hólm, P. Bauer, A. Beljaars, G. A. Kelly, A. P. McNally, A. J. Simmons, J.-N. Thépaut, and Tompkins, A. M.: Analysis and forecast impact of the main humidity observing systems, *Q. J. Roy. Meteor. Soc.*, 133, 1473-1485, doi:10.1002/qj.112, 2007.
- 25 Bonavita, M., Isaksen, L., and Hólm, E., 'On the use of EDA background error variances in the ECMWF 4D-Var,' ECMWF Tech Memo 664. Available at <http://old.ecmwf.int/publications/library/do/references/show?id=90381> (last access: 21 April 2015), 2012.

- Buehner, M., Houtekamer, P. L., Charette, C., Mitchell, H. L., and He, B.: Intercomparison of variational data assimilation and the ensemble Kalman filter for global deterministic NWP. Part II: One-month experiments with real observations, *Mon. Wea. Rev.*, 138, 1567-1586, doi:10.1175/2009MWR3158.1, 2010.
- Buehner, M., Morneau, J., and Charette, C.: Four-dimensional ensemble-variational data assimilation for global deterministic weather prediction, *Nonlin. Processes Geophys.*, 20, 669-682, doi:10.5194/npg-20-669-2013, 2013.
- Buehner, M., McTaggart-Cowan, R., Beaulne, A., Charette, C., Garand, L., Heilliette, S., Lapalme, E., Laroche, S., Macpherson, S. R., Morneau, J., and Zadra, A.: Implementation of deterministic weather forecasting systems based on ensemble-variational data assimilation at Environment Canada. Part I: The global system, *Mon. Wea. Rev.*, 143, 2532-2559, doi:10.1175/MWR-D-14-00354.1, 2015.
- 10 Clayton, A. M., Lorenc, A. C., and Barker, D. M.: Operational implementation of a hybrid ensemble/4D-Var global data assimilation system at the Met Office, *Q. J. R. Meteorol. Soc.*, 139, 1445–1461, doi:10.1002/qj.2054, 2013.
- Daley, R.: *Atmospheric Data Analysis*, Cambridge University Press, Cambridge, pp 150-185, 1991.
- Daley, R.: Estimating the wind-field from chemical-constituent observations - experiments with a one-dimensional extended Kalman filter, *Mon. Wea. Rev.*, 123, 181-198, doi: 10.1175/1520-0493(1995)123<0181:ETWFFC>2.0.CO;2, 1995.
- 15 Daley, R.: Recovery of the one and two dimensional windfields from chemical constituent observations using the constituent transport equation and an extended Kalman filter, *Meteorol. Atmos. Phys.*, 60, 119-136, 1996.
- Daley, R., and Barker, E.: NAVDAS Source Book 2001, Naval Research Laboratory Publication NRL/PU/7530—01-441, 163 pp., available at <http://www.dtic.mil/docs/citations/ADA396883>, 2001.
- Dragani, R., and McNally, A. P.: Operational assimilation of ozone-sensitive infrared radiances at ECMWF, *Q. J. Roy. Meteor. Soc.*, 139, 2068-2080, doi:10.1002/qj.2106, 2013.
- 20 Evensen, G.: The ensemble Kalman filter: theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343-367, doi:10.1007/s10236-003-0036-9, 2003.
- Frolov, S., and C. Bishop: Localized ensemble-based tangent linear models and their use in propagating hybrid error covariance models, *Mon. Wea. Rev.*, [144, 1383-1405](#), doi:10.1175/MWR-D-15-0130.1, [in press](#), 2016.
- 25 Gaspari, G., and Cohn, S. E.: Construction of correlation functions in two and three dimensions, *Q. J. Roy. Meteor. Soc.*, 125, 723-757, doi:10.1002/qj.49712555417, 1999.
- Han, W., and McNally, A. P.: The 4D-Var assimilation of ozone-sensitive infrared radiances measured by IASI, *Q. J. Roy. Meteor. Soc.*, 136, 2025-2037, doi:10.1002/qj.708, 2010.

- Houtekamer, P. L., and Mitchell, H. L.: Data assimilation using an ensemble Kalman filter technique, *Mon. Wea. Rev.*, 126, 796-811, doi: 10.1175/1520-0493(1998)126<0796:Dauaek>2.0.Co;2, 1998.
- Houtekamer, P. L., and Mitchell, H. L.: A sequential ensemble Kalman filter for atmospheric data assimilation, *Mon. Wea. Rev.*, 129, 123-137, doi: 10.1175/1520-0493(2001)129<0123:Asekff>2.0.Co;2, 2001.
- 5 Kepert, J. D.: Covariance localisation and balance in an ensemble Kalman filter, *Q. J. Roy. Meteor. Soc.*, 135, 1157-1176, doi:10.1002/qj.443, 2009.
- Kleist, D. T., and Ide, K.: An OSSE-based evaluation of hybrid variational-ensemble data assimilation for the NCEP GFS. Part II: 4DEnVar and hybrid variants, *Mon. Wea. Rev.*, 143, 452-470, doi:10.1175/MWR-D-13-00350.1, 2015.
- Kuhl, D. D., Rosmond, T. E., Bishop, C. H., McLay, J., and Baker, N. L.: Comparison of Hybrid Ensemble/4DVar and
10 4DVar within the NAVDAS-AR data assimilation framework, *Mon. Wea. Rev.*, 141, 2740-2758, doi:10.1175/MWR-D-12-00182.1, 2013.
- Lorenc, A. C.: The potential of the ensemble Kalman filter for NWP – a comparison with 4D-Var, *Q. J. Roy. Meteor. Soc.*, 129, 3183-3203, doi:10.1256/qj.02.132, 2003.
- Lorenc, A. C., Bowler, N. E., Clayton, A. M., Pring, S. R., and Fairbairn, D.: Comparison of Hybrid-4DEnVar and Hybrid-
15 4DVAR data assimilation methods for global NWP, *Mon. Wea. Rev.*, 142, 212-229, doi:10.1175/MWR-D-14-00195.1, 2015.
- Milewski, T., and Bourqui, M. S.: Assimilation of stratospheric temperature and ozone with an ensemble Kalman filter in a chemistry-climate model, *Mon. Wea. Rev.*, 139, 3389-3404, doi:10.1175/2011mwr3540.1, 2011.
- [Machenhauer, B.: On the dynamics of gravity oscillations in a shallow water model, with applications to normal mode initialization, *Contrib. Atmos. Phys.*, 50, 253-271, 1977.](#)
- 20 Peubey, C., and McNally, A. P.: Characterization of the impact of geostationary clear-sky radiances on wind analyses in a 4D-Var context, *Q. J. Roy. Meteor. Soc.*, 135, 1863-1876, doi: doi:10.1002/qj.500, 2009.
- Peuch, A., Thépaut, J. N., and Pailleux, J.: Dynamical impact of total-ozone observations in a four-dimensional variational assimilation, *Q. J. Roy. Meteor. Soc.*, 126, 1641-1659, doi: 10.1002/qj.49712656605, 2000.
- Riishøjgaard, L. P.: On four-dimensional variational assimilation of ozone data in weather-prediction models, *Q. J. Roy. Meteor. Soc.*, 122, 1545-1571, doi:10.1002/qj.49712253505, 1996.
- 25 Meteor. Soc., 122, 1545-1571, doi:10.1002/qj.49712253505, 1996.
- Rosmond, T., and Xu, L.: Development of NAVDAS-AR: non-linear formulation and outer loop tests, *Tellus*, 58A, 45-58, 2006.
- Semane, N., Peuch, V. H., Pradier, S., Desroziers, G., El Amraoui, L., Brousseau, P., Massart, S., Chapnik, B., and Peuch, A.: On the extraction of wind information from the assimilation of ozone profiles in Meteo-France 4-D-Var operational
30 NWP suite, *Atmos. Chem. Phys.*, 9, 4855-4867, doi:10.5194/acp-9-4855-2009, 2009.

Xu, L., Rosmond, T., and Daley, R.: Development of NAVDAS-AR: formulation and initial tests of the linear problem, *Tellus*, 57A, 546-559, 2005.

Experiment	L (km)	WEP-EnKF (%)	WEP-4DVar (%)	WEP-Hybrid (%)	α (unitless)
Z			67.6 (69.1)		
25 members	6000 (6000)	62.5 (63.9)		70.0 (71.0)	0.1 (0.1)
50 members	7500 (7500)	70.5 (71.1)		72.5 (73.8)	0.4 (0.9)
100 members	14000 (13500)	75.3 (75.7)		75.5 (76.5)	0.8 (0.8)
1518 members	15000 (15000)	77.5 (77.8)		78.3 (79.3)	1.0 (1.0)
Z/O_3			77.1 (78.7)		
25 members	3500 (5000)	73.9 (79.1)		81.3 (84.8)	0.3 (0.5)
50 members	5500 (6000)	78.3 (81.8)		82.2 (84.8)	0.2 (0.4)
100 members	7500 (8000)	84.9 (87.8)		86.3 (90.0)	0.5 (0.5)
1518 members	20000 (20000)	89.2 (90.3)		89.1 (90.7)	0.7 (1.0)

Table 1. Results for the optimal runs (i.e., maximum wind extraction potential (WEP), in %), for each experiment. The localization length (L) is provided along with WEP. Results with NMI applied as post-processing of the analysis fields are provided in parentheses.




Fig. 1. Schematic diagram of hybrid system. The 4DVar uses a 6-h window, while the offline EnKF uses a 20-min window. At the beginning of the analysis window, information is passed from the EnKF to 4DVar by blending the covariances.

5|




Fig. 2. (a) Global RMSE of the zonal wind (in ms^{-1}), and (b) Wind Extraction Potential (in %) for the optimal 100 member hybrid 4DVar assimilation of Z and O₃. Black circles indicate the values at the start of the assimilation experiment.




Fig. 23. Wind extraction potential (WEP, in %) as a function of localization length scale calculated from offline EnKF experiments assimilating Z (top row) and Z/O₃ (bottom row) for ensembles with 25, 50, 100, and 1500 members (columns 1, 2, 3, and 4, respectively). Solid (dotted) lines indicate results without (with) NMI post-processing. Red lines denote the length scale that resulted in the maximum WEP (without NMI post-processing).




Fig. 34. Wind extraction potential (in %) versus blending coefficient for Z assimilation (top row) and Z/O_3 assimilation (bottom row) for ensembles with 25, 50, 100, and 1500 members (columns 1, 2, 3, and 4, respectively). Solid (dotted) lines indicate results without (with) NMI post-processing. Red lines denote the blending coefficients that resulted in the maximum WEP (without NMI post-processing). Dashed lines indicate range of blending coefficients that resulted in WEP values within 0.5% of the maximum.




Fig. 45. Optimal blending coefficient, α (unitless), as a function of ensemble size for Z assimilation (top) and Z/O_3 assimilation (bottom). The range values in red indicate hybrid experiments with WEP values within 0.5% of the maximum WEP for each ensemble size.

5




Fig. 56. Imbalance (in m) versus blending coefficient for Z assimilation (top row) and Z/O_3 assimilation (bottom row) for ensembles with 25, 50, 100, and 1500 members (columns 1, 2, 3, and 4, respectively).




Fig. 67. Maximum WEP (in %) as function of ensemble size for Z assimilation (top) and Z/O₃ (bottom) assimilation. WEP is shown for 4DVar (red), EnKF (black), and hybrid (blue).








Fig. 78. Global RMS vector wind error differences between experiments that assimilate Z assimilation only and experiments that assimilate both and Z and O₃ assimilation. For each ensemble size, the final vector wind errors are calculated for each experiment and then the differences are taken. Left: absolute difference in ms⁻¹. Right: fractional difference (i.e., the difference divided by the vector wind error for Z only assimilation) in %. Red is 4DVar, and blue is hybrid, and black is EnKF. Solid (dotted) lines indicate results without (with) NMI post-processing. Note that larger positive numbers indicate smaller errors due to adding O₃.







Fig. 89. Zonal wind errors (in ms^{-1}) as a function of latitude for initial conditions (thin black), 4DVar (red), hybrid (blue), EnKF (thick black), and 4DVar (red) for Z assimilation (top row) and Z/O₃ assimilation (bottom row) for ensembles with 25, 50, 100, and 1500 members (columns 1, 2, 3, and 4, respectively). Solid (dotted) lines indicate results without (with) NMI post-processing.






Fig. 910. Same as Fig. 89, but for meridional wind.