Colorado air quality impacted by long-range transported aerosol: A set of case studies during the 2015 Pacific Northwest fires

- 8 ²NOAA Earth System Research Laboratory, Physical Sciences Division, Boulder, CO, 80305, USA
- 9 ³NOAA Earth System Research Laboratory, Chemical Sciences Division, Boulder, CO, 80305, USA
- ⁴HORIBA Instruments Inc., Process and Environmental, Irvine, CA, 92618, USA
- 11 *Correspondence to*: Jessie M. Creamean (jessie.creamean@noaa.gov)

12 Abstract. Biomass burning plumes containing aerosols from forest fires can be transported long distances, which can 13 ultimately impact climate and air quality in regions far from the source. Interestingly, these fires can inject aerosols 14 other than smoke into the atmosphere, which very few studies have evidenced. Here, we demonstrate a set of case 15 studies of long-range transport of mineral dust aerosols in addition to smoke from numerous forest fires in the Pacific 16 Northwest to Colorado, U.S. These aerosols were detected in Boulder, Colorado along the Front Range using Betaray attenuation and energy dispersive X-ray fluorescence spectroscopy, and corroborated with satellite-borne lidar 17 18 observations of smoke and dust. Further, we examined the transport pathways of these aerosols using air mass 19 trajectory analysis and regional and synoptic scale meteorological dynamics. Three separate events with poor air 20 quality and increased mass concentrations of metals from biomass burning (S and K) and minerals (Al, Si, Ca, Fe, and 21 Ti) occurred due to the introduction of smoke and dust from regional and synoptic scale winds. Cleaner time periods 22 with good air quality and lesser concentrations of biomass burning and mineral metals between the haze events were 23 due to the advection of smoke and dust away from the region. Dust and smoke present in biomass burning haze can

- 24 have diverse impacts on visibility, health, cloud formation, and surface radiation. Thus, it is important to understand
- 25 how aerosol populations can be influenced by long-range transported aerosols, particularly those emitted from large
- 26 source contributors such as wildfires.
- 27 **Keywords.** Aerosol transport, air quality, mineral dust, biomass burning, remote sensing, in situ observations

28 1 Introduction

- 29 Wildfires in both forested and agricultural regions serve as a steady source of pollutants into the atmosphere. Gas
- 30 phase constituents such as methane (CH_4), carbon monoxide (CO_2), carbon dioxide (CO_2), sulphur dioxide (SO_2) and
- 31 nitrogen oxides $(NO_x; NO + NO_2)$ can be produced from burning of biofuels (Gadi et al., 2003; Radojevic, 2003), in
- 32 addition to precursors that induce ozone production (Jaffe and Wigder, 2012). Additionally, wildfires produce large
- 33 concentrations of aerosols which are injected into the atmosphere or formed in the smoke plume via secondary

Jessie M. Creamean^{1,2*}, Paul. J. Neiman², Timothy Coleman^{1,2}, Christoph J. Senff^{1,3}, Guillaume
 Kirgis^{1,3}, Raul J. Alvarez³, and Atsushi Yamamoto⁴

 ¹University of Colorado at Boulder, Cooperative Institute for Research in Environmental Sciences, Boulder, CO,
 80309, USA

34 processes and include carbonaceous species (elemental and organic carbon) (Park et al., 2003; Spracklen et al., 2007) 35 and biogenic heavy metals (including but not limited to Fe, Mn, Cd, Cu, Pb, Cr, and Ni) (Nriagu, 1989; Radojevic, 36 2003). Soluble inorganic species such as sulphate, nitrate, ammonium, and chloride are found in fire emissions and 37 partitioned to the particle phase through heterogeneous reactions with the gas phase species released during the 38 combustion process (Pio et al., 2008). Strong, turbulent winds inside combustion zones from controlled and wild 39 vegetation fires can introduce considerable amounts of dust particles into the free troposphere, which can subsequently 40 be transported over thousands of kilometres with the smoke (Clements et al., 2008; Ansmann et al., 2009; Baars et al., 41 2011). Forest fires tend to be much larger than agricultural fires, and enable injection of smoke high into the free 42 troposphere (Colarco et al., 2004; Peterson et al., 2014). Yet, smoke from agricultural, and shrub and grassland fires 43 can still be transported long distances. However, few studies have documented how wildfires from any of the 44 aforementioned biofuel sources inject mineral dust into the atmosphere (Gaudichet et al., 1995; Chalbot et al., 2013; 45 Yang et al., 2013; Nisantzi et al., 2014), particularly in heavily forested or agricultural regions such as the Pacific 46 Northwest of the U.S. where dust sources are limited relative to arid regions in Africa, the Middle East, and Asia. 47 Prescribed burning (i.e., slash-and-burn techniques) and wildfires are common in these arid "dust belt" regions, 48 inducing the simultaneous emission of dust and smoke (Streets et al., 2003; Pinker et al., 2010).

49

50 Aerosols produced directly from wildfires (i.e., carbonaceous and soluble inorganic particulates) or injected into the 51 free troposphere from smoke plume dynamics (i.e., mineral dust) have diverse effects on climate and air quality. For 52 instance, absorbing aerosols such as soot from fires enhance the semi-direct effect that affect cloud and atmospheric 53 lapse rate, particularly when the absorbing aerosols are above cloud (Ge et al., 2014). Further, hygroscopic organic 54 aerosol, sulphate, and nitrate can enable aerosols to serve as cloud condensation nuclei (CCN) (Cruz and Pandis, 55 1997), whereas mineral dust and black carbon are effective ice nucleating particles (INPs) at sub-freezing temperatures 56 (DeMott et al., 1999; DeMott et al., 2003; Vali et al., 2015). Both of these aerosol nuclei modify cloud radiative 57 properties, lifetime, and impact precipitation formation, and have been shown to originate from prescribed burns and 58 wildfires (Eagan et al., 1974; McCluskey et al., 2014). Enhanced pollutants from fires also severely influence air 59 quality, and can prompt adverse health effects (Bravo et al., 2002; Phuleria et al., 2005; Wiedinmyer et al., 2006). For 60 instance, smoke plumes from wildfires have been linked to childhood mortality (Jayachandran, 2008), asthma 61 (Bowman and Johnston, 2005), and various respiratory illness and diseases (Mott et al., 2002; Moore et al., 2006). 62 These effects are additionally complicated by aging from biogenic gases in the smoke plume during transport. Further, 63 previous air quality studies on the East Coast of the U.S. have shown that enhanced aerosol optical depths associated 64 with both wildfires and anthropogenic sources can cause large errors in meteorological models used to forecast poor 65 air quality events (Zamora et al., 2005). Overall, the aerosol species emitted or formed from wildfire plumes are 66 complex in nature and possess several diverse climate and health effects, thus demonstrating the need to better 67 understand the various types, sources, and transport pathways of these emissions.

68

Air quality is strongly dependent not only on emission sources such as wildfires, but also on weather and climate change (Jacob and Winner, 2009). Regions with complex topography such as the Front Range of Colorado, U.S. (see 71 Figure 1) have unique meteorological phenomena such as upslope/downslope flows that serve as agents for focusing 72 or cleaning out local air pollution from the Denver metropolitan area (Haagenson, 1979). Typically, this region is 73 characterized by good air quality in terms of particulate matter (PM) relative to other larger urban and industrial areas, 74 although it experiences occasional pollution episodes due to modulation of the mountain slope dynamics, oil and 75 natural gas production, and wildfires (Watson et al., 1998; Sibold and Veblen, 2006; Brown et al., 2013). Here, we 76 show that the Front Range air quality was severely impacted by long-range transported wildfire emissions from the 77 Pacific Northwest during August 2015. A reoccurring influx of smoke aerosols infiltrated the Front Range region due 78 to shifts in regional and synoptic scale meteorology. Interestingly, mineral dust was also transported with the smoke 79 plume to the Front Range from the wildfires. This complex mixture of aerosols can have numerous climate and health 80 effects in the region, and should be evaluated to develop a better understanding of future influences from wildfire 81 emissions, especially considering a warmer and drier climate will potentially lead to more frequent wildfires 82 (Westerling et al., 2006; Liu et al., 2010).

83 2 Methods

84 **2.1 Satellite observations**

85 The source of aerosols from the fires was determined using imagery from the Moderate Resolution Imaging 86 Spectroradiometer (MODIS) on board the Terra satellite. MODIS is a multi-spectral sensor with 36 spectral bands, 87 ranging in wavelength from 0.4 to 14.2 µm. Aerosol optical depth (AOD) data at 550 nm from MODIS were acquired 88 from the Giovanni data server (http://giovanni.gsfc.nasa.gov/giovanni/) for daily AOD at a 1° spatial resolution using 89 a domain of 82°W to 163°W and 26°N to 59°N (MOD08 D3 051). MODIS AOD is retrieved from three spectral 90 channels (0.47 µm, 0.66 µm, and 2.1 µm) using the algorithm described by Kaufman et al. (1997) in cloud-free pixels 91 (10 km x 10 km grid box) (Ackerman et al., 1998). Fire and surface thermal anomaly data were also acquired from 92 the MODIS Terra satellite using brightness temperature measurements in the 4-um and 11-um channels 93 (https://earthdata.nasa.gov/labs/worldview/) (Giglio, 2010). The fire detection strategy is based on absolute detection 94 of a fire (when the fire strength is sufficient to detect), and on detection relative to its background (to account for 95 variability of the surface temperature and reflection by sunlight) (Giglio et al., 2003). The algorithms include masking 96 of clouds, bright surfaces, glint, and other potential false alarms (Giglio et al., 2003). Swaths from overpasses over 97 the Pacific Northwest were used to determine the locations of fires on a daily basis.

98

99 In order to evaluate the types of aerosols present in enhanced AOD plumes over the western U.S., aerosol subtype 100 data were retrieved from Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) on board Cloud-Aerosol Lidar 101 and Infrared Pathfinder Satellite Observations (CALIPSO). Level-2 ValStage1 V.30 Vertical Feature Mask data 102 obtained from NASA's Earth Observing System Data and Information System (EOSDIS: 103 https://search.earthdata.nasa.gov/) contain vertically-resolved data of aerosol layer sub-type, including but not limited 104 to smoke, dust, and polluted dust (i.e., dust mixed with smoke) (Vaughan et al., 2004; Omar et al., 2009; Winker et 105 al., 2009). CALIPSO was launched on 28 Apr 2006 and flies in an orbital altitude of 705 km as part of the sun-

- 106 synchronous "A-train" satellite constellation. CALIOP is an elastic backscatter lidar operating at 532 nm and 1064
- 107 nm, completed with a depolarization channel at 532 nm to enable detection of aerosols and clouds. Granule data were
- 108 acquired from orbital swaths that passed over the north-western U.S. (domain includes Washington, Oregon, northern
- 109 California, Idaho, Nevada, Montana, Wyoming, Utah, and Colorado) from 15 Aug to 2 Sep 2015 and processed using
- 110 modified Python code developed by the Hierarchical Data Format (HDF) group at the University of Illinois, Urbana-
- 111 Champaign (http://hdfeos.org/). Aerosol sub-types were also examined off the U.S. West Coast across the central
- 112 North Pacific Ocean, in the context of air mass trajectory analysis, to ensure mineral dust and smoke were transported
- 113 to Colorado from the Pacific Northwest fires rather than from deserts or fires overseas.

114 **2.2 Colorado air quality data**

- All air quality data were acquired from the Colorado Department of Public Health and Environment (CDPHE; 115 116 http://www.colorado.gov/airquality/report.aspx) from 15 Aug to 2 Sep 2015 at various sites throughout the Colorado Front Range (see Figure 1). The DESCI site (Denver Visibility Station; 39.73°N, 104.96°W; 1,633 m MSL) is 117 118 highlighted in blue, near downtown Denver, where horizontal atmospheric extinction (km⁻¹) data measured with a transmissometer are available through CPDHE. These data provide a quantitative measure of "haziness" indicated 119 120 throughout the text. Table 1 provides the site latitudes, longitudes, elevations, and which PM measurements were 121 available at each site. Hourly measurements included mass concentrations (µg m⁻³) of particulate matter for particles 122 with diameters $\leq 2.5 \ \mu m \ (PM_{2.5})$ and $\leq 10 \ \mu m \ (PM_{10})$. All times shown are coordinated universal time [UTC; local 123 time or mountain daylight time (MDT) + 6].

124 2.3 In situ aerosol observations at Boulder, Colorado

Real-time, hourly ambient aerosol samples were analysed for PM_{2.5} mass concentrations (µg m⁻³) and concentrations 125 126 of various metals (ng m⁻³) using the HORIBA, Ltd. PX-375 continuous particle mass and elemental speciation monitor 127 (http://www.horiba.com/process-environmental/products/ambient/details/continuous-particulate-monitor-with-x-rayfluorescence-px-375-27871/) from 26 Aug to 2 Sep 2015 at the National Oceanic and Atmospheric Administration 128 129 (NOAA) David Skaggs Research Centre (DSRC) located in Boulder, Colorado (39.99°N, 105.26°W, and 1672 m MSL; see Figure 1). The PX-375 draws in air at 16.7 L min⁻¹ through a U.S. Environmental Protection Agency (EPA) 130 Louvered PM₁₀ inlet, then subsequently passes through a BGI Very Sharp Cut Cyclone (VSCCTM) to filter for particles 131 132 smaller than 2.5 µm in diameter. Air is pulled through a nozzle for 60 minutes per hourly sample, where particles are subsequently deposited in a 100-mm diameter spot on TeflonTM PTFE fabric filter tape for analysis. Once the sample 133 134 is collected for 60 minutes, beta-ray attenuation and energy dispersive X-ray fluorescence spectroscopy (EDXRF) 135 analyses are conducted for 60 minutes and 1000 seconds, respectively, per hourly sample, simultaneous to the 136 collection of the subsequent sample. Beta-ray attenuation analysis is used to measure total PM2.5 mass concentrations 137 and EDXRF is used to analyse concentrations of Ti, V, Cr, Mn, Fe, Ni, Cu, Zn, As, Pb, Al, Si, S, K, and Ca. The 138 EDXRF unit contains a CMOS camera for sample images. Calibration material used for X-ray intensity is NIST SRM 139 2783. Lower detection limits (LDLs) are shown in Table 2 and error was calculated to be $\pm 2\%$ for hourly metal 140 concentrations. Hourly total PM_{2.5} mass concentrations had an LDL of 2.00 µg m⁻³.

141 2.4 Aerosol and ozone remote sensing observations at Boulder, Colorado

142 The Tunable Optical Profiler for Aerosol and oZone (TOPAZ) lidar was operated at the DSRC on 9 days from 14 Aug through 2 Sep 2015 and it collected about 62 hours of ozone and aerosol profile data, primarily between mid-morning 143 144 and early evening local time. TOPAZ is a state-of-the-art, tunable ozone differential absorption lidar. It emits pulsed 145 laser light at three ultraviolet wavelengths between 285 and 295 nm and measures ozone as well as aerosol backscatter and extinction profiles with high temporal and spatial resolutions (Alvarez et al., 2011). The TOPAZ lidar is mounted 146 147 in a truck with a rooftop two-axis scanner. This scanner permits pointing the lidar beam at elevation angles between 148 -5 and 30 degrees at a fixed but changeable azimuth angle. To achieve zenith operation the scanner mirror is moved 149 out of the beam path. Typical TOPAZ operation consists of a scan sequence at 2, 6, 20, and 90 degrees elevation, 150 repeated approximately every five minutes. The range-resolved ozone and aerosol observations at the shallow 151 elevations angles are projected onto the vertical and spliced together with the zenith observations, resulting in 152 composite vertical ozone and aerosol profiles from about 15 m to 2-3 km above ground level (AGL) at five minutes 153 time resolution (Alvarez et al., 2012). In this study, we only used the lidar aerosol extinction profiles measured at a wavelength of 294 nm. The aerosol profile retrieval requires assumptions about the lidar calibration constant and the 154 155 aerosol extinction-to-backscatter or lidar ratio. For this study we used an altitude-constant lidar ratio of 40 sr, which is a good approximation for continental and urban aerosols. The lidar signal at the aerosol wavelength of 294 nm is 156 157 also affected by ozone absorption. Therefore, uncertainties in the ozone observations can cause biases in the aerosol 158 retrieval. This, combined with uncertainties in the calibration constant and lidar ratio, can lead to errors in the aerosol 159 extinction coefficient profiles of up to about 30%. The precision of the 5-minute aerosol extinction measurements is

160 typically better than 10%.

161 **2.5 Meteorological data and analysis**

A gridded perspective of synoptic-scale conditions across North America was provided using the NOAA/National Centres for Environmental Prediction (NCEP) Rapid Refresh numerical data package [RAP; http://rapidrefresh.noaa.gov/ (Benjamin et al., 2016)]. The RAP is an operational assimilation/modelling system updated hourly, with 13-km horizontal resolution and 50 vertical levels.

166

Air mass backward trajectory analyses were conducted using HYSPLIT 4 (Draxler and Rolph, 2011) and data from 167 168 the NOAA/NCEP Global Data Assimilation System (GDAS) (Kalnay et al., 1996). HYSPLIT trajectories do not include processes that may affect particle concentrations such as convective transport, wet removal, or dry removal, 169 170 and are only intended to highlight the possible transport pathways. To study the potential for transport from the Pacific 171 Northwest fires region, and to eliminate potential contribution from aerosol sources overseas, we used an ensemble of 172 backward trajectories initiated at multiple altitudes and times ending above the NOAA building in Boulder. Ten-day 173 back trajectories were initiated every 6 hours (at 00:00, 06:00, 12:00, and 18:00 UTC) during 15 Aug-2 Sep 2015 at 174 500, 1000, and 2000 m AGL (corresponds to 2172, 2672, and 3672 m MSL).

- 176 A 449-MHz wind profiler (White et al., 2013), deployed near the Boulder Atmospheric Observatory in Erie, Colorado
- 177 (BAO; 40.05N, 105.01°W, and 1577 m MSL; location shown in Figure 1), provided hourly-averaged profiles of
- horizontal wind. The high (low) mode extended from 145 m (195 m) to 10074 m (5059 m) AGL with a vertical
- resolution of 200 m (100 m). The wind-profiler data were edited objectively using the vertical-temporal continuity
- 180 method of Weber et al. (1993) and then subjected to additional manual editing as needed. For the purpose of this study,
- 181 we utilized only the low-mode observations.

182 3 Results and discussion

183 **3.1 Haze events induced poor air quality along Colorado's Front Range**

The shift in air quality was evident during three August haze events in the Denver metro area. Figure 2 shows photos 184 185 of notable air quality transitions in Denver looking westward towards the foothills of the Rocky Mountains and Figure 186 3 shows the atmospheric extinction measurements from DESCI. Higher values of extinction indicate hazier conditions. 187 The image on 15 Aug shows typical, clean conditions, where the foothills were visible west of Denver. Extinction was 188 also relative low on 15 Aug. On 17 Aug, a haze settled in the region, creating a low-level pollution plume that masked 189 the view of the foothills. This haze continued to infiltrate the Denver metro area, reaching the poorest visibility (i.e., 190 highest extinction) on 23 Aug. This haze persisted in the Denver metro area until 27 Aug, when clear conditions were re-established and the foothills were once again visible. However, the air quality deteriorated again by 29 Aug, with 191 192 hazy conditions obscuring the foothills. This haze event was shorter lived, clearing out on 31 Aug. The cleaner 193 conditions persisted until the end of the measurement period on 2 Sep. The qualitative observations of the three 194 separate haze events were corroborated by in situ air quality measurements along the Front Range. Figure 3 also shows hourly and daily averaged PM_{2.5} mass concentrations (herein, simply called "PM_{2.5}") at the sites provided in Table 1. 195 196 Overall, three separate haze events occurred along the Front Range with the worst days visually observed (Figure 2) 197 on the 17, 23, and 29 Aug (events 1, 2, and 3, respectively), when extinction was highest, $PM_{2.5}$ reached maximum concentrations, and a cold front passed through (discussed in section 3.3). Prior to each of these events, PM_{2.5} was 198 suppressed then slowly increased to each event's maximum concentrations on 17, 23, and 29 Aug. PM_{2.5} slowly 199 200 decreased following each of these haze events. PM₁₀ (not shown) did not follow similar increases and decreases as 201 the $PM_{2.5}$, suggesting the smaller particles contributing to $PM_{2.5}$ originated from different, likely more distant sources 202 as compared to coarser particles contributing to the PM₁₀, which are likely from more local sources (VanCuren, 2003; 203 Neff et al., 2008).

204 **3.2 Biomass burning plume propagates towards Colorado**

During the 15 Aug–2 Sep time period, fires in high elevation (>3000 feet MSL) forested areas and to some extent in shrub and grasslands in the Pacific Northwest were prominent, while few fire hotspots were located in low elevation agricultural land (see Figure S1 of the Supporting Information). Figures 4–6 show MODIS retrievals of fire hotspots and aerosol optical depth (AOD) during the first, second, and third haze events in Colorado, when numerous fires

209 were detected in Washington, Oregon, northern California, northern Idaho, and north-western Montana. Three cases

are defined as the time periods surrounding and including the haze event days: Case 1 (15–18 Aug), Case 2 (20–23

- 211 Aug), and Case 3 (26–29 Aug).
- 212

213 On 15 Aug, prior to the onset of the first haze event in Colorado, the plume of enhanced AOD propagating from the 214 fires in the Pacific Northwest remained north of Colorado in Montana and southern Canada (Figure 4). The air above 215 the Denver/Boulder area contained relatively diminished AOD (0.12, averaged from the domain of 39.5°N, 104.5°W, 216 40.5°N, and 105.5°W). Although the core of the plume remained north of Colorado, its more diffuse southern region 217 drifted south-eastward on 16 Aug. By 17 Aug, enhanced AOD was observed along the Front Range in northcentral 218 Colorado near Denver/Boulder (0.37). The AOD slightly decreased on 18 Aug over Denver/Boulder (0.25), which is 219 supported by the decrease of PM_{2.5} starting on 18 Aug from the CDPHE data (Figure 3). AOD increased in value and 220 spatial extent on 20 Aug during the second haze event, when more fires were detected in the Pacific Northwest (see 221 increase in number of MODIS hotspots in Figure 5). This plume contained a high density of aerosols that travelled 222 over the northcentral U.S. The southern periphery of this plume impacted Colorado east of the Continental Divide 223 starting on 20 Aug, as corroborated by the CDPHE air quality measurements in Figure 3. Although the AOD values 224 were not as enhanced over Colorado as compared to the core of the AOD plume, AOD values over the Front Range 225 were enhanced as compared to before the long-range transport of this plume. Enhanced AOD was observed around 226 Denver/Boulder and the Front Range the following three days (0.26-0.35), with the largest values in this four-day 227 period observed on 23 Aug. The third haze event (Figure 6) followed a similar evolution to the first two. The AOD 228 plume remained north of Colorado on 26–27 Aug, then infiltrated the northern and eastern part of the state on 28–29 229 Aug. The AOD values over Denver/Boulder during this event (0.26–0.45) were considerably larger than the two 230 previous events. It is important to note that AOD is a column measurement, thus the largest aerosol concentrations 231 may be elevated in the atmosphere as compared to what is observed on the ground. However, the AOD observations 232 still provide information regarding the spatial extent of the plume of aerosols emitted from the fires and that Colorado 233 was indeed impacted by air transported from the Pacific Northwest fires.

234

Further, the satellite retrievals generally corroborate the air quality observations on the ground along the Front Range in terms of when large concentrations of aerosols might be expected. More fires were detected across the Pacific Northwest by MODIS during the second event (678 fires, on average) when $PM_{2.5}$ was largest as compared to the first event (231 fires, on average), which had the smallest maximum $PM_{2.5}$ out of the three haze events. The third event had $PM_{2.5}$ values in between the first and second, while also having 607 fires on average. Thus, the number of fires likely influenced the relative amount of smoke produced and transported to the Front Range. However, meteorological

conditions as described below also played a vital role in enabling transport of the smoke.

242 **3.3** Synoptic and regional scale meteorology fuel long-range aerosol transport from the Pacific Northwest

243 The transport of the enhanced AOD plume from the Pacific Northwest to Colorado during each of the three events,

and the relationship between the AOD column and ground-based in situ observations, are supported by the

245 meteorological features present on both the synoptic and regional scales. Plan-view synoptic analyses aloft and at the

246 surface during the first air quality event along Colorado's Front Range on 17-18 Aug 2015 are shown in Figure 7. At 247 500 hPa (Figure 7a and b), a transient shortwave trough embedded in baroclinic zonal flow aloft migrates eastward 248 across the northern Rocky Mountains (i.e., north of Colorado), with westerly (north-westerly) flow preceding 249 (following) the passage of the trough axis. These flow patterns are corroborated by the HYSPLIT air mass back 250 trajectories during the first event, shown in Figure 7e. On average, air mass back trajectories passed over the fire 251 plume region 40% of the time, i.e., 19 of the 48 trajectories passed over regions of enhanced AOD and fire hotspot 252 locations from MODIS. At the surface, high pressure and shallow cool air initially resides primarily north of Colorado 253 at 0600 UTC 17 Aug (Figure 7c). However, by 2100 UTC 17 Aug (Figure 7d), the shallow cool air has moved 254 southward across eastern Colorado. A companion time-height section of hourly wind profiles at BAO (Figure 7f) 255 shows low-level southerly flow ahead of the frontal passage at ~1100 UTC 17 Aug and generally westerly to 256 northwesterly flow aloft for the duration of the plot. The observed flow aloft is represented in many of the back 257 trajectories, which show west to northwest flow reaching Boulder during this event. Following the frontal passage at 258 the wind profiler, the shallow cool air mass deepens to ~3 km MSL by 1800 UTC 17 Aug in generally northerly-259 component flow. Thereafter, the depth of the cool air decreases as the low-level flow shifts to south-easterly. Operational rawinsonde data from Denver (not shown) captures the top of the frontal inversion at 2.1 km MSL at 1200 260 261 UTC 17 Aug and at 2.7 km MSL at 0000 UTC 18 Aug, consistent with the wind-profiler analysis of the time-varying frontal altitude at BAO. For plan-view context, the times of the synoptic analyses are marked on the time-height 262 263 section. The high PM_{2.5} values (Figure 3) on 17 Aug are corroborated by the transition of air arriving from enhanced 264 AOD regions (see air mass backward trajectories in Figure 7e) over and off the coast of the Pacific Northwest and 265 northern California (Figure 4c). PM_{2.5} increased markedly after the passage of the shallow front, thus suggesting the 266 post-frontal air mass—which originated over Wyoming downstream of the Pacific Northwest fires—contains a large 267 concentration of particulates from those fires.

268

The evolution of the shallow cold front described above is typical of southward propagating cold fronts more generally across eastern Colorado, and the frontal propagation is influenced heavily by the complex regional topography depicted in Figure 1. Specifically, the blocking effect of the Rocky Mountains accelerates cold air southward along the eastern side of the high terrain (e.g., Colle and Mass, 1995; Neiman et al., 2001). Additionally, the postfrontal northerly-component airstream flowing across the west-east-oriented Cheyenne Ridge in south eastern Wyoming induces an anticyclonic gyre to the lee (south) of this ridge, subsequently shifting the postfrontal flow from northerly to easterly and driving the front westward against Colorado's Front Range (e.g., Davis, 1997; Neiman et al., 2001).

276

The meteorology during the second air quality event, on 22–23 Aug (Figure 8), is qualitatively similar to its predecessor, although the transient shortwave trough aloft is more amplified during the latter event (Figure 8a and b). Consequently, during the second event, the terrain-trapped cold front and its trailing shallow cool air mass east of the Rockies surges much farther southward across eastern New Mexico (Figure 8c and d). The corresponding air mass back trajectories (Figure 8e) traveled south-eastward from the Pacific Northwest fires to Colorado and passed over the fire plume region 96% of the time, leading to the worst event along the Front Range in terms of PM_{2.5} and total-

283 column extinction (Figure 3). The wind-profiler analysis at BAO (Figure 8f) shows an abrupt low-level wind shift from westerly to easterly with the frontal passage at 1900 UTC 22 Aug, followed by a rapid deepening of the shallow 284 285 cool air mass to nearly 3 km MSL. Thereafter, the depth of this air mass ranges between ~2.2 and 3.4 km MSL. Nearby 286 rawinsonde observations at Denver from 0000 UTC 23 Aug to 0000 UTC 24 Aug (not shown) document a strong 287 frontal inversion ranging between 3.3 and 3.8 km MSL, consistent with the wind-profiler analysis. Above the shallow 288 cool air mass, the profiler shows westerly flow aloft, shifting to north-westerly with the passage of the transient 289 shortwave trough. The largest PM_{2.5} values observed during this event, on 23 Aug, corresponds to the most direct 290 transport of air (Figure 8e) from over the enhanced AOD regions over the Pacific Northwest fires (Figure 5). As with 291 the previous case, the PM_{2.5} increased markedly with the passage of the shallow front (Figure 3). Significantly, air 292 quality is considerably poorer with the second event, perhaps due partly to a stronger cold-frontal push across 293 Colorado's Front Range that originated near the smoke source region and partly due to north-westerly (rather than 294 westerly) flow aloft that could transport the smoke through a deeper layer toward Colorado. Further, more fires were 295 detected during the second event (678, on average) compared to the first event (231 fires, on average), thus the larger 296 number of fires could result in more smoke production and thus a denser smoke plume transported to the Front Range.

297

298 The synoptic-scale conditions on 27–28 Aug (Figure 9) associated with the third air quality case differ considerably 299 from those of the two earlier events. Most significantly, a broad ridge aloft covers the intermountain West for the 300 duration of this final event, while an embedded weak shortwave trough migrates eastward through the ridge from 301 Wyoming/Colorado to the Great Plains (Figure 9a and b). A surface reflection of the upper-level shortwave trough is 302 manifest as a weak low-pressure centre over western Nebraska and Kansas at 1800 UTC 27 Aug (Figure 9c). This low 303 migrates eastward during the subsequent 24 h (Figure 9d) in tandem with the upper-level shortwave. Because this 304 surface low resides beneath a mean ridge aloft, the temperature contrast across this trailing cold front is weaker than 305 its earlier counterparts (not shown). Nevertheless, the southward migration of the front east of the Rockies suggests 306 that terrain blocking may have influenced its evolution. The air mass back trajectories show parcels originating from 307 the region of the fires and enhanced AOD 85% of the time, similar to the trajectories from the earlier two events 308 (Figure 9e). Companion observations from the BAO wind profiler (Figure 9f) capture the shallow frontal passage at 309 2000 UTC 27 Aug, when westerly flow shifts abruptly to northerly. Above 3 km MSL, the wind field exhibits a more 310 gradual transition from westerly to north-westerly as the weak shortwave trough moves across the wind profiler. The 311 Denver rawinsondes at 0000 and 1200 UTC 28 Aug observed a frontal inversion at ~2.1 km MSL (not shown). It is 312 less prominent than the frontal inversions during the earlier events, largely because the temperature contrast across 313 this front is weaker than its predecessors. The subsequent rawinsonde profile at 0000 UTC 29 Aug (not shown) 314 captures a deep, dry-convective boundary layer extending up to 4 km MSL, despite persistent low-level northerly 315 flow. Sensible heating eroded the remnant low-level cool air east of the Rockies. PM_{2.5} increases following the initial 316 shallow cold-frontal passage at 2000 UTC 27 Aug and continues to increase for the remainder of the wind-profiler 317 time-height section, as deep northerly-component flow behind the weak shortwave trough transports smoke 318 particulates across Colorado.

319 **3.4 Mineral dust and smoke arrive along the Front Range**

320 The types of aerosols present in the enhanced AOD plumes that were transported towards the Front Range via the aforementioned synoptic conditions were evaluated using additional satellite-based measurements and support the 321 322 interpretation of transport of aerosols from the wildfires in the Pacific Northwest to Colorado. Figure 10 shows aerosol 323 subtype data from the CALIPSO satellite in planar (a panel) and vertical-profile (b panel) views during Event 1. 324 CALIPSO data were strikingly similar for Events 2 and 3 and are provided in the Supporting Information. Only on 325 the day prior to, or on the worst day of, each haze event are shown, although aerosol subtype data were examined 326 anytime CALIPSO passed over the Pacific Northwest or Colorado from 15 Aug to 2 Sep. CALIPSO demonstrates the 327 presence of smoke, dust, or polluted dust (dust mixed with smoke in each profile) during times that intersect the 328 enhanced AOD plume propagating from the Pacific Northwest or when over Colorado. Dust and smoke plumes from 329 the fires extended up to 10 km MSL over the western U.S. The mineral dust and smoke detected by CALIPSO in 330 transit to the Front Range was also detected with the TOPAZ lidar and the in situ aerosol particle mass and speciation 331 monitor at the DSRC. Figure 11 shows aerosol extinction profiles from the surface to 2.5 km AGL measured with the 332 TOPAZ lidar on 9 days during the smoke episodes. The time resolution of the extinction profiles is 5 minutes and the 333 vertical resolution is 1 m at the lowest altitudes, increasing to 6 m above 500 m AGL. The observations on 14 Aug 334 and 2 Sep, which bracket the smoke episodes, indicate very clean conditions with AOD from the surface up to 2.5 km 335 AGL (AOD_{2.5km}) of 0.05 and 0.04, respectively. Aerosol extinction coefficients and AOD_{2.5km} were significantly larger during the smoke episodes with an approximately 7-fold increase in AOD_{2.5km} on 20 and 21 Aug. This time 336 337 period also corresponds to increasing extinction at DESCI (Figure 3). Aerosol extinction was enhanced over the entire 338 2.5 km column, but the largest aerosol extinction values were observed in the boundary layer in the lowest few hundred 339 meters up to 1.5 km AGL. Also, the lidar measurements reveal that on most days aerosol extinction varied significantly 340 over the course of the day (e.g., 20 Aug). The largest aerosol extinction values around 1-1.5 km AGL observed on 19 341 Aug were primarily due to swelling of aerosol particles in the moist relative humidity environment beneath cumulus 342 clouds at the top of the boundary layer. However, aerosol extinction in the lower part of the boundary was still 343 significantly larger than on 14 Aug, which is consistent with the larger aerosol particle concentrations in the smoke 344 plumes. The lidar measurements are consistent with the atmospheric extinction measurements from DESCI, and the 345 in situ PM_{2.5} and MODIS AOD observations. When comparing lidar AOD_{2.5km} with MODIS AOD one has to be cognizant of the fact that the TOPAZ observations only cover a portion of the atmospheric column and that the two 346 347 AOD measurements were made at different wavelengths. A comparison between the near-surface TOPAZ and DESCI 348 extinction observations also needs to take into account that the measurements were made at different wavelengths. 349

Figure 12 shows the time series of $PM_{2.5}$, soil mass concentrations, and elemental mass concentrations (data from the PX-375 was not available prior to this time period due to instrumental complications). Soil concentrations were calculated by following the Interagency Monitoring of Protected Visual Environments (IMPROVE) convention using concentrations of specific metals: SOIL = 2.2[AI] + 2.49[Si] + 1.63[Ca] + 2.42[Fe] + 1.94[Ti] (Malm et al., 1994; Hand et al., 2011). Both PM_{2.5} and soil mass concentrations increased during the worst haze event days (i.e., 26 and 29 Aug), when the Pacific Northwest fires were influencing air along the Front Range and when CALIPSO showed the presence of smoke and dust over the western U.S. The diurnal pattern is likely caused by the upslope/downslope

- 357 flow patterns due to proximity from the base of the foothills, which is particularly pronounced in the summer (Toth
- and Johnson, 1985). Further, select metals also increased in concentration during haze events, particularly those
- typically sourced from mineral dust (i.e., in the IMPROVE soil convention equation) and S and K, which are metal
- tracers that have been observed in smoke or biomass burning aerosols originating from fires (Artaxo et al., 1994;
- Gaudichet et al., 1995; Yamasoe et al., 2000; Pachon et al., 2013). It is important to note that K may also originate
- from soil. We calculated the soil K and non-soil K based on the methods of Kreidenweis et al. (2001), which is shown
- 363 in the Supporting Information. Concentrations of both soil K and non-soil K were highest during the influence from
- the fires. Additionally, IMPROVE measurements at the Rocky Mountain National Park location showed higher concentrations of soil, S, and K during event days in August, corroborating our measurements (see Supporting Information).
- 367

368 Figure 13 shows the average concentrations of mineral dust or biomass burning metal tracers from the PX-375 from 369 26 Aug to 2 Sep, during conditions influenced by the Pacific Northwest fires (days with enhanced PM_{2.5}; 29–30 Aug) and days with cleaner, normal Front Range conditions (days with low PM2.5; remaining days during this time period). 370 371 PM_{2.5} and soil mass, biomass burning metal (S and K), and mineral dust marker (Al, Si, Fe, and Ca) concentrations 372 were all larger, on average, during influences from the Pacific Northwest fires, corroborating the CALIPSO 373 observations. It is important to note the possibility that some small concentration of Ca, Al and Fe could also originate 374 from biomass burning, although the apportionment of this source remains in question and their contribution from 375 biomass burning aerosol are likely minor in comparison to their concentrations in mineral dust (Chang-Graham et al., 376 2011). Also included are metals that are typical of industrial tracers As and Pb (Figure 13e) (Paciga and Jervis, 1976; 377 Hutton and Symon, 1986; Thomaidis et al., 2003), which were actually lower in concentration during influences from 378 wildfires and enhanced during normal, regionally-sourced influences. The average PM_{2.5} mass concentration from the 379 CDPHE data was almost 3 times larger on 29–30 Aug as compared to the remaining days in the 26 Aug-2 Sep time period (15.9 versus 5.7 µg m⁻³, respectively). This result demonstrates how influences from typical, regional industrial 380 381 sources is disrupted by the synoptic conditions that introduced the long-range transported biomass burning plumes. Although Zn and Cu have been shown to originate from wildfires (Yamasoe et al., 2000), the averages were similar— 382 383 within 1 ng m⁻³—thus a distinct comparison could not be made within certainty. Further, these metals can also be 384 derived from vehicular emissions, thus their concentrations may additionally be influenced by local traffic (Sternbeck 385 et al., 2002). These results demonstrate the transport of mineral dust and biomass burning aerosol species to the Front Range, which were indeed larger in concentration during poor air quality/haze events. Interestingly, mineral dust 386 387 mixed within a smoke plume from fires has predominantly been observed originating from more arid regions along 388 the global dust belt, and using modelling or remote sensing data only (e.g., Radojevic, 2003; Tesche et al., 2009; Yang 389 et al., 2013; Nisantzi et al., 2014). To our knowledge, this co-lofting of dust and smoke has not been shown to occur 390 in the U.S., particularly in a region as densely covered in vegetation as the Pacific Northwest.

391 4 Conclusions

392 We have demonstrated the transport of mineral dust and smoke/biomass burning aerosols from wildfires in the Pacific 393 Northwest to the Colorado Front Range using a combination of in situ, remote sensing, and air parcel modelling 394 techniques (Severijnen, 2015). These aerosols were transported under synoptic conditions that contributed to three 395 different haze events, inducing poor air quality in the Denver metro area. Three separate poor air quality events with enhanced PM_{2.5} were likely dependent on the number of fires and observed to occur with cold frontal passages along 396 397 Colorado's Front Range, enabling the enhanced AOD plumes originating from the Pacific Northwest wildfires to 398 propagate south-eastward to Colorado's Front Range. Air masses were shown to originate from over the region dense 399 with wildfires, and followed through satellite-detected aerosol plumes, which were rich in a mixture of mineral dust 400 and smoke. Tracers for these aerosol types were also detected in situ along the Front Range, and were shown to be 401 enhanced during periods of influence from the fires.

402

403 Overall, these unique observations were demonstrated using a complete suite of in situ and remote sensing aerosol 404 measurements in the context of in situ meteorological observations and air mass trajectory modelling. In tandem, we 405 utilized a real-time X-ray fluorescence spectroscopy technique using the novel and field-portable PX-375 from 406 HORIBA, Ltd. demonstrating the utility of the instrument. Although the haze events were short lived, they 407 demonstrate how quickly (i.e., on the order of 2 to 3 days from the fire region to the Front Range) aerosols can be 408 transported long distances and affect air quality in regions thousands of kilometres away. Interestingly, mineral dust 409 was observed to be co-lofted and transported within the smoke plumes, an observation not previously reported for 410 vegetated regions such as the Pacific Northwest.

411

412 Mineral dust and smoke aerosols have disparate implications for health and climate, particularly at the levels observed 413 along the Front Range. These unique observations should be taken into account when developing health standards, 414 seeing as not only regional urban and industrial emissions contribute to poor air quality conditions. Additionally, dust 415 and smoke are efficient cloud forming nuclei—which impacts cloud lifetime, radiative effects, and precipitation 416 formation mechanisms—particularly when orographically lifted along barriers such as the Front Range into the upper 417 atmosphere, where cloud formation is prominent. Thus, transport of these aerosols from wildfires has broad 418 implications for altering aerosol composition in regions far from the source.

Author contribution. J. M. C. analysed XRF data, compiled CDPHE and MODIS data, ran HYSPLIT simulations,
 and wrote the manuscript. P. J. N. conducted meteorological analysis and interpretation. T. C. compiled and analysed

- 421 CALIPSO data. C. J. S., G. K., and R. A. analysed and supplied TOPAZ data. A. Y. provided PX-375 for usage. All
- 422 co-authors contributed to the writing of or provided comments for manuscript.
- 423 **Acknowledgements.** The authors would like to acknowledge the many agencies and organization from which data
- 424 were acquired, including the CDPHE for air quality data, NASA for MODIS and CALIPSO observations, NOAA for
- 425 HYSPLIT and HMT meteorological data, and the HDF group for providing example code to process CALIPSO data.

426 **References**

- Ackerman, S. A., Strabala, K. I., Menzel, W. P., Frey, R. A., Moeller, C. C., and Gumley, L. E.: Discriminating clear sky from
 clouds with MODIS, J Geophys Res-Atmos, 103, 32141-32157, 1998.
- Alvarez, R. J., Senff, C. J., Langford, A. O., Weickmann, A. M., Law, D. C., Machol, J. L., Merritt, D. A., Marchbanks, R. D.,
 Sandberg, S. P., Brewer, W. A., Hardesty, R. M., and Banta, R. M.: Development and Application of a Compact, Tunable,
 Solid-State Airborne Ozone Lidar System for Boundary Layer Profiling, J Atmos Ocean Tech, 28, 1258-1272, 2011.
- Alvarez, R. J., Senff, C. J., Weickmann, A. M., Sandberg, S. P., Langford, A. O., Marchbanks, R. D., Brewer, W. A., and Hardesty,
 R. M.: Reconfiguration of the NOAA TOPAZ lidar for ground-based measurement of ozone and aerosol backscatter,
 Proceedings of the 26th International Laser Radar Conference, Porto Heli, Greece, 2012,
- Ansmann, A., Baars, H., Tesche, M., Muller, D., Althausen, D., Engelmann, R., Pauliquevis, T., and Artaxo, P.: Dust and smoke
 transport from Africa to South America: Lidar profiling over Cape Verde and the Amazon rainforest, Geophys Res Lett,
 36, 2009.
- Artaxo, P., Gerab, F., Yamasoe, M. A., and Martins, J. V.: Fine Mode Aerosol Composition at 3 Long-Term Atmospheric
 Monitoring Sites in the Amazon Basin, J Geophys Res-Atmos, 99, 22857-22868, 1994.
- Baars, H., Ansmann, A., Althausen, D., Engelmann, R., Artaxo, P., Pauliquevis, T., and Souza, R.: Further evidence for significant smoke transport from Africa to Amazonia, Geophys Res Lett, 38, 2011.
- Benjamin, S. G., Weygandt, S. S., Brown, J. M., Hu, M., Alexander, C., Smirnova, T. G., Olson, J. B., James, E., Dowell, D. C.,
 Grell, G. A., Lin, H., Peckham, S. E., Smith, T. L., and Moninger, W. R.: A North American hourly assimilation and
 model forecast cycle: The rapid refresh, Mon. Wea. Rev., 144, in review, 2016.
- 445 Bowman, D. J. S., and Johnston, F.: Wildfire Smoke, Fire Management, and Human Health, EcoHealth, 2, 76-80, 2005.
- Bravo, A. H., Sosa, E. R., Sanchez, A. P., Jaimes, P. M., and Saavedra, R. M. I.: Impact of wildfires on the air quality of Mexico City, 1992-1999, Environ Pollut, 117, 243-253, 2002.
- Brown, S. S., Thornton, J. A., Keene, W. C., Pszenny, A. A. P., Sive, B. C., Dube, W. P., Wagner, N. L., Young, C. J., Riedel, T.
 P., Roberts, J. M., VandenBoer, T. C., Bahreini, R., Ozturk, F., Middlebrook, A. M., Kim, S., Hubler, G., and Wolfe, D.
 E.: Nitrogen, Aerosol Composition, and Halogens on a Tall Tower (NACHTT): Overview of a wintertime air chemistry
 field study in the front range urban corridor of Colorado, J Geophys Res-Atmos, 118, 8067-8085, 2013.
- Chalbot, M. C., Nikolich, G., Etyemezian, V., Dubois, D. W., King, J., Shafer, D., da Costa, G. G., Hinton, J. F., and Kavouras, I.
 G.: Soil humic-like organic compounds in prescribed fire emissions using nuclear magnetic resonance spectroscopy,
 Environ Pollut, 181, 167-171, 2013.
- Chang-Graham, A. L., Profeta, L. T. M., Johnson, T. J., Yokelson, R. J., Laskin, A., and Laskin, J.: Case Study of Water-Soluble
 Metal Containing Organic Constituents of Biomass Burning Aerosol, Environ Sci Technol, 45, 1257-1263, 2011.
- Clements, C. B., Zhong, S. Y., Bian, X. D., Heilman, W. E., and Byun, D. W.: First observations of turbulence generated by grass
 fires, J Geophys Res-Atmos, 113, 2008.
- Colarco, P. R., Schoeberl, M. R., Doddridge, B. G., Marufu, L. T., Torres, O., and Welton, E. J.: Transport of smoke from Canadian forest fires to the surface near Washington, D. C.: Injection height, entrainment, and optical properties, J Geophys Res-Atmos, 109, 2004.
- 462 Colle, B. A., and Mass, C. F.: The Structure and Evolution of Cold Surges East of the Rocky-Mountains, Mon Weather Rev, 123, 463 2577-2610, 1995.
- 464 Cruz, C. N., and Pandis, S. N.: A study of the ability of pure secondary organic aerosol to act as cloud condensation nuclei, Atmos
 465 Environ, 31, 2205-2214, 1997.
- 466Davis, C. A.: Mesoscale anticyclonic circulations in the Lee of the central Rocky Mountains, Mon Weather Rev, 125, 2838-2855,4671997.
- DeMott, P. J., Chen, Y., Kreidenweis, S. M., Rogers, D. C., and Sherman, D. E.: Ice formation by black carbon particles, Geophys
 Res Lett, 26, 2429-2432, 1999.
- DeMott, P. J., Sassen, K., Poellot, M. R., Baumgardner, D., Rogers, D. C., Brooks, S. D., Prenni, A. J., and Kreidenweis, S. M.:
 African dust aerosols as atmospheric ice nuclei, Geophys Res Lett, 30, 2003.
- 472 Draxler, R. R., and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA
 473 ARL READY Website (<u>http://ready.arl.noaa.gov/HYSPLIT.php</u>), 2011.

- Eagan, R. C., Hobbs, P. V., and Radke, L. F.: Measurements of Cloud Condensation Nuclei and Cloud Droplet Size Distributions
 in Vicinity of Forest Fires, J Appl Meteorol, 13, 553-557, 1974.
- Gadi, R., Kulshrestha, U. C., Sarkar, A. K., Garg, S. C., and Parashar, D. C.: Emissions of SO2 and NOx from biofuels in India, Tellus B, 55, 787-795, 2003.
- Gaudichet, A., Echalar, F., Chatenet, B., Quisefit, J. P., Malingre, G., Cachier, H., Buatmenard, P., Artaxo, P., and Maenhaut, W.:
 Trace-Elements in Tropical African Savanna Biomass Burning Aerosols, J Atmos Chem, 22, 19-39, 1995.
- Ge, C., Wang, J., and Reid, J. S.: Mesoscale modeling of smoke transport over the Southeast Asian Maritime Continent: coupling
 of smoke direct radiative effect below and above the low-level clouds, Atmos Chem Phys, 14, 159-174, 2014.
- 482 Giglio, L., Descloitres, J., Justice, C. O., and Kaufman, Y. J.: An enhanced contextual fire detection algorithm for MODIS, Remote
 483 Sens Environ, 87, 273-282, 2003.
- 484 Giglio, L.: MODIS Collection 5 Active Fire Product User's Guide Version 2.4, Science Systems and Applications, Inc., 2010.
- 485 Haagenson, P. L.: Meteorological and Climatological Factors Affecting Denver Air-Quality, Atmos Environ, 13, 79-85, 1979.
- Hand, J. L., Copland, S. A., Dillner, A. M., Indresand, H., Malm, W. C., McDade, C. E., Moore, C. T., Pitchford, M. L., Schichtel,
 B. A., and Watson, J. G.: Spatial and Seasonal Patterns and Temporal Variability of Haze and its Constituents in the
 United States Report V Cooperative Institute for Research in the Atmosphere, 2011.
- Hutton, M., and Symon, C.: The Quantities of Cadmium, Lead, Mercury and Arsenic Entering the Uk Environment from Human Activities, Sci Total Environ, 57, 129-150, 1986.
- Jacob, D. J., and Winner, D. A.: Effect of climate change on air quality, Atmos Environ, 43, 51-63, 2009.
- Jaffe, D. A., and Wigder, N. L.: Ozone production from wildfires: A critical review, Atmos Environ, 51, 1-10, 2012.
- Jayachandran, S.: Air quality and early-life mortality: Evidence from Indonesia's wildfires, J. Human Resources, 44, 916-954,
 2008.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y.,
 Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds,
 R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, B Am Meteorol Soc, 77, 437-471, 1996.
- Kaufman, Y. J., Tanre, D., Remer, L. A., Vermote, E. F., Chu, A., and Holben, B. N.: Operational remote sensing of tropospheric
 aerosol over land from EOS moderate resolution imaging spectroradiometer, J Geophys Res-Atmos, 102, 17051-17067,
 1997.
- Kreidenweis, S. M., Remer, L. A., Bruintjes, R., and Dubovik, O.: Smoke aerosol from biomass burning in Mexico: Hygroscopic
 smoke optical model, J Geophys Res-Atmos, 106, 4831-4844, 2001.
- Liu, Y. Q., Stanturf, J., and Goodrick, S.: Trends in global wildfire potential in a changing climate, Forest Ecol Manag, 259, 685-697, 2010.
- Malm, W. C., Sisler, J. F., Huffman, D., Eldred, R. A., and Cahill, T. A.: Spatial and Seasonal Trends in Particle Concentration
 and Optical Extinction in the United-States, J Geophys Res-Atmos, 99, 1347-1370, 1994.
- McCluskey, C. S., DeMott, P. J., Prenni, A. J., Levin, E. J. T., McMeeking, G. R., Sullivan, A. P., Hill, T. C. J., Nakao, S., Carrico,
 C. M., and Kreidenweis, S. M.: Characteristics of atmospheric ice nucleating particles associated with biomass burning
 in the US: Prescribed burns and wildfires, J Geophys Res-Atmos, 119, 2014.
- Moore, D., Copes, R., Fisk, R., Joy, R., Chan, K., and Brauer, M.: Population Health Effects of Air Quality Changes Due to Forest
 Fires in British Columbia in 2003: Estimates from Physician-visit Billing Data, Canadian Journal of Public Health /
 Revue Canadienne de Sante'e Publique, 97, 105-108, 2006.
- Mott, J. A., Meyer, P., Mannino, D., Redd, S. C., Smith, E. M., Gotway-Crawford, C., and Chase, E.: Wildland forest fire smoke:
 health effects and intervention evaluation, Hoopa, California, 1999, Western Journal of Medicine, 176, 157-162, 2002.
- Neff, J. C., Ballantyne, A. P., Farmer, G. L., Mahowald, N. M., Conroy, J. L., Landry, C. C., Overpeck, J. T., Painter, T. H.,
 Lawrence, C. R., and Reynolds, R. L.: Increasing eolian dust deposition in the western United States linked to human activity, Nat Geosci, 1, 189-195, 2008.
- Neiman, P. J., Ralph, F. M., Weber, R. L., Uttal, T., Nance, L. B., and Levinson, D. H.: Observations of nonclassical frontal propagation and frontally forced gravity waves adjacent to steep topography, Mon Weather Rev, 129, 2633-2659, 2001.
- Nisantzi, A., Mamouri, R. E., Ansmann, A., and Hadjimitsis, D.: Injection of mineral dust into the free troposphere during fire
 events observed with polarization lidar at Limassol, Cyprus, Atmos Chem Phys, 14, 12155-12165, 2014.
- 522 Nriagu, J. O.: A Global Assessment of Natural Sources of Atmospheric Trace-Metals, Nature, 338, 47-49, 1989.
- Omar, A. H., Winker, D. M., Kittaka, C., Vaughan, M. A., Liu, Z. Y., Hu, Y. X., Trepte, C. R., Rogers, R. R., Ferrare, R. A., Lee,
 K. P., Kuehn, R. E., and Hostetler, C. A.: The CALIPSO Automated Aerosol Classification and Lidar Ratio Selection
 Algorithm, J Atmos Ocean Tech, 26, 1994-2014, 2009.
- Pachon, J. E., Weber, R. J., Zhang, X. L., Mulholland, J. A., and Russell, A. G.: Revising the use of potassium (K) in the source apportionment of PM2.5, Atmos Pollut Res, 4, 14-21, 2013.
- 528 Paciga, J. J., and Jervis, R. E.: Multielement Size Characterization of Urban Aerosols, Environ Sci Technol, 10, 1124-1128, 1976.
- Park, R. J., Jacob, D. J., Chin, M., and Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, J Geophys Res-Atmos, 108, 2003.
- Peterson, D., Hyer, E., and Wang, J.: Quantifying the potential for high-altitude smoke injection in the North American boreal
 forest using the standard MODIS fire products and subpixel-based methods, J Geophys Res-Atmos, 119, 3401-3419,
 2014.
- Phuleria, H. C., Fine, P. M., Zhu, Y. F., and Sioutas, C.: Air quality impacts of the October 2003 Southern California wildfires, J
 Geophys Res-Atmos, 110, 2005.

- Pinker, R. T., Liu, H., Osborne, S. R., and Akoshile, C.: Radiative effects of aerosols in sub-Sahel Africa: Dust and biomass burning,
 J Geophys Res-Atmos, 115, 2010.
- Pio, C. A., Legrand, M., Alves, C. A., Oliveira, T., Afonso, J., Caseiro, A., Puxbaum, H., Sanchez-Ochoa, A., and Gelencser, A.:
 Chemical composition of atmospheric aerosols during the 2003 summer intense forest fire period, Atmos Environ, 42, 7530-7543, 2008.
- Radojevic, M.: Chemistry of forest fires and regional haze with emphasis on Southeast Asia, Pure Appl Geophys, 160, 157-187,
 2003.
- 543 Severijnen, M.: Detecting source of unexpected peak of PM10 over Flanders (Belgium): a walk through public available 544 information, personal communication, 2015.
- 545 Sibold, J. S., and Veblen, T. T.: Relationships of subalpine forest fires in the Colorado Front Range with interannual and 546 multidecadal-scale climatic variation, J Biogeogr, 33, 833-842, 2006.
- Spracklen, D. V., Logan, J. A., Mickley, L. J., Park, R. J., Yevich, R., Westerling, A. L., and Jaffe, D. A.: Wildfires drive interannual
 variability of organic carbon aerosol in the western US in summer, Geophys Res Lett, 34, 2007.
- 549 Sternbeck, J., Sjodin, A., and Andreasson, K.: Metal emissions from road traffic and the influence of resuspension results from 550 two tunnel studies, Atmos Environ, 36, 4735-4744, 2002.
- 551 Streets, D. G., Yarber, K. F., Woo, J. H., and Carmichael, G. R.: Biomass burning in Asia: Annual and seasonal estimates and 552 atmospheric emissions, Global Biogeochem Cy, 17, 2003.
- Tesche, M., Ansmann, A., Muller, D., Althausen, D., Engelmann, R., Freudenthaler, V., and Gross, S.: Vertically resolved
 separation of dust and smoke over Cape Verde using multiwavelength Raman and polarization lidars during Saharan
 Mineral Dust Experiment 2008, J Geophys Res-Atmos, 114, 2009.
- Thomaidis, N. S., Bakeas, E. B., and Siskos, P. A.: Characterization of lead, cadmium, arsenic and nickel in PM2.5 particles in the
 Athens atmosphere, Greece, Chemosphere, 52, 959-966, 2003.
- Toth, J. J., and Johnson, R. H.: Summer Surface Flow Characteristics over Northeast Colorado, Mon Weather Rev, 113, 1458 1469, 1985.
- Vali, G., DeMott, P. J., Mohler, O., and Whale, T. F.: Technical Note: A proposal for ice nucleation terminology, Atmos Chem
 Phys, 15, 10263-10270, 2015.
- VanCuren, R. A.: Asian aerosols in North America: Extracting the chemical composition and mass concentration of the Asian continental aerosol plume from long-term aerosol records in the western United States, J Geophys Res-Atmos, 108, 2003.
- Vaughan, M., Young, S., Winker, D., Powell, K., Omar, A., Liu, Z. Y., Hu, Y. X., and Hostetler, C.: Fully automated analysis of
 space-based lidar data: an overview of the CALIPSO retrieval algorithms and data products, Bba Lib, 5575, 16-30, 2004.
- Watson, J. G., Fujita, E., Chow, J. C., Zielinska, B., Richards, L. W., Neff, W., and Dietrich, D.: Northern Front Range Air Quality
 Study Final Report, 1998.
- Weber, B. L., Wuertz, D. B., and Welsh, D. C.: Quality Controls for Profiler Measurements of Winds and Rass Temperatures, J
 Atmos Ocean Tech, 10, 452-464, 1993.
- Westerling, A. L., Hidalgo, H. G., Cayan, D. R., and Swetnam, T. W.: Warming and earlier spring increase western US forest
 wildfire activity, Science, 313, 940-943, 2006.
- White, A. B., Anderson, M. L., Dettinger, M. D., Ralph, F. M., Hinojosa, A., Cayan, D. R., Hartman, R. K., Reynolds, D. W.,
 Johnson, L. E., Schneider, T. L., Cifelli, R., Toth, Z., Gutman, S. I., King, C. W., Gehrke, F., Johnston, P. E., Walls, C.,
 Mann, D., Gottas, D. J., and Coleman, T.: A Twenty-First-Century California Observing Network for Monitoring
 Extreme Weather Events, J Atmos Ocean Tech, 30, 1585-1603, 2013.
- Wiedinmyer, C., Quayle, B., Geron, C., Belote, A., McKenzie, D., Zhang, X. Y., O'Neill, S., and Wynne, K. K.: Estimating
 emissions from fires in North America for air quality modeling, Atmos Environ, 40, 3419-3432, 2006.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y. X., Powell, K. A., Liu, Z. Y., Hunt, W. H., and Young, S. A.: Overview of the
 CALIPSO Mission and CALIOP Data Processing Algorithms, J Atmos Ocean Tech, 26, 2310-2323, 2009.
- Yamasoe, M. A., Artaxo, P., Miguel, A. H., and Allen, A. G.: Chemical composition of aerosol particles from direct emissions of
 vegetation fires in the Amazon Basin: water-soluble species and trace elements, Atmos Environ, 34, 1641-1653, 2000.
- Yang, Z. F., Wang, J., Ichoku, C., Hyer, E., and Zeng, J.: Mesoscale modeling and satellite observation of transport and mixing of
 smoke and dust particles over northern sub-Saharan African region, J Geophys Res-Atmos, 118, 12139-12157, 2013.
- Zamora, R. J., Dutton, E. G., Trainer, M., McKeen, S. A., Wilczak, J. M., and Hou, Y. T.: The accuracy of solar irradiance
 calculations used in mesoscale numerical weather prediction, Mon Weather Rev, 133, 783-792, 2005.



588 Figure 1. Map of monitoring locations, including NOAA DSRC in Boulder, which housed the PX-375 and

589 TOPAZ lidar instruments, the BAO where the 449-MHz wind profiler was deployed, downtown Denver, the 590 CDPHE DESCI site where atmospheric extinction/visibility is measured, and the CDPHE sites where PM_{2.5}

591 and PM₁₀ are monitored (see Table 1 for site descriptions). The approximate area encompassing the Colorado

592 Front Range is highlighted by the dashed line. The Cheyenne Ridge in Wyoming is also notated.



593

594 Figure 2. Images of downtown Denver facing west taken at 1400 UTC (0800 MDT). Images acquired from the CDPHE

595 Visibility Station (DESCI; 39.73°N, 104.96°W; 1633 m MSL). Only days of significant meteorological and visibility

596 transitions in August 2015 are shown. Days in red are those which correspond to the haziest days during the study time

597 period. In panels (a), (d), and (f), the visibility of the foothills (and background high terrain) is highlighted.



599

Figure 3. Top panel shows atmospheric extinction measured at the CDPHE DESCI site (see Figure 1). Bottom
 panel shows hourly and daily averaged PM_{2.5} mass concentrations at CDPHE sites. The pairs of red dashed

602 lines shows the times before "B" and after "A" cold-frontal passages at BAO during or prior to each haze event.

603 The daily averaged PM_{2.5} in red represent the haziest days during or following cold front al passages (i.e.,

604 Events 1, 2, and 3 on 17, 23, and 29 Aug 2015, respectively).



605

606 Figure 4. Daily averaged aerosol optical depth (AOD; colour bar lower right) at 550 nm and fire hotspots (black

- 607 markers) detected by MODIS during the first major haze case study between 15 and 18 Aug 2015. The haziest
- 608 day from the CDPHE data is labelled in red (i.e., Event 1).



- 610 Figure 5. Same as Figure 4, but for the second major haze event between 20 and 23 Aug 2015. The haziest day
- 611 from the CDPHE data is labelled in red (i.e., Event 2).



- 613 Figure 6. Same as Figure 4, but for the third major haze event between 26 and 29 Aug 2015. The haziest day
- 614 from the CDPHE data is labelled in red (i.e., Event 3).



Figure 7. Meteorological analysis for Event 1 (17 Aug 2015). Top row shows 13-km resolution RAP gridded 616 617 dataset of 500-hPa geopotential heights (black contours) with 500-hPa wind velocities (flags = 25 m s^{-1} , barbs = 5 m s^{-1} , half-barbs = 2.5 m s⁻¹) from before (a) and after (b) the passage of a cold front at 0600 and 2100 UTC, 618 619 respectively. Middle row shows mean sea-level pressure (black contours) with near-surface wind velocities 620 (flags and barbs as above) from before (c) and after (d) the cold-front passage. Standard frontal notation is used. (e) 10-day air mass backward trajectories initiated every 6 hours at 500, 1000, and 2000 m MSL during 621 622 the time period surrounding Event 1 (15-18 Aug). Trajectories in red correspond to the haziest day (17 Aug) 623 and the blue dashed trajectories show the remaining. (f) Time-height section of hourly-averaged wind profiles from the 449-MHz wind profiler at BAO between 0600 UTC 17 Aug and 0600 UTC 18 Aug (flags and barbs 624 625 are as above). The bold black line denotes the approximate frontal shear boundary. The pair of red dashed lines shows the RAP analysis times before "B" and after "A" the cold-frontal passage at BAO. Time increases 626 627 from right to left to portray the advection of upper-level synoptic features from west to east.



628

629 Figure 8. Same as Figure 7, but for Event 2 (23 Aug 2015). Before and after the cold-frontal passage correspond

630 to 1800 UTC 22 Aug and 1200 UTC 23 Aug, respectively. Trajectories were initiated for the time period

631 surrounding Event 2 (20–23 Aug). Time-height section measurements were between 1700 UTC 22 Aug and

632 0100 UTC 24 Aug.



Figure 9. Same as Figure 7, but for Event 3 (29 Aug 2015). Before and after the cold-frontal passage correspond
to 1800 UTC 27 Aug and 1800 UTC 28 Aug, respectively. Trajectories were initiated for the time period
surrounding Event 3 (26–29 Aug). Time-height section measurements were between 1300 UTC 27 Aug and

2100 UTC 28 Aug.



Figure 10. CALIPSO swath data from the night prior to Event 1. Swath data contained in CAL_LID_L2_VFM_ValState1-V3-30 file are from 16 Aug 2015 09:57:00 UTC. (a) Map showing CALIPSO coverage, with the purple markers representing locations in the column measurement where dust, smoke, or polluted dust were observed. (b) Vertical profile (in km MSL) for all aerosol subtypes of the swath corresponding to (a).



644

Figure 11. Aerosol extinction profiles at 294 nm observed with the TOPAZ lidar on 9 days during the smoke pollution episodes. The numbers next to each day's observations represent the daily mean AOD from the surface up to 2.5 km AGL computed from the lidar measurements.



649

Figure 12. (a) Time series of hourly PM_{2.5} and soil mass concentrations as measured by PX-375 between 27 Aug and 2 Sep 2015 and (b) hourly mass concentrations of select individual metals relative to their maximum concentration observed during the study time period, including an error of $\pm 2\%$. Only data higher than the LDLs are shown. PX-375 data overlapped with Event 3.





Figure 13. Averages of (a) PM_{2.5} and soil concentrations and (b) – (e) select metal mass concentrations during non-event days (i.e., cleaner conditions) compared to averages from haze event days (i.e., influence from fires haze) for 26 Aug–2 Sep 2015. "Low" and "high" correspond to the PM_{2.5} concentration values. Error bars represent the 90% confidence intervals. Concentration averages were statistically significant based on t-tests of two samples of unequal variances.

Table 1. CDPHE sites used for particulate data within the Colorado Front Range. Each site has an 'x' for each measurement it maintained throughout the current work. Elevation is provided in meters above mean sea level (m MSL).

City/Site Name	Site ID	Latitude (degrees N)	Longitude (degrees W)	Elevation (m MSL)	PM 2.5	PM 10
Boulder - CU/Athens	BOU	40.01	105.27	1,621	Х	
Chatfield Park	CHAT	39.53	105.05	1,685	Х	
Colorado College	CCOL	38.85	104.83	1,833	Х	
Commerce City/Alsup Elementary	COMM	39.83	104.94	1,565	Х	
Denver - Continuous Air Monitoring site	CAMP	39.68	104.99	1,610	Х	Х
Denver - National Jewish Health	NJH	39.74	104.94	1,615	Х	
Fort Collins - CSU Facilities	FTCF	40.57	105.08	1,525	Х	Х
Greeley - Hospital	GREH	40.42	104.71	1,439	Х	
I-25 - Denver	I-25	39.73	105.02	1,586	Х	Х
La Casa	CASA	39.78	105.01	1,601	Х	Х
Longmont - Municipal	LNGM	40.16	105.10	1,517	Х	
Welby	WBY	39.84	104.95	1,554		х

 $PM_{2.5}$ = particulate matter with diameters $\leq 2.5 \ \mu m$

5 PM_{10} = particulate matter with diameters $\leq 10 \ \mu m$

Table 2. Lower detection limits (LDLs, ng m⁻³) for metals measured by the PX-375 during 15 Aug-2 Sep 2015. Concentrations less than the LDLs were excluded from analysis.

Species	LDL		
Ti	2.29		
V	0.23		
Cr	0.61		
Mn	0.93		
Fe	1.51		
Ni	0.33		
Cu	0.78		
Zn	1.21		
As	0.02		
Pb	0.80		
Al	32.2		
Si	5.17		
S	1.11		
K	4.37		
Ca	1.18		