

July 25, 2016

Dear Prof. Heard,

Enclosed please find our revised manuscript acp-2016-292 entitled "*Observation of atmospheric peroxides during Wangdu Campaign 2014 at a rural site in the North China Plain*". We have completed the revisions according to your comments. We have uploaded all of the materials to the ACP website.

Many thanks for your comments and time.

Sincerely yours,

Zhongming Chen and co-authors

Comments to the Author:

Thank you for the responses to the comments of reviewer (1).

There are a few things I would like you to expand on a little. Mainly it is when you give a response, but the MS does not seem to have been modified to reflect the response. See below.

The reviewer asks:

In the model description it seems to suggest that the uncertainty in the model is just due to the uncertainty in the observational constraints (linee184-5). What about uncertainties in chemical mechanism - rate coefficients, reaction products, reaction simplifications?

Your reply to this:

"The uncertainties in chemical mechanism was calculated to be less than 20% according to the method described in Lu et al. (2012)."

This is somewhat vague. Can you please indicate which parts of the chemical mechanism are likely to have the largest uncertainties and give a little more detail on those areas of the mechanism which contribute most to the overall 20% that Lu et al. report.

Reply: Yes. The largest uncertainties in chemical mechanism are possibly from the rate constants of peroxy radical self/cross reaction related to the secondary formation of H₂O₂, MHP and PAA. The uncertainties in the rate constants of the reaction between HO₂ radicals, the reaction between CH₃O₂ radicals and HO₂ radicals and the reaction between CH₃C(O)O₂ radicals and HO₂ radicals are calculated to be ~15%

with an accuracy of 30% on the rate constant in MCMv3.3.1. The detail on the areas of mechanism that contribute most to the overall uncertainty in Lu et al. (2012) is not available as we just employed the method described in Lu et al. (2012) to estimate our model uncertainty.

Can you please indicate the fraction of peroxides which are lost by dry deposition. You state it is not major, whereas the referee states that dry deposition is major - can you please provide some evidence that dry deposition is minor.

Reply: Yes. Dry deposition is the marginal removal pathway that contributed less than 20% to the destruction of H₂O₂, MHP and PAA, as shown in Fig. 6 in the MS. The simulated mean lifetime of H₂O₂, MHP and PAA between 18:00 and 24:00 LT in the Phase I can be about 14 h, 16 h and 21 h, respectively, supposing that the loss of H₂O₂, MHP and PAA is merely due to photolysis, OH-initiated reaction and dry deposition, which is strongly overestimated compared with the mean lifetime of H₂O₂, MHP and PAA obtained from the field observation between 18:00 and 24:00 LT in the Phase I (<6 h). Heterogeneous uptake by aerosols was the predominant sink for atmospheric peroxides during Wangdu Campaign 2014.

Referee said:

Lines 211-13: If MHP and PAA being about 20% and 5% of the total peroxides in Wangdu is similar to the results from other rural sites in China, then presumably it could also be stated that with 70% of the peroxides in Wangdu being H₂O₂ that Wangdu is also similar to other rural sites in China?

Your reply is "yes" but can you make sure that this is stated in the MS - you did not say you had done this.

Reply: Yes. We have stated this in the MS as "H₂O₂ accounted for ~70% of total detected peroxides (H₂O₂ + MHP + PAA) similar to those determined at other rural sites in China".

Please state in the paper the criteria for when there is a "haze day". You state this nicely in the reply but not clear if stated in the MS - please ensure that it is.

It is very important that a clear definition of a haze day is contained in the MS.

Reply: Yes. We have stated this in the MS as "the haze day is defined as a day with daily-averaged PM_{2.5} concentration over 75 μg m⁻³".

The referee states:

Lines 299-305: The text here implies that there were modifications made to the chemical mechanism, with the new reactions given in Table 2. However, it is not clear what they have changed from. New reactions or just new rates?

In your reply you say the new reaction between CH_3O_2 and HO_2 was incorporated into your model. Please state clearly in the text exactly what you have done - state the old rate constants and what you have changed them to.

Reply: Sorry. This is a mistake in our response to reviewer (1). For CH_3O_2 related chemistry, the new reaction between CH_3O_2 radicals and OH radicals was incorporated in our box model. Please see the former revised manuscript.

The referee says:

When discussing Fig. 3. I would like to see more comment on the shape of the diurnal cycle. The modelled and measured shapes are often very different.

You reply to this, but you do not seem to say you have changed the MS for this. Please put more comment on the shape of the diurnal cycle in the MS as the referee asks.

Reply: Yes. We have stated this in Sect. 3.2 in the MS as "the calculated values in the model base case showed a general tendency to strongly overestimate the observed values (Fig. 3). The modelled and measured shapes of the diurnal cycle of atmospheric peroxides were different", "with the inclusion of heterogeneous reactions on aerosol particles, the simulated concentrations of atmospheric peroxides were apparently improved and the modelled shape of the diurnal cycle of H_2O_2 was closer to the measured shape" and in Sect. 3.4 in the MS as "The modelled and measured shape of the diurnal cycle of H_2O_2 in the Phase I and the Phase III are similar". Please see the revised manuscript.

The referee says:

What is the impact of including primary sources of the peroxides on the chemistry? E.g. does it have an impact on HO_x ? Are the modelled HO_x values comparable to observed values and does the addition of primary sources change this comparison?

You say the impact on HO_x is marginal. Can you please give a precise % value for this and put this in the MS.

Reply: Yes. We have stated this in the MS as "The impact of primary sources of the peroxides on HO_x radicals was limited with the increase of OH radicals not more than 10% and the increase of HO_2 radicals not more than 5%".

Referee question and response:

Lines 483-498: Given that OH concentrations and photolysis rates will be reduced in the evenings, the lifetimes calculated will be lower than daytime values. Are the modelled values 24 hour averages? It needs to be clear exactly what you are comparing and if they are for the same times of day. How sensitive are the modelled relative losses (Fig. 6) to uncertainties in the boundary layer height?

Reply: Yes. The modelled values were 24 hour averages. The comparison was done for the same times of day. The sensitivity of modelled relative loss of dry deposition to the uncertainties in the planetary boundary layer height was low, for instance, the contribution of dry deposition to the loss of H₂O₂ in Phase I decreased from 17% to 10% with the PBL height doubled. Heterogeneous uptake by aerosols was still the predominant sink for atmospheric peroxides.

Thanks for detailed response, but please make sure this is all contained in the revised MS.

Reply: Yes. We have stated this in the MS as "Although dry deposition is thought to dominate the atmospheric lifetime of peroxides in previous studies (Reeves and Penkett, 2003), its role in the lifetime of atmospheric peroxides is insignificant during Wangdu Campaign 2014. The sensitivity of modelled relative loss of dry deposition to the uncertainties in the planetary boundary layer height was low as the contribution of dry deposition to the loss of H₂O₂ in Phase I decreased no more than 10% with the PBL height doubled". We have also clarified the statement further by modifying "the lifetime" to "the daily-averaged lifetime". Please see the revised manuscript.

Fig 2, 3 replace "the grey shade" with "the grey shaded area...". Same for "orange shade"

Reply: Yes. We have changed "shade" to "shaded area".

References

Reeves, C. E., and Penkett, S. A.: Measurements of peroxides and what they tell us, Chem. Rev., 103, 5199–5218, 2003.

Observation of atmospheric peroxides during Wangdu Campaign 2014 at a rural site in the North China Plain

**Yin Wang, Zhongming Chen, Qinqin Wu, Hao Liang, Liubin Huang, Huan Li,
Keding Lu, Yusheng Wu, Huabin Dong, Limin Zeng, and Yuanhang Zhang**

State Key Laboratory of Environmental Simulation and Pollution Control, College of
Environmental Sciences and Engineering, Peking University, Beijing 100871, China

Correspondence to: Zhongming Chen (zmchen@pku.edu.cn)

Abstract

Measurements of atmospheric peroxides were made during Wangdu Campaign 2014 at Wangdu, a rural site in the North China Plain (NCP) in summer 2014. The predominant peroxides were detected to be hydrogen peroxide (H₂O₂), methyl hydroperoxide (MHP) and peroxyacetic acid (PAA). The observed H₂O₂ reached up to 11.3 ppbv, which was the highest value compared with previous observations in China at summer time. A box model simulation based on the Master Chemical Mechanism and constrained by the simultaneous observations of physical parameters and chemical species was performed to explore the chemical budget of atmospheric peroxides. Photochemical oxidation of alkenes was found to be the major secondary formation pathway of atmospheric peroxides, while contributions from alkanes and aromatics were of minor importance. The comparison of modelled and measured peroxide concentrations revealed an underestimation during biomass burning events and an overestimation on haze days, which were ascribed to the direct production of peroxides from biomass burning and the heterogeneous uptake of peroxides by aerosols, respectively. The strengths of the primary emissions from biomass burning were on the same order of the known secondary production rates of atmospheric peroxides during the biomass burning events. The heterogeneous process on aerosol particles was suggested to be the predominant sink for atmospheric peroxides. The atmospheric lifetime of peroxides on haze days in summer in the NCP was about 2–3 hours, which is in good agreement with the laboratory studies. Further comprehensive


investigations are necessary to better understand the impact of biomass burning and heterogeneous uptake on the concentration of peroxides in the atmosphere.

1 Introduction

Atmospheric peroxides, including hydrogen peroxide (H_2O_2) and organic peroxides (ROOH), are vital oxidants present in the gaseous, aqueous and particulate phase in the atmospheric chemical processes. They serve as temporary reservoirs for HO_x radicals, contributing to the atmospheric oxidation capacity (Reeves and Penkett, 2003). Peroxides also participate in the conversion of S(IV) to S(VI) in the aqueous phase, leading to the acid precipitation and the formation of secondary sulfate (SO_4^{2-}) aerosols in the troposphere (Calvert et al., 1985; Stein and Saylor, 2012). Furthermore, atmospheric peroxides are considered as the key components of secondary organic aerosol (SOA), which play a significant role in the formation and duration of haze pollution (Kroll and Seinfeld, 2008; Ziemann and Atkinson, 2012; Li et al., 2016). In addition, it has been suggested that atmospheric peroxides are toxic to ecosystem and may be the critical pollutants of forest decline (Hellpointner and G ¨ub, 1989; Chen et al., 2010). More importantly, peroxides in the particle phase have been found to act as reactive oxygen species (ROS) and result in adverse influence on human health (Ayres et al., 2008).

The concentrations of atmospheric peroxides are determined by their production and destruction. The known formation pathways of peroxides in the atmosphere are primary emissions, for instance, biomass burning (Lee et al., 1997, 1998; Yokelson et al., 2009), and secondary sources such as peroxy radical self/cross reactions and the ozonolysis of unsaturated volatile organic compounds (VOCs), as shown in Reaction (R1, R2) and (R3, R4), respectively (Hewitt and Kok, 1991; Neeb et al., 1997; Sauer et al., 2001; Chao et al., 2015; Winiberg et al., 2016). Additionally, atmospheric aqueous reactions in the bulk solution or on the surface of wet particles coupled with subsequent release to the gas phase could also generate peroxides in the troposphere (Wang et al., 2012; Liang et al., 2013a; Zhao et al., 2013a). The typical removal pathways of peroxides in the atmosphere are photolysis (R5, R6), reaction with OH

radicals (R7, R8) and physical deposition (Atkinson et al., 2006; Sander et al., 2011; Nguyen et al., 2015). Heterogeneous uptake by atmospheric aerosols is recognized as another significant sink for peroxides in the troposphere, especially in dusty and polluted urban areas (Zhao et al., 2013b; Wu et al., 2015).



In the past years, a number of field observations, laboratory studies and modelling research have been carried out to investigate the abundance and behavior of peroxides in the atmosphere (Chen et al., 2008; Mao et al., 2010; Huang et al., 2013; Liang et al., 2013a; Sarwar et al., 2013; Epstein et al., 2014; Fischer et al., 2015; Khan et al., 2015). Hydrogen peroxide (H_2O_2), hydroxymethyl hydroperoxide (HMHP, HOCH_2OOH), methyl hydroperoxide (MHP, CH_3OOH) and peroxyacetic acid (PAA, $\text{CH}_3\text{C}(\text{O})\text{OOH}$) are generally determined to be the principal peroxide compounds in the troposphere with their concentrations ranging from pptv (parts per trillion by volume) to ppbv (parts per billion by volume) (Lee et al., 2000; He et al., 2010; Zhang et al., 2010, 2012). However, to date, there have been limited studies concerned with atmospheric peroxides in the regions primarily affected by anthropogenic sources such as the North China Plain (NCP), which is a typical region with frequent biomass burning and suffering from serious haze pollution in China (Tao et al., 2012; Huang et al., 2014). Few numerical simulations focused on atmospheric peroxides in the NCP are conducted to examine whether the models can reproduce the observations of peroxides (Liang et al., 2013a). The impact of biomass burning and high aerosol loading on the atmospheric chemistry of peroxides over such a polluted region is poorly understood. Therefore, this work was carried out in order to make an endeavor

to fill in these research gaps.

In this study, we present a novel dataset of atmospheric speciated peroxides and explore their atmospheric chemistry at a rural site, Wangdu, which represents regional air pollution conditions of the NCP during Wangdu Campaign 2014. Given the diversity of emission sources and chemical transformation of atmospheric peroxides over this region, it is challenging to analyze the phenomena and understand the primary emission and secondary formation of peroxides in the atmosphere during this field observation. However, with the continuous measurements of atmospheric peroxides, physical parameters and other chemical species performed simultaneously, a quantitative assessment of the budget of atmospheric peroxides can be carried out employing the zero-dimensional model based on Master Chemical Mechanism (MCM) and constrained by observed meteorological parameters and trace gases, which provides a good opportunity to comprehensively facilitate our knowledge of the chemistry of atmospheric peroxides in the NCP. As far as we know, this is the first study to test whether current atmospheric peroxides related chemistry could explain the field observation in the rural area of the NCP. Through the comparison between measurement and simulation, our aim is to investigate the role of biomass burning and heterogeneous uptake on aerosols in the concentration of atmospheric peroxides, which helps to develop more robust mechanism in the model.

2 Experiments

2.1 Measurement site

Measurements of atmospheric peroxides were performed at Wangdu site (38.66 °N, 115.20 °E) in Baoding city, Hebei Province, a rural supersite for the Wangdu Campaign 2014 situated in the northwest of the NCP, about 200 km southwest of the mega-city Beijing. The surrounding regions of Wangdu site are mainly agricultural fields. There are almost no industries near this site. During the summer season, the air pollution is caused by the primary emission from biomass burning and secondary formation including photochemical and heterogeneous processes. The instruments were placed in a container with the sampling inlet approximately 5 m above the

ground. The continuous observation of atmospheric peroxides was conducted from 4 June to 7 July 2014.

2.2 Measurement methods

2.2.1 Measurement method for atmospheric peroxides

Atmospheric peroxide concentrations were investigated by an automated on-site high performance liquid chromatography (HPLC) with post-column enzyme derivatization and detected by fluorescence spectroscopy. Air samples were pumped through a glass scrubbing coil maintained at a controlled temperature of about 4°C to collect the peroxides in the atmosphere. The flow rate of air samples was set to be 2.7 standard L min⁻¹. The stripping solution, 5×10⁻³ M H₃PO₄ in water was delivered into the scrubbing coil collector. The flow rate of stripping solution was set to be 0.2 mL min⁻¹. Once the air samples mixed with the stripping solution in the collector, the mixture was carried by the mobile phase containing 5×10⁻³ M H₃PO₄ at 0.5 mL min⁻¹ and injected into HPLC. The peroxide components were separated after the mixture passed through HPLC column. With the catalysis of Hemin at ~40 °C, the derivatization reaction between peroxide components and para-hydroxyphenylacetic acid (PHPAA) produced the fluorescent matter that can be quantified by fluorescence detector. In this work, atmospheric peroxides were measured every 20 min. The collection efficiencies for hydrogen peroxide and organic peroxides were determined to be 100% and 85%, respectively. The detection limit of peroxides in the gas phase was about 10 pptv.

The interference of SO₂ on the sampling was estimated using the theoretical thermodynamic and kinetic analysis presented in Hua et al. (2008). Considering the rate constant for reaction between peroxides and S(IV) reported by Ervens et al. (2003) and the mean level of SO₂ was 7.0±7.0 ppbv during the campaign, the negative artifact caused by SO₂ interference for peroxides was calculated to be less than 15%. The influence of ambient relative humidity (RH) on the measurement of atmospheric peroxides was calculated following the method introduced by Liang et al. (2013b). The change of the concentration of atmospheric peroxides after this calibration is less

than 10%. Here, we did not correct the observational data for any artifacts due to the uncertainties from the theoretical estimation of peroxides loss that possibly result in new errors. The uncertainty of our observational data is estimated to be ~15%. Further details about our measurement method for atmospheric peroxides can be obtained from Hua et al. (2008).

2.2.2 Measurement methods for other pollutants and parameters

During Wangdu Campaign 2014, SO₂, CO, NO/NO₂ and O₃ were measured concurrently at this supersite using a suite of commercial instruments (Thermo 43i, 42i, 48i and 49i). HONO was measured every 2 min with a LOng Path Absorption Photometer (LOPAP) (Liu et al., 2016). C₂–C₁₀ non-methane hydrocarbons (NMHCs) were analyzed with a time resolution of 60 min by a custom-built online VOC analyzer using automated gas chromatography (GC) coupled with flame ionization detector (FID) or mass spectrometry (MS) technique (Wang et al., 2014). OH and HO₂ radicals were measured by laser-induced fluorescence (LIF) spectroscopy (Tan et al., 2016). Size distributions of aerosols (PM₁₀) were determined every 10 min with a Twin Differential Mobility Particle Sizer (TDMPS) and an Aerodynamic Particle Sizer (APS) to calculate dry particle surface area concentrations (S_a). Hygroscopic growth factor, f (RH), which is defined as the ratio of scattering coefficient for ambient aerosol to scattering coefficient for dry aerosol, was derived from the integrating nephelometer (Liu, 2015). Measurements of the mass concentration of PM_{2.5} were obtained by TEOM 1400A analyzer. Water-soluble ions (i.e., NH₄⁺, K⁺, Cl⁻, SO₄²⁻ and NO₃⁻) in PM_{2.5} was measured every 30 min with a Gas and Aerosol Collector (GAC) (Ye, 2015). Photolysis frequencies were derived from a spectro-radiometer (Bohn et al., 2008). Meteorological parameters including ambient temperature, relative humidity (RH), pressure, wind speed, wind direction and rainfall were monitored continuously by a weather station. The uncertainties (1σ) in these measurements are estimated as 5% for NO, O₃, and CO, 10% for H₂O, NO₂, HONO, NMHCs, and solar radiation, and 20% for S_a .

2.3 Model description

A zero-dimensional box model using a near-explicit mechanism, MCM Version 3.3.1 (<http://mcm.leeds.ac.uk/MCM/>) (Jenkin et al., 1997, 2003; Saunders et al., 2003; Jenkin et al., 2015) was employed to examine the influence of biomass burning and heterogeneous uptake on the budget of atmospheric peroxides. MCMv3.3.1 describes the degradation of 143 VOCs, leading to about 5800 species and 17000 reactions. In the current study, we extracted a subset of MCMv3.3.1 containing the reactions of atmospheric oxidants with measured VOCs and subsequent chemical products. Measurements of NO/NO₂, CO, O₃, HONO, NMHCs, temperature, pressure and H₂O were used as inputs to constrain the model calculations. The model ran with a 5-min time step and a spin-up time of 2 days to reach a steady state.

Photolysis frequencies were calculated by the Tropospheric Ultraviolet and Visible (TUV, version 5.2) model (Madronich, 2002), and further rescaled with the measured $j(\text{NO}_2)$. Dry deposition velocities of trace gases in our box model were parameterized as V_d/h (Seinfeld and Pandis, 2006), where V_d is the dry deposition rate of species and h is the height of planetary boundary layer (PBL). Dry deposition rates of HNO₃, PANs, organic nitrates, H₂O₂, organic peroxides and aldehydes incorporated in the model were set as $2.0 \times 10^{-5} \text{ s}^{-1}$, $5.0 \times 10^{-6} \text{ s}^{-1}$, $1.0 \times 10^{-5} \text{ s}^{-1}$, $1.0 \times 10^{-5} \text{ s}^{-1}$, $5.0 \times 10^{-6} \text{ s}^{-1}$ and $1.0 \times 10^{-5} \text{ s}^{-1}$, respectively at the PBL height of 1 km (Zhang et al., 2003; Emmerson et al., 2007; Lu et al., 2012; Guo et al., 2014; Li et al., 2014c; Liu et al., 2015; Nguyen et al., 2015). The PBL height over Wangdu during this campaign was derived from the hybrid single-particle lagrangian integrated trajectory (HYSPLIT) model (Draxler and Rolph, 2012), which varied between about 300 m at midnight and over 3000 m at noon.

The uncertainty of our model calculation derives from the uncertainty of observational data, PBL height and reaction rate coefficients in chemical mechanism. The total uncertainty in the model was estimated from the errors of all input parameters using error propagation, which is similar to the method that can be found in Hofzumahaus et al. (2009). On average, the modelled concentration of atmospheric peroxides had an uncertainty of approx. 60%.

In the present study, to explore the impact of the heterogeneous process on the

concentration of atmospheric peroxides, our box model is extended with the aerosol uptake of peroxides. The pseudo-first-order rate constant for the heterogeneous uptake of peroxides on ambient aerosols is parameterized as follows:

$$k = \frac{1}{4} \gamma \cdot v \cdot S_{aw} \quad (1)$$

(Jacob, 2000), where γ is the uptake coefficient, v is the mean molecular velocity, S_{aw} is the aerosol surface concentration corrected by the measured hygroscopic factor, $f(\text{RH})$ that could be expressed as $S_{aw} = S_a \times f(\text{RH})$.

3 Results and Discussion

3.1 General observations

The concentrations of peroxides in the atmosphere were measured continuously from 4 June to 7 July 2014. The predominant peroxides over Wangdu included H_2O_2 , MHP and PAA. Time series for atmospheric peroxides during Wangdu Campaign 2014 are illustrated in Fig. 1. The statistical data about the observed concentration of atmospheric peroxides are summarized and given in Table 1. It should be noted that values below the detection limit (D.L.) of our instrument were replaced by half of the D.L. in Fig. 1, Fig. 2 and statistical calculations. In this study, H_2O_2 accounted for ~70% of total detected peroxides ($\text{H}_2\text{O}_2 + \text{MHP} + \text{PAA}$) [similar to those determined at other rural sites in China](#). However, in our previous work, H_2O_2 contributed not more than 30% of total peroxides in the atmosphere over urban Beijing at the summer time of 2010 and 2011 (Liang et al., 2013b). This might be caused by the difference on the production and destruction of atmospheric peroxides between two sites. MHP and PAA were determined to be about 20% and 5% of total peroxides over Wangdu, respectively, which is similar to the results of other rural sites in China from our previous investigations (Zhang et al., 2010, 2012).

In the present work, on the basis of the latest national Ambient Air Quality Standards of China (GB3095-2012), the haze [day is defined as a day with daily-averaged \$\text{PM}_{2.5}\$ concentration over \$75 \mu\text{g m}^{-3}\$](#) . The haze pollution episode is defined as the event that a set of continuous days with daily-averaged $\text{PM}_{2.5}$

删除的内容:).

concentration exceeding $75 \mu\text{g m}^{-3}$, which has been used to distinguish non-haze and haze episode in the literature (Che et al., 2014; Zhang et al., 2016; Zheng et al., 2015; Zheng et al 2016). During this campaign, there were four haze pollution episodes at Wangdu site as follows: Episode 1 (4 June–6 June), Episode 2 (12 June–17 June), Episode 3 (29 June–3 July) and Episode 4 (5 July–7 July) with elevated average $\text{PM}_{2.5}$ concentrations ($75 \mu\text{g m}^{-3}$, $92 \mu\text{g m}^{-3}$, $79 \mu\text{g m}^{-3}$ and $99 \mu\text{g m}^{-3}$, respectively). In Episode 1, H_2O_2 , MHP and PAA were observed up to 11.3 ppbv, 0.9 ppbv and 1.5 ppbv, respectively. The maximum H_2O_2 concentration on 5 June was the highest value so far among the previously reported observations in urban, suburban and rural areas in China at summer time. The possible reason for this peak concentration at Wangdu site could be the primary emission from biomass burning combined with the secondary formation by the intense photochemical process. Nevertheless, owing to the lack of supporting data for other pollutants and parameters, it is difficult to identify the relative contributions of biomass burning versus photochemical formation to the burst of atmospheric peroxides on 5 June. In Episode 2, there was widespread and intensive biomass burning in the NCP as this observation period covered the local wheat harvest season. The evidence for biomass burning from the measurement of K^+ in $\text{PM}_{2.5}$ was illustrated in Fig. 2. The sudden raise of atmospheric peroxides was observed and further discussed in Sect. 3.3. In Episode 3, there was a substantial decline of H_2O_2 level during this typical haze event compared with the foregoing two episodes, which can be ascribed to the heterogeneous uptake of peroxides on atmospheric aerosols on haze days over Wangdu (See Sect. 3.4). In Episode 4, Wangdu was significantly impacted by the regional transport (Ye, 2015). The concentrations of atmospheric peroxides remained relatively low compared with Episode 1 and Episode 2. In addition to the above-mentioned episodes, it was relatively clear between 8 June and 11 June and 27 June and 28 June, with mean $\text{PM}_{2.5}$ concentrations under $40 \mu\text{g m}^{-3}$. The intermittent thunderstorm activities occurred from 19 June to 25 June that caused the electric power failure and several data gaps.

删除的内容: exceeds

3.2 Peroxide simulation

In this study, we employed a box model based on the MCMv3.3.1 to simulate H₂O₂, MHP and PAA concentrations. Here, to explore the atmospheric chemistry of peroxides on non-haze, biomass burning and haze days, the observational data from 8 June to 11 June (Phase I), from 15 June to 17 June (Phase II) and from 29 June to 3 July (Phase III) in 2014 were selected as phase of interest and analyzed in detail using box model in the following sections. The temporal variations of meteorological parameters, chemical species and atmospheric peroxides for the whole campaign are displayed in Fig. 2. The observed and calculated levels of atmospheric peroxides for the three phases are illustrated in Fig. 3. During these case study phases, 75% of the wind speed data were $\leq 2.2 \text{ m s}^{-1}$ and the mean value was 1.6 m s^{-1} . It has been shown that the atmospheric lifetimes of peroxides are on the order of several hours as reported previously (He et al., 2010; Wu et al., 2015), implying that the effect of regional transport or dilution on the concentrations of atmospheric peroxides was of little significance over Wangdu. Hence, the regional-scale transport can be excluded in our box model and the budgets of peroxides are, to a large extent, dependent on local chemical processes during the observation.

In the Phase I, as shown in Fig. 4, the model base case prediction of H₂O₂ level had good performance in the daytime (06:00–18:00 local time), which was 1–2 times higher than the measurement results. This seems to be explained by the model-measurement uncertainty. Similarly, a previous observation carried out at a suburban site also showed reasonable model-measurement agreement in H₂O₂ level on sunny days (Guo et al., 2014). The excellent description yielded by the model base case indicated that the production and destruction of H₂O₂ in the atmosphere on non-haze days were calculated correctly based on the current understanding of atmospheric peroxide related chemistry. However, the simulation in the nighttime (18:00–06:00 local time) during the Phase I demonstrated an obvious overestimation compared to the observation by a factor of 4–6 and up to an order of magnitude. This large discrepancy between calculated and observed results is speculated to be resulted

from the underestimation of sink terms as the key precursors governing the formation of atmospheric peroxides are constrained by the observation and the overestimation of source terms can be ruled out. It is consistent with the comparison of the simulated and observed H₂O₂ concentration over urban Beijing, in which the explanation for the overprediction of H₂O₂ level on haze days was thought to be the heterogeneous processes on liquid or solid particles that were missing from the current atmospheric chemistry model (Liang et al., 2013b). Considering the high aerosol loading in the NCP and the higher aerosol surface area concentration at nighttime (1158 μm² cm⁻³) than that at daytime (773 μm² cm⁻³) in the Phase I, we believe that the missing sink for atmospheric peroxides in the model base case is probably heterogeneous uptake of peroxides occurring on aerosols. The strengths of the missing sinks for H₂O₂, MHP and PAA quantified by the difference between modelled and measured peroxide concentrations were about 0.24 ppbv h⁻¹, 0.09 ppbv h⁻¹ and 0.03 ppbv h⁻¹ on average, respectively, which were on the same order of magnitude as the known loss rates of atmospheric peroxides during the Phase I.

In the Phase II, the comparison of the modelled and measured peroxide concentrations in Fig. 3 displays that the observed magnitude of atmospheric peroxides was unexpectedly large, indicating a missing source for peroxides. Such a strong imbalance was found only in the Phase II during the whole campaign. In the past, the higher-than-expected concentrations of atmospheric peroxides have also been reported by Lee et al. (1997), in which H₂O₂, MHP, PAA and other organic peroxides levels elevated near biomass burning plumes. Given the frequent fire emissions in the NCP during the Phase II that are quite similar to the conditions in Lee et al. (1997), it appears that the significant mismatch can be attributed to the direct production from biomass burning (See Sect. 3.3).

In the Phase III, the calculated values in the model base case showed a general tendency to strongly overestimate the observed values (Fig. 3). [The modelled and measured shapes of the diurnal cycle of atmospheric peroxides were different.](#) The haze arose on 29 June with the elevated PM_{2.5} concentration. The diffusion condition was poor as the CO concentration was enhanced. The precursors of atmospheric

peroxides also accumulated on 29 June and 30 June. The modelled peroxide concentrations over 10 times higher than the measured peroxide concentrations. On 1 July and 3 July, the daily-averaged $PM_{2.5}$ concentration was 1.6 times higher than those on 29 June and 30 June. However, the photolysis frequencies and the PBL height on 1 July and 3 July were about half of those on 29 June and 30 June, which weakened the secondary formation of atmospheric peroxides and strengthened the loss of atmospheric peroxides via dry deposition. Although the haze on 1 July and 3 July was more serious than that on 29 June and 30 June, the ratios of modelled to measured peroxide concentrations on 1 July and 3 July were much lower than those on 29 June and 30 June. As there was a typical haze event during the Phase III, the model-measurement imbalance was probably due to the missing sink for atmospheric peroxides, which was the same deficiency in the model as that in the Phase I. It can be seen in Fig. 3 that with the inclusion of heterogeneous reactions on aerosol particles, the simulated concentrations of atmospheric peroxides were apparently improved [and the modelled shape of the diurnal cycle of \$H_2O_2\$ was closer to the measured shape](#), which is further quantified in Sect. 3.4.

Before exploring the impact of biomass burning and heterogeneous uptake on the chemistry of atmospheric peroxides, we performed a model test by implementing the newly proposed chemical mechanisms for $CH_3C(O)O_2$ and CH_3O_2 related chemistry in MCMv3.3.1, as listed in Table 2. The rate constant and the branching ratios of the $CH_3C(O)O_2 + HO_2$ reaction that was the major pathway for the formation of PAA in this model scenario were modified according to the recent laboratory study conducted by Winiberg et al. (2016). Additionally, we also incorporated the reaction between CH_3O_2 radicals and OH radicals, which has as yet seldom been involved in atmospheric chemistry model. The reaction between CH_3O_2 radicals and OH radicals is recognized as an important sink for CH_3O_2 radicals with non-negligible effect on subsequent formation of MHP under remote conditions (Bossolasco et al., 2014; Fittschen et al., 2014). As shown in Fig.3, the model run containing newly-proposed mechanisms did not have a remarkable influence on the simulated results of H_2O_2 in comparison to the model base case. But a slight difference of up to ~20% between

calculated and observed MHP can be noted at night, resulting from the additional removal pathway of CH_3O_2 radicals from the noon to the sunset. The increase of over 70% in rate constant and the reduction of about 10% in the branching ratio of the reaction $\text{CH}_3\text{C}(\text{O})\text{O}_2 + \text{HO}_2 \rightarrow \text{CH}_3\text{C}(\text{O})\text{OOH}$ generated systematically 1.5 times higher PAA concentration in this model scenario than that in the model base case. Nevertheless, although the modelled PAA during the Phase II can be raised close to the level of the observation, the concentrations of atmospheric peroxides were not fully captured by the model with the implementation of newly proposed mechanisms (Fig. 3). The additional chemical mechanisms embedded in the model only have a marginal impact that is not sufficient to match the observed peroxides in the atmosphere. The efficient source or sink for the reproduction of the observation will be deeply investigated below.

As outlined in the introduction, the sources of H_2O_2 , MHP and PAA are the direct emission from biomass burning and the photochemical oxidation of VOC precursors via HO_2 , CH_3O_2 and $\text{CH}_3\text{C}(\text{O})\text{O}_2$ formation. However, it is still difficult to determine the contributions of VOC precursors at a species level. Here, to gain further insight into the secondary chemical transformation of atmospheric peroxides at Wangdu site, the sensitivity study was conducted to track out the major VOC precursors of atmospheric peroxides. An indirect approach referring to the relative incremental reactivity (RIR) concept for ozone formation in Cardelino and Chameides (1995) was adopted for the sensitivity study using the numerical model with the application of the MCMv3.3.1. MCM describes the explicit degradations of individual VOC species, and hence facilitates to quantify the role of VOC in the secondary formation of atmospheric peroxides at a species level. In this work, the definition of RIR is the ratio of reduction in the production rates of atmospheric peroxides to the reduction of VOC precursor abundances by 25% compared to the model base case, which can be regarded as a proxy for the influence of a specific VOC on the *in-situ* formation of atmospheric peroxides. Phase I and Phase III were selected for the analysis, while the Phase II was precluded from the analysis as it was affected by the local emission not included in the model base case.

Fig. 5 displays the average RIRs of H₂O₂, MHP and PAA for alkane, alkene, aromatic and NO_x classes as well as the seven most important individual VOC precursors. The results demonstrate that the formation of H₂O₂ was sensitive to alkenes and insensitive to alkanes, aromatics and NO_x. The production of MHP and PAA shows a strong dependence on alkenes and NO_x, while it is relatively independent of aromatics and alkanes other than methane. Isoprene and trans-2-butenes turn out to be the key VOC species controlling the formation of atmospheric peroxides. Moreover, cis-2-butene, cis-2-pentene, propene and 1,2,4-trimethylbenzene also seem to be the major individual VOC precursors as evidence by Fig. 5. Methane is noticed to be an important contributor to the formation of MHP. Such a list of VOC species is not consistent with our previous studies over urban Beijing that suggested aromatics (i.e., toluene and dialkylbenzenes) as the dominant VOC precursor of atmospheric peroxides (Zhang et al., 2010; Liang et al., 2013b). It reflects that the relative significance of individual VOC precursors varies from place to place. The distinction between the two sites is attributable to the relatively more abundant isoprene, anthropogenic alkenes and much less reactive aromatics at the rural site in the NCP than those at the urban site, Beijing.

With the identification of a small class of key VOC precursors contributing to the formation of peroxides in the atmosphere of NCP, the effective control strategies for mitigating the pollution resulted from atmospheric peroxides can be formulated. In the NCP, it has been revealed that the vehicular exhaust is the predominant source responsible for the VOC species such as propene, trans/cis-2-butenes and trimethylbenzenes in the surrounding areas of the observation site (Yuan et al., 2009; Ran et al., 2011; Li et al., 2014b; Li et al., 2015; Wu et al., 2016), while the vegetation governs the release of isoprene. It is recommended to take measures for vehicle emission control and land use management (e.g. modifying the amount and types of vegetation) in order to mitigate the pollution of atmospheric peroxides in the NCP and hence alleviate their potential harmful effects on air quality, human health and ecosystem.

3.3 Direct production of peroxides from biomass burning

In the Phase II, the levels of H₂O₂, MHP and PAA were highly elevated in comparison with the other phases, which could not be explained by the photochemical process in the model base case alone. It provides us a hint that an additional formation pathway is required to improve the results of model simulation. In Sect. 3.2, we hypothesized that the direct production of peroxides from biomass burning should serve as an essential source for the unexpected burst of atmospheric peroxides. Here, we tested the hypothesis by means of the box model and linear regression with the observation data from three events mentioned below during the Phase II. It is well known that CO and K⁺ can be used as the reference for the biomass combustion (Koppmann et al., 2005; Reid et al., 2005; Li et al., 2007; Sullivan et al., 2008; Cheng et al., 2013, 2014; Li et al., 2014a; Wang et al., 2015). The averaged CO levels were 0.42±0.16 ppmv, 0.79±0.20 ppmv and 0.61±0.20 ppmv for the Phase I, Phase II and Phase III, respectively. The mean K⁺ concentrations were about 0.64±1.19 µg m⁻³ for the Phase I, 2.51±1.53 µg m⁻³ for the Phase II and 0.26±0.21 µg m⁻³ for the Phase III. The abundances of CO and K⁺ during the Phase II were higher than that during the Phase I and Phase III, which is consistent with the observed intensive biomass burning activities at Wangdu site (Ye, 2015). Nevertheless, in addition to the biomass burning, CO level in the NCP was also affected by anthropogenic activities with the regional transport of polluted air masses, for example, the urban plumes. It has been proved that airborne K⁺ is acceptable as the tracer for biomass burning during summertime in the NCP (Cheng et al., 2013; Wang et al., 2015). The concentrations of CH₃CN, another tracer for biomass burning, measured by Proton-transfer-reaction mass spectrometry (PTR-MS) exhibited similar temporal variation to the concentrations of K⁺ during the Wangdu campaign 2014 except on 10 June (X. Huang, personal communication, 2015). Therefore, K⁺ might be a better indicator of biomass burning than CO here. In the Phase II, we identified several biomass burning events with concentrations of K⁺ more than twice the mean value of that in the Phase I and Phase III. Considering the availability of the observation data for atmospheric

peroxides, we focused our analysis on three events as follows: Event I (17:00–20:00 on 15 June), Event II (16:00–19:00 on 16 June) and Event III (12:00–15:00 on 17 June) with durations of over 3 hours.

As illustrated in Fig. 3, the model base case cannot reproduce the measurements for atmospheric peroxides in the three events. To match the observations, the primary sources for H₂O₂, MHP and PAA were applied to our model. The strengths of the primary sources for H₂O₂, MHP and PAA quantified by the difference between modelled and measured peroxide concentrations were about 0.25–1.80 ppbv h⁻¹, 0.24–0.44 ppbv h⁻¹ and 0.02–0.16 ppbv h⁻¹, respectively. These values are on the order of the known secondary production rates of atmospheric peroxides during the three events. [The impact of primary sources of the peroxides on HO_x radicals was limited with the increase of OH radicals not more than 10% and the increase of HO₂ radicals not more than 5%.](#) It should be pointed out that the estimation was associated with large uncertainties since it did not include the heterogeneous uptake of peroxides by aerosols in the model here. In view of the possible additional sink for atmospheric peroxides as discussed in Sect. 3.4 below, the primary sources for H₂O₂, MHP and PAA might represent the lower limit. The effect of biomass burning on the levels of atmospheric peroxides might be underestimated as well. We underscore that there might exist even larger missing sources for H₂O₂, MHP and PAA due to the scarcity of some important removal pathways of atmospheric peroxides in the model in this section.

The results of linear regression involving correlation coefficients and their statistical significance of H₂O₂, MHP and PAA to CO and K⁺ were listed in Table 3 for the three biomass burning events. The relationships between atmospheric peroxides and biomass burning indicators were analyzed separately for each event owing to the variability of fire emissions. A notable trend between atmospheric peroxides and K⁺ was found with correlation coefficients exceeding over the significance threshold, which provided a convincing evidence for the direct production of peroxides from biomass burning as the additional source. Moreover, it was noticed that CO agreed well with K⁺ for the Event I and Event II, exhibiting

excellent correlation with atmospheric peroxides (Table 3). The enhancement ratios of H₂O₂, MHP and PAA relative to CO were calculated to be at the magnitude of 10⁻³, which are similar to the enhancement signals of atmospheric peroxides to CO obtained near biomass fires from flights published by Lee et al. (1997).

It is noteworthy that several other chemical processes, for example, secondary formation via the photooxidation of potential unmeasured short-lived VOC species emitted from biomass fires prior to our sampling of the plume at the observational site are alternatives to the direct production from biomass burning as the missing source of atmospheric peroxides in the model. Thus, it appears necessary and desirable to further distinguish the extent to which atmospheric peroxides are generated via the direct production or secondary formation from biomass burning in future research. Laboratory studies are required to simulate the biomass fires in the NCP using combustion chamber to critically characterize the emission factors of atmospheric peroxides to CO and determine their generation mechanisms. Also, more reliable aircraft and ground-based field measurements for the variation of atmospheric peroxides during the harvest seasons in China need to be carried out and would help to shed some light on the role of biomass burning in the abundance of peroxides in the atmosphere.

3.4 Heterogeneous uptake of peroxides by aerosol

In Sect. 3.2, heterogeneous uptake on atmospheric particles was considered as a suitable explanation for the missing sink for H₂O₂, MHP and PAA during the Phase I and Phase III in view of substantial aerosol loading in the NCP that provided considerable sites for heterogeneous reactions. Here, we make an attempt to implement a parameterization of heterogeneous uptake by aerosols in our box model to resolve the deviation between the simulated and observed data (See Sect. 2.3). Using the uptake coefficient of 1×10^{-3} for H₂O₂, MHP and PAA, a good agreement between the modelled and measured temporal variation of atmospheric peroxides can be obtained in Phase I and Phase III by taking into account the combined model-measurement error that is conservatively assumed to be ~50% (Fig. 3). [The](#)

带格式的：缩进：首行缩进：1 字符，定义网格后自动调整右缩进，段落间距段前：0 磅，段后：0 磅

[modelled and measured shape of the diurnal cycle of H₂O₂ in the Phase I and the Phase III are similar.](#)

The calculated H₂O₂, MHP and PAA with the coupling of the heterogeneous reaction were on average decreased by about 75% compared to the results in the model base case during the Phase III. The uptake coefficient of 1×10^{-3} approached the upper limit of the laboratory measured value for H₂O₂ on mineral dust (9×10^{-4}) reported by Pradhan et al. (2010), but a little higher than the previous measured values on ambient PM_{2.5} of $(1-5) \times 10^{-4}$ during the summertime over urban Beijing (Wu et al., 2015). It is reasonable as Wu et al. (2015) pointed out that the uptake coefficients for H₂O₂ and organic peroxides on ambient PM_{2.5} are in the same range and show no obvious differences between daytime and nighttime or between non-hazy and hazy conditions.

With the adoption of heterogeneous uptake coefficients of 1×10^{-3} , we evaluated the sinks of atmospheric peroxides in the Phase I and Phase III that represented non-haze and haze conditions, respectively. The mean surface area concentration that was corrected for the hygroscopic growth of aerosol was measured to be $968 \mu\text{m}^2 \text{cm}^{-3}$ for Phase I and $1491 \mu\text{m}^2 \text{cm}^{-3}$ for Phase III. Fig. 6 demonstrated that the destruction of atmospheric peroxides during the two phases originated from a diversity of sinks, including photolysis, OH-initiated reaction, dry deposition and heterogeneous uptake. It has been reported that the heterogeneous reaction is the most important sink for H₂O₂ in urban (Liang et al., 2013b) and suburban areas (Guo et al., 2014). In contrast, OH-initiated reaction and dry deposition were regarded as the major removal pathways of organic peroxides in rural (Zhang et al., 2012) and forests areas (Nguyen et al., 2015). Here, heterogeneous uptake by aerosols turned out to be the predominant sink for atmospheric peroxides in the NCP, accounting for more than 60% of the total loss, while dry deposition became the marginal removal pathway that contributed ~10% to the destruction of H₂O₂, MHP and PAA. The role of OH-initiated reaction in the total loss varied between the speciated peroxides with no more than 30%. Photolysis only represented a minor contribution (<3%). The most prominent feature on haze days was the larger loss of atmospheric peroxides via heterogeneous process,

demonstrating the enhanced impact of aerosols on the sink of peroxides during the haze episode compared to that during the non-haze episode.

On the basis of the analysis above, we investigated the atmospheric lifetime of peroxides in the NCP with the integration of observation and modelling. The lifetime of H₂O₂, MHP and PAA were estimated with the concentration-to-time curves between 18:00 and 24:00 LT as the formation of atmospheric peroxides was weak and negligible during this phase. The average lifetime obtained from the field observation between 18:00 and 24:00 LT in the Phase I was around 4.0 h, 5.6 h and 3.1 h for H₂O₂, MHP and PAA, respectively, which was similar to the values [between 18:00 and 24:00 LT in the Phase I](#) of 3.4 h, 4.3 h and 5.2 h for H₂O₂, MHP and PAA, respectively, given by our modeling simulation. The [daily-averaged](#) lifetime of atmospheric peroxides in the Phase III was ~40% smaller than that in the Phase I. Using the box model, the [daily-averaged lifetime of](#) atmospheric H₂O₂, MHP and PAA during the whole of Phase I and Phase III were calculated to be about 2.1 h, 2.3 h and 3.0 h, respectively. This is comparable to the literature results with the inclusion of heterogeneous reaction (Liang et al., 2013b; Wu et al., 2015), but notably shorter than the recent studies conducted by Khan et al. (2015) and Nguyen et al. (2015) without the coupling of the heterogeneous process. [Although dry deposition is thought to dominate the atmospheric lifetime of peroxides in previous studies \(Reeves and Penkett, 2003\), its role in the lifetime of atmospheric peroxides is insignificant during Wangdu Campaign 2014. The sensitivity of modelled relative loss of dry deposition to the uncertainties in the planetary boundary layer height was low as the contribution of dry deposition to the loss of H₂O₂ in Phase I decreased no more than 10% with the PBL height doubled. The simulated daily-averaged](#) lifetime of atmospheric peroxides can be over 10 h by supposing that the loss of H₂O₂, MHP and PAA is merely due to photolysis, OH-initiated reaction and dry deposition. It emphasizes that heterogeneous uptake on aerosols determines the atmospheric lifetime of peroxides.

It is worth noting that the heterogeneous uptake of peroxides by aerosols in the atmospheric chemical model is still controversial as it is possibly that the aerosol

删除的内容: lifetimes of

删除的内容: The simulated

uptake of HO₂ radicals is the explanation for the missing sink. This raises an interesting question of whether HO₂ uptake or peroxide uptake is responsible for the imbalance between observation and modelling. It has been inferred by formerly published literature that aerosol uptake of HO₂ radicals is the major reason for the overprediction of the levels of atmospheric peroxides in the model (de Reus et al., 2005; Mao et al., 2013; Guo et al., 2014). Nevertheless, it is apparent that the extent of HO₂ heterogeneous degradation depends on the atmospheric environment, especially the concentration and property of aerosol particles that vary under different conditions. The measured HO₂ concentrations at Wangdu site are close to the modelled HO₂ concentrations by the box model merely with the gas-phase regional atmospheric chemical mechanism (RACM) comprised (K. Lu, personal communication, 2015). Hence, the impact of aerosol uptake of HO₂ radicals on the concentration of atmospheric peroxides is insignificant during Wangdu Campaign 2014 and not taken into account in our model, while heterogeneous uptake of atmospheric peroxides by aerosols is exclusively adopted to improve the reproduction of the observation in the two phases above.

It has been inferred that heterogeneous uptake of peroxides on ambient PM_{2.5} probably results from solid surface reactions and aerosol aqueous reactions (Wu et al., 2015), for instance, “Fenton-like” reactions between peroxides and transition metal ions, which is supported by the laboratory studies (Chevallier et al., 2004; Deguillaume et al., 2005) and field observation (Liang et al., 2013b; Guo et al., 2014). Nevertheless, the detailed heterogeneous mechanism containing individual reaction channels was not included in the present work owing to the chemical complexity of the ambient aerosol. Given the potential importance of atmospheric peroxide compounds on the generation of HO_x radicals and aerosol ROS, the aging of mineral dust and SOA and the formation of haze (Huang et al., 2015; Pöschl and Shiraiwa, 2015; Zhang et al., 2015; Li et al., 2016), more comprehensive investigations including laboratory, field and modelling studies on the heterogeneous uptake processes of H₂O₂, MHP, PAA and other peroxides are indispensable to provide

concrete evidence to elucidate the chemical budget of atmospheric peroxides in the future.

4 Conclusions

Atmospheric peroxides including H₂O₂, MHP and PAA were measured at a rural site during the Wangdu Campaign 2014. The maximum H₂O₂ concentration was observed to be 11.3 ppbv, which was the highest value compared with previous observations in China. The concentrations of atmospheric peroxides were highly elevated during the biomass burning activities, but underwent substantial decline during the haze events. With the application of an observation-based model combining measured meteorological parameters and trace gases, we analyzed the chemical budget of peroxides under biomass burning, non-haze and haze conditions. Photochemical formation of atmospheric peroxides was attributed to a small class of alkenes, while it was insensitive to alkanes and aromatics. The key VOC precursors controlling the formation of peroxide compounds were identified to be isoprene, trans/cis-2-butenes, cis-2-pentene, propene and trimethylbenzene.

The base model simulation (MCMv3.3.1) underpredicted the levels of atmospheric peroxides during biomass burning events compared with the measurement. The direct production from biomass burning was regarded as the explanation for the unexpected burst of peroxides. To improve the simulated concentrations, the strengths of the primary emissions from biomass burning should be on the same order of the known secondary production rates of atmospheric peroxides. Moreover, the model base case also overpredicted the concentrations of atmospheric peroxides on haze days in comparison with the observation. The heterogeneous uptake by aerosols was suggested to be responsible for the attenuation of peroxides. The model could reproduce the observed values with the introduction of heterogeneous process using the uptake coefficient of 1×10^{-3} for atmospheric peroxides. According to the closure between observed and calculated concentrations, the heterogeneous uptake on aerosol particles was found to be the predominant sink for atmospheric peroxides, accounting for more than 60% of the total loss, followed by the OH-initiated reaction (<30%) and

dry deposition (~10%). The mean atmospheric lifetime of peroxides in summer in the NCP was estimated to be around several hours that was in good agreement with previous laboratory studies for the aerosol uptake of peroxides, indicating that heterogeneous reaction determines the atmospheric lifetime of peroxides.

In view of the importance of peroxides in tropospheric oxidation capacity and formation potential of secondary aerosols, more reliable investigations focused on the biomass burning emission factors and detailed heterogeneous mechanism of speciated peroxides are urgently required to further quantitatively evaluate the role of biomass burning and heterogeneous uptake in the abundance as well as budget of atmospheric peroxides and facilitate our knowledge of the formation of haze pollution.

Acknowledgements. This work was funded by the National Natural Science Foundation of China (grants 41275125, 21190051, 21190053, 21477002, and 41421064). The authors would like to thank Min Shao group (Peking University) for their VOCs data and Alfred Wiedensohler group (Leibniz Institute for Tropospheric Research) for their particle surface area concentrations data. The authors wish to gratefully thank the entire Wangdu Campaign 2014 team for the support and collaboration at Wangdu site.

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and Subcommittee, I.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II—gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, 2006.
- Ayres, J. G., Borm, P., Cassee, F. R., Castranova, V., Donaldson, K., Ghio, A., Harrison, R. M., Hider, R., Kelly, F., Kooter, I. M., Maranok, F., Maynard, R. L., Mudwaym, I., Nel, A., Sioutas, C., Smith, S., Baeza-Squibank, A., Chon, A., Duggan, S., and Froines, J.: Evaluating the toxicity of airborne particulate matter and nanoparticles by measuring oxidative stress potential—a workshop report and consensus statement, *Inhal. Toxicol.*, 20, 75–99, 2008.
- Bohn, B., Corlett, G. K., Gillmann, M., Sanghavi, S., Stange, G., Tensing, E.,

- Vrekoussis, M., Bloss, W. J., Clapp, L. J., Kortner, M., Dorn, H.P., Monks, P. S., Platt, U., Plass-Dulmer, C., Mihalopoulos, N., Heard, D. E., Clemitshaw, K. C., Meixner, F. X., Prevot, A. S. H., and Schmitt, R.: Photolysis frequency measurement techniques: results of a comparison within the ACCENT project, *Atmos. Chem. Phys.*, 8, 5373–5391, 2008.
- Bossolasco, A., Faragó E. P., Schoemaeker, C., and Fittschen, C.: Rate constant of the reaction between CH_3O_2 and OH radicals, *Chem. Phys. Lett.*, 593, 7–13, 2014.
- Calvert, J. G., Lazrus, A., Kok, G. L., Heikes, B. G., Walega, J. G., Lind, J., and Cantrell, C. A.: Chemical mechanisms of acid generation in the troposphere, *Nature*, 317, 27–35, 1985.
- Cardelino, C. A., and Chameides, W. L.: An observation-based model for analyzing ozone precursor relationships in the urban atmosphere, *J. Air Waste Manage. Assoc.*, 45, 161–180, 1995.
- Chao, W., Hsieh, J. T., and Chang, C. H.: Direct kinetic measurement of the reaction of the simplest Criegee intermediate with water vapor, *Science*, 347, 751–754, 2015.
- Che, H., Xia, X., Zhu, J., Li, Z., Dubovik, O., Holben, B., Goloub, P., Chen, H., Estelles, V., Cuevas-Agulló E., Blarel, L., Wang, H., Zhao, H., Zhang, X., Wang, Y., Sun, J., Tao, R., Zhang, X. and Shi, G.: Column aerosol optical properties and aerosol radiative forcing during a serious haze-fog month over North China Plain in 2013 based on ground-based sunphotometer measurements, *Atmos. Chem. Phys.*, 14, 2125–2138, 2014.
- Chen, X., Aoki, M., Takami, A., Chai, F. H., and Hatakeyama, S.: Effect of ambient-level gas-phase peroxides on foliar injury, growth, and net photosynthesis in Japanese radish (*Raphanus sativus*), *Environ. Pollut.*, 158, 1675–1679, 2010.
- Chen, Z. M., Wang, H. L., Zhu, L. H., Wang, C. X., Jie, C. Y., and Hua, W.: Aqueous-phase ozonolysis of methacrolein and methyl vinyl ketone: a potentially important source of atmospheric aqueous oxidants, *Atmos. Chem. Phys.*, 8, 2255–2265, 2008.
- Cheng, Y., Engling, G., He, K. B., Duan, F. K., Ma, Y. L., Du, Z. Y., Liu, J. M., Zheng,

- M., and Weber, R. J.: Biomass burning contribution to Beijing aerosol, *Atmos. Chem. Phys.*, 13, 7765–7781, 2013.
- Cheng, Y., Engling, G., He, K. B., Duan, F. K., Du, Z. Y., Ma, Y. L., Liang, L. L., Lu, Z. F., Liu, J. M., Zheng, M., and Weber, R. J.: The characteristics of Beijing aerosol during two distinct episodes: Impacts of biomass burning and fireworks, *Environ. Pollut.*, 185, 149–157, 2014.
- Chevallier, E., Jolibois, R. D., Meunier, N., Carlier, P., and Monod, A.: “Fenton-like” reactions of methylhydroperoxide and ethylhydroperoxide with Fe^{2+} in liquid aerosols under tropospheric conditions, *Atmos. Environ.*, 38, 921–933, 2004.
- de Reus, M., Fischer, H., Sander, R., Gros, V., Kormann, R., Salisbury, G., Van Dingenen, R., Williams, J., Zöllner, M., and Lelieveld, J.: Observations and model calculations of trace gas scavenging in a dense Saharan dust plume during MINATROC, *Atmos. Chem. Phys.*, 5, 1787–1803, 2005.
- Deguillaume, L., Leriche, M., Desboeufs, K., Mailhot, G., George, C., and Chaumerliac, N.: Transition metals in atmospheric liquid phases: sources, reactivity, and sensitive parameters, *Chem. Rev.*, 105, 3388–3431, 2005.
- Draxler, R. R., and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) model access via NOAA ARL READY website (<http://www.arl.noaa.gov/ready/hysplit4.html>), NOAA Air Resources Laboratory, Silver Spring, MD, 2012.
- Emmerson, K. M., Carslaw, N., Carslaw, D. C., Lee, J. D., McFiggans, G., Bloss, W. J., Gravestock, T., Heard, D. E., Hopkins, J., Ingham, T., Pilling, M. J., Smith, S. C., Jacob, M., and Monks, P. S.: Free radical modelling studies during the UK TORCH Campaign in Summer 2003, *Atmos. Chem. Phys.*, 7, 167–181, 2007.
- Epstein, S. A., Blair, S. L., and Nizkorodov, S. A.: Direct photolysis of α -pinene ozonolysis secondary organic aerosol: effect on particle mass and peroxide content, *Environ. Sci. Technol.*, 48, 11251–11258, 2014.
- Ervens, B., George, C., Williams, J. E., Buxton, G. V., Salmon, G. A., Bydder, M., Wilkinson, F., Dentener, F., Mirabel, P., Wolke, R. and Herrmann, H.: CAPRAM 2.4 (MODAC mechanism): an extended and condensed tropospheric aqueous phase

- mechanism and its application, *J. Geophys. Res.*, 108, 4426, 2003.
- Fischer, H., Pozzer, A., Schmitt, T., Jöckel, P., Klippel, T., Taraborrelli, D., and Lelieveld, J.: Hydrogen peroxide in the marine boundary layer over the South Atlantic during the OOMPH cruise in March 2007, *Atmos. Chem. Phys.*, 15, 6971–6980, 2015.
- Fittschen, C., Whalley, L. K., and Heard, D. E.: The reaction of CH_3O_2 radicals with OH radicals: a neglected sink for CH_3O_2 in the remote atmosphere, *Environ. Sci. Technol.*, 48, 7700–7701, 2014.
- Guo, J., Tilgner, A., Yeung, C., Wang, Z., Louie, P. K. K., Luk, C. W. Y., Xu, Z., Yuan, C., Gao, Y., Poon, S., Herrmann, H., Lee, S., Lam, K. S. and Wang, T.: Atmospheric peroxides in a polluted subtropical environment: seasonal variation, sources and sinks, and importance of heterogeneous processes, *Environ. Sci. Technol.*, 48, 1443–1450, 2014.
- He, S. Z., Chen, Z. M., Zhang, X., Zhao, Y., Huang, D. M., Zhao, J. N., Zhu, T., Hu, M., and Zeng, L. M.: Measurement of atmospheric hydrogen peroxide and organic peroxides in Beijing before and during the 2008 Olympic Games: chemical and physical factors influencing their concentrations, *J. Geophys. Res.*, 115, D17307, 2010.
- Hellpointner, E., and Gäb, S.: Detection of methyl, hydroxymethyl and hydroxyethyl hydroperoxides in air and precipitation, *Nature*, 631–634, 1989.
- Hewitt, C. N., and Kok, G. L.: Formation and occurrence of organic hydroperoxides in the troposphere: laboratory and field observations, *J. Atmos. Chem.*, 12, 181–194, 1991.
- Hofzumahaus, A., Rohrer, F., Lu, K. D., Bohn, B., Brauers, T., Chang, C. C., Fuchs, H., Holland, F., Kita, K., Kondo, Y., Li, X., Lou, S. R., Shao, M., Zeng, L. M., Wahner, A. and Zhang, Y. H.: Amplified trace gas removal in the troposphere, *Science*, 324, 1702–1704, 2009.
- Hua, W., Chen, Z. M., Jie, C. Y., Kondo, Y., Hofzumahaus, A., Takegawa, N., Chang, C. C., Lu, K. D., Miyazaki, Y., Kita, K., Wang, H. L., Zhang Y. H., and Hu, M.: Atmospheric hydrogen peroxide and organic hydroperoxides during

- PRIDE-PRD'06, China: their concentration, formation mechanism and contribution to secondary aerosols, *Atmos. Chem. Phys.*, 8, 6755–6773, 2008.
- Huang, D., Chen, Z. M., Zhao, Y., and Liang, H.: Newly observed peroxides and the water effect on the formation and removal of hydroxyalkyl hydroperoxides in the ozonolysis of isoprene, *Atmos. Chem. Phys.*, 13, 5671–5683, 2013.
- Huang, L. B., Zhao, Y., Li, H., and Chen, Z. M.: Kinetics of heterogeneous reaction of sulfur dioxide on authentic mineral dust: effects of relative humidity and hydrogen peroxide, *Environ. Sci. Technol.*, 49, 10797–10805, 2015.
- Huang, R. J., Zhang, Y. L., Bozzetti, C., Ho, K. F., Cao, J. J., Han, Y. M., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbazade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prévôt A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, 2014.
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, 34, 2131–2159, 2000.
- Jenkin, M. E., Saunders, S. M., Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, 1997.
- Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 181–193, 2003.
- Jenkin, M. E., Young, J. C., and Rickard, A. R.: The MCM v3. 3. 1 degradation scheme for isoprene, *Atmos. Chem. Phys.*, 15, 11433–11459, 2015.
- Khan, M. A. H., Cooke, M. C., Utembe, S. R., Xiao, P., Morris, W. C., Derwent, R. G., Archibald, A. T., Jenkin, M. E., Percival, C. J., and Shallcross, D. E.: The global budgets of organic hydroperoxides for present and pre-industrial scenarios, *Atmos. Environ.*, 110, 65–74, 2015.

- Koppmann, R., Czapiewski, K. V., and Reid, J. S.: A review of biomass burning emissions, part I: gaseous emissions of carbon monoxide, methane, volatile organic compounds, and nitrogen containing compounds, *Atmos. Chem. Phys. Discuss.*, 5, 10455–10516, 2005.
- Kroll, J. H., and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593–3624, 2008.
- Lee, M., Heikes, B. G., Jacob, D. J., Sachse, G., and Anderson, B.: Hydrogen peroxide, organic hydroperoxide, and formaldehyde as primary pollutants from biomass burning, *J. Geophys. Res.*, 102, 1301–1309, 1997.
- Lee, M., Heikes, B. G., and Jacob, D. J.: Enhancements of hydroperoxides and formaldehyde in biomass burning impacted air and their effect on atmospheric oxidant cycles, *J. Geophys. Res.*, 103, 13201–13212, 1998.
- Lee, M., Heikes, B. G., and O'Sullivan, D. W.: Hydrogen peroxide and organic hydroperoxide in the troposphere: a review, *Atmos. Environ.*, 34, 3475–3494, 2000.
- Li, H., Chen, Z. M., Huang, L. B., and Huang, D.: Organic peroxides' gas-particle partitioning and rapid heterogeneous decomposition on secondary organic aerosol, *Atmos. Chem. Phys.*, 16, 1837–1848, 2016.
- Li, J. F., Song, Y., Mao, Y., Mao, Z. C., Wu, Y. S., Li, M. M., Huang, X., He, Q. C., and Hu, M.: Chemical characteristics and source apportionment of PM_{2.5} during the harvest season in eastern China's agricultural regions, *Atmos. Environ.*, 92, 442–448, 2014a.
- Li, L. Y., Xie, S. D., Zeng, L. M., Wu, R. R., and Li, J.: Characteristics of volatile organic compounds and their role in ground-level ozone formation in the Beijing-Tianjin-Hebei region, China, *Atmos. Environ.*, 113, 247–254, 2015.
- Li, M., Zhang, Q., Streets, D. G., He, K. B., Cheng, Y. F., Emmons, L. K., Huo, H., Kang, S. C., Lu, Z., Shao, M., Su, H., Yu, X., Zhang, Y.: Mapping Asian anthropogenic emissions of non-methane volatile organic compounds to multiple chemical mechanisms, *Atmos. Chem. Phys.*, 14, 5617–5638, 2014b.

- Li, X. H., Wang, S. X., Duan, L., Hao, J. M., Li, C., Chen, Y. S., and Yang, L.: Particulate and trace gas emissions from open burning of wheat straw and corn stover in China, *Environ. Sci. Technol.*, 41, 6052–6058, 2007.
- Li, X., Rohrer, F., Brauers, T., Hofzumahaus, A., Lu, K. D., Shao, M., Zhang, Y. H., and Wahner, A.: Modeling of HCHO and CHOCHO at a semi-rural site in southern China during the PRIDE-PRD2006 campaign, *Atmos. Chem. Phys.*, 14, 12291–12305, 2014c.
- Liang, H., Chen, Z. M., Wu, Q. Q., Huang, D., and Zhao, Y.: Do aerosols influence the diurnal variation of H₂O₂ in the atmosphere?, *AGU Fall Meeting Abstracts*, 2013a.
- Liang, H., Chen, Z. M., Huang, D., Zhao, Y., and Li, Z. Y.: Impacts of aerosols on the chemistry of atmospheric trace gases: a case study of peroxides and HO₂ radicals, *Atmos. Chem. Phys.*, 13, 11259–11276, 2013b.
- Liu, H. J.: Measurement of aerosol light scattering enhancement factor and study on hygroscopicity parameter, Ph. D, thesis, Peking University, China, 2015.
- Liu, Y., Yuan, B., Li, X., Shao, M., Lu, S. H., Li, Y., Chang, C. C., Wang, Z. B., Hu, W. W., Huang, X. F., He, L. Y., Zeng, L. M., Hu, M., and Zhu, T.: Impact of pollution controls in Beijing on atmospheric oxygenated volatile organic compounds (OVOCs) during the 2008 Olympic Games: observation and modeling implications, *Atmos. Chem. Phys.*, 15, 3045–3062, 2015.
- Liu, Y. H., Lu, K. D., Dong, H. B., Li, X., Cheng, P., Zou, Q., Wu, Y. S., Liu, X. G., and Zhang, Y. H.: In situ monitoring of atmospheric nitrous acid based on multi-pumping flow system and liquid waveguide capillary cell, *J. Environ. Sci.*, 43, 273–284, 2016.
- Lu, K. D., Rohrer, F., Holland, F., Fuchs, H., Bohn, B., Brauers, T., Chang, C. C., Haeseler, R., Hu, M., Kita, K., Kondo, Y., Li, X., Lou, S. R., Nehr, S., Shao, M., Zeng, L. M., Wahner, A., Zhang, Y. H., and Hofzumahaus, A.: Observation and modelling of OH and HO₂ concentrations in the Pearl River Delta 2006: a missing OH source in a VOC rich atmosphere, *Atmos. Chem. Phys.*, 12, 1541–1569, 2012.
- Madronich, S.: The Tropospheric visible Ultra-violet (TUV) model web page,

available at: <http://www.acd.ucar.edu/TUV>, 2002.

- Mao, J. Q., Jacob, D. J., Evans, M. J., Olson, J. R., Ren, X. R., Brune, W. H., St Clair, J. M., Crouse, J. D., Spencer, K. M., Beaver, M. R., Wennberg, P. O., Cubison, M. J., Jimenez, J. L., Fried, A., Weibring, P., Walega, J. G., Hall, S. R., Weinheimer, A. J., Cohen, R. C., Chen, G., Crawford, J. H., McNaughton, C., Clarke, A. D., Jaeglé L., Fisher, J. A., Yantosca, R. M., Le Sager, P., and Carouge, C.: Chemistry of hydrogen oxide radicals (HO_x) in the Arctic troposphere in spring, *Atmos. Chem. Phys.*, 10, 5823–5838, 2010.
- Mao, J. Q., Fan, S., Jacob, D. J., and Travis, K. R.: Radical loss in the atmosphere from Cu-Fe redox coupling in aerosols, *Atmos. Chem. Phys.*, 13, 509–519, 2013.
- Neeb, P., Sauer, F., Horie, O., and Moortgat, G. K.: Formation of hydroxymethyl hydroperoxide and formic acid in alkene ozonolysis in the presence of water vapour, *Atmos. Environ.*, 31, 1417–1423, 1997.
- Nguyen, T. B., Crouse, J. D., Teng, A. P., Clair, J. M. S., Paulot, F., Wolfe, G. M., and Wennberg, P. O.: Rapid deposition of oxidized biogenic compounds to a temperate forest, *Proc. Nat. Acad. Sci.*, 112, E392–E401, 2015.
- Pöschl, U., and Shiraiwa, M.: Multiphase chemistry at the atmosphere-biosphere interface influencing climate and public health in the anthropocene, *Chem. Rev.*, 115, 4440–4475, 2015.
- Pradhan, M., Kyriakou, G., Archibald, A. T., Papageorgiou, A. C., Kalberer, M., and Lambert, R. M.: Heterogeneous uptake of gaseous hydrogen peroxide by Gobi and Saharan dust aerosols: a potential missing sink for H_2O_2 in the troposphere, *Atmos. Chem. Phys.*, 10, 7127–7136, 2010.
- Ran, L., Zhao, C. S., Xu, W. Y., Lu, X. Q., Han, M., Lin, W. L., Yan, P., Xu, X. B., Deng, Z. Z., Ma, N., Liu, P. F., Yu, J., Liang, W. D., and Chen, L. L.: VOC reactivity and its effect on ozone production during the HaChi summer campaign, *Atmos. Chem. Phys.*, 11, 4657–4667, 2011.
- Reeves, C. E., and Penkett, S. A.: Measurements of peroxides and what they tell us, *Chem. Rev.*, 103, 5199–5218, 2003.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass

- burning emissions part II: intensive physical properties of biomass burning particles, *Atmos. Chem. Phys.*, 5, 799–825, 2005.
- Sander, S. P., Abbatt, J., Barker, J. R., Burkholder, J. B., Friedl, R. R., Golden, D. M., Huie, R. E., Kolb, C. E., Kurylo, M. J., Moortgat, G. K., Orkin, V. L., and Wine, P. H.: Chemical kinetics and photochemical data for use in atmospheric studies, Evaluation No, 17, JPL Publication 10–6, Jet Propulsion Laboratory, Pasadena, CA, USA, available at: <http://jpldataeval.jpl.nasa.gov>, 2011.
- Sarwar, G., Godowitch, J., Henderson, B. H., Fahey, K., Pouliot, G., Hutzell, W. T., Mathur, R., Kang, D., Goliff, W. S., and Stockwell, W. R.: A comparison of atmospheric composition using the Carbon Bond and Regional Atmospheric Chemistry Mechanisms, *Atmos. Chem. Phys.*, 13, 9695–9712, 2013.
- Sauer, F., Beck, J., Schuster, G., and Moortgat, G. K.: Hydrogen peroxide, organic peroxides and organic acids in a forested area during FIELDVOC'94, *Chemosphere*, 3, 309–326, 2001.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, 2003.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, John Wiley & Sons, 2006.
- Stein, A. F., and Saylor, R. D.: Sensitivities of sulfate aerosol formation and oxidation pathways on the chemical mechanism employed in simulations, *Atmos. Chem. Phys.*, 12, 8567–8574, 2012.
- Sullivan, A. P., Holden, A. S., Patterson, L. A., McMeeking, G. R., Kreidenweis, S. M., Malm, W. C., Hao, W. M., Wold, C. E., and Collett, J. L.: A method for smoke marker measurements and its potential application for determining the contribution of biomass burning from wildfires and prescribed fires to ambient PM_{2.5} organic carbon, *J. Geophys. Res.*, 113, D22302, 2008.
- Tan, Z. F., Fuchs, H., Lu, K. D., Bohn, B., Broch, S., Haeseler, R., Hofzumahaus, A., Holland, F., Li, X., Liu, Y., Rohrer, F., Shao, M., Wang, B. L., Wang, M., Wu, Y. S.,

- Zeng, L. M., Wahner, A. and Zhang, Y. H.: Observation and modelling of the OH, HO₂ and RO₂ radicals at a rural site (Wangdu) in the North China Plain in summer 2014, *Geophysical Research Abstracts*, pp. EGU2016-5459, 2016.
- Tao, M. H., Chen, L. F., Su, L., and Tao, J. H.: Satellite observation of regional haze pollution over the North China Plain, *J. Geophys. Res.*, 117, D12203, 2012.
- Wang, H. L., Huang, D., Zhang, X., Zhao, Y., and Chen, Z. M.: Understanding the aqueous phase ozonolysis of isoprene: distinct product distribution and mechanism from the gas phase reaction, *Atmos. Chem. Phys.*, 12, 7187–7198, 2012.
- Wang, L. L., Xin, J. Y., Li, X. R., and Wang, Y. S.: The variability of biomass burning and its influence on regional aerosol properties during the wheat harvest season in North China, *Atmos. Res.*, 157, 153–163, 2015.
- Wang, M., Zeng, L. M., Lu, S. H., Shao, M., Liu, X. L., Yu, X. N., Chen, W. T., Yuan, B., Zhang, Q., Hu, M., and Zhang, Z. Y.: Development and validation of a cryogen-free automatic gas chromatograph system (GC-MS/FID) for online measurements of volatile organic compounds, *Anal. Methods*, 6, 9424–9434, 2014.
- Winiberg, F. A. F., Dillon, T. J., Orr, S. C., Groß, C. B., Bejan, I., Brumby, C. A., Evans, M. J., Smith, S. C., Heard, D. E., and Seakins, P. W.: Direct measurements of OH and other product yields from the HO₂ + CH₃C(O)O₂ reaction, *Atmos. Chem. Phys.*, 16, 4023–4042, 2016.
- Wu, Q. Q., Huang, L. B., Liang, H., Zhao, Y., Huang, D., and Chen, Z. M.: Heterogeneous reaction of peroxyacetic acid and hydrogen peroxide on ambient aerosol particles under dry and humid conditions: kinetics, mechanism and implications, *Atmos. Chem. Phys.*, 15, 6851–6866, 2015.
- Wu, R. R., Bo, Y., Li, J., Li, L. Y., Li, Y. Q., and Xie, S. D.: Method to establish the emission inventory of anthropogenic volatile organic compounds in China and its application in the period 2008–2012, *Atmos. Environ.*, 127, 244–254, 2016.
- Ye, N. N.: Observations and budget analysis of ambient nitrous acid (HONO) in Wangdu, a rural site in North China Plain, Master thesis, Peking University, China, 2015.
- Yokelson, R. J., Crounse, J. D., DeCarlo, P. F., Karl, T., Urbanski, S., Atlas, E.,

- Campos, T., Shinozuka, Y., Kapustin, V., Clarke, A. D., Weinheimer, A., Knapp, D. J., Montzka, D. D., Holloway, J., Weibring, P., Flocke, F., Zheng, W., Toohey, D., Wennberg, P. O., Wiedinmyer, C., Mauldin, L., Fried, A., Richter, D., Walega, J., Jimenez, J. L., Adachi, K., Buseck, P. R., Hall, S. R., and Shetter, R.: Emissions from biomass burning in the Yucatan, *Atmos. Chem. Phys.*, 9, 5785–5812, 2009.
- Yuan, Z. B., Lau, A. K. H., Shao, M., Louie, P. K. K., Liu, S. C., and Zhu, T.: Source analysis of volatile organic compounds by positive matrix factorization in urban and rural environments in Beijing, *J. Geophys. Res.*, 114, D00G15, 2009.
- Zhang, L., Brook, J. R., and Vet, R.: A revised parameterization for gaseous dry deposition in air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, 2003.
- Zhang, R. Y., Wang, G. H., Guo, S., Zamora, M. L., Ying, Q., Lin, Y., Wang, W. G., Hu, M., and Wang, Y.: Formation of urban fine particulate matter, *Chem. Rev.*, 115, 3803–3855, 2015.
- Zhang, X., Chen, Z. M., He, S. Z., Hua, W., Zhao, Y., and Li, J. L.: Peroxyacetic acid in urban and rural atmosphere: concentration, feedback on PAN–NO_x cycle and implication on radical chemistry, *Atmos. Chem. Phys.*, 10, 737–748, 2010.
- Zhang, X., He, S. Z., Chen, Z. M., Zhao, Y., and Hua, W.: Methyl hydroperoxide (CH₃OOH) in urban, suburban and rural atmosphere: ambient concentration, budget, and contribution to the atmospheric oxidizing capacity, *Atmos. Chem. Phys.*, 12, 8951–8962, 2012.
- Zhang, Y., Huang, W., Cai, T. Q., Fang, D. Q., Wang, Y. Q., Song, J., Hu, M. and Zhang, Y. X.: Concentrations and chemical compositions of fine particles (PM_{2.5}) during haze and non-haze days in Beijing, *Atmos. Res.*, 174, 62–69, 2016.
- Zhao, R., Lee, A. K. Y., Soong, R., Simpson, A. J., and Abbatt, J. P. D.: Formation of aqueous-phase α -hydroxyhydroperoxides (α -HHP): potential atmospheric impacts, *Atmos. Chem. Phys.*, 13, 5857–5872, 2013a.
- Zhao, Y., Chen, Z. M., Shen, X. L., and Huang, D.: Heterogeneous reactions of gaseous hydrogen peroxide on pristine and acidic gas-processed calcium carbonate particles: Effects of relative humidity and surface coverage of coating, *Atmos. Environ.*, 67, 63–72, 2013b.

- Zheng, G. J., Duan, F. K., Ma, Y. L., Zhang, Q., Huang, T., Kimoto, T. K., Cheng, Y. F., Su, H. and He, K. B.: Episode-based evolution pattern analysis of haze pollution: method development and results from Beijing, China, *Environ. Sci. Technol.*, 50, 4632–4641, 2016.
- Zheng, G. J., Duan, F. K., Su, H., Ma, Y. L., Cheng, Y., Zheng, B., Zhang, Q., Huang, T., Kimoto, T., Chang, D., Poschl, U., Cheng, Y. F. and He, K. B.: Exploring the severe winter haze in Beijing: the impact of synoptic weather, regional transport and heterogeneous reactions, *Atmos. Chem. Phys.*, 15, 2969–2983, 2015.
- Ziemann, P. J., and Atkinson, R.: Kinetics, products, and mechanisms of secondary organic aerosol formation, *Chem. Soc. Rev.*, 41, 6582–6605, 2012.

Table 1. Summary of the concentrations of atmospheric peroxides during Wangdu Campaign 2014.

		H ₂ O ₂ (ppbv)	MHP (ppbv)	PAA (ppbv)
	D.L. ^a	0.01	0.01	0.01
24 h	N ^b	1797	1797	1797
	Mean	0.51	0.16	0.03
	S.D. ^c	0.90	0.21	0.11
	Median	0.19	0.11	0.01
	Maximum	11.3	1.25	1.49
		N ^b	829	829
Daytime (06:00–18:00 LT ^d)	Mean	0.55	0.16	0.03
	S.D. ^c	0.83	0.18	0.12
	Median	0.24	0.12	0.01
	Maximum	10.2	1.20	1.49
		N ^b	968	968
Nighttime (18:00–06:00 LT ^d)	Mean	0.48	0.17	0.04
	S.D. ^c	0.96	0.23	0.11
	Median	0.15	0.11	0.01
	Maximum	11.3	1.25	1.47

^a D.L.: detection limit.

^b N: number of samples.

^c S.D.: standard deviation.

^d LT: local time.

Table 2. Chemical mechanisms for CH₃C(O)O₂ and CH₃O₂ related chemistry modified or added to MCMv3.3.1.

Reactions	Rate constants (cm ³ molecule ⁻¹ s ⁻¹)	Reference
CH₃C(O)O₂ chemistry		
CH ₃ C(O)O ₂ + HO ₂ → CH ₃ C(O)OOH + O ₂	2.40 × 10 ⁻¹¹ × 0.37	Winiberg et al. (2016)
CH ₃ C(O)O ₂ + HO ₂ → CH ₃ C(O)OH + O ₃	2.40 × 10 ⁻¹¹ × 0.12	Winiberg et al. (2016)
CH ₃ C(O)O ₂ + HO ₂ → CH ₃ + CO ₂ + OH + O ₂	2.40 × 10 ⁻¹¹ × 0.51	Winiberg et al. (2016)
CH₃O₂ chemistry		
CH ₃ O ₂ + OH → PRODUCT	2.80 × 10 ⁻¹⁰	Fittschen et al. (2014)

Table 3. Linear regression of atmospheric peroxide species to CO and K⁺ for three biomass burning events during the Phase II (15 June–17 June). Correlation coefficients shown in italic and bold indicate statistical significance ($p < 0.05$) and higher statistical significance ($p < 0.01$), respectively.

Species	Slope ^a	Correlation coefficient		N ^b	Critical correlation coefficient
		CO	K ⁺		
<i>Event I</i>					
H ₂ O ₂	2.17×10^{-3}	0.8144	0.8432	10	0.7646 ($p < 0.01$), 0.6319 ($p < 0.05$)
MHP	1.23×10^{-3}	<i>0.6873</i>	<i>0.7624</i>	10	
PAA	7.16×10^{-4}	0.8378	0.9515	10	
<i>Event II</i>					
H ₂ O ₂	N/A ^c	N/A ^c	0.9394	12	0.7079 ($p < 0.01$), 0.5760 ($p < 0.05$)
MHP	N/A ^c	N/A ^c	0.9491	12	
PAA	N/A ^c	N/A ^c	0.9449	12	
<i>Event III</i>					
H ₂ O ₂	N/A ^c	N/A ^c	0.9632	9	0.7977 ($p < 0.01$), 0.6664 ($p < 0.05$)
MHP	N/A ^c	N/A ^c	0.8741	9	
PAA	N/A ^c	N/A ^c	0.8436	9	

^a Slope: enhancement ratio of speciated peroxides relative to CO.

^b N: number of samples.

^c N/A: missing data.




Figure 1. Temporal profile for atmospheric peroxides over the entire Wangdu Campaign 2014.




Figure 2. Time series of meteorological parameters, chemical species and atmospheric peroxides. The orange shaded area represents the Phase I (8 June–11 June), Phase II (15 June–17 June) and Phase III (29 June–3 July). The grey shaded area indicates three biomass burning events.

删除的内容: shade

删除的内容: shade




Figure 3. Observed and modelled concentrations of atmospheric peroxides for Phase I (8 June–11 June), Phase II (15 June–17 June) and Phase III (29 June–3 July). The green circles represent observed concentrations. The blue, orange and red lines indicate the modelled concentrations from three different scenarios: base case, new gas-phase reaction case and heterogeneous uptake case, respectively. The grey shaded area indicates three biomass burning events.

删除的内容: shade




Figure 4. Comparisons between measured and predicted concentrations of atmospheric peroxides in the model base case for daytime and nighttime during the Phase I (8 June–11 June). The solid lines represent the 1:1 ratio of observed to modelled values.




Figure 5. Sensitivity of production rate of atmospheric peroxides to major VOC precursor groups and individual VOC species for Phase I and Phase III.




Figure 6. Contributions of each sink to H₂O₂, MHP and PAA destruction in the box model with the heterogeneous uptake by aerosols added during Phase I and Phase III.