

Temporal and spectral cloud screening of polar-winter aerosol optical depth (AOD): impact of homogeneous and inhomogeneous clouds and crystal layers on climatological-scale AODs

Norman T. O'Neill¹, K. Baibakov¹, S. Hesaraki¹, L. Ivanescu¹, R. V. Martin², C. Perro², J. P. Chaubey¹, A. Herber³, T. J. Duck²

¹ Centre d'Applications et de Recherches en Télédétection, Université de Sherbrooke, Sherbrooke, Canada

² Dept. of Physics and Atmospheric Science, Dalhousie University, Halifax, Canada

³ Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

Correspondence to: Jai Prakash Chaubey (jaiprakash.spl@gmail.com)

Abstract

We compared star photometry-derived, polar winter aerosol optical depths (AODs), acquired at Eureka, Nunavut, Canada and Ny-Ålesund, Svalbard with GEOS-Chem (GC) simulations as well as ground-based lidar and CALIOP retrievals over a sampling period of two polar winters. The results indicate significant cloud and/or low-altitude ice crystal (LIC) contamination which is only partially corrected using temporal cloud screening. Spatially homogeneous clouds and LICs that remain after temporal cloud screening represent an inevitable systematic error in the estimation of AOD: this error was estimated to vary from 78% to 210% at Eureka and from 2% to 157% at Ny-Ålesund. Lidar analysis indicated that LICs appeared to have a disproportionately large influence on the homogenous coarse mode optical depths that escape temporal cloud screening. In principle, spectral cloud screening (to yield fine mode or sub-micron AODs) reduces pre-cloud-screened AODs to the aerosol contribution if one assumes that coarse mode (super-micron) aerosols are a minor part of the AOD. Large, low frequency, differences between these retrieved values and their GC analogue appeared to be often linked to strong, spatially extensive planetary boundary layer events whose presence at either site was inferred from CALIOP profiles. These events were either not captured or significantly underestimated, by the GC simulations. High frequency AOD variations of GC fine mode aerosols at Ny-Ålesund were attributed to sea-salt (SS) while low frequency GC variations at Eureka and Ny-Ålesund were attributable to sulfates. CALIOP profiles and AODs were invaluable as spatial and temporal redundancy support (or, alternatively, as insightful points of contention) for star photometry retrievals and GC estimates of AOD.

1 Introduction

The importance of understanding aerosol mechanisms driving the direct and indirect effects is of particular significance over the Arctic where climate change impacts are known to be amplified (IPCC, 2013). This is very important during the polar winter when aerosol variability, generally associated with the Arctic haze phenomenon, is typically stronger than during the polar summer (see Di Pierro et al., 2013 for example) and when the number and nature of Arctic haze aerosols can have significant (indirect) effects on thin ice cloud properties and their radiative forcing budget (c.f. Garrett & Zhao, 2006 and Blanchard et al., 1994 respectively).

In order to properly evaluate aerosol processes and emission representation in chemical transport models one needs to develop a reliable and varied measurement system to exercise as many of the aerosol functionalities as possible. Ground and satellite based remote sensing (RS) measurements are arguably the key components of such a measuring system since they provide the front-line, robust parameters that define the first order comparative constraints that models must necessarily satisfy. There are currently only a few instances of aerosol RS measurements during the polar winter: (a) satellite-based, polar orbit, lidar profiles and their derived aerosol optical depths (AODs) (b) ground-based lidar profiles and derived AODs as well as star photometer (and some moonphotometer) AOD measurements acquired at a few Arctic sites.

Star photometry is currently the defacto reference for all polar winter AOD measurements since it is a direct extinction measurement¹. In the same way that RS parameters should be front line model comparison parameters, an AOD climatology (or, at least, a multi-year statistical analysis) should be a necessary basis of comparison in parallel to more spatially and temporally demanding (meteorological scale) evaluations. The AOD contamination impact of clouds and other sources of starphotometry error as well as the AOD computation impact of model limitations such as spatial resolution and time-step resolution are often dampened by carrying out comparisons at climatological scales.

In the Arctic, the process of cloud-screening raw star photometry AODs (of rejecting raw AODs, deemed to be cloud contaminated) is critical, given the relative weakness of AOD amplitudes as well as the occurrence of cloud and low-altitude ice crystal (LIC) events during the polar winter. Lesin's et al. (2009), studied LIC events at Eureka during 2006 and observed that 19.1% of lidar events were due to clear-night or cloudy-night LICs at an average altitude of 450 ± 100 m (average of the Dec., Jan., Feb., March period of 2006). Cloud-screening may be temporal in nature (rejection is based on sufficiently rapid changes in optical depth where the assumption is that only clouds go through high frequency changes in optical depth) or of a spectral nature (rejection is based on the assumption that only cloud optical depths are spectrally neutral). The former approach suffers from errors of commission and omission (elimination of high frequency aerosol data and the inability to identify homogeneous cloud events respectively) while the latter approach may, for example, exclude super-micron aerosols (i.e. in addition to the cloud events which it is expected to exclude). If relevant comparisons are to be made with models then proper cloud screening is critical.

For our purposes, the current role of lidars in such climatologies or multi-year analyses is more of a supportive nature: ground-based lidars provide fundamental supporting data for AOD measurements in terms of the interpretation of the vertical contributions to the AOD (as well as the vertical contributions of cloud contamination) and the correlative coherence of their estimated AODs (Baibakov et al., 2015) while a satellite-based lidar provides critical interpretative information on the horizontal extent of these contributions and, their integrated AOD estimate.

High Arctic, near sea-level, star photometers at the AWI (Alfred Wegner Institute) base in Ny-Ålesund, Svalbard (79°N, 12°E) and the PEARL (Polar Environmental Atmospheric Research Laboratory) site at Eureka, Nunavut, Canada (80°N, 86°W) were employed to acquire a common, 2-year ensemble of polar winter AODs (Baibakov, 2014; Ivanescu et al., 2014). The simulated polar winter AODs of the GEOS-Chem (GC) model were compared with the star photometer AODs in order to quantitatively evaluate the relative temporal agreement of the star photometer and model over the 2-year reference period. Pan-Arctic AOD map products from the CALIOP lidar aboard the CALIPSO polar orbiting satellite (Winker et al., 2013) were also used in this study. AOD animations for all daily orbit lines were compared with daily GC AOD maps to achieve a qualitative measure of the relative spatiotemporal agreement between the model and CALIOP animations and to better understand the extent of major AOD events during the polar winter.

2 Methodological considerations

In the text that follows we discuss specific issues related to the AODs derived from the measurements and model simulations. The symbol and acronym glossary allows for a centralized reference concerning the different types of AODs (whether measured or simulated) and other key parameters. As part of this study, we processed individual AODs and analyzed daily averaged and monthly averaged AODs.

2.1 Star photometer measurements

2.1.1 AODs generated by the star photometer

A brief description of the star photometer along with retrieval, calibration and logistical issues related to star photometer measurements is given in Baibakov et al. (2015). In that paper, we carried out an event level analysis of synchronized star photometer and Raman lidar measurements for a sampling of the data set employed in the present analysis. That communication was the first paper in which we reported on the optical coherency of passive / active, polar winter measurements subdivided into total, fine, coarse (optical) modes. It confirmed the relevance of extracting total, fine and coarse mode AODs (τ_a , τ_f and τ_c at a reference wavelength of 500 nm for the passive measurements and 532 nm for the active measurements) and motivated us to create a preliminary (testbed) AOD climatology which

¹ as opposed to the backscatter measurements provided by elastic and inelastic lidars which require, respectively, a knowledge of the transfer ratio from backscattering to extinction and an evaluation of the attenuation of the molecular signal.

99 could be compared with AODs derived from GC simulations and CALIOP extinction profiles (see the symbol and
acronym glossary for more details).

102 2.1.2 Spectral and temporal cloud-screening

105 As in Baibakov et al. (2015), raw AOD spectra were processed through the SDA (Spectral Deconvolution algorithm)
to yield estimates of τ_a , τ_f and τ_c . The τ_f component is of particular relevance because it represents the contribution of
aerosols that remain after the removal of the contribution of coarse mode clouds, coarse mode LICs and coarse mode
108 aerosols. This is what we call spectral cloud-screening: if the coarse mode aerosol contribution to τ_c is relatively small
(and this is supported, for example by GC (aerosol) ratios of $\tau_{c, GC} / \tau_{a, GC}$ being $< \sim 10\%$ for the two stations across
our climatological period) then one can argue that τ_f is representative of aerosols in the Arctic and that τ_c is
predominantly due to cloud or LIC contamination.

111 Baibakov et al. (2015) employed star photometry and lidar data to illustrate the utility of spectral cloud screening in
the presence of temporally and spatially inhomogeneous clouds (their Fig. 8) as well as the effectiveness of both
114 temporal and spectral cloud screening in the presence of inhomogeneous LICs embedded in what appeared to be a
background environment of more homogeneously distributed LICs (their Figure 9). They noted that the two cloud-
screening approaches gave similar results in the presence of relatively inhomogeneous LICs while indicating that the
117 remaining difference was arguably due to temporally (spatially) homogenous coarse mode particles (which, given the
argument above, would be predominantly due to homogeneous LIC layers or homogeneous clouds).

120 If one divides the raw AOD data ensemble (and their derived SDA component ensembles) into temporally cloud-
screened (accepted) and rejected raw AODs ensembles ("cs" and "rej"), then, for daily means ($x = a, f, \text{ or } c$) one can
show that;

$$123 \quad \tau_x = \gamma \tau_{x, cs} + (1 - \gamma) \tau_{x, rej} \quad (1)$$

126 with $\gamma = N_{cs} / (N_{cs} + N_{rej})$ and where N_{cs} and N_{rej} are the number of AODs in the cloud-screened and rejected
ensembles. Equation (1) one can be re-arranged to yield a sum of homogeneous and inhomogeneous components;

$$129 \quad \tau_x = \tau_{x, hom} + \tau_{x, inh} \quad (2)$$

132 where $\tau_{x, cs}$ has been renamed $\tau_{x, hom}$ in order to achieve a more intuitive vocabulary and where the inhomogeneous
component (the perturbation above the low frequency, cloud-screened, homogeneous component) is,

$$132 \quad \tau_{x, inh} = (1 - \gamma) [\tau_{x, rej} - \tau_{x, hom}] \quad (3)$$

135 The algebraic manipulation used to isolate and label $\tau_{x, hom}$ in equation (2) (and, as a consequence, $\tau_{x, inh}$) is not
a subjective choice². The daily average of all optical depths that are not rejected ($\tau_{x, hom}$) is the daily average that
would be reported as the result of temporal cloud screening : it is precisely this quantity that should be evaluated in
138 terms of the effectiveness of temporal cloud screening.

141 The fine mode AOD can be considered approximately homogeneous ($\tau_f \cong \tau_{f, hom}$; this is largely the basis of
temporal cloud-screening). Appealing to equation (2) and the propagation of $\tau_a = \tau_f + \tau_c$ across averages applied
to any of the data ensembles (see the acronym and symbol glossary for a brief overview of that propagation), $\tau_{f, inh} \cong$
0 and thus $\tau_{a, inh} \cong \tau_{c, inh}$. For $x = a$, equation (2) can then be expanded ;

$$144 \quad \tau_a = \tau_{a, hom} + \tau_{a, inh} \quad (4)$$

² Subjective in the sense that one could have erroneously argued, for example, that the first term of equation (1)
should have been labelled as $\tau_{x, hom}$

$$\begin{aligned}
&= \tau_{f, hom} + \tau_{c, hom} + \tau_{f, inh} + \tau_{c, inh} \\
147 \quad &\cong \tau_{f, hom} + \tau_{c, hom} + \tau_{c, inh}
\end{aligned} \tag{5}$$

150 Equation (5) approximately represents the components of the total AOD while, in comparison with equation (4), reminds us that the cloud-screened AOD ($\tau_{a, hom}$) is divided into homogeneous components ($\tau_{f, hom}$ and $\tau_{c, hom}$) and that $\tau_{a, inh} \cong \tau_{c, inh}$. Equations (2), (4) and (5) propagate into monthly averages (maintain the same form).

153 2.2 GEOS-Chem simulations

The model that we employed for our comparisons was the GEOS-Chem global chemical transport model (GC) version 9-# (<http://acmg.seas.harvard.edu/geos/>). It is driven by GEOS-5 assimilated meteorological fields from the NASA Goddard Modeling and Assimilation Office (GMAO). The GC simulation has a 15 minute time step for transport and a 60 minute time step for chemistry and emissions. The lat / log grid size over the Arctic was 2° by 2.5° (approximately 220 km x 50 km respectively at the high Arctic latitudes of Eureka and Ny-Ålesund) with 47 vertical levels up to 0.01 hPa.

162 An overview of the aerosol physics and chemistry in GC is given in Park et al., (2004). We divided GC AODs into their fine and coarse mode components ($\tau_{f, GC}$ and $\tau_{c, GC}$) using the species by species segregation provided by GC (fine mode organic carbon, sulfate and black carbon along with fine and coarse mode sea-salt (SS) and mineral dust). The GC aerosol simulation includes the sulfate-nitrate-ammonium system (Park et al., 2004; Pye et al., 2009), primary (Park et al., 2003) and secondary (Henze et al., 2006; Henze et al., 2008; Liao et al., 2007; Fu et al., 2008) organics, mineral dust (Fairlie et al., 2007), and sea salt (Jaegle et al., 2011). AOD is calculated at 550 nm using RH-dependent aerosol optical properties (see Martin et al., 2003 for an overview of the optical processing employed for GC aerosols).

168 2.3 AODs generated from CALIOP profiles

171 The CALIOP processing algorithm generates attenuated backscatter coefficient profiles and, after the application of an aerosol classification algorithm, estimates of tropospheric AOD along a given CALIPSO orbit line. A discussion of CALIOP extinction coefficient and AOD retrievals and their sources of variability within an Arctic night context can be found in Di Pierro et al. (2013). The AODs are, even in the significantly more optimal environment of nighttime conditions, very sensitive to the vagaries of aerosol vs cloud classification in conditions of weak backscatter return typical of the relatively low concentrations of Arctic aerosols under or mixed with thin clouds or LICs, etc.. Di Pierro et al. (2013) suggest, for example, that sub 2-km "diamond dust" may have been misclassified as aerosols and thus may have been responsible for very high values of aerosol extinction coefficient (and thus of AOD) from CALIOP retrievals (in 5% of the multi-year, December to February, Arctic-scale cases that they sampled).

180 With these considerations in mind, we employed CALIOP profiles and CALIOP AOD animations to gain insights into the spatio-temporal dynamics of aerosol events which might have influenced measurements at Eureka and Ny-Ålesund. We also employed averages of near-Eureka and near-Ny-Ålesund CALIOP AODs (i.e. spatial averages of all CALIOP AODs within a specified radial distance from Eureka and Ny Alesund) as an auxiliary AOD context in our temporal comparisons of GC AODs with star photometer AODs at Eureka and Ny-Ålesund. We chose 500 km as the radius of the near-site CALIOP averages since this case generally displayed the least amount of day to day variance in comparison with smaller radii choices (reduction in standard deviation of about a factor of 3 when increasing the radius from 100 to 500 km). The AODs were retrieved from the CALIOP "Column_Optical_Depth_Aerosols_532" product associated with the "5km Aerosol Profile".

189 2.3.1 Impact of differences in wavelength

192 For reasons of historical consistency we chose to retain the standard output wavelength that we employ for starphotometry retrievals (500 nm), the 532 nm lidar wavelength of CALIOP and the 550 nm GC standard. As an indicator of the impact of these wavelength differences (for the case of the fine mode where the decrease from 500 to 532 to 550 nm would be at its largest), we performed a 2009 to 2011 survey of τ_f values for 5 Arctic AERONET stations. The results indicated that the global 550 nm average was less than 0.01 below the global 500 nm average.

3 Results

198 3.1 GC and CALIOP spatial comparisons

201 Spatial comparisons between CALIOP and GC AODs were spotty at best. CALIOP sampling represents a rather
extreme statistical challenge with generally modest signal to noise for the weak aerosol optical properties typical of
the Arctic and strong cloud / LIC layer interference coupled with a highly irregular, spatial sampling grid. In spite of
204 these limitations we frequently observed strong, spatially expansive, PBL³ backscatter structures of low DR⁴ that
were characterized as aerosol layers by the CALIOP processing algorithm. These structures were often not captured
by GC in the sense that the simulated AOD amplitude was typically much smaller than the computed CALIOP AODs.
207 Strong GC AOD events, on the other hand, are often unsupported by any CALIOP evidence simply because the
atmosphere in the region of interest is cloud dominated (although there can be relatively small, tantalizing windows
of cloudless sky which suggest a, difficult to substantiate, spatial correlation between the model and the
measurements).

210 3.2 Climatological-scale analysis of star photometer AODs

Daily statistics

213 Figures 1 and 2 show star photometer and GC AOD comparisons for, respectively, daily averages at Eureka and Ny-
Ålesund during the polar winters of 2010/2011 and 2011/2012. Each graph includes estimates of non cloud-screened
216 (raw) AODs (τ_a in grey), cloud-screened AODs ($\tau_{a, hom}$ in black), fine mode AOD (τ_f in light red), filtered fine
mode AODs (τ_f^* in dark red) and GC-estimated fine mode AODs ($\tau_{f, GC}$ dark red dashes). τ_f^* represents our best
attempt at producing climatological-scale AODs: to ensure the survival of only the most robust estimates of τ_f , we
219 allow ourselves the luxury of eliminating \mathcal{G} values for which $\mathcal{G} / \tau_a < 0.3$ (for which the risk of errors due to residual
cloud contamination is greatest).

222 The most striking feature of these curves, in particular for Eureka, is the notable variation in the AODs, before and
after temporal or spectral cloud screening (we leave the detailed discussion of these notable variations to the section
below on temporal and spectral cloud-screening). The cloud screening (in particular the τ_f^* spectral cloud-screening)
225 tends to reduce magnitudes towards the $\tau_{f, GC}$ values. We have confidence in the τ_f^* estimations based on our lidar
/ star photometer event level comparisons of Baibakov et al. (2015) and based on our detailed analysis of the diurnal
variation of individual τ_f retrievals: in general the τ_f^* values in Figures 1 and 2 that were significantly higher than
228 the $\tau_{f, GC}$ values were associated with robust and diurnally smooth variations of individual retrievals (see Fig. S1 of
the supporting information for starphotometer illustrations of robust and moderately robust fine mode events).

231 High frequency variations of $\tau_{f, GC}$ for Ny-Ålesund (in particular the late winter variations of 2012 seen in Figure 2d)
are predominantly due to fine mode SS aerosols associated with the yearly winter depression and strong winds
southeast of Greenland (see, for example. Ma et al., 2008). It is noteworthy that virtually all large-amplitude, high
234 frequency variation of $\tau_{f, GC}$ at Ny-Ålesund is due to SS: outside of these peaks the dominant species is generally
sulfate (an affirmation based on a component by component analysis of $\tau_{f, GC}$ values). It is difficult if not impossible
to demonstrate any degree of correlative agreement between the sparse τ_f^* points and the high frequency $\tau_{f, GC}$
237 spikes. Fig. S2 of the supporting information shows an example of apparent coherence between GC AODs (dominated
by fine-mode, SS aerosols) and CALIOP AODs (the largest $\tau_{f, GC}$ peak of Fig 2d corresponds to the same day as this
illustration). However such examples were frustratingly rare given the frequent appearance of strong SS plumes in
240 GC imagery (of which Fig. S2 is one of many examples): this is no doubt partly due to cloud-contamination of
CALIOP profiles but it conceivably might also be GC overestimates of SS AODs. Attempts to relate $\tau_{f, GC-SS}$ to

³ Planetary boundary layer

⁴ CALIOP depolarization ratio (see the symbol and acronym glossary for details)

243 NaCl mass concentration measurements⁵ acquired at the Ny-Ålesund, Zeppelin observatory (475 m.a.s.l.) were
inconclusive in the sense of achieving any kind of significant correlation (and we note that no better correlation was
achieved if GC mass concentrations at the Zeppelin elevation were employed instead of $\tau_{f, GC-SS}$).

246 A notable Ny-Ålesund star photometry feature was what appeared to be a continuity of strong τ_f^* values from the
last week in November, 2011 to the first week in January, 2012 where τ_f^* was ~ 3 times the $\tau_{f, GC}$ values (Figure
2c). We believe that this difference is real because of the robustness of individual τ_f variations mentioned above and
249 because the CALIOP vertical profiles of this period were often dominated by strong PBL events of low DR. These
vertical profiles were associated with spatially broad and robust $\tau_{a, CALIOP}$ features that were either not captured or
significantly underestimated by the GC simulations (see two examples of these PBL events in Fig. S3 and Fig. S4 of
252 the supporting information). The predominance of PBL aerosol events during the polar winter was, in particular, noted
by Di Pierro et al. (2013) as part of their 6 year Arctic climatology using CALIOP profiles. A sampling of the CALIOP
and GC vertical profiles for the event of Fig. S3 showed that GC appeared to capture the general vertical form of the
255 PBL feature but with $\tau_{a, GC}$ (largely sulphate dominated) values that were much weaker than the $\tau_{a, CALIOP}$ values.
In this context of negatively biased $\tau_{a, GC}$ values, Di Pierro (2013) also found a negative, fine mode, polar winter GC
258 bias and suggested that an important fine mode component during the polar winter (and currently not included in GC)
is dry SS particles that result from the sublimation of crystals from wind blown snow events.

A prominent Eureka event was the largest τ_f^* value of Fig. 1a (Mar. 1, 2011). This corresponded to a strong value
261 of $\tau_{a, CALIOP}$ and what appeared to be a spatially broad, PBL CALIOP event of low DR whose spatial continuity was
inferred to be frequently hidden by higher altitude clouds. A second notable Eureka event was the largest τ_f^* value of
Fig. 1b (Mar. 29, 2012). This was a very stable fine mode event ($\tau_f \gg \tau_c$ with low frequency diurnal variation typical
264 of aerosol events) which, however, only lasted for about 2½ hours (a duration which, at this late date of Mar. 29, is
the result of the star photometer's inability to track stars in the presence of competitive or dominant, sunlight-induced
background radiance). CALIOP data did not support this strong value but the $\tau_{a, CALIOP}$ maps were very spotty with
267 strong cloud contamination in the vertical profiles (and Eureka overpasses were all daylight overpasses so that the S/N
advantages of the polar winter were largely lost at this late date).

270 *Temporal and spectral cloud screening*

Fig. 3a shows monthly averaged starphotometer AODs ($\langle \tau_a \rangle$) partitioned into grey and black, $\langle \tau_{a, inh} \rangle$ and $\langle \tau_{a, hom} \rangle$
273 components (in support of equation (4)) as well as $\langle \tau_a \rangle$ partitioned into $\langle \tau_{f, hom} \rangle$, $\langle \tau_{c, hom} \rangle$
, and $\langle \tau_{c, inh} \rangle$ components (in support of equation (5)⁶). The need for temporal cloud screening (the significant
276 amplitude of $\langle \tau_{a, inh} \rangle$) relative to ($\langle \tau_{a, hom} \rangle$) is evident (especially for Eureka). It is also evident that a
significant fraction of homogeneous coarse mode values have circumvented the temporal cloud screening process
dark blue ($\langle \tau_{c, hom} \rangle$ values have been accepted as legitimate AODs). This (the unavoidable failure to reject raw
AODs associated with homogeneous clouds or LICs) is a cloud / LIC detection error of the temporal cloud screening
279 process (given, as indicated above, the GC-driven assumption that coarse mode aerosols are a small fraction of the
AOD in the Arctic). An estimate of the relative (%) error, due to this error of omission is $\langle \tau_{c, hom} \rangle / \langle \tau_{f, hom} \rangle$
: this yields values that range from 78% to 210% for Eureka and from 2% to 157% for Ny-Ålesund.

282 In order to better understand the large temporal cloud screening errors of the Eureka starphotometry data, we
performed an analogous partitioning of lidar-derived coarse mode optical depths (τ_c') into inhomogeneous and
285 homogeneous components above and below a nominal LIC upper limit ($h_{LIC} = 600$ m using the statistical results of
Lesins et al., 2013). The details of the partitioning process are given in Appendix A. The results, shown in Figure 3b,
are colour coded to match the inhomogeneous / homogeneous colour coding of the Figure 3a starphotometry results
288 as well as being sub-divided into segments above and below h_{LIC} . The correspondence in terms of inhomogeneous
and homogeneous partitioning is reasonable given the differences in sampling strategies of the two instruments as
well as specific instrumental idiosyncracies such as the overlap function associated with the lidar data (see Appendix

⁵ <http://ebas.nilu.no/default.aspx>, link provided by Ove Hermansen, 2015

⁶ The fact that the two columns don't have the same height is a reflection of the approximate nature of equation (5) (that $\langle \tau_{f, inh} \rangle$ is not negligible)

291 A for more details). What is of particular interest is that the homogeneous contribution within the presumed LIC layer
averages ~ 50% of the homogeneous total : a disproportionate amount in terms of vertical distance in the atmosphere
294 (i.e. LICs appear to have an important influence on the homogenous coarse mode optical depths that escape temporal
cloud screening). At the same time we note the expected result that the inhomogeneous component is dominated by
contributions above h_{LIC} .

297 *Monthly statistics*

300 Fig. 4a shows month to month variations, along with standard deviations of $\langle \tau_{a, CALIOP} \rangle$, $\langle \tau_f \rangle$, $\langle \tau_f^* \rangle$ and, for the
specific case of Ny-Ålesund, the monthly, 9-year star photometry climatology of Herber et al. 2002 ($\langle \tau_{a, Herber} \rangle$). The
variability (standard deviation) of $\langle \tau_{a, CALIOP} \rangle$ is generally greater than the variability of spectrally cloud-screened
303 data ($\langle \tau_f \rangle$ and $\langle \tau_f^* \rangle$). The differences in variability can be ascribed to differences due to orbit distance from our
two sites, statistical anomalies due to the sparse and irregular nature of CALIOP AODs, and expected challenges in
comparing two inevitably different methods of discriminating clouds and aerosols. The difference of
306 $\langle \tau_f \rangle - \langle \tau_f^* \rangle$ is largely small and positive with the biggest positive difference being ~ 0.03 for Eureka in March
of 2011. $\langle \tau_f^* \rangle$ is ~ $\langle \tau_{a, Herber} \rangle$ at Ny-Ålesund with certain months (Dec., 2011, Jan. 2012 and Mar., 2012) when it is
significantly higher.

309 The standard deviations of Figure 4a aside, the estimates of τ_f and τ_f^* are not all equal in terms of estimated SDA
inversion errors. In Appendix B we show that the monthly averaged SDA retrieval errors ($\langle \Delta \tau_f \rangle$) were inordinately
312 large for the Ny-Ålesund data of 2011/2012 and that these large errors were associated with unphysically large spectral
curvature values (large values of the monthly averaged 2nd derivative, $\langle \alpha' \rangle$). While the retrieval errors were generally
~ the standard deviations for Eureka and the 2010/2011 season at Ny-Ålesund they were ~ 3 to 8 times the standard
315 deviations of the 2011/2012 season.

318 Figure 4b shows a scale zoom (relative to Figure 4a) for the component selected for comparison with GC simulations
($\langle \tau_f^* \rangle$), alongside the $\langle \tau_f, GC \rangle$ predictions. The former is largely greater than the latter, in keeping with the results
of Figure 1. The larger differences are frequently significant in terms of the standard deviations of the two data sets.
These differences are most likely due to model underestimation, if only on the basis of the persistence of this apparent
321 problem in the literature (Di Pierro, 2013; Breider et al., 2014). Potential sources of systematic bias in GC estimations
could be ascribed to a missing fine mode component (such as Di Pierro's hypothesis concerning the lack of a modelled
SS, fine mode aerosol ascribed to blown snow), emission underestimation, transport pathway errors, etc. Potential
324 sources of systematic bias in the starphotometry estimates include the frequently sporadic temporal sampling of the
star photometer as constrained by cloud and / or LIC conditions, unacceptable levels of background sunlight in the
late winter, star photometer calibration errors and errors in the SDA retrieval algorithm (there is also the wavelength
327 difference bias, mentioned above, which would increase the $\langle \tau_f, GC \rangle$ values by ~ 0.01 if those values had been
computed at 500 nm). All measured and modelling cases in Fig. 4b, except for Eureka in 2011, show an increase from
February to March. This increase is likely attributable to the late winter influence of Arctic haze (Herber et al., 2002)
while the 2011 springtime increase in $\langle \tau_f, GC \rangle$ at Ny-Ålesund is primarily attributable to fine mode SS.

330 **4 Conclusions**

333 We performed a climatological-scale analysis of polar winter AODs measured at two high-Arctic sites in comparison
with GC simulations and CALIOP retrievals. The results indicate significant cloud / LIC contamination which is only
partially corrected with a temporal cloud screening algorithm. Temporal cloud screening eliminates raw AODs due to
336 inhomogeneous (temporally and spatially variable) clouds and LICs. Homogeneous clouds and LICs that remain after
temporal cloud screening represent an inevitable systematic error in the estimation of AOD which varies from 78% to
210% at Eureka and from 2% to 157% for Ny-Ålesund. In principle, spectral cloud screening (to obtain fine mode
339 AODs) reduces raw AODs to the aerosol contribution if one assumes (supported by GC simulations) that coarse mode
aerosols are a minor part of the total AOD. Lidar analysis indicated, for the case of Eureka, that LICs appeared to have
a disproportionately large influence on the homogenous coarse mode optical depths that escape temporal cloud
342 screening.

345 The SDA filtered parameter $\langle \tau_f^* \rangle$ was chosen as the most conservative approach for climatological-scale estimates
of AOD. These values, typically larger than $\tau_{f, GC}$ estimates, are believed to be robust representations of τ_f variations:
348 an important consideration in a context of weak amplitude and weakly varying signal embedded in an environment of
large amplitude and strongly varying cloud and LIC signal. Large, low frequency, differences between τ_f^* and $\tau_{f, GC}$
351 appeared to often coincide with strong PBL events whose presence at either site was inferred from spatially expansive,
low-DR, PBL events in CALIOP profiles. These events were either not captured or, more likely, significantly
354 underestimated, by the GC simulations. High frequency $\tau_{f, GC}$ variations at Ny-Ålesund were attributed to SS while
low frequency variations at Eureka and Ny-Ålesund were attributable to sulfates. CALIOP profiles and AODs were
invaluable as spatial and temporal redundancy support (or, alternatively, as insightful points of contention) for star
photometry retrievals and GC estimates of AOD. Estimates of
 $\langle \tau_{a, CALIOP} \rangle$ were found however to be significantly more variable than their fine mode counterparts from star
photometry and GC simulations.

Appendix A – lidar based partition into homogeneous and inhomogeneous coarse mode contributions

357 Coarse mode optical depths (τ_c') derived from CANDAC Raman Lidar (CRL) profiles were computed as discussed
in Baibakov (2015) for an ensemble of profiles characterized by a sampling interval of approximately 10 minutes. In
360 a similar fashion to the homogeneous / inhomogeneous starphotometer AOD development above, the τ_c' values can
be divided into homogeneous and inhomogeneous ensembles. The process first involves, computing the temporal
363 derivative between pairs of τ_c' values and then discriminating homogeneous and inhomogeneous excursions by
comparing the absolute value of each temporal derivative ($|d\tau_c'/dt|$) with a threshold value. This values was chosen
366 to be 0.006 min^{-1} , the threshold discussed in Baibakov (2015) for star photometer sampling intervals of approximately
5 minutes (although the actual threshold employed in that paper was strategically chosen to be roughly equivalent in
performance to the 0.006 min^{-1} threshold where the effective sampling interval was increased to an hour in order to
369 better reject less inhomogeneous clouds). The lidar and star photometer were run fairly independently during the 2010-
2011 and 2011-2102 seasons and there was no strategic effort to have them collect synchronized data sets; the result
was a certain amount of commonality in their acquisition periods but also periods when one or the other was making
measurements alone. This yielded monthly average statistics for which $\langle \tau_c' \rangle$ was significantly greater than the
starphotometer average. We accordingly filtered the values with a maximum τ_c' cutoff filter so that their monthly
372 average was equal to the starphotometer average ($\langle \tau_c \rangle$) for each of the 4 months of Eureka data acquisition employed
in our comparisons.

375 In a similar fashion to equation (1) above, monthly averages of τ_c' can be expressed as;

$$378 \langle \tau_c' \rangle_- = \gamma_{hom-} \langle \tau_c', hom \rangle_- + (1 - \gamma_{hom-}) \langle \tau_c', inh \rangle_- \quad (A1)$$

$$\langle \tau_c' \rangle_+ = \gamma_{hom+} \langle \tau_c', hom \rangle_+ + (1 - \gamma_{hom+}) \langle \tau_c', inh \rangle_+ \quad (A2)$$

381 for integrations below and above h_{LIC} (the assumed upper limit of LICs). We note that these averages are carried out
over individual lidar profiles and thus that there is no daily averaging (i.e there is no use of a bold font as in equation
(1)). The parameter γ_{hom-} is given by $\gamma_{hom-} = N_{hom-} / (N_{hom-} + N_{inh-})$ where N_{hom-} and N_{inh-} are the
384 number of coarse mode optical depths in the homogeneous (accepted) and inhomogeneous (rejected) sub-ensembles
for integrations below h_{LIC} (analogous expressions exist for the "+" case above h_{LIC}). We note that the γ factors are
conservative ($\gamma_{hom\pm} + \gamma_{inh\pm} = 1$) because the total number of lidar-derived optical depths over the averaging period
387 of a month;

$$390 N = N_{hom-} + N_{inh-} = N_{hom+} + N_{inh+}$$

(a given lidar-derived optical depth must be in one of the two sub-ensembles). The lidar-derived average for the total
profile is given by;

393

$$\langle \tau'_c \rangle = \frac{\sum_{i=1}^N \tau'_c}{N} = \frac{\sum_{i=1}^N (\tau'_{c-} + \tau'_{c+})}{N}$$

396

$$= \langle \tau'_c \rangle_- + \langle \tau'_c \rangle_+ \quad (\text{A3})$$

Substituting equations (A1) and (A2) into (A3) yields;

399

$$\langle \tau'_c \rangle = \gamma_{hom-} \langle \tau'_{c, hom} \rangle_- + (1 - \gamma_{hom-}) \langle \tau'_{c, inh} \rangle_-$$

402

$$+ \gamma_{hom+} \langle \tau'_{c, hom} \rangle_+ + (1 - \gamma_{hom+}) \langle \tau'_{c, inh} \rangle_+ \quad (\text{A4})$$

thus partitioning $\langle \tau'_c \rangle$ values into their homogeneous and inhomogeneous components, below and above h_{LIC} .

405 Appendix B - SDA retrieval errors

408 An error model for all retrieved parameters of the SDA (in particular $\Delta\tau_f$) is given in O'Neill et al. (2003). Two
 409 important influences on $\Delta\tau_f$, at least within the context of an empirical analysis of Eureka and Ny-Ålesund star
 410 photometry retrievals are the amplitude of τ_f and the second derivative of τ_a (α'). Both influences can be approximated
 411 by a simple expression. In the first instance one has the pure differential in terms of τ_a and the fine mode fraction
 ($\eta = \tau_f/\tau_a$);

$$d\tau_f = \eta d\tau_a + \tau_a d\eta \quad (\text{B1})$$

414

Empirically one finds that rms errors associated with rms errors in the input AOD spectra are approximated by;

417

$$\Delta\tau_f \cong \tau_a \Delta\eta \quad (\text{B2})$$

420

The uncertainty $\Delta\eta$ is a strong function of the curvature at least for positive α' (which is generally true for cases where
 η is reasonably large). Thus;

$$\Delta\tau_f \propto \tau_a \alpha' \quad (\text{B3})$$

423

In the presence of comparatively strong variations in α' , $\Delta\tau_f$ will be roughly proportional to α' . For the 13 monthly
 426 averages of Figure 3c we obtained the results shown in Figure B1. Curvature values were excessive in the Ny-Ålesund
 data of 2011-2012 and this produced the quite large values of $\langle \Delta\tau_f \rangle$ seen in the figure. These excessive values
 correspond to unphysical spectral AOD variations, involving spectral changes (often non-physical valleys and peaks)
 429 which cannot be described by Mie theory. The second order spectral polynomial that we fit to AOD spectra before the
 application of the SDA tends to smooth out these artifactual variations but there will nonetheless be a residual
 influence.

5 Acknowledgements

432

We would like to thank NSERC for CCAR funding via the PAHA and NETCARE projects, the NSERC training
 program in Arctic Atmosphere Science as well as NSERC DG funding, the CSA, and the CFI for their financial

435 support. The contributions of the PEARL, AWI and CALIOP operations and processing staff are gratefully
acknowledged.

References

- 438 Baibakov, K., Pan-Arctic Characterization of Polar Winter Aerosols, Ph.D. thesis, Université de Sherbrooke and
University of Bremen, April, 2014.
- 441 Baibakov, K., O'Neill, N. T., Ivanescu, L., Duck, T. J., Perro, C., Herber, A., Schulz, K.-H., and Schrems, O.:
Synchronous polar winter starphotometry and lidar measurements at a High Arctic station, *Atmos. Meas. Tech.*,
8, 3789-3809, doi:10.5194/amt-8-3789-2015, 2015.
- 444 Blanchet, J.-P., E. Girard, Arctic 'greenhouse effect', *Nature*, Vol. 371, p. 383, 1994.
- 444 Di Pierro, M., L. Jaeglé, E. W. Eloranta, and S. Sharma, Spatial and seasonal distribution of Arctic aerosols observed
by the CALIOP satellite instrument (2006–2012), *Atmospheric Chemistry and Physics*, 13 (14), 7075–7095,
2013, doi:10.5194/acp-13-7075-2013.
- 447 Di Pierro, Modelling of the spatial and vertical distribution of Arctic aerosols, chap. 3, Ph.D. thesis, University of
Washington, 2013.
- 450 Fairlie, T. D., D.J. Jacob, and R.J. Park, The impact of transpacific transport of mineral dust in the United States,
Atmos. Environ., 1251-1266, 2007.
- 450 Fu, Tzung-May, et al. "Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of
secondary organic aerosols." *Journal of Geophysical Research: Atmospheres* 113.D15, 2008.
- 453 Garrett, T. J., C. Zhao, Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes,
Nature Letters, Vol 440, 2006, 687-789, doi:10.1038/nature04636.
- 456 Henze, D. K., and J. H. Seinfeld, Global secondary organic aerosol from isoprene oxidation, *Geophys. Res. Lett.*, 33,
L09812, 2006.
- 456 Henze, D. K., et al., Global modeling of secondary organic aerosol formation from aromatic hydrocarbons: High- vs.
low-yield pathways, *Atmos. Chem. Phys.*, 8, 2405-2420, 2008.
- 459 Herber, A., Thomason, L. W., Gernandt, H., Leiterer, U., Nagel, D., Schulz, K.-H., Kaptur, J., Albrecht, T. and
Notholt, J.: Continuous day and night aerosol optical depth observations in the Arctic between 1991 and 1999,
J. Geophys. Res., 107, 2002.
- 462 IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner,
465 M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University
Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- 465 Ivanescu, L., K. Baibakov, N. T. O'Neill, J.-P. Blanchet, Y. Blanchard, A. Saha, M. Rietzec and K.-H. Schulz,
Challenges in operating an Arctic telescope, *Proc. SPIE 9145, Ground-based and Airborne Telescopes V*,
468 914549, 2014; doi:10.1117/12.2071000.
- 468 Jaeglé, L., P.K. Quinn, T. Bates, B. Alexander, and J.-T. Lin, Global distribution of sea salt aerosols: New constraints
from in situ and remote sensing observations, *Atmos. Chem. Phys.*, 11, 3137-3157, doi:10.5194/acp-11-3137-
471 2011, 2011.
- 471 Lesin, G., L. Bourdages, T. J. Duck, J. R. Drummond, E.W. Eloranta, and V. P. Walden, Large surface radiative
forcing from topographic blowing snow residuals measured in the High Arctic at Eureka, *Atmos. Chem. Phys.*,
474 9, 1847–1862, 2009.
- 474 Liao, H., et al., Biogenic Secondary Organic Aerosol over the United States: Comparison of Climatological
Simulations with Observations, *J. Geophys. Res.*, 112, 2007.
- 477 Martin, R.V., Jacob, D.J., Yantosca, R.M., Chin, M. and Ginoux, P., Global and regional decreases in tropospheric
oxidants from photochemical effects of aerosols. *Journal of Geophysical Research*, 108, 4097, doi:
10.1029/2002JD002622, 2003.
- 480 Ma, X., K. Von Salzen, J. Li, Modelling sea salt aerosol and its direct and indirect effects on climate, *Atmos. Chem.*
Phys., 8, 1311–1327, 2008.
- 483 O'Neill, N. T., Eck, T. F., Smirnov, A., Holben, B. N., and Thulasiraman, S.: Spectral discrimination of coarse and
fine mode optical depth, *J. Geophys. Res.*, 108, 15, 4559, doi:10.1029/2002jd002975, 2003.
- Park, R. J., D. J. Jacob, M. Chin and R. V. Martin, Sources of carbonaceous aerosols over the United States and
implications for natural visibility, *J. Geophys. Res.*, 108(D12), 4355, doi:10.1029/2002JD003190, 2003.



- 486 Park, R. J., D. J. Jacob, B. D. Field, R. M. Yantosca, and M. Chin, Natural and transboundary pollution influences on
sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *J. Geophys. Res.*, 109,
D15204, doi:10.1029/2003JD004473, 2004.
- 489 Pye, H. O. T., H. Liao, S. Wu, L. J. Mickley, D.J. Jacob, D. K. Henze, and J. H. Seinfeld, Effect of changes in climate
and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States, *J. Geophys. Res.*, 114,
D01205, 2009.
- 492 Winker, D. M. A. Vaughan, A. Omar, Y. Hu, K. A. Powell, Z. Liu, W. H. Hunt, S. A. Young, Overview of the
CALIPSO Mission and CALIOP Data Processing Algorithms, *Jour. Atmos. Oceanic Tech.*, 26, 2310–2323,
2009.
- 495 Winker, D. M. J. L. Tackett, B. J. Getzewich, Z. Liu, M. A. Vaughan, and R. R. Rogers, The global 3-D distribution
of tropospheric aerosols as characterized by CALIOP, *Atmos. Chem. Phys.*, 13, 3345–3361, 2013.

498


501

504

507




510 **Figure 1** - Comparison of measured and cloud screened AODs (daily averages) derived from star photometry data with GEOS-
 513 Chem simulations over the polar winters of 2010/2011 and 2011/2012 at Eureka. The grey and black curves represent raw and
 cloud-screened AODs (τ and τ_{hom} respectively), while the light red and dark red curves represent the results of spectral cloud
 screening (τ^* and τ^* respectively). In order to be included in Fig. 1, all points required at least 10 raw AOD measurements per day.
 516 The simulated GC estimates of fine mode AOD (τ_{GC}) are shown as dashed red curves (see nomenclature details in the symbol and
 acronym glossary).




519

Figure 2 As per the legend of Figure 1 but for Ny-Ålesund




522

525 **Figure 3 - (a)** Comparison of temporal and spectral cloud screening (partitioned according to equations (4) and (5) respectively)
 528 for monthly AOD averages computed for Eureka and Ny-Ålesund during the polar winters of 2010/2011 and 2011/2012, **(b)**
 partitioning of lidar-derived coarse mode optical depths into homogeneous and inhomogeneous contributions above and below the
 nominal maximum altitude of low-altitude ice crystal layers (h_{LIC}) at Eureka.



531

534 **Figure 4** - (a) $\langle \tau_a, \text{CALIOP} \rangle$, $\langle \tau_f \rangle$, $\langle \tau_f^* \rangle$, and the 9-year AOD climatology of Herber et al. (2002). For our purposes we
 537 simply repeated Herber's values that belonged to the same calendar month, (b) Zoom of the $\langle \tau_f^* \rangle$ values of (a) compared with
 $\langle \tau_f, \text{GC} \rangle$.



540 **Figure B1** : Variation of the monthly averaged error in the SDA fine-mode AOD ($\langle \Delta\tau_f \rangle$) as a function of the monthly averaged
 spectral curvature ($\langle \alpha' \rangle$), the second derivative of the spectral AOD; c.f. O'Neill et al., 2003 for details). These monthly averages
 were computed using individual measurements rather than daily averages (and thus $\Delta\tau_f$ and α' are not in bold)

543

Symbol and acronym glossary	
High-level definitions	
AOD	The community uses the acronym "AOD" to represent a variety of concepts. These range from nominal aerosol optical depth which hasn't been cloud-screened to the conceptual (theoretical) interpretation of aerosol optical depth. In this paper we use AOD in the latter sense and apply adjectives as required (see for example "raw AOD" below).
PBL	Planetary boundary layer.
raw AOD	Nominal AOD derived before temporal or spectral cloud screening.
SS	Sea-salt.
SDA	Spectral Deconvolution Algorithm : τ_x retrieval that employs AOD spectra as input (method described in O'Neill et al., 2003).
τ_x	τ_a , τ_f , or τ_c for total, fine and coarse mode AODs. Without explicit subscript qualification to another data source (CALIOP, GC, etc.) this nomenclature is reserved for outputs of the SDA (at a reference wavelength of 500 nm) applied to raw AOD spectra. τ_f , or τ_c are conserved in the sense that $\tau_a = \tau_f + \tau_c$. This conservation expression, propagates through daily and monthly averages of the non cloud-screened, homogeneous and rejected data ensembles (the data sets corresponding to raw AODs, cloud-screened AODs and rejected raw AODs),.
$\bar{\tau}_x$	Daily average of τ_x (in bold : this avoids an awkward nomenclature of $\langle\langle\tau_x\rangle\rangle$ for the monthly average of daily averages
$\langle\bar{\tau}_x\rangle$	Monthly average of $\bar{\tau}_x$.
UT	Universal Time : the time standard (with respect to 24 hour clock) employed throughout this study.
Lower level definitions	
DR	CALIOP Depolarization Ratio (see Winker et al., 2009 for a definition and a discussion on the particulate typing capabilities of the DR).
GC	GEOS-Chem, version 9.01.03. FlexAOD (Flexible AOD) is employed to perform offline calculations of AOD.
h_{LIC}	assumed upper limit of LICs at Eureka (assumed to be 600 m).
LIC	low altitude ice crystals.
$\tau_{x, cs}$	τ_x values whose raw AOD inputs have been cloud-screened (have survived the cloud screening process). See $\tau_{x, hom}$ entry.
$\tau_{x, hom}$	τ_x values associated with homogeneous conditions, defined as $\tau_{x, hom} = \tau_{x, cs}$.
$\tau_{x, rej}$	τ_x values whose raw AOD inputs were rejected by the temporal cloud-screening process (see Baibakov et al., 2015 for details).
$\tau_{x, inh}$	In homogeneous cloud or LIC contribution to τ_x (see equation (3))
$\bar{\tau}_a, CALIOP$	Daily averaged (532 nm) values of the CALIOP AOD product within 500 km of Eureka / Ny-Ålesund.
$\bar{\tau}_f, GC$	Daily averaged GEOS-Chem, τ_f at 550 nm. In the supplementary information file, $\bar{\tau}_f, GC$ values are used for comparisons with $\bar{\tau}_a, CALIOP$ (i.e. since the CALIOP AOD product is not divided into fine and coarse mode contributions).
$\tau_a, Herber$	AODs at 532 nm from the 10-year AOD climatology at Ny-Ålesund of Herber et al. (2002). Some simple interpolation was employed to estimate tropospheric AODs for months that were not given in their Table 3 (Oct., Nov., Jan. and Mar.). Total AOD values were computed by adding a stratospheric AOD of 0.01 (derived from the 525 nm case of their Figure 5).
$\bar{\tau}_f^*$	$\bar{\tau}_f$ values on days for which $\bar{\tau}_f / \bar{\tau}_a$ values < 0.3 have been excluded.