

# 1    **Impacts of historical climate and land cover changes on fine particulate** 2    **matter (PM<sub>2.5</sub>) air quality in East Asia between 1980 and 2010**

3    *Yu Fu<sup>1,\*</sup>, Amos P. K. Tai<sup>2,\*</sup>, Hong Liao<sup>3</sup>*

4    <sup>1</sup> *Climate Change Research Center (CCRC), Chinese Academy of Sciences, Beijing 100029, China*

5    <sup>2</sup> *Earth System Science Programme and Graduate Division of Earth and Atmospheric Sciences, Faculty of*  
6    *Science, Chinese University of Hong Kong, Hong Kong*

7    <sup>3</sup> *School of Environmental Science and Engineering, Nanjing University of Information Science & Technology,*  
8    *Nanjing, 210044, China*

9    *\*Corresponding authors: Yu Fu ([fuyu@mail.iap.ac.cn](mailto:fuyu@mail.iap.ac.cn)), Amos P. K. Tai ([amostai@cuhk.edu.hk](mailto:amostai@cuhk.edu.hk))*

10

11    **Abstract.** To examine the effects of changes in climate, land cover and land use (LCLU), and  
12    anthropogenic emissions on fine particulate matter (PM<sub>2.5</sub>) between the 5-year periods 1981-1985  
13    and 2007-2011 in East Asia, we perform a series of simulations using a global chemical transport  
14    model (GEOS-Chem) driven by assimilated meteorological data and a suite of land cover and land  
15    use data. Our results indicate that climate change alone could lead to a decrease in wintertime  
16    PM<sub>2.5</sub> concentration by 4.0-12.0  $\mu\text{g m}^{-3}$  in northern China, but to an increase in summertime PM<sub>2.5</sub>  
17    by 6.0-8.0  $\mu\text{g m}^{-3}$  in those regions. These changes are attributable to the changing chemistry and  
18    transport of all PM<sub>2.5</sub> components driven by long-term trends in temperature, wind speed and  
19    mixing depth. The concentration of secondary organic aerosol (SOA) is simulated to increase by  
20    0.2-0.8  $\mu\text{g m}^{-3}$  in both summer and winter in most regions of East Asia due to climate change  
21    alone, mostly reflecting higher biogenic volatile organic compound (VOC) emissions under  
22    warming. The impacts of LCLU change alone on PM<sub>2.5</sub> (-2.1 to +1.3  $\mu\text{g m}^{-3}$ ) are smaller than that  
23    of climate change, but among the various components the sensitivity of SOA and thus organic  
24    carbon to LCLU change (-0.4 to +1.2  $\mu\text{g m}^{-3}$ ) is quite significant especially in summer, which is  
25    driven mostly by changes in biogenic VOC emissions following cropland expansion and changing  
26    vegetation density. The combined impacts show that while the effect of climate change on PM<sub>2.5</sub>  
27    air quality is more pronounced, LCLU change could offset part of the climate effect in some  
28    regions but exacerbate it in others. As a result of both climate and LCLU changes combined, PM<sub>2.5</sub>  
29    levels are estimated to change by -12.0 to +12.0  $\mu\text{g m}^{-3}$  across East Asia between the two periods.  
30    Changes in anthropogenic emissions remain the largest contributor to deteriorating PM<sub>2.5</sub> air  
31    quality in East Asia during the study period, but climate and LCLU changes could lead to a  
32    substantial modification of PM<sub>2.5</sub> levels.

33

## 1 Introduction

Over the recent decades atmospheric particulate matter (PM, or aerosols) has received considerable attention in East Asian countries due to its impacts on regional air quality, human health and climate change. A number of projection studies have examined the effects of changes in anthropogenic emissions and climate on future PM air quality globally and in East Asia (Fiore et al., 2012; Jiang et al., 2013), but there still exist large uncertainties arising from the complex interactions between aerosol chemistry, meteorology and the underlying land cover, especially for East Asia which is expected to undergo tremendous land use change in the next few decades (Hurt et al., 2011). A better understanding of how PM formation and removal have historically been shaped by meteorological and land surface conditions in East Asia would be useful to help to better project the future evolution of PM air quality. In this work, we use a chemical transport model driven by past meteorological and land surface data to evaluate the individual and combined effects of climate and land cover changes in East Asia over 1980-2010, and compare these effects with that of increasing anthropogenic emissions. This attribution of East Asia air quality trends in the past would shed light on their potential course of evolution in the coming few decades, and provide valuable information for policymaking concerning public health, land use and climate management.

Of particular public health concern is fine particulate matter ( $PM_{2.5}$ ), defined to be suspended liquid or solid particles with a diameter of 2.5  $\mu m$  or less.  $PM_{2.5}$  has been shown to have detrimental effects on human health, leading to increased mortality related to cardiovascular diseases and lung cancer (Krewski et al., 2009; Silva et al., 2013; Fang et al., 2013).  $PM_{2.5}$  is also associated with poor visibility and haze (Wang et al., 2014), and plays a significant role in modifying the Earth's energy budget (IPCC, 2013).  $PM_{2.5}$  has a variety of sources that depend on the chemical components, which include sulfate, nitrate, ammonium, black carbon (BC), organic carbon (OC), sea salt, and mineral dust. Air quality degradation with elevated  $PM_{2.5}$  in East Asia is primarily attributable to increasing anthropogenic emissions of their precursors. For instance, Wang et al. (2013) reported that annual mean sulfate-nitrate-ammonium concentrations increased by 60% in China from 2000 to 2006, which mainly resulted from the 60% and 80% increases in sulfur dioxide ( $SO_2$ ) and nitrogen oxides ( $NO_x$ ) emissions, respectively. Yang et al. (2015) found that the decadal trends of aerosol outflow from East Asia were dominated by the trends and variations in anthropogenic emissions, which could account for about 86% of the decadal trend in  $PM_{2.5}$ .

outflow over 1986-2006.

Air pollution associated with  $PM_{2.5}$  is also strongly sensitive to weather conditions and therefore influenced by climate change (Jacob and Winner, 2009; Tai et al., 2010; Fiore et al., 2012). Meteorological conditions affect the production, transport and deposition of  $PM_{2.5}$  components and their precursors. The effects of climate change on  $PM_{2.5}$  are complex due to the widely varying sensitivities of its components to different meteorological factors. For example, higher temperature can enhance sulfate concentration due to faster  $SO_2$  oxidation (Dawson et al., 2007; Jacob and Winner, 2009), while nitrate and OC concentrations decrease because higher temperature shifts more of these semivolatile components from the particle to gas phase (Liao et al., 2006; Kanakidou and Tsigaridis, 2007). This is further complicated by the covariation of temperature with cold-frontal passages, which are an important ventilating mechanism for air pollutants (Leibensperger et al., 2008; Tai et al., 2012a; 2012b). In general, changes in ventilation (e.g., wind speed and direction, mixing depth) significantly modify aerosol dispersion and transport. Zhu et al (2012) suggested that the decadal-scale weakening of the East Asian summer monsoon could have increased aerosol concentrations in eastern China mostly due to changes in circulation patterns. Furthermore, higher humidity generally promotes the formation of ammonium nitrate (Dawson et al. 2007; Tai et al., 2010), and all  $PM_{2.5}$  components are very sensitive to precipitation, which provides a major sink via scavenging (Dawson et al. 2007).

The tropospheric concentrations of  $PM_{2.5}$  are also influenced by land cover and land use (LCLU) changes. Vegetation represents an important source of biogenic volatile organic compounds (VOC), especially isoprene and monoterpenes, which are major precursors of secondary organic aerosols (SOA). SOA can be the major contributor to aerosols especially in remote regions far away from industrial influence (Carslaw et al., 2010), but can also be significant in many urban areas due to high year-round anthropogenic and summertime biogenic VOC emissions. For instance, Ding et al. (2014) investigated the origins of SOA in various Chinese regions based on observations from 14 sites during the summer of 2012, and found that biogenic isoprene was the major contributor ( $46 \pm 14\%$ ) to secondary OC in every site. In addition, vegetation and land surface characteristics may further modulate atmospheric aerosols by controlling soil  $NO_x$  emissions and the dry deposition of both gases and particles within the planetary boundary layer.

The impacts of climate and land cover changes on  $PM_{2.5}$  on a multidecadal scale have been quantified to various extents using [chemical transport models \(CTM\)](#) driven by assimilated meteorological data or simulated meteorological fields from general circulation

models (GCM). Jeong and Park (2013) found that sulfate-nitrate-ammonium concentrations decreased by 4% in summer but increased by 7% in winter in eastern China over 1985-2006 as a result of meteorological changes alone. Jiang et al. (2013) showed that following climate change alone under the IPCC A1B scenario, different aerosol species over China would generally be altered by  $-1.5$  to  $+0.8 \mu\text{g m}^{-3}$ , and  $\text{PM}_{2.5}$  concentration is projected to change by 10-20% in eastern China. Tai et al. (2013) reported that in China, climate change together with climate- and  $\text{CO}_2$ -driven natural vegetation changes would change annual mean surface SOA by  $-0.4$  to  $+0.1 \mu\text{g m}^{-3}$  over 2000-2050 under the IPCC A1B scenario, while anthropogenic land use change can increase SOA by up to  $0.4 \mu\text{g m}^{-3}$  over the same period. Wu et al. (2012) also predicted that changes in natural vegetation and anthropogenic land use over 2000-2050 would lead to higher summertime SOA over East Asia. Most of these studies, except Jeong and Park (2013), are concerned with the effects of future climate change and/or land cover change on aerosols. Previous studies that focus on the effects of historical LCLU change on East Asian aerosols are few. Fu and Liao (2014) suggested that surface SOA might decrease by as much as  $0.4 \mu\text{g m}^{-3}$  (-20%) between the late 1980s and mid-2000s over China due to changes in biogenic emissions induced by LCLU (mainly) and climate (to a lesser extent) changes. However, the overall role of LCLU change in controlling regional  $\text{PM}_{2.5}$  and its composition via biogenic emissions and deposition, especially under the simultaneous influence of climate change and  $\text{CO}_2$  fertilization, is still poorly understood.

Because of climate change, rapid economic development and other human activities, LCLU in East Asia has undergone remarkable changes in the past 30 years. Especially in China, some major economic reforms and land use policies have been implemented since December 1978, which together with simultaneous changes in climate have resulted in a whirlwind of changes in the terrestrial environments of China. Based on satellite-derived images and surveys, LULC changes in China from late 1980s to the mid-2000s are characterized by an expansion of urban areas, deserts and bare lands, and an overall decrease in vegetation coverage (Fu and Liao, 2014). In this study, we use the GEOS-Chem global chemical transport model (CTM) driven by past land cover data and meteorological fields from a single and coherent set of assimilated meteorology (MERRA), which covers the period 1979-present. We quantify the impacts of historical changes in climate, land cover and land use on  $\text{PM}_{2.5}$  air quality in East Asia between two 5-year periods: historical period, 1981-1985 (referred to as “1980”), and the present day, 2007-2011 (referred to as “2010”). We consider 5-year averages in each of these periods to account for interannual variability.

We also compare the effects of climate and LCLU changes with the contribution from anthropogenic emissions over the same time frame. The findings would shed light on the possible climate and land use “penalties” or benefits that might have exacerbated or offset the effect of anthropogenic emissions in the past, and provide a constraint for future air quality projections, which currently have large uncertainties for East Asia.

## 2 Methods and model description

We perform a series of model experiments to simulate aerosols using the GEOS-Chem global CTM (version 9-02) with assimilated meteorology and integrated historical land cover data. The modeling framework in this study is the same as that used in Fu and Tai (2015). In brief, GEOS-Chem performs fully coupled simulations of ozone-NO<sub>x</sub>-VOC (Bey et al., 2001) and aerosol chemistry (Park et al., 2003, 2004; Pye et al., 2010). In this study, GEOS-Chem is driven by the assimilated meteorological data from Modern Era Retrospective-analysis for Research and Applications (MERRA) with a horizontal resolution of 2.0 ° latitude by 2.5 ° longitude, and a vertical resolution of 47 levels. Aerosol species simulated include sulfate, nitrate, ammonium, organic carbon, black carbon, sea salt, and mineral dust. Inorganic aerosol thermodynamic equilibrium calculations are based on the ISORROPIA II scheme of Fountoukis and Nenes (2007). SOA formation is based on the reversible gas-aerosol partitioning of VOC oxidation products (Chung and Seinfeld, 2002, Liao et al., 2007) with precursors including isoprene, monoterpenes, alcohols, and aromatic hydrocarbons (Henze et al., 2008). The wet deposition scheme for water-soluble aerosol species is described by Liu et al. (2001). Model details for other relevant modules and emission inventories can be found in Fu and Tai (2015).

The land cover dependence of atmospheric chemistry is mainly encapsulated in two land cover inputs, [namely leaf area index \(LAI\) and land](#) or plant functional types (as a single categorical value or as fractional coverage in each grid cell), mostly through their effects on biogenic VOC and soil NO<sub>x</sub> emissions, and dry deposition velocities. The emissions of biogenic VOC species in each grid cell are determined by the canopy-scale emission factors multiplied by various activity factors that account for variations in temperature, light, leaf age and LAI, using [online](#) the Model of Emissions of Gases and Aerosols from Nature (MEGAN) module [\(Guenther et al., 2012\) embedded in GEOS-Chem with the emissions time-step set to 30 minutes](#). Soil NO<sub>x</sub> emission follows the parameterization of Yienger and Levy (1995) and Hudman et al. (2012), which includes a physical representation of key soil

processes derived from field measurements, with dependence on vegetation types, temperature, precipitation history, fertilizer use, and a canopy reduction factor. The reservoir of nitrogen associated with manure and chemical fertilizer remains unchanged between 1980 and 2010 by using the fixed inventory for fertilizer and manure emissions from Potter et al. (2010). Dry deposition follows the resistance-in series scheme of Wesely (1989) as implemented by Wang et al. (1998), and is dependent on species properties, land cover types and meteorological conditions. It uses the Olson land cover classes with 76 land types (Olson, 1992) reclassified into 11 land types. Aerosol dry deposition follows Zhang et al. (2001) as described by Pye et al. (2009). The land cover inputs used for this study are derived from a fusion of various datasets, including the Moderate Resolution Imaging Spectroradiometer (MODIS) land cover product (MCD12Q1), the National Land Cover Dataset (NLCD) for China, harmonized historical land use for Representative Concentration Pathways (RCP) from Hurtt et al. (2011), and the global LAI product from Liu et al. (2012). See Fu and Tai (2015) for detailed description of these land cover inputs.

We conduct a 5-year simulation in the present-day period (2007-2011) with the corresponding meteorological variables, emissions, and land cover and land use as the control simulation [*CTRL*]. Sensitivity simulations are conducted for: (1) [*S\_CLIM*]: historical (1981-1985) climate with present-day land cover inputs and emissions (scaled to 2005 levels) used in [*CTRL*]; (2) [*S\_LCLU*]: historical land cover inputs with present-day climate and emissions used in [*CTRL*]; (3) [*S\_COMB*]: historical climate and land cover inputs but with present-day emissions used in [*CTRL*]; and (4) [*S\_ATH*]: historical emissions scaled to 1985 levels but with present-day climate and land cover inputs. Additional sensitivity simulations perturbing certain meteorological variables while keeping the rest at present-day levels are conducted to examine which meteorological factors have been the most important for shaping different PM<sub>2.5</sub> components.

### 3 Simulated spatiotemporal variations of PM<sub>2.5</sub> concentrations

Simulated seasonal mean surface concentrations of sulfate (SO<sub>4</sub><sup>2-</sup>), nitrate (NO<sub>3</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>), black carbon (BC), organic carbon (OC), and total PM<sub>2.5</sub> (sum of sulfate, nitrate, ammonium, BC and OC) in East Asia averaged over 2007-2011 from the [*CTRL*] simulation are shown in Fig. 1. Simulated PM<sub>2.5</sub> is high over the eastern regions of East Asia (east of 110°E), especially in eastern and central China, where PM<sub>2.5</sub> is in the range of 60-90 μg m<sup>-3</sup> in all seasons. Nitrate (~33%), sulfate (~24%), ammonium (~19%) and OC (~22%)



are the major components of annual mean PM<sub>2.5</sub> in eastern China (Table S1). Based on the measurements from 16 sites in China, Zhang et al. (2012) reported that sulfate (~16%), OC (~15%), nitrate (~7%), ammonium (~5%) and mineral aerosol (~35%) are majorities of the total PM<sub>10</sub> concentration. The contribution of each species to total PM<sub>2.5</sub> is generally overestimated compared with nationwide measurements from Zhang et al. (2012), probably due to the exclusion of mineral dust and sea salt in this work, and the inadequacy of current emission inventories.

Among all seasons, simulated sulfate is the highest in summer (JJA) mainly due to enhanced photochemical oxidation of SO<sub>2</sub>. The maximum summertime sulfate concentration of 25-35  $\mu\text{g m}^{-3}$  is found in the North China Plain, while the largest sulfate concentration in other seasons is within the range of 15-25  $\mu\text{g m}^{-3}$  over much of eastern and central China (Fig. 1). Nitrate shows a different seasonal variation with a maximum in winter (DJF) and minimum in summer, as the low wintertime temperature promotes ammonium nitrate formation. The maximum wintertime nitrate concentration is in the range of 30-40  $\mu\text{g m}^{-3}$  around central China (Fig. 1). The spatial distribution of seasonal ammonium concentration is similar to that of nitrate and sulfate, within a range of 5-20  $\mu\text{g m}^{-3}$  in all four seasons over the domain of study. The concentrations of OC and BC in the eastern regions of East Asia are the largest in winter, with maximum values of 20-25  $\mu\text{g m}^{-3}$  and 5-10  $\mu\text{g m}^{-3}$ , respectively, reflecting higher anthropogenic emissions associated with domestic heating in winter. Our simulated aerosol concentrations and distributions show general agreement with previous studies in East Asia (Fu et al., 2012; Wang et al., 2013; Jeong and Park, 2013; Lou et al., 2014), demonstrating the ability of GEOS-Chem to capture the spatial variations of different PM<sub>2.5</sub> species despite biases in the absolute concentrations. The model biases of simulated annual mean sulfate, nitrate, ammonium, BC and OC in East Asia are -10%, +31%, +35%, -56% and -76%, respectively (Fu et al., 2012; Wang et al., 2013).

The distribution of simulated seasonal surface SOA concentration is shown in Fig. 2. Total SOA is the sum of SOA from the oxidation of biogenic VOC (including isoprene, monoterpenes, sesquiterpenes) and anthropogenic VOC (including benzene, toluene and xylene). In winter, simulated SOA is less than 0.5  $\mu\text{g m}^{-3}$  over the northern regions of East Asia, where biogenic emissions, temperature, and radiation are lower than in southern regions (south of 25°N). In summer, simulated SOA over the eastern parts of East Asia are within the range of 2.0-7.0  $\mu\text{g m}^{-3}$  with the highest concentration in the lower and middle reaches of Yangtze River and east of Sichuan province. SOA distributions in spring (MAM) and fall (SON) are similar, within the range of 2.0-5.0  $\mu\text{g m}^{-3}$  over central and southeastern

China. Figure 2 also shows that SOA from biogenic emissions is the major contributor to total SOA in East Asia, larger than the anthropogenic contribution by nearly an order of magnitude in all seasons.

Of the total OC in China, about 6-27% is attributable to SOA (Table S1), which is lower than the fraction (~55-60%) found by Zhang et al. (2012). Fig. 3a compares the simulated and observed July-September mean SOA concentrations in 14 sites over China. The observed concentrations are taken from Ding et al. (2014), which are measured at 5 urban sites, 7 rural or sub-urban sites, and 2 remote sites around China in 2012. Simulated SOA is underestimated by about 60%. The simulated secondary organic carbon (SOC) to OC ratios are also compared with the observed ratios from Zhang et al. (2012) (Fig. 3b), which are measured at 14 sites in China during 2006-2007. The simulated seasonal mean SOC contributions range from 14% to 54% over China, with relatively higher contribution (54%) in summer but much lower (14%) in winter, while the observed ratios exhibit little seasonality between 51-57%. The simulated SOC/OC ratios agree generally well with the measurements in summer with a low-bias of 5%, while that in winter, spring, and autumn are underestimated by about 74%, 56%, and 41%, respectively (Fig. 3b). This suggests that SOA is underestimated in these seasons, likely reflecting **uncertainties not only in** the VOC emission inventories but also SOA formation mechanism. In this study, we consider SOA formation from absorptive partitioning of semivolatile organic compounds, but previous studies have suggested the potential importance of heterogeneous uptake of dicarbonyls (Fu et al., 2009) and oxidation of gas-phase semivolatile primary organic compounds and intermediate VOC (e.g., naphthalene) (Pye et al., 2010), which may constitute potentially large SOA sources but are not included in our study.

#### **4 Impacts of climate change alone on PM<sub>2.5</sub>**

Climate change alone ( $[CTRL] - [S\_CLIM]$ ) can substantially alter the simulated concentrations of PM<sub>2.5</sub> and its components in East Asia between 1980 and 2010. Here we focus on the changes in winter and summer (Fig. 4 and Fig. 5, respectively). The most significant decrease of PM<sub>2.5</sub> occurs in winter (Fig. 4), with the maximum decrease in the North China Plain by up to 12.0  $\mu\text{g m}^{-3}$ , while wintertime PM<sub>2.5</sub> over southeastern and central-western China (30 °-40 °N; 100 °-110 °E) increases by up to 4.0  $\mu\text{g m}^{-3}$ , due to climate change alone. We find that the spatial pattern of changes in summertime PM<sub>2.5</sub> due to climate change alone is mostly reversed (Fig. 5), being enhanced by as much as 8.0  $\mu\text{g m}^{-3}$  in the



North China Plain and central-western China, but reduced by up to  $8.0 \mu\text{g m}^{-3}$  in central and southern China.

Climate-driven changes in wintertime  $\text{PM}_{2.5}$  are primarily driven by the changing concentrations of sulfate, nitrate and ammonium induced by meteorological changes (Fig. 4). Wintertime sulfate increases by up to  $3.0 \mu\text{g m}^{-3}$  over southeastern and central-western China, but decreases by up to  $2.0 \mu\text{g m}^{-3}$  in the North China Plain. In southeastern and central-western China, the simulated increase in wintertime sulfate is mostly a result of regionally reduced surface wind speed and planetary boundary layer (PBL) (which reduce ventilation and mixing), but also in part due to increased temperature (which accelerates  $\text{SO}_2$  oxidation). In other regions, especially within the North China Plain, the simulated decrease in sulfate reaches a maximum of  $-2.0 \mu\text{g m}^{-3}$ , likely reflecting increased surface wind speed. Simulated nitrate and ammonium concentrations decrease in most of eastern parts of China, which can be explained by the elevated temperature, decreased RH, enhanced wind speed and PBL between the two periods (Fig. 4). The spatial patterns of changes and our sensitivity simulations suggest that increased temperature has contributed the most broadly to the reduction in ammonium nitrate formation, but [depending on region, changes in wind speed](#) and PBL might have either substantially enhanced (most of northeastern, northern and central China) or partly counteracted (e.g., southeastern China) [the reduction in ammonium nitrate](#), with RH playing only a minor role. OC and BC changes generally follow the same patterns as [those](#) for sulfate, reflecting influence from the same suite of meteorological variables.

Climate-driven changes in summertime total  $\text{PM}_{2.5}$  are also dominated by changes in the inorganic components, which generally show an opposite sign of changes to that in winter. Summertime sulfate, nitrate and ammonium increase over northern China, the North China Plain, and part of northwestern and eastern China, but decrease elsewhere (Fig. 5). In much of central, southern and northeastern China, decreased sulfate, nitrate and ammonium are attributable to the significantly increased PBL, which enhances mixing and dilution despite reduced wind speed. Increased temperature also [partly](#) contributes to lower ammonium nitrate. In northern and northwestern China, however, the large increase in sulfate is likely driven by regionally reduced PBL and wind speed. The significantly enhanced nitrate and ammonium concentrations in much of northern and eastern China are shaped by less ventilation driven by wind speed in the North China Plain, further modulated by regional cooling and increased RH around Shandong province (Fig. 5). We also find that simulated summertime OC and BC increase in much of eastern China, reflecting a combination of

increased temperature and reduced wind speed, except in south-central China where OC is reduced likely by enhanced PBL mixing.

Climate change alone could lead to increased SOA concentration in both winter and summer in East Asia by as much as  $+0.8 \mu\text{g m}^{-3}$  (Fig. 6). The climate-driven changes in SOA are primarily due to changes in temperature that influence biogenic VOC emissions (Liao et al., 2006). In winter, with enhanced isoprene and monoterpene emissions due to warming, SOA concentration in the southern parts of China increases by up to  $0.4 \mu\text{g m}^{-3}$ . In summer, the simulated SOA concentration changes within the range of  $-0.5$  to  $+0.8 \mu\text{g m}^{-3}$ , mostly attributable to the biogenic emission changes but also partly modulated by transport changes, similar to the pattern of OC changes (Fig. 5).

## 5 Impacts of land cover and land use change alone on $\text{PM}_{2.5}$

Figure 7 represents the impacts of 1980-2010 land cover and land use change alone on  $\text{PM}_{2.5}$  concentrations ( $[CTRL] - [S\_LCLU]$ ). We find that although the LCLU change effects on all  $\text{PM}_{2.5}$  components are generally smaller than the climate change effects, LCLU change can in part modify (either exacerbate or offset) the sensitivity of  $\text{PM}_{2.5}$  to climate change. In most of the eastern parts of East Asia, wintertime  $\text{PM}_{2.5}$  concentration increases by up to  $1.3 \mu\text{g m}^{-3}$  as a result of LCLU change alone (Fig. 7a). In contrast, LCLU change alone leads to a decrease in  $\text{PM}_{2.5}$  by up to  $-2.1 \mu\text{g m}^{-3}$  over much of China in summer except in some of the southern parts (Fig. 7b). Such changes are mostly attributable to changes in nitrate, ammonium and OC; BC is largely unaffected by LCLU change.

We find that with LCLU change alone, nitrate in winter increases by up to  $0.6 \mu\text{g m}^{-3}$  around central China ( $\sim 30^\circ\text{N}$ ). Such an increase is largely driven by reduced  $\text{HNO}_3$  and  $\text{NO}_2$  dry deposition following a decrease in wintertime LAI (Fig. S1 and Fig. S2 in supplementary materials). The changes in ammonium follow the changes of nitrate, with which they are chemically linked, and are partly due to reduced  $\text{NH}_3$  dry deposition. In summer, the sign of changes in nitrate and ammonium are mostly reversed. Nitrate decreases by as much as  $-1.2 \mu\text{g m}^{-3}$  in the North China Plain, mostly driven by the enhanced  $\text{HNO}_3$  and  $\text{NO}_2$  dry deposition resulting from enhanced summertime LAI (thus vegetation density) (Fig. S1 and Fig. S3), overshadowing the effect of increased soil  $\text{NO}_x$  emission from cropland expansion. See Fu and Tai (2015) for more discussion on East Asian land cover change.

LCLU change effects on OC are relatively minor in winter ( $-0.1$  to  $+0.2 \mu\text{g m}^{-3}$ ), but are significant in summer ( $-0.4$  to  $+1.0 \mu\text{g m}^{-3}$ ) since both LAI and plant functional type (PFT)

changes can significantly affect the emissions of biogenic VOCs, which are the major precursors to SOA especially in summer. In much of the North China Plain, central and northeastern China where deforestation and cropland expansion (in terms of PFT changes) have been the most rapid (Fig. S1), the effect of cropland expansion appears to dominate over that of enhanced summertime grid cell-averaged LAI in modifying biogenic emissions, leading to a decrease in OC that largely reflects a reduction in biogenic emissions (Fig. 8). The concentration of OC increases elsewhere, especially in southwestern and southern China where reforestation has been observed and increased summertime LAI further enhances the increase in biogenic emissions (Fig. S1). Figure 8 shows the contribution to surface SOA concentration from LCLU change alone, which largely follows the spatial pattern of OC changes and reflects the underlying changes in biogenic VOC emissions. Summertime SOA in summer increases by more than  $1.0 \mu\text{g m}^{-3}$  in southern and southwestern China between 1980 and 2010, but decreases by up to  $0.4 \mu\text{g m}^{-3}$  in other parts of China. In winter, LCLU change increases SOA by up to  $0.4 \mu\text{g m}^{-3}$  around Guizhou province, but leads to only negligible decreases in SOA in much of the rest of East Asia due to the small biogenic VOC emissions in winter.

## **6 Combined impacts of climate, land cover and land use changes vs. anthropogenic emissions**

With anthropogenic emissions fixed at present-day levels, the changes in wintertime  $\text{PM}_{2.5}$  resulting from both climate and LCLU changes combined between 1980 and 2010 are in the range of  $-12.0$  to  $+6.0 \mu\text{g m}^{-3}$  in East Asia, with the maximum decrease found in the North China Plain (Fig. 9a). In summer, changes in  $\text{PM}_{2.5}$  are within the range of  $-8.0$  to  $+12.0 \mu\text{g m}^{-3}$  under the combined effects of climate and LCLU changes, with an enhancement of  $4.0$ - $12.0 \mu\text{g m}^{-3}$  in the North China Plain. The changes of  $\text{PM}_{2.5}$  and its components are largely driven by climate change. For SOA alone, the combined effects of climate change and LCLU changes modify summertime SOA by  $-0.6$  to  $+1.2 \mu\text{g m}^{-3}$ , reflecting comparable contribution from both climate and LCLU changes, which can either exacerbate or offset each other depending on the region. For instance, in southwestern China, climate change alone might decrease SOA (Fig. 6), but LCLU change could more than offset the climate effect there (Fig. 9b).

Fig. 9c shows the effects on total  $\text{PM}_{2.5}$  and SOA of changes in anthropogenic emissions alone ( $[CTRL] - [S\_ANTH]$ ), which we find as expected to be the dominant factor shaping

PM<sub>2.5</sub> air quality in East Asia over 1980-2010. In both summer and winter, PM<sub>2.5</sub> is simulated to increase on average by 37% and 54% in East Asia, respectively, resulting from changes in anthropogenic emissions. From 1985 to 2005, anthropogenic emissions of NO<sub>x</sub>, CO, SO<sub>2</sub>, NH<sub>3</sub>, OC, and BC have increased by 180%, 143%, 52%, 50%, 36% and 46%, respectively, over East Asia (Table S2). Such emission-driven changes in PM<sub>2.5</sub> would be partially offset in winter but substantially enhanced in summer by climate- and LCLU-driven changes in the most polluted regions (e.g., in the vicinity of the North China Plain) between 1980 and 2010.

## 7 Conclusions and discussion

We simulate the effects of changes in climate, land cover and land use (LCLU), and anthropogenic emissions between the two 5-year periods 1981-1985 (historical) and 2007-2011 (present-day) on the surface concentrations of total PM<sub>2.5</sub> and its components including sulfate, nitrate, ammonium, organic carbon (OC), and black carbon (BC) in East Asia using the GEOS-Chem chemical transport model driven by assimilated meteorological data and a suite of satellite- and survey-derived LCLU data. GEOS-Chem is shown to capture the spatial and seasonal variations of different PM<sub>2.5</sub> species in the present day despite some significant biases in the absolute concentrations. The present-day secondary organic aerosol (SOA) concentration is underestimated in comparison with measurements in China. The volatile organic compound (VOC) emission inventory and SOA formation mechanism might represent the major sources of uncertainty for SOA simulation in the model.

With anthropogenic emissions fixed at present-day levels, the effects of climate change alone on the concentrations of different PM<sub>2.5</sub> species display substantial seasonal differences and spatial variability between the two periods. In winter, climate change alone is found to decrease PM<sub>2.5</sub> concentration by as much as 12.0  $\mu\text{g m}^{-3}$  in the North China Plain, but increase by up to 4.0  $\mu\text{g m}^{-3}$  in southeastern, northwestern and southwestern China. These changes are mostly attributable to the changing chemistry and transport of different species driven by changes in temperature, surface wind speed and planetary boundary layer (PBL) depth. In summer, however, the changes of PM<sub>2.5</sub> display a generally opposite pattern with increases (+6.0 to +8.0  $\mu\text{g m}^{-3}$ ) found in the North China Plain, and reductions (more than -4.0  $\mu\text{g m}^{-3}$ ) found in most places of central and southern China, reflecting changes in the same suite of meteorological variables but with varying relative importance. Climate change alone leads to an increase in SOA concentration both in winter and summer (0.2-1.0

$\mu\text{g m}^{-3}$ ) in most of the eastern parts of China, primarily driven by enhanced biogenic VOC emissions resulting from warming.

The impacts of LCLU change alone on total  $\text{PM}_{2.5}$  ( $-2.1$  to  $+1.3 \mu\text{g m}^{-3}$ ) is generally smaller than that of climate change alone, but the impacts on SOA and thus OC can be quite significant ( $-0.4$  to  $+1.2 \mu\text{g m}^{-3}$ ), reflecting the effects of deforestation, cropland expansion, reforestation as well as climate- and  $\text{CO}_2$ -driven changes in leaf area index (LAI). Changes in anthropogenic emissions from 1985 to 2005 levels are still the largest contributor to worsening  $\text{PM}_{2.5}$  air quality in both summer and winter, leading to an increase in  $\text{PM}_{2.5}$  by 54% on average in winter and 37% in summer over East Asia.

Our results show that the annual mean concentrations for  $\text{PM}_{2.5}$  are in the range of 10-70  $\mu\text{g m}^{-3}$  in East Asia over 2007-2011 (Fig. S4). In many places of eastern and central China, and part of southwestern China, annual mean  $\text{PM}_{2.5}$  well exceeds  $35 \mu\text{g m}^{-3}$ , which is the limit value specified in the current air quality standards of the Ministry of Environmental Protection of China for all the cities nationwide except some special regions. The  $\text{PM}_{2.5}$  air quality guideline set by the World Health Organization (WHO) is that annual mean concentration must not exceed  $10 \mu\text{g m}^{-3}$ , which is even stricter (WHO, 2006). Beyond this level, the morbidity and premature mortality of health risks such as lung cancer, cardiovascular and respiratory diseases would increase in response to long-term exposure to  $\text{PM}_{2.5}$ . From this perspective, our results indicate that the effects of climate change would partly counteract the emission-driven increase in  $\text{PM}_{2.5}$  in winter by a substantial fraction in most of northeastern, northern, eastern and central China especially in the North China Plain, imposing a so-called “climate benefit” for  $\text{PM}_{2.5}$  air quality and thus public health. However, climate change could substantially exacerbate  $\text{PM}_{2.5}$  pollution in summer in the North China Plain, northern and northwestern China, imposing a “climate penalty” instead. We also find that LCLU change might partially offset the increase in summertime  $\text{PM}_{2.5}$  but further enhance wintertime  $\text{PM}_{2.5}$  in the model through modifying the dry deposition of various  $\text{PM}_{2.5}$  precursors and biogenic VOC emissions, which also act as important factors in modulating air quality.

There are various sources of uncertainties in this study. Previous work by Tai et al. (2013) suggested that the inclusion of  $\text{CO}_2$  inhibition effect could reduce the sensitivity of surface SOA to climate and land cover changes in regions where isoprene emission is important, but this effect is not considered here. However, experimental data for  $\text{CO}_2$ -isoprene relationship at lower  $\text{CO}_2$  levels are generally scarce and not robust enough to be included in our model periods. In addition, as pointed out by Fu and Tai (2015), vegetation

composition and resistance values for each vegetation or land type in this work are assumed to remain unchanged between 1980 and 2010, which may yield part of the uncertainties. The changes in manure and chemical fertilizer associated with the changes in agriculture practices and land use are also not taken into account in this study, which may affect soil  $\text{NO}_x$  emission and contribute to the formation of inorganic  $\text{PM}_{2.5}$ , which may be particularly important in the future as anthropogenic  $\text{NO}_x$  emissions are expected to decline. Furthermore, the deposition of  $\text{PM}_{2.5}$  might also affect the terrestrial ecosystems and crops in various manners, e.g., via the acidification of soils that may lead to more leaching of mineral nutrients, and the introduction of excess nitrogen that may fertilize the soils or disrupt the soil nitrogen cycle. These processes would induce feedback effects that can further modify the land cover but are not explicitly taken into account in this study. All these issues remain poorly understood and warrant further investigation in future studies.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China under grants 41405138 given to Yu Fu, and a start-up grant from the Croucher Foundation and The Chinese University of Hong Kong (6903601) given to Amos P. K. Tai. MERRA data used in this study/project have been provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center through the NASA GES DISC online archive. We acknowledge the MODIS LAI product provided by Land Processes Distributed Active Archive Center (LP DAAC). We also thank the Information Technology Services Centre (ITSC) at The Chinese University of Hong Kong for their devotion in providing the necessary computational services for this work.

## References

- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B., Fiore, A. M., Li, Q., Liu, H., Mickley, L. J., and Schultz, M.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23,073-23,096, doi:10.1029/2001JD000807, 2001.
- Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth system, *Atmos. Chem. Phys.*, 10, 1701-1737, doi: 10.5194/acp-10-1701-2010, 2010.



- Chung, S. H., and Seinfeld, J. H.: Global distribution and climate forcing of carbonaceous aerosols, *Journal of Geophysical Research*, 107, 4407, doi: 10.1029/2001jd001397, 2002.
- Dawson, J. P., Adams, P. J., and Pandis, S. N.: Sensitivity of ozone to summertime climate in the eastern USA: A modeling case study, *Atmos. Environ.*, 41, 1494-1511, doi: 10.1016/j.atmosenv.2006.10.033, 2007.
- Ding, X., He, Q.-F., Shen, R.-Q., Yu, Q.-Q., and Wang, X.-M.: Spatial distributions of secondary organic aerosols from isoprene, monoterpenes,  $\beta$ -caryophyllene, and aromatics over China during summer, *J. Geophys. Res.*, 119, 2014JD021748, doi: 10.1002/2014JD021748, 2014.
- Fang, Y., Mauzerall, D., Liu, J., Fiore, A., and Horowitz, L.: Impacts of 21<sup>st</sup> century climate change on global air pollution-related premature mortality, *Climatic Change*, 121, 239-253, doi: 10.1007/s10584-013-0847-8, 2013.
- Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsoren, S., Eyring, V., Folberth, G. A., Ginoux, P., Horowitz, L. W., Josse, B., Lamarque, J.-F., MacKenzie, I. A., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S. T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T., and Zeng, G.: Global air quality and climate, *Chem. Soc. Rev.*, 41, 6663-6683, doi: 10.1039/C2CS35095E, 2012.
- Fountoukis, C., and A. Nenes, ISORROPIA II: A computationally efficient thermodynamic equilibrium model for  $K^+$ - $Ca^{2+}$ - $Mg^{2+}$ - $NH_4^+$ - $Na^+$ - $SO_4^{2-}$ - $NO_3^-$ - $Cl^-$ - $H_2O$  aerosols, *Atmos. Chem. Phys.*, 7(17), 4639-4659, doi: 10.5194/acp-7-4639-2007, 2007.
- Fu, T.-M., Cao, J. J., Zhang, X. Y., Lee, S. C., Zhang, Q., Han, Y. M., Qu, W. J., Han, Z., Zhang, R., Wang, Y. X., Chen, D., and Henze, D. K.: Carbonaceous aerosols in China: top-down constraints on primary sources and estimation of secondary contribution, *Atmos. Chem. Phys.*, 12, 2725-2746, doi: 10.5194/acp-12-2725-2012, 2012.
- Fu, T.-M., Jacob, D. J., and Heald, C. L.: Aqueous-phase reactive uptake of dicarbonyls as a source of organic aerosol over eastern North America, *Atmos. Environ.*, 43, 1814-1822, doi: 10.1016/j.atmosenv.2008.12.029, 2009.
- Fu, Y., and Liao, H.: Impacts of land use and land cover changes on biogenic emissions of volatile organic compounds in China from the late 1980s to the mid-2000s: implications for tropospheric ozone and secondary organic aerosol, *Tellus B*, 66, 24987, doi: 10.3402/tellusb.v66.24987, 2014.
- Fu, Y., and Tai, A. P. K.: Impact of climate and land cover changes on tropospheric ozone air

quality and public health in East Asia between 1980 and 2010, *Atmos. Chem. Phys.*, 15, 10093-10106, doi: 10.5194/acp-15-10093-2015, 2015.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, doi: 10.5194/gmd-5-1471-2012, 2012.

Henze, D. K., Seinfeld, J. H., Ng, N. L., Kroll, J. H., Fu, T.-M., Jacob, D. J., and Heald, C. L.: Global modeling of secondary organic aerosol formation from aromatic hydrocarbons: High- vs. low-yield pathways, *Atmos. Chem. Phys.*, 8, 2405-2420, doi: 10.5194/acp-8-2405-2008, 2008.

Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C., and Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions: implementation and space based-constraints, *Atmos. Chem. Phys.*, 12, 7779-7795, doi: 10.5194/acp-12-7779-2012, 2012.

Hurtt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117-161, doi: 10.1007/s10584-011-0153-2, 2011.

IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp. pp., 2013.

Jacob, D. J., and Winner, D. A.: Effect of climate change on air quality, *Atmos. Environ.*, 43, 51-63, doi: 10.1016/j.atmosenv.2008.09.051, 2009.

Jiang, H., Liao, H., Pye, H.O.T., Wu, S., Mickley, L.J., Seinfeld, J.H., Zhang, X.Y. Projected effect of 2000–2050 changes in climate and emissions on aerosol levels in China and associated transboundary transport. *Atmos. Chem. Phys.* 13, 7937-7960, doi: 10.5194/acp-13-7937-2013, 2013.

Jeong, J. I., and Park, R. J.: Effects of the meteorological variability on regional air quality in East Asia, *Atmos. Environ.*, 69, 46-55, doi: 10.1016/j.atmosenv.2012.11.061, 2013.

- Kanakidou, M., and Tsigaridis, K.: Secondary organic aerosol importance in the future atmosphere, *Atmos. Environ.*, 41, 4682-4692, doi: 10.1016/j.atmosenv.2007.03.045, 2007.
- Krewski, D., Jerrett, M., Burnett, R. T., Ma, R., Hughes, E., Shi, Y., Turner, M. C., Pope, C. A., III, Thurston, G., Calle, E. E., Thun, M. J., Beckerman, B., DeLuca, P., Finkelstein, N., Ito, K., Moore, D. K., Newbold, K. B., Ramsay, T., Ross, Z., Shin, H., and Tempalski, B.: Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality, HEI (Health Effects Institute) Research report 140, Health Effects Institute, Boston, MA, 5-114, 2009.
- Leibensperger, E. M., Mickley, L. J., and Jacob, D. J.: Sensitivity of US air quality to mid-latitude cyclone frequency and implications of 1980–2006 climate change, *Atmos. Chem. Phys.*, 8, 7075-7086, doi: 10.5194/acp-8-7075-2008, 2008.
- Liao, H., Chen, W. T., and Seinfeld, J. H.: Role of climate change in global predictions of future tropospheric ozone and aerosols, *J. Geophys. Res.*, 111, D12304, doi: 10.1029/2005jd006852, 2006.
- Liao, H., Henze, D. K., Seinfeld, J. H., Wu, S., and Mickley, L. J.: Biogenic Secondary Organic Aerosol over the United States: Comparison of Climatological Simulations with Observations, *J. Geophys. Res.*, 112, doi: 10.1029/2006JD007813, 2007.
- Liu, H., Jacob, D. J., Bey, I., and Yantosca, R. M.: Constraints from <sup>210</sup>Pb and <sup>7</sup>Be on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, *J. Geophys. Res.*, 106, 12109-12128, doi: 10.1029/2000jd900839, 2001.
- Liu, Y., Liu, R., and Chen, J. M.: Retrospective retrieval of long-term consistent global leaf area index (1981–2011) from combined AVHRR and MODIS data, *J. Geophys. Res.-Biogeo.*, 117, G04003, doi: 10.1029/2012JG002084, 2012.
- Lou, S. J, Liao, H., Zhu, B.: Impacts of aerosols on surface-layer ozone concentrations in China through heterogeneous reactions and changes in photolysis rates. *Atmos. Environ.* 85, 123-138, doi: 10.1016/j.atmosenv.2013.12.004, 2014.
- Olson, J.: World Ecosystems (WE1.4). Digital Raster Data on a 10-minute Geographic 1080x2160 grid, in: *Global Ecosystems Database, Version 1.0: Disc A.*, edited by: NOAA National Geophysical Data Center, Boulder, Colorado, 1992.
- Park, R. J., Jacob, D. J., Chin, M., and Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, *J. Geophys. Res.*, 108(D12), 4355, doi: 10.1029/2002JD003190, 2003.

- Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: implications for policy, *J. Geophys. Res.*, 109, D15204, doi: 10.1029/2003JD004473, 2004.
- Potter, P., Ramankutty, N., Bennett, E. M., and Donner, S. D.: Characterizing the Spatial Patterns of Global Fertilizer Application and Manure Production, *Earth Interact.*, 14, 1-22, doi:10.1175/2009EI288.1, 2010.
- Pye, H. O. T., Liao, H., Wu, S., Mickley, L. J., Jacob, D. J., Henze, D. K., and Seinfeld, J. H.: Effect of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States, *J. Geophys. Res.: Atmospheres*, 114, D01205, doi: 10.1029/2008JD010701, 2009.
- Pye, H. O. T., and Seinfeld, J. H.: A global perspective on aerosol from low-volatility organic compounds, *Atmos. Chem. Phys.*, 10, 4377-4401, doi: 10.5194/acp-10-4377-2010, 2010.
- Silva, R. A., West, J. J., Zhang, Y., Anenberg, S. C., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S., Skeie, R., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V., Josse, B., MacKenzie, I. A., Plummer, D., Righi, M., Stevenson, D. S., Strode, S., Szopa, S., and Zeng, G.: Global premature mortality due to anthropogenic outdoor air pollution and the contribution of past climate change, *Environ. Res. Lett.*, 8, 034005, doi: 10.1088/1748-9326/8/3/034005, 2013.
- Tai, A. P. K., Mickley, L. J., and Jacob, D. J.: Correlations between fine particulate matter ( $PM_{2.5}$ ) and meteorological variables in the United States: Implications for the sensitivity of  $PM_{2.5}$  to climate change, *Atmos. Environ.*, 44, 3976-3984, doi: 10.1016/j.atmosenv.2010.06.060, 2010.
- Tai, A. P. K., Mickley, L. J., Jacob, D. J., Leibensperger, E. M., Zhang, L., Fisher, J. A., and Pye, H. O. T.: Meteorological modes of variability for fine particulate matter ( $PM_{2.5}$ ) air quality in the United States: implications for  $PM_{2.5}$  sensitivity to climate change, *Atmos. Chem. Phys.*, 12, 3131-3145, doi: 10.5194/acp-12-3131-2012, 2012a.
- Tai, A. P. K., Mickley, L. J., and Jacob, D. J.: Impact of 2000–2050 climate change on fine particulate matter ( $PM_{2.5}$ ) air quality inferred from a multi-model analysis of meteorological modes, *Atmos. Chem. Phys.*, 12, 11329-11337, doi: 10.5194/acp-12-11329-2012, 2012b.
- Tai, A. P. K., Mickley, L. J., Heald, C. L., and Wu, S. L.: Effect of  $CO_2$  inhibition on

- biogenic isoprene emission: Implications for air quality under 2000 to 2050 changes in climate, vegetation, and land use, *Geophys. Res. Lett.*, 40, 3479-3483, doi: 10.1002/Grl.50650, 2013.
- Wang, Y., Jacob, D. J., and Logan, J. A.: Global simulation of tropospheric O<sub>3</sub>-NO<sub>x</sub>-hydrocarbon chemistry-1. Model formulation, *J. Geophys. Res.*, 103, 10713-10725, doi: 10.1029/98JD00158, 1998.
- Wang, Y., Zhang, Q. Q., He, K., Zhang, Q., and Chai, L.: Sulfate-nitrate-ammonium aerosols over China: response to 2000–2015 emission changes of sulfur dioxide, nitrogen oxides, and ammonia, *Atmos. Chem. Phys.*, 13, 2635-2652, doi: 10.5194/acp-13-2635-2013, 2013.
- Wang, Y., Yao, L., Wang, L., Liu, Z., Ji, D., Tang, G., Zhang, J., Sun, Y., Hu, B., and Xin, J.: Mechanism for the formation of the January 2013 heavy haze pollution episode over central and eastern China, *Sci. China Earth Sci.*, 57, 14-25, doi: 10.1007/s11430-013-4773-4, 2014.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.* (1967), 23, 1293-1304, doi: 10.1016/0004-6981(89)90153-4, 1989.
- [World Health Organization \(WHO\): WHO air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide-Global update 2005-Summary of risk assessment. WHO, Geneva, Switzerland, 9-13 pp., 2006.](#)
- Wu, S., Mickley, L. J., Kaplan, J. O., and Jacob, D. J.: Impacts of changes in land use and land cover on atmospheric chemistry and air quality over the 21<sup>st</sup> century, *Atmos. Chem. Phys.*, 12, 1597-1609, doi: 10.5194/acp-12-1597-2012, 2012.
- Yang, Y., Liao, H., and Lou, S.: Decadal trend and interannual variation of outflow of aerosols from East Asia: Roles of variations in meteorological parameters and emissions, *Atmos. Environ.*, 100, 141-153, doi: 10.1016/j.atmosenv.2014.11.004, 2015.
- Yienger, J. J., and Levy, H., II: Empirical model of global soil-biogenic NO<sub>x</sub> emissions, *J. Geophys. Res.*, 100, 11447-11464, doi: 10.1029/95jd00370, 1995.
- Zhang, L., Gong, S., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environ.*, 35, 549-560, doi: 10.1016/S1352-2310(00)00326-5, 2001.
- Zhang, X. Y., Wang, Y. Q., Niu, T., Zhang, X. C., Gong, S. L., Zhang, Y. M., and Sun, J. Y.: Atmospheric aerosol compositions in China: spatial/temporal variability, chemical signature, regional haze distribution and comparisons with global aerosols, *Atmos.*

638 Chem. Phys., 12, 779-799, doi: 10.5194/acp-12-779-2012, 2012.  
639 Zhu, J., Liao, H., and Li, J.: Increases in aerosol concentrations over eastern China due to the  
640 decadal-scale weakening of the East Asian summer monsoon, Geophys. Res. Lett., 39,  
641 L09809, doi: 10.1029/2012gl051428, 2012.  
642



## Table and Figure Captions

### Table 1. Summary of the simulations conducted in this study.

Figure 1. Seasonal mean surface concentrations of total PM<sub>2.5</sub>, sulfate (SO<sub>4</sub><sup>2-</sup>), nitrate (NO<sub>3</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>), organic carbon (OC), and black carbon (BC) in East Asia from the control ([CTRL]) simulation, averaged over 2007-2011.

Figure 2. Seasonal mean surface concentrations of total secondary organic aerosols (SOA), biogenic SOA, and anthropogenic SOA in East Asia from the control ([CTRL]) simulation, averaged over 2007-2011.

Figure 3. (a) Simulated vs. observed mean July-September SOA concentration. Observations are from Ding et al. (2014). Also shown is the 1:1 line (solid line) and linear fit (dashed), NMB is the normalized mean bias between simulated and observed concentrations; (b) Simulated vs. observed ratio of secondary organic carbon (SOC) to total organic carbon (OC) in China. The observed ratios are from Zhang et al. (2012). Also shown are the 1:1 line (solid line), 2:1 line and 1:2 line (dashed).

Figure 4. Simulated changes of wintertime (DJF) surface concentrations for PM<sub>2.5</sub>, sulfate, nitrate, ammonium, organic aerosol, black carbon, surface temperature, total precipitation at the ground, relative humidity, surface wind speed, planetary boundary layer depth (PBL), and cloud fraction in East Asia arising from 1980-2010 changes in climate alone ([CTRL] – [S\_CLIM]).

Figure 5. Simulated changes of summertime (JJA) surface concentrations for PM<sub>2.5</sub>, sulfate, nitrate, ammonium, organic aerosol, black carbon, surface temperature, total precipitation at the ground, relative humidity, surface wind speed, planetary boundary layer depth (PBL), and cloud fraction in East Asia arising from 1980-2010 changes in climate alone ([CTRL] – [S\_CLIM]).

Figure 6. Changes in surface secondary organic aerosol (SOA) concentration, isoprene emission, and monoterpene emission in winter (DJF) and summer (JJA) across East Asia arising from changes in climate alone ([CTRL] – [S\_CLIM]) over 1980-2010.

Figure 7. Changes in seasonal mean surface concentrations of total PM<sub>2.5</sub>, sulfate (SO<sub>4</sub><sup>2-</sup>), nitrate (NO<sub>3</sub><sup>-</sup>), ammonium (NH<sub>4</sub><sup>+</sup>) and organic carbon (OC) in East Asia arising

674 from 1980-2010 changes in land cover and land use alone ( $[CTRL] - [S\_LCLU]$ ).

675 Figure 8. Changes in surface secondary organic aerosol (SOA) concentration, isoprene  
676 emission, and monoterpene emission in winter (DJF) and summer (JJA) across  
677 East Asia arising from 1980-2010 changes in land cover and land use alone  
678 ( $[CTRL] - [S\_LCLU]$ ).

679 Figure 9. Changes in seasonal (DJF and JJA) and annual (ANN) mean surface concentrations  
680 of  $PM_{2.5}$  and SOA in East Asia resulting from the combined effects of 1980-2010  
681 changes in climate, land cover and land use ( $[CTRL] - [S\_COMB]$ ), and 1980-  
682 2010 changes in anthropogenic emissions alone ( $[CTRL] - [S\_ANTH]$ ).

683

Table 1 Summary of the simulations conducted in this study.

| Simulations | MERRA       | Vegetation parameters |                 | Anthropogenic emissions |
|-------------|-------------|-----------------------|-----------------|-------------------------|
|             | Meteorology | Land cover *          | Leaf area index |                         |
| CTRL        | 2007-2011   | 2010                  | 2010            | 2005                    |
| S_CLIM      | 1981-1985   | 2010                  | 2010            | 2005                    |
| S_LCLU      | 2007-2011   | 1980                  | 1982            | 2005                    |
| S_COMB      | 1981-1985   | 1980                  | 1982            | 2005                    |
| S_ANTH      | 2007-2011   | 2010                  | 2010            | 1985                    |

\* Land cover in terms of land or plant functional types.




Fig. 1. Seasonal mean surface concentrations of total PM<sub>2.5</sub>, sulfate ( $\text{SO}_4^{2-}$ ), nitrate ( $\text{NO}_3^-$ ), ammonium ( $\text{NH}_4^+$ ), organic carbon (OC), and black carbon (BC) in East Asia from the control ([CTRL]) simulation, averaged over 2007-2011.





Fig. 2. Seasonal mean surface concentrations of total secondary organic aerosols (SOA), biogenic SOA, and anthropogenic SOA in East Asia from the control ([CTRL]) simulation, averaged over 2007-2011.

700



701

702

703

704

705

706

707

708

Fig. 3. (a) Simulated vs. observed mean July-September SOA concentration. Observations are from Ding et al. (2014). Also shown is the 1:1 line (solid line) and linear fit (dashed), NMB is the normalized mean bias between simulated and observed concentrations; (b) Simulated vs. observed ratio of secondary organic carbon (SOC) to total organic carbon (OC) in China. The observed ratios are from Zhang et al. (2012). Also shown are the 1:1 line (solid line), 2:1 line and 1:2 line (dashed).







Fig. 4. Simulated changes of wintertime (DJF) surface concentrations for  $PM_{2.5}$ , sulfate, nitrate, ammonium, organic aerosol, black carbon, surface temperature, total precipitation at the ground, relative humidity, surface wind speed, planetary boundary layer depth (PBL), and cloud fraction in East Asia arising from 1980-2010 changes in climate alone ( $[CTRL] - [S_{CLIM}]$ ).



718

719

720

721

722

723

724

Fig. 5. Simulated changes of summertime (JJA) surface concentrations for PM<sub>2.5</sub>, sulfate, nitrate, ammonium, organic aerosol, black carbon, surface temperature, total precipitation at the ground, relative humidity, surface wind speed, planetary boundary layer depth (PBL), and cloud fraction in East Asia arising from 1980-2010 changes in climate alone ([CTRL] – [S\_CLIM]).




Fig. 6. Changes in surface secondary organic aerosol (SOA) concentration, isoprene emission, and monoterpene emission in winter (DJF) and summer (JJA) across East Asia arising from changes in climate alone ( $[CTRL] - [S\_CLIM]$ ) over 1980-2010.






Fig. 7. Changes in seasonal mean surface concentrations of total  $PM_{2.5}$ , sulfate ( $SO_4^{2-}$ ), nitrate ( $NO_3^-$ ), ammonium ( $NH_4^+$ ) and organic carbon (OC) in East Asia arising from 1980-2010 changes in land cover and land use alone ( $[CTRL] - [S\_LCLU]$ ).




Fig. 8. Changes in surface secondary organic aerosol (SOA) concentration, isoprene emission, and monoterpene emission in winter (DJF) and summer (JJA) across East Asia arising from 1980-2010 changes in land cover and land use alone ( $[CTRL] - [S\_LCLU]$ ).




Fig. 9. Changes in seasonal (DJF and JJA) and annual (ANN) mean surface concentrations of  $PM_{2.5}$  and SOA in East Asia resulting from the combined effects of 1980-2010 changes in climate, land cover and land use ( $[CTRL] - [S\_COMB]$ ), and 1980-2010 changes in anthropogenic emissions alone ( $[CTRL] - [S\_ANTI]$ ).