Seasonal Variability of Stratospheric Methane: Implications for Constraining Tropospheric Methane Budgets Using Total Column Observations

K. M. Saad¹, D. Wunch^{1,2}, N. M. Deutscher^{3,4}, D. W. T. Griffith³, F. Hase⁵, M. De Mazière⁶, J. Notholt⁴, D. F. Pollard⁷, C. M. Roehl¹, M. Schneider⁵, R. Sussmann⁸, T. Warneke⁴, and P. O. Wennberg¹

¹California Institute of Technology, Pasadena, California, USA
²University of Toronto, Toronto, Ontario, Canada
³University of Wollongong, Wollongong, NSW, Australia
⁴University of Bremen, Bremen, Germany
⁵Karlsruhe Institute of Technology, IMK-ASF, Karlsruhe, Germany
⁶Royal Belgian Institute for Space Aeronomy, Brussels, Belgium
⁷National Institute of Water and Atmospheric Research, Omakau, New Zealand
⁸Karlsruhe Institute of Technology, IMK-IFU, Garmisch-Partenkirchen, Germany

Correspondence to: K. M. Saad (katsaad@caltech.edu)

Abstract. Global and regional methane budgets are markedly uncertain. Conventionally, estimates of methane sources are derived by bridging emissions inventories with atmospheric observations employing chemical transport models. The accuracy of this approach requires correctly simulating advection and chemical loss such that modeled methane concentrations scale with surface fluxes. When total column measurements are assimilated into this framework, modeled stratospheric methane

- 5 introduces additional potential for error. To evaluate the impact of such errors, we compare Total Carbon Column Observing Network (TCCON) and GEOS-Chem total and tropospheric column-averaged dry-air mole fractions of methane. We find that the model's stratospheric contribution to the total column is insensitive to perturbations to the seasonality or distribution of *tropospheric* emissions or loss. In the Northern Hemisphere, we identify disagreement between the measured and modeled stratospheric contribution, which increases as the tropopause altitude decreases, and a temporal phase lag in the model's tro-
- 10 pospheric seasonality driven by transport errors. Within the context of GEOS-Chem, we find that the errors in tropospheric advection partially compensate for the stratospheric methane errors, masking inconsistencies between the modeled and measured tropospheric methane. These seasonally-varying errors alias into source attributions resulting from model inversions. In particular, we suggest that the tropospheric phase lag error leads to large misdiagnoses of wetland emissions in the high latitudes of the Northern Hemisphere.

15 1 Introduction

Identifying the processes that have driven changes in atmospheric methane (CH_4) , a potent radiative forcing agent and major driver of tropospheric oxidant budgets, is critical for understanding future impacts on the climate system. Methane's growth rate, which had been decreasing through the 1990s from about 10 to 0 ppb per year, began to increase again in 2006 and over

the past decade has averaged 5 ppb per year (Dlugokencky et al., 2011). Developing robust constraints on the global CH_4 budget is integral for understanding which processes produced these decadal trends (e.g., Bergamaschi et al., 2013; Wecht et al., 2014a, b; Turner et al., 2015).

- One common approach to quantifying changes in the spatial distribution of sources are atmospheric inversions, which in-5 corporate surface fluxes estimated by bottom-up inventories as boundary conditions for a chemical transport model (CTM). The modeled CH_4 concentrations are compared to observations within associated grid boxes, and prior emissions are scaled to minimize differences with measured dry-air mole fractions (DMFs), producing posterior estimates. The accuracy of these optimized emissions depends on how well the CTM simulates atmospheric transport and CH_4 sinks, which are generally prescribed.
- Pressure-weighted total column-averaged DMFs (X_{gas}) provide a relatively new constraint and have previously been shown to improve estimates of regional and interhemispheric gradients in trace gases (Yang et al., 2007). Infrared spectrometers can measure CH₄ DMFs (X_{CH_4}) from ground-based sites, such as those in the Total Carbon Column Observing Network (TCCON) and Network for the Detection of Atmospheric Composition Change (NDACC), and satellites, including SCanning Imaging Absorption spectroMeter for Atmospheric CartograpHY (SCIAMACHY) (Bergamaschi et al., 2007), Green-
- 15 house gases Observing SATellite (GOSAT) (Parker et al., 2011), and the upcoming TROPOspheric Monitoring Instrument (TROPOMI) (Butz et al., 2012). These observations complement surface measurements because they add information about the vertically-averaged profile and are sensitive in the free troposphere (Yang et al., 2007). Additionally, they complement aircraft observations by measuring trace gases at higher temporal frequency, although they share the limitation of not measuring in inclement weather. Satellite measurements add global coverage that can fill in gaps where in situ observations are sparse.
- Fraser et al. (2013) found assimilating GOSAT CH_4 columns into the GEOS-Chem CTM with an ensemble Kalman filter reduced posterior emissions uncertainties by 9 - 48% for individual source categories and by more than three times those of inversions that only assimilated surface data for most regions. Wecht et al. (2014b) determined from their analysis of observing system simulation experiments (OSSEs) that TROPOMI's daily frequency and global coverage performs similarly to aircraft campaigns on sub-regional scales and could provide a constraint on California's CH_4 emissions similar to CalNex aircraft
- 25 observations (Santoni et al., 2014; Gentner et al., 2014).

Incorporating total columns into modeling assessments can also be used to diagnose systematic issues with model transport. For example, comparing carbon dioxide (CO₂) from TCCON and TransCom (Baker et al., 2006), Yang et al. (2007) found that most models included in the comparison lack sufficiently strong vertical exchange between the planetary boundary layer (PBL) and the free troposphere, thereby dampening the seasonal cycle amplitude of X_{CO_2} . The limitations of models to accurately

30 represent vertical transport can lead to radically different spatial distributions of fluxes; Stephens et al. (2007) found, for example, that the northern terrestrial carbon land sink and tropical emissions were overestimated by 0.9 and 1.7 PgC·year⁻¹, respectively, when comparing models to aircraft CO_2 profiles. More recent studies attribute to model transport errors the tendency of simulated CH_4 in the Southern Hemisphere to be higher at the surface than the free troposphere, in contrast with measurements (Fraser et al., 2011; Patra et al., 2011). Tropospheric CH_4 typically does not vary radically with height above the PBL; above the tropopause, however, the vertical profile of CH_4 exhibits a rapid decline with altitude as a result of its oxidation and the lack of any source beyond advection from the troposphere. Fluctuations in stratospheric dynamics, including the height of the tropopause, change the contribution of the stratosphere to the total column. CH_4 profiles with similar tropospheric values can thus have significant differences in X_{CH_4} (Saad et al., 2014; Washenfelder et al., 2003; Wang et al., 2014).

Provided that simulations replicate seasonal and zonal variability of stratospheric CH_4 loss, tropopause heights, and vertical exchange across the upper troposphere and lower stratosphere (UTLS), posterior flux estimates from inversions incorporating X_{CH_4} measurements would not be sensitive to stratospheric processes. However, most models do not accurately represent stratospheric transport, producing low age of air values and zonal gradients in the subtropical lower stratosphere that are

- 10 less steep than observations (Waugh and Hall, 2002). The TransCom- CH_4 CTM intercomparison assessment of transport using sulfur hexafluoride SF_6 showed a strong correlation between the stratosphere-troposphere exchange (STE) rate and the model's CH_4 budget and a weaker correlation between the CH_4 growth rate and vertical gradient in the model's equatorial lower stratosphere (Patra et al., 2011). These forward model dependencies of CH_4 concentrations on vertical transport, both within the troposphere and across the tropopause, have the potential to introduce substantial errors in atmospheric inversions.
- 15 As temporal and spatial biases in a model's vertical profile will alias into posterior emissions, inversions that incorporate total column measurements must ensure that the stratosphere is sufficiently well described so as to not introduce spurious seasonal, zonal and interhemispheric trends in CH_4 concentrations and consequently emissions.

In this analysis, we identify systematic model errors in the seasonal cycle and spatial distribution of CH_4 DMFs by comparing TCCON total and tropospheric columns (Saad et al., 2014) to vertically integrated profiles derived from the GEOS-Chem

20 CTM (Bey et al., 2001; Wang et al., 2004; Wecht et al., 2014a). We assess the impact of errors in the characterization of stratospheric processes on the assimilation of X_{CH_4} and resulting posterior emissions estimates. In Section 2 we describe the TCCON column measurements and GEOS-Chem set up and characteristics. In Section 3 we present the results of the measurement-model comparison. In Section 4 we compare the base case simulation to one in which emissions do not vary within each year and quantify the sensitivity of source attribution of the biggest seasonal emissions sector, wetlands, to the tropospheric seasonal delay.

2 Methods

5

2.1 Tropospheric Methane Columns

TCCON has provided precise measurements of X_{CH4} and other atmospheric trace gases for over ten years (Wunch et al., 2011a, 2015). Developed to address open questions in carbon cycle science, the earliest sites are located in Park Falls, Wisconsin,
United States and Lauder, New Zealand at about 45° North and South, respectively. Since 2004, the ground-based network of Fourier transform spectrometers has expanded greatly. X_{CH4} are processed with the current version of the TCCON software, GGG2014, to be consistent, and thereby comparable, across sites. Total column retrievals are generated with the GFIT nonlinear least-squares fitting algorithm, which calculates the best spectral fit of the solar absorption signal to an a priori vertical profile



Figure 1. Map of TCCON sites used in this analysis. Site colors are on a spectral color scale in order of latitude, with Northern Hemisphere sites designated by cool colors and Southern Hemisphere sites designated by warm colors.

and outputs a scaling factor. The pressure-weighted integration of the scaled a priori profile produces column abundances, which are then divided by the dry air column, calculated using concurrently retrieved oxygen (O_2) columns (Wunch et al., 2010, 2011a, 2015). Trace gas a priori profiles are derived with empirical models, which are generated incorporating aircraft and balloon in situ and satellite measurements (see Wunch et al., 2015, for a complete list), and for CH₄ include a secular increase

- 5 of 0.3% per year and an interhemispheric gradient in the altitude dependence of the vertical profiles (Toon and Wunch, 2014). These models are fit to daily noontime National Centers for Environmental Protection and National Center for Atmospheric Research (NCEP/NCAR) reanalysis pressure grids (Kalnay et al., 1996), interpolated to the surface pressure measured realtime on site. Because the profile of CH_4 drops off rapidly in the stratosphere, the accuracy of the a priori shape, and thus the retrieved column, depends on correctly determining the tropopause.
- 10 Tropospheric columns have been shown to represent the magnitude and seasonality of in situ measurements (Saad et al., 2014; Washenfelder et al., 2003; Wang et al., 2014). The tropospheric CH_4 column-averaged DMFs ($X_{CH_4}^t$) are derived by the HF-proxy method described in Saad et al. (2014), which uses the relationship between CH_4 and HF in the stratosphere, derived from ACE-FTS satellite measurements (Bernath, 2005; De Mazière et al., 2008; Mahieu et al., 2008; Waymark et al., 2014), to calculate and remove the stratospheric contribution to X_{CH_4} . The $X_{CH_4}^t$ used in this analysis have been processed consistently
- 15 with the GGG2014 TCCON products, with airmass dependence and calibration factors calculated for and applied to $X_{CH_4}^t$ (Wunch et al., 2010, 2015). Additional details about the tropospheric CH₄ measurements can be found in Appendix A.

With the exception of Eureka and Sodankylä, which are highly influenced by the stratospheric polar vortex, all TCCON sites that provide measurements before December 2011 are included in this analysis (Fig. 1). Table 1 lists locations and data collection start dates for each of the sites.

Site	Latitude (°)	Longitude (°)	Elevation (km)	Start Date	Location	Data Reference
Bialystok	53.2	23.0	0.18	Mar 2009	Bialystok, Poland	Deutscher et al. (2014)
Bremen	53.1	8.9	0.03	Jan 2007	Bremen, Germany	Notholt et al. (2014)
Karlsruhe	49.1	8.4	0.11	Apr 2010	Karlsruhe, Germany	Hase et al. (2014)
Orleans	48.0	2.1	0.13	Aug 2009	Orleans, France	Warneke et al. (2014)
Garmisch	47.5	11.1	0.75	Jul 2007	Garmisch, Germany	Sussmann and Rettinger (2014)
Park Falls	45.9	-90.3	0.47	Jan 2005	Park Falls, WI, USA	Wennberg et al. (2014b)
Lamont	36.6	-97.5	0.32	Jul 2008	Lamont, OK, USA	Wennberg et al. (2014c)
JPL	34.2	-118.2	0.39	Jul 2007	Pasadena, CA, USA	Wennberg et al. (2014d, a)
Saga	33.2	130.3	0.01	Jul 2011	Saga, Japan	Kawakami et al. (2014)
Izaña	28.3	-16.5	2.37	May 2007	Tenerife, Canary Islands	Blumenstock et al. (2014)
Darwin	-12.4	130.9	0.03	Aug 2005	Darwin, Australia	Griffith et al. (2014a)
Réunion Island	-20.9	55.5	0.09	Sep 2011	Saint-Denis, Réunion	De Maziere et al. (2014)
Wollongong	-34.4	150.9	0.03	Jun 2008	Wollongong, Australia	Griffith et al. (2014b)
Lauder	-45.0	169.7	0.37	Jan 2005	Lauder, New Zealand	Sherlock et al. (2014a, b)

Table 1. TCCON sites, coordinates, altitudes, start date of measurements and locations used in this analysis.

2.2 GEOS-Chem Model

Model comparisons use the offline CH₄ GEOS-Chem version 9.02 at 4° × 5° horizontal resolution on a reduced vertical grid (47L). CH₄ loss is calculated on 60 minute intervals and is set by annually-invariable monthly 3D fields: hydroxyl radical (OH) concentrations in the troposphere (Park et al., 2004) and parameterized CH₄ loss rates per unit volume in the stratosphere (Considine et al., 2008; Allen et al., 2010; Murray et al., 2012). Emissions are released at 60-minute time steps and are provided by the GEOS-Chem development team for 10 sectors: gas and oil, coal, livestock, waste, biofuel, and other anthropogenic annual emissions from EDGAR v4.2 (European Commission Joint Research Centre, Netherlands Environmental Assessment Agency, 2011; Wecht et al., 2014a); other natural annual emissions from (Fung et al., 1991); rice agriculture (European Commission Joint Research Centre, Netherlands Environmental Assessment Agency, 2011) and wetland (PickettHeaps et al., 2011) monthly emissions, which incorporate GEOS5 annual and monthly mean soil moisture values; and biomass

10 Heaps et al., 2011) monthly emissions, which incorporate GEOS5 annual and monthly mean soil moisture values; and biomass burning daily emission from GFED3 estimates (Mu et al., 2011; van der Werf et al., 2010). Loss via soil absorption (Fung et al., 1991), set annually, is subtracted from the total emissions at each time step.

2.2.1 Model Set Up

We initialized zonal CH₄ distributions with GGG2014 data version a priori profiles (Toon and Wunch, 2014) produced at
horizontal grid centers, which we adjusted vertically to match the zonally averaged daily mean model's tropopause, derived from the National Aeronautics and Space Administration Global Modeling and Assimilation Office (NASA/GMAO) Goddard

Earth Observing System Model, Version 5 (GEOS5). The model was run from December 2003, the first month in which GEOS5 meteorological data were available, to June 2004, the beginning of the TCCON time series; we then ran the model repeatedly over the June 2004-May 2005 time frame, which allowed us to make comparisons with the TCCON data at Park Falls and Lauder, until CH_4 concentrations reached equilibrium. A number of perturbation experiments were run in this way

to quantify the sensitivity of CH_4 distribution and seasonality to the offline OH fields, prescribed emissions, and tropopause 5 levels (Table 2). These model experiments are described in greater detail in Appendix B1.

Using CH_4 fields for 1 January 2005 from the equilibrium simulation as initial conditions, model daily mean CH_4 mole fractions were computed through 2011. These were converted to dry mole fractions, as described in Appendix B2. In addition to the default emissions scheme, an aseasonal simulation setup, in which rice, wetland, and biomass burning emissions were

- disabled and aseasonal emissions scaled up such that total annual zonal fluxes approximate those in the base simulation, was 10 similarly run to equilibrium and used as initial conditions for the 2005-2011 run. The model infrastructure posed difficulties for setting the seasonally-varying fluxes constant throughout each year; thus we implement this scaling technique as an alternative to assess first-order impacts of emission seasonality. The resulting changes to the spatial distribution of CH_4 emissions are shown in Fig. 13.
- 15 For comparisons with column measurements, model vertical profiles were smoothed with corresponding TCCON CH_4 averaging kernels, interpolated for the daily mean solar zenith angles, and prior profiles, scaled with daily median scaling factors, following the methodology in Rodgers and Connor (2003) and Wunch et al. (2010). Averaging kernels and prior profiles were interpolated to the model's pressure grid, and all terms in the smoothing equation were interpolated to daily mean surface pressures measured at each site. Tropospheric columns were integrated in the same manner as the total columns up to
- the grid level completely below the daily mean tropopause, consistent with how GEOS-Chem partitions the atmosphere in the 20 offline CH_4 simulation. To test the dependence of our results on the chosen vertical integration level, tropospheric columns were also calculated assuming the tropopause was one and two grid cells above this level. While $X_{CH_4}^t$ changed slightly, by a median of about 1 and 5 ppb for a one and two-level increase respectively, shifting the tropopause did not alter the findings discussed in this paper. A description of the model smoothing methodology and assumptions is provided in Appendix B3. The stratospheric contribution to the total column, which is calculated as the residual between the $X_{CH_4}^t$ and X_{CH_4} , is the amount

by which the stratosphere attenuates X_{CH_4} via stratospheric loss and transport (see Appendix C for the derivation).

25

Table 2.	Sensitivity	Experiments
I abit -	Sensitivity	Experimento

Run Name	Description	CH ₄ Lifetime (years)	Final CH ₄ Burden (Tg)	
Base	Default OH and Emissions	9.55	4825	
Aseasonal	Constant Monthly Emission Rates	9.57	4872	
Updated OH	Monthly OH fields from Standard Chemistry + Biogenic VOCs	8.53	4828	



Figure 2. Seasonality of the difference between base and aseasonal CH_4 for tropospheric, total and stratospheric contribution to total columns. Site colors are as in Fig. 1.

2.2.2 Model Features

The seasonal amplitude of the differences between base and aseasonal simulations are small-within ± 4 ppb-for all vertical levels in the Southern Hemisphere (Fig. 2). In the Northern Hemisphere, however, the difference is much larger and primarily impacts the troposphere, where it varies between -10 and +13 ppb. The insensitivity of the stratosphere to the seasonality of emissions is due to the common source of stratospheric air in the tropics (Boering et al., 1995) and the loss of seasonal information as the age of air increases (Mote et al., 1996).

Due to the relatively short photochemical lifetime of CH_4 in the stratosphere, about 22 months in the base simulation, stratospheric CH_4 concentrations stabilize much more quickly than in the troposphere (Fig. 3a). This rapid response time of the stratosphere occurs regardless of perturbations to the troposphere, such as the seasonality of emissions (Fig. 3b) or tropo-



Figure 3. Smoothed daily mean $X_{CH_4}^t$ and stratospheric contribution to X_{CH_4} at Park Falls (blue) and Lauder (red) for (a) base equilibrium simulation and the difference between the base and (b) aseasonal and (c) updated OH simulations.

spheric OH fields (Fig. 3c). In both hemispheres the differences between the base and experimental simulations asymptotically approach steady state with seasonal variability over a decade in the troposphere, but oscillate seasonally around a constant mean in the stratosphere. Stratospheric differences between simulations are considerably smaller than the seasonal amplitude of the base run: within six and one ppb, respectively, versus a seasonal range of 30 ppb at Park Falls. By contrast, $X_{CH_4}^t$ have differences within 30 and 10 ppb, respectively, versus a seasonal range of 20 ppb at Park Falls. The stratosphere at Lauder is even less sensitive to tropospheric perturbations.

3 Measurement-Model Comparison

5

The TCCON daily median and GEOS-Chem daily mean CH₄ column-averaged DMFs demonstrate a strong interhemispheric difference for X^t_{CH4} and X_{CH4} in both the base and aseasonal simulations (Fig. 4). The Northern Hemisphere X^t_{CH4} slope
deviates from the one-to-one line more than the X_{CH4} slope (0.60±0.02 versus 0.86±0.03), and the correlation coefficients are equivalent (R² = 0.41), which indicates that the poorer agreement between measurements and models in the troposphere drive the scatter in the total column.

The stratospheric contribution comparison between TCCON and the base simulation for the Northern Hemisphere sites has an equivalent slope (0.60 ± 0.1) and higher correlation coefficient $(R^2 = 0.68)$ compared to $X_{CH_4}^t$ (Fig. 4c). GEOS-Chem's

15 larger stratospheric contribution to the total column, coupled with lower tropospheric values, depresses X_{CH_4} . Because this effect on X_{CH_4} occurs more at higher latitudes, zonal errors in the model's stratosphere balances those in the troposphere. The result is better measurement-model agreement in the total columns.



Figure 4. Daily median TCCON and smoothed daily mean GEOS-Chem base (top) and aseasonal (bottom) DMFs for (a) $X_{CH_4}^t$, (b) X_{CH_4} , and (c) stratospheric contribution. Site colors are as in Fig. 1. Northern Hemisphere least squares regression equations are in the top left, and Southern Hemisphere least squares regression equations are in the bottom right of each plot. Dashed lines mark the one-to-one lines.

The aseasonal simulation produces lower slopes and correlation coefficients for, $X_{CH_4}^t$ (slope=0.42±0.02, $R^2 = 0.32$), X_{CH_4} (slope=0.60±0.03, $R^2 = 0.26$), and the stratospheric contribution (slope=0.52±0.01, $R^2 = 0.66$) in the Northern Hemisphere. Removing the seasonality of emissions increases both measurement-model differences and scatter, as we would expect given the seasonality of Northern Hemisphere emissions noted in bottom-up studies (Kirschke et al., 2013). The aseasonal simulation also reduces the offset between TCCON and GEOS-Chem, whereby modeled $X_{CH_4}^t$ and X_{CH_4} are systematically low. TransCom-CH₄ showed that GEOS-Chem CH₄ concentrations tend to be lower than the model median, and much lower than the range of other models when using the same OH fields (Patra et al., 2011). The aseasonal emissions used in this analysis likely reduce this documented imbalance with the model's tropospheric OH fields.

5

The X_{CH_4} and $X_{CH_4}^t$ regression equations across Southern Hemisphere sites are nearly equivalent, which suggests that the 10 Southern Hemisphere is not as impacted by the STE errors as the Northern Hemisphere. This consistency between X_{CH_4} and $X_{CH_4}^t$ could also be a function of the zonal dependence of the stratospheric error: whereas more than half of the Northern Hemisphere sites are north of 45°N, the most poleward site in the Southern Hemisphere is located at 45°S. The increased scatter associated with the slightly lower $X_{CH_4}^t R^2$ value of 0.63, compared to the $X_{CH_4} R^2$ value of 0.88, does indicate that the Southern Hemisphere is not exempt from model errors associated with emissions, the OH distribution, or transport. The lower

15 $X_{CH_4}^t$ slope of the aseasonal simulation (1.1 versus 1.3) illustrates the influence of emissions: removing their seasonality leads to better measurement-model agreement, evidenced by a slope closer to both the one-to-one line and the zero-intercept. We

hypothesize that either the seasonality of Southern Hemispheric emissions is too strong or, more likely, errors in the Northern Hemispheric seasonality of emissions drive measurement-model mismatch in the Southern Hemisphere via interhemispheric transport. If this effect was solely due to a changed emissions distribution, we would expect the X_{CH_4} slope to also change for the Southern Hemisphere sites, if only slightly; instead the slope is equivalent to the base simulation $X_{CH_4}^t$ and X_{CH_4} slopes, and $R^2 = 0.87$, only marginally less than the base simulation X_{CH_4} correlation coefficient.

5

20

The stratospheric contribution regression equations differ only slightly between the base and aseasonal simulations: $(0.64 \pm$ (0.02)x + 14, $R^2 = 0.68$, versus $(0.62 \pm 0.02)x + 15$, $R^2 = 0.67$. The insensitivity of both the stratospheric contribution and the total columns in the Southern Hemisphere to perturbations in the seasonality of tropospheric emissions could be driven by the smaller vertical gradient across the UTLS that results from the influence of Northern Hemispheric air both in the free

- troposphere (Fraser et al., 2011) and the stratosphere (Boering et al., 1995). This effect would also support the interpretation of 10 Northern Hemispheric emission errors driving disagreement between observations and the model in the Southern Hemisphere. In the troposphere, CH₄ increases from south to north; the stratospheric contribution of CH₄, however, increases from the equator to the poles due to the zonal gradient in troppause height. In the Northern Hemisphere total column, the zonal gradient largely disappears: at high latitudes, the larger tropospheric emissions balances the larger stratospheric contribution.
- 15 By contrast, zonal gradients in the Southern Hemisphere troposphere and stratosphere are additive, and greater south to north differences are apparent in the total column.

Figure 5 illustrates how the model differs from ACE-FTS CH_4 measurements in the stratosphere over boreal spring (March-April-May) and fall (September-October-November). Except above the tropical troppause, CH_4 is considerably lower in the ACE-FTS climatology (v. 2.2, Jones et al., 2012) compared to GEOS-Chem. The difference varies both with altitude and latitude, especially in the Northern spring poleward of 40° N. The vertical gradient is the least pronounced in Lauder, where the stratospheric contributions of TCCON and GEOS-Chem fall most closely to the one-to-one line (Fig. 4). The low CH_4 in the

tropical mid and upper stratosphere in GEOS-Chem could be a result of too weak vertical ascent to the stratosphere; however, the ACE-FTS data gaps in the tropical troposphere make this hypothesis difficult to test.

3.1 **Dependence on Tropopause Height**

- In the Northern Hemisphere, the measurement-model mismatch of the stratospheric contribution increases as the tropopause 25 altitude shifts downward (Fig. 6). As the model's stratospheric portion of the pressure-weighted total column increases, the error in stratospheric CH₄ is amplified, causing a larger disagreement with measurements. Because the tropopause height decreases with latitude, and this gradient increases during winter and spring, this introduces both zonal and seasonal biases. The disagreement exhibits a large spread for relatively few tropopause pressure heights because the model's effective tropopause.
- that is, the pressure level at which the model divides the troposphere from the stratosphere in GEOS-Chem, is defined at 30 discrete grid level pressure boundaries.

The tropospheric mismatch ($\Delta X_{CH_4}^t$), by contrast, decreases with tropopause height for the majority of days and exhibits a much weaker correlation to tropopause height, 0.099 versus 0.22 for the stratospheric contribution. Thus, as expected, the tropopause height explains less of the variance in the measurement-model mismatch in $X_{CH_4}^t$: the upper troposphere is gener-



Figure 5. Zonally averaged ACE minus GEOS-Chem climatological CH_4 mole fractions for boreal spring and fall. Black line represents the mean zonal tropopause level. Site colors of squares on the x-axis are as in Fig. 1.



Figure 6. TCCON minus GEOS-Chem CH_4 column-averaged DMFs as a function of the effective GEOS-Chem tropopause height, shown for Northern Hemisphere sites. Site colors are as in Fig. 1.

ally well-mixed, and chemical loss does not vary with altitude as much as in the lower stratosphere. This weaker relationship also demonstrates that the choice of tropopause used in the tropospheric profile integration does not strongly impact $\Delta X_{CH_4}^t$.

The relationship between $\Delta X_{CH_4}^t$ and tropopause height has a clear zonal component that indicates that the correlation is instead a result of another parameter that varies with latitude. The tropospheric slope is dominated by high-latitude sites; the subtropical sites exhibit a much weaker correlation. At Izaña which is in the sub-tropics at an altitude of 2.4 km, the correlation

5 subtropical sites exhibit a much weaker correlation. At Izaña, which is in the sub-tropics at an altitude of 2.4 km, the correlation between $\Delta X_{CH_4}^t$ and tropopause position is weak: the slope of -0.035 ± 0.03 is nearly flat within error, and R^2 is 0.025. By contrast, the stratospheric relationship at Izaña corresponds more closely with the other Northern Hemisphere sites: the slope is -0.088 ± 0.02 , and $R^2 = 0.36$.



Figure 7. Detrended seasonality of TCCON (black diamonds), GEOS-Chem base (red circles), and GEOS-Chem aseasonal (blue squares) CH_4 column-averaged DMFs, averaged across Northern Hemisphere sites, excluding Saga, which has less than one year of measurements prior to 2012. Error bars denote the 1σ standard deviation across sites.

3.2 Seasonal Agreement

5

The tropospheric difference between TCCON and GEOS-Chem, $\Delta X_{CH_4}^t$, has a periodic trend indicating that the model error has a strong seasonal component in the troposphere. To isolate stable seasonal patterns from the cumulative influence of emissions, we calculate the detrended seasonal mean column-averaged DMFs for each site. In the Southern Hemisphere, the measurements and model agree well. Across the Northern Hemisphere sites, however, the seasonality differs (Fig. 7). The seasonal amplitude of GEOS-Chem $X_{CH_4}^t$ is about equal to that of TCCON, but the TCCON $X_{CH_4}^t$ seasonal minimum is in

Seasonal amplitude of GEOS-Chem X_{CH_4} is about equal to that of TCCON, but the TCCON X_{CH_4} seasonal minimum is in June/July while the GEOS-Chem seasonal minimum is in September/October. Additionally, while TCCON $X_{CH_4}^t$ begins to decrease in January, GEOS-Chem shows some persistence into the spring.

The seasonal delay also appears in comparisons of GEOS-Chem surface CH_4 with National Oceanic and Atmospheric Administration (NOAA) surface flask measurements at the LEF site in Park Falls (Fig. 8). The seasonality of GEOS-Chem's surface is regulated more by emissions than transport: CH_4 peaks in the summer, when wetland emissions are highest (Fig. 10). This contrasts with the flask measurements, which reach a minimum in the summer (Fig. 8). The seasonality covaries remarkably closely with respect to other features: the late winter decrease, spring persistence, and local minimum in October. The spring plateau lasts twice as long as seen in observations, however, and matches $X_{CH_4}^t$, indicating that feature is not the result

15 of vertical transport between the PBL and free troposphere.

Not surprisingly, a time lag does not occur in the stratosphere; the TCCON stratospheric seasonal amplitude is less than half but in phase with that of GEOS-Chem (Fig. 7). The vertical inconsistency of the seasonality produces unusual features in the model total column. From January through April, the TCCON and GEOS-Chem X_{CH_4} are consistent because the model's bias in the troposphere is balanced by the larger stratospheric contribution. Starting in May, however, the model diverges from

20 the measurements as the higher tropopause limits the stratosphere's influence, and the phase lag in the troposphere dominates. This balancing effect is also demonstrated by the greater variance across sites in the model $X_{CH_4}^t$ and stratospheric contribution compared to measurements, but about the same variance in X_{CH_4} .



Figure 8. NOAA surface flask (black) and GEOS-Chem surface level (red) seasonality of CH₄ DMFs over 2005-2011 at Park Falls, WI, USA and Baring Head, NZ. Lower and upper bounds denote the 25th and 75th percentiles, respectively, of detrended data for each month.

For the aseasonal simulation, the tropospheric seasonal cycle amplitude and variance across sites increase (Fig. 7). The greatest model differences, from August through October, are a result of dampening the large wetland fluxes in the base simulation that balance higher OH concentrations. The seasonal amplitude does not increase as drastically in the sub-tropics, where the total emissions are not as impacted by seasonally-varying sources, leading to the greater variance across sites. The second largest difference between simulation amplitudes occurs in the spring, and OH loss could potentially be driving in these months also. The aseasonal simulation spreads the wetland fluxes so as to introduce emissions in the winter and spring, when the OH concentrations are lowest. Another possibility is that the model could be subject to errors that are in phase with the base simulation seasonal emissions, which would then have an ameliorating effect that produces the reasonable seasonal cycle amplitude. The stratospheric contribution does not change, however, further demonstrating that the stratosphere is insensitive to perturbations to Northern Hemisphere emissions.

10

5

The impact of a static stratosphere and changing troposphere is to make the seasonality of the aseasonal simulation X_{CH_4} bimodal: the October local minimum in the base simulation becomes a fall absolute minimum. The aseasonal X_{CH_4} agrees with TCCON in late winter, masking the greater disagreement in the troposphere. Notably, the main tropospheric features of the

base simulation, the seasonal phase lag and spring persistence, are still apparent. Thus, the seasonality of emissions prescribed in the forward model is not the driver of the discrepancies between measurement and model $X_{CH_4}^t$ seasonalities. OH is not likely the driver of these features, as the Northern Hemisphere phase shift also occurs in simulations performed with large changes in OH (Fig. 15, in Appendix B1). Transport is thus the most likely driver of these tropospheric trends in the model.

5 4 Discussion

The stratospheric insensitivity to changes in emissions and tropospheric loss has significant implications for flux inversions. Model inversions use the sensitivity of trace gas concentrations at a given location to perturbations of different emission sources to adjust those emissions so as to match observations at that location. The response of modeled CH_4 DMFs to changing emissions depends on the model's transport and chemical loss, as well as assumptions about the seasonal and spatial distribution

- 10 of emissions relative to each other. Thus the model sensitivity kernel, the linear operator that maps emissions to CH_4 concentrations, implicitly includes uncertainties in these terms. The model's stratospheric response to emission perturbations differs from that of the troposphere and is subject to different transport and loss errors. Because the tropospheric transport errors covary with emissions, they alias into the resulting source attribution.
- Comparing measurement and model stratospheric CH₄ as a fraction of the total column provides a normalized comparison
 that isolates differences in the vertical structure from those caused by initial conditions and unbalanced sources and sinks.
 Figure 9 illustrates the error associated with the normalized stratospheric column and the associated stratospheric contribution to X_{CH4} at Park Falls. Although the stratosphere accounts for less than 30% of X_{CH4}, a relatively small error can produce significant seasonal differences; the springtime error of 4.5 × 10¹⁷ molec·cm⁻² (23 ppb) is more than twice the seasonal cycle amplitude. Winter and spring are also when X^t_{CH4} is least sensitive to seasonal emissions; by contrast, the error is about 15
 ppb in the summer, when seasonal emissions have the greatest influence (Fig. 9, top panel). The seasonality of the stratospheric error will therefore distort the inversion mechanism and thus posterior emissions estimates.

Additional bias is introduced by differences in the seasonal patterns of $\Delta X_{CH_4}^t$ and ΔX_{CH_4} . Wetlands are the largest seasonal source of CH₄ in models and the largest natural source in flux inventories, and their emissions are very uncertain: estimates range between 142 and 284 TgC·year⁻¹ for the 2000-2009 time period (Kirschke et al., 2013). A priori GEOS-

- 25 Chem CH_4 emissions from northern high-latitude wetlands are extremely variable, with large fluxes in June, July and August, moderate fluxes in May and September, and almost no fluxes the remainder of the year (Fig. 10a). Surface CH_4 concentrations in models depend on the assumed seasonally-varying emissions. Patra et al. (2011) found that correlations between the seasonal cycles of the forward model averages and in situ observations of CH_4 DMFs at the surface varied for a given site by up to 0.78 ± 0.4 depending on wetland and biomass burning fields used. Model inversions that scale emissions in a given grid box
- 30 based on the incorrect seasonality will invariably change the posterior attribution of seasonal emissions. Fraser et al. (2013) found that optimized wetland emissions from inversions that assimilate surface data only are smaller than the priors, while those from inversions that assimilate GOSAT total columns are larger, even if surface measurements are also assimilated. From



Figure 9. Top: Seasonally-averaged fraction of model emissions from seasonally-varying sources, north of 40° N. Bottom: Seasonally-averaged normalized model stratospheric column error (teal) and the difference between base and aseasonal simulation tropospheric columns (orange) at Park Falls.

this we infer that the transport errors in the model's free troposphere lead to an "optimization" of the prior fluxes of opposite sign to that of the emission errors that the inversion attempts to correct.

A two to three-month shift in the phase of the $X_{CH_4}^t$ seasonality will produce a strong under- or overestimation of posterior wetland fluxes in late spring through early fall. In an inversion, prior emissions are adjusted in proportion to the deviation

- 5 of the model's CH_4 DMFs from observed values. Attribution of these posterior emissions to different sectors depends on a priori information and assumptions about how they vary in time and location relative to one another. Thus, an increase in posterior emissions relative to the prior in the northern mid and high latitudes during winter will not change emissions from wetlands. For example, Fig. 10b illustrates the sensitivity of posterior wetland emissions to a three-month lag in the Northern Hemisphere. The change in posterior emissions is derived by calculating the total emissions required to produce an increase
- 10 of 1 ppb of CH_4 in each tropospheric column and scaling those emissions according to the a priori contribution of wetlands, estimated as the fractional contribution of wetlands to the total monthly mean emissions. The difference between this change in wetland emissions and the value in the same location three months prior produces the sensitivity of wetland emissions to the



Figure 10. (a) GEOS-Chem monthly zonal mean wetland emissions, in Gg. (b) The Northern Hemisphere sensitivity of GEOS-Chem wetland emission attribution caused by a 3-month lag for each 1 ppb increase of CH_4 in the tropospheric column, in Gg.

tropospheric phase lag. This approach provides an alternative to the computationally expensive calculation of the gain matrix over the entire time series but does not include information about model transport.

The tropics and subtropics are less sensitive to a phase shift, but polewards of 40° N, both the magnitude and seasonality of the difference is significant. Large differences between measured and modeled $X_{CH_4}^t$ are concurrent with low emissions from

5

5 seasonal sources. The adjustments to prior emissions produced by larger measurement-model disagreement that occur when seasonal sources are a small fraction of total emissions will overestimate posterior emissions from aseasonal sources. Thus these seasonal errors will bias source apportionment toward emissions that do not vary on timescales shorter than annually.

5 Conclusions

Assimilation of total column measurements into CTMs can improve constraints on the global CH_4 budget; however, the model's treatment of stratospheric chemistry and dynamics must be carefully considered. This work has compared TCCON and GEOS-Chem pressure-weighted total and tropospheric column-averaged CH_4 DMFs, X_{CH_4} and $X_{CH_4}^t$ respectively, parsing

5 out the seasonality of the troposphere and stratosphere and the resulting impacts on X_{CH_4} (Fig. 9a). The Southern Hemisphere measurement-model agreement is robust to changes in emissions or tropospheric OH. In the Northern Hemisphere the model's stratospheric contribution is larger than that of the measurements, and the mismatch increases as the tropopause altitude decreases. The result is greater model error at high-latitude sites, with the magnitude of this error varying seasonally. Moreover, in the Northern Hemisphere the GEOS-Chem $X_{CH_4}^t$ exhibits a 2-3 month phase lag. The combined tropospheric and stratospheric

10 errors smooth the model X_{CH_4} such that they may agree with total column measurements despite having an incorrect vertical distribution.

Model transport errors coupled with spatial and seasonal measurement sparsity can limit the accuracy of the location and timing of emissions scaling. The differences in the seasonality mismatch across vertical levels amplifies the error uncertainty because the timing of optimized fluxes will be especially susceptible to limitations in model transport. The stronger influence

15 of the stratosphere at higher latitudes due to lower tropopause heights, together with the higher temporal variability of the stratospheric fraction of the total column due to the stronger seasonal cycle of the tropopause, also impacts the seasonality of the meridional gradient of X_{CH_4} .

The influence of stratospheric variability on emissions is not unique to the model chosen for this analysis. Bergamaschi et al. (2013) ran TM5-4DVAR inversions using SCIAMACHY column and NOAA surface measurements and found that the mean

- ²⁰ biases between the optimized CH_4 profiles and aircraft measurements differ between the PBL, free troposphere, and UTLS. Seasonal emissions from wetlands and biomass burning vary by ± 10 and ± 7 TgCH₄, respectively, from year to year, and the zonal partitioning of posterior emissions is sensitive to the wetland priors chosen. Moreover, the larger changes to emissions and sensitivity to assumptions in the Northern Hemisphere indicate that TM5 is also subject to the strong hemispheric differences found in GEOS-Chem. The TransCom-CH₄ model comparison found that the interhemispheric exchange time in GEOS-
- 25 Chem was near the model median over the 1996-2007 time series (Patra et al., 2011), which suggests that GEOS-Chem's interhemispheric transport, and thus associated errors, is not particularly distinct. Ostler et al. (2016) found that ACTM and other CTMs used in TransCom-CH₄ are subject to transport errors that impact emissions optimization. Furthermore, ACTM profiles show a similar over-estimation of stratospheric CH₄, zonally-varying measurement-model mismatch dependent on tropopause height, and a smaller seasonal cycle for Northern Hemisphere X_{CH_4} compared to TCCON.
- 30 In this analysis we have used TCCON $X_{CH_4}^t$ derived with the HF-proxy method; however, $X_{CH_4}^t$ calculated using other stratospheric tracers such as nitrous oxide (N₂O) (Wang et al., 2014) would provide an additional constraint on models' representations of the stratosphere, as N₂O is not subject to the spectral interference with water vapor that impacts HF. Information about the vertical tropospheric CH₄ profile directly retrieved from NDACC spectra (Sepúlveda et al., 2014) can also be used to assess whether transport errors differ at different levels of the free troposphere. Ideally, information from these tropospheric

products could be integrated to overcome the limitations of each: the sensitivity of $X_{CH_4}^t$ to prior assumptions of stratospherictropospheric exchange and the sensitivity of profile retrievals to UTLS variability (Ostler et al., 2014).

A limitation of the aseasonal simulation was that the distribution of emissions was not identical to that of the base simulation due to the scaling approach we employed. Ideally, the aseasonal emissions for each sector would have been fluxes calculated

- 5 for each grid box from the base simulation annual emissions. The robustness of the model's tropospheric phase shift that was apparent regardless of the emissions used demonstrates that this feature is not a product of the chosen emissions fields. However, more nuanced analysis on smaller spatial scales would benefit from simulations that prescribe the annual mean for each of the seasonal sources. The most recent version of GEOS-Chem has a much more flexible emissions scheme (Keller et al., 2014) that allows these more nuanced experiments to be performed and analyzed.
- 10 The insensitivity of model stratospheres to tropospheric change allows for a straightforward solution: prescribed stratospheric CH_4 fields based on satellite observations from ACE-FTS, MIPAS (von Clarmann et al., 2009), or a compilation of remote sensing instruments (Buchwitz et al., 2015). As the representation of tropical convection and exchange across the UTLS advances in models and reduce stratospheric isolation, chemical loss, and transport mechanisms would need to be improved. The output from more accurate stratospheric models over the time period of interest could be used to set the stratospheric
- 15 component in the offline CH_4 simulation. For instance, the Universal tropospheric-stratospheric Chemistry eXtension (UCX) mechanism, which has been added to more recent versions of GEOS-Chem, updates the stratospheric component of the standard full chemistry simulation such that CH_4 has more sophisticated upwelling, advection and chemical reaction schemes (Eastham et al., 2014). Models that account for interannual variability in both stratospheric and tropospheric dynamics can then assimilate total column measurements to develop more accurate global CH_4 budgets.

20 Appendix A: Updates to Tropospheric Methane Data

The TCCON $X_{CH_4}^t$ data used in this analysis were developed as in Saad et al. (2014) with several adjustments to both the parameters used and methodology.

The HF-proxy method for determining X^t_{CH4} incorporates the relationship between CH4 and HF in the stratosphere, which is calculated using ACE-FTS data. These CH4-HF slopes now use updated ACE-FTS version 3.5 measurements with v.1.1 flags
(Boone et al., 2013; Sheese et al., 2015). The data quality flags are provided for profile data on a 1 km vertical grid, which uses a piecewise quadratic method to interpolate from the retrievals (Boone et al., 2013). Additionally, the CH4 and HF measurement errors are now considered in the pressure-weighted linear regression that determines the slopes. All other data processing to produce the CH4-HF slopes followed methods described in Saad et al. (2014). Figure 11 shows the updated annual zonal values used to calculate X^t_{CH4} with Washenfelder et al. (2003) and MkIV (retrieved from http://mark4sun.jpl.nasa.gov/m4data.html)

The derivation of the tropospheric column in Washenfelder et al. (2003), Saad et al. (2014), and Wang et al. (2014) implicitly assumed that the CH_4 profile is continuous across the tropopause; however, the boundary condition for stratospheric CH_4 is

³⁰

values included for reference (c.f. Saad et al., 2014, Fig. 2). These updates altered $\hat{X}_{CH_4}^t$ for the sites and time period covered in this paper by less than 2 ppb.



Figure 11. Long-term CH₄-HF slopes from Washenfelder et al. (2003), MkIV, and updated ACE-FTS measurements. Inset: Time series of zonal pressure-weighted ACE-FTS slopes (β) used to calculate $\hat{X}_{CH_4}^t$, with error bars denoting the 2σ standard error. Zonal slopes are offset each year for visual clarity.

rather set by tropospheric air transported through the tropical tropopause (Brewer, 1949; Dobson, 1956). Boering et al. (1996) showed that the concentration of CO₂ directly above the tropopause can be approximated by introducing a two-month phase lag to the average concentration at northern and southern tropical surface sites: Mauna Loa, Hawaii (MLO) and Tutuila, American Samoa (SMO), respectively. As the CH_4 entering the stratosphere originates in both hemispheres (Boering et al.,

- 1995), stratospheric CH_4 exhibits a smaller interhemispheric gradient than in the troposphere: about 20ppb, as calculated 5 from ACE-FTS measurements, versus about 50 ppb, taken as the difference at MLO and SMO. To calculate the stratospheric boundary condition for CH₄ we remove the seasonal component of the mean of CH₄ DMFs at MLO and SMO, which are made available through 2014 by the NOAA Earth System Research Laboratory (ESRL) Global Monitoring Division (Dlugokencky et al., 2016). To capture the interhemispheric gradient observed in ACE stratospheric CH_4 measurements, we add and subtract
- 10 ppb, in the northern and southern extratropics respectively, the limits of which we choose as the Tropic of Cancer $(23^{\circ}N)$ 10 and the Tropic of Capricorn (23°S). A constant value is chosen in each hemisphere to reflect the rapid mixing time of air from the extra-tropics in the region directly above the tropopause, which Boering et al. (1996) found to be less than one month. Within the tropics, we interpolate the boundary condition as a linear function of altitude such that $x_{CH_4}(P^t) = \bar{x}_{CH_4}^s + \frac{10}{23}\lambda$, where $x_{CH_4}(P^t)$ is the boundary condition at the tropopause, $\bar{x}_{CH_4}^s$ is the mean DMF of CH₄ at the surface, and λ is the
- 15 latitude of the site.

Assuming hydrostatic equilibrium, the tropospheric column of CH_4 , $c_{CH_4}^t$, can be calculated as the integral of the vertical profile, $x_{CH_4} \equiv x_{CH_4}(P)$, from the surface, P^s , to the tropopause, P^t :

$$c_{\rm CH_4}^t = \int_{P^t}^{P^s} x_{\rm CH_4} \frac{dP}{gm} = X_{\rm CH_4}^t \frac{P^s - P^t}{g_*^t m}$$
(A1)

where P is the pressure height, g is the gravitational acceleration, g_*^t is the pressure-weighted tropospheric value of g, and m 20

is the mean molecular mass of CH₄ (Washenfelder et al., 2006). The profile of CH₄ in the stratosphere can be expressed as a linear function of pressure altitude, $x_{CH_4}(P) = x_{CH_4}(P^t) + \delta \cdot P$, where $\delta = \frac{dx_{CH_4}}{dP}$ is the stratospheric loss of CH₄. This stratospheric loss term is estimated by the HF-proxy method to produce the retrieved tropospheric column-averaged DMF, $\hat{X}_{CH_4}^t$, such that

5
$$\hat{X}_{CH_4}^t \frac{P^s}{g_*m} = \hat{c}_{CH_4}^t = \int_0^{P^s} x_{CH_4} \frac{dP}{gm} - \int_0^{P^t} \delta \cdot P \frac{dP}{gm}$$
 (A2)

where g_* is the pressure-weighted column average of g. The stratospheric boundary condition can thus be related to the retrieved tropospheric column as

$$\int_{0}^{P^{t}} x_{\mathrm{CH}_{4}} \frac{dP}{gm} = \int_{0}^{P^{t}} x_{\mathrm{CH}_{4}}(P^{t}) \frac{dP}{gm} - \hat{c}_{\mathrm{CH}_{4}}^{t} + \int_{0}^{P^{s}} x_{\mathrm{CH}_{4}} \frac{dP}{gm}.$$
(A3)

10 Given the total column integration is the sum of the tropospheric and stratospheric partial columns, and substituting Equation A3:

$$\int_{P^t}^{P^s} x_{\mathrm{CH}_4} \frac{dP}{gm} = \int_{0}^{P^s} x_{\mathrm{CH}_4} \frac{dP}{gm} - \int_{0}^{P^t} x_{\mathrm{CH}_4} \frac{dP}{gm}$$
(A4)

$$= \int_{0}^{P^{s}} x_{\mathrm{CH}_{4}} \frac{dP}{gm} - \int_{0}^{P^{t}} x_{\mathrm{CH}_{4}}(P^{t}) \frac{dP}{gm} + \hat{c}_{\mathrm{CH}_{4}}^{t} - \int_{0}^{P^{s}} x_{\mathrm{CH}_{4}} \frac{dP}{gm}$$
(A5)

$$= \hat{c}_{\mathrm{CH}_4}^t - \int\limits_0^F x_{\mathrm{CH}_4}(P^t) \frac{dP}{gm}$$
(A6)

15
$$X_{CH_4}^t \frac{P^s - P^t}{g_*^t m} = \hat{X}_{CH_4}^t \frac{P^s}{g_* m} - x_{CH_4} (P^t) \frac{P^t}{g_*^0 m}$$
 (A7)

where g_*^0 is the pressure-weighted average of g from the tropopause to the top of the atmosphere. While the molecular mass of air changes as a function of water vapor and thus altitude and gravity changes as a function of both altitude and latitude, assuming constant values of g and m changes $X_{CH_4}^t$ by less than 2 ppb. Thus, to good approximation these variables can be canceled out:

$$X_{CH_4}^t[P^s - P^t] = \hat{X}_{CH_4}^t \cdot P^s - x_{CH_4}(P^t) \cdot P^t$$
(A8)

20

$$X_{\rm CH_4}^t = \frac{\ddot{X}_{\rm CH_4}^t \cdot P^s - x_{\rm CH_4}(P^t) \cdot P^t}{P^s - P^t}.$$
(A9)

The surface pressure is measured at each site, and the tropopause pressure is calculated from the TCCON prior temperature

profiles. The uncertainties associated with the interpolated value of the tropopause height are determined by calculating $X_{CH_4}^t$ for $\pm 30\%$ of P^t and adding these confidence intervals in quadrature to the precision error of $\hat{X}_{CH_4}^t$. The aforementioned deseasonalization of $x_{CH_4}(P^t)$ is an approximation that adds another uncertainty. The signal of the tropospheric seasonal cycle of a trace gas entering the stratosphere is apparent directly above the tropopause and both dampens in amplitude and shifts in

5 time with increasing altitude (Mote et al., 1996). Thus, the stratospheric boundary condition is not truly constant throughout the column, but rather the pressure-weighted sum of these attenuated signals. Calculating $x_{CH_4}(P^t)$ without removing the seasonality, which provides the maximum impact of this uncertainty, decreases $X_{CH_4}^t$ by an average of 1 ppb and 4 ppb in the Northern and Southern Hemispheres, respectively, and does not alter the seasonal cycle of $X_{CH_4}^t$. Moreover, as described below, the mismatch between the calibrated TCCON $X_{CH_4}^t$ and the in situ aircraft $X_{CH_4}^t$ does not correlate with season ($R^2 = 0.017$).

10 Thus, we retain the simpler computation of deseasonalized $x_{CH_4}(P^t)$ in Equation A9.

Airmass-dependent artifacts were derived for updated values consistently with the total column CH₄ (Wunch et al., 2015). Removing these artifacts, the $X_{CH_4}^t$ was then calibrated with in situ aircraft profiles using the same methodology described in Wunch et al. (2010) and including the updates delineated in Wunch et al. (2015) to produce a calibration correction factor of 0.9700 (Fig. 12). The covariance between the difference between the calibrated TCCON and aircraft $X_{CH_4}^t$ and several

15 parameters were assessed to ensure biases were not introduced into the measurements. These differences had an uncertaintyweighted correlation coefficient of 0.1 for solar zenith angle and uncertainty-weighted correlation coefficients of less that 0.02 for tropopause and surface pressures, year, and season. Measurement precisions and errors were determined as in Saad et al. (2014), with the additional uncertainties mentioned in this section included. Individual TCCON sites have median $X_{CH_4}^t$



Figure 12. Calibration curve of TCCON $X_{CH_4}^t$ (c.f. Wunch et al., 2015, Fig. 8). Site colors are as in Fig. 1. Aircraft campaigns are described in Table 6 of Wunch et al. (2015).

precisions in the range of 0.1-0.8%, and mean and median precisions are 0.3 and 0.2%, respectively, for all sites through May 2016.

Appendix B: GEOS-Chem Simulations

B1 Equilibrium Sensitivity Experiments

- 5 All equilibrium runs for a given simulation have identical meteorology, emissions, and OH fields over June 2004-May 2005. Initial conditions for each year are set by the restart files of the previous run. To calculate columns at each site, GEOS-Chem monthly mean mole fractions are adjusted for the monthly medians of the site's daily mean surface pressures and smoothed with the monthly median scaled prior profiles and averaging kernels, interpolated using the monthly median solar zenith angle daily means. Because Park Falls and Lauder are the only TCCON sites that had started taking measurements over this time
 10 period, they are the only sites used to generate smoothed columns for the comparisons to the experimental simulations.
- Emissions in the aseasonal simulation were derived by running a two-dimensional regression on the annual emissions to determine the scale factors that would produce the smallest residual of total emissions and the interhemispheric gradient. Figure 13 illustrates the difference in total emissions between the base and aseasonal simulations for each zonal band.

The updated OH simulation used OH output from a 2012 GEOS-Chem standard chemistry simulation with extensive updates to the photochemical oxidation mechanisms of biogenic volatile organic compounds (VOCs), described in Bates et al. (2016) and references therein. These were converted to 3D monthly mean OH concentrations to conform to the infrastructure of the GEOS-Chem offline CH₄ tropospheric loss mechanism. The OH was then scaled by 90% to keep the lifetime above 8 years,



Figure 13. Monthly averages of the difference in total CH_4 emissions between the base and aseasonal GEOS-Chem simulations, summed over each zonal band, in Tg·mo⁻¹.



Figure 14. Zonal averages of the difference in total column OH (molec \cdot cm⁻²) between the base and updated monthly OH fields.

and emissions were scaled by 112% to maintain the same balance between sources and sinks in the base simulation. Figure 14 provides zonal averages of the difference between the base and updated OH columns.

The full list of simulations run is provided in Table 3, with descriptions and the CH_4 emissions, tropospheric OH, and total chemical loss lifetimes. Figure 15 shows each simulation's seasonality of $X_{CH_4}^t$ at Park Falls, with TCCON seasonality plotted for reference, as well as the seasonality of the difference between the base and each simulation.

B2 Derivation of Dry Gas Values

5

15

Versions of GEOS-Chem prior to v.10 have inconsistencies in wet versus dry definitions of pressure, temperature, and air mass, which propagate into model diagnostics and conversions calculated using these terms. As a consequence, CH_4 concentrations are output assuming air masses that include water vapor but calculated with the molar mass of dry air. For all comparisons in

10 this analysis CH_4 DMFs are calculated taking into account the GEOS-5 specific humidity, q_s (in units of $g_{H2O} \cdot kg_{air}^{-1}$), such that

$$x_{\rm CH_4, dry} = \frac{x_{\rm CH_4}}{1 - q_s \times 10^{-3}} \tag{B1}$$

where x_{CH_4} is the model profile in mole fractions. Dry air profiles were derived by subtracting the water vapor mole fraction, also calculated from the GEOS-5 specific humidity, from the total air mass at each pressure level, as in Wunch et al. (2010); Geibel et al. (2012).

B3 Model Smoothing for Measurement Comparisons

Base and aseasonal daily runs were initialized using CH_4 fields from their respective 34th equilibrium cycles. Daily CH_4 mole fractions averaged over both 24-hour and 10-14 local time were output to test whether TCCON's daytime-only observations



Figure 15. Seasonality of tropospheric methane $(X_{CH_4}^t)$ at Park Falls for TCCON (black solid line), GEOS-Chem (red solid line), and the difference from the base simulation (dotted red line) for each of the sensitivity experiments, in ppb.

would introduce a bias in the comparisons. Measurement-model differences were not sensitive to averaging times. Comparison of measurements to model columns produced using the 24-hour and 10-14 local time averages produce equivalent slopes and only slightly different intercepts and correlation coefficients. The seasonality of 10-14 local time column-averaged DMFs does not differ, except that the fall seasonal maximum of the adjusted troposphere and stratospheric contribution at Park Falls in October, one month later than the 24-hour column-averaged DMF seasonality.

 CH_4 dry vertical profiles for each grid box associated with a TCCON site, $x_{CH_4}^m$, were smoothed with corresponding FTS column averaging kernels, a_{CH_4} , and scaled priors for each day and vertically integrated using pressure-weighted levels:

5

$$X_{\rm CH_4}^s = \gamma_{\rm CH_4} \cdot X_{\rm CH_4}^a + \boldsymbol{a}_{\rm CH_4}^{\$} (\boldsymbol{x}_{\rm CH_4}^m - \gamma_{\rm CH_4} \boldsymbol{x}_{\rm CH_4}^a)$$
(B2)

where $X_{CH_4}^s$ is the smoothed GEOS-Chem column-averaged DMF, γ_{CH_4} is the TCCON daily median retrieved profile scal-10 ing factor, and $x_{CH_4}^a$ and $X_{CH_4}^a$ are respectively the a priori profile and column-integrated CH₄ DMFs (Rodgers and Connor, 2003). The pressure weighting function, h, was applied such that $X = h^T x$. TCCON priors were interpolated to the GEOS-Chem pressure grid, and GEOS-Chem pressure and corresponding gas profiles were adjusted using daily mean surface pressures local to each site (Wunch et al., 2010; Messerschmidt et al., 2011). The averaging kernels were interpolated for the local daily mean solar zenith angle and the GEOS-Chem pressure grid so that it could be applied to the difference between the

15 GEOS-Chem and TCCON profiles as $a^{\$}x = \sum_{i=1}^{N} a_i h_i x_i$ from the surface to the highest level, *N*, at *i* pressure levels (Connor et al., 2008; Wunch et al., 2011b). Figure 16 shows how the smoothed column compares to the column that only uses the dry gas correction.



Figure 16. GEOS-Chem smoothed versus dry integrated CH_4 DMFs for base simulation tropospheric columns, total columns, and stratospheric contribution. Site colors are as in Fig. 1. Dashed lines mark the one-to-one lines.

Appendix C: Derivation of Stratospheric Contribution

Considering the CH₄ profile integration as in Equation A4, and substituting the profile of CH₄ in the stratosphere, $x_{CH_4}(P) = x_{CH_4}(P^t) + \delta \cdot P$, described in Appendix A, the total column is calculated as:

$$\int_{0}^{P^s} x_{\mathrm{CH}_4} \frac{dP}{gm} = \int_{P^t}^{P^s} x_{\mathrm{CH}_4} \frac{dP}{gm} + \int_{0}^{P^t} [x_{\mathrm{CH}_4}(P^t) + \delta \cdot P] \frac{dP}{gm}$$
(C1)

5
$$X_{CH_4} \cdot P^s = X_{CH_4}^t [P^s - P^t] + x_{CH_4} (P^t) \cdot P^t + c_{CH_4}^{\delta}$$
 (C2)

where $c_{CH_4}^{\delta}$, is the pressure-weighted column average of CH_4 loss in the stratosphere. Rearranging terms, Equation C2 becomes:

$$[X_{\rm CH_4} - X_{\rm CH_4}^t]P^s = [x_{\rm CH_4}(P^t) - X_{\rm CH_4}^t]P^t + c_{\rm CH_4}^\delta$$
(C3)

$$X_{CH_4}^t - X_{CH_4} = [X_{CH_4}^t - x_{CH_4}(P^t)] \frac{P^t}{P^s} - \frac{c_{CH_4}^o}{P^s}$$
(C4)

- 10 such that the difference between the tropospheric and total column-averaged DMFs is a function of the two terms governing the stratospheric contribution to the total column: the gradient across the tropopause, $x_{CH_4}(P^t) - X_{CH_4}^t$, and stratospheric CH_4 loss, $c_{CH_4}^{\delta}$. The stratospheric contribution is thus a proxy for the impact of stratospheric variability on the total column of CH_4 : given a constant tropospheric column, as the stratospheric contribution becomes larger the total column-averaged DMF becomes smaller.
- 15 Acknowledgements. This work was supported by NASA Headquarters under the NASA Earth and Space Science Fellowship Program grant NNX14AL30H and NASA's Carbon Cycle Science program. Park Falls, Lamont, and JPL are funded by NASA grants NNX14AI60G, NNX11AG01G, NAG5-12247, NNG05-GD07G, and NASA Orbiting Carbon Observatory Program; we are grateful to the DOE ARM program and Jeff Ayers for their technical support in Lamont and Park Falls, respectively. Darwin and Wollongong are funded by NASA grants NAG5-12247 and NNG05-GD07G and the Australian Research Council grants DP140101552, DP110103118, DP0879468 and LP0562346,
- 20 and Nicholas Deutscher is supported by an Australian Research Council Fellowship, DE140100178; we are grateful to the DOE ARM program for technical support in Darwin. Bremen, Bialystok, and Orleans are funded by the EU projects InGOS and ICOS-INWIRE and by the Senate of Bremen. Réunion Island is funded by the EU FP7 project ICOS-INWIRE, the national Belgian support to ICOS and the AGACC-II project (Science for Sustainable Development Program), the Université de la Réunion, and the French regional and national organizations (INSU, CNRS). From 2004 to 2011 the Lauder TCCON program was funded by the New Zealand Foundation of Research Science and
- 25 Technology contracts CO1X0204, CO1X0703 and CO1X0406. We thank Shuji Kawakami for his technical support in Saga. We thank Peter Bernath, Kaley Walker, and Chris Boone for their guidance using the ACE-FTS data, which were obtained through the Atmospheric Chemistry Experiment (ACE) mission, primarily funded by the Canadian Space Agency. We are grateful to Geoff Toon for his continuous efforts developing the GGG software, for providing the MkIV data, and his input on the manuscript. We thank Arlyn Andrews for providing the LEF surface flask data, which were generated by NOAA-ESRL, Carbon Cycle Greenhouse Gases Group. Baring Head NIWA surface data
- 30 were provided courtesy of Gordon Brailsford, Dave Lowe and Ross Martin. We also acknowledge the contributions of in situ vertical profiles

from the AirCore, HIPPO, IMECC, INTEX, Learjet, and START08 campaigns. We are grateful to Kelvin Bates for providing monthly OH fields for the GEOS-Chem Updated OH sensitivity experiments. Lastly, we thank the two anonymous reviewers who provided comments and suggestions.

References

- Allen, D., Pickering, K., Duncan, B., and Damon, M.: Impact of lightning NO emissions on North American photochemistry as determined using the Global Modeling Initiative (GMI) model, Journal of Geophysical Research: Atmospheres, 115, 2010.
- Baker, D., Law, R., Gurney, K., Rayner, P., Peylin, P., Denning, A., Bousquet, P., Bruhwiler, L., Chen, Y., Ciais, P., Fung, I., Heimann, M.,
- 5 John, J., Maki, T., Maksyutov, S., Masarie, K., Prather, M., Pak, B., Taguchi, S., and Zhu, Z.: TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO2 fluxes, 1988-2003, Global Biogeochemical Cycles, 20, doi:10.1029/2004GB002439, 2006.
 - Bates, K. H., Nguyen, T. B., Teng, A. P., Crounse, J. D., Kjaergaard, H. G., Stoltz, B. M., Seinfeld, J. H., and Wennberg, P. O.: Production and Fate of C4 Dihydroxycarbonyl Compounds from Isoprene Oxidation, The Journal of Physical Chemistry A, 120, 106–117,
- 10 doi:10.1021/acs.jpca.5b10335, http://dx.doi.org/10.1021/acs.jpca.5b10335, 2016.
 - Bergamaschi, P., Frankenberg, C., Meirink, J. F., Krol, M., Dentener, F., Wagner, T., Platt, U., Kaplan, J. O., Körner, S., Heimann, M., Dlugokencky, E. J., and Goede, A.: Satellite chartography of atmospheric methane from SCIAMACHY on board ENVISAT: 2. Evaluation based on inverse model simulations, Journal of Geophysical Research: Atmospheres, 112, n/a–n/a, doi:10.1029/2006JD007268, http: //dx.doi.org/10.1029/2006JD007268, d02304, 2007.
- 15 Bergamaschi, P., Houweling, S., Segers, A., Krol, M., Frankenberg, C., Scheepmaker, R. A., Dlugokencky, E., Wofsy, S. C., Kort, E. A., Sweeney, C., Schuck, T., Brenninkmeijer, C., Chen, H., Beck, V., and Gerbig, C.: Atmospheric CH4 in the first decade of the 21st century: Inverse modeling analysis using SCIAMACHY satellite retrievals and NOAA surface measurements, Journal of Geophysical Research: Atmospheres, 118, 7350–7369, doi:10.1002/jgrd.50480, http://dx.doi.org/10.1002/jgrd.50480, 2013.
- Bernath, P. F.: Atmospheric Chemistry Experiment (ACE): Mission overview, Geophysical Research Letters, 32, L15S01,
 doi:10.1029/2005GL022386, http://doi.wiley.com/10.1029/2005GL022386, 2005.
- Bey, I., Jacob, D., Yantosca, R., Logan, J., Field, B., Fiore, A., Li, Q., Liu, H., Mickley, L., and Schultz, M.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, Journal of Geophysical Research-Atmospheres, 106, 23 073– 23 095, doi:10.1029/2001JD000807, 2001.
- Blumenstock, T., Hase, F., Schneider, M., Garcia, O., and Sepulveda, E.: TCCON data from Izana, Tenerife, Spain, Release GGG2014R0,
 http://dx.doi.org/10.14291/tccon.ggg2014.izana01.R0/1149295, doi:10.14291/tccon.ggg2014.izana01.R0/1149295, TCCON data archive,
 - hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
 - Boering, K., Wofsy, S., Daube, B., Schneider, H., Loewenstein, M., Podolske, J., and Conway, T.: Stratospheric Mean Ages and Transport Rates from Observations of Carbon Dioxide and Nitrous Oxide, Science, 274, 1340–1343, doi:10.1126/science.274.5291.1340, 1996.
 - Boering, K. A., Hintsa, E. J., Wofsy, S. C., Anderson, J. G., Daube, B. C., Dessler, A. E., Loewenstein, M., McCormick, M. P., Podolske, J. R., Weinstock, E. M., and Yue, G. K.: Measurements of stratospheric carbon dioxide and water vapor at northern midlatitudes: Implications
- 30 Weinstock, E. M., and Yue, G. K.: Measurements of stratospheric carbon dioxide and water vapor at northern midlatitudes: Implications for troposphere-to-stratosphere transport, Geophysical Research Letters, 22, 2737–2740, doi:10.1029/95GL02337, http://dx.doi.org/10. 1029/95GL02337, 1995.
- Boone, C. D., Walker, K. A., and Bernath, P. F.: Version 3 Retrievals for the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS), in: The Atmospheric Chemistry Experiment ACE at 10: A Solar Occultation Anthology, edited by Bernath, P. F.,
 pp. 103–127, A. Deepak Publishing, Hampton, Virginia, U.S.A., 2013.
 - Brewer, A.: Evidence for a world circulation provided by the measurements of helium and water vapour distribution in the stratosphere, Quarterly Journal of the Royal Meteorological Society, 75, 351–363, 1949.

- Buchwitz, M., Reuter, M., Schneising, O., Boesch, H., Guerlet, S., Dils, B., Aben, I., Armante, R., Bergamaschi, P., Blumenstock, T., Bovensmann, H., Brunner, D., Buchmann, B., Burrows, J. P., Butz, A., Chédin, A., Chevallier, F., Crevoisier, C. D., Deutscher, N. M., Frankenberg, C., Hase, F., Hasekamp, O. P., Heymann, J., Kaminski, T., Laeng, A., Lichtenberg, G., De Mazière, M., Noël, S., Notholt, J., Orphal, J., Popp, C., Parker, R., Scholze, M., Sussmann, R., Stiller, G. P., Warneke, T., Zehner, C., Bril, A., Crisp, D., Griffith, D. W. T.,
- 5 Kuze, A., O'Dell, C., Oshchepkov, S., Sherlock, V., Suto, H., Wennberg, P., Wunch, D., Yokota, T., and Yoshida, Y.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): Comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, Remote Sensing of Environment, 162, 344–362, 2015.
 - Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., and Aben, I.: TROPOMI aboard Sentinel-5 Precursor: Prospective performance of {CH4} retrievals for aerosol and cirrus loaded atmospheres, Remote Sensing of Environment, 120, 267–276,
- 10 doi:http://dx.doi.org/10.1016/j.rse.2011.05.030, http://www.sciencedirect.com/science/article/pii/S003442571200082X, the Sentinel Missions New Opportunities for Science, 2012.
 - Connor, B. J., Boesch, H., Toon, G., Sen, B., Miller, C., and Crisp, D.: Orbiting Carbon Observatory: Inverse method and prospective error analysis, Journal of Geophysical Research, 113, 1–14, doi:10.1029/2006JD008336, 2008.

Considine, D. B., Logan, J. A., and Olsen, M. A.: Evaluation of near-tropopause ozone distributions in the Global Modeling Initiative
 combined stratosphere/troposphere model with ozonesonde data, Atmospheric Chemistry and Physics, 8, 2365–2385, 2008.

De Mazière, M., Vigouroux, C., Bernath, P. F., Baron, P., Blumenstock, T., Boone, C., Brogniez, C., Catoire, V., Coffey, M., Duchatelet, P., Griffith, D., Hannigan, J., Kasai, Y., Kramer, I., Jones, N., Mahieu, E., Manney, G. L., Piccolo, C., Randall, C., Robert, C., Senten, C., Strong, K., Taylor, J., Tétard, C., Walker, K. A., and Wood, S.: Validation of ACE-FTS v2.2 methane profiles from the upper troposphere to the lower mesosphere, Atmospheric Chemistry and Physics, 8, 2421–2435, doi:10.5194/acp-8-2421-2008, http://www.atmos-chem-phys.

20 net/8/2421/2008/, 2008.

- De Maziere, M., Sha, M. K., Desmet, F., Hermans, C., Scolas, F., Kumps, N., Metzger, J.-M., Duflot, V., and Cammas, J.-P.: TCCON data from Reunion Island (La Reunion), France, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.reunion01.R0/1149288, doi:10.14291/tccon.ggg2014.reunion01.R0/1149288, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- 25 Deutscher, N., Notholt, J., Messerschmidt, J., Weinzierl, C., Warneke, T., Petri, C., Grupe, P., and Katrynski, K.: TC-CON data from Bialystok, Poland, Release GGG2014R1, http://dx.doi.org/10.14291/tccon.ggg2014.bialystok01.R1/1183984, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.

 Dlugokencky, E., Lang, P., Crotwell, A., Mund, J., Crotwell, M., and Thoning, K.: Atmospheric Methane Dry Air
 30 Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1983-2015, ftp://aftp.cmdl.noaa.gov/data/trace_gases/ch4/flask/surface/, 2016.

- Dlugokencky, E. J., Nisbet, E. G., Fisher, R., and Lowry, D.: Global atmospheric methane: budget, changes and dangers, Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences, 369, 2058–2072, doi:10.1098/rsta.2010.0341, http://rsta.royalsocietypublishing.org/content/369/1943/2058, 2011.
- 35 Dobson, G.: Origin and distribution of the polyatomic molecules in the atmosphere, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, 236, 187–193, 1956.

- Eastham, S. D., Weisenstein, D. K., and Barrett, S. R.: Development and evaluation of the unified tropospheric–stratospheric chemistry extension (UCX) for the global chemistry-transport model GEOS-Chem, Atmospheric Environment, 89, 52–63, doi:http://dx.doi.org/10.1016/j.atmosenv.2014.02.001, http://www.sciencedirect.com/science/article/pii/S1352231014000971, 2014.
 European Commission Joint Research Centre, Netherlands Environmental Assessment Agency: Emission Database for Global Atmospheric
- 5 Research (EDGAR), http://edgar.jrc.ec.europa.eu, available at http://edgar.jrc.ec.europa.eu, 2011.
- Fraser, A., Chan Miller, C., Palmer, P. I., Deutscher, N. M., Jones, N. B., and Griffith, D. W. T.: The Australian methane budget: Interpreting surface and train-borne measurements using a chemistry transport model, Journal of Geophysical Research: Atmospheres, 116, n/a–n/a, doi:10.1029/2011JD015964, http://dx.doi.org/10.1029/2011JD015964, d20306, 2011.
- Fraser, A., Palmer, P. I., Feng, L., Boesch, H., Cogan, A., Parker, R., Dlugokencky, E. J., Fraser, P. J., Krummel, P. B., Langenfelds, R. L.,
- 10 O'Doherty, S., Prinn, R. G., Steele, L. P., van der Schoot, M., and Weiss, R. F.: Estimating regional methane surface fluxes: the relative importance of surface and GOSAT mole fraction measurements, Atmospheric Chemistry and Physics, 13, 5697–5713, doi:10.5194/acp-13-5697-2013, http://www.atmos-chem-phys.net/13/5697/2013/, 2013.
 - Fung, I., John, J., Lerner, J., Matthews, E., Prather, M., Steele, L., and Fraser, P.: Three-dimensional model synthesis of the global methane cycle, Journal of Geophysical Research: Atmospheres, 96, 13 033–13 065, 1991.
- 15 Geibel, M. C., Messerschmidt, J., Gerbig, C., Blumenstock, T., Chen, H., Hase, F., Kolle, O., Lavrič, J. V., Notholt, J., Palm, M., Rettinger, M., Schmidt, M., Sussmann, R., Warneke, T., and Feist, D. G.: Calibration of column-averaged CH₄ over European TCCON FTS sites with airborne in-situ measurements, Atmospheric Chemistry and Physics, 12, 8763–8775, doi:10.5194/acp-12-8763-2012, http://www.atmos-chem-phys.net/12/8763/2012/, 2012.
- Gentner, D. R., Ford, T. B., Guha, A., Boulanger, K., Brioude, J., Angevine, W. M., de Gouw, J. A., Warneke, C., Gilman, J. B., Ryerson,
- 20 T. B., Peischl, J., Meinardi, S., Blake, D. R., Atlas, E., Lonneman, W. A., Kleindienst, T. E., Beaver, M. R., St Clair, J. M., Wennberg, P. O., VandenBoer, T. C., Markovic, M. Z., Murphy, J. G., Harley, R. A., and Goldstein, A. H.: Emissions of organic carbon and methane from petroleum and dairy operations in California's San Joaquin Valley, Atmospheric Chemistry and Physics, 14, 4955–4978, doi:10.5194/acp-14-4955-2014, 2014.

Griffith, D. W. T., Deutscher, N., Velazco, V. A., Wennberg, P. O., Yavin, Y., Aleks, G. K., Washenfelder, R., Toon, G. C., Blavier, J.-

- F., Murphy, C., Jones, N., Kettlewell, G., Connor, B., Macatangay, R., Roehl, C., Ryczek, M., Glowacki, J., Culgan, T., and Bryant, G.: TCCON data from Darwin, Australia, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.darwin01.R0/1149290, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014a.
- Griffith, D. W. T., Velazco, V. A., Deutscher, N., Murphy, C., Jones, N., Wilson, S., Macatangay, R., Kettlewell,
 G., Buchholz, R. R., and Riggenbach, M.: TCCON data from Wollongong, Australia, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.wollongong01.R0/1149291, doi:10.14291/tccon.ggg2014.wollongong01.R0/1149291, TC-CON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014b.

Hase, F., Blumenstock, T., Dohe, S., Gross, J., and Kiel, M.: TCCON data from Karlsruhe, Germany, Release GGG2014R1,

- 35 http://dx.doi.org/10.14291/tccon.ggg2014.karlsruhe01.R1/1182416, doi:10.14291/tccon.ggg2014.karlsruhe01.R1/1182416, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
 - Jones, A., Walker, K. A., Jin, J. J., Taylor, J. R., Boone, C. D., Bernath, P. F., Brohede, S., Manney, G. L., McLeod, S., Hughes, R., and Daffer, W. H.: Technical Note: A trace gas climatology derived from the Atmospheric Chemistry Experiment Fourier Transform Spectrometer

(ACE-FTS) data set, Atmospheric Chemistry and Physics, 12, 5207–5220, doi:10.5194/acp-12-5207-2012, http://www.atmos-chem-phys. net/12/5207/2012/, 2012.

- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., and Zhu, Y.: The NCEP/NCAR 40-year reanalysis project, Bulletin of the American Meteorological Society, 77, 437–471, 1996.
- 5 Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Taura, C., Fukamachi, T., and Sakashita, M.: TCCON data from Saga, Japan, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.saga01.R0/1149283, doi:10.14291/tccon.ggg2014.saga01.R0/1149283, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
 - Keller, C. A., Long, M. S., Yantosca, R. M., Da Silva, A. M., Pawson, S., and Jacob, D. J.: HEMCO v1.0: a versatile, ESMF-compliant
- 10 component for calculating emissions in atmospheric models, Geoscientific Model Development, 7, 1409–1417, doi:10.5194/gmd-7-1409-2014, http://www.geosci-model-dev.net/7/1409/2014/, 2014.
 - Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., et al.: Three decades of global methane sources and sinks, Nature Geoscience, 6, 813–823, 2013.

Mahieu, E., Duchatelet, P., Demoulin, P., Walker, K. A., Dupuy, E., Froidevaux, L., Randall, C., Catoire, V., Strong, K., Boone, C. D.,

- 15 Bernath, P. F., Blavier, J.-F., Blumenstock, T., Coffey, M., De Mazière, M., Griffith, D., Hannigan, J., Hase, F., Jones, N., Jucks, K. W., Kagawa, A., Kasai, Y., Mebarki, Y., Mikuteit, S., Nassar, R., Notholt, J., Rinsland, C. P., Robert, C., Schrems, O., Senten, C., Smale, D., Taylor, J., Tétard, C., Toon, G. C., Warneke, T., Wood, S. W., Zander, R., and Servais, C.: Validation of ACE-FTS v2.2 measurements of HCl, HF, CCl₃F and CCl₂F₂ using space-, balloon- and ground-based instrument observations, Atmospheric Chemistry and Physics, 8, 6199–6221, doi:10.5194/acp-8-6199-2008, http://www.atmos-chem-phys.net/8/6199/2008/, 2008.
- 20 Messerschmidt, J., Geibel, M. C., Blumenstock, T., Chen, H., Deutscher, N. M., Engel, A., Feist, D. G., Gerbig, C., Gisi, M., Hase, F., Katrynski, K., Kolle, O., Lavrič, J. V., Notholt, J., Palm, M., Ramonet, M., Rettinger, M., Schmidt, M., Sussmann, R., Toon, G. C., Truong, F., Warneke, T., Wennberg, P. O., Wunch, D., and Xueref-Remy, I.: Calibration of TCCON column-averaged CO₂: the first aircraft campaign over European TCCON sites, Atmospheric Chemistry and Physics, 11, 10765–10777, doi:10.5194/acp-11-10765-2011, http://www.atmos-chem-phys.net/11/10765/2011/, 2011.
- 25 Mote, P. W., Rosenlof, K. H., McIntyre, M. E., Carr, E. S., Gille, J. C., Holton, J. R., Kinnersley, J. S., Pumphrey, H. C., Russell, J. M., and Waters, J. W.: An atmospheric tape recorder: The imprint of tropical tropopause temperatures on stratospheric water vapor, Journal of Geophysical Research: Atmospheres, 101, 3989–4006, doi:10.1029/95JD03422, http://dx.doi.org/10.1029/95JD03422, 1996.

Mu, M., Randerson, J. T., van der Werf, G. R., Giglio, L., Kasibhatla, P., Morton, D., Collatz, G. J., DeFries, R. S., Hyer, E. J., Prins, E. M., Griffith, D. W. T., Wunch, D., Toon, G. C., Sherlock, V., and Wennberg, P. O.: Daily and 3-hourly variability in global fire

- 30 emissions and consequences for atmospheric model predictions of carbon monoxide, Journal of Geophysical Research: Atmospheres, 116, doi:10.1029/2011JD016245, http://dx.doi.org/10.1029/2011JD016245, d24303, 2011.
 - Murray, L. T., Jacob, D. J., Logan, J. A., Hudman, R. C., and Koshak, W. J.: Optimized regional and interannual variability of lightning in a global chemical transport model constrained by LIS/OTD satellite data, Journal of Geophysical Research: Atmospheres, 117, 2012.
- Notholt, J., Petri, C., Warneke, T., Deutscher, N., Buschmann, M., Weinzierl, C., Macatangay, R., and Grupe, P.: TC CON data from Bremen, Germany, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.bremen01.R0/1149275, doi:10.14291/tccon.ggg2014.bremen01.R0/1149275, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.

Ostler, A., Sussmann, R., Rettinger, M., Deutscher, N. M., Dohe, S., Hase, F., Jones, N., Palm, M., and Sinnhuber, B.-M.: Multistation intercomparison of column-averaged methane from NDACC and TCCON: impact of dynamical variability, Atmospheric Measurement Techniques, 7, 4081–4101, doi:10.5194/amt-7-4081-2014, http://www.atmos-meas-tech.net/7/4081/2014/, 2014.

Ostler, A., Sussmann, R., Patra, P. K., Houweling, S., De Bruine, M., Stiller, G. P., Haenel, F. J., Plieninger, J., Bousquet, P., Yin, Y., Saunois,

M., Walker, K. A., Deutscher, N. M., Griffith, D. W. T., Blumenstock, T., Hase, F., Warneke, T., Wang, Z., Kivi, R., and Robinson, J.: Evaluation of column-averaged methane in models and TCCON with a focus on the stratosphere, Atmospheric Measurement Techniques, 9, 4843–4859, doi:10.5194/amt-9-4843-2016, http://www.atmos-meas-tech.net/9/4843/2016/, 2016.

Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrateammonium aerosols in the United States: Implications for policy, Journal of Geophysical Research: Atmospheres, 109, 2004.

10 Parker, R., Boesch, H., Cogan, A., Fraser, A., Feng, L., Palmer, P. I., Messerschmidt, J., Deutscher, N., Griffith, D. W. T., Notholt, J., Wennberg, P. O., and Wunch, D.: Methane observations from the Greenhouse Gases Observing SATellite: Comparison to ground-based TCCON data and model calculations, Geophysical Research Letters, 38, n/a–n/a, doi:10.1029/2011GL047871, http://dx.doi.org/10.1029/ 2011GL047871, 115807, 2011.

Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield, M. P., Corbin,

- 15 K., Fortems-Cheiney, A., Fraser, A., Gloor, E., Hess, P., Ito, A., Kawa, S. R., Law, R. M., Loh, Z., Maksyutov, S., Meng, L., Palmer, P. I., Prinn, R. G., Rigby, M., Saito, R., and Wilson, C.: TransCom model simulations of CH₄ and related species: linking transport, surface flux and chemical loss with CH₄ variability in the troposphere and lower stratosphere, Atmospheric Chemistry and Physics, 11, 12813–12837, doi:10.5194/acp-11-12813-2011, http://www.atmos-chem-phys.net/11/12813/2011/, 2011.
- Pickett-Heaps, C. A., Jacob, D. J., Wecht, K. J., Kort, E. A., Wofsy, S. C., Diskin, G. S., Worthy, D. E. J., Kaplan, J. O., Bey, I., and Drevet, J.:
- 20 Magnitude and seasonality of wetland methane emissions from the Hudson Bay Lowlands (Canada), Atmospheric Chemistry and Physics, 11, 3773–3779, doi:10.5194/acp-11-3773-2011, http://www.atmos-chem-phys.net/11/3773/2011/, 2011.
 - Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, Journal of Geophysical Research, 108, 4116, doi:10.1029/2002JD002299, 2003.

Saad, K. M., Wunch, D., Toon, G. C., Bernath, P., Boone, C., Connor, B., Deutscher, N. M., Griffith, D. W. T., Kivi, R., Notholt, J., Roehl,

- 25 C., Schneider, M., Sherlock, V., and Wennberg, P. O.: Derivation of tropospheric methane from TCCON CH₄ and HF total column observations, Atmospheric Measurement Techniques, 7, 2907–2918, doi:10.5194/amt-7-2907-2014, http://www.atmos-meas-tech.net/7/ 2907/2014/, 2014.
 - Santoni, G. W., Daube, B. C., Kort, E. A., Jimenez, R., Park, S., Pittman, J. V., Gottlieb, E., Xiang, B., Zahniser, M. S., Nelson, D. D., McManus, J. B., Peischl, J., Ryerson, T. B., Holloway, J. S., Andrews, A. E., Sweeney, C., Hall, B., Hintsa, E. J., Moore, F. L., Elkins,
- 30 J. W., Hurst, D. F., Stephens, B. B., Bent, J., and Wofsy, S. C.: Evaluation of the airborne quantum cascade laser spectrometer (QCLS) measurements of the carbon and greenhouse gas suite - CO2, CH4, N2O, and CO - during the CalNex and HIPPO campaigns, Atmospheric Measurement Techniques, 7, 1509–1526, doi:10.5194/amt-7-1509-2014, 2014.
 - Sepúlveda, E., Schneider, M., Hase, F., Barthlott, S., Dubravica, D., García, O. E., Gomez-Pelaez, A., González, Y., Guerra, J. C., Gisi, M., Kohlhepp, R., Dohe, S., Blumenstock, T., Strong, K., Weaver, D., Palm, M., Sadeghi, A., Deutscher, N. M., Warneke, T., Notholt, J., Jones,
- 35 N., Griffith, D. W. T., Smale, D., Brailsford, G. W., Robinson, J., Meinhardt, F., Steinbacher, M., Aalto, T., and Worthy, D.: Tropospheric CH₄ signals as observed by NDACC FTIR at globally distributed sites and comparison to GAW surface in situ measurements, Atmospheric Measurement Techniques, 7, 2337–2360, doi:10.5194/amt-7-2337-2014, http://www.atmos-meas-tech.net/7/2337/2014/, 2014.

- Sheese, P. E., Boone, C. D., and Walker, K. A.: Detecting physically unrealistic outliers in ACE-FTS atmospheric measurements, Atmospheric Measurement Techniques, 8, 741–750, doi:10.5194/amt-8-741-2015, http://www.atmos-meas-tech.net/8/741/2015/, 2015.
- Sherlock, V., Connor, B., Robinson, J., Shiona, H., Smale, D., and Pollard, D.: TCCON data from Lauder, New Zealand, 120HR, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.lauder01.R0/1149293, doi:10.14291/tccon.ggg2014.lauder01.R0/1149293, TC-
- 5 CON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014a.
 - Sherlock, V., Connor, B., Robinson, J., Shiona, H., Smale, D., and Pollard, D.: TCCON data from Lauder, New Zealand, 125HR, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.lauder02.R0/1149298, doi:10.14291/tccon.ggg2014.lauder02.R0/1149298, TC-CON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee,
- 10 U.S.A., 2014b.
 - Stephens, B. B., Gurney, K. R., Tans, P. P., Sweeney, C., Peters, W., Bruhwiler, L., Ciais, P., Ramonet, M., Bousquet, P., Nakazawa, T., Aoki, S., Machida, T., Inoue, G., Vinnichenko, N., Lloyd, J., Jordan, A., Heimann, M., Shibistova, O., Langenfelds, R. L., Steele, L. P., Francey, R. J., and Denning, A. S.: Weak Northern and Strong Tropical Land Carbon Uptake from Vertical Profiles of Atmospheric CO2, Science, 316, 1732–1735, doi:10.1126/science.1137004, http://science.sciencemag.org/content/316/5832/1732, 2007.
- 15 Sussmann. R. and Rettinger. M.: TCCON data from Garmisch. Germany. Release GGG2014R0. http://dx.doi.org/10.14291/tccon.ggg2014.garmisch01.R0/1149299, doi:10.14291/tccon.ggg2014.garmisch01.R0/1149299, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
- Toon, G. C. stand-alone profile GGG2014, and Wunch. D.: А priori generation tool for а 20 http://dx.doi.org/10.14291/tccon.ggg2014.priors.R0/1221661, doi:10.14291/tccon.ggg2014.priors.R0/1221661, 2014.
- Turner, A. J., Jacob, D. J., Wecht, K. J., Maasakkers, J. D., Lundgren, E., Andrews, A. E., Biraud, S. C., Boesch, H., Bowman, K. W., Deutscher, N. M., Dubey, M. K., Griffith, D. W. T., Hase, F., Kuze, A., Notholt, J., Ohyama, H., Parker, R., Payne, V. H., Sussmann, R., Sweeney, C., Velazco, V. A., Warneke, T., Wennberg, P. O., and Wunch, D.: Estimating global and North American methane emissions with high spatial resolution using GOSAT satellite data, Atmospheric Chemistry and Physics, 15, 7049–7069, doi:10.5194/acp-15-7049-2015, http://www.function.org/10.5194/acp-15-7049-2015/
- 25 http://www.atmos-chem-phys.net/15/7049/2015/, 2015.
 - van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), Atmospheric Chemistry and Physics, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, http://www.atmos-chem-phys.net/10/11707/2010/, 2010.
- 30 von Clarmann, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P., and Versick, S.: Retrieval of temperature, H₂0, O₃, HNO₃, CH₄, N₂0, ClONO₂ and ClO from MIPAS reduced resolution nominal mode limb emission measurements, Atmospheric Measurement Techniques, 2, 159–175, doi:10.5194/amt-2-159-2009, http://www.atmos-meas-tech.net/2/159/2009/, 2009.
- Wang, J., Logan, J., McElroy, M., Duncan, B., Megretskaia, I., and Yantosca, R.: A 3-D model analysis of the slowdown and interannual
 variability in the methane growth rate from 1988 to 1997, GLOBAL BIOGEOCHEMICAL CYCLES, 18, doi:10.1029/2003GB002180, 2004.

- Wang, Z., Deutscher, N. M., Warneke, T., Notholt, J., Dils, B., Griffith, D. W. T., Schmidt, M., Ramonet, M., and Gerbig, C.: Retrieval of tropospheric column-averaged CH₄ mole fraction by solar absorption FTIR-spectrometry using N₂O as a proxy, Atmospheric Measurement Techniques, 7, 3295–3305, doi:10.5194/amt-7-3295-2014, http://www.atmos-meas-tech.net/7/3295/2014/, 2014.
- Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N., Petri, C., Grupe, P., Vuillemin, C., Truong,
- 5 F., Schmidt, M., Ramonet, M., and Parmentier, E.: TCCON data from Orleans, France, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.orleans01.R0/1149276, doi:10.14291/tccon.ggg2014.orleans01.R0/1149276, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014.
 - Washenfelder, R. A., Wennberg, P. O., and Toon, G. C.: Tropospheric methane retrieved from ground-based near-IR solar absorption spectra,
- 10 Geophysical Research Letters, 30, 2226, doi:10.1029/2003GL017969, http://doi.wiley.com/10.1029/2003GL017969http://dx.doi.org/10. 1029/2003GL017969, 2003.
 - Washenfelder, R. A., Toon, G. C., Blavier, J.-F., Yang, Z., Allen, N. T., Wennberg, P. O., Vay, S. A., Matross, D. M., and Daube, B. C.: Carbon dioxide column abundances at the Wisconsin Tall Tower site, Journal of Geophysical Research: Atmospheres, 111, D22 305, doi:10.1029/2006JD007154, http://dx.doi.org/10.1029/2006JD007154, 2006.
- 15 Waugh, D. and Hall, T.: Age of Stratospheric Air: Theory, Observations, and Models, Reviews of Geophysics, 40, 1–1–1–26, doi:10.1029/2000RG000101, http://dx.doi.org/10.1029/2000RG000101, 1010, 2002.
 - Waymark, C., Walker, K., Boone, C., and Bernath, P.: ACE-FTS version 3.0 data set: validation and data processing update, Annals of Geophysics, 56, http://www.annalsofgeophysics.eu/index.php/annals/article/view/6339, 2014.
- Wecht, K. J., Jacob, D. J., Frankenberg, C., Jiang, Z., and Blake, D. R.: Mapping of North American methane emissions with high
- 20 spatial resolution by inversion of SCIAMACHY satellite data, Journal of Geophysical Research: Atmospheres, 119, 7741–7756, doi:10.1002/2014JD021551, http://dx.doi.org/10.1002/2014JD021551, 2014JD021551, 2014a.
 - Wecht, K. J., Jacob, D. J., Sulprizio, M. P., Santoni, G. W., Wofsy, S. C., Parker, R., Bösch, H., and Worden, J.: Spatially resolving methane emissions in California: constraints from the CalNex aircraft campaign and from present (GOSAT, TES) and future (TROPOMI, geostationary) satellite observations, Atmospheric Chemistry and Physics, 14, 8173–8184, doi:10.5194/acp-14-8173-2014,
- 25 http://www.atmos-chem-phys.net/14/8173/2014/, 2014b.
 - Wennberg, P. O., Roehl, C., Blavier, J.-F., Wunch, D., Landeros, J., and Allen, N.: TCCON data from Jet Propulsion Laboratory, Pasadena, California, USA, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.jpl02.R0/1149297, doi:10.14291/tccon.ggg2014.jpl02.R0/1149297, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014a.
- 30 Wennberg, P. O., Roehl, C., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R., Keppel-Aleks, G., Allen, N., and Ayers, J.: TC-CON data from Park Falls, Wisconsin, USA, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.parkfalls01.R0/1149161, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014b.
- Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., Allen, N., Dowell, P., Teske, K., Martin, C., and Martin, J.:
- 35 TCCON data from Lamont, Oklahoma, USA, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.lamont01.R0/1149159, doi:10.14291/tccon.ggg2014.lamont01.R0/1149159, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014c.

- Wennberg, P. O., Wunch, D., Yavin, Y., Toon, G. C., Blavier, J.-F., Allen, N., and Keppel-Aleks, G.: TCCON data from Jet Propulsion Laboratory, Pasadena, California, USA, Release GGG2014R0, http://dx.doi.org/10.14291/tccon.ggg2014.jpl01.R0/1149163, doi:10.14291/tccon.ggg2014.jpl01.R0/1149163, TCCON data archive, hosted by the Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2014d.
- 5 Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jiménez, R., Keppel-Aleks, G., Kort, E. a., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. a.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, Atmospheric Measurement Techniques, 3,
- 10 1351–1362, doi:10.5194/amt-3-1351-2010, http://www.atmos-meas-tech.net/3/1351/2010/, 2010.
 - Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. a., Notholt, J., Connor, B. J., Griffith, D. W. T., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, Philosophical transactions. Series A, Mathematical, physical, and engineering sciences, 369, 2087–2112, doi:10.1098/rsta.2010.0240, http://www.ncbi.nlm.nih.gov/pubmed/21502178, 2011a.

Wunch, D., Wennberg, P. O., Toon, G. C., Connor, B. J., Fisher, B., Osterman, G. B., Frankenberg, C., Mandrake, L., O'Dell, C., Ahonen, P.,

- Biraud, S. C., Castano, R., Cressie, N., Crisp, D., Deutscher, N. M., Eldering, a., Fisher, M. L., Griffith, D. W. T., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyrö, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C. E., Morino, I., Notholt, J., Oyafuso, F. a., Rettinger, M., Robinson, J., Roehl, C. M., Salawitch, R. J., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D. R., Uchino, O., Warneke, T., and Wofsy, S. C.: A method for evaluating bias in global measurements of CO2 total columns from space, Atmospheric Chemistry and Physics, 11, 12317–12337, doi:10.5194/acp-11-12317-2011, http://www.atmos-chem-phys.net/11/12317/
- 20 2011/acp-11-12317-2011.htmlhttp://www.atmos-chem-phys.net/11/12317/2011/, 2011b.
- Wunch, D., Toon, G. C., Sherlock, V., Deutscher, N. M., Liu, C., Feist, D. G., and Wennberg, P. O.: The Total Carbon Column Observing Network's GGG2014 Data Version, Tech. rep., Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A. doi: 10.14291/tccon.ggg2014.documentation.R0/1221662, doi:10.14291/tccon.ggg2014.documentation.R0/1221662, http://dx.doi.org/10.14291/tccon.ggg2014.documentation.R0/1221662, 2015.
- 25 Yang, Z., Washenfelder, R. a., Keppel-Aleks, G., Krakauer, N. Y., Randerson, J. T., Tans, P. P., Sweeney, C., and Wennberg, P. O.: New constraints on Northern Hemisphere growing season net flux, Geophysical Research Letters, 34, L12 807, doi:10.1029/2007GL029742, http://doi.wiley.com/10.1029/2007GL029742, 2007.

Run Name	Description	CH ₄ Li	fetime (years) with re	ears) with respect to	
		Emissions	Tropospheric OH	Total Loss	Burden (Tg)
Base	Default OH and Emissions	9.6	10.7	9.7	4825
Aseasonal	Constant Monthly Emission Rates	9.6	10.7	9.7	4872
Updated OH	Monthly OH fields from Standard	8.5	9.4	8.6	4828
	Chemistry + Biogenic VOCs, scaled				
	down by 10%				
Unscaled Updated OH	Monthly OH fields from Standard	7.7	8.4	7.8	4917
	Chemistry + Biogenic VOCs				
90% OH	Default OH scaled down by 10%	10.5	11.9	10.7	5296
110% OH	Default OH scaled up by 10%	8.8	9.7	8.8	4425
Scaled Rice Emissions	Rice Emissions Increased by 20%	9.6	10.7	9.6	4780
No Wetlands	Wetlands Wetland Emissions Turned Off		10.6	9.5	3768
Scaled Livestock Emissions	Iled Livestock Emissions Scale livestock emissions by 50%		10.7	9.6	4359
MERRA	MERRA meteorology fields	9.6	10.7	9.6	4849
Tropopause Level	Set top of troposphere 2 vertical levels	9.6	10.6	9.6	4855
	higher				

Table 3. List of Sensitivity Experiments