Answer to Dr. Lamont Poole

The authors are grateful for the time and thought that Lamont Poole put into the review and comments regarding our paper. We incorporate most of the comments into the revised manuscript, which has led to substantial improvements. Detailed responses to all comments follow below. The original comments from Lamont Poole are in italics and our responses as well as changes in the manuscript in plain text.

This is an interesting paper describing cirrus cloud occurrence frequencies, vertical distributions, and optical depths derived from lidar measurements at Zurich and Jungfraujoch Research Station, Switzerland, and Jülich, Germany. These results are compared with those from some earlier studies and are also used in a simple radiative transfer model to compute shortwave, longwave, and net cirrus radiative forcings. The paper is generally well written and the results are presented rather clearly. I do have a number of specific comments that the authors need to address before the paper is published in ACP.

Specific comments

The authors either are not aware of or have ignored some earlier papers describing groundbased lidar measurements of cirrus clouds obtained during the ECLIPS (Experimental Cloud Lidar Pilot Study) program. These papers include Platt et al., Bull.Amer.Met.Soc., 75, p.1635, 1994; Vaughan and Winker, Atmos.Res., 34, p.117, 1994; and Pal et al., J. Appl.Met., 34, p.2388, 1995. The authors should also mention how their new results compare with findings from these papers.

Response:

Thanks for your comment. We were not aware of these particular publications. We have added some of the results from these papers to our manuscript and compared them to our measurements.

Changes in the revised manuscript are marked in blue:

Lines 24 on page 2:

Lidar (LIght Detection And Ranging) measurements can be used to establish long time series of aerosol or cloud measurements (e.g., Platt et al., 1994).

Lines 2-3 on page 12:

agreeing well with the CALIPSO measurements discussed by Sassen et al. (2008) and being slightly smaller than the 18-19% measured during the ECLIPS campaign by Winker and Vaughan (1994).

Lines 11-12 on page 15:

These τ values agree well with the ECLIPS-campaign (Pal et al., 1995), where most detected cirrus clouds had optical depths smaller than 0.1.

Pages 4-5: There is no discussion of the possibility of cross-talk between the co-polarized and cross-polarized channels of the lidar and the effect that might have on any results.

Response:

We have added information about the possibility of cross-talk to the manuscript.

Changes in the revised manuscript on page 5, lines 1-6 are marked in blue:

We assume an ideal lidar system, which means that there is no cross-talk between the copolarized and the cross-polarized channels. Rolf (2012) has examined this for the lidar used in Jülich. He found that for the parallel detector every 2000th detected photon is actual perpendicular polarized and for the perpendicular detector about every 570 detected photon is parallel polarized. While this justifies our assumption of an ideal system for the Jülich lidar, we found considerable cross-talk in the Swiss lidar, depending on certain maintenance conditions. However, cross-talk influences in particular the perpendicular channel which we use mainly for cloud detection but not for optical depth retrieval.

Page 5, line 26: The particulate lidar ratio can also be determined directly from high-spectral resolution lidar (HSRL) measurements.

Response:

Thanks for this remark, we have added this information to the manuscript.

Changes in the revised manuscript on page 6, lines 2-3 are marked in blue:

It can be obtained directly from Raman lidars that allow for an independent measurement of particle extinction and backscatter coefficients (Cooney, 1972; Giannakaki et al., 2007; Radlach et al., 2008; Reichardt et al., 2002; Achtert et al., 2013) as well as from high-spectral resolution lidar (HSRL) measurements (e.g., Burton et al., 2012).

Page 6, lines 17-18: It is not clear how the total uncertainty is computed. I don't think it should be the "sum" of the individual contributions as stated here. Is it the square root of the sum of the squares (RSS) of the individual contributions?

Response:

We combined the uncertainties in such a way that we assess the "worst case" of uncertainty. We have added this information to the paper with the remark, that a Gaussian error would be smaller.

Changes in the revised manuscript on page 6, lines 25-27 are marked in blue:

To assess the total, maximum uncertainty, we combine the individual contributions to provide an upper bound of the uncertainty. We calculate the largest possible error, which usually is larger than the error calculated by a Gaussian error (square root of the sum of the squares (RSS) of the individual contributions).

Page 7, line 1: Is the boxcar filter a moving average boxcar?

Response:

Yes, this information has been added to the manuscript.

Changes in the revised manuscript on page 7, line 9 are marked in blue:

"further smoothed using a moving average boxcar filter"

Page 7, line 24: What is meant by "a set of lidar ratios (5:5:40)"?

Response:

We use lidar ratios between 5 and 40, with steps of 5 in between. The sentence has been changed in the manuscript.

Changes in the revised manuscript on page 7, line 32 are marked in blue:

"lidar ratios between 5 and 40 sr, in steps of 5 sr ..."

Page 7, line 31: Why is the temperature -38° C used to ensure pure ice clouds? Can the authors provide references?

Response:

We chose this threshold as it is the threshold for homogeneous nucleation to take place. We ensure that we detect pure ice clouds and exclude the mixed-phase clouds. We have added references to the manuscript.

Changes in the revised manuscript on page 8, lines 8-9 are marked in blue:

"Temperature has to be lower than -38°C (e.g., Pruppacher and Klett, 1997; Koop et al., 2000; Krämer et al., 2016) to ensure pure ice clouds and avoid detecting mixed-phase clouds (this..."

Page 12, Table 2, footnote (6): The text is confusing as written. Did the authors intend to say that relative uncertainties in their mean optical depths are comparable to "monthly mean values of 10-20% from ISCCP"?

Response:

The mean values of tau compare reasonably well with monthly mean tau values of 0.1-0.2 from ISCCP. We have changed the sentence accordingly in the manuscript.

Changes in the revised manuscript on page 13, table 2, footnote 6 are marked in blue:

"Mean values of τ compare reasonably well with monthly mean τ values of 0.1-0.2 from ISCCP..."

Page 14, lines 9-19: It would be good if the authors did some statistical analysis on the optical depth distributions in Figure 4 and could state whether the various distributions are significantly different from a statistical point of view.

Response:

We have used a Wilcoxon rank sum test to test this. We find p-values smaller than 1e-9, which clearly indicates that the distributions are significantly different.

Changes in the revised manuscript on page 15, lines 12-13 are marked in blue:

A Wilcoxon Rank sum test reveals that the optical depth distributions of the different sites are significantly different from each other.

Page 20, line 3: From Table 3, I conclude that CRFSW at 50°N from ISCCP is about an order of magnitude than the present results, but the CRFLW at 50° N from ISCCP is only a factor of 1.5-3 larger.

Response:

Thank you for pointing this out. We have provided this information to the manuscript and also added some explanations.

Changes in the revised manuscript on page 21, lines 15-16 and page 22, line 1 are marked in blue:

"1.5-3 times larger radiative forcing in the longwave, CRF_{LW} , and one order of magnitude larger radiative forcing in the shortwave, CRF_{SW} . The difference in the CRF_{SW} can only be explained in terms of a..."

Page 23, lines 8-10: I don't understand what is meant by "radiative forcing of the lateral boundary" of cirrus clouds? It would be good if the authors could provide a brief explanation.

Response:

In this cited paper the authors (Li et al., 2014) discussed the radiative effect of observed cirrus cloud edges. With lateral boundary they describe the first 10 km horizontal from outside cirrus clouds with optical depth of 0 towards small optical depth less than 0.3. In this transition region, they found still a CRF_{LW} of 10 W/m2.

This information has been added to the manuscript.

Changes in the revised manuscript on page 23, lines 29-31 are marked in blue:

The radiative effect of observed cirrus cloud edges is discussed. In the transition region of large cirrus, defined as their optically thin rim ($\tau < 0.3$), which is often missed by satellite passive optical sensors such as MODIS, the CRFLW found to be still substantial (~ 10 Wm⁻²).

Page 23, lines 22-23: I don't understand the last sentence of this paragraph. What did the "close examination of CRFNET" with respect to cloud τ show?

Response:

Thank you for pointing to this ambiguity. We have removed the sentence, since it belongs to the paragraph below and is repeated there.

Changes in the revised manuscript on page 25, lines 8-9 are marked in blue:

The sentence was removed.

Technical Corrections

Page 1, line 2: It would be better to say that cirrus "...affect the water vapor budget ..." not determine it.

Page 1, line 15: Reword to say "... thus enabling lidar measurements of higher ..."

Page 2, line 3: The word "subvisible" is misspelled.

Page 20, line 9: Reword sentence to say "Cirrostratus clouds with $\tau < 3.6$ occur particularly in this altitude range."

Page 26, line 1: The word "subvisible" is misspelled again.

Response:

These corrections have all been implemented.

Answer to Anonymous Referee 2

The authors are grateful for the time and thought that Anonymous Referee 2 put into the review and comments regarding our paper. We incorporate most of those comments into our revised manuscript, which has led to substantial improvements. Detailed responses to all comments follow below. The original comments from Anonymous Referee 2 are in italics and our responses as well as changes in the manuscript in plain text.

In this article, an algorithm is developed (FLICA) to retrieve cirrus cloud properties based on lidar measurements at three stations in Europe. Using these retrievals, a cirrus climatology and cirrus radiative forcing in each station are presented. Differences of cirrus at three locations are discussed. Subvisual, thin and opaque cirrus are analyzed. Results are also compared to previous studies, and the differences with results by Chen et al. (2000) are particularly emphasized. This paper is generally completed and well written. My main comments/questions relate to the section of comparisons with previous studies, and methods to calculate ice cloud radiative forcing.

Specific comments.

1. Title of this study, 'Radiative properties of ...', since a climatology of cirrus is also an important part of this study, is it better to say 'Climatological and radiative properties of...'

Response:

This is an excellent idea. We have changed the title accordingly.

2. Aerosols:

How the algorithm distinguish aerosols and cirrus in this article? Cirrus clouds with small optical depth (e.g. tau<0.001) look more like aerosols?

Response:

FLICA uses a criterion for depolarization to help preventing that aerosols are classified as cirrus clouds. As we have already stated in the manuscript, to ensure that the highest cirrus clouds observed above Jungfraujoch were not volcanic ash particles (which would also depolarize the backscattered light), we have examined satellite measurements and found no indication for volcanic influence. Furthermore, our temperature threshold of -38°C excludes most aerosols because most aerosol layers are located below 6 km. Even though this combination of criteria might not be perfect, we believe that cirrus clouds are clearly distinguished from aerosols in the very most cases.

Changes in the manuscript:

No changes have been applied to the manuscript.

3. Page 15. Line 5. How do you make sure such a small optical depth (< 5x10-4) is not resulted from noises or from aerosols?

Response:

Areas of 3×3 pixels are examined, with the pixel to be checked for cloudiness in the center. At least 8 of the 9 examined pixels have to have a volume depolarization larger than 0.007(0.006) and a BSR larger than 1.08(1.03) for day(night)-time measurements. As mentioned above, the output of the cloud detection scheme was in addition visually inspected for individual days and was found not to show any apparent artifacts (such as aerosol layers or noise).

Changes in the manuscript:

No changes have been applied to the manuscript.

4. Page 17, line 25, The mean solar zenith angle for three locations is 60°. However, JUL and JFJ are about 4 degrees latitude off, and thus the mean length of daytime (mean SZA) in the two locations should also be quite different, which will cause radiative flux biases. Have you ever check the differences?

Response:

Thank you for pointing out this issue. The differences in latitude between the three stations have two consequences. First, the length of daytime is different for the stations. This value is however taken directly from the measurements (see line 28 on page 19). Second, the differences in the solar angle affect the incident radiation. The overall difference in the incident solar radiation is 6%. Nevertheless, the radiation model of Corti & Peter has an accuracy that "is typically better than 20% when comparing with the accurate results from the Fu and Liou (1992, 1993) radiative transfer model", thus we consider this error to be of minor relevance for the model results. In addition, addressing as well item 5 below, we want to examine the radiative properties of the cirrus clouds as a function of optical depth and temperature under otherwise comparable conditions.

Changes in the manuscript:

No changes have been applied to the manuscript.

5. Page 17, line 25, An albedo of 0.3 is globally average planetary albedo. The mean surface albedo is about 0.15 (Kiehl and Trenberth, 1997). Also, surface albedo varies from different locations and in different seasons. In particular, JFJ is located at a high altitude and has a cold climate. How many days of this location will be covered by snow in a year? Surface albedo covered by snow is large (> 50%).

Response:

Jungfraujoch lies on top of a glacier and is all year covered by snow. We have made calculations using albedos of snow (0.65) for Jungfraujoch. If we use a snow-albedo, the radiative effect of the cirrus clouds disappear as all radiation is scattered back by the snow surface. As already mentioned in reply to item Ulrich Schumann, we chose a value of 0.3 to demonstrate the global average effect of cirrus clouds such as those detected above the three locations.

Changes in the manuscript:

No changes have been applied to the manuscript.

6. Page 17. Line 29: The extinction coefficient can be derived from radar backscattering and then optical depth is obtained as shown in equation 7 in this paper. The tau values are used to calculate ice cloud radiative effect. Have you look at how different cirrus radiative effect

will be if the extinction coefficient profile is used? The profiles characterize the vertical details of a cloud, which are more accurate to produce radiative fluxes (Chen 2000).

Response:

For this work, we have chosen to focus on the model of Corti and Peter, which uses the optical depth and cloud temperature as input (besides the surface albedo. This model has an accuracy better than 20% as compared to the model of Fu and Liou. We added this information to the manuscript.

Sentence added to the manuscript on page 19, line 13-14 in blue:

The accuracy of Corti and Peter (2009) is better than 20 % in comparison with the Fu-Liou model.

7. How the asymmetric factor and single-scattering albedo of the clouds are determined?

Response:

The asymmetry factor determines the forward scattering of a cloud and thus also the reflectivity. The reflectivity is approximated depending on optical depth and a fixed value as a result of radiative transfer calculations using the Fu-Liou code. For more detail see Corti and Peter, 2009. p. 5755.

Our value for single scattering albedo is $\omega_0 = 0.55$, which is a realistic estimate for longwave radiation (Stephens et al., 1990). The exact value is not decisive for our parameterization, since variations of 10 % in this parameter increases the mean error of CRF_{LW} in comparision with radiative transfer calculations only by about 1%. Again, for more detail see Corti and Peter, 2009. p. 5754.

Changes in the manuscript:

No changes have been applied to the manuscript.

8. Section 4.2, Comparisons with previous results

Several paragraphs in Section 4.2 describe how results in this study differ from Chen et al. 2000. I doubt the way of comparisons for the following reasons. 1) different definitions of cirrus as stated in this article, and thus radiative forcing of cirrus is different. 2) different resolutions, it is 280 km resolution in Chen et al. 2000, while in this study, lidar has a small field of view (1.5 mrad by 0.3 mrad); 3) time period is very different (four days in Chen's study and five years in this study); 4) More importantly, although the three stations are located around 50°N, comparing ice cloud properties to zonally average values at 50°N provided by Chen et al.2000 is unreasonable since ice clouds vary significantly around 50° (e.g. Sassen et al. 2008). I'm confused why this section is necessary. Do you want to check that your results are correct? If so, why not compare to CERES fluxes? Could you justify the reasons why such comparison is necessary in this study?

Response:

We wanted to compare our data with similar measurements. There are only few publications assessing the radiative effect of mid-latitude cirrus clouds and the paper by Chen et al. is one of them. We are aware (and mention extensively in section 4.2) that it is very difficult to compare our and Chen's results due to the reasons you mention. Concerning a comparison

with CERES fluxes, we are not aware of a publication discussing cloud radiative forcings derived from the CERES data for mid-latitude cirrus. Conversely, establishing CRF for cirrus from CERES data is beyond the scope of the present investigation.

Changes in the manuscript:

No changes have been applied to the manuscript.

Besides Chen et al. 2000, this article also lists a series of related studies in Section 4.2 (page 21, lines9-30). It may be better to move these paragraphs to introduction.

Response:

After consultation with all co-authors we decided to keep the information at its origin position in the article as we think it is most suitable where it is.

9. Page 25, line 11: ' cirrus clouds, which remain undetected by satellites (requiring typically tau>0.2)...', CALIPSO lidar can detect ice clouds with tau< 0.2. It would be better to revise the satellites as ' passive remote sensing'. It'll be better if a related reference added after tau> 0.2.

Response:

We agree and have adapted this in the manuscript, reading now "passive remote-sensing satellites".

Changes in the manuscript on page 28, line 4 in blue:

"by passive remote-sensing satellites"

Reply to the discussion comment submitted by Prof. Ulrich Schumann

The authors are grateful for the time and thought that Ulrich Schumann put into the comments regarding our paper. Subsequently we show the original comments from Ulrich Schumann in italics and our responses as well as changes in the manuscript in plain text.

We appreciate that Mr. Schumann highlights the occurrence of contrail cirrus as an important point, especially over the European continent, which is subject of strong air traffic. Concerning this aspect, we of course have to admit that the developed data analysis scheme, FLICA, is not intended to separate contrail from natural cirrus. Such a separation would require – according to our understanding – the incorporation of side information from either detailed flight maps or satellite imagery to identify periods of strong contrail formation. Such an analysis was outside the scope of this study. In a possible future follow-up study a separation from natural and contrail cirrus could become an important component.

We decided to mention the topic in the conclusions of the manuscript, highlighting the potential of such long-term datasets for the characterization of anthropogenic effects on cirrus cloud properties. We also incorporated some of the references provided by Prof. Schumann.

This is a nice study of thin cirrus over 3 stations in the Alps and Northern Germany.

1) Which fraction of the thin cirrus originates from contrail cirrus? Liou et al. [1990], e.g., noted a strong increase of thin cirrus over Salt Lake City since about the late 1960's in correlation with increases in jet traffic. The stations are located in regions where line-shaped contrails are ubiquitous [Mannstein et al., 1999; Meyer et al., 2002]. The stations are located near the routes from London to the Near East or the routes from or across Paris to the Far East etc. (see contrail cover results and major traffic routes in Fig. 7 in [Schumann, 2005]). Often aged contrail cirrus might have gotten advected from, e.g., the routes over Lyon to the central Alps. The observed optical depth is fully consistent with optical depth for contrail cirrus from other sources [Immler et al., 2008; Iwabuchi et al., 2012; Vázquez-Navarro et al., 2015]. The computed cover and RF values are consistent with contrail cirrus calculations [Schumann et al., 2015]. Hence, it is very likely that contrails contributed a large fraction to the observed thin cirrus. So far, your nice paper, not even mentions this possibility. I think, at least that needs to be changed.

Response:

We find clear evidence of detected contrails only on one of the days. However, an uncertain number of the detected cirrus clouds may have evolved from contrails. How many is impossible to say without further, elaborate calculations. Especially, aged contrails in a water vapor supersaturated environment are hard to discriminate from natural cirrus clouds when only a stationary lidar is available. Due to turbulent mixing of contrail air with surrounding air masses and further growth of contrail ice crystals their microphysical properties become indistinguishable from those of natural cirrus. We have added a paragraph plus some references about this.

Paragraph added in revised manuscript on page 27, lines 11-20 in blue:

Owing to the central location of the three measurement sites in Europe, a significant fraction of the thin cirrus observed within the present study might actually have originated from contrails. Clear indications for the occurrence of contrails were found on at least one day, given the optically and geometrically very thin cirrus. Liou et al. (1990) noted an increase of thin cirrus with increases in jet traffic. Our measurement sites are located in a region, where line-shaped contrails are ubiquitous (Mannstein et al., 1999; Meyer et al., 2002) as many flight routes cross this area. The observed optical depths are consistent with optical depths of

contrail cirrus (Immler et al., 2008; Iwabuchi et al., 2012; Vázquez-Navarro et al., 2015). Furthermore, the cirrus cloud cover determined in the present study is consistent with contrail cirrus calculations by Schumann et al. (2015). Therefore, it is likely that contrails contributed a fraction of the observed cirrus. The determination of the actual contribution of contrails to the cirrus cloud dataset is, however, not subject of this study, considering that the applied data analysis algorithm FLICA cannot distinguish natural and contrail cirrus.

2) How important for longwave radiative forcing (RF) from thin cirrus for otherwise clear sky is the water vapor in the atmosphere below the cirrus? The longwave RF of thin cirrus correlates far better with the brightness temperature of the atmosphere than with surface temperature, see Fig. 15.4 in [Schumann et al., 2012a]. The brightness temperature is related to the outgoing longwave radiation (OLR) at top of the atmosphere, as available, e.g., from Numerical Weather Prediction(NWP) data, e.g. from COSMOS. Also: how important is the difference between Earth surface albedo and effective albedo of the Earth-Atmosphere system, e.g. when clouds are nearby the location of observations or when the mountains are snow covered or when there is any dust or haze (derivable from known solar direct radiation and from reflected shortwave radiation, RSR, also available from NWP data), as discussed in these papers? Perhaps you can quantify these effects?

Response:

We have added some remarks about the brightness temperature and its relation to the longwave radiative forcing to the manuscript and added the suggested reference.

As also mentioned in our response to reviewer 2, we chose a value of 0.3 for the underlying albedo to demonstrate, which radiative effect the detected cirrus would have, if they were located above the "mean" of the planet. Jungfraujoch is located on top of a glacier and is all year covered by snow. We have made calculations using albedos of snow (0.65) for Jungfraujoch. In that case, the radiative effect of the cirrus clouds disappears as all radiation is scattered back by the snow surface.

Changes in revised manuscript on page 19, lines 18-20 in blue:

The CRF_{LW} further correlates well with the brightness temperature of the atmosphere, which is related to the outgoing longwave radiation at top of atmosphere (Schumann et al., 2012a). This correlation has not been considered in the model of Corti and Peter (2009).

3) Why not to test the differences between the nice and simple Corti&Peter parametrization and that which we developed in parallel (see my comment of May2009 on the ACPD paper by Corti and Peter and [Schumann et al., 2012b])? The input needed (OLR and RSR) is available form COSMO and other NWP models. The model could be used to test the influence of various assumptions on particle habits and particle sizes [Markowicz and Witek, 2011]. The quantitative results may well change by50 %, and hence change your conclusions.

Response:

Thank you for this interesting remark. We have added a remark on this in the manuscript and provided the suggested references, but refrain from performing additional computations at this stage of the paper and leave this to potential follow-up projects.

Changes in revised manuscript on page 27, lines 33-35 and page 28, line 1 in blue:

Besides the radiation model of Corti and Peter (2009) used for this study, other approaches exist that can be used to investigate the effect of other cloud properties besides optical depth on the cirrus radiative forcing. For instance, the radiation model of Schumann et al. (2012b) could be used to test the influence of various assumptions on particle habits and particle sizes (Markowicz and Witek, 2011).

4) Does the Lidar signal (e,g., depolarization) allow to discriminate, perhaps together with other data, contrails from cirrus? Perhaps there are some ideas which could fit into your outlook?

Response:

We have not examined this so far. We think that it might be possible to distinguish fresh contrails from cirrus clouds as the contrails have a large number of small particles that are rather round due to the rapid cooling they were exposed to (Jensen, 1998). This would result in different depolarization values than for natural cirrus clouds. In a supersaturated environment, contrails can stay persistent for a long time and are more similar to natural cirrus clouds after growth. In a subsaturated environment the contrails will evaporate very quickly.

Changes in revised manuscript:

No changes have been made in the manuscript.

References implemented in the revised version of the paper:

Immler, F., R. Treffeisen, D. Engelbart, K. Krüger, and O. Schrems (2008), Cirrus, contrails, and ice supersaturated regions in high pressure systems at northern mid latitudes, Atmos. Chem. Phys., 8, 1689–1699, doi:10.5194/acp-8-1689-2008.

Iwabuchi, H., P. Yang, K. N. Liou, and P. Minnis (2012), Physical and optical properties of persistent contrails: Climatology and interpretation, J. Geophys. Res., 117, D06215, doi:10.1029/2011JD017020.

Liou, K. N., S. C. Ou, and G. Koenig (1990), An investigation of the climatic effect of contrail cirrus. In: Air Traffic and the Environment – Background, Tendencies and Potential Global Atmospheric Effects. U. Schumann (Ed.), Lecture Notes in Engineering, Springer Berlin, 154-169.

Mannstein, H., R. Meyer, and P. Wendling (1999), Operational detection of contrails from NOAA-AVHRR data, Int. J. Remote Sensing, 20, 1641-1660, doi: 10.1080/014311699212650.

Markowicz, K. M., and M. Witek (2011), Sensitivity study of global contrail radiative forcing due to particle shape, J. Geophys. Res., 116, D23203, doi:10.1029/2011JD016345.

Meyer, R., H. Mannstein, R. Meerkötter, U. Schumann, and P. Wendling (2002), Regional radiative forcing by line-shaped contrails derived from satellite data, J. Geophys. Res., 107, ACL 17-11 - ACL 17-15, 10.1029/2001jd000426.

Schumann, U., K. Graf, H. Mannstein, and B. Mayer (2012a), Contrails: Visible aviation induced climate impact, in Atmospheric Physics – Background - Methods - Trends, edited by U. Schumann, pp. 239-257, Springer, Berlin, Heidelberg, DOI: 10.1007/978-

3-642-30183-4_15.

Schumann, U., B. Mayer, K. Graf, and H. Mannstein (2012b), A parametric radiative forcing model for contrail cirrus, J. Appl. Meteorol. Clim., 51, 1391-1406, doi: 10.1175/JAMC-D-11-0242.1.

Schumann, U., J. E. Penner, Y. Chen, C. Zhou, and K. Graf (2015), Dehydration effects from contrails in a coupled contrail-climate model, Atmos. Chem. Phys., 15, 11179-11199, doi:10.5194/acp-15-11179-2015.

Vázquez-Navarro, M., H. Mannstein, and S. Kox (2015), Contrail life cycle and properties from 1 year of MSG/SEVIRI rapid-scan images, Atmos. Chem. Phys., 15, 8739-8749, doi:10.5194/acp-15-8739-2015.

Climatological and radiative properties of mid-latitude cirrus clouds derived by automatic evaluation of lidar measurements

Erika Kienast-Sjögren^{1,2}, Christian Rolf³, Patric Seifert⁴, Ulrich K. Krieger¹, Bei P. Luo^{1,5}, Martina Krämer³, and Thomas Peter¹

¹Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland
²Now at: Fed. Office of Meteorology and Climatology, MeteoSwiss, Zurich Airport, Operation Center 1, CH-8058 Zurich, Switzerland
³Institute for Energy and Climate Research, Stratosphere, Forschungszentrum Jülich, Jülich, Germany
⁴Institute for Tropospheric Research (TROPOS), Leipzig, Germany
⁵Physical Meteorological Observatory Davos, PMOD WRC, CH-7260 Davos, Switzerland

Correspondence to: Erika Kienast-Sjögren (Erika.Kienast@meteoswiss.ch)

Abstract. Cirrus, i.e. high thin clouds that are fully glaciated, play an important role in the Earth's radiation budget as they interact with both long- and shortwave radiation and determine affect the water vapor budget of the upper troposphere and stratosphere. Here, we present a climatology of mid-latitude cirrus clouds measured with the same type of ground-based lidar at three mid-latitude research stations: at the Swiss high alpine Jungfraujoch station (3580 m a.s.l.), in Zürich (Switzerland, 510

- 5 m a.s.l.) and in Jülich (Germany, 100 m a.s.l.). The analysis is based on 13'000 hours of measurements from 2010 2014. To automatically evaluate this extensive data set, we have developed the "Fast LIdar Cirrus Algorithm" (FLICA), which combines a pixel-based cloud-detection scheme with the classic lidar evaluation techniques. We find mean cirrus optical depths of 0.12 on Jungfraujoch and of 0.14 and 0.17 in Zürich and Jülich, respectively.
- 10 Above Jungfraujoch, subvisible cirrus clouds ($\tau < 0.03$) have been observed during 7% of the observation time, whereas above Zürich and Jülich significantly less. From Jungfraujoch, clouds with $\tau < 10^{-3}$ can be observed three times more often than over Zürich and Jülich, and clouds with $\tau < 2 \times 10^{-4}$ even ten times more often. Above Jungfraujoch, cirrus have been observed to altitudes of 14.4 km a.s.l., whereas only to about 1 km lower at the other stations. These features highlight the advantage of the high-altitude station Jungfraujoch, which is often in the free troposphere above the polluted boundary layer,
- 15 thus allowing to perform enabling lidar measurements of thinner and higher clouds. In addition, the measurements suggest a change in cloud morphology at Jungfraujoch above \sim 13 km, possibly because high particle number densities form in the observed cirrus clouds, when many ice crystals nucleate in the high supersaturations following rapid uplifts in lee waves above mountainous terrain.
- 20 The retrieved optical properties are used as input for a radiative transfer model to estimate the net cloud radiative forcing, CRF_{NET} , for the analysed cirrus clouds. All cirrus detected here have a positive CRF_{NET} . This confirms that these thin, high cirrus have a warming effect on the Earth's climate, whereas cooling clouds typically have lower cloud edges too low in altitude

to satisfy the FLICA criterion of temperatures below -38° C. We find $CRF_{NET} = 0.9 \text{ Wm}^{-2}$ for Jungfraujoch and 1.0 Wm⁻² (1.7 Wm⁻²) for Zürich (Jülich). Further, we calculate that subvisible cirrus ($\tau < 0.03$) contribute about 5%, thin cirrus (0.03 $< \tau < 0.3$) about 45% and opaque cirrus ($0.3 < \tau$) about 50% of the total cirrus radiative forcing.

1 Introduction

- 5 One of the large challenges in climate modeling, characterized by a low level of scientific understanding, are clouds and their effects on climate (Dessler and Yang, 2003; Solomon et al., 2007; Boucher et al., 2013). This concerns also the microphysical processes leading to cirrus formation. These processes are subject to uncertainties in the understanding and parametrization of homogeneous and heterogeneous nucleation (e.g., Cirisan et al., 2014). For any specific cloud scene, unless there are in situ measurements, there is either no or incomplete knowledge of the number of ice nuclei (IN), the intensity of small-scale temper-
- 10 ature fluctuations or the corresponding accurate values of upper tropospheric humidity (e.g., Ickes et al., 2015; Kienast-Sjögren et al., 2015).

Cloud properties such as cloud particle number, size and ice particle shape determine ice water content and optical depth, which together with the temperature of the cirrus cloud top determines whether the net cloud radiative forcing, CRF_{NET} , is

- 15 positive or negative, i.e. whether a particular cirrus cloud is warming or cooling (Platt and Harshvardhan, 1988; Ebert and Curry, 1992; Lin et al., 1998a; Chen et al., 2000; Corti and Peter, 2009). The fact that liquid clouds contain spherical particles helps estimating their microphysical and radiative properties. Conversely, the different shapes and orientations (Pruppacher and Klett, 1997) of ice particles affect the extinction of light, complicating the estimation of the cirrus climate effect (Fu and Liou, 1993; Liou, 2002). Previous studies of the radiative effect of cirrus (e.g., Chen et al., 2000; Fusina et al., 2007; Cziczo
- and Froyd, 2014) have identified a range of several watts per square meter (Wm^{-2}) depending on the ice crystal number in a cirrus as compared to having an ice-free supersaturated region.

Lidar (LIght Detection And Ranging) measurements can be used to establish long time series of aerosol or cloud measurements (e.g., Platt et al., 1994). From the co- and cross-polarized components of the backscattered light the profile of the
depolarization ratio can be obtained providing information about the sphericity of the retrieved particles, and thus on their liquid or solid state. Several lidar stations have applied their measurements of elastically backscattered light to investigate the properties of mid-latitude cirrus clouds. See Table 1 for an overview.

Here we present a cirrus cloud climatology based on 13'000 hours of lidar measurements from three mid-latitude sites;
Jungfraujoch, Zürich and Jülich. The lidar technique is briefly described in Section 2.1. In Section 2.2, the newly developed evaluation algorithm FLICA is presented. Using FLICA we are able to analyze extensive lidar measurements automatically. The climatology of this data is presented in Section 3. We then apply the radiative transfer model of Corti and Peter (2009) to estimate the cloud radiative forcing caused by the detected cirrus clouds in Section 4. The results are compared to previous

Measurement	Location	Altitude	Observation	Wavelength	Hours	References
site		[m a.s.l.]	Period	[mn]	of data	
Salt Lake City	42°N, 68°W	1726	1986-1996	694	2200	Sassen and Campbell (2001); Sassen and Benson (2001)
USA						Sassen and Comstock (2001); Sassen et al. (2003, 2007)
Punta Arenas	53°S, 71°W	126	03-04 2000	355, 532	71	Immler and Schrems (2002)
Chile						
Prestwick	56°N, 5°W	7	9-10 2000	355, 532	74	Immler and Schrems (2002)
Scotland						
Haute Provence	44°N, 6°E	679	1997-2012	532,1064	~ 7000	Goldfarb et al. (2001); Hoareau et al. (2013)
France						
Rome Tor Vergata	42°N, 13°E	107	2007-2010	532	500	Dionisi et al. (2013)
Italy						
Clermont-Ferrand	46°N, 3°E	420	2008-2014	355	~ 2000	Fréville et al. (2015)
France						
Seoul	37°N, 127°E	116	2006-2009	532, 1064	~ 1000	Kim et al. (2014)
South Korea						
Jülich	51° N, 6° E	95	2011-2013	355	3274	This work, also Rolf (2012)
Germany						
Zürich	47°N, 9°E	509	2010-2013	355	4678	This work
Switzerland						
Jungfraujoch	47° N, 8° E	3580	2010-2014	355	5170	This work
Switzerland						

Table 1. Lidar stations that have been used for systematic climatological studies of cirrus clouds in the mid-latitudes

studies in Subsection 4.2. The influence of the thinnest, subvisible, cirrus clouds on the cirrus radiative forcing is examined in Subsection 4.3. Finally, the main findings are summarized in Section 5.

2 Lidar

2.1 Lidar technique

5 This work uses the commercially available elastic backscatter lidar Leosphere ALS 450. This lidar emits linearly polarized laser pulses with an energy of 16 mJ at a wavelength of 355 nm and a repetition rate of 20 Hz. The full-angle field of view of the receiver telescope and the laser beam divergence are 1.5 mrad and 0.3 mrad, respectively.

The Nd:YAG laser of the ALS 450 is powered by a flash lamp. The flash lamp has lifetime corresponding to 5×10^7 shots or

- 10 694 hours or a month of continuous operation. In order to save flash lamp lifetime, the ALS 450 operated at Zürich and on the Jungfraujoch was coupled to a Vaisala Ceilometer CL31, which is a simple, low-maintenance elastic backscatter lidar (with a pulse energy about three orders of magnitude lower than the ALS 450). We use the ceilometer to detect thick clouds at low altitudes. Once thick clouds are present at an altitude lower than 1 km above the station, the lidar is automatically switched off (this is the case at roughly 30-40% of the time), and it is automatically switched back on once the low-level clouds are gone. In
- 15 Jülich, where no ceilometer was available, the ALS 450 was operated manually and switched off and on after visual inspection.

The range-corrected signal $r^2 P(r)$ detected by the ALS 450 can be described with the lidar equation (Kovalev and Eichinger, 2004; Wandinger, 2005):

$$r^{2}P(r) = C \times O(r)[\beta_{m}(r) + \beta_{p}(r)] \exp\left(-2\int_{r_{0}}^{r} [\alpha_{m}(r') + \alpha_{p}(r')] dr'\right).$$
(1)

- 20 where β_m and β_p describe the backscatter from molecules and particles and α_m and α_p specify molecular and particulate extinctions, i.e. light attenuation by scattering and absorption, and take changes of scatterer density with altitude into account. Instrumental properties are described by the constant *C*. O(r) is the overlap function which describes the overlap between the laser footprint and the telescope field of view. For the ALS 450 the complete overlap is achieved at a distance of 450 m from the lidar. As we analyze cirrus clouds that occurred entirely at greater heights above the lidar, we do not need to consider the 25 overlap function
- 25 overlap function.

The Leosphere ALS 450 measures the co- and cross-polarized components of the return signal. In order to solve equation 1 it is required to obtain the total signal from these two components. We calculate the total signal based on both channels as described by Rolf (2012).

30

From the detected co- and cross-polarized signal components the depolarization ratio can be obtained (Schotland et al., 1971).

We assume an ideal lidar system, which means that there is no cross-talk between the co-polarized and the cross-polarized channels. Rolf (2012) has examined this for the lidar used in Jülich. He found that for the parallel detector every 2000th detected photon is actual perpendicular polarized and for the perpendicular detector about every 570 detected photon is parallel polarized. While this justifies our assumption of an ideal system for the Jülich lidar, we found considerable cross-talk in the

- 5 Swiss lidar, depending on certain maintenance conditions. However, cross-talk influences in particular the perpendicular channel which we use mainly for cloud detection but not for optical depth retrieval. Light that is scattered back by non-spherical particles changes its polarization state, whereas spherical particles do not change the state of polarization of the returned light. Therefore, the depolarization ratio provides information about the sphericity of the detected particles (Schotland et al., 1971; Kovalev and Eichinger, 2004). The cross-polarized signal from aspherical ice particles in thin cirrus often provides the better
- 10 contrast than the parallel signal, a property we will use in our cloud retrieval algorithm. The lidar is pointed to 5° off-zenith to avoid the effect of specular reflections of horizontally oriented ice crystal plates on the measured backscatter signal and depolarization ratio (Platt et al., 1978; Westbrook et al., 2010).

For our cloud detection scheme elaborated in Subsection 2.2, we use the backscatter ratio (BSR) defined as:

15 BSR =
$$\frac{\beta_p + \beta_m}{\beta_m}$$
. (2)

To solve the lidar equation (Eq. 1) with four unknowns (β_m , β_p , α_m and α_p) and only one measurement $r^2 P(r)$, we need to make use of best current knowledge. The molecular quantities β_m and α_m are calculated from analysis data of the numerical weather prediction model (NWP) COSMO-2. We use pressure (p) and temperature (T) from COSMO-2 (COSMO, 2015) to calculate the molecular density of air and determine β_m and α_m using Rayleigh theory (Bucholtz, 1995).

For the solution of Eq. (1) we use a lidar retrieval as described in Kovalev and Eichinger (2004). To ensure stable solutions, we use a far end boundary condition (Klett, 1981). Further, we need to define the extinction-to-backscatter ratio (hereafter referred to as lidar ratio). We derive $\epsilon = 0.234$, the anisotropy of the molecules present in the atmosphere, from Eq. (6) in She (2001) and Table 1 in Bucholtz (1995) for our lidar wavelength of 355 nm. The lidar ratio of the molecular part is evaluated as:

25
$$L_m = \frac{8\pi}{3} \times \frac{180 + 40\epsilon}{180 + 7\epsilon} \approx 8.7,$$
 (3)

where σ^R , given by Eq. (6) of She (2001), is divided by the expression for σ_{π}^C , provided in Eq. (4) of She (2001), as the receiver optical bandpass spectral width of 0.3 nm (<24 cm⁻¹ at 28170 cm⁻¹) suppresses the rotational Raman wing spectral contribution (Arshinov and Bobrovnikov, 1999). Note that our ϵ is called R_A by She (2001). The particulate lidar ratio is defined as:

$$L_p = \frac{\alpha_p}{\beta_p} \tag{4}$$

Several studies have been performed to measure the particulate lidar ratio of cirrus clouds (e.g., Ackermann, 1998; Immler and Schrems, 2002; Larchevêque et al., 2002; Seifert et al., 2007). It can be obtained directly from Raman lidars that allow for an

²⁰

independent measurement of particle extinction and backscatter coefficients (Cooney, 1972; Giannakaki et al., 2007; Radlach et al., 2008; Reichardt et al., 2002; Achtert et al., 2013) as well as from high-spectral resolution lidar (HSRL) measurements (e.g., Burton et al., 2012). In our retrieval we determine the lidar ratio such that BSR=1 above and below the cirrus cloud (e.g., Rolf, 2012).

5

The lidar equation (Eq. 1) assumes single-scattering of the emitted light in the direction 180° to the emitted direction only. In reality, this is not strictly the case. As seen in Fig. 1 in Wandinger (1998), cloud particles produce strong forward scattering. This causes some of the scattered photons to remain within the field of view of the lidar, where they can be scattered back to the lidar receiver during a subsequent scattering process. These additional backscattered photons cause an underestimation of the

10 particle extinction. The strength of multiple scattering depends mainly on the laser divergence, the telescope field of view and the effective radius of the scattering particles. In order to provide extinction values that are comparable to other lidar systems and cloud conditions, the measured apparent, multiple-scattering affected extinction coefficient α_p^{obs} needs to be corrected with the correction factor γ to obtain single-scattering related values α_p^{single} , such that

$$\alpha_{\rm p}^{\rm single} = \frac{\alpha_{\rm p}^{\rm obs}}{\gamma}.$$
(5)

15 We use the multiple scattering model by Hogan (2008) as described by Wandinger (1998) and Seifert et al. (2007) to derive γ . The effective radius of the cirrus particles is taken from a climatology provided by Wang and Sassen (2002). For particles much larger than the detection wavelength, as it is the case for ice crystals observed with lidar, about 50% of the scattering occurs into the forward direction. In this study we find an average value for γ of 0.56 for Jungfraujoch and 0.52 (0.54) for Zürich (Jülich).

The lidar retrieval poses several uncertainties. Using NWP-data to calculate the molecular properties results in a maximal error of 2 %. However, there are uncertainties pertained to the data themselves. The lidar detector counts photons, and we calculate the counting error by means of poisson statistics. The assumed lidar ratio is also an error source. Here, we use lidar ratios that deviate ± 5 sr from the determined lidar ratios to assess for the uncertainty caused by determining a lidar ratio. To assess

25 the total, maximum uncertainty, we sum up combine the individual contributions to provide an upper bound of the uncertainty. We calculate the largest possible error, which usually is larger than the error calculated by a Gaussian error (square root of the sum of the squares (RSS) of the individual contributions). Seifert et al. (2007) estimated the error in the multiple-scattering correction in the order of 10%. The signal-to-noise ratio (SNR) is determined by the variation within the 5-min-average profiles. As the Leosphere lidar does not allow to retrieve the photon counts directly from the data, we calculate the SNR from the

$$SNR = \sqrt{N} \cdot \frac{\text{mean}\left(r^2 P(r)\right)}{\text{std}\left(r^2 P(r)\right)},\tag{6}$$

where mean $(r^2 P(r))$ is the mean range-corrected signal and std $(r^2 P(r))$ the standard deviation of the originally retrieved lidar profiles over N = 600 shots (following a suggestion by P. Royer, Leosphere, private communication 11.8.2014).

²⁰

2.2 Cirrus Detection Algorithm FLICA

For an efficient evaluation of this extensive data set, the automated data evaluation algorithm FLICA (Fast LIdar Cirrus Algorithm) was developed. The algorithm is based on a classical lidar retrieval (e.g., Klett, 1981; Kovalev and Eichinger, 2004) combined with a cloud detection scheme. FLICA analyzes profiles over 5 minutes (i.e. averages over 5 min \times 60 s/min \times 20 shots/s = 6000 shots). This time range was chosen to be short enough to ensure that all clouds could be detected by the algorithm while long enough to provide profiles smooth enough for the lidar retrieval to function. The 5-min profiles of the lidar measurements are further smoothed using a moving average boxcar filter in the vertical coordinate over 150 m and 5 profiles

10 in time to reduce the noise level and hence simplify the automatic evaluation by FLICA.

The output of the cloud detection scheme has been visually inspected for individual days and was found not to show any apparent artifacts. There is a trade-off between detecting cloud structures small enough and avoiding misclassifying noise as a cloud, especially for daytime measurements. The combination of the criteria below represents a rather conservative approach, which might result in missing some particularly small/thin clouds. The conservative approach ensures that no noise is misclassify

sified as a cirrus cloud. An example of the resulting cloud detection can be seen in Fig. 1.

The FLICA algorithm contains the following steps:

20

15

I **Cloud top detection.** The cloud top is needed as an upper boundary for the subsequent lidar retrieval. Our cloud top detection averages individual lidar profiles so that the resolution of one pixel is 5 min in time and 30 m in altitude. Areas of 3×3 pixels are examined, with the pixel to be checked for cloudiness in the center. At least 8 of the 9 examined pixels have to have a volume depolarization larger than 0.007(0.006) for day(night)-time measurements. At least 8 of 9 pixels also have to have larger co- and cross-polarized raw signals than empirical thresholds.

- II Setting the far-end boundary condition for the lidar retrieval. The mean of the co-polarized signal at altitudes from detected cloud top to 500 m above cloud top is computed for each profile individually and used as far-end boundary condition for the lidar retrieval as described in Klett (1981). At this boundary, we assume a BSR of 1. This assumption introduces no error if the aerosol density above the cloud is the same as the one of the interstitial aerosol. If these densities are different we estimate from our in-situ observations (http://www.iac.ethz.ch/groups/peter/research/Balloon_soundings/ COBALD sensor) that the error introduced is of the order of 1 2 %.
- 30 III Lidar retrieval. The lidar retrieval is performed as described in Chapter 5 in Kovalev and Eichinger (2004) to solve the lidar equation (Eq. 1) and calculate the extinction coefficients and BSR of the cirrus cloud. The retrieval is performed for a set of lidar ratios between 5 and 40 sr, in steps of 5 sr. The best choice was determined such that BSR is closest to 1

below the cirrus cloud. The BSR is corrected during the retrieval such that the mean BSR in the range 500 m above the cloud top is equal to 1.

- IV **Cirrus cloud detection.** The cloud detection scheme is based on the retrieved BSR and volume depolarization as follows: Resolution of one pixel: 5 min in time, 30 m in altitude.
- 5 Areas of 3×3 pixels are examined, with the pixel to be checked for cloudiness in the center. At least 8 of the 9 examined pixels have to have a volume depolarization larger than 0.007(0.006) and a BSR larger than 1.08(1.03) for day(night)time measurements.

Temperature has to be lower than -38° C (e.g., Pruppacher and Klett, 1997; Koop et al., 2000; Krämer et al., 2016) to ensure pure ice clouds and avoid detecting mixed-phase clouds (this is checked using COSMO-2 or COSMO-7 analysis data).

10

20

The detection is applied to each pixel at each time independently.

Clouds extending less than 150 m in altitude during daytime conditions are not further taken into account (as noise-limiting measure), whereas nighttime clouds are allowed to be as thin as 3×30 m = 90 m.

Cloud pixels separated vertically by less than 150 m are merged into one cloud layer.

- 15 The detected cloud top and cloud base, h_{top} and h_{base} , are stored.
 - VI **Multiple scattering correction.** The single-scattering extinction coefficients are derived from the apparent, multiplescattering affected extinction coefficients as described in Section 2.1. We use the multiple scattering model of Hogan (2008) as described in Wandinger (1998) and Seifert et al. (2007).
 - VII **Optical depth.** The optical depth τ of the detected cirrus cloud is calculated by integrating over the retrieved extinction profiles.

$$\tau = \int_{h_{\text{base}}}^{h_{\text{top}}} \alpha_p(r) dr \tag{7}$$

VIII **Radiative effect.** The optical depth τ combined with temperatures from COSMO-2 or COSMO-7 are used to calculate the radiative effect of the cirrus cloud by means of the model of Corti and Peter (2009).

3 Lidar cirrus climatology

25 3.1 Measurement sites

Here we present the retrieved lidar cirrus climatology. First, a description of the different measurement sites shown in Fig. 2 is provided. Subsequently, we present the climatology of the cirrus cloud properties. The section ends with a comparison of our data with previous mid-latitude climatology studies.





Figure 2. Location of the measurements sites Jülich(JUL), Jungfraujoch(JFJ) and Zürich(ZRH). Color-coded: Topography in COSMO-7. Black lines: National borders. Source: National Geophysical Data Center, 1993. 5-minute Gridded Global Relief Data (ETOPO5). National Geophysical Data Center, NOAA. doi:10.7289/V5D798BF [13.08.2015]

3.1.1 Jungfraujoch

Jungfraujoch is the highest measurement site used in this study. It is located in the Swiss Alps (46.55° N, 7.99° E) at 3580 m a.s.l. at the top of the Aletsch glacier. Due to its high elevation, the research station is frequently situated in the free troposphere (Zieger et al., 2012), which is a great advantage for lidar measurements. The sphinx observatory, where our measurements

5 took place, is one of the Global Atmospheric Watch (GAW) research stations. Therefore, long time series of meteorological measurement data are available for this site. Due to its high location and cold climate, the site poses challenges for instruments being able to run continuously. The Leosphere ALS 450 used in this study was built into a ventilated, temperature-controlled and regulated containment.

3.1.2 Zürich

- 10 Zürich (47.37° N, 8.55° E), the largest city in Switzerland, is situated in the northern part of Switzerland at 408 m a.s.l., within the Swiss Plateau. The Swiss Plateau is surrounded by the Alps and Jura mountains, which create a basin through which air masses originating from the Atlantic Ocean are funneled. Therefore, the predominant wind direction in Zürich is from the southwest. Although the Swiss plateau is a large basin, it is still hilly. Lake Zürich is a basin itself within the Swiss plateau, and the city of Zürich is situated on the lake's northern shore. Lidar measurements were taken from the roof of ETH's Institute for
- 15 Atmospheric and Climate Science (IAC), which is 500 m above sea level and located in the middle of Zürich. Aerosol particles in and around Zürich arise from industry, transportation and housing, and the large airport nearby. In contrast to Jungfraujoch, such additional aerosol sources cause low level extinction of the emitted laser pulse.

3.1.3 Jülich

The Research Center Jülich is located 91 m a.s.l. in the western part of middle Germany (50.91° N, 6.40° E) between the larger cities Aachen and Köln in North Rhine-Westphalia. Due to its low elevation and location close to the Netherlands, the weather fronts arrive more or less directly from the Atlantic Ocean without moderation by orography. The terrain around Jülich is relatively flat. The Research Center itself is located in a rural area and therefore the lidar measurements might be less influenced by boundary layer aerosol than the Zürich measurements, despite nearby brown coal industry activity.

3.2 Cirrus Climatology

- 25 Following Section 2.2 we present the climatological evaluation of more than 13'000 hours of lidar measurements within the period 2010-2014 from the three mid-latitude measurement sites. The main information on the measurement statistics for the three sites is compiled in Table 2. More measurements are available from Jungfraujoch and Zürich than from Jülich. For Jungfraujoch, most of the data was retrieved in the spring, whereas during the summer only very limited data is available. In Zürich, on the other hand, a large number of the measurements took place during the summer, while the other seasons show
- 30 similar coverage. The Jülich lidar was running predominantly during spring and summer, while the autumn and winter data is sparse. The amount of data has to be considered when judging seasonal variability. The retrieved cirrus properties listed

in Table 2, indicate a temporal cirrus cloud coverage between 9 and 15 % for all stations, agreeing well with the CALIPSO measurements discussed by Sassen et al. (2008) and being slightly smaller than the 18-19% measured during the ECLIPS campaign by Winker and Vaughan (1994).

Table 2. Properties of the cirrus clouds detected between 2010-2014.

	JFJ	Zürich	Jülich
General Properties			
Hours of measurements ⁽¹⁾	5170	4678	3274
Number of cirrus detected ⁽²⁾	10295	6021	7184
Cirrus cloud coverage in $\%^{(3)}$	14	9	15
Low cloud coverage in $\%^{(4)}$	15	8	26
Clear sky in $\%^{(5)}$	71	83	59
Fraction of measurement time by season in $\%$			
DJF	24	17	18
MAM	40	15	39
JJA	14	48	31
SON	22	20	12
Cloud occurrence frequencies in categories			
according to Sassen and Cho (1992) (expressed			
as fraction of "number of cirrus detected")			
Subvisible cirrus ($\tau < 0.03$) in %	43	35	32
Thin cirrus (0.03 $< \tau <$ 0.3) in %	46	52	51
Opaque cirrus (0.3 < τ) in %	11	13	17
Mean $ au^{(6)}$	$0.12\substack{+.02 \\06}$	$0.14\substack{+.02 \\08}$	$0.17\substack{+.02 \\08}$

⁽¹⁾ Refers to the number of hours lidar measurements with the ALS 450. Not included are times times when the ceilometer detected low-level clouds closer than 1.5 km.

⁽²⁾ According to the specifications of the FLICA algorithm, see Subsection 2.2.

⁽³⁾ This compares reasonably well with 11 % zonal average by Chen et al. (2000).

⁽⁴⁾ Refers only to clouds at least 1 km above the lidar.

⁽⁵⁾ As observed by the ALS 450.

5

⁽⁶⁾ Uncertainties as described in the last paragraph of Subsection 2.1. Mean values of τ compare reasonably well with monthly mean τ values of 10-20% 0.1-0.2 from ISCCP (Soden and Donner (1994); also http://www.gfdl.noaa.gov/ice-clouds-in-the-skyhi-general-circulation-model).

The seasonal dependence of the observed cirrus coverage is displayed in Fig. 3. The most striking feature is the difference between the wintertime measurements in Zürich and Jungfraujoch showing a cirrus coverage of around 12%, while in Jülich this is about 33%. This is in qualitative agreement with geographical maps of high cloud amount (cloud pressure smaller



Figure 3. Seasonal cycle of cirrus cloud coverage for the three measurement sites. Dashed lines: annual means.

than 440 hPa) for January observed by the TIROS-N Operational Vertical Sounder, TOVS, averaged over 8 years, 1987-1995 (http://ara.abct.lmd.polytechnique .fr/index.php?page=clouds). For January, these data suggest decreasing amounts of high clouds when the air passes from the North Sea towards the Alps. Also, a time series of 40 years of measurements of turbidity in Jülich confirm the high cloud coverage during wintertime (personal communication A. Knaps, Forschungszentrum Jülich). However, there are large uncertainties in this part of our climatology, as the number of hours of measurements with the Leosphere ALS 450 available for the Jülich winter is small (Table 2), the specific winter might have had a particularly high cirrus cloud coverage, and the applied manual operation of the lidar might add bias. Another remarkable feature is that autumn maximum in cirrus coverage observed above Jungfraujoch. Also this feature is in qualitative agreement with the seasonal cycle suggested by the TOVS data set, but again interannual variability might be important but cannot be derived from our measure-

10 ments.

5

The first property of interest is the distribution of the optical depths of the detected cirrus clouds, which we classify according to Sassen and Cho (1992) (cf. Table 2): clouds with an optical depth $\tau < 0.03$ are not visible to the naked eye, and hence termed subvisible. Cirrus clouds with an optical depth τ in the range $0.03 \le \tau < 0.3$ are termed thin, while clouds with $\tau \ge 0.3$ are referred to as opaque. The upper limit of detection for our lidars is $\tau \approx 3$, as for larger optical depths the light is almost fully extinguished within the cloud. Under these conditions no molecular signal from above the cloud can be detected (Immler and Schrems, 2002), as would be required for an inversion. Therefore, we are not able to specify the optical thickness

5 of the thickest cirrus clouds. Chen et al. (2000) classified clouds with tops above 440 hPa (≈ 6500 m) and optical depths larger than 3.6 as cirrostratus. These cirrus clouds may have a negative cloud radiative effect CRF_{NET}, but cannot be considered here because of the detection limits of our lidar instrument.

Figure 4b shows the optical depth of the retrieved cirrus clouds for different seasons. The grey dashed lines indicate the

- 10 categories defined by Sassen and Cho (1992). The optical depth averaged over the whole data sets for each measurement site is displayed in Figure 4a. These τ values agree well with the ECLIPS-campaign (Pal et al., 1995), where most detected cirrus clouds had optical depths smaller than 0.1. A Wilcoxon Rank sum test reveals that the optical depth distributions of the different sites are significantly different from each other. The occurrence frequency of subvisible cirrus clouds is larger for Jungfraujoch than for the other two sites. Two reasons may be responsible for the observed differences. First, Jungfraujoch
- 15 is located in the central Alps, where orography-driven lifting of air masses leads frequently to mountain-wave (lenticularis) cirrus. These clouds are thicker than large-scale cirrus clouds, but thinner than the cirrus formed as outflow of anvils or in warm conveyor belts. The second reason is the enhanced detectability of optically thin clouds at Jungfraujoch as a result of improved signal-to-noise ratios (SNR, see Eq 6). The alpine site is located at an altitude of 3500 m asl, 3000 m above Zürich and 3400 m above Jülich. According to Eq. 1 the received signal depends on the inverse of the squared range between lidar and target. In addition, the Jungfraujoch is frequently above the boundary layer. Therefore, measurements from Jungfraujoch avoid strong
- beam extinction due to boundary layer aerosols.

Figure 5 provides vertical profiles of the cloud-mean signal-to-noise ratio (SNR) of the three stations, where the noise is obtained from Eq. 6. From the profiles it can be seen that the SNR of Jungfraujoch extends to greater heights by about 3 km.
This suggests that the increased detection rate of thin and subvisible cirrus clouds is a result of the increased signal-to-noise ratio. Furthermore, the SNR at Jungfraujoch increases at heights above 13 km a.s.l. This suggests that the morphology of the clouds at these heights differs from the morphology of the highest clouds observed at Jülich and Zürich. The backscattering efficiency appears to be enhanced in these clouds, possibly because a large amount of small crystals formed in the observed cirrus clouds, when many ice crystals nucleated in the high supersaturations in rapid uplifts as they occur in lee waves above mountainous terrain (Lin et al., 1998a, b; Kärcher, 2003).

The number of detected subvisible cirrus as function of optical depth and cloud top altitude is depicted in Fig. 6. As expected, Jungfraujoch displays a larger fraction of subvisible cirrus as well as higher cirrus cloud tops. Therefore, we have evidence for both:



Figure 4. Optical depths of the three measurement sites. (a): Means across whole data set. (b): Seasonal cycle of optical depth. Horizontal line in box: median. Boxes: the upper and lower quartile. Whisker: extremes. Gray horizontal lines: Cirrus categories by Sassen and Cho (1992).



Figure 5. Vertical distribution of the signal-to-noise ratio of the detected cirrus clouds for Jungfraujoch, Jülich, and Zürich. The center line of each box plot represents the median. The left and right limits of the box plots mark the 25% and 75% percentiles, respectively.

- (a) the advantage of location of the higher Jungfraujoch as evidenced by the ability to measure thinner subvisible clouds (by about a factor 5);
- (b) the special conditions above Jungfraujoch caused by orographic forcing, which affects the morphology of the cirrus as evidenced by the enhanced SNR at high altitudes.
- 5 It is interesting to see that at Jungfraujoch the lower detection limit in optical depth of a few times 10^{-5} is approached in a few cases. However, by far the most subvisible cirrus stay clearly above $\tau = 10^{-4}$ proving that physical mechanisms prevent clouds so thin to survive for appreciable times. Nucleation is one such mechanism. In case these clouds nucleate homogeneously, this is likely to happen in nucleation bursts, which will provide the newly formed clouds immediately a minimum optical depth. The same is true for heterogeneous nucleation, if the nucleation barrier and the number of nuclei are at all significant. One
- 10 mechanism that might lead to extremely low ice crystal number densities is the formation of fall-streaks and subsequent dispersion of the particles. The rare occurrence of clouds with $\tau < 5 \times 10^{-4}$ suggests that such mechanisms do not often lead to the formation of so such thin clouds. The height distribution of the detected cirrus clouds in Fig. 6 agrees well with the cirrus clouds measured during the ECLIPS campaign (Winker and Vaughan, 1994).
- 15 To ensure that the highest cirrus clouds observed above Jungfraujoch were not volcanic particles, we have examined satellite measurements and found no indication for volcanic influences. The effect of the high altitude of Jungfraujoch can also be seen in the cloud tops at the different measurement sites (see Fig. 7a). The cloud tops are higher above Jungfraujoch than above



Figure 6. Scatter plots of cloud optical depths and cloud top altitudes for the cirrus detected above Jungfraujoch (JFJ), Zürich (ZRH) and Jülich (JUL). The red lines provide an indication of the range of data accessible by the lidar measurements: $AOD_{min} = 4 \times 10^{-5}$, $AOD_{max} = 2.6$, and $Alt_{min} = 5.8$ km. The lower edge of the accessible altitude is determined from T < -38°C. Thicker clouds are more likely to extend into lower, warmer levels and therefore are more likely to be excluded from the analysis.

Zürich and Jülich. The retrieved cloud tops agree well with the observations by Sassen and Campbell (2001) in Salt Lake City (40° N, 12° W, 1520 m a.s.l.) as well as by Hoareau et al. (2013) in Haute Provence (44° N, 6° E, 679 m a.s.l.). The data from Salt Lake City and from Haute Provence were evaluated using evaluation schemes differing from FLICA and differing amongst themselves, which may influence the results. However, the cloud top altitudes are very similar for the five mid-latitude

5 stations. As Salt Lake City is located further south than the other sites, the slightly higher cloud tops may be a result of a higher tropopause being present over Salt Lake City compared to the other sites. Similarly, we see in Fig. 7b that the tropopause over Jülich, which is located further north than Zürich and Jungfraujoch, generally is lower. Between Zürich and Jungfraujoch, the tropopause reaches similar altitudes with a larger spread over Jungfraujoch (possibly due to the Alpine heat low affecting the Jungfraujoch frequently during summer time).

10 4 Cirrus Radiative Forcing (CRF)

4.1 Method of Calculation

To quantify the net radiative effect, CRF_{NET} , for the cirrus clouds observed here we use the radiation model of Corti and Peter (2009), which is a simplified model based on the more sophisticated Fu-Liou radiative model (Fu and Liou, 1992, 1993). The accuracy of Corti and Peter (2009) is better than 20 % in comparison with the Fu-Liou model. The cloud radiative forcing due

- 15 to shortwave radiation, CRF_{SW} , is dependent on the surface albedo, the solar zenith angle as well as the cirrus cloud optical depth τ (see Eq. 13 of Corti and Peter (2009) for details). The longwave cloud radiative forcing, CRF_{LW} , is mainly determined by the temperature difference between the Earth's surface and the cirrus cloud top temperature and by the cloud optical depth τ (see Eq. 6 of Corti and Peter (2009) for details). The CRF_{LW} further correlates well with the brightness temperature of the atmosphere, which is related to the outgoing longwave radiation at top of atmosphere (Schumann et al., 2012a). This correlation
- 20 has not been considered in the model of Corti and Peter (2009). The net cloud radiative forcing, CRF_{NET} , is calculated as a superposition of these two effects (i.e. $CRF_{NET}=CRF_{SW}+CRF_{LW}$). The following parameters are needed as input for the calculation of the radiative effect:
 - Solar constant S
 - Solar zenith angle Z
- 25 The surface albedo α
 - The cloud optical depth τ
 - The surface temperature $T_{\rm srf}$
 - The cloud top temperature T_{cld}

The values of the solar constant S, multiplied by the fraction of the day that the sun is above the horizon, and the mean solar 30 zenith angle Z are set to 684 Wm⁻² and 60° (daily mean conditions with zero incoming flux at nighttime), respectively, as



Figure 7. (a) Cloud tops (in 500-m steps) at the three sites of this study as well as in Salt Lake City by Sassen and Campbell (2001) and Haute Provence by Hoareau et al. (2013). (b) Tropopause derived from COSMO regional weather forecast model analyses (2.2 km horizontal resolution for JFJ and ZRH, 6.6 km resolution for JUL).

Table 3. Cirrus radiative forcing at the Top of Atmosphere in Wm^{-2} for Jungfraujoch, Zürich and Jülich, as compared to 50° N zonally averaged and globally averaged values provided by Chen et al. (2000). Small numbers in CRF-values indicate uncertainty ranges according to the last paragraph of Subsection 2.1. Small numbers in global cloud coverage indicates variability in zonal averages.

	JFJ	Zürich	Jülich	50° N	global
Cirrus coverage in %	14	9	15	11	13^{+7}_{-8}
Overcast CRF_{NET}	$6.2\substack{+0.7 \\ -3.0}$	$10.6^{+1.5}_{-5.3}$	$11.0^{+1.4}_{-4.9}$	2.0	5.4
CRF _{LW}	$7.2^{+1.0}_{-3.6}$	$12.3^{+1.8}_{-6.1}$	$13.3^{+1.6}_{-6.0}$	20.1	30.7
CRF _{SW}	$-1.0^{+0.5}_{-0.3}$	$-1.7^{+0.8}_{-0.3}$	$-2.4^{+1.1}_{-0.2}$	-18.1	-25.3
All sky CRF_{NET}	$0.9\substack{+0.1 \\ -0.4}$	$1.0\substack{+0.1 \\ -0.5}$	$1.6\substack{+0.2 \\ -0.7}$	0.5	1.3
CRF_{LW}	$1.0\substack{+0.1 \\ -0.5}$	$1.1\substack{+0.2 \\ -0.6}$	$2.0\substack{+0.2 \\ -0.9}$	3.0	5.5
CRF _{sw}	$-0.1^{+0.1}_{-0.0}$	$-0.2^{+0.1}_{-0.0}$	$-0.3^{+0.2}_{-0.0}$	-2.5	-4.2

suggested by the online version of the radiation model (Corti and Peter, 2009b). This results in an incident solar flux $I = 684 \times 0.5 = 342 \text{ Wm}^{-2}$. The amplitude of the solar background noise in the lidar signal profiles is used to distinguish between day and night time. We use an albedo of 0.3 (corresponding to the global average value). The cloud optical depth τ is automatically calculated in the FLICA for 5-min profiles as described in Section 2.2. The temperatures needed for the radiation model (T_{srf} and T_{cld}) are extracted from the COSMO-2 (for Jungfraujoch and Zürich) and COSMO-7 (for Jülich) model. The radiation model of Corti and Peter (2009) is well suitable to be used with lidar data, as the model does not require further information, such as ice crystal sizes or shapes, which the lidar measurements could not provide.

4.2 Comparison of CRFs with previous studies

5

We compare our computational results to satellite data, which have been averaged zonally at 50°N or globally and combined 10 with a radiative transfer model (Chen et al., 2000). The results of this comparison are listed in Table 3 together with maximum possible uncertainty ranges (see last paragraph of Subsection 2.1). The "overcast values" (i.e. taking only cloudy values into account) consider the radiative effect under conditions with cirrus clouds, while the "all sky values" include also conditions without cirrus by considering the cirrus occurrence frequency. While the cirrus cloud coverage at 50°N from the satellite-based climatology ISCCP (Chen et al., 2000) is similar to our observations, the ISCCP category of cirrus clouds show much larger

15 eloud radiative forcings in the short- and longwave, CRF_{SW} and CRF_{LW} (by more than one order of magnitude) 1.5-3 times larger radiative forcing in the longwave, CRF_{LW} , and one order of magnitude larger radiative forcing in the shortwave, CRF_{SW} .

This The difference in the CRF_{SW} can only be explained in terms of a much larger optical depth τ of the clouds observed by the satellites. The CRF_{SW} is mostly linear dependent on the optical depth tau, whereas the CRF_{LW} depends linear on the temperature difference between ground and cloud as well as tau (see Corti and Peter, 2009). Clouds with an increased optical depth are mostly found at lower altitudes where the air is more humid. Thus the temperature difference is smaller which results in

5 a smaller increase of the the CRF_{LW} compared to the CRF_{SW} . Thus, warmer clouds with increased optical depth have larger negative CRF_{NET} . This explains the difference between the CRF_{SW} and CRF_{LW} in Chen et al. (2000), where more clouds with larger optical depth are included.

The reason for this at first sight surprising differences are the different definitions of "cirrus". First, FLICA detects only

- 10 clouds with lower cloud edge colder than -38°C, which is typically above 7-8 km. Chen et al. (2000) instead used a pressure threshold of 440 hPa to separate clouds, which corresponds to an altitude of 6.3 km (standard atmosphere). The clouds in the range 6.3-7.5 km are missing in our study. Especially in this altitude range thick cirrus stratus (still with $\tau < 3.6$) Cirrostratus clouds with $\tau < 3.6$ occur particularly in this altitude range. Second, although our criteria allow for thick clouds up to $\tau = 3.6$ at altitudes clearly above lower edge, they cut clouds once their lower edge gets warmer than -38°C, which is more likely for
- 15 thicker clouds (see rounded edge in the red boxes in Figure 6). Third, while we count vertically distinct cirrus layers as separate clouds, the geostationary ISCCP weather satellites add the signal of vertically staggered layers, which increases τ . Furthermore, it should be noted that satellite data reveal discrepancies amongst themselves (ISCCP, MISR, MODIS) with differences of 20-30% in coverage of cirrus with $\tau < 3.6$ (Marchand et al., 2010). Finally, the distribution of thicker cirrus with $\tau > 0.3$ is zonally inhomogeneous, with clouds preferentially occurring at the continental east coasts.

20

In our study, we want to address only cirrus clouds and not mixed phase clouds. Therefore, we have chosen a conservative limit towards lower, thicker clouds. Also, a temperature-based selection criterion is a better for separating different cloud types than a pressure-based criterion, because temperature is the main microphysical parameter for cloud formation. As ISCCP is based on the analysis of weather satellite images, clouds still must have optical depths $\tau \ge 0.2$ in order to be reliably detected by such satellites (Rossow and Schiffer, 1999). Large uncertainties can also be traced to different approaches to partly cloudy

25 by such satellites (Rossow and Schiffer, 1999). Large uncertainties can also be traced to different approaches to partly cloudy pixels, which are 30 km x 30 km for ISCCP and are treated as homogeneous, i.e. either cloud free or filled with a thinned homogeneous cloud (Pincus et al., 2012)

The overcast and all sky CRF_{NET} are significantly higher in Jülich than at Jungfraujoch, which is also clearly reflected in
overcast and all sky optical depths found in the ISCCP data (Soden and Donner, 1994). This may be related to the frequent low-pressure systems and fronts rolling in from the northwest across the North Sea. The related cirrus clouds are weaken with distance from the coast.

The effect of the optically thicker clouds above Jülich compared to the Jungfraujoch is also evident in Fig. 8. The magenta lines indicate positive (i.e. warming) cirrus cloud radiative forcing (in Wm^{-2}) as a function of altitude and optical depth cal-

culated by the model of Corti and Peter (2009) with mean temperature profiles from COSMO-2 (Jungfraujoch and Zürich) and COSMO-7 (Jülich) during the time period of our measurements. The blue isolines indicate negative (i.e. cooling) cirrus radiative forcing. A zero net effect, $CRF_{NET}=0$ is indicated by a cyan line. Color-coded is the occurrence frequency of the cirrus clouds measured at the different sites. The occurrence frequency is categorized by 40 logarithmically spaced bins in optical

- 5 depth between 10^{-4} and 10 and 500-m bins in cloud top altitude. From Fig. 8 we clearly see that the cirrus clouds observed in this study have all a positive (warming) net radiative effect CRF_{NET}. It is important to note, that with the FLICA algorithm we do not find cirrostratus or cumulonimbus outflow clouds, i.e. no clouds with $\tau > 3.6$. Of course, such clouds do exist also above our measurement sites. However, such clouds always have lower edges warmer than -38°C and thus are not considered.
- 10 The pattern of cirrus cloud occurrence is quite similar above Jungfraujoch and Zürich, although the Jungfraujoch cirrus clouds show a slightly broader distribution in optical depths. Most cirrus layers are present at 11 km a.s.l. with optical depths between 0.01(0.04) and 0.2(0.4) above Jungfraujoch (in Zürich). Cirrus clouds above Jülich are frequent at altitudes between 8 and 12 km a.s.l. with optical depths ranging from 0.02 to 0.7. This wider distribution of high occurrence frequencies in altitude is likely related to the high frequency of frontal systems crossing Jülich. The lower CRF_{NET} above Jungfraujoch is visible in the shift

15 towards thinner clouds at Jungfraujoch as compared to the other two measurement sites. Due to the lower SNR over Zürich and Jülich at cirrus altitude, these two sites underestimate the amount of subvisible cirrus clouds as compared to Jungfraujoch.

Beside Chen et al. (2000) other studies indicate also a general net warming effect of cirrus clouds in the mid-latitudes. Oreopoulos and Rossow (2011) investigated the overall cloud radiative forcing based on a 24 year long data set from the Inter-

- 20 national Satellite Cloud Climatology Project (ISCCP). For cases with frequent occurrence of high clouds, a positive net cloud radiative forcing (warming) was obtained whereas it was negative (cooling) for cases with frequent occurrence of low-level clouds. This confirms our results where cirrus clouds create a positive net radiative forcing. A case study of Katagiri et al. (2010) found a cirrus cloud radiative forcing of 13.2 Wm⁻² at the Fukue observatory (32°N, Japan) by a combination of MODIS satellite and ground-based observations. This value is similar to what we found for Zürich and Jülich and obviously
- 25 larger than the respective value reported by Chen et al. (2000). Another study of Min et al. (2010) found a radiative forcing of cirrus of 36.5 Wm⁻² over China using also CALIOP and MODIS satellite data. The authors ascribe the high value to cirrus observations above the Tibertian plateau where very thick cirrus clouds with a mean optical depth of 1 are observed frequently. For the other parts of China lower values of 20 Wm⁻² are found. The radiative forcing of the lateral boundary of cirrus clouds observed with CALIOP is investigated by Li et al. (2014). The radiative effect of observed cirrus cloud edges is discussed.
- 30 In the transition region of large cirrus, defined as their optically thin rim ($\tau < 0.3$), which is often missed by satellite passive optical sensors such as MODIS, the CRFLW found to be still substantial ($\sim 10 \text{ Wm}^{-2}$). The lateral boundary with optical depths less than 0.3 is found to have still a radiative forcing of 10 Wm⁻². This value is similar to our mean overcast radiative forcing of Zürich and Jülich and demonstrates also the sensitivity of cirrus cloud inhomogeneity on cloud forcing as also found by Gu and Liou (2006). However, all these studies investigate single cases or different regions compared to our study. There is
- 35 no study which investigates the cirrus radiative forcing over Europe.



Figure 8. Cloud Radiative Forcing (CRF) in Wm^{-2} for the different sites. Magenta/cyan/blue isolines: positive/zero/negative values in Wm^{-2} from the CRF model (Corti and Peter, 2009). Color coding: Occurrence frequency of cirrus clouds as function of optical depth and cloud top altitude. First row: CRF_{LW}. Second row: CRF_{SW}. Third row: CRF_{NET}.

In a study of Dupont et al. (2010) cirrus cloud observations over 2 years from the CALIPSO satellite lidar CALIOP are compared to four ground based lidar stations (two sites in the US and two in France) for their consistency of macrophysical and optical properties. They found larger discrepancies in the frequency distributions of cloud base, top and thickness. They

- 5 point out that the significant part of the deviations can be attributed to different sampling (seasonal, irregular sampling of ground based stations, opaque low level clouds). However, they found that for high cirrus clouds the optical depth distribution $\tau > 0.1$ from ground stations and CALIOP is consistent within 10% using the same retrieval method. This shows that our optical depth distribution of all three stations is most likely not or only less affected by sampling issues of ground base lidar compared to satellite measurements. Further, we more closely examined the CRF_{NET} categorized by their optical thickness τ
- 10 as in Sassen and Cho (1992).

4.3 Influence of subvisible cirrus on the net radiative forcing by cirrus clouds

Subvisible cirrus clouds generally are not considered in numerical weather prediction models as their optical depths are considered to be too small. The overcast net radiative effect CRF_{NET} , divided into the categories defined by Sassen and Cho (1992), is shown in Fig. 9. We see that the subvisible cirrus clouds indeed have an effect on the net cirrus cloud radiative forcing CRF_{NET} .

- 15 On average they contribute about 4 % of the total CRF_{NET} of cirrus clouds at Jungfraujoch and in Zürich, and 3 % in Jülich. The maximal effect of 6 % is reached in Zürich during spring. As seen in Fig. 9, the thin and opaque cirrus clouds are the main contributors to CRF_{NET} of cirrus clouds above each of the three stations, both by roughly equal shares (with a small dominance of opaque clouds).
- Jungfraujoch displays the lowest CRF_{NET}-values throughout the whole year. This pattern is also seen in the optical depths shown in Fig. 4a. Generally, thinner clouds are detected above Jungfraujoch than at the other two sites. This influences the CRF_{NET}: as more subvisible cirrus are observed at Jungfraujoch (cf. Table 2) and their contribution to CRF_{NET} is smaller than the contribution by thin and opaque cirrus, this leads to a smaller CRF_{NET} on Jungfraujoch. Summing up the percentages listed in Table 2, Jungfraujoch shows a fraction of 57 % thin and opaque cirrus while Zürich and Jülich both occurrence frequencies
 of 65 and 68% thin and opaque clouds, respectively. These different sums result in the CRFs listed in Table 3 (taking note of

the fact that the thresholds for the cirrus cloud categories (Sassen and Cho, 1992) are on a logarithmic scale).

5 Conclusions

We have presented a cirrus climatology based on 13'000 hours of lidar measurements at the three different mid-latitude sites Jungfraujoch, Zürich and Jülich from 2010-2014. This extensive data set was evaluated using the newly developed FLICA

30 algorithm, which combines a pixel-based cloud detection scheme with a classic lidar retrieval. With FLICA, the lidar data have been automatically evaluated. The retrieved backscatter coefficients are converted into extinction coefficients, which are corrected for multiple scattering to establish single scattering extinction and then converted into optical depths.



Figure 9. Cirrus radiative forcing under cloudy conditions, CRF_{NET} (overcast), for the different seasons on Jungfraujoch (blue), in Zürich (pink) and Jülich (green). Light shading: subvisible cirrus ($\tau < 0.03$). Medium shading: thin cirrus ($0.03 < \tau < 0.3$). Dark shading: opaque cirrus ($0.3 < \tau$).

We find mean optical depths of 0.12 for the cirrus measured over Jungfraujoch and of 0.14 and 0.17, respectively, for Zürich and Jülich. While the cirrus coverage over Jungfraujoch and Jülich are almost equal, the amount of subvisible clouds detected over Jungfraujoch is significantly larger (cf. Table 2). Due to its unique location at 3580 m a.s.l., Jungfraujoch is an excellent site

- 5 to measure subvisible cirrus clouds with a much improved SNR at cirrus altitude in comparison with the lower-lying stations. The mean cloud tops detected were located at 10.7 km in Zürich and on Jungfraujoch and at 10.3 km in Jülich, consistent with previous studies of Sassen and Campbell (2001) and Hoareau et al. (2013). Further, we have measured a temporal cirrus cloud coverage of 9-15 % with a mean value of 13 %. This is consistent with the evaluation of the global CALIPSO-measurements of Sassen et al. (2008).
- 10

Owing to the central location of the three measurement sites in Europe, a significant fraction of the thin cirrus observed within the present study might actually have originated from contrails. Clear indications for the occurrence of contrails were found on at least one day, given the optically and geometrically very thin cirrus. Liou et al. (1990) noted an increase of thin cirrus with increases in jet traffic. Our measurement sites are located in a region, where line-shaped contrails are ubiquitous

- 15 (Mannstein et al., 1999; Meyer et al., 2002) as many flight routes cross this area. The observed optical depths are consistent with optical depths of contrail cirrus (Immler et al., 2008; Iwabuchi et al., 2012; Vázquez-Navarro et al., 2015). Furthermore, the cirrus cloud cover determined in the present study is consistent with contrail cirrus calculations by Schumann et al. (2015). Therefore, it is likely that contrails contributed a fraction of the observed cirrus. The determination of the actual contribution of contrails to the cirrus cloud dataset is, however, not subject of this study, considering that the applied data analysis algorithm
 20 ELICA connect distinguish natural and contrail cirrus.
- 20 FLICA cannot distinguish natural and contrail cirrus.

The evaluated cirrus cloud properties are used together with the radiation model of Corti and Peter (2009) to estimate the cloud radiative forcing of the cirrus clouds. The optical depth as well as the cloud top temperature are the most important quantities determining the CRF_{NET} , and this dependence has been displayed in Fig. 8. Our results clearly confirm the warming

- effect of mid-latitude cirrus clouds with optical depths below 3, corroborating previous studies. Using the radiation model of Corti and Peter (2009), we find a net effect of 0.9 Wm⁻² for Jungfraujoch and 1.0/1.6 Wm⁻² for Zürich/Jülich. These values are larger by factors of 2-3 than the 50°N zonally averaged CRF_{NET} derived by Chen et al. (2000) from satellite measurements in combination with a radiative transfer model. Even stronger deviations–but with opposite sign–are found for CRF_{SW} and CRF_{LW}, where the zonally averaged data are higher than our CRF by up to one order of magnitude. This is due to different
- 30 cloud definitions used by Chen et al. (2000) and us, and to the fact that the satellite-based zonal average includes regions with more pronounced thick cirrus (e.g. the continental east coasts).

Besides the radiation model of Corti and Peter (2009) used for this study, other approaches exist that can be used to investigate the effect of other cloud properties besides optical depth on the cirrus radiative forcing. For instance, the radiation model

of Schumann et al. (2012b) could be used to test the influence of various assumptions on particle habits and particle sizes

The actual purpose of this work is the investigation of the thin $(0.03 < \tau < 0.3)$ and subvisible $\tau < 0.03$ cirrus clouds, which remain undetected by passive remote-sensing satellites (requiring typically $\tau > 0.2$) and have so far not yet been sys-

- 5 tematically characterized in a climatological manner. The present study presents more than 13'000 hours of elastic backscatter lidar data, comprising more than 23'000 individual cirrus clouds. Of these clouds about 40% were subvisible, 50% thin, and 10% opaque cirrus. In terms of fraction of cloud coverage, subvisible cirrus were observed during about 6%, thin cirrus during about 7% and opaque cirrus during about 1.5% of the observation time. Seasonal variability in cirrus coverage shows characteristic autumn and spring maxima in agreement with satellite climatologies. Finally, in terms of cloud radiative forcing, all
- 10 clouds discussed here show a positive, i.e. warming, effect. We calculate that subvisible cirrus contribute about 5%, thin cirrus about 45% and opaque cirrus about 50% of the total cirrus radiative forcing. In order to exert a negative forcing, i.e. a cooling effect, clouds need to be either optically much thicker or in altitude much lower, or both, but we excluded these clouds by demanding that the lower edge of the cloud needs to be colder than -38°C (cf. Fig. 8).
- 15 One important difference between the high ice clouds measured at Jungfraujoch compared to Jülich (with Zürich intermediate) is the possibility to measure thinner clouds above Jungfraujoch, which emphasizes the enhanced suitability of the high alpine measurement station to achieve a high signal-to-noise ratio. Reasons for this are that the objects of interest are closer (and the backscattered signals scales with the square of the distance) and that the polluted boundary layer stays often below the Jungfraujoch station. The Jungfraujoch data show that the lower detection limit in optical depth of a few times 10^{-5} is
- approached in a few cases, but by far the most subvisible cirrus stay clearly above $\tau = 10^{-4}$. We argue that this indicates that physical mechanisms prevent clouds to become and stay so thin for appreciable times. After formation, clouds will typically grow quickly and assume higher optical thicknesses. Conversely, evanescent clouds–once having become so thin–will evaporate quickly, not leaving much time for their detection. This leads us to speculate that the Jungfraujoch measurements enable us to explore the very onset of cirrus formation and to possibly learn from the lidar measurements about the relative importance
- 25 of homogeneous and heterogeneous ice nucleation.

Acknowledgements. We are very grateful to Frank Wienhold for providing scripts for lidar evaluation and for input to an early version of the manuscript. We thank Uwe Weers, Marco Vecellio and Edwin Hausammann for technical support with the lidar. We are particularly grateful to Joan and Martin Fischer as well as Maria and Urs Otz for excellent local support at the Jungfraujoch. Thanks also to Albert Ansmann for very helpful scientific input, and to Andrew Huisman, Laura Revell as well as Silke Gegenbauer for proof reading of an early stage of the

³⁰ manuscript. This work has been funded by GAW-CH, the Swiss branch of the Global Atmosphere Watch (GAW) programme, coordinated by the GAW-CH Office at MeteoSwiss in Switzerland.

References

- Achtert, P., Khaplanov, M., Khosrawi, F. and Gumbel, J.: Pure rotational-Raman channels of the Esrange lidar for temperature and particle extinction measurements in the troposphere and lower stratosphere, Atmospheric Measurement Techniques, 6,1,91–98,doi: 10.5194/amt-6-91-2013, 2013.
- 5 Ackermann, J.: The extinction-to-backscatter ratio of tropospheric aerosol: A numerical study, J. Atmos. Oceanic Technol., 15, 1043–1050, 1998.
 - Ansmann, A., Wandinger, U., Riebesell, M., Weitkamp, C. and Michaelis, W.: Independent measurement of extinction and backscatter profiles in cirrus clouds by using a combined Raman elastic-backscatter lidar, Appl. Opt., 31, 7113–7131,1992.

Arshinov, Y. and Bobrovnikov, S.: Use of a Fabry-Perot interferometer to isolate pure rotational Raman spectra of diatomic molecules, Appl.

- Boucher, O., Randal, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S.K., Sherwood, S., Stevens, B. and Zhang, X.Y.: Clouds and Aerosols. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P.M. (eds.)]. Cambridge University
- Press, Cambridge, United Kingdom and New York, NY, USA, pp. 571–658, doi:10.1017/CBO9781107415324.016,2013.
 Bucholtz, A.: Rayleigh-scattering calculations for the terrestrial atmosphere, Appl. Opt., 34, 2765-2773, 1995.
 - Burton, S.P., Ferrare, R.A., Hostetler, C.A., Hair, J.W., Rogers, R.R., Obland, M.D., Butler, C.F., Cook, A.L., Harper, D.B. and Froyd, K.D: Aerosol classification using airborne High Spectral Resolution Lidar measurements – methodology and examples, Atmos. Meas. Tech., 5, 73-98, doi:10.5194/amt-5-73-2012, 2012.
- 20 Chen, T., Rossow, W.B. and Zhang, Y.: Radiative Effects of Cloud-Type Variations, J. Climate, 13, 264–286, 2000.
- Cirisan, A., Luo, B. P., Engel, I., Wienhold, F. G., Krieger, U. K., Weers, U., Romanens, G., Levrat, G., Jeannet, P., Ruffieux, D., Philipona, R., Calpini, B, Spichtinger, P., and Peter, T.: Balloon-borne match measurements of mid-latitude cirrus clouds. Atmos. Chem. Phys., 14, 7341–7365, doi:10.5194/acp-14-7341-2014, 2014.

Cooney, J.: Measurement of Atmospheric Temperature Profiles by Raman Backscatter, J. Appl. Meteor., 11, 108-112, 1972.

- Corti, T. and Peter, T.: A simple model for cloud radiative forcing, Atmos. Chem. Phys., 9, 5751-5857, 2009.
 Corti, T. and Peter, T.: A simple model for cloud radiative forcing. http://iacweb.ethz.ch/php/toa/toa_high.php as per 16.07.2015.
 COSMO: Consortium for Small-scale Modeling. http://www.cosmo-model.org/ as per 16.07.2015.
 - Cziczo, D. J., Froyd, K. D., Hoose, C., Jensen, E. J., Diao, M., Zondlo, M. A., Smith, J. B., Twohy, C. H. and Murphy, D. M.: Clarifying the Dominant Sources and Mechanisms of Cirrus Cloud Formation, Science, Vol. 340, 6138, 1320–1324, doi:10.1126/science.1234145, 2013
- 30 2013.

Cziczo, D. and Froyd, K.D.: Sampling the composition of cirrus ice residuals, Atmos. Res., 142, 15–31, 2014.

Dessler, A. E., and Yang, P.: The distribution of tropical thin cirrus clouds inferred from Terra MODIS data, J. Clim., 16, 1241–1247, doi:10.1175/1520-0442, 2003.

Dionisi, D., Keckhut, P., Liberti, G. L., Cardillo, F. and Congeduti, F.: Midlatitude cirrus classification at Rome Tor Vergata through a

35 multichannel Raman-Mie-Rayleigh lidar, Atmos. Chem. Phys., 13,23, 11853–11868, doi:10.5194/acp-13-11853-2013, 2013.

¹⁰ Opt., 21, 4635–4638, 38, doi:10.1364/AO.38.004635, 1999.

- Dupont, J. C., Haeffelin, M., Morille, Y., Noel, V., Keckhut, P., Winker, D., Comstock, J., Chervet, P., and Roblin, A.: Macrophysical and optical properties of midlatitude cirrus clouds from four ground-based lidars and collocated CALIOP observations, Journal of Geophysical Research-atmospheres, 115, D00H24, doi:10.1029/2009JD011943, 2010.
- Ebert, E. and Curry, J.: A Parameterization of Ice Cloud Optical Properties for Climate Models, J. Geophys. Res., 97, 3831-3836, 1992.
- 5 Fréville, P., Montoux, N., Baray, J.-L., Chauvigné, A., Réveret, F., Hervo, M., Dionisi, D., Payen, G. and Sellegri, K.: LIDAR Developments at Clermont-Ferrand-France for Atmospheric Observation, Sensors (Basel, Switzerland), 15(2), 3041-3069, doi:10.3390/s150203041, 2015.
 - Freudenthaler, V., Esselborn, M., Wiegner, M., Heese, B., Tesche, M., Ansmann, A., Müller, D., Althausen, D., Wirth, M., Fix, A., Ehret, G., Knippertz, P., Toledano, C. Gasteiger, J. Garhammer, M. and Seefeldner, M.: Depolarization ratio profiling at several wavelengths in pure Saharan dust during SAMUM 2006, TellusB, 61, 165-179, 2009.
- Fu, Q. and Liou, K. N.: On the Correlated K-Distribution Method for Radiative-Transfer in Nonhomogeneous Atmospheres, J. Atmos. Sci., 49, 2139–2156, 1992.
 - Fu, Q. and Liou, K. N.: Parameterization of the Radiative Properties of Cirrus Clouds, J. Atmos. Sci., 50, 2008-2025, 1993.
 - Fusina, F., Spichtinger, P. and Lohmann, U.: Impact of ice supersaturated regions and thin cirrus on radiation in the midlatitudes, J. Geophys.
- 15 Res., 112, D24S14, doi:10.1029/2007JD008449, 2007.

10

- Giannakaki, E., Balis, D. S., Amiridis, V., and Kazadzis, S.: Optical and geometrical characteristics of cirrus clouds over a Southern European lidar station, Atmos. Chem. Phys., 7, 5519-5530, doi:10.5194/acp-7-5519-2007, 2007.
 - Goldfarb, L. and Keckhut, P. and Chanin, M.-L. and Hauchecorne, A.: Cirrus climatological results from lidar measurements at OHP (44°N, 6°E), Geophysical Research Letters, 28, 9,1687-1690, doi:10.1029/2000GL012701, 2001.
- 20 Gu, Y. and Liou, K. N.: Cirrus cloud horizontal and vertical inhomogeneity effects in a GCM, Meteorology and Atmospheric Physics, 91, 223–235, doi:10.1007/s00703-004-0099-2, 2006.
 - Heese, B., Flentje, H., Althausen, D., Ansmann, A., and Frey, S.: Ceilometer lidar comparison: backscatter coefficient retrieval and signalto-noise ratio determination, Atmos. Meas. Tech., 3, 1763–1770, doi:10.5194/amt-3-1763-2010, 2010.

Hogan, R. J.: Fast Lidar and Radar Multiple-Scattering Models. Part I: Small-Angle Scattering Using the Photon Variance-Covariance

- 25 Method, Journal of the Atmospheric Sciences, 65, 3621–3635, doi:10.1175/2008JAS2642.1, 2008
 - Hoareau, C., Keckhut, P., Noel, V., Chepfer, H., and Baray, J.-L.: A decadal cirrus clouds climatology from ground-based and spaceborne lidars above the south of France (43.9° N–5.7° E), Atmos. Chem. Phys., 13, 6951-6963, doi:10.5194/acp-13-6951-2013, 2013.
 - Ickes, L., Welti, A., Hoose, C. and Lohmann, U.: Classical nucleation theory of homogeneous freezing of water: thermodynamic and kinetic parameters, Phys. Chem. Chem. Phys., 17, 8, 5514-5537, The Royal Society of Chemistry, doi:10.1039/C4CP04184D, 2015.
- 30 Immler, F. and Schrems, O.: LIDAR measurements of cirrus clouds in the northern and southern midlatitudes during INCA (55°N, 53°S): A comparative study, Geophysical Research Letters, 29, 16, 56-1–56-4,doi:10.1029/2002GL015077, 2002.
 - Immler, F., Treffeisen, R., Engelbart, D., Krüger, K. and Schrems, O.:Cirrus, contrails, and ice supersaturated regions in high pressure systems at northern mid latitudes, Atmos. Chem. Phys., 8, 1689–1699, doi:10.5194/acp-8-1689-2008,2008.
 - Iwabuchi, H., Yang, P., Liou, K. N. and Minnis, P.: Physical and optical properties of persistent contrails: Climatology and interpretation, J.
- **35** Geophys. Res., 117, D06215, doi:10.1029/2011JD017020, 2012.
 - Kärcher, B.: Simulating gas-aerosol-cirrus interactions: Process-oriented microphysical model and applications. Atmos. Chem. Phys, 3, 1645–1664, 2003.

- Katagiri, S., Kikuchi, N., Nakajima, T. Y., Higurashi, A., Shimizu, A., Matsui, I., Hayasaka, T., Sugimoto, N., Takamura, T., and Nakajima, T.: Cirrus Cloud Radiative Forcing Derived from Synergetic Use of MODIS Analyses and Ground-Based Observations, Sola, 6, 25–28, doi:10.2151/sola.2010-007, 2010.
- Kienast-Sjögren, E., Miltenberger, A. K., Luo, B. P. and Peter, T.:Sensitivities of Lagrangian modeling of mid-latitude cirrus clouds to trajectory data quality, Atmos. Chem. Phys. Disc., 15,5, 7535-7584, doi:10.5194/acpd-15-7535-2015, 2015.
- Kim, Y., Kim, S.-W., Kim, M.-H. and Yoon, S.-C.: Geometric and optical properties of cirrus clouds inferred from three-year ground-based lidar and CALIOP measurements over Seoul, Korea, Atmospheric Research, 139, 27-35, 2014.

Klett, J.: Stable analytical inversion solution for processing lidar returns, Appl. Opt., 20, 211–220, 1981.

Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions. Nature, 406, 611–614, doi:10.1038/35020537, 2000.

Kovalev, V. and Eichinger, W.: Elastic Lidar, Wiley Online Library, 2004.

5

10

35

- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Zöger, M., Smith, J., Herman, R., Buchholz, B., Ebert, V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds – Part 1: Cirrus types, Atmos. Chem. Phys., 16, 3463-3483, doi:10.5194/acp-16-3463-2016, 2016.
- 15 Larchevêque, G., Balin, I., Nessler, R., Quaglia, P., Simeonov, V., van den Bergh, H. and Calpini, B.: Development of a multiwavelength aerosol and water-vapor lidar at the Jungfraujoch Alpine Station (3580 m above sea level) in Switzerland, Appl. Opt., 41,15,2781–2790, doi:10.1364/AO.41.002781, 2002.
 - Larroza, E. G., Nakaema, W. M., Bourayou, R., Hoareau, C., Landulfo, E., and Keckhut, P.: Towards an automatic lidar cirrus cloud retrieval for climate studies, Atmos. Meas. Tech., 6, 3197-3210, doi:10.5194/amt-6-3197-2013, 2013.
- 20 Li, R., Cai, H. K., Fu, Y. F., Wang, Y., Min, Q. L., Guo, J. C., and Dong, X.: The optical properties and longwave radiative forcing in the lateral boundary of cirrus cloud, Geophysical Research Letters, 41, 3666–3675, doi:10.1002/2014GL059432, 2014.
 - Lin, H., Noone, K. J., Ström, J. and Heymsfield, A.J.: Small Ice Crystals in Cirrus Clouds: A Model Study and Comparison with In Situ Observations. J. Atmos. Sci., 55, 1928–1939, doi: http://dx.doi.org/10.1175/1520-0469(1998)055<1928:SICICC>2.0.CO;2, 1998

Lin, H. and Noone, K. J. and Ström, J. and Heymsfield, A. J: Dynamical Influences on Cirrus Cloud Formation Process, J. Atmos. Sci., 55,

- 25 1928-1939, http://dx.doi.org/10.1175/1520-0469(1998)055<1940:DIOCCF>2.0.CO;2, 1998.
- Liou, K. N., Ou, S. C. and Koenig, G.: An investigation of the climatic effect of contrail cirrus. In: Air Traffic and the Environment Background, Tendencies and Potential Global Atmospheric Effects. U. Schumann (Ed.), Lecture Notes in Engineering, Springer Berlin, 154-169, 1990.
 - Liou, K.N.: An Introduction to Atmospheric Radiation, Second Edition, Academic Press, Elsevier Science, San Diego, USA, 2002.
- 30 Mannstein, H., Meyer, R. and Wendling, P.: Operational detection of contrails from NOAA-AVHRR data, Int. J. Remote Sensing, 20, 1641-1660, doi: 10.1080/014311699212650, 1999.
 - Marchand, R., Ackerman, T., Smyth, M. and Rossow, W. B.: A review of cloud top height and optical depth histograms from MISR, ISCCP, and MODIS, J. Geophys. Res., 115, D16206, doi:10.1029/2009JD013422, 2010.
 - Markowicz, K. M., and Witek, M.: Sensitivity study of global contrail radiative forcing due to particle shape, J. Geophys. Res., 116, D23203, doi:10.1029/2011JD016345, 2011.
 - Mattis, I., Tesche, M., Grein, M., Freudenthaler, V., and Müller, D.: Systematic error of lidar profiles caused by a polarization-dependent receiver transmission: quantification and error correction scheme, Applied Optics, 48, 2742–2751, doi:10.1364/AO.48.002742, 2009.

- Meyer, R., Mannstein, H., Meerkötter, R., Schumann, U.and Wendling, P.: Regional radiative forcing by line-shaped contrails derived from satellite data, J. Geophys. Res., 107, ACL 17-11 ACL 17-15, 10.1029/2001jd000426, 2002.
- Min, M., Wang, P. C., Campbell, J. R., Zong, X. M., and Li, Y.: Midlatitude cirrus cloud radiative forcing over China, Journal of Geophysical Research-atmospheres, 115, D20 210, doi:10.1029/2010JD014161, 2010.
- 5 Oreopoulos, L. and Rossow, W. B.: The cloud radiative effects of International Satellite Cloud Climatology Project weather states, Journal of Geophysical Research-atmospheres, 116, D12 202, doi:10.1029/2010JD015472, 2011.
 - Pal, S.R., Carswell, A.I., Gordon, I. and Fong, A.: Lidar-Derived Cloud Optical Properties Obtained during the ECLIPS Program, Journal of Applied Meteorology, Vol. 34, 2388-2399, 1995.

Pincus, R., Platnick, S., Ackerman, S. A., Hemler, R. S. and Hofmann, R. J. P.: Reconciling simulated and observed views of clouds: modis,

10 isccp, and the limits of instrument simulators. J. Climate, 25, 4699–4720, doi: http://dx.doi.org/10.1175/JCLI-D-11-00267.1, 2012.

Platt, C. M. R., Abshire, N. L. and McNice, G. T.: Some microphysical properties of an ice cloud from lidar observation of horizontally oriented crystals, Journal of Applied Meteorology, 17, 1220–1224, 1978.

Platt, C. M., Young, S. A., Carswell, A. I., Pal, S. R., McCormick, M. P., Winker, D. M., DelGuasta, M., Stefanutti, L., Eberhard, W. L., Hardesty, M., Flamant, P. H., Valentin, R., Forgan, B., Gimmestad, G.G., Jäger, H., Khmelevtsov, S.S., Kolev, I., Kaprieolev, B.,

- 15 Lu, Da-ren, Sassen, K., Shamanaev, V.S., Uchino, O., Mizuno, Y., Wandinger, U., Weitkamp, C., Ansmann, A. and Woolridge, C.: The Experimental Cloud Lidar Pilot-Study (ECLIPS) for Cloud-Radiation Research, Bulletin of the American Meteorological Society, 75, 9, 1635-1654, 1994.
 - Platt, C.M.R. and Harshvardhan: Temperature Dependence of Cirrus Extinction: Implications for Climate Feedback, J. Geophys. Res., 93, D9, 11051–11058, 1988
- 20 Pruppacher, H. R. and Klett, J. D.: Microphysics und Clouds and Precipitation, Kluwer Academic Publishers New York, ISBN: 0-7923-4211-9, 1997
 - Radlach, M.,Behrendt, A. and Wulfmeyer, V.: Scanning rotational Raman lidar at 355 nm for the measurement of tropospheric temperature fields, Atmos. Chem. Phys., 8,2,159–169, doi 10.5194/acp-8-159-2008, 2008.

Reichardt, J., Reichardt, S., Hess, M. and McGee, T. J.:Correlations among the optical properties of cirrus-cloud particles: Microphysical

interpretation, J. Geophys. Res.: Atmospheres, 107, D21, 2156-2202, doi:10.1029/2002JD002589, 2002.
 Rolf, C., Krämer, M., Schiller, C., Hildebrandt, M., and Riese, M.: Lidar observation and model simulation of a volcanic-ash-induced cirrus cloud during the Eyiafjallajökull eruption, Atmos. Chem. Phys., 12, 10281–10294, doi:10.5194/acp-12-10281-2012.

Rolf, C.: Lidar observations of natural and volcanic-ash-induced cirrus clouds, Bergische Universität Wuppertal, 2012.

- Rossow, W. B. and Schiffer, R. A.: Advances in understanding clouds from ISCCP, Bull. Amer. Meteor. Soc., 80, 2261–2287. doi: http://dx.doi.org/10.1175/1520-0477(1999)080<2261:AIUCFI>2.0.CO;2, 1999.
 - Sassen, K. and Cho, B.S.: Subvisual-Thin Cirrus Lidar Dataset for Satellite Verification and Climatological Research. J. Appl. Meteor., 31, 1275–1285. doi: http://dx.doi.org/10.1175/1520-0450(1992)031<1275:STCLDF>2.0.CO;2, 1992.

Sassen, K. and Campbell, J.R.: A Midlatitude Cirrus Cloud Climatology from the Facility for Atmospheric Remote Sensing. Part I: Macrophysical and Synoptic Properties. J. Atmos. Sci., 58, 481-496, doi:http://dx.doi.org/10.1175/1520-0460(2001)058-0481:AMCCCE>2.0 CO:2.2001

- 35 0469(2001)058<0481:AMCCCF>2.0.CO;2, 2001.
 - Sassen, K and Benson, S.: A Midlatitude Cirrus Cloud Climatology from the Facility for Atmospheric Remote Sensing. Part II: Microphysical Properties Derived from Lidar Depolarization. J. Atmos. Sci., 58, 2103–2112, doi: http://dx.doi.org/10.1175/1520-0469(2001)058<2103:AMCCCF>2.0.CO;2, 2001.

- Sassen, K. and Comstock, J.M.: A Midlatitude Cirrus Cloud Climatology from the Facility for Atmospheric Remote Sensing. Part III: Radiative Properties, Journal of the Atmospheric Sciences, Vol. 58, 2001.
- Sassen, K., Zhu, J. and Benson, S.: Midlatitude cirrus cloud climatology from the Facility for Atmospheric Remote Sensing. IV. Optical displays, Appl. Opt. 42, 332-341, 2003.
- 5 Sassen, K., Wang, L., Starr, D. O'C., Comstock, J.M. and Quante, M.: A Midlatitude Cirrus Cloud Climatology from the Facility for Atmospheric Remote Sensing. Part V: Cloud Structural Properties. J. Atmos. Sci., 64, 2483–2501, doi: http://dx.doi.org/10.1175/JAS3949.1, 2007.
 - Sassen, K. and Wang, Z. and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) measurements, J. Geophys. Res., 118, D00A12, doi:10.1029/2008JD009972, 2008.
- 10 Schotland, R.M., Sassen, K. and Stone, R.:Observations by Lidar of Linear Depolarization Ratios for Hydrometeors, J. Appl. Meteor., 10, 1011–1017, http://dx.doi.org/10.1175/1520-0450(1971)010<1011:OBLOLD>2.0.CO;2, 1971.
 - Schumann, U., Graf, K., Mannstein, H. and Mayer, B.: Contrails: Visible aviation induced climate impact, in Atmospheric Physics Background - Methods - Trends, edited by U. Schumann, pp. 239-257, Springer, Berlin, Heidelberg, DOI: 10.1007/978-3-642-30183-415, 2012.
 - Schumann, U., Mayer, B., Graf, K. and Mannstein, H.:A parametric radiative forcing model for contrail cirrus, J. Appl. Meteorol. Clim., 51, 1391-1406, doi: 10.1175/JAMC-D-11-0242.1, 2012.

Schumann, U., Penner, J. E., Chen, Y., Zhou, C. and Graf, K.: Dehydration effects from contrails in a coupled contrail-climate model, Atmos.

- 15 Chem. Phys., 15, 11179-11199, doi:10.5194/acp-15-11179-2015, 2015.
 - Seifert, P. and Ansmann, A. and Müller, D. and Wandinger, U. and Althausen, D. and Heymsfield, A. J. and Massie, S. T. and Schmitt, C.: Cirrus optical properties observed with lidar, radiosonde, and satellite over the tropical Indian Ocean during the aerosolpolluted northeast and clean maritime southwest monsoon, Journal of Geophysical Research: Atmospheres, 112, D17, 2156-2202, http://dx.doi.org/10.1029/2006JD008352, doi:10.1029/2006JD008352, 2007.
- 20 She, S.: Spectral structure of laser light scattering revisited: bandwidths of nonresonant scattering lidars, Appl. Opt., 40, 4875-4884, 2001. Soden, B. J., and Donner, L. J.: Evaluation of a GCM cirrus parameterization using satellite observations, J. Geophys. Res., 99(D7), 14401–14413, doi:10.1029/94JD00963, 1994.
 - Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., and Miller, H.L. (eds.): IPCC 2007: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on
- 25 Climate Change, Cambridge University Press, 2007
 - Vázquez-Navarro, M., Mannstein, H. and Kox, S.: Contrail life cycle and properties from 1 year of MSG/SEVIRI rapid-scan images, Atmos. Chem. Phys., 15, 8739- 8749, doi:10.5194/acp-15-8739-2015, 2015.
 - Wandinger, U.: Multiple-scattering influence on extinction- and backscatter-coefficient measurements with Raman and high-spectral-resolution lidars, Appl. Opt., 37, 417–427, 1998.
- 30 Wandinger, U.: Raman Lidar in LIDAR: Range–Resolved Optical Remote Sensing of the Atmosphere,241–271, Springer,2005.
 Wandinger, U., Tesche, M., Seifert, P., Ansmann, A., Müller, D. and Althausen, D.: Size matters: Influence of multiple scattering on CALIPSO light-extinction profiling in desert dust, Geophysical Research Letters,37,10,1944-8007, http://dx.doi.org/10.1029/2010GL042815, 2010.
 - Wang, Z., and Sassen, K.: Cirrus cloud microphysical property retrieval using lidar and radar measurements. Part II: Midlatitude cirrus microphysical and radiative properties, J. Atmos. Sci., 59, 2291–2302, 2002.
- 35 Wernli, H. and Davies, H. C.: A lagrangian-based analysis of extratropical cyclones. I: The method and some applications, Q. J. Roy. Meteor. Soc., 123, Issue 538, 467–489, January 1997 Part B, doi:10.1002/qj.49712353811, 1997.

- Westbrook C.D., Illingworth A.J., O'Connor E.J., Hogan R.J: Doppler lidar measurements of oriented planar ice crystals falling from supercooled and glaciated layer clouds, Q. J. R. Meteorol. Soc., 136, 260 – 276. DOI:10.1002/qj.528, 2010.
- Winker, D.M. and Vaughan, M.A.: Vertical distribution of clouds over Hampton, Virginia observed by lidar under the ECLIPS and FIRE ETO programs, Atmospheric Research, 34, 117-133, 1994.
- 5 Yorks, J.E., Hlavka, D.L., Hart, W.D. and McGill, M.J.: Statistics of Cloud Optical Properties from Airborne Lidar Measurements. J. Atmos. Oceanic Technol., 28, 869–883,2011. doi: http://dx.doi.org/10.1175/2011JTECHA1507.1
 - Zieger, P., Kienast-Sjögren, E., Starace, M., von Bismarck, J., Bukowiecki, N., Baltensperger, U., Wienhold, F. G., Peter, T., Ruhtz, T., Collaud Coen, M., Vuilleumier, L., Maier, O., Emili, E., Popp, C., and Weingartner, E.: Spatial variation of aerosol optical properties around the high-alpine site Jungfraujoch (3580 m a.s.l.), Atmos. Chem. Phys., 12, 7231–7249, doi:10.5194/acp-12-7231-2012, 2012.