

## **Response Letter to Reviewers' comments:**

We would like to thank both reviewers for their comments and recommendations. We believe that we have corrected and improved the paper by incorporating their comments, in the revised version. The figure proposed by the first reviewer was a very good idea where we had to clarify several points of 'our story' to provide sufficient context. We reran the simulations at higher resolution, replaced figures and modified the discussion, accordingly. The main changes are the following:

Following both Reviewers' comment, regarding the model's estimation of the simulated new particle formation, we reran the model by ignoring NPF process.

In the revised manuscript, section 3.5 is divided in 2 sections: 3.5 is called "*Impact of NPF events on CCN production*" and 3.6 "*Impact of NPF events on cloud droplet number.*"

We followed reviewer's suggestion to use for the two types of northern flow the terms: Etesian Flow (EF) and Moderate Surface Flow (MSF), in order to have a more concise wording. We also followed the same formalism in the revised Tables and Figure captions.

## **Reviewer #1:**

### **General Comments:**

*Kalkavouras and co-authors present results from an intriguing experiment in the Aegean Sea. The nature of pollution arriving at the long-standing Finokalia measurement platform is investigated directly with observations at Santorini, a site strategically located along the trajectory from the European mainland. The observations target aerosol size distributions, relevant primary and secondary pollutant concentrations, and detection of new particle formation events. The authors identify two events representing characteristic flow from the north, albeit of two distinct types. The Etesian flow example is marked with NPF events at both sites, although the events are stronger at the Santorini site. The authors extend their observations of particle number and composition to predictions of CCN at various supersaturations. They also go further to predict the total effect of the NPF events on cloud droplet number, taking into account the impact of constrained available water vapor. The paper starts with a nice scientific idea and goes into good detail into the results. What I see lacking is a little more connective tissue linking the observations at each site with each other and the mainland into a cohesive story. The material is already there, but it is somewhat buried and could be highlighted with a figure, for instance. I would like to see the following points addressed before recommending publication:*

### **Specific Comments:**

*1. I recommend the authors adopt two shorthand names for the distinct periods (22/7- 24/7 and 25/7-27/7). They could be referred to as "Etesian Flow" and "Moderate Surface Flow," for example. Small changes like this could help the*

*readability of the paper. A useful addition to this paper would be a two-panel cartoon, each overlaid on the Greek domain map in Fig. 1, for example, that describes the factors at play in these two periods. They could identify generally where they expect emissions, mixing, oxidation, NPF, and aging of new particles to be happening.*

To be more concise, we used for the two types of northern flow the terms: Etesian Flow (EF) and Moderate Surface Flow (MSF). We also followed the same formalism in the revised Tables and Figure captions. Figure 1 is replaced with a two-panel cartoon in order to identify the locations where we expect/identify the various processes.

*2. The WRF-Chem aerosol module configuration, as documented by the authors, is problematic for this particular application. It is quite likely that NPF events and subsequent processing are not captured realistically at all by the model. The sulfuric acid/water pathway parameterized by Kulmala et al. (1998) is likely not strong enough to enhance particles near the surface and lower troposphere to the levels observed at the Santorini site. It is now well-documented that other reagents play important roles in this process (e.g. NH<sub>3</sub> and organics), and these pollutants have been identified by the authors to be present and significant components of the aerosol. My guess is that most of the Aitken-mode particles in the model originate from direct emissions, not from secondary generation. A related issue is the lack of a dedicated nucleation mode in the model. Without this mode, any NPF events will artificially broaden the Aitken mode distribution and give unrealistic lifetimes against deposition and coagulation. It will also affect the growth rates predicted by the model.*

*The authors astutely sidestep relying on the model to predict size distributions and use their own observations when possible for calculating CCN and cloud drop number concentrations. However, since they include an entire section (2.2) detailing the regional modeling they performed, it is a good idea to explicitly state the limitations of this analysis for particle size distributions, and remind the reader that they are using the WRF-Chem output for its knowledge of advection flows and chemical composition, not microphysical processing.*

We agree with the Reviewer's comment. In version 3.3 the aerosol models are not appropriate to simulate the NPF events realistically. Luo and Yu (2011), discuss the need to improve the representation of the nucleation process in earlier versions of WRF-Chem. We believe that more research is needed regarding the nucleation modeling in the area, which we plan to perform in a separate paper in the near future. Nevertheless, we conducted another simulation ignoring the nucleation parameterization in order to comprehend/emphasize the spatial extent of these processes and present them in Fig. 1. In the revised manuscript, the relevant discussion on model limitations (due to lack of a dedicated nucleation mode, nucleation parameterization) is presented in section 2.3 'Regional modeling' (page 7, lines 6-21).

To further elucidate the conditions under which NPF events take place in our region, we reran the model by ignoring the NPF process. In the revised manuscript our

hypothesis is not based on the simulated Aitken-mode particles but on the number concentration differences considering and ignoring NPF process. The relevant discussion is presented in section 3.4 ‘Spatial extent of NPF event (pages 13-16).

***3. The authors conclude that the NPF events observed at Santorini are regional in nature with a spatial scale of 250 km and characteristic transport time of 4.5 hours. They also assert that Finokalia does not see the bursts because it is 3 hours away and particle populations age before they arrive there. The authors do note that this second observation demonstrates how site-to-site variability can be important during a regional event. I am not sure that this totally addresses the issue though. Why are the events sort of regional and sort of not-regional? Is this an issue of using up the NPF precursors before the air mass gets to Santorini and then shifting to chemical conditions that favor condensation to available surface area? Or is something else at play here?***

We used the regional characterization, mainly because the number of particles remained high for several hours at Santorini (Kulmala et al., 2012). In addition, the NPF event was found to extend over hundreds of kilometers. Thus, as the reviewer points out, we tried to relate these fine aerosols to regional sources of pollution transferred by long-range transport (LRT) during Etesian flow conditions. Despite that, we observed local variability at sub-regional scales, due to the differences in geographical and atmospheric conditions between stations along the same trajectory. This is the case between Santorini and Finokalia stations. We also expect that local variability is unavoidable at smaller scales, over locations at the same distance from the sources. This is based on the simulations that show that the spatial differences of chemical and physical properties in the initial steps of the formation, under the stable Etesian flow, produce streams with different characteristics, especially upwind of Santorini.

***4. The paragraph beginning line 13 on page 9 describes an interesting hypothesis for how pollutants are transported to the middle of the Aegean Sea with limited aging. However, I'm not convinced there is enough evidence to warrant the detailed discussion that is given to this possibility or the certainty with which it is treated in the conclusions section. As described in my first comment, any model data related to the size distribution of Aitken-mode particles probably cannot be trusted in this case. If I understand correctly, the main assertion here is that the particles were formed over the Turkish mainland and transported quickly before they have a chance to be significantly coagulated away. Why could the enhanced number concentrations not come instead from oxidation and NPF over the water during transport, where there may be enhanced photochemistry, complex interactions with clouds, interesting boundary layer phenomena, etc? If I'm not understanding the meat of the argument correctly, please explain it to me and consider rephrasing it in the text to be clearer. What insight do the model CO concentrations help to provide regarding the stratification and mixing of distinct layers downwind of the continent?***

The revised section 3.4 ‘Spatial extent of NPF event’ (pages 13-16) explains more clearly now the processes taking place. The discussion is mainly based on the number concentration differences considering and ignoring NPF process and not on the Aitken-mode simulated particles. Although the model severely underestimates the NPF, the decisive role of the Etesian flow on the evolution of the phenomenon over the Aegean Sea is evident (from page 13 line 24 to page 15 line 1).

The atmospheric conditions under a similar Etesian event have been studied thoroughly in a separate paper that has been submitted to BLM. In particular, the heat fluxes simulated and calculated from airborne measurements over the AS (Tombrou et al., 2015) varied from  $-25 \text{ W m}^{-2}$  (over the northeastern AS) to  $25 \text{ W m}^{-2}$  (over the southeastern AS).

Furthermore, vertical cross-sections of measured CO concentrations along the eastern AS under an Etesian flow, are shown in Fig. 7 by Tombrou et al. (2015). The strong gradient of stratification and mixing downwind of AS, is apparent. In particular, at  $40^\circ$  latitude, where the plume leaves the Turkey continent, the vertical mixing extends up to 500-600m height according to the CO vertical extent. Above the Cyclades complex (lat  $36.5^\circ - 38^\circ$ ) the mixing extends up to 1km and gradually increases up to 2km, upwind of Crete (Finokalia at  $35^\circ$ ).

***5. I recommend separating the paragraphs detailing the Nd calculations (starting on Page 12) into their own section, perhaps called “Impact of NPF events on cloud droplet number.” Then section 3.5 would be called “Impact of NPF events on CCN production.”***

Done

***6. How is the partial sensitivity of cloud droplet number to chemical composition and vertical velocity determined? Can the equations be provided? What is the uncertainty associated with this? Please document it if possible.***

The reviewer raises a good question. The sensitivity is derived from the parameterization using either a direct sensitivity or finite difference approach, as described in Karydis et al. (2012). Here we use the finite difference implementation. This information is now given on page 6; Ln 16-18.

The accuracy of the method, i.e. the ability of the parameterization to capture the sensitivity of droplet number to each parameter examined was explored in detail by Morales and Nenes (2014). Given that the parameterization has been shown to give cloud droplet closure in ambient clouds to within experimental uncertainty (Meskhidze et al., 2005; Fountoukis et al., 2007; Hoyle et al., 2016), and that the same parameterization also reproduces the droplet number and sensitivities of the detailed numerical simulation with high fidelity (Morales and Nenes, 2014), we expect the sensitivities and attribution calculations presented here to be representative of the ambient clouds in the study region.

## Minor Changes/Typos:

***Pg 2, Ln 25: The phrase “without any particular seasonal preference about their occurrence” is difficult to understand. Can the authors please reword this to be more specific?***

The phrase has been replaced by (Pg 3, Ln 11-12): “Most of these ground-based observations indicate that the mass of fine aerosols presents a summer maximum, however the frequency of the events is season independent.”

***Pg 3, Ln 4: “prior to reaching”***

Done

***Pg 5, Ln 11-13: This is technically not a sentence.***

Replaced by (page 8, lines 6-8): “Air mass origin and trajectories were determined by HYSPLIT4 (Hybrid Single-Particle Lagrangian Integrated Trajectory; [www.arl.noaa.gov/ready/hysplit4.html](http://www.arl.noaa.gov/ready/hysplit4.html)) back-trajectory analysis (Draxler and Rolph, 2015).”

***Fig. 3 and caption: “open circles” not “cycles”.***

Done.

***Also, please indicate on the figure that the solid lines describe wind speed and the circles describe direction. It is hinted at in the figure and explained in the caption, but it would be quicker for the reader to have it identified visually, with an arrow or something.***

Figure 3 is replaced by a new one, where the abbreviations ‘ws’ and ‘wd’ are now included indicating wind speed and wind direction respectively. The caption is rephrased accordingly: ‘Time series of the wind speeds (ws, solid lines on left axis) and wind directions (wd, open circles on right axis) at Santorini (simulations by the WRF-Chem model) and at Finokalia (measurements). The second period of the EF is shaded with yellow and the MSF period with grey.’

***Pg 6, Ln 16-17: The “less pronounced” diurnal cycle at Finokalia for ozone is not obvious to me from Fig. S4. Please include a plot of the actual diurnally averaged profiles or report the daily minima and maxima to demonstrate this point.***

We refer to the Etesian period (EF) that corresponds to the yellow panel in Fig. S4. The mean diurnal range at Finokalia station (from 21 to 24 July) is 8 ppbv, while at Santorini, for the same period is 18 ppbv (Fig. S4).

This information is now included in the text (page 9, lines 22-24)

***Pg 6, Ln 28-29: I would not characterize -21% or -15% under-prediction as “small”. Either establish what they are small compared to, or please get rid of this qualification.***

We agree with the reviewer’s comment, therefore, we decided to delete the word ‘small’.

***Pg 6, Ln27-30: Please break this sentence up. It is long and confusing.***

Replaced by (page 10, lines 9-13): Simulations confirm that the air masses received at both stations during the prevailing strong northern wind are of the same origin, and representative of EF conditions (Fig. S3) albeit with an O<sub>3</sub> under-prediction (average bias during afternoon hours up to -21% on 23 and -15% on 24 July, Fig. 5). During the MSF period, simulations indicate an O<sub>3</sub> increase, especially in the southern AS, but also underpredicted (average bias during afternoon hours up to -24% on 26 July, Fig. 5).

***Pg 6, Ln 31-32: Is there a more recent or relevant reference than McKeen et al. (1991)? Anything that specifically identifies this model scenario or modern European scenarios in general as suffering from ozone boundary condition issues?***

The chemical boundary conditions used in this modeling study are hardcoded in the WRF-Chem model. The values are based on an idealized, northern hemispheric, mid-latitude, clean environmental, vertical profile from the NOAA Aeronomy Lab Regional Oxidant Model (NALROM) (Liu et al. 1996; Peckham et al. 2011). This information has been added to section 2.3 ‘Regional modeling’ (page 7, lines 25-27 while the reference of McKeen et al. (1991) was omitted.

***Pg 7, Ln 5-6: In what way did the inorganic and organic mass concentrations show “similar behavior” to that of ozone? Are the authors just identifying them all as secondary pollutants? Please provide an estimate of the correlation coefficient or index of agreement for a statement like this.***

We identify all of them as secondary pollutants driven by the same meteorological conditions. The R<sup>2</sup> of O<sub>3</sub> to Organics and O<sub>3</sub> to inorganics is 0.5 and 0.59 respectively (included in the revised manuscript on page 10, line 22). Also, the simulated spatial patterns of O<sub>3</sub> and sulfates are similar, for each period (EF and MSF).

***Pg 7, Ln 9: Please remove the comma after the parentheses.***

Done

***Pg 7, Ln 20: Please refer to some quantitative statistics to back up this claim.***

An extended evaluation of WRF-Chem model against airborne and ground observations over the AS during the Etesians is presented in Bossioli et al. (2016). In that study biomass burning emissions were also included.

After the reviewer’s suggestion some statistics have been added in the revised manuscript:

For EF period (Page 11, lines 3-4): “...on average during EF underprediction of 30% for sulfates and 60% for ammonium”

For MSF period (Page 11, lines 12-15): “(simulated and observed concentrations correlate during both periods R<sup>2</sup>=0.8), however they are lower than the measured values at Finokalia station (on average underprediction of 50% for sulfates and 75% for ammonium).

***Pg 10, Ln 6-8: This sentence is worded in a confusing way.***

The discussion has been revised (page 15, lines 10-17). The specific sentence has been revised to “*The nucleation-mode particles are significantly reduced as they have shifted gradually towards larger sizes (Aitken-mode), before reaching Finokalia (Fig. 4).*”

*Pg 10, Ln 25 – Pg 11, Ln 19: Most of this material would be better-placed in the methods section (2.3 maybe). This goes for the second paragraph a page 12 as well.*

Done

## References:

Bossioli, E., Tombrou, M., Kalogiros, J., Allan, J., Bacak, A., Bezantakos, S., Biskos, G., Coe, H., Jones, B. T., Kouvarakis, G., Mihalopoulos, N., and Percival, C. J.: Atmospheric composition in the Eastern Mediterranean: Influence of biomass burning during summertime using the WRF-Chem model, *Atmos. Environ.*, 132, 317–331, 2016

Draxler, R.R. and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY Website (<http://ready.arl.noaa.gov/HYSPLIT.php>). NOAA Air Resources Laboratory, Silver Spring, MD., 2015

Fountoukis, C., Nenes, A., Meskhidze, N., Bahreini, R., Brechtel, F., Conant, W. C., Jonsson, H., Murphy, S., Sorooshian, A., Varutbangkul, V., R. C. Flagan, and J. H. Seinfeld Aerosol–cloud drop concentration closure for clouds sampled during ICARTT, *J.Geoph.Res.*, 112, D10S30, doi:10.1029/2006JD007272, 2007

Hoyle, C.R., Webster, C.S., Rieder, H.E., Nenes, A., Hammer, E., Herrmann, E., Gysel, M., Bukowiecki, N., Weingartner, E., Steinbacher, M., and U. Baltensperger Chemical and physical influences on aerosol activation in liquid clouds: a study based on observations from the Jungfraujoch, Switzerland, *Atmos.Chem.Phys.*, 16, 4043–4061, 2016

Karydis, V. A., Capps, S. L., Russell, A. G., and Nenes, A.: Adjoint sensitivity of global cloud droplet number to aerosol and dynamical parameters, *Atmos. Chem. Phys.*, 12, 9041–9055, doi:10.5194/acp-12-9041-2012, 2012

Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V.-M.: Measurement of the nucleation of atmospheric aerosol particles. *Nature Protocols* 7, 1651–1667, 2012.

Liu, S. C., McKeen, S. A., Hsie, E-Y., Lin, X., Kelly, K. K., Bradshaw, J. D., Sandholm, S. T., Browell, E. V., Gregory, G. L., Sachse, G. W., Bandy, A. R., Thornton, D. C., Blake, D. R., Rowland, F. S., Newell, R., Heikes, B. G., Singh, H., Talbot, R. W.: Model study of tropospheric trace species distributions during PEM-West A, *J. Geophys. Res.*, 101(D1), 2073–2085, doi:10.1029/95JD02277, 1996

Luo, G. and Yu, F.: Simulation of particle formation and number concentration over the Eastern United States with the WRF-Chem + APM model, *Atmos. Chem. Phys.*, 11, 11521-11533, doi:10.5194/acp-11-11521-2011, 2011.

Meskhidze, N., A. Nenes, Conant, W. C., and Seinfeld, J.H.: Evaluation of a new Cloud Droplet Activation Parameterization with In Situ Data from CRYSTAL-FACE and CSTRIFE, *J.Geoph.Res.*, 110, D16202, doi:10.1029/2004JD005703, 2005

Morales Betancourt, R., and Nenes, A.: Aerosol Activation Parameterization: The population splitting concept revisited, *Geosci.Mod.Dev.*, 7, 2345–2357, 2014

Peckham, S., G. A. Grell, S. A. McKeen, M. Barth, G. Pfister, C. Wiedinmyer, J. D. Fast, W. I. Gustafson, R. Zaveri, R. C. Easter, J. Barnard, E. Chapman, M. Hewson, R. Schmitz, M. Salzmann, S. Freitas: WRF/Chem Version 3.3 User's Guide. NOAA Technical Memo., 98 pp., 2011

Tombrou, M., Bossioli, E., Kalogiros, J., Allan, J. D., Bacak, A., Biskos, G., Coe, H., Dandou, A., Kouvarakis, G., Mihalopoulos, N., Percival, C. J., Protonotariou, A. P., and Szabó-Takács, B.: Physical and chemical processes of air masses in the Aegean Sea during Etesians: Aegean-GAME airborne campaign, *Sci. Total Environ.*, 506-507, 201–216, doi:10.1016/j.scitotenv.2014.10.098, 2015



## Reviewer#2

*The manuscript presents measurements of the number size distribution and chemical composition of submicron aerosols at two islands in the Eastern Mediterranean. The analysis is based on a measurement period over two weeks in the summer 2013, during persistent transport of continental airmasses from north to the sites. A chemical transport model and airmass back-trajectories are used to identify the source areas and transport routes of aerosols to the sites. Using case studies of two new particle formation (NPF) events the contribution of NPF to both the cloud condensation nuclei (CCN) and cloud droplet (Nd) concentrations is assessed. The results for CCN and Nd are based on Köhler theory and parameterizations.*

*I agree with the comments presented by the anonymous referee #1, and would like the authors to address my further comments below. After addressing these comments I can recommend the manuscript for publication in Atmospheric Chemistry and Physics.*

### General comments:


*Page 5, lines 6–7: Is it known what are the possible reasons for the underestimation of organic matter concentrations in the model results; could it be due to underestimation of primary emissions or underestimation of SOA formation in the model?*

The biases are probably related to the underestimated POA emissions but also to the limitation of the RADM2 mechanism regarding the treatment of monoterpene emissions (Tuccella et al., 2012). This information is now included on page 8, lines 4–5.

WRF-Chem simulations over the Aegean Sea during Etesian flow revealed that the simulated SOA, formed from anthropogenic and biogenic emissions, contributes respectively to less than 5% and almost negligibly to the OM (Bossioli et al., 2016). The importance of secondary aerosols over the area has been pointed out in earlier works (Athanasopoulou et al., 2015; Fountoukis et al., 2011)

*Page 5, lines 8–12: Care should be taken when using the HYSPLIT model with the GDAS 0.5\_ input data: the back-trajectory results might differ from those obtained with GDAS 1\_ input data due to the differences in the airmass vertical advection calculation method between these two datasets (see e.g. Su et al., 2015). Perhaps the authors could check that their back-trajectories shown in Figure 2 remain the same if using the GDAS with 1\_ resolution as input meteorological data.*

There are no significant differences, especially at low levels. The differences are mainly noticed on the 24th, but they do not change the hypothesis that air masses are better mixed throughout the boundary layer, covering a broader area over Asian Turkey.



*Page 8, lines 6–7: Why are coagulation losses not included in the calculation of the formation rate of nucleation mode particles? This should be fairly straightforward to calculate based on the measured size distributions, and including the coagulation losses would make the calculated formation rates more readily comparable to literature values (which typically account for coagulation).*

Both coagulation flux and condensational growth are now included in the calculations. The text has been modified accordingly (page 12, lines 10-14).

*Page 10, line 4: Where does the 3 hour difference in the comparison between particle observations at Santorini and Finokalia come from? Based on the particle size distribution data in Fig. 8 the particle formation at both stations seems to start at 9 a.m. on 23 July, and the only appreciable difference in the particle concentrations in Fig. 4 seems to be in the nucleation mode concentration (i.e. intensity of particle formation). Regarding the discussion on the CCN-sized particles and the calculated hygroscopicity parameters, it would be interesting to see how the results differ on days without new particle formation. This type of comparison between NPF and non-NPF days would put the results presented in the manuscript better into context with regard to the importance of NPF to CCN and cloud droplet number at the Aegean Sea. Where there during the campaign any such non-NPF days for which the parameters of Table 3 could be calculated and reported for comparison with the two NPF days?*

We agree with the reviewer that this was not clear in the text. The air masses spent 3-4 h to reach Finokalia after Santorini, according to HYSPLIT (Fig. S3 left panel), on 23 July. The 3-h transit timescale is in agreement with the prevailing wind speed (about  $10 \text{ m s}^{-1}$ ; Fig. S1) and the 120 km distance between Santorini and Finokalia. For this reason, we claim that the air masses reaching Finokalia earlier (Fig. 4) are probably due to a local nucleation event initiated at Heraklion (Crete).

Throughout the non-NPF events (MSF period), the CCN concentrations decrease by almost 48% and 23% at Santorini and Finokalia respectively, compared to the levels during the NPF events. We have added this information on page 17 (lines 22-24) but we decided not to change Table 3.

#### **Minor and technical comments:**

***Page 2, line 30: The sentence starting with “Short-lived events of small number young Aitken particles” is difficult to understand, consider revising it. Does “small number” refer to low concentrations?***

This sentence has been replaced by (page 3, lines 18-19): “A few short-lived particle formation events (18–25 nm) were first recorded at Finokalia by Kalivitis et al. (2008), arriving with low speed from the west, during autumn.”

***Page 3, line 4: should be “prior to reaching ”***

Done

***Page 7, line 2: “non-refractive” should be “non-refractory”***

Done

***Page 8, line 21: A more recent reference for NPF event classification is Kulmala et al. (2012).***

Done

***Page 10, line 3: In the sentence “ : : : have trace a lower number of : : : ” the word “trace” should be omitted.***

Done

***Page 10, line 21: As also suggested by the other referee, Section 3.5 could be divided into two parts, one dealing with CCN concentrations and another dealing with cloud droplet concentrations. That would make this section more readable.***

Done

***Page 13, line 30: “: : : have a similar to ozone behavior : : : ” should be “: : : behave similarly to ozone : : : ”***

Done

#### **References:**

Athanasopoulou E., E.P. Protonotariou, E. Bossioli, A. Dandou, M. Tombrou, J.D. Allan, H. Coel, N. Mihalopoulos, J. Kalogiros, A. Bacak, J. Sciare, G. Biskos: ‘Aerosol chemistry above an extended Archipelago of the Eastern Mediterranean basin during strong northern winds’, Atmospheric Chemistry and Physics, 15, 8401-8421, doi: 10.5194/acp-15-8401-2015, 2015

Bossioli, E., Tombrou, M., Kalogiros, J., Allan, J., Bacak, A., Bezantakos, S., Biskos, G., Coe, H., Jones, B. T., Kouvarakis, G., Mihalopoulos, N., and Percival, C. J.: Atmospheric composition in the Eastern Mediterranean: Influence of biomass burning

during summertime using the WRF-Chem model, *Atmos. Environ.*, 132, 317–331, 2016

Fountoukis, C., P.N. Racherla, H.A.C. Denier van der Gon, P. Polymeneas, P.E. Haralabidis, A. Wiedensohler, C. Pilinis, and S.N. Pandis, 2011: Evaluation of a three-dimensional chemical transport model (PMCAMx) in the European domain during the EUCAARI May 2008 campaign. *Atmos. Chem. Phys.*, 11, 10331-10347, doi:10.5194/acp-11-10331-2011, 2011

Kalivitis, N., Birmili, W., Stock, M., Wehner, B., Massling, A., Wiedensohler, A., Gerasopoulos, E. and Mihalopoulos, N.: Particle size distributions in the Eastern Mediterranean troposphere, *Atmos. Chem. Phys.*, 8, 6729–6738, doi:10.5194/acp-8-6729-2008, 2008

# New Particle Formation in the South Aegean Sea during the Etesians: importance for CCN production and cloud droplet number

5 Kalkavouras P.<sup>1</sup>, Bossioli E.<sup>1</sup>, Bezantakos S.<sup>2</sup>, Bougiatioti A.<sup>3,7</sup>, Kalivitis N.<sup>3</sup>, Stavroulas I.<sup>3</sup>, Kouvarakis G.<sup>3</sup>, Protonotariou A. P.<sup>1</sup>, Dandou A.<sup>1</sup>, Biskos G.<sup>4,5</sup>, Mihalopoulos N.<sup>3,5,6</sup>, Nenes A.<sup>6,7,8,9</sup>, Tombrou M.<sup>1\*</sup>

<sup>1</sup> Department of Physics, Univ. of Athens, 15784 Athens, Greece

<sup>2</sup> Dept. of Environment, Univ. of the Aegean, Mytilene 81100, Greece

<sup>3</sup> Env. Chemical Processes Lab., Dept. of Chemistry, Univ. of Crete, Heraklion 71003, Greece

10 <sup>4</sup> Fac. of Civil Engineering and Geosciences, Delft Univ. of Technology, Delft 2628 CN, The Netherlands

<sup>5</sup> Energy Environment and Water Research Center, The Cyprus Institute, Nicosia 2121, Cyprus

<sup>6</sup> Inst. of Env. Research & Sustainable Development, Nat. Observatory of Athens, Greece

<sup>7</sup> School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

15 <sup>8</sup> School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA, USA

<sup>9</sup> Institute for Chemical Engineering Science, Foundation for Research and Technology Hellas, Patra, Greece

*Correspondence to:* Maria Tombrou ([mtombrou@phys.uoa.gr](mailto:mtombrou@phys.uoa.gr))

20 **Abstract.** This study examines how new particle formation (NPF) in the summertime Eastern Mediterranean affect CCN concentrations and cloud droplet formation. For this, the concentration and size distribution of submicron aerosol particles along with the concentration of trace gases and meteorological variables were studied over the central (Santorini) and south Aegean Sea (Finokalia, Crete) from 15 to 28 July 2013, a period that includes Etesian events and moderate surface northern  
25 winds. Particle nucleation bursts were recorded during the Etesian flow at both stations, with those observed at Santorini reaching up to  $1.5 \times 10^4$  particles  $\text{cm}^{-3}$ ; the fraction of nucleation-mode particles over Crete was relatively diminished, but a higher number of Aitken-mode was observed as a result of aging. Aerosol and photochemical pollutants covaried throughout the measurement period; lower concentrations were observed during the period of Etesian flow (e.g. 43 - 70 ppbv for ozone, 1.5 -5.7  $\mu\text{g}$   
30  $\text{m}^{-3}$  for sulfate), but were substantially enhanced during the period of moderate surface winds (i.e.,

increase of up to 32% for ozone, and 140% for sulfate). We find that NPF can double CCN number (at 0.1% supersaturation) but the resulting strong competition for water vapor in cloudy updrafts decreases maximum supersaturation by 14% and augments the potential droplet number only by 12%. Therefore, although NPF events may strongly elevate CCN numbers, the relative impacts on cloud droplet number  
5 (compared to pre-event levels) is eventually limited by water vapor availability and depends on the prevailing cloud formation dynamics and the aerosol levels associated with the background in the region.

**Keywords.** Ozone concentrations, Aerosol particle concentrations, Particle size distributions, Chemical  
10 composition, New particle formation, CCN production, droplet number, WRF-Chem simulations

## 1 Introduction

During summer and early autumn (warm period), the circulation over the Eastern Mediterranean (EM) is dominated by a persistent northerly flow known as Etesians (Tyrlis and Lelieveld, 2013). Under the  
15 prevalence of the Etesians, the advection of the air masses is pronounced over the EM rendering the atmospheric conditions as the most important factor for high concentrations of gases and aerosol particles even in remote areas. The scientific interest over the Aegean Sea (AS), which is part of the EM, has led to a number of experimental campaigns, during the warm period (Mihalopoulos et al., 1997; Paronis et al., 1998; Formenti et al., 2002a, b; Kouvarakis et al., 2002; Lelieveld et al., 2002;  
20 Zerefos et al., 2002), focusing initially on the interpretation of ozone (O<sub>3</sub>) enhancement under the Etesian regime. Apart from the simultaneous contribution of local and distant sources in the area, in the presence of enhanced photochemistry, strong subsidence was also identified in most of these events (Kallos et al., 1998, 2007; Lelieveld et al., 2002; Salisbury et al., 2003; Kalabokas et al., 2007, 2008, 2013; Kanakidou et al., 2011; Bossioli et al., 2016). Airborne measurements performed during an  
25 Etesian outbreak (Aegean-GAME campaign; Tombrou et al., 2015) have clearly shown that neutral to

stable atmospheric conditions prevailed over the north and central AS, with reduced friction velocities and absolute turbulent fluxes (momentum or heat) cumulating the concentrations below the planetary boundary layer (PBL) and mainly inside the shallow Marine Atmospheric Boundary Layer (MABL). Unstable conditions were found only over southeast AS, in the vicinity of Crete, resulting in enhanced friction velocities and large positive values of sensible heat flux.

Long-term aerosol observational studies in the EM have been mainly performed based on ground measurements collected at Finokalia, a remote coastal site in the northeastern part on the island of Crete (Bardouki et al., 2003; Eleftheriadis et al., 2006; Lazaridis et al., 2006; Gerasopoulos et al., 2007; Kalivitis et al., 2008, 2014, 2015; Koulouri et al., 2008; Querol et al., 2009; Pikridas et al., 2010, 2012) with a few more at Akrotiri station on western Crete (Lazaridis et al., 2008; Kopanakis et al., 2012, 2013). Most of these ground-based observations indicate that the mass of fine aerosols presents a summer maximum, however the frequency of the events is season independent. These fine aerosols have been related to regional sources of pollution enhanced by long-range transport (LRT) during the Etesian flow. In particular, a mixture of anthropogenic (Koçak et al., 2011), biogenic (Im and Kanakidou, 2012) and biomass burning emissions (Sciare et al., 2008; Bougiatioti et al., 2014) originating mainly from the Balkans and the central and Eastern Europe, result in enhanced aerosol concentrations in the southern AS (Kalivitis et al., 2014).

A few short-lived particle formation events (18–25 nm) were first recorded at Finokalia by Kalivitis et al. (2008), arriving with low speed from the west, during autumn. Thereafter, new particle formation (NPF) events have been frequently observed at Finokalia (Manninen et al., 2010; Ždímal et al., 2011; Pikridas et al., 2012; Kalivitis et al., 2014, 2015) and Akrotiri (Kopanakis et al., 2013) stations during different periods of the year, but more frequently during winter than summer. According to the literature, the NPF events are favored when airmasses are enriched by a reactant (e.g.,  $\text{NH}_3$ ), prior to reaching the site of Finokalia (Pikridas et al., 2012; Kalivitis et al., 2014). During Etesian flow conditions, the particle size distributions were centered on the lower end of the accumulation-mode size range (Kalivitis et al., 2014). This was partly attributed to the production of sufficient sulfuric acid to increase the condensation sink and suppress NPF events during the summer (Pikridas et al., 2012). It has been only recently shown that NPF events could occur at Finokalia during Etesians (Kalivitis et al.,

2015). A large number of PM<sub>1</sub> particles (of the order of 10<sup>4</sup> cm<sup>-3</sup>) were also observed at the northeastern AS during an Etesian outbreak (Tombrou et al., 2015), whereas high number concentrations of nucleation-mode particles observed in the north AS by Triantafyllou et al. (2016), have been associated with polluted air masses transported from Istanbul.

5 A natural question therefore, is to understand the history of the air masses as they transect the Aegean before arriving at Finokalia. In particular, we need to elucidate the atmospheric and chemical processes that affect ageing of the air masses passing over the AS maritime area between the Cyclades and Crete, and furthermore, examine whether NPF events observed at Finokalia would be stronger over the central Aegean during the northern Etesian flow. Bougiatioti et al. (2009, 2011) have observed high  
10 CCN concentrations at Finokalia, from air masses coming from the Balkans, during a period representative of an Etesian regime while Kalivitis et al., (2015) have recently demonstrated that the NPF events are associated with an increase in the concentration of cloud condensation nuclei (CCN) production in the EM atmosphere. However, few studies to date have focused on understanding the increase in cloud droplet number that results from NPF, which is the true microphysical link between  
15 NPF and the aerosol indirect effect.

Driven by the above arguments, we chose to perform measurements at a remote site on Santorini, which is located within the same path of airmasses reaching the station of Finokalia, during the Etesians. Our aim is to elucidate both atmospheric and chemical processes that affect ageing of the air masses passing over the AS before reaching its southern edge, the island of Crete. Continuous  
20 ground measurements of particle properties, concentration of gaseous species, and meteorological variables were simultaneously collected on Santorini and Crete. During this short-term campaign (15-28 July 2013) intense bursts of nucleation-mode particles were observed at both sites. The synoptic wind flow and boundary layer dynamics as well as the atmospheric chemical composition that favor the enhanced NPF events during the Etesian flow are examined in this study. To understand how NPF could  
25 affect cloud formation throughout its evolution, we quantify its impact on CCN levels, cloud droplet number concentration (CDCN) and supersaturation formed in clouds that develop before, during and after NPF events at both ground sites. Complementary to this analysis, wind patterns and atmospheric chemical composition based on WRF-Chem mesoscale model simulations, are presented.



## 2 Methodology

### 2.1 Experimental Observations

Ground level measurements were conducted simultaneously at two remote coastal areas (cf. Fig. 1), from 15 to 28 July 2013: on the island of Santorini (at Ag. Artemios; 36° 26' N, 25° 26' E) and at the  
5 monitoring station of Finokalia on the island of Crete (35° 20' N, 25° 40' E; <http://finokalia.chemistry.uoc.gr>; Mihalopoulos et al., 1997). Ag. Artemios (hereafter referred to as Santorini) is located at an elevation of 153 m above sea level (asl), while Finokalia on the top of a hill at  
10 260 m asl. Both measuring sites are far from any large city or anthropogenic activity, and are close to the sea; Finokalia is facing the sea within a sector of 270° to 90°, whereas the station on Santorini within  
10 a sector of 340° to 120°.

The Finokalia monitoring station houses a suite of instruments for measuring the meteorological parameters, the concentrations of gaseous species, as well as the physical properties and chemical composition of atmospheric particles. We used an O<sub>3</sub> analyzer (Thermo Electron model 49I), a Scanning Mobility Particle Sizer (SMPS; TROPOS Type, Wiedensohler et al., 2012) with a TSI-3772  
15 condensation particle counter (CPC) for measuring the size distribution of aerosol particles having diameters from 9 to 848 nm (scanned range), and an Aerodyne Research Inc. Aerosol Chemical Speciation monitor (ACSM; Ng et al., 2011), for measuring the mass and chemical composition ( SO<sub>4</sub><sup>2-</sup>,  
NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, Cl<sup>-</sup>, and organics) of non-refractory submicron aerosol particles. A TSI SMPS (Model 3034) measured the size distribution of particles having diameters from 10 to ca. 500 nm at Santorini.  
20 The concentrations of gaseous species were also measured using an O<sub>3</sub> analyzer (Photometric M400E), a dual channel chemiluminescence analyzer for nitrogen oxides (NO, NO<sub>2</sub>; Photometric M200E) and a fluorescence analyzer for sulfur dioxide (SO<sub>2</sub>; Photometric M100E). An overview of the instruments used for the measurements is provided in Table 1.

## 2.2 CCN and droplet number calculations

CCN concentrations are calculated using the observations of size distribution and chemical composition as follows. First, Köhler theory (Köhler, 1936; Seinfeld and Pandis, 2006; Petters and Kreidenweis, 2007) is applied to determine, based on knowledge of aerosol composition, the minimum dry size of particles,  $d_c$ , that can activate at a given level of supersaturation,  $s$ . Then, the CCN concentration is determined from the observed size distributions, by calculating the concentration of particles with sizes above  $d_c$  (Seinfeld and Pandis, 2006).  $s$  is either prescribed or determined from a cloud parameterization, both of which are used here. Chemical composition is expressed in terms of the hygroscopicity parameter,  $\kappa$ , (Petters and Kreidenweis, 2007).

Thereafter, we calculate the droplet number ( $N_d$ ) and supersaturation for clouds forming in the vicinity of both sites during all NPF events. The droplet parameterization used is based on the “population splitting concept” of Nenes and Seinfeld (2003), later improved by Barahona et al. (2010) and Morales Betancourt and Nenes (2014). In the calculations of droplet number, the size distribution is represented by the sectional approach, derived directly from the SMPS distribution files. The updraft velocity has been calculated from high-resolution airborne measurements performed over this region of AS, under similar atmospheric conditions (Tombrou et al., 2015). The partial sensitivity of cloud droplet number to chemical composition and vertical velocity is derived from the finite difference approach (Karydis et al., 2012).

## 2.3 Regional modeling

The WRF-Chem version-3.3 mesoscale model (Grell et al., 2005) is used to understand the dominant meteorological regimes and the regional characteristics of the aerosol during the sampling period. Simulations were performed by applying triple nesting: the outmost 1<sup>st</sup> domain covers the extended area of Europe (spatial resolution  $0.5^\circ \times 0.5^\circ$ ); the first level of nesting 2<sup>nd</sup> nested domain covers the extended area of Greece and Italy ( $0.167^\circ \times 0.167^\circ$ ) and the innermost nesting is centered on the extended area of Greece ( $0.056^\circ \times 0.056^\circ$ ).

The RADM2 chemical mechanism is used to simulate the gas phase chemistry (Stockwell et al., 1990). Aerosol dynamics are treated with the Modal Aerosol Dynamics Model for Europe (MADE; Ackermann et al., 1998). Aerosols in MADE are represented by two lognormal size distributions, that correspond to the Aitken and accumulation modes. Supermicrometer particles are represented by a coarse mode (Schell et al., 2001). New particle formation in MADE is treated with the Kulmala et al. (1998) parameterization of sulfuric acid nucleation although it is now well-documented that other reagents (e.g. NH<sub>3</sub> and organics) play important roles in this process (Kulmala et al., 2004). New particles are assigned to the Aitken mode with a diameter of 3.5 nm, and the size distribution parameters are adjusted to retain the lognormal shape of the distribution. Condensation rates of low-vapor-pressure gas-phase species onto existing particles, are determined by Binkowski and Shankar (1995). The Secondary Organic Aerosol (SOA) Model (SORGAM) (Schell et al., 2001) is used to simulate organic aerosol respectively. The aerosol species treated by the modules are the main inorganic ions (SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, Na<sup>+</sup>, Cl<sup>-</sup>), elemental carbon (EC), primary organic aerosols (POA), SOA, a primary unspciated PM<sub>2.5</sub> fraction covering all the unspciated/unknown fine particles (PM<sub>2.5-uns</sub>), and three species for the coarse mode (i.e., anthropogenic, marine, and soil derived aerosols). For the fine particles fraction, each model species has an Aitken-mode and an accumulation-mode. The lack of a dedicated nucleation-mode in the model neglects the actual processes of particle formation and growth towards the Aitken-mode and eventually leads to unrealistic lifetimes against deposition and coagulation as well as to unrealistic growth rates. In the framework of this study we use the results of WRF-Chem mainly to investigate the flow advection and chemical composition.

For anthropogenic emissions from Europe (1<sup>st</sup> and 2<sup>nd</sup> domains) we use the EMEP database while for Greece (3<sup>rd</sup> domain) we employ the national emission inventory (Tombrou et al., 2009). Natural (biogenic and sea-salt) emissions are calculated on-line within the WRF-Chem model. Biomass burning emissions are not considered. The chemical boundary conditions used in this study are based on an idealized, northern hemispheric, mid-latitude, clean environmental, vertical profile from the NOAA Aeronomy Lab Regional Oxidant Model (NALROM) (Liu et al., 1996; Peckham et al., 2011). Simulations were performed from 12 to 29 July. An extended evaluation of WRF-Chem model against airborne and ground observations over the AS during the Etesians is presented in Bossioli et al. (2016).

Under long-range transport conditions, the model successfully simulates CO, O<sub>3</sub>, sulfate, and ammonium concentrations while it underestimates the aerosol carbonaceous fraction, which is mostly organic matter. The biases were mainly attributed to underestimated POA emissions and limitation of the RADM2 mechanism regarding the treatment of monoterpene emissions (Tuccella et al., 2012).

5 Air mass origin and trajectories were determined by HYSPLIT4 (Hybrid Single-Particle Lagrangian Integrated Trajectory; [www.arl.noaa.gov/ready/hysplit4.html](http://www.arl.noaa.gov/ready/hysplit4.html)) back-trajectory analysis (Draxler and Rolph, 2015). The back-trajectories, initialized with meteorological conditions from GDAS (0.5°), were computed at several heights. All three-dimensional trajectories were computed with an end point either at Santorini or Finokalia station.

## 10 3 Results and Discussion

### 3.1 Prevailing atmospheric and air quality conditions

Northern winds prevailed over the AS throughout the entire campaign. Based on the simulated wind patterns at 100 m above ground level (agl) throughout Greece (cf. Fig. S1 in the supplementary material) and on the sea level pressure fields (NCEP/NCAR; Fig. S2), 17 - 18 July and 22 - 24 July are periods of strong Etesian winds (Brody and Nestor, 1985; Kotroni et al., 2001; Anagnostopoulou et al., 2014). Hereafter, we refer only to the second period, as higher aerosol number concentrations were measured at both stations, but also because there were no O<sub>3</sub> measurements at Santorini, during the first period. Immediately after the second period, another characteristic period followed (25 - 27 July), having a similar pressure pattern with the previous two; the pressure gradient over the Dardanelles was weaker. Back trajectory analysis of the air masses sampled at both stations indicates almost the same source regions, for both periods (Figs. 2, S3). However, different conditions prevailed during these two periods altering mainly the last part of the journey of the airmasses, over the AS. From 22 to 24 July, strong northern wind speeds prevailed ( $> 10 \text{ m s}^{-1}$ ) with the wind direction forming the characteristic 'ring-shape' (Fig. S1) of the Etesian flow around Turkey (Tyrlis and Lelieveld, 2013). From 25 to 27 July, moderate surface wind speeds (up to ca.  $8 \text{ m s}^{-1}$ ) with northeasterly surface flow displayed over the central and southern AS, while stagnant conditions prevailed at the north (Fig. S1, S2). Hereafter, the two periods will be referred to as Etesian Flow (EF) and Moderate Surface Flow (MSF).

The measured wind speeds at Finokalia station exceeded  $9 \text{ m s}^{-1}$ , and the wind direction was mainly from west - southwestern during the daytime hours (Fig. 3) owing to topographic features that steer the prevailing direction towards the west/southwest direction. Capturing this local feature is a known challenge for regional models (e.g., Gauss et al., 2011; Im et al., 2011; Hodnebrog et al., 2012).  
5 At the same time, the simulated wind direction at Santorini station exhibited a northern direction, with wind speeds exceeding  $8 \text{ m s}^{-1}$  during the daytime hours (Fig. 3).

The number concentrations for the three particle modes (nucleation, Aitken and accumulation) together with the  $\text{O}_3$  concentrations are shown for both periods at Santorini and Finokalia stations in Fig. 4. Simultaneous routine meteorological measurements, such as surface temperature and relative  
10 humidity, are also provided for each station. Apart from the region-wide differences, intense bursts in the concentration of nucleation-mode particles having diameters smaller than 25 nm were observed at both stations during the period of EFs (Fig. 4, shaded with yellow); it should be noted that these events were not observed at any of the stations during the period of MSF (Fig. 4, shaded with grey). In the subsequent sections the different characteristics and processes prevailing under EF or MSF are explored  
15 aiming to elucidate the interconnection between the two stations.

### 3.2 Ozone concentrations

During the EF period,  $\text{O}_3$  levels at Santorini and Finokalia stations ranged between 38 and 66 ppbv and 43 to 70 ppbv, respectively (Figs. 4, S4, Table 2); these levels are consistent with previous measurements ( $57 \pm 4$  ppbv) inside the MABL for EF carried out during the Aegean-GAME campaign.  
20 The values also agree with the climatological values recorded over the greater area during summer (Gerasopoulos et al., 2005; Kalabokas et al., 2007, 2013). During EF, the less pronounced diurnal cycle at Finokalia station (from 21 to 24 July the mean diurnal range is 8 ppbv; Fig. S4), compared to Santorini (18 ppbv; Fig. S4), is attributed to a shallower and more stable MABL over Santorini compared to Crete (Tombrou et al., 2015) that favors higher primary concentrations and thus  $\text{O}_3$   
25 scavenging at Santorini, especially when the MABL collapses during nighttime. In the vicinity of Crete, the MABL becomes less stable, due to larger sea surface temperatures (SST) existing southern of Santorini. This fact, together with the topography (i.e., Crete forms a mass of land that is located

perpendicular to the EF), enhances the mixing and the downward transport from the above rich in O<sub>3</sub> concentrations layer. During the MSF, high O<sub>3</sub> levels (the highest concentrations of the summer in 2013) were measured at both stations, ranging between 50 and 99 ppbv (Figs. 4, S4, Table 2). At both stations the highest values were observed on 26 July. The lower winds over the northern AS contributed to O<sub>3</sub> accumulation at this area, explaining the high O<sub>3</sub> concentrations at both stations. The maximum O<sub>3</sub> concentration observed (but simulated as well) at Finokalia had a 4-h delay compared to that observed at Santorini.

Simulations confirm that the air masses received at both stations during the prevailing strong northern wind are of the same origin, and representative of EF conditions (Fig. S3) albeit with an O<sub>3</sub> underprediction (average bias during afternoon hours up to -21% on 23 and -15% on 24 July, Fig. 5). During the MSF period, simulations indicate an O<sub>3</sub> increase, especially in the southern AS, but also underpredicted (average bias during afternoon hours up to -24% on 26 July, Fig. 5). In case the chemical boundary conditions, including stratosphere-troposphere exchange processes are represented realistically from a global chemical transport model, WRF-Chem simulates a significant O<sub>3</sub> increase inside the PBL (up to 40%), during Etesians (Bossioli et al. 2016). Furthermore, inaccuracies of the emissions inventory could also have impact to the results.

### 3.3 Aerosol mass and number concentrations

Figure 6 shows the non-refractory submicron aerosol concentrations measured at Finokalia during the whole experimental period. In general the inorganic and organic mass concentrations behave similar to ozone ( $R^2$  of O<sub>3</sub> to organics and inorganics is 0.5 and 0.59, respectively) during most of the experimental period (Figs S4, 6). During EF, the PM<sub>1</sub> mass concentrations were reduced roughly by a factor of two compared to those during the MSF period (Table 2), and were in the range of concentrations measured in the framework of the Aegean-GAME campaign. However, despite that the concentrations of all four species (SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup> and organics) were substantially decreased during EF (23-24 July), the organic fraction exhibited a relative increase, especially at the beginning of this period. Due to lack of data at Santorini, simulated PM<sub>2.5</sub> mass concentrations are used for the analysis. The modeled concentrations for sulfate are about 2 μg m<sup>-3</sup>, at both stations at 09:00 LST (Fig. 7) quite

close to the measured values at Finokalia (Fig. 6, on average during EF underprediction of 30% for sulfates and 60% for ammonium). Similar to the case of O<sub>3</sub>, the two stations are located along the less polluted airflow over the AS.

During the MSF period, the aerosol mass concentrations at Finokalia were substantially higher (Table 2; Fig. 6). The increased concentrations were retained until noon of 27 July for sulfate and ammonium, while those of organics continued to increase further until the end of the campaign. The modeled spatial distribution of sulfate concentrations was nearly uniform over the AS, while as for ozone their concentrations increased offshore of the northeastern coast of Crete due to the ageing of air masses in combination with the strong impact of the topography (Fig. 7). The simulated mass concentrations of secondary inorganic fine aerosols also increased (simulated and observed concentrations correlate during both periods R<sup>2</sup>=0.8); however, they are lower than the measured values at Finokalia station (average underprediction of 50% for sulfates and 75% for ammonium).

In contrast to the fine aerosol mass concentrations, their total number concentrations were substantially increased, reaching continental levels during Etesian flow conditions (from  $1.5 \times 10^3$  to  $1.5 \times 10^4$  cm<sup>-3</sup> at Santorini and from  $2.4 \times 10^3$  to  $7.5 \times 10^3$  cm<sup>-3</sup> at Finokalia; Table 2, Fig. S5). The Aitken-mode particles followed a similar diurnal variation at both stations, ranging from  $4.4 \times 10^2$  to  $7.7 \times 10^3$  cm<sup>-3</sup> and peaking around noon. Accumulation-mode particles were higher at Finokalia. The total particle number concentration measured within the MABL of eastern AS during Aegean-GAME campaign under similar atmospheric conditions were on average  $8 \times 10^3$  cm<sup>-3</sup>, with almost 20% ( $1.4 \pm 1.2 \times 10^3$  cm<sup>-3</sup>) being in the 20-50 nm size range (Tombrou et al., 2015). Greater differences were observed for the nucleation-mode particles (i.e. particles having diameters smaller than 25 nm), with sudden concentration bursts observed at both stations (Fig. 4). On 23 July (EF), a nucleation-mode burst was recorded, reaching number concentrations up to  $1.3 \times 10^4$  cm<sup>-3</sup> at Santorini and almost  $1.4 \times 10^3$  cm<sup>-3</sup> at Finokalia. A second event, but of lower intensity, was recorded on 24 July (EF). It is worth mentioning that apart from the strong winds and lower temperatures, this period is considered humid (relative humidity values reaching up to 80% at Finokalia station) in comparison to the MSF period (Fig. 4). The nucleation-mode particles shift gradually towards larger sizes in a banana-shape pattern at

both stations, as shown in Fig. 8. The number of particles remained high for several hours at Santorini (cf. Fig. 8), indicating regional NPF (Kulmala et al., 2012).

The associated growth rates ( $GR$ ) for particles that increased in size from 10 to 25 nm were estimated to be  $3.06 \text{ nm h}^{-1}$  at Santorini and  $2.05 \text{ nm h}^{-1}$  at Finokalia on 23 July, and  $2.08 \text{ nm h}^{-1}$  and  $1.76 \text{ nm h}^{-1}$ , respectively, on 24 July. The average  $GR$  for particles increasing in size from 7 to 20 nm at Finokalia was reported to be substantially higher ( $7.5 \pm 5.8 \text{ nm h}^{-1}$ ) by Pikridas et al. (2012), with the highest daily  $GR$ s observed during the hottest months of the year (May to July 2008). It should be mentioned, however, that the nucleation events reported in that study were mainly related to air masses spending most of the time over the island of Crete, which is not the case for the observations reported here. The formation rates of nucleation mode particles,  $J_D$ , have been computed according to Kulmala et al. (2012) considering both the coagulation flux and the condensational growth as sinks. For the two consequent events at Santorini,  $J_D$  for particles having diameters from 10 to 25 nm ranged between  $4.82 \text{ cm}^{-3} \text{ sec}^{-1}$  (23 July) and  $2.77 \text{ cm}^{-3} \text{ sec}^{-1}$  (24 July; Fig. 8). At the station of Finokalia,  $J_D$  was lower for particles between 9 and 25 nm, ranging between  $2.27$  (23 July) and  $2.25 \text{ cm}^{-3} \text{ sec}^{-1}$  (24 July; Fig. 8). The similarity between the  $J_D$  rates at the two sites on 24 July indicate a region-wide NPF event has occurred, yet the rates taken a day earlier are markedly different and thus, indicating a local event. However, we will show later (section 3.4) that this is not the case and more information needs to be taken into account.

Under MSF conditions, the total fine aerosol number concentrations were considerably lower than those during the EFs (from  $1.4 \times 10^3$  to  $2.9 \times 10^3 \text{ cm}^{-3}$  at Santorini and from  $2.6 \times 10^3$  to  $5.1 \times 10^3 \text{ cm}^{-3}$  at Finokalia; Fig. S5). Particles in the nucleation mode were absent, while the concentrations in the Aitken mode were substantially lower at both stations, varying from  $3.2 \times 10^2$  to  $4.1 \times 10^3 \text{ cm}^{-3}$  (Fig. S5). The particle concentrations in the accumulation mode at Santorini had a comparable variation to that of the Aitken-mode, while they were apparently always higher at Finokalia.

### 25 3.4 Spatial extent of NPF event

The synoptic wind flow and boundary layer dynamics as well as the chemical atmospheric background conditions that favor the enhanced NPF events during the EF are further examined here. This type of



event could be characterized as “type A” according to Boy and Kulmala (2002), owing to the sudden appearance of nucleation-mode particles and their consistent growth for at least 1 hour. The horizontal scale of this event was estimated based on air mass back-trajectory analysis (Hussein et al., 2009), taking into account the time during which measurements at the site indicate a distinct nucleation mode.

5 Following Birmili et al. (2003), HYSPLIT4 back-trajectory calculations started at the time when a nucleation mode was first distinguishable from the Aitken mode at Santorini and were performed for each subsequent hour until the two modes merged (nucleation duration). Following Crippa and Pryor (2013), the duration of NPF was based on the geometric mean diameter of particles with sizes between 10 and 100 nm and from 30 to 100 nm; an event is said to initiate when the difference between the two

10 geometric mean diameters becomes maximum and ends when this difference is less than 15% (Fig. S6). Assuming a linear *GR* (Lehtinen and Kulmala, 2003), this approach showed that the ca. 10-nm particles (the smallest particles we could detect with our instrumentation) were able to grow up to 60 nm within 4.5 h of initial detection. This *GR* was then used to calculate the minimum spatial scale. On 23 July, the distance covered by the back-trajectories within 4.5 h (starting when the nucleation-mode burst was first

15 recorded at Santorini) spans at least over 250 km to the northeast of Santorini in the center of AS, upwind of the Cyclades complex (Fig. 2; red line in Fig. 1, left panel). A couple of hours before the sunrise these back-trajectories (both below and above 500 m agl) are observed over the northwestern Asian forest peninsula of Turkey (area marked as black ellipse in Fig. 1, left panel; Fig. 2), having previously passed (at higher altitudes > 1.5 km agl) from the Greater Area of Istanbul (GAI) and the

20 west coast of the Black Sea (from even higher altitudes > 3 km agl). A similar spatial extent also occurs during the less intense EF event on 24 July, although this starts with two hours delay (Fig. 2). Air masses are better mixed throughout the boundary layer covering a broader area over Asian Turkey on 24 July.

25 Despite the limitations of the model (absence of a nucleation mode, binary homogeneous nucleation parameterization only included), the conditions under which NPF events take place and their overall impact can still be estimated by conducting another simulation that deactivates the nucleation parameterization (nucleation-off run). According to the simulations, a wide stream of clean air masses of low preexisting aerosol particles (number concentrations <  $2.5 \times 10^3$  particles  $\text{cm}^{-3}$ , not shown) but of

sufficient  $\text{H}_2\text{SO}_4$  ( $\sim 10^7$  molecules  $\text{cm}^{-3}$  from high altitudes, not shown), overpass the urban mixing height (at 1-2 km) over the GAI during the previous evening on 22 July (20:00 LST, Fig. 1, left panel), avoiding mixing with the local emissions. Thereafter, they penetrate at lower levels (due to the EF structure) over northwestern Turkey (Fig. 2, left panel). At this forested area (black ellipse in Fig. 1, left panel) they find favorable conditions for NPF, such as low relative humidity,  $\text{H}_2\text{SO}_4$  and availability of biogenic emissions (not shown) that endorse further the NPF efficiency. In Figure S7, the number concentration of new particles (nucleation-on – nucleation off), at 1 km at 6:00 LST, for both EF (left panel) and MSF (right panel) periods, is presented. Although severely underestimated (simulated NPF contribution up to  $\sim 400$   $\text{cm}^{-3}$ ), the critical role of the EF on the NPF event over northern AS is revealed.

10 Closer to the surface, the air masses have a substantial number of primary particles (emitted by the various activities of the GAI, Fig. S8), providing more surface available for condensation (NPF contribution up to 200  $\text{cm}^{-3}$ , not shown). According to the simulations, a plume with large particulate load in the Aitken mode ( $\sim 9 \times 10^3$  particles  $\text{cm}^{-3}$ ) is transported over northwestern Turkey (Fig. S8, left panel). Our results agree with previous observations during an Etesian event, where number concentrations up to  $\sim 1.2 \times 10^4$   $\text{cm}^{-3}$  were observed at the northeastern AS with the Aitken-mode particles dominating by up to 70% (Tombrou et al., 2015). The less intense event on 24 July is associated with a narrow stream of low preexisting particles over the GAI (concentrations  $< 2.5 \times 10^3$  particles  $\text{cm}^{-3}$ , not shown).

20 The plume, after crossing the Turkish mainland overnight, is transported over the AS, with most of the new particles above the stable MABL (Fig. S7, top-left panel; dashed purple lines in Fig. 1, left panel). The plume is moving fast with rather negligible mixing, especially above the MABL, thereby affecting areas located further away, such as the central AS, within a couple of hours after sunrise on 23 July (around 9:00 LST) and around noon on 24 July. The rapid advection above MABL, in combination with the low number of pre-existing particles there (Fig. S7, bottom-left panel), seems to leave almost intact the majority of the newly formed particles. Thereafter, we mainly consider that while the part of the plume above the MABL passes over the Cyclades complex, the wakes on the lee side of the islands enhance vertical mixing, enabling its entrainment into the MABL (area indicated with a white dashed line in Fig. 1, left panel). This assumption does not reject the fact that oxidation enhanced by

photochemistry over the AS may also contribute to NPF process. The freshly nucleated particles that remained constantly inside the well-mixed MABL, suffered an early ageing (i.e. growth by condensation and coagulation). The concentrations at both nucleation and Aitken modes jump almost simultaneously accompanied by concurrent increases in O<sub>3</sub>, NO<sub>2</sub> and SO<sub>2</sub> concentrations, at Santorini station, during these two consequent events (Fig. 4). This could be an indication that this station receives masses simultaneously from different layers (inside and above the MABL), in line with a number of cases where maximum rate of change of ultra-fine particle concentrations close to the surface was always preceded by breakdown of the nocturnal inversion and enhancement of vertical mixing (Crippa et al., 2012).

The air masses arrive 3-h later (after 13:00 LST; Fig. S3 left panel) at Finokalia, on 23 July (Fig. 4). The 3-h transit timescale is in agreement with the prevailing wind speed (about 10 m s<sup>-1</sup>; Fig. S1) and the 120 km distance between Santorini and Finokalia. The nucleation-mode particles are significantly reduced as they have shifted gradually towards larger sizes (Aitken-mode), before reaching Finokalia (Fig. 4). The measured nucleated concentrations measured previously at Finokalia are probably due to a local nucleation event initiated at Heraklion (Crete). The current simultaneous measurements along the same flow stream, show that both stations are under the influence of regional NPF events, during the Etesians.

During the MSF period, on 26 July, the air masses arriving at lower levels (below 500 m agl) at Santorini station (Fig. 1, 2, both right panel) have mainly passed from low altitudes over continental areas (< 1 km), and have been substantially enriched by anthropogenic emissions, while those at higher levels have covered longer distances over Eastern Europe at the same time (exact opposite behavior of EF). Over the GAI, the simulated particle number concentration is much higher compared to EF conditions ( $5-7 \times 10^3$  particles cm<sup>-3</sup>, not shown) limiting the NPF event (Fig. S7). These atmospheric conditions promote the mixing of air masses with local anthropogenic and natural emissions favoring photochemical production of secondary pollutants such as O<sub>3</sub> (Fig. 5, right panel) and higher secondary aerosols (e.g. SO<sub>4</sub> shown in Fig. 7).

### 3.5 Impact of NPF events to CCN production

Understanding how NPF affects cloud formation requires quantification of its impact on the CCN levels that develop for cloud-relevant supersaturations. As CCN concentrations were not measured, they have been calculated using the observations of size distribution and chemical composition as already described in section 2.2. The presentation of the results and the relevant discussion are based on the periods before and after the NPF events.

CCN concentrations are calculated for prescribed values of  $s$  between 0.2 and 0.8%, corresponding to supersaturations found in relatively pristine stratiform to convective clouds (Seinfeld and Pandis, 2006).  $\kappa$  is calculated from the  $PM_{10}$  chemical composition observed at Finokalia as follows:  $\kappa = \varepsilon_{inorg} \kappa_{inorg} + \varepsilon_{org} \kappa_{org}$ , where  $\kappa_{inorg} = 0.6$  is the value for ammonium sulfate (Petters and Kreidenweis, 2007), and  $\kappa_{org} = 0.16$  corresponds to the organic fraction (Bougiatioti et al., 2009), and  $\varepsilon_{inorg}$ ,  $\varepsilon_{org}$  are the volume fractions of each constituent measured at Finokalia. The volume fractions range from 0.45 to 0.76 for inorganics and from 0.24 to 0.55 for organics, similar to the values measured under comparable atmospheric conditions by Bougiatioti et al. (2009, 2011) and Bezantakos et al. (2013). Throughout the measurement period, the aerosol exhibited predicted values of hygroscopicity from 0.20 to 0.39, which is also consistent with the values determined by Bougiatioti et al. (2009, 2011) and Bezantakos et al. (2013). The aerosol hygroscopicity follows a diurnal cycle being minimum just before noon and becoming maximum late in the afternoon, owing to a higher sulfate-to-organic mass ratios (Fig. 6). Consequently, average  $\kappa$  values were estimated to be higher after the NPF events compared to the period before (increase by ~35% on 23 July and up to 15% on 24 July). Given a lack of  $PM_{10}$  chemical composition measurements at Santorini, the chemical composition at Finokalia is applied instead to the Santorini size-distribution observations. The WRF simulations support this assumption, as a similar chemical behavior is simulated for both stations (Figs. 5,7). The model systematically underestimates the organic fraction at both stations (organic volume fraction does not exceed 0.2), but with minimal impact on resulting  $\kappa$  values, as they do not differ from measurements for more than  $\pm 6\%$  throughout the simulation period. From long-term measurements in the study area, the relative contribution of the main  $PM_{10}$  constituents, including ammonium, is quite consistent over the years (Sciare et al., 2003; Koulouri et

al., 2008, Bougiatioti et al., 2013). Thus, a sensitivity test of CCN concentration at Santorini to shifts in  $\kappa$  by  $\pm 20\%$ , is also carried out.

The resulting CCN timeseries during Etesian flow are shown in Figure 9. Average values of  $\kappa$ ,  $d_c$ , and CCN concentrations at both stations, before and after the NPF events are provided in Table 3. The  
5 calculated CCN number concentrations follow a diurnal cycle and tend to be maximum during the afternoon, after the NPF events, following the increase of  $\kappa$  values. Most particles are CCN-active for  $s \geq 0.6\%$ , as they converge towards the total CN timeseries. Bougiatioti et al. (2009) observed similar behavior at Finokalia for polluted air masses with a similar origin (Balkans). For  $s = 0.6\%$ ,  $d_c$  varies from  
10 43 to 51 nm and CCN concentrations reach up to  $\sim 6 \times 10^3 \text{ cm}^{-3}$  following the Aitken-mode concentrations at both stations (Figs. 4, 9). The higher CCN number concentrations at Finokalia, compared to those observed at Santorini (Table 3), is a result of higher number of accumulation-mode particles passing previously from Santorini (that are too small to be CCN at Santorini, but have grown to CCN-relevant sizes by the time they arrive at Finokalia, section 3.3). Accordingly, the higher activation fractions (CCN/CN) are observed at the station of Finokalia, with larger and more aged aerosol particles, while at  
15 Santorini this is observed at the end of the events, when the smaller particles drop in concentration because they grow to larger sizes. On 23 July, the NPF event increases the CCN concentrations by 157% at Santorini and 106% at Finokalia, compared to their pre-event values; while in some moments the increase can reach up to a factor of 6. During the second less intense event, on 24 July, the CCN increase is lower at both stations (31% at Santorini and 53% at Finokalia). The lower increase is also due to the  
20 pre-event background, characterized by higher CCN concentrations. Throughout the MSF period, the CCN concentrations decrease by almost 48% and 23% at Santorini and Finokalia respectively, compared to the levels during the NPF events. Changes in chemical composition, as described above exhibit a relative low variation in CCN concentrations (at  $s = 0.6\%$ ) up to 10%, following the same diurnal behavior. As expected, lowering the supersaturation at 0.2% leads to the activation of larger particles  
25 with  $d_c$  ranging from 91 to 106 nm that is consistent with the observations reported by Kalivitis et al. (2015). At  $s = 0.2\%$ , both NPF events contribute up to 50% to the increase of the CCN concentrations at both stations. However, the higher CCN production at Finokalia on 24 July is associated with the accumulation-mode particles at the end of both events.

### 3.6 Impact of NPF events on droplet number

Studying the impact of NPF on CCN concentrations at prescribed levels of supersaturation is a simple and frequently used approach for observational studies of NPF (e.g., Kalivitis et al., 2015 and references therein). It however provides an incomplete description of NPF impacts on cloud droplet number, as it does not consider the feedback of CCN on cloud supersaturation that develops in cloudy updrafts. Mechanistic cloud droplet formation parameterizations (Ghan et al., 2011; Morales Betancourt and Nenes, 2014) can capture this complexity by efficiently calculating the maximum supersaturation ( $s_{max}$ ) that forms in a cloud given knowledge of the aerosol size distribution, composition and updraft velocity. Observations suggest that the distribution of vertical velocities in the boundary layer display a spectral dispersion of  $\sigma_w = 0.2-0.3 \text{ m s}^{-1}$  around a zero average value, which is consistent with vertical velocities observed in marine boundary layers (e.g., Meskhidze et al., 2005; Ghan et al., 2011). When applying the droplet parameterization, we employ the “characteristic velocity” approach of Morales and Nenes (2010) to obtain velocity PDF-averaged values of cloud droplet number concentration (CDNC) and  $s_{max}$ . As a sensitivity test, we also consider calculations for a convective boundary layer ( $\sigma_w = 0.6 \text{ m s}^{-1}$ ).

The calculation of PDF-averaged values of  $N_d$  and  $s_{max}$  is carried out for every distribution of aerosol number and composition measured for all NPF events. The resulting timeseries are shown in Figure 10 for Santorini (top panel) and Finokalia (bottom panel).  $s_{max}$  is negatively correlated with  $N_d$  at both stations, owing to the increased competition for water vapor by the growing droplets when CCN increase. As a result,  $N_d$  responds sublinearly to CCN increases – the degree to which this depends on the level of aerosol concentrations before and during the NPF event. At Santorini, the CCN levels are much lower than at Finokalia (Table 3), so we expect the relative increase in  $N_d$  from NPF to be higher there. Assuming  $\sigma_w = 0.3 \text{ m s}^{-1}$ , the NPF events are associated with  $s_{max}$  decreases at both stations, compared to the period before the events. On 23 July, the decrease is on average 12% at Santorini and 9% at Finokalia. As a result,  $N_d$  concentrations during the NPF event increases by 13% to  $124 \pm 8 \text{ cm}^{-3}$  at Santorini, compared to the period before the event. At Finokalia, however, aerosol levels are much higher and  $N_d$  remains virtually the same before and after the NPF event (Table 3). The effect of the less intense

NPF event on 24 July is higher;  $N_d$  concentration increases by 36% at Santorini and 4% at Finokalia compared to pre-event values. The decrease of  $s_{max}$  is also higher on this day, 17% at Santorini and even higher 36.4% (at 0.06-0.08%) at Finokalia (Table 3) owing to the higher accumulation particle concentrations compared to the previous events. The variance of  $N_d$  during the event period, for  $\sigma_w$  equal to 0.3 m s<sup>-1</sup>, is 475 cm<sup>-3</sup> at Santorini and 37 cm<sup>-3</sup> at Finokalia, while for  $\sigma_w$  equal to 0.6 m s<sup>-1</sup> the variance is 865 cm<sup>-3</sup> and 20 cm<sup>-3</sup>, respectively. Altogether, this clearly shows that when NPF particles age (e.g., arrive at Finokalia) their competition for water vapor can reduce cloud supersaturation to very low levels.

The larger updraft velocity ends in larger values of  $s_{max}$ , which allow smaller particles to activate into cloud droplets. In particular,  $N_d$  exhibits a substantial increase for  $\sigma_w = 0.6$  m s<sup>-1</sup>, but with a similar pattern to that with the lower velocity, especially at Santorini indicating that the impact of mean vertical velocity on the CDNC is higher at this station. In this case, the average  $N_d$  concentration is 217±15 cm<sup>-3</sup> at Santorini and 619±109 cm<sup>-3</sup> at Finokalia (increase relative to  $\sigma_w = 0.3$  m s<sup>-1</sup> by 75% and 52%), after the event on 23 July and 286±15 and 786±11, respectively (increase relative to  $\sigma_w = 0.3$  m s<sup>-1</sup> by ~76%, for both stations), on 24 July. It is interesting to note that for  $\sigma_w = 0.6$  m s<sup>-1</sup> two  $N_d$  peaks are observed at Finokalia, from which the first is attributed to local processes as it is observed much earlier than the NPF event at Santorini. The stronger variation of  $N_d$  at Finokalia, under the higher vertical wind, compared to Santorini, indicates that vertical velocity variations likely dominate the variance of droplet number for clouds in the region of Finokalia. Furthermore, from the partial sensitivity of  $N_d$  to the total aerosol number, and to  $\kappa$ , the relative contribution of chemical composition and total aerosol number to the variance of  $N_d$  is attributed. We find that in most cases the predicted  $N_d$  variability is almost exclusively governed by the aerosol number variation (> 98%, Table 3) and to a lesser extent by the chemical composition (< 2%). The relative contribution of chemical composition becomes more significant at Finokalia only after the intense NPF event on 23 July (10% for  $\sigma_w = 0.3$  m s<sup>-1</sup> and 19% for  $\sigma_w = 0.6$  m s<sup>-1</sup>). This can be attributed to the more "aged" nature of the sampled aerosol at Finokalia, compared to the one at Santorini. This is consistent with the lower  $s_{max}$  predicted for Finokalia, leading to the activation of larger particle sizes that have been subject to longer atmospheric processing, during their transition to more unstable conditions after Santorini. Altogether, although NPF events may strongly elevate CCN

numbers, the relative impacts on cloud droplet number (compared to pre-event levels) is eventually limited by water vapor availability and depends on the aerosol levels associated with the background.

#### 4 Conclusions

Concentrations of chemically- and size-resolved submicron aerosol particles along with concentrations of trace gases and meteorological variables have been simultaneously measured at Santorini (central AS) and Finokalia on Crete (southern AS) from 15 to 28 July 2013. Two well-distinguished periods are identified: the first with strong wind speeds and wind directions forming the characteristic 'ring-shape' of the Etesian flow (EF) around Turkey, and the second with moderate surface wind speeds and northerly direction over the AS (MSF). The two periods exhibited intense differences on air quality levels.

10 During EF, the mass concentrations were reduced roughly by a factor of two compared to those during the MSF period. The total number concentration of aerosol particles was increased during the EF, varying from  $1.5 \times 10^3$  to  $1.5 \times 10^4$  particles  $\text{cm}^{-3}$  at Santorini and from  $2.4 \times 10^3$  to  $7.5 \times 10^3$  particles  $\text{cm}^{-3}$  at Finokalia. Furthermore, intense burst of nucleation-mode particles have been recorded at both stations, with more intense those observed at Santorini. At Finokalia, the fragment of nucleated particles is diminished, and a higher number concentration of the Aitken-mode particles is observed, attributed to atmospheric mixing, growth process and photochemistry. The nucleation-mode particles are gradually shifting towards larger sizes at both stations, however, at Santorini the number of particles remains high for several hours, indicating regional NPF. During the MSF period, the total number concentration of the particles reaches lower values, while nucleation-mode particles are not detected at any of the stations.

20 The observed NPF events have been initiated at least 250 km (covered within 4.5 hours) to the northeast of Santorini in the center of AS, upwind of Cyclades complex, under favorable meteorological conditions, under a strong-channeled northeastern wind flow received by both stations. Based on the simulation, it seems that what contributes to the NPF events is the clean air masses of low preexisting aerosol particles but of sufficient  $\text{H}_2\text{SO}_4$ , from high altitudes. In contrast to the non-NPF period, the air masses overpass the greater Istanbul area, avoiding mixing with the local emissions. Thereafter, they penetrate at lower levels (due to the EF structure) over northwestern Turkey, while in case of non- NPF they suffer a strong mixing during their longer journey over the Turkey mainland. Without excluding the



role of photochemistry in NPF, we have shown by both measurements and simulations that the plume over AS is moving fast with rather negligible mixing, especially above the MABL. The fast advection above MABL as well as the low number of pre-existing concentrations inside this plume, prevent the subsequent growth of the nucleated particles towards central Aegean. The wakes on the lee side of the islands, however, enhance vertical mixing, enabling its subsequent entrainment into the MABL, in the central Aegean. The freshly nucleated particles that remained constantly inside the well-mixed MABL, suffered an early ageing (i.e. growth by condensation and coagulation).

To understand the impact of NPF on CCN levels, using the  $\kappa$  of particles and in conjunction with a typical supersaturation for the area, we calculated the number concentration of particles which act as CCN at both stations. NPF was found to augment CCN concentrations considerably during early afternoon (87% on average for both stations and both events), with concentration levels at Finokalia being higher due to particle growth and atmospheric processing. Calculations of droplet number generated in clouds within the observed airmasses indicate that NPF augments droplet number, but to a much lesser extent (12%) than implied by the variations in CCN. This behavior demonstrates there is a limit in the amount of droplets that NPF can contribute because the supersaturation in cloud depresses (here, by roughly 14%) as additional CCN are added from NPF. The pre-NPF aerosol levels and prevailing dynamics of the clouds determine the degree of water vapor competition and precondition clouds to be sensitive - or not - to further CCN increases from NPF.

## 20 Acknowledgments

AN acknowledges support from a Georgia Power Faculty Scholar Chair and a Cullen-Peck Faculty Fellowship. This work was supported by computational time granted from the Greek Research & Technology Network (GRNET) in the National HPC facility - ARIS

## References

- Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S., and Shankar, U.: Modal Aerosol Dynamics model for Europe: Development and first applications, *Atmos. Environ.*, 32, 2981–2999, 1998.
- 5 Anagnostopoulou, C., Zanis, P., Katragkou, E., Tegoulas, I., and Tolika, K.: Recent past and future patterns of the Etesian winds based on regional scale climate model simulations, *Clim. Dyn.*, 42, 1819-1836, doi:10.1007/s00382-013-1936-0, 2014.
- Barahona, D., West, R. E. L., Stier, P., Romakkaniemi, S., Kokkola, H., and Nenes, A.: Comprehensively Accounting for the Effect of Giant CCN in Cloud Activation Parameterizations, *Atmos.Chem.Phys.*, 2467-2473, doi:10.5194/acp-10-2467-2010, 2010.
- 10 Bardouki, H., Liakakou, H., Economou, C., Sciare, J., Smolík, J., Ždímal, V., Eleftheriadis, K., Lazaridis, M., Dye, C. and Mihalopoulos, N.: Chemical composition of size-resolved atmospheric aerosols in the eastern Mediterranean during summer and winter, *Atmos. Environ.*, 37, 195-208, 2003.
- 15 Bezantakos, S., Barmounis, K., Giamarelou, M., Bossioli, E., Tombrou, M., Mihalopoulos, N., Eleftheriadis, K., Kalogiros, J., Allan, J. D., Bacak, A., Percival, C. J., Coe, H., and Biskos, G.: Chemical composition and hygroscopic properties of aerosol particles over the Aegean Sea, *Atmos. Chem. Phys.*, 13, 11595–11608, doi:10.5194/acp-13-11595-2013, 2013.
- Binkowski, F. S., and Shankar, U.: The Regional Particulate Matter Model: 1. Model description and preliminary results, *J. Geophys. Res.*, 100(D12), 26191–26209, doi:10.1029/95JD02093, 1995.
- 20 Birmili, W., Berresheim, H., Plass-Dülmer, C., Elste, T., Gilge, S., Wiedensohler, A. and Uhrner, U.: The Hohenpeissenberg aerosol formation experiment (HAFEX): A long-term study including size-resolved aerosol, H<sub>2</sub>SO<sub>4</sub>, OH, and monoterpenes measurements, *Atmos. Chem. Phys.*, 3, 361-376, doi:10.5194/acp-3-361-2003, 2003.
- 25 Bossioli, E., Tombrou, M., Kalogiros, J., Allan, J., Bacak, A., Bezantakos, S., Biskos, G., Coe, H., Jones, B. T., Kouvarakis, G., Mihalopoulos, N., and Percival, C. J.: Atmospheric composition in the Eastern Mediterranean: Influence of biomass burning during summertime using the WRF-Chem model, *Atmos. Environ.*, 132, 317–331, 2016.

- Bougiatioti, A., Fountoukis, C., Kalivitis, N., Pandis, S. N., Nenes, A., and Mihalopoulos, N.: Cloud condensation nuclei measurements in the marine boundary layer of the Eastern Mediterranean: CCN closure and droplet growth kinetics, *Atmos. Chem. Phys.*, 9, 7053–7066, doi:10.5194/acp-9-7053-2009, 2009.
- 5 Bougiatioti, A., Nenes, A., Fountoukis, C., Kalivitis, N., Pandis, S. N., and Mihalopoulos, N.: Size-resolved CCN distributions and activation kinetics of aged continental and marine aerosol, *Atmos. Chem. Phys.*, 11, 8791–8808, doi:10.5194/acp-11-8791-2011, 2011.
- Bougiatioti, A., Zarnpas, P., Koulouri, E., Antoniou, M., Theodosi, C., Kouvarakis, G., Saarikoski, S., Mäkelä, T., Hillamo, R., and Mihalopoulos, N.: Organic, elemental and water-soluble organic carbon  
10 in size segregated aerosols, in the marine boundary layer of the Eastern Mediterranean, *Atmos. Environ.*, 64, 251-262, 2013.
- Bougiatioti, A., Stavroulas, I., Kostenidou, E., Zarnpas, P., Theodosi, C., Kouvarakis, G., Canonaco, F., Prévôt, A. S. H., Nenes, A., Pandis, S. N., and Mihalopoulos, N.: Processing of biomass-burning aerosol in the eastern Mediterranean during summertime, *Atmos. Chem. Phys.*, 14, 4793–4807,  
15 doi:10.5194/acp-14-4793-2014, 2014.
- Boy, M. and Kulmala, M.: Nucleation events in the continental boundary layer: Influence of physical and meteorological parameters, *Atmos. Chem. Phys.*, 2, 1-16, doi:10.5194/acp-2-1-2002, 2002.
- Brody, L. R. and Nestor, M. J. R.: Regional forecasts for the Mediterranean basin, Technical Report, No. 80-110, Naval Environmental Prediction Research Facility, Monterey, California, USA, 1985.
- 20 Crippa, P., Petäjä, T., Korhonen, H., El Afandi, G. S., and Pryor, S. C.: Evidence of an elevated source of nucleation based on model simulations and data from the NIFTy experiment, *Atmos. Chem. Phys.*, 12, 8021-8036, doi:10.5194/acp-12-8021-2012, 2012.
- Crippa, P. and Pryor, S. C.: Spatial and temporal scales of new particle formation events in eastern North America, *Atmos. Environ.*, 75, 257–264, doi:10.1016/j.atmosenv. 2013.04.051, 2013.
- 25 Draxler, R.R. and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) Model access via NOAA ARL READY Website (<http://ready.arl.noaa.gov/HYSPLIT.php>). NOAA Air Resources Laboratory, Silver Spring, MD., 2015.

- Eleftheriadis, K., Colbeck, I., Housiadas, C., Lazaridis, M., Mihalopoulos, N., Mitsakou, C., Smolík, J. and Ždímal, V.: Size distribution, composition and origin of the submicron aerosol in the marine boundary layer during the eastern Mediterranean “SUB-AERO” experiment, *Atmos. Environ.*, 40, 6245–6260, doi:10.1016/j.atmosenv.2006.03.059, 2006.
- 5 Formenti, P., Reiner, T., Sprung, D., Andreae, M. O., Wendisch, M., Wex, H., Kindred, D., Dewey, K., Kent, J., Tzortziou, M., Vasaras, A. and Zerefos, C.: STAAARTE-MED 1998 summer airborne measurements over the Aegean Sea 1. Aerosol particles and trace gases, *J. Geophys. Res.*, 107(D21), 4450, doi:10.1029/2001JD001337, 2002a.
- 10 Formenti, P., Boucher, O., Reiner, T., Sprung, D., Andreae, M. O., Wendisch, M., Wex, H., Kindred, D., Tzortziou, M., Vasaras, A. and Zerefos, C.: STAAARTE-MED 1998 summer airborne measurements over the Aegean Sea 2. Aerosol scattering and absorption, and radiative calculations, *J. Geophys. Res.*, 107(D21), 4451, doi:10.1029/2001JD001536, 2002b.
- 15 Gauss, M., Benedictow, A. C., Fagerli, H. and B.M. Steensen, EMEP/MSC-W model performance for acidifying and eutrophying components and photo-oxidants in 2009, Supplementary material to EMEP Status Report 1/2011, METEOROLOGISK INSTITUTT, Norwegian Meteorological Institute, EMEP July 2011.
- Gerasopoulos, E., Kouvarakis, G., Vrekoussis, M., Kanakidou, M. and Mihalopoulos, N.: Ozone variability in the marine boundary layer of the eastern Mediterranean based on 7-year observations, *J. Geophys. Res. D: Atmospheres*, 110, D15309, doi:10.1029/2005JD005991, 2005.
- 20 Gerasopoulos, E., Koulouri, E., Kalivitis, N., Kouvarakis, G., Saarikoski, S., Mäkelä, T., Hillamo, R. and Mihalopoulos, N.: Size-segregated mass distributions of aerosols over Eastern Mediterranean: seasonal variability and comparison with AERONET columnar size-distributions, *Atmos. Chem. Phys.*, 7, 2551-2561, doi:10.5194/acp-7-2551-2007, 2007.
- 25 Ghan, S. J., Abdul-Razzak, H., Nenes, A., Ming, Y., Liu, X., Ovchinnikov, M., Shipway, B., Meskhidze, N., Xu, J., and Shi, X.: Droplet Nucleation: Physically-based Parameterization and Comparative Evaluation, *J. Adv. Model. Earth Syst.*, 3, M10001, 33 pp. doi:10.1029/2011MS000074, 2011.

- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock W. C. and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39, 6957–6975, 2005.
- Hodnebrog, Ø., Solberg, S., Stordal, F., Svendby, T. M., Simpson, D., Gauss, M., Hilboll, A., Pfister, G. G., Turquety, S., Richter, A., Burrows, J. P. and Denier Van Der Gon, H. A. C.: Impact of forest fires, biogenic emissions and high temperatures on the elevated Eastern Mediterranean ozone levels during the hot summer of 2007, *Atmos. Chem. Phys.*, 12, 8727-8750, doi:10.5194/acp-12-8727-2012, 2012.
- Hussein, T., Junninen, H., Tunved, P., Kristensson, A., Dal Maso, M., Riipinen, I., Aalto, P. P., Hansson, H.-C., Swietlicki, E. and Kulmala, M.: Time span and spatial scale of regional new particle formation events over Finland and Southern Sweden, *Atmos. Chem. Phys.*, 9, 4699-4716, doi:10.5194/acp-9-4699-2009, 2009.
- Im, U., Markakis, K., Poupkou, A., Melas, D., Unal, A., Gerasopoulos, E., Daskalakis, N., Kindap, T. and Kanakidou, M.: The impact of temperature changes on summer time ozone and its precursors in the Eastern Mediterranean, *Atmos. Chem. Phys.*, 11, 3847-3864, doi:10.5194/acp-11-3847-2011, 2011.
- Im, U., and Kanakidou, M. Impacts of East Mediterranean megacity emissions on air quality. *Atmos. Chem. Phys.*, 12, 6335-6355, doi:10.5194/acp-12-6335-2012, 2012.
- Kalabokas, P. D., Volz-Thomas, A., Brioude, J., Thouret, V., Cammas, J.-P. and Repapis, C. C.: Vertical ozone measurements in the troposphere over the Eastern Mediterranean and comparison with Central Europe, *Atmos. Chem. Phys.*, 7, 3783–3790, doi:10.5194/acp-7-3783-2007, 2007.
- Kalabokas, P. D., Mihalopoulos, N., Ellul, R., Kleanthous, S. and Repapis, C. C.: An investigation of the meteorological and photochemical factors influencing the background rural and marine surface ozone levels in the Central and Eastern Mediterranean, *Atmos. Environ.*, 42, 7894–7906, doi:10.1016/j.atmosenv.2008.07.009, 2008.
- Kalabokas, P. D., Cammas, J.-P., Thouret, V., Volz-Thomas, A., Boulanger, D. and Repapis, C. C.: Examination of the atmospheric conditions associated with high and low summer ozone levels in the lower troposphere over the Eastern Mediterranean, *Atmos. Chem. Phys. Discuss.*, 13, 2457–2491, doi:10.5194/acpd-13-2457-2013, 2013.

- Kalivitis, N., Birmili, W., Stock, M., Wehner, B., Massling, A., Wiedensohler, A., Gerasopoulos, E. and Mihalopoulos, N.: Particle size distributions in the Eastern Mediterranean troposphere, *Atmos. Chem. Phys.*, 8, 6729–6738, doi:10.5194/acp-8-6729-2008, 2008.
- Kalivitis, N., Kouvarakis G., Bougiatioti A., Stavroulas I., Wiedensohler A., and Mihalopoulos N.:  
5 Five-years of atmospheric aerosol number size distribution measurements in Eastern Mediterranean, *Geophys. Res. Abstracts*, 16, EGU2014-15860-1, 2014.
- Kalivitis, N., Kerminen, V.-M., Kouvarakis, G., Stavroulas, I., Bougiatioti, A., Nenes, A., Manninen, H. E., Petäjä, T., Kulmala, M., and Mihalopoulos, N.: Atmospheric new particle formation as source of CCN in the Eastern Mediterranean marine boundary layer, *Atmos. Chem. Phys.*, 15, 9203-9215,  
10 doi:10.5194/acp-15-9203-2015, 2015.
- Kallos, G., Kotroni, V., Lagouvardos, K., Papadopoulos, A., Varinou, M., and Kakaliagou, O.: Temporal and spatial scales for transport and transformation processes in the Eastern Mediterranean, *Proc. Of the 22nd NATO/CCMS Int. Tchn. Meeting on Air pollution Modelling and its Application*, Clermont-Ferrand, France, 6-11 June; 1998.
- 15 Kallos, G., Astitha, M., Katsafados, P., and Spyrou, C.: Long-Range Transport of Anthropogenically and Naturally Produced Particulate Matter in the Mediterranean and North Atlantic: Current State of Knowledge, *J. Appl. Meteorol. Climatol.*, doi: 10.1175/JAM2530.1, 1230–1251, 2007.
- Kanakidou, M., Mihalopoulos, N., Kindap, T., Im, U., Vrekoussis, M., Gerasopoulos, E., Dermitzaki, E., Unal, A., Koçak, M., Markakis, K., Melas, D., Kouvarakis, G., Youssef, A. F., Richter, A.,  
20 Hatzianastassiou, N., Hilboll, A., Ebojie, F., Wittrock, F., von Savigny, C., Burrows, J. P., Ladstaetter-Weissenmayer, A. and Moubasher, H.: Megacities as hot spots of air pollution in the East Mediterranean, *Atmos. Environ.*, 45, 1223–1235, doi:10.1016/j.atmosenv.2010.11.048, 2011.
- Karydis, V. A., Capps, S. L., Russell, A. G., and Nenes, A.: Adjoint sensitivity of global cloud droplet number to aerosol and dynamical parameters, *Atmos. Chem. Phys.*, 12, 9041-9055, doi:10.5194/acp-  
25 12-9041-2012, 2012
- Koçak, M., Theodosi, C., Zarnpas, P., Im, U., Bougiatioti, A., Yenigun, O., and Mihalopoulos, N.: Particulate matter (PM<sub>10</sub>) in Istanbul: Origin, source areas and potential impact on surrounding regions, *Atmos. Environ.*, 45, 6891-6900, 2011.

- Köhler, H.: The nucleus in the growth of hygroscopic droplets. *Trans. Faraday Soc.* 32, 1152, 1936.
- Kopanakis, I., Eleftheriadis, K., Mihalopoulos, N., Lydakis-Simantiris, N., Katsivela, E., Pentari, D., Zampas, P. and Lazaridis, M.: Physico-chemical characteristics of particulate matter in the Eastern Mediterranean, *Atmos. Res.*, 106, 93–107, doi:10.1016/j.atmosres.2011.11.011, 2012.
- 5 Kopanakis, I., Chatoutsidou, S. E., Torseth, K., Glytsos, T. and Lazaridis, M.: Particle number size distribution in the eastern Mediterranean: Formation and growth rates of ultrafine airborne atmospheric particles, *Atmos. Environ.*, 77, 790–802, doi:10.1016/j.atmosenv.2013.05.066, 2013.
- Kotroni, V., Lagouvardos, K. and Lalas, D.: The effect of the island of Crete on the Etesian winds over the Aegean Sea, *Q.J.R. Meteorol. Soc.*, 127(576), 1917–1937, doi:10.1002/qj.49712757604, 2001.
- 10 Koulouri, E., Saarikoski, S., Theodosi, C., Markaki, Z., Gerasopoulos, E., Kouvarakis, G., Mäkelä, T., Hillamo, R. and Mihalopoulos, N.: Chemical composition and sources of fine and coarse aerosol particles in the Eastern Mediterranean, *Atmos. Environ.*, 42(26), 6542–6550, doi:10.1016/j.atmosenv.2008.04.010, 2008.
- Kouvarakis, G., Doukelis, Y., Mihalopoulos, N., Rapsomanikis, S., Sciare, J. and Blumthaler, M.:  
15 Chemical, physical, and optical characterization of aerosols during PAUR II experiment, *J. Geophys. Res.*, 107(D18), 8141, doi:10.1029/2000JD000291, 2002.
- Kulmala, M., Laaksonen, A. and Pirjola, L.: Parameterizations for sulfuric acid/water nucleation rates, *J. Geophys. Res.*, 103(D7), 8301–8307, doi:10.1029/97JD03718, 1998.
- Kulmala, M., Kerminen, V.-M., Anttila, T., Laaksonen, A., and O'Dowd, C. D.: Organic aerosol  
20 formation via sulphate cluster activation, *J. Geophys. Res.*, 109, D04205, doi:10.1029/2003JD003961, 2004.
- Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H.E., Lehtipalo, K., Dal Maso, M., Aalto, P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K.E.J., Laaksonen, A., Kerminen, V.-M.:  
25 Measurement of the nucleation of atmospheric aerosol particles, *Nature Protocols*, 1651-1667, doi:10.1038/nprot.2012.091, 2012.
- Lazaridis, M., Eleftheriadis, K., Smolik, J., Colbeck, I., Kallos, G., Drossinos, Y., Zdimal, V., Vecera, Z., Mihalopoulos, N., Mikuska, P., Bryant, C., Housiadas, C., Spyridaki, A., Astitha, M. and

- Havranek, V.: Dynamics of fine particles and photo-oxidants in the Eastern Mediterranean (SUB-AERO), *Atmos. Environ.*, 40, 6214–6228, doi:10.1016/j.atmosenv.2005.06.050, 2006.
- Lazaridis, M., Dzumbova, L., Kopanakis, I., Ondracek, J., Glytsos, T., Aleksandropoulou, V., Voulgarakis, A., Katsivela, E., Mihalopoulos, N. and Eleftheriadis, K.: PM10 and PM2.5 levels in the eastern Mediterranean (Akrotiri research station, Crete, Greece), *Water, Air, and Soil Pollution*, 189(1-4), 85–101, doi:10.1007/s11270-007-9558-y, 2008.
- Lehtinen, K. E. J. and Kulmala, M.: A model for particle formation and growth in the atmosphere with molecular resolution in size, *Atmos. Chem. Phys.*, 3, 251-257, doi:10.5194/acp-3-251-2003, 2003.
- Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Korrmann, R., Lawrence, M. G., Levin, Z., Markowicz, K. M., Mihalopoulos, N., Minikin, A., Ramanathan, V., Reus, M. de, Roelofs, G. J., Scheeren, H. A., Sciare, J., Schlager, H., Schultz, M., Siegmund, P., Steil, B., Stephanou, E. G., Stier, P., Traub, M., Warneke, C., Williams, J. and Ziereis, H.: Global Air Pollution Crossroads over the Mediterranean, *Science*, 298(5594), 794–799, doi:10.1126/science.1075457, 2002.
- Liu, S. C., McKeen, S. A., Hsie, E-Y., Lin, X., Kelly, K. K., Bradshaw, J. D., Sandholm, S. T., Browell, E. V., Gregory, G. L., Sachse, G. W., Bandy, A. R., Thornton, D. C., Blake, D. R., Rowland, F. S., Newell, R., Heikes, B. G., Singh, H., Talbot, R. W.: Model study of tropospheric trace species distributions during PEM-West A, *J. Geophys. Res.*, 101(D1), 2073–2085, doi:10.1029/95JD02277, 1996.
- Manninen, H. E., Nieminen, T., Asmi, E., Gagné, S., Häkkinen, S., Lehtipalo, K., Aalto, P., Vana, M., Mirme, A., Mirme, S., Hörrak, U., Plass-Dülmer, C., Stange, G., Kiss, G., Hoffer, A., Töro, N., Moerman, M., Henzing, B., De Leeuw, G., Brinkenberg, M., Kouvarakis, G. N., Bougiatioti, A., Mihalopoulos, N., O’Dowd, C., Ceburnis, D., Arneth, A., Svenningsson, B., Swietlicki, E., Tarozzi, L., Decesari, S., Facchini, M. C., Birmili, W., Sonntag, A., Wiedensohler, A., Boulon, J., Sellegri, K., Laj, P., Gysel, M., Bukowiecki, N., Weingartner, E., Wehrle, G., Laaksonen, A., Hamed, A., Joutsensaari, J., Petäjä, T., Kerminen, V.-M. and Kulmala, M.: EUCAARI ion spectrometer measurements at 12 European sites-analysis of new particle formation events, *Atmos. Chem. Phys.*, 10, 7907–7927, doi:10.5194/acp-10-7907-2010, 2010.



- Meskhidze, N., A. Nenes, Conant, W. C., and Seinfeld, J. H.: Evaluation of a new Cloud Droplet Activation Parameterization with In Situ Data from CRYSTAL-FACE and CSTRIFE, *J. Geophys. Res.*, 110, D16202, doi:10.1029/2004JD005703, 2005.
- Mihalopoulos, N., Stephanou, E., Kanakidou, M., Pilitsidis, S. and Bousquet, P.: Tropospheric aerosol ionic composition in the Eastern Mediterranean region, *Tellus, Series B: Chemical and Physical Meteorology*, 49, 314–326, 1997.
- Morales Betancourt, R., and Nenes, A.: Aerosol Activation Parameterization: The population-splitting concept revisited, *Geosci. Mod. Dev.*, 7, 2345–2357, 2014.
- Morales R. and Nenes, A.: Characteristic updrafts for computing distribution-averaged cloud droplet number, autoconversion rate and effective radius, *J. Geophys. Res.*, 115, D18220, doi:10.1029/2009JD013233, 2010.
- Nenes, A. and Seinfeld, J. H.: Parameterization of cloud droplet formation in global climate models *J. Geophys. Res.*, 108 (D7), 4415, doi: 10.1029/2002JD002911, 2003.
- Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L. and Jayne, J. T.: An Aerosol Chemical Speciation Monitor (ACSM) for routine monitoring of the composition and mass concentrations of ambient aerosol, *Aerosol Science and Technology*, 45, 770–784, doi:10.1080/02786826.2011.560211, 2011.
- Paronis, D., Dulac, F., Chazette, P., Hamonou, E. and Liberti, G. L.: Aerosol optical thickness monitoring in the Mediterranean, *Journal of Aerosol Science*, 29(SUPPL.2), S671–S672, 1998.
- Peckham, S., Grell, G. A., McKeen, S. A., Barth, M., Pfister, G., Wiedinmyer, C., Fast, J. D., Gustafson, W. I., Zaveri, R., Easter, R. C., Barnard, J., Chapman, E., Hewson, M., Schmitz, R., Salzmann, M., Freitas, S.: WRF/Chem Version 3.3 User's Guide. NOAA Technical Memo., 98 pp, 2011.
- Petters, M.D. and Kreidenweis, S.M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, doi:10.5194/acp-7-1961-2007, 2007.
- Pikridas, M., Bougiatioti, A., Hildebrandt, L., Engelhart, G. J., Kostenidou, E., Mohr, C., Prévôt, A. S. H., Kouvarakis, G., Zampas, P., Burkhardt, J. F., Lee, B.-H., Psichoudaki, M., Mihalopoulos, N.,

- Pilinis, C., Stohl, A., Baltensperger, U., Kulmala, M. and Pandis, S. N.: The Finokalia Aerosol Measurement Experiment – 2008 (FAME-08): an overview, *Atmos. Chem. Phys.*, 10(14), 6793–6806, doi:10.5194/acp-10-6793-2010, 2010.
- Pikridas, M., Riipinen, I., Hildebrandt, L., Kostenidou, E., Manninen, H., Mihalopoulos, N., Kalivitis, N., Burkhart, J. F., Stohl, A., Kulmala, M. and Pandis, S. N.: New particle formation at a remote site in the eastern Mediterranean, *J. Geophys. Res.: Atmospheres*, 117(12), doi:10.1029/2012JD017570, 2012.
- Querol, X., Pey, J., Pandolfi, M., Alastuey, A., Cusack, M., Pérez, N., Moreno, T., Viana, M., Mihalopoulos, N., Kallos, G. and Kleanthous, S.: African dust contributions to mean ambient PM<sub>10</sub> mass-levels across the Mediterranean Basin, *Atmos. Environ.*, 43, 4266–4277, doi:10.1016/j.atmosenv.2009.06.013, 2009.
- Salisbury, G., Williams, J., Holzinger, R., Gros, V., Mihalopoulos, N., Vrekoussis, M., Sarda-Estève, R., Berresheim, H., von Kuhlmann, R., Lawrence, M. and Lelieveld, J.: Ground-based PTR-MS measurements of reactive organic compounds during the MINOS campaign in Crete, July–August 2001, *Atmos. Chem. Phys.*, 3, 925–940, doi:10.5194/acp-3-925-2003, 2003.
- Schell, B., Ackermann, I. J., Hass, H., Binkowski, F. S., Ebel, A.: Modeling the formation of secondary organic aerosol within a comprehensive air quality model system, *J. Geophys. Res. Atmos.*, 106, 28275–28293, 2001.
- Sciare, J., H. Bardouki, C. Moulin, and Mihalopoulos N.: Aerosol sources and their contribution to the chemical composition of aerosols in the Eastern Mediterranean Sea during summertime, *Atmos. Chem. Phys.*, 3, 291-302, doi:10.5194/acp-3-291-2003, 2003.
- Sciare, J., Oikonomou, K., Favez, O., Liakakou, E., Markaki, Z., Cachier, H. and Mihalopoulos, N.: Long-term measurements of carbonaceous aerosols in the Eastern Mediterranean: Evidence of long-range transport of biomass burning, *Atmos. Chem. Phys.*, 8, 5551-5563, doi:10.5194/acp-8-5551-2008, 2008.
- Seinfeld, J., and Pandis, S. (Eds): *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2<sup>nd</sup> edition, John Wiley, Hoboken, N. J., ISBN: 978-0-471-72018-8, 1232 pp., 2006.

- Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling. *J. Geophys. Res. Atmos.*, 95, 16343–16367, 1990.
- 5 Tombrou, M., Bossioli, E., Protonotariou, A. P., Flocas, H., Giannakopoulos, C., and Dandou, A.: Coupling GEOS-CHEM with a regional air pollution model for Greece, *Atmos. Environ.*, 43, 4793–4804, doi:10.1016/j.atmosenv.2009.04.003, 2009.
- Tombrou, M., Bossioli, E., Kalogiros, J., Allan, J. D., Bacak, A., Biskos, G., Coe, H., Dandou, A., Kouvarakis, G., Mihalopoulos, N., Percival, C. J., Protonotariou, A. P., and Szabó-Takács, B.: Physical and chemical processes of air masses in the Aegean Sea during Etesians: Aegean-GAME  
10 airborne campaign, *Sci. Total Environ.*, 506-507, 201–216, doi:10.1016/j.scitotenv.2014.10.098, 2015.
- Triantafyllou, E., M. Giamarelou, E., Bossioli, P., Zarmas, C., Theodosi, C., Matsoukas, M., Tombrou, N., Mihalopoulos, and G., Biskos.: Particulate Pollution Transport Episodes from Eurasia to a Remote Region of Northeast Mediterranean, *Atmos. Environ.*, 128, 45–52,  
15 doi:10.1016/j.atmosenv.2015.12.054, 2015.
- Tuccella, P., Curci, G., Visconti, G., Bessagnet, B., Menut, L., Park, R.J., 2012. Modeling of gas and aerosol with WRF/Chem over Europe: evaluation and sensitivity study. *J. Geophys. Res. Atmos.* 117.
- Tyrlis, E., Lelieveld, J., Climatology and Dynamics of the Summer Etesian Winds over the Eastern  
20 Mediterranean. *J. Atmos. Sci.* ,70, 3374–96, doi: 10.1175/JAS-D-13-035.1, 2013.
- Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Gruning, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G., O’ Dowd, C. D., Marinoni, A., Horn, H. G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Löschau, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term  
25

observations of atmospheric particle number size distributions, *Atmos. Meas. Tech.*, 5, 657–685, doi:10.5194/amt-5-657-2012, 2012.

5 Ždímal, V., Smolík, J., Eleftheriadis, K., Wagner, Z., Housiadas, C., Mihalopoulos, N., Zbynek Vecera, P. M., Kopanakis, I., and Lazaridis, M.: Dynamics of Atmospheric Aerosol Number Size Distributions in the Eastern Mediterranean During the SUB-AERO Project, *Water Air Soil Pollut.*, doi:10.1007/s11270-010-0410-4, 2010.

10 Zerefos, C. S., Kourtidis, K. A., Melas, D., Balis, D., Zanis, P., Katsaros, L., Mantis, H. T., Repapis, C., Isaksen, I., Sundet, J., Herman, J., Bhartia, P.K., and Calpini, B.: Photochemical activity and solar ultraviolet radiation (PAUR) modulation factors: an overview of the project, *J. Geophys. Res.*, 107 (D18), 8134, 2002.

## Tables

**Table 1.** Summary of the variables and operation characteristics of the instruments at Santorini and 5 Finokalia stations.

<b>Santorini</b>	<b>Instrument</b>	<b>Resolution</b>	<b>Period Operation</b>
<i>Aerosols</i>			
Aerosol Number Distribution (10 - 500 nm)	TSI 3034 SMPS	3 min	15 – 28 July
<i>Gaseous Species</i>			
O <sub>3</sub>	M400E Photometric ozone analyzer	1 min	18 – 28 July
SO <sub>2</sub>	M100E UV Fluorescence analyzer	1 min	15 – 28 July
NO, NO <sub>2</sub> , NO <sub>x</sub>	M200E Nitrogen Oxide analyzer	1 min	15 – 28 July
<b>Finokalia</b>			
<i>Aerosols</i>			
Aerosol Number Distribution (9 – 848 nm)	TROPOS type SMPS	5 min	16 – 29 July
Chemical composition (SO <sub>4</sub> <sup>2-</sup> , NO <sub>3</sub> <sup>-</sup> , NH <sub>4</sub> <sup>+</sup> , Cl <sup>-</sup> , organics)	Aerodyne Research Inc. Aerosol Chemical Speciation Monitor (ACSM)	30 min	15 – 28 July
<i>Gaseous Species</i>			
O <sub>3</sub>	Thermo electron Model 49I	3 min	15 – 28 July
<i>Meteorology</i>			
Relative humidity, Temperature	MP101A Humidity- Temperature	5 min	15 – 28 July
Wind Speed, direction	05103 Wind Monitor	5 min	15 – 28 July

**Table 2.** Average ( $\pm$  standard deviation) of O<sub>3</sub> concentrations and aerosol mass and number concentrations during the two examined periods; a) EF (22 - 24 July) and b) MSF (25 - 27 July).

5

Tracer	Santorini	Finokalia	Santorini	Finokalia
	EF period		MSF period	
O <sub>3</sub> (ppbv)	51.4 $\pm$ 6.2	53.8 $\pm$ 4.5	70.0 $\pm$ 8.5	71.0 $\pm$ 7.9
Sulfate ( $\mu\text{g m}^{-3}$ )	N/A	3.1 $\pm$ 1.2	N/A	7.3 $\pm$ 1.5
Ammonium ( $\mu\text{g m}^{-3}$ )	N/A	1.4 $\pm$ 0.6	N/A	3.1 $\pm$ 0.6
Organics ( $\mu\text{g m}^{-3}$ )	N/A	4.2 $\pm$ 1.3	N/A	8.6 $\pm$ 1.2
Nitrate ( $\mu\text{g m}^{-3}$ )	N/A	0.38 $\pm$ 0.12	N/A	0.8 $\pm$ 0.1
Total number conc. ( $\text{cm}^{-3}$ )	3.6 $\pm$ 2.1 $\times 10^3$	.9 $\pm$ 1.2 $\times 10^3$	2.0 $\pm$ 0.6 $\times 10^3$	3.6 $\pm$ 0.5 $\times 10^3$
Aitken mode ( $\text{cm}^{-3}$ )	2.2 $\pm$ 1.4 $\times 10^3$	.5 $\pm$ 1.5 $\times 10^3$	1.2 $\pm$ 0.5 $\times 10^3$	1.6 $\pm$ 0.5 $\times 10^3$
Accumulation mode ( $\text{cm}^{-3}$ )	9.6 $\pm$ 3.5 $\times 10^2$	.6 $\pm$ 0.9 $\times 10^3$	1.0 $\pm$ 0.5 $\times 10^3$	2.1 $\pm$ 0.6 $\times 10^3$

**Table 3:** Average ( $\pm$  standard deviation) of calculated  $\kappa$  using the PM<sub>1</sub> chemical composition at Finokalia, the  $d_c$  (as described in the text), and the estimated CCN concentration particles at both stations, on 23 and 24 July (EF period). Here  $S_{max}$  is the maximum supersaturation in the cloud,  $N_{total}$  is the total particle number concentration, and  $N_d$  is the potential cloud droplet number concentration calculated according to the approach described in the main text. Two probability density function (PDF) of the characteristic updraft velocity are used with  $\sigma_w = 0.3 \text{ m s}^{-1}$  and  $\sigma_w = 0.6 \text{ m s}^{-1}$ . Time is in LST.

	Santorini		Finokalia		Santorini		Finokalia	
	23/7				24/7			
	<i>Before</i> 00:00 - 8:00	<i>After</i> 15:00 - 21:00	<i>Before</i> 00:00 - 10:00	<i>After</i> 17:00 - 21:00	<i>Before</i> 00:00 - 10:00	<i>After</i> 18:00 - 21:00	<i>Before</i> 00:00 - 11:00	<i>After</i> 17:00 - 21:00
$\kappa$	0.29 $\pm$ 0.01	0.36 $\pm$ 0.03	0.28 $\pm$ 0.02	0.38 $\pm$ 0.02	0.29 $\pm$ 0.01	0.34 $\pm$ 0.01	0.30 $\pm$ 0.01	0.34 $\pm$ 0.01
$d_c$ (nm) ( $s=0.2\%$ )	104 $\pm$ 2	95 $\pm$ 2	104 $\pm$ 2	94 $\pm$ 1	101 $\pm$ 1	96 $\pm$ 1	102 $\pm$ 2	97 $\pm$ 1
$d_c$ (nm) ( $s=0.6\%$ )	50 $\pm$ 1	46 $\pm$ 1	50 $\pm$ 1	45 $\pm$ 1	49 $\pm$ 1	46 $\pm$ 1	49 $\pm$ 1	46 $\pm$ 0
CCN <sub>0.2</sub> (cm <sup>-3</sup> )	536 $\pm$ 27	794 $\pm$ 145	1002 $\pm$ 76	1420 $\pm$ 383	682 $\pm$ 66	1028 $\pm$ 61	1062 $\pm$ 156	1822 $\pm$ 154
CCN <sub>0.6</sub> (cm <sup>-3</sup> )	1225 $\pm$ 90	3155 $\pm$ 789	2111 $\pm$ 196	4343 $\pm$ 1119	1535 $\pm$ 66	2004 $\pm$ 224	2191 $\pm$ 270	3346 $\pm$ 399
$N_{total}$ (cm <sup>-3</sup> )	1777 $\pm$ 421	4621 $\pm$ 1986	3506 $\pm$ 699	5710 $\pm$ 779	2306 $\pm$ 154	2557 $\pm$ 351	3198 $\pm$ 384	3921 $\pm$ 404
$\sigma_w=0.3 \text{ m s}^{-1}$								
$S_{max}$ (%)	0.25	0.22	0.11	0.10	0.23	0.19	0.11	0.07
$N_d$ (cm <sup>-3</sup> )	110 $\pm$ 4	124 $\pm$ 8	423 $\pm$ 4	407 $\pm$ 19	121 $\pm$ 5	165 $\pm$ 9	423 $\pm$ 3	440 $\pm$ 5
Activation Fr. (%)	6.5 $\pm$ 1.5	3.1 $\pm$ 1.1	12.5 $\pm$ 2.4	7.2 $\pm$ 0.6	6.4 $\pm$ 0.4	7.9 $\pm$ 1.2	13.4 $\pm$ 1.6	11.3 $\pm$ 1.1
Contribution of $\kappa$ (%)	1.4	2.7	2.6	10.2	0.7	1.9	0.9	0.3
Contribution of $N_{aer}$ (%)	98.6	97.3	97.4	89.8	99.3	98.1	99.1	99.7

$\sigma_w=0.6 \text{ m s}^{-1}$									
$S_{max}$ (%)	0.32±0.01	0.28±0.01	0.14±0.01	0.14±0.01	0.29±0.01	0.23±0.01	0.14±0.01	0.11±0.01	
$N_d$ (cm <sup>-3</sup> )	192±6	217±15	627±67	619±109	213±7	286±15	621±73	786±11	
Activation Fr. (%)	11.4±2.6	5.4±1.9	18.8±5.1	10.8±0.7	11.3±0.7	13.7±2.1	19.7±3.1	20.2±1.9	
Contribution of $\kappa$ (%)	1.2	1.9	3.8	19.0	0.6	1.6	0.7	0.2	
Contribution of $N_{act}$ (%)	98.8	98.1	96.2	81.0	99.4	98.4	99.3	99.8	



## Figure captions

**Fig. 1.** The extended area of study where the major routes (arrows) of air masses passing through Santorini and Finokalia, during 23 July (EF - left panel) and 26 July (MSF - right panel), are indicated.

5 On the left panel, the areas of NPF (black ellipse), the spatial extent of NPF event (red line), flow entrainment into MABL (white dashed line) and the condensation (yellow dashed ellipse), are shown. The major traffic (green) and urban (red) emission sources, are also shown. The marine traffic is shown in the middle panel.

**Fig. 2.** HYSPLIT4 back-trajectories computed with an end point at the Santorini station (from the heights of 100, 500 and 1000 m), on 23 (left panel), 24 (central panel) (both during EF period) and 26 July (MSF - right panel), 2013.

**Fig. 3.** Timeseries of the wind speeds (ws, solid lines on left axis) and wind directions (wd, open circles, right axis) at Santorini (simulations by the WRF-Chem model) and at Finokalia (measurements). The second period of the EF is shaded with yellow and the MSF with grey.

**Fig. 4.** Aerosol modal number concentrations, meteorological and pollutant ( $O_3$ ,  $NO_2$  when available) concentrations, along with meteorological parameters of relative humidity (RH) and surface temperature (T) at Santorini (top panel) and Finokalia (bottom panel). Note that  $SO_2$  is shown at Santorini, while  $SO_4$  is shown for Finokalia.

**Fig. 5.** Spatial distribution of  $O_3$  concentration (ppb) and wind speed at 400 m asl over the extended area of Greece as simulated by WRF-Chem at 15:00 LST for 23 July (EF - left panel), and 26 July (MSF - right panel).

25

**Fig. 6.** Mass concentrations of submicron aerosol measured at Finokalia station from 13 to 30 of July 2013.


**Fig. 7.** As in Fig. 5, but for sulfate concentration ( $\mu\text{g m}^{-3}$ ) at 09:00 LST.

**Fig. 8.** Diurnal evolution of the aerosol size-distribution on 23 and 24 July (EF) at Santorini (top panel) and Finokalia (bottom panel). The white dots stand for nucleation, the black dots for Aitken and the purple dots for accumulation geometric mean diameter.

**Fig. 9.** Timeseries of the CN and estimated CCN concentration particles, for various supersaturations, at Santorini (top panel) and Finokalia (bottom panel), during 23 and 24 July (EF). Time is in LST.

10 **Fig. 10.** Timeseries of the estimated cloud droplet number concentrations ( $N_d$ ), and maximum supersaturation in the cloud ( $s_{max}$ ) for updraft velocities ( $\sigma_w$ ) of  $0.3 \text{ m s}^{-1}$  and  $0.6 \text{ m s}^{-1}$ , at Santorini (top panel) and Finokalia (bottom panel), during 23 and 24 July (EF). Blue and red lines correspond to updraft velocity ( $\sigma_w$ ) equal to  $0.3 \text{ m s}^{-1}$ , while orange and green to  $0.6 \text{ m s}^{-1}$ .

# Figures



**Fig. 1.**

5




Fig. 2







Fig. 3




**Fig. 4**




5

**Fig. 5**




**Fig. 6**



5



**Fig. 7**




**Fig. 8**



5 Fig. 9





**Fig. 10**