

Interactive comment on “Case Studies of the Impact of Orbital Sampling on Stratospheric Trend Detection and Derivation of Tropical Vertical Velocities: Solar Occultation versus Limb Emission Sounding” by L. F. Millán et al.

L. F. Millán et al.

lmillan@jpl.nasa.gov

Received and published: 8 June 2016

We thank the reviewer for his/her comments. Below are our responses in blue.

1 Reviewer 2 comments

Summary

C1

This paper analyzes the impact of orbital sampling on remotely sensed data and investigates implications of related sampling biases and drifts. Although this is a methodical and partly technical paper, I recommend publication in ACP rather than any of the more technical journals, because the readership addressed are scientists working in atmospheric sciences rather than scientists working in more technical fields.

The methodology chosen is adequate; results are interesting; the presentation is concise without any unnecessary length but still complete; the authors have taken care to put their work adequately into the context of existing work. I recommend publication in ACP and I have only a list of minor comments which may help to improve the paper.

Comments

p.2 l24-32: While I usually insist that results must be summarized in the abstract (as correctly done in this paper), I am not sure if it is adequate to summarize the results already in the introduction. Usually it is announced in the introduction what will be investigated, without anticipating the results.

The lines detailing the results will be deleted.

p3 l18: It should probably read “long-term O3 trends” here? *Ok*

p4 l4: I miss some information about the HALOE altitude resolution here. *This sentence will be added: The vertical resolution of this dataset is about 2.3 km.*

p4 l9: Same for ACE-FTS. *This sentence will be added: The vertical resolution of this dataset is about 3 km.*

C2

p4 l13: Same for MLS. This sentence will be added: The vertical resolution of this dataset is, in general, about 3 km.

p4 l21: It might not even be necessary to assume that the vertical resolutions are the same. If the true fields are smooth enough in the vertical domain, even different vertical resolutions cause only marginal differences. Possibly the assumption of equal vertical resolutions can be challenged, and since a much weaker assumption (that all vertical resolutions are good enough to resolve the fields) might be sufficient, I suggest to formulate the argument with this weaker assumption. We will add: Given that our focus is in horizontal/temporal sampling, all satellite measurements are assumed to have the same vertical resolution as CMAM30-SD, however we want to emphasize that the vertical resolution of these instruments is in general good enough to resolve the model fields.

p5 l12-22: Some discussion of the distributions of the biases might be useful. Where are the largest biases and why? Is this consistent with Toohey et al.? Although we do not discuss Figure 3 sampling biases we do discuss the RMS sampling biases in Figure 4 and compare it against / agreeing with Toohey et al. We believe that a discussion of Figure 3 sampling biases will be redundant.

p4 l22: On page 3 it is not stated how fine the vertical sampling of the model is in the lower and middle stratosphere but I guess it is much better than the vertical resolution of ACE-FTS, which is often reported to be around 3 km, if I remember correctly. Unfortunately to my knowledge neither for ACE-FTS nor for HALOE averaging kernels are available. But if you smooth the model fields using a triangular or Gaussian function of 3 km full width at half maximum, and if the model fields change only marginally by

C3

this, then you are on the safe side. At least Fig 2. suggests that there seem to exist no major vertical resolution related problems. we will modify this sentence to: That is, the impact of the averaging kernels is not addressed in this study, however, we note that for the parameters studied here a 3km averaging kernel does not impact significantly their values in the upper troposphere/stratosphere.

p5 l25 Temperature: why capital "T"? it will be changed to lowercase

p6 l4: RMS and standard deviation are often not adequately distinguished, and I am not sure if centered RMDd is a widely known concept. I would not mind to see these terms defined by equations here. After careful consideration we decided against including more equations that might add complexity to the study, we feel that the reader could easily go to any statistic books and to Taylor (2001) for detail descriptions. Furthermore, reviewer 2 did not find this issue problematic.

p6 l5 "everything" sounds a bit vague. "all data" or something similar would sound a bit more "intellectual" but would say essentially the same. as suggested, we will use "all data"

p6 l15 shows the raw model (remove one "the") . Ok

p7 l1: It would be interesting to know if the autocorrelation is assumed constant over the entire time series or if it has some time-dependence in itself. the autocorrelation is assumed constant over the entire time series, we will add: computed following Tiao et al. (1990), so the reader can see how it was estimated. Also, we will add on page 7 line 20: ϕ is computed for the raw as well as the satellite-sampled data.

C4

p7 l4: It would be interesting to see if the sampling artifacts somehow show up as autocorrelation. Can we learn anything about this by analysis of the values of the various instruments? The reviewer points out an excellent point. The figures/analysis shown uses the autocorrelation derived from the raw model as well as the satellite-sampled data. However, using the autocorrelation from the raw-model instead of the satellite-sampled data did not produce significantly different results. We will add the following sentence at the end of page 7 (original document): We also performed this analysis using only the autocorrelation computed for the raw model data and found no significant differences.

p7 l10: Trends in state variables can have latitudinal structure (see, e.g. Eckert et al., ACP, 2014, for ozone). I do not challenge the adequacy of trend analysis in more global terms but a caveat about latitudinal structure might be useful. That is partly why we included 11. We will add the following sentence at the end of section 4: As shown, the ability to detect trends depends upon the natural variability and the correlation of the data. These in turn vary upon which parameter is being studied as well as which location and height. Studies of natural variability and autocorrelation of the data will help identify where to monitor to find more readily detectable trends, however, such a study is outside the scope of this paper.

p7 l21-23: I am not happy with the term “noise” here, because this can easily be understood as “measurement noise”. But here the residuum is meant, which contains real physical effects, like QBO or whatever. I suggest “The autocorrelation of the residual between the data points and the trend model” ... “The standard deviation of the residual, which corresponds ...” Ok

l7 l25: I do not like the term “model” here, because there are so many models (trend models, CMAM, etc). I would prefer a more specific term here. It will be change to

C5

trend model (equation 4)

p8 l10-12: The dichotomy “solar occultation vs. microwave emission” does not exist. The HALOE and ACE-FTS data gaps due to clouds are due to the spectral region (IR), not to the measurement geometry (occultation). The IR instrument MIPAS, e.g., has also data gaps due to clouds although it is an emission instrument. The statement made here is certainly correct (except for the dichotomy-like wording) but it seems to me to have nothing to do with orbital sampling. I think that the discussion of this issue detracts from the main issue of this paper. If you want to discuss sampling artifacts due to clouds, the fact that IR sounders have a sampling bias towards the cloud-free atmosphere would be interesting and deserves an investigation, particularly in the context of water vapor, but this is clearly beyond the scope of this paper. We will delete that sentence.

Sect 4 general: Unless I have got things completely wrong, the following question remains unanswered: MLS provides both denser sampling and more uniform sampling. Both characteristics contribute to the better MLS trends. Which of the sampling attributes (data density or uniform sampling) is the primary cause of the better MLS trends? That is correct, that question remains unanswered and a more detailed study, presumably with fake sampling patterns, will have to be carried out. Could analysis of the values help to find this out? We do not think so, see comments about ϕ above.

If I haven't missed anything, the data points have equal weight in the trend fit. Is this adequate? Since orbits converge with higher latitudes, each data point represents a smaller area at higher latitudes. Thus, high latitudes are over-represented if no area weighting is applied. This, however, is a kind of sampling bias in itself. As the reviewer points out this will affect primarily at higher latitudes and we only use latitudes between 60S and 60N where we do not expect any significant difference between equal weight

C6

and area weighted trends.

p9 l13 ff: there appear a lot of abbreviations and acronyms (QBO, ENSO, QSI, MEI). Please make sure that they all are defined. Some of them are but I have not checked all of them. [they are defined](#)

p10 l26: Here the better performance of instruments with MLS sampling pattern is attributed to the data density. This is intuitive but I am not sure if it has really been shown that this is not due to uniformity. The ϕ is not necessarily the same for the instruments under investigation. [We will modify the sentence to: This is because the sparse **non-uniform** sampling leads to an increase ... See comments about \$\phi\$ above.](#)

p11 l7: This reads a bit as if the increase in the circulation is a fact but to my knowledge it is an expectation based on model calculations. To my knowledge the increase of circulation has not yet been empirically confirmed, at least not beyond any doubt. I would prefer “such as a possible increase in the circulation”. For my personal taste, the last three lines are a bit too general and I would prefer a concluding remark which is more focused on the sampling issue. [we will modify the sentence to include “such as a possible increase in the circulation”](#)

p11 l10: Finally a fussy one: (Do not take it too seriously because as a non-native English speaker I may be wrong here) It sounds funny to me that a laboratory does the work. I thought it is the researchers who do the work. The researchers are listed in the authors' list, along with their affiliations. Thus it seems redundant to me to mention this again in the acknowledgements. [it will be changed to: Work at the Jet Propulsion Laboratory, California Institute of Technology, was done under contract with the National Aeronautics and Space Administration.](#)

C7

Fig Cap 5: “either K or %” sounds a bit vague. Better: ...as a function of pressure for temperature (in Kelvin) and O₃... (in %). Again: why capital “T” for temperature” ? [Ok, the new sentence will be: Mean \(thin lines\) and maximum \(thicker lines\) RMS sampling bias over all latitudes for 2005 as a function of pressure for temperature \(in Kelvin\), and O₃, CH₃Cl, H₂O, CO, HCl, N₂O and HNO₃ \(in %\). The vertical grid indicates values of 0.5, 1, 5, 10, 50, 100.](#)

Interactive comment on Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2016-356, 2016.

C8