



# Are Atmospheric Updrafts a Key to Unlocking Climate Forcing and Sensitivity?

Leo J. Donner<sup>1</sup>, Travis A. O'Brien<sup>2</sup>, Daniel Rieger<sup>3</sup>, Bernhard Vogel<sup>3</sup>, and William F. Cooke<sup>4</sup>

<sup>1</sup>NOAA/GFDL, Princeton University Forrestal Campus, 201 Forrestal Road, Princeton, New Jersey 08540 USA

<sup>2</sup>Lawrence Berkeley National Laboratory, Berkeley, California USA

<sup>3</sup>Karlsruhe Institute of Technology, Karlsruhe, Germany

<sup>4</sup>UCAR/GFDL, Princeton, New Jersey USA

*Correspondence to:* L.J. Donner (leo.j.donner@noaa.gov)

**Abstract.** Both climate forcing and climate sensitivity persist as stubborn uncertainties limiting the extent to which climate models can provide actionable scientific scenarios for climate change. A key, explicit control on cloud-aerosol interactions, the largest uncertainty in climate forcing, is the vertical velocity of cloud-scale updrafts. Model-based studies of climate sensitivity indicate that convective entrainment, which is closely related to updraft speeds, is an important control on climate sensitivity.

5 Updraft vertical velocities also drive many physical processes essential to numerical weather prediction.

Vertical velocities and their role in atmospheric physical processes have been given very limited attention in models for climate and numerical weather prediction. The relevant physical scales range down to tens of meters and are thus frequently sub-grid and require parameterization. Many state-of-science convection parameterizations provide mass fluxes without specifying vertical velocities, and parameterizations which do provide vertical velocities have been subject to limited evaluation  
10 against what have until recently been scant observations. Atmospheric observations imply that the distribution of vertical velocities depends on the areas over which the vertical velocities are averaged. Distributions of vertical velocities in climate models may capture this behavior, but it has not been accounted for when parameterizing cloud and precipitation processes in current models.

New observations of convective vertical velocities offer a potentially promising path toward developing process-level cloud  
15 models and parameterizations for climate and numerical weather prediction. Taking account of scale-dependence of resolved vertical velocities offers a path to matching cloud-scale physical processes and their driving dynamics more realistically, with a prospect of reduced uncertainty in both climate forcing and sensitivity.

## 1 Introduction

Uncertainties in both anthropogenic climate forcing and climate sensitivity continue to limit our understanding of climate  
20 change and the precision with which scenarios for future climate change can be constructed. As had Kiehl (2007) for an earlier generation of climate models, Forster et al. (2013) found that CMIP5 models able to successfully simulate observed global warming over the pre-industrial to present-day period did so by balancing a range of anthropogenic climate forcings and climate sensitivities. Both forcing and sensitivity have proved resistant to reducing their uncertainties. Understanding the




relative roles of forcing and sensitivity, along with variability, is essential to actionable estimates of future climate change. Given the long lifetime of greenhouse gases relative to atmospheric aerosols, aerosol "masking" of warming by greenhouse gases over the pre-industrial to present-day period will become less important. The future corresponding to a "low net-forcing, high-sensitivity" twentieth century differs from that for "high net-forcing, low sensitivity." The importance of this question 5 is cast into stark relief by assessments of the extent to which the Intended Nationally Determined Contributions, emissions reductions pledged at the recent Paris COP21 conference, will meet the COP21 goal of holding increases in globally averaged temperature below 2° C by 2100. The relationship between temperature increases, other elements of human-induced climate change, and emissions remains uncertain within a range implied by ongoing uncertainty in sensitivity. Reducing this uncertainty would be of great value as emissions goals are revised going forward.

10 This review presents the perspective that the vertical velocities, updrafts on all scales, are among the keys to understanding and simulating climate forcing and are plausibly also important for climate sensitivity. Vertical velocities have received limited attention in climate (and even cloud) models and have only recently become a focus of observational studies. Sub-grid clouds are parameterized in climate models, traditionally based on mass fluxes (product of vertical velocity, density, and area) without vertical velocity specifically. The dependence and realism of explicitly resolved vertical velocities, even in higher-resolution 15 models, has not been extensively examined.

## 2 Vertical Velocity and Climate Forcing

Aerosol-cloud interactions are the largest source of uncertainty in climate forcing, with estimates ranging from close to zero to  $-1.3 \text{ W m}^{-2}$ , in contrast to forcing by carbon dioxide of  $1.7 \pm 0.4 \text{ W m}^{-2}$  (Stocker et al., 2013). Rosenfeld et al. (2013) and Rosenfeld et al. (2014) discuss elements of aerosol-cloud interactions leading to this uncertainty. Here, we emphasize 20 that the updraft speeds at which cloud liquid and ice are activated are among the primary controls on cloud drop sizes and number concentrations, which are in turn related to cloud optical properties, precipitation, and macrophysical properties. Fig. 1 shows the sensitivities of drop effective size (Feingold, 2003) and number concentration (McFiggans and Co-Authors, 2006) to vertical velocity, aerosol number concentration and size distribution, and aerosol composition. Drop sizes and number concentrations are more sensitive to vertical velocity than aerosol composition, and, as aerosol concentrations increase from 25 clean to polluted, vertical velocity becomes increasingly important relative to aerosol number and size. For homogeneous freezing, ice crystal number concentrations are often controlled more by vertical velocity than aerosol number concentration (Fig. 2, (Kay and Wood, 2008)).

In general, then, physically based simulation of aerosol-cloud interactions requires knowledge of the updraft speeds at which 30 these interactions occur, along with requisite information on aerosol composition and size distributions. It is important to note that these updrafts occur on a wide range of scales, down to large eddies with horizontal and vertical scales of tens of meters. As a matter of simulation, this implies parameterization of vertical velocities in models whose scales are coarser than models




**Figure 1.** Sensitivities of (a) drop effective radius ( $r_e$ ) (Feingold, 2003) and (b) drop number concentration ( $N_d$ ) (McFiggans and Co-Authors, 2006) to aerosol number concentration ( $N_a$ ), aerosol median size ( $r_g$ ), breadth parameter ( $\sigma$ ) for the unimodal lognormal distribution assumed for aerosol sizes, vertical velocity ( $w$ ), and aerosol mass fraction ( $\epsilon$ ) of ammonium sulfate. The drop effective radius  $r_e$  is the ratio of the number-weighted radius cubed to the number-weighted radius squared.

that resolve large eddies, and, even in large-eddy simulations, attention to the realism of distributions of vertical velocities will be important.

### 3 Vertical Velocity and Climate Sensitivity

The possible relationship of vertical velocity to climate sensitivity, broadly defined as the response of a climate measure such as global, annual-mean surface temperature to a change in climate forcing, is less obvious. The strongest suggestions of a link emerge from several studies showing that the rates of entrainment in cumulus convection in climate models are related to the sensitivity of these models (Stainforth et al. (2005), Rougier et al. (2009), Sanderson et al. (2010), Zhao (2014)). This result is perhaps surprising, given that low- and midlevel-cloud feedback is the major uncertainty in climate sensitivity (Zelinka et al., 2012). One possible explanation is provided by Sherwood et al. (2014), who trace climate sensitivity to convective mixing, in turn related to dehydration of low-cloud layers. In their perturbed parameter experiments with a climate model, Klocke et al. (2011) found that the entrainment rate for shallow convection explained most of the variation in its sensitivity, with no sensitivity to entrainment rate for deep convection. As a mechanism for forming low clouds and an agent for dehydrating the



**Figure 2.** Aerosol sensitivity (derivative of logarithm of ice crystal number concentration with respect to logarithm of aerosol concentration) and ice number concentration (contoured) as functions of vertical velocity and aerosol number concentration for homogeneous freezing, from Kay and Wood (2008). Shaded aerosol sensitivities are for a parcel lifted at  $-50^\circ\text{C}$  and 250 hPa, with the line indicating an aerosol sensitivity of 0.5 at  $-80^\circ\text{C}$  and 100 hPa. Observations (10-90% percentile) from the INCA (INterhemispheric differences in Cirrus properties from Anthropogenic emissions) field campaign fall within the light circle.

lower atmosphere, shallow convection quite plausibly is important for climate sensitivity. Heat released in deep convection drives the Hadley and Walker circulations, and, as noted by Su et al. (2014), changes in cloud radiative effect in the tropics are closely related to patterns of strengthening and weakening of the Hadley circulation. Taken together, these results point to the important role played by deep cumulus convection in determining the dynamic and thermodynamic environments in which stratocumulus and shallow cumulus form. Though low and midlevel clouds are the proximate agent determining climate sensitivity, deep convection is among their remote controls and thereby also possibly important to climate sensitivity. The roles of these mechanisms likely exhibit model dependence, as suggested, for example, by Klocke et al. (2011)'s insensitivity to




deep convective entrainment. Entrainment and detrainment in convection are closely related to their vertical profiles of mixing, heating, moistening, and drying. Through its influence on buoyancy, entrainment bears a close relationship to vertical velocity. Though less directly tied to climate sensitivity than climate forcing, vertical velocities through these mechanisms are quite plausibly correlated to climate sensitivity.

## 5 4 Modeling Implications and Prospective Breakthroughs

The implications of strong dependencies of climate forcing and sensitivity on vertical velocities present a challenge to current climate model development but also hold the promise of possible breakthroughs. Since the physically relevant scales for vertical velocity are not only resolved explicitly in climate models but also sub-grid, both the vertical velocities from the model dynamical cores and those in parameterizations for sub-grid processes are important. Neither has been given much attention  
10 in model development to date. Indeed, many parameterizations of sub-grid processes, e.g., mass-flux parameterizations for cumulus convection, have not even provided vertical velocities. Therein lies the promise for possible breakthroughs, as attention can be turned to these issues. There are numerous challenges, related especially to scale awareness for physical processes and realism of parameterized and resolved vertical velocities.


Empirically and theoretically, there is strong reason to suspect that vertical velocity should depend on resolution (Rauscher  
15 et al., 2016). Consider the discretized continuity equation:  $\Delta_x u_D / \Delta x + \Delta_y v_D / \Delta y + \Delta_p \omega / \Delta p = 0$ , where  $u_D$  and  $v_D$  represent the horizontally divergent wind components,  $\omega$  represents vertical velocity in pressure coordinates,  $\Delta_*$  represents a finite difference operator, and  $\Delta x \sim \Delta y$  represents the horizontal differencing distance. Scale analysis of this equation implies that vertical velocity scales as  $|\omega| \propto |\Delta_x u_D| / \Delta x$ . Inspection of this relationship reveals that  $|\Delta_x u_D|$  is equivalent to the first order structure function of wind, which has been demonstrated to exhibit power-law behavior in nature:  $|\Delta_x u_D| \propto x^H$  (e.g., Cho  
20 and Lindborg, 2001). Hence the combination of mass continuity and the scaling properties of the wind field imply that vertical velocities should change with averaging distance or, in a model, resolution:  $|\omega| \propto \Delta x^{H-1}$ . For typical structure function exponents of  $H \sim 1/3$ , this implies that vertical velocity increases with resolution like  $\Delta x^{-2/3}$ . Fig. 3 shows that the structure function from a high-resolution forecast model indeed closely follows this  $\Delta x^{-2/3}$  relationship over the range of resolutions represented in the CMIP3 and CMIP5 model archives.

25 Relatedly, the probability distribution of updraft sizes in the same forecast output also closely follows a powerlaw of updraft area over all observable scales in the model (Fig. 3). However, the current paradigm for the subgrid representation of updrafts assumes a separation of scales: that convection occurs at scales  $\mathcal{O}(10\text{km})$  or less. This implies that there is a continuum of updraft sizes below the resolution of contemporary climate models that are not represented by current parameterizations. Increasing model resolution will increase the representation of updrafts in the unparameterized continuum, represented by the  
30 gray swath in Fig. 3. So unless subgrid parameterizations compensate accordingly, this will naturally result in an increase in vertical transport as resolution increases.



**Figure 3.** Probability distribution of updraft area  $P(A)$  (black curve; left axis) and the structure function of the zonal, divergent wind component  $|\Delta_x U_D|$  divided by the structure function distance  $\Delta x$  (blue curve; right axis). Both  $P(A)$  and the structure function are derived from 1.5 years of 500 hPa output from the ECMWF Year of Tropical Convection (YoTC) T799 operational forecast output (European Centre for Medium-Range Weather Forecasts, 2012). The shaded golden and blue regions correspond to the interquartile range of CMIP3 and CMIP5 models respectively, and the shaded rose region corresponds to the scales at which convection occurs.  $A^{-5/3}$  and  $\Delta x^{-2/3}$  powerlaws are provided for reference (gray dashed and gray dash-dotted lines respectively).

As discussed previously, vertical velocity is a critical control on droplet and ice activation. This suggests that climate forcing could exhibit a dependence on model resolution, even if vertical velocities were simulated realistically for a given resolution. Indeed, Ma et al. (2015) found that aerosol indirect forcing decreased by about 30% in the Northern Hemisphere midlatitudes, where most anthropogenic emissions occur, and 15% globally in CAM5 as horizontal grid spacing was decreased from  $2^{\circ}$  to  $0.25^{\circ}$ . Resolved vertical velocities are among the factors to which aerosol indirect forcing is related in their study. To examine the dependence of ice activation on model resolution, we modeled ice activation by heterogeneous freezing in a massive Saharan dust plume which advected to central Europe on 3 April 2014 using the ICON (ICOsehedral Non-hydrostatic) numerical weather prediction and climate model (Zängl et al., 2015), developed at the Deutscher Wetterdienst and Max Planck Institute for Meteorology and extended as ICON-ART to include aerosols and their interactions with clouds (Rieger et al., 2015). Fig. 4a shows the vertical velocities at which heterogeneous freezing occurs depend strongly on the ICON-ART horizontal resolution. Figs. 4b-d show that this dependence leads to large changes in the number of ice particles produced by heterogeneous freezing, with overall increases in ice crystals formed at finer resolutions. These results show the distributions of cloud microphysical, and thereby radiative and precipitation, properties depend on model resolution. The power-law dependence of the observed structure functions, shown in Fig. 3, suggests that such dependencies could possibly be taken into account by scaling the



**Figure 4.** (a) Relative frequency of vertical velocity in the heterogeneous freezing regime ( $235 \text{ K} < T < 273.15 \text{ K}$ ) for ICON-ART horizontal resolutions of  $\Delta x = 40, 20, 10$ , and  $5 \text{ km}$ . (b)-(d) Frequency distribution of the ratio of the formation rates of ice crystal number produced by heterogeneous freezing. The ratio is defined as  $R_{\Delta x_2, \Delta x_1} = \log(\bar{n}_{het, \Delta x_2} / n_{het, \Delta x_1})$  with the number of formed ice particles at coarser resolution,  $n_{het, \Delta x_1}$ , and the mean number of formed ice particles in the corresponding spatial volume (model grid cells) at finer resolution,  $\bar{n}_{het, \Delta x_2}$ . The results are for the period 12 UTC, 3 April 2014, to 0 UTC, 4 April 2014, for a circular domain with a radius of  $8^\circ$  centered at  $6^\circ\text{W}, 46^\circ\text{N}$ . ICON-ART was initialized with a 15 March 2014 European Centre for Medium-Range Weather Forecasts Integrated Forecast System analysis to spin-up background dust, with daily forecasts from 29 March until 4 April. Domains ranged from global for 40-km resolution to central Europe for 10 and 5-km resolutions.




vertical velocities used to calculate activation with model resolution. Cusack et al. (1999) exploit this scaling to estimate unresolved variance of saturation in developing a cloud parameterization.

Updraft scales extend down to large eddies (tens of meters), so modeling cloud processes depending on vertical velocities requires their sub-grid scales to be parameterized. In climate models, resolutions are still coarser than resolved convection, 5 leaving the possible dependence of climate sensitivity on some aspects of convection also to be parameterized. Realistic vertical velocities for parameterized convection would satisfy an important constraint related to entrainment, detrainment, and convective mixing. Many cumulus treatments in climate models parameterize only mass fluxes and do not provide its factors (vertical velocity, area, and density) independently, but some cumulus parameterizations do, e.g., the parameterizations for shallow (Bretherton et al., 2004) and deep (Donner, 1993) convection in the Geophysical Fluid Dynamics Laboratory Climate 10 Model-3 (Donner et al., 2011). Even for these parameterizations, limited attention has been directed to the realism of their vertical velocities. Donner (1993)'s parameterization was calibrated using observations of convective vertical velocities from the Global Atmospheric Research Program Atlantic Tropical Experiment in the 1970s. Only recently have observations from other field campaigns permitted independent evaluation of the manner in which vertical velocities are calculated in that parameterization. Fig. 5 shows the parameterization captures the basic shape of the updraft profile observed in the Tropical Warm 15 Pool-International Cloud Experiment (TWP-ICE). The updraft speeds for the strongest updrafts are generally within a factor of two of observations with the parameterized updrafts stronger than observed in one case and weaker in the other. (Varble et al., 2014).

In the longer-term future, global models with resolutions fine enough to explicitly resolve deep cumulus clouds are envisioned, eliminating the need for parameterizations of deep cumulus convection. Very short time integrations of models with 20 horizontal resolutions of 0.87 km have already been reported (Miyamoto et al., 2013). In these models, and in limited-domain cloud-system models, the preceding discussions imply a corresponding concern with the realism of their resolved vertical velocities. Evaluations of cloud-system models with horizontal resolutions as fine as 1 km show that many of these models produce vertical velocities that are too strong. These models show some similarities to the parameterized velocities in Fig. 5 (Varble et al., 2014). Vertical velocities in these models can depend strongly on the method used to model their microphysics, 25 and including more physically based microphysics in these models may improve the simulated vertical velocities (Fan et al., 2015). The methods used to model turbulence below even the fine resolutions in these models are also important and offer another path forward (Bogenschutz and Krueger, 2013).

We have claimed that updraft speeds are among the controls on climate forcing and provide an observable constraint related to climate sensitivity. These relationships among vertical velocities, climate forcing, and climate sensitivity emerge through 30 the effects of vertical velocities on physical processes. Resolution dependence of vertical velocities and limited attention to their sub-grid parameterization in models for climate and numerical weather prediction could consequently limit the realism of these models. Taking account of these issues could open promising new paths toward reducing model uncertainties in climate forcing and sensitivity. Sub-grid parameterizations should include vertical velocities where the underlying physical processes



**Figure 5.** Probability distribution functions of vertical velocities in deep convective updrafts using the cumulus parameterization in Donner et al. (2011) and observed using dual-Doppler radar (Varble et al., 2014). The panels show two periods in January 2006 for which radar observations are available, along with model-generated updrafts for the same periods. Both panels show radar-observed and model-generated updrafts whose vertical extent is at least 5 km with a minimum speed of  $1 \text{ m s}^{-1}$  or larger. The percentiles are the fraction of the updrafts with vertical velocities less than the plotted velocities, as functions of height.

depend on them. Some parameterizations already do so (e.g., Fig. 5; Donner (1993), Donner et al. (2011), and Chikira and Sugiyama (2010) for deep cumulus convection; Golaz et al. (2002) and Bretherton et al. (2004) for boundary-layer clouds). Aerosol and cloud microphysical processes depend nonlinearly on vertical velocities, and physically based parameterizations which include these dependencies will fail if driven even by realistically averaged velocity fields, which may even smooth away such phenomena as small-scale updrafts necessary for aerosol activation. Until models explicitly resolve these scales, it is worth investigating simple scaling of resolved vertical velocities when using them to drive parameterizations. Fig. 3 implies that resolved vertical velocities will scale with a power-law dependence on resolution of  $\Delta x^{-2/3}$ , suggesting a scaling of vertical velocity for a physical parameterization  $(\frac{\Delta x_{param}}{\Delta x})^{-2/3}$ .  $\Delta x_{param}$  is the scale at which a parameterization is physically realistic or becomes “scale aware” by design. For example, for a cloud system with uniform mesoscale ascent at a physical scale of around 10 km in a model with a resolution of 100 km, the vertical velocities for aerosol activation would be scaled from 100 km to 10 km. This approach would further have the advantage of requiring physical parameterizations to identify the scales at which their physics applies. It would also introduce at least a crude “scale awareness” into parameterizations which lack them. The vertical-velocity scaling also implies that resolved, advected fields, such as water vapor, will also have



resolution dependencies. So, scaling vertical velocities used in parameterizations would introduce consistency issues not easily reconciled, underscoring the importance of pursuing resolutions as close as possible to the physical scales of climate processes.

## 5 Outlook and Challenges

The atmosphere sustains a broad spectrum of vertical motions. The magnitudes of upward motions are an important control 5 on the formation of liquid and ice particles in clouds and, consequently, anthropogenic climate forcing by cloud-aerosol interactions. In climate models, convective entrainment and mixing, which are among the key governors of vertical velocity, are related to climate sensitivity. Recent observations of convective vertical velocities could provide an important observational constraint for both anthropogenic climate forcing and climate sensitivity. Insightful analysis of these observations in the context of climate models could reduce two of the major uncertainties in climate change.

10 In climate models, both resolved and sub-grid vertical velocities are important. Modeling strategies should include (1) recognizing the dependence of vertical velocities on resolution and exploiting this dependence to scale resolved vertical velocities to process scales, (2) parameterizing sub-grid vertical velocities where the cloud-scale processes depend on them, and (3) explicitly taking into account scale and scale dependence for physical processes, both for resolved and parameterized processes. Preliminary observational studies and new approaches in parameterization are providing the means for doing so.

15 Neither high-resolution climate models, nor those with advanced parameterizations, will satisfactorily deal with connections among vertical velocities, climate forcing, and climate sensitivity if not grounded in realistic cloud-resolving and large-eddy models. Current cloud-resolving models require further development, based on preliminary comparison of their vertical velocities with observations. An intersection of resolution, microphysics, and turbulence will likely bring these models and observations into agreement. A high research priority is to focus on these issues in cloud-resolving and large-eddy modeling.

20 Observationally, field observations of atmospheric updrafts at cloud scale remain limited and should be expanded to sample a wider range of synoptic settings. If a high degree of confidence can be established in cloud-resolving and large-eddy models based on these observational studies, the models can be used to explore the many contexts in which cloud systems develop in the climate system.

Both anthropogenic climate forcing by aerosols and climate sensitivity are extremely difficult problems which have challenged 25 climate scientists for decades. We raise the prospect here that new observations of an element of the climate system, its spectrum of updrafts on all scales, could provide important new clues. The argument that updrafts are a key to unlocking climate forcing and sensitivity is nuanced (especially for sensitivity). It is critical to determine the extent to which vertical velocities control climate forcing and constrain climate sensitivity. If vertical velocities were indeed to provide a breakthrough on this problem, they would do so through a satisfying unification of observational, theoretical, and modeling across the scales 30 and phenomena that comprise the very broad field of contemporary atmospheric science.



*Acknowledgements.* Analysis of convective vertical velocities has been supported by the US Department of Energy Atmospheric System Research program through interagency agreement DE-SC0004534-NOAA. Reviews by Levi Silvers and Yi Ming and comments by Graham Feingold on controls on drop activation in warm clouds are appreciated. Charles Seman participated in figure preparation.



## References

- Bogenschutz, P. and Krueger, S.: A simplified PDF parameterization of sub-grid scale clouds and turbulence for cloud-resolving models, *J. Adv. Model. Earth Syst.*, 5, 195–211, doi: 10.1002/jame.20018, 2013.
- Bretherton, C., McCaa, J., and Grenier, H.: A new parameterization for shallow cumulus parameterization and its application to marine 5 subtropical cloud-topped boundary layers, *Mon. Wea. Rev.*, 132, 864–882, 2004.
- Chikira, M. and Sugiyama, M.: A cumulus parameterization with state-dependent entrainment rate. Part I: Description and sensitivity to temperature and humidity profiles, *J. Atmos. Sci.*, 67, 2171–2193, doi: 10.1175/2010JAS3316.1, 2010.
- Cho, J. and Lindborg, E.: Horizontal velocity structure functions in the upper troposphere and lower stratosphere. 1. Observations, *J. Geophys. Res.*, 106, 10,223–10,232, 2001.
- Cusack, S., Edwards, J., and Kershaw, R.: Estimating the subgrid variance of saturation, and its parameterization for use in a GCM cloud scheme, *Q. J. R. Meteorol. Soc.*, 125, 3057–3076, 1999.
- Donner, L.: A cumulus parameterization including mass fluxes, vertical momentum dynamics, and mesoscale effects, *J. Atmos. Sci.*, 50, 889–906, 1993.
- Donner, L. J., Wyman, B., Hemler, R., Horowitz, L., Ming, Y., Zhao, M., Golaz, J.-C., Ginoux, P., and Coauthors: The dynamical core, 15 physical parameterizations, and basic simulation characteristics of the atmospheric component of the GFDL global coupled model CM3, *J. Climate*, 24, 3484–3519, doi: 10.1175/2011JCLI3955.1., 2011.
- European Centre for Medium-Range Weather Forecasts: WCRP and WWRP THORPEX YOTC (Year of Tropical Convection) Project, Single Parameter 6-Hourly Pressure Level Analysis Time Series, Transformed to a Regular 1600 by 800 (N400) Gaussian Grid, 2012.
- Fan, J., Liu, Y.-C., Xu, K.-M., North, K., Collis, S., Dong, X., Zhang, G., Chen, Q., and Coauthors: Improving representation of convective 20 transport for scale-aware parameterization: 1. Convection and cloud properties simulated with spectral bin and bulk microphysics, *J. Geophys. Res. Atmos.*, 120, 3485–3509, doi:10.1002/2014JD022142, 2015.
- Feingold, G.: Modeling of the first indirect effect: Analysis of measurement requirements, *Geophys. Res. Lett.*, 30, doi:10.1029/2003GL017967, 2003.
- Forster, P. M., Andrews, T., Good, P., Gregory, J., Jackson, L., and Zelinka, M.: Evaluating adjusted forcing and model spread for historical 25 and future scenarios in the CMIP5 generation of climate models, *J. Geophys. Res. Atmos.*, 118, 1139–1150, doi: 10.1002/jgrd.50174, 2013.
- Golaz, J.-C., Larson, V., and Cotton, W.: A PDF based model for boundary layer clouds. Part I: Method and model description, *J. Atmos. Sci.*, 59, 3540–3551, 2002.
- Kay, J. and Wood, R.: Timescale analysis of aerosol sensitivity during homogeneous freezing and implications for upper tropospheric water 30 vapor budgets, *Geophys. Res. Lett.*, 35, L10 809, doi: 10.1029/2007GL032628., 2008.
- Kiehl, J. T.: Twentieth century climate model response and climate sensitivity, *Geophys. Res. Lett.*, 34, L22 710, doi:10.1029/2007GL031383, 2007.
- Klocke, D., Pincus, R., and Quaas, J.: On constraining estimates of climate sensitivity with present-day observations through model weighting, *J. Climate*, 24, 6092–6099, doi:10.1175/2011JCLI4193.1, 2011.
- Ma, P.-L., Rasch, P., Wang, M., Wang, H., Ghan, S., Easter, R., Gustafson, W., Liu, X., and Coauthors: How does increasing horizontal resolution in a global climate model improve the simulation of aerosol-cloud interactions?, *Geophys. Res. Lett.*, 42, 5058–35 Ma, P.-L., Rasch, P., Wang, M., Wang, H., Ghan, S., Easter, R., Gustafson, W., Liu, X., and Coauthors: How does increasing horizontal resolution in a global climate model improve the simulation of aerosol-cloud interactions?, *Geophys. Res. Lett.*, 42, 5058–5065, doi:10.1002/2015GL064183, 2015.



McFiggans, G. and Co-Authors: The effect of physical and chemical aerosol properties on warm cloud droplet activation, *Atmos. Chem. Phys.*, 6, 2593–2649, 2006.

Miyamoto, Y., Kajikawa, Y., Yoshida, R., Yamaura, T., Yashiro, H., and Tomita, H.: Deep moist atmospheric convection in a subkilometer global simulation, *Geophys. Res. Lett.*, 40, doi:10.1002/grl.50944, 2013.

5 Rauscher, S., O'Brien, T., Piani, C., Coppola, E., Giorgi, F., Collins, W., and Lawston, P.: A multi-model intercomparison of resolution effects on precipitation: Simulations and theory, *Clim. Dyn.*, p. in press., 2016.

Rieger, D., Bangert, M., Bischoff-Gauss, I., Förstner, J., Lundgren, K., Reinert, D., Schröter, J., Vogel, H., Zängl, G., Ruhnke, R., and Vogel, B.: ICON-ART 1.0 - A new online-coupled model system from global to regional scale, *Geoscientific Model Development*, 6, 1659–1676, doi:10.5194/gmd-8-1659-2015, <http://www.geosci-model-dev.net/8/1659/2015>, 2015.

10 Rosenfeld, D., Wood, R., Donner, L., and Sherwood, S.: Aerosol cloud-mediated radiative forcing: Highly uncertain and opposite effects from shallow and deep clouds, in: *Climate Science for Serving Society: Research, Modeling, and Prediction Priorities*, Springer, 2013.

Rosenfeld, D., Sherwood, S., Wood, R., and Donner, L.: Climate effects of aerosol-cloud interactions, *Science*, 343, 379–380, doi: 10.1126/science.1247490, 2014.

15 Rougier, J., Sexton, D., Murphy, J., , and Stainforth, D.: Analyzing the climate sensitivity of the HadSM3 climate model using ensembles from different but related experiments, *J. Climate*, 22, 3540–3557, doi:10.1175/2008JCLI2533.1, 2009.

Sanderson, B., Shell, K., and Ingram, W.: Climate feedbacks determined using radiative kernels in a multi-thousand member ensemble of AOGCMs, *Clim. Dyn.*, 35, 1219–1236, 2010.

Sherwood, S., Bony, S., and Dufresne, J.-L.: Spread in model climate sensitivity traced to atmospheric convective mixing, *Nature*, 505, 37–42, doi:10.1038/nature12829, 2014.

20 Stainforth, D., Alna, T., Christensen, C., Collins, M., Faull, N., Frame, D., Kettleborough, J., Knight, S., Martin, A., and Co-Authors: Uncertainty in predictions of the climate response to rising levels of greenhouse gases, *Nature*, 433, 403–406, 2005.

Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., and Coauthors: Summary for policymakers, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. Cambridge University Press, Cambridge/New York, 2013.

25 Su, H., Jiang, J., Zhai, C., Shen, J., Neelin, J., Stephens, G., and Yung, Y.: Weakening and strengthening structures in the Hadley Circulation change under global warming and implications for cloud response and climate sensitivity, *J. Geophys. Res.*, 119, 5787–5805, doi:10.1002/2014JD021642, 2014.

Varble, A., Zipser, E., Fridland, A., Zhu, P., Ackerman, A., Chaboureau, J.-P., Collis, S., Fan, J., and Co-Authors: Evaluation of cloud-resolving and limited area model intercomparison simulations using TWP-ICE observations. 1. Deep convective updraft properties, *J. Geophys. Res. Atmos.*, 119, 13,891–13,918,doi: 10.1002/2013JD021371, 2014.

Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOahedral Non-hydrostatic) modeling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Q. J. R. Meteorol. Soc.*, 141, 563–579, 2015.

Zelinka, M., Klein, S., and Hartmann, D.: Computing and partitioning cloud feedbacks using cloud property histograms: Part I. Cloud radiative kernels, *J. Climate*, 25, 3715–3735, doi:10.1175/JCLI-D-11-00248.1, 2012.

35 Zhao, M.: An investigation of the connections among convection, clouds, and climate sensitivity in a global climate model, *J. Climate*, 27, doi:10.1175/JCLI-D-13-00145.1, 2014.