

Answer to Reviewer 1

We thank reviewer 1 for his careful review and the useful annotated paper. We apologize for the large number of English errors and the new manuscript has been carefully edited by one of the English speaking co-author. Find below the answers to the questions raised by the reviewer.

The paper meets the ACP criteria and can be published after minor revision and technical corrections. An important feature of this paper is that it is not clear if only nighttime tropospheric O₃ measurements are performed. The authors have to clarify this everywhere needed in the manuscript (i.e. in Fig. 17, it is not mentioned; and also in all Figs., this should be mentioned).

There is no figure 17 but we specify now that the ozone lidar performances are given for daytime measurements in section 2. It is true for both the ALTO lidar and the NASA airborne lidar. Regarding the questions about daytime versus nighttime lidar measurements. All the aircraft flights are during the day and the ozone measurement performances are given for daytime measurement. It is said more clearly in section 2.

The annotated paper is attached to this report. The main remarks in the annotated are:

- Need to rephrase line 18-20 p. 2 and line 6-7 p. 4

The two new pieces of text are now:

“ The ozone distributions over North America have been discussed for the spring period at high latitude using the Tropospheric Ozone Production about the Spring Equinox (TOPSE) and Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) data set in several publications (Browell 2003, Wang 2003, Olson 2012, Koo 2012) showing: (i) frequent occurrence of ozone depletion events (ODE) in the planetary boundary layer (PBL), (ii) a net O₃ photochemical production rate equal to zero throughout most of the troposphere, (iii) a latitudinal increase of the tropospheric ozone concentrations due to transport from mid-latitudes and Stratosphere-Troposphere Exchange (STE)”

“The system is fully described in Ancellet and Ravetta (1998), and the instrument performances for different examples of daytime airborne measurements are discussed in Ancellet and Ravetta (2003) (urban pollution in the boundary layer, ozone in the UTLS, long-range transport in the free troposphere)”

- Need of a rephrased conclusions showing more clearly the outcome (p 21 line 20 and 26)

The rephrased conclusion is now:

“The purpose of this work is to provide a complete picture of Arctic ozone using measurements available during the summer 2008 POLARCAT campaigns over Canada and Greenland, where 3 aircraft were deployed and 7 ozonesonde stations intensified their ozone vertical profiling. This is the first case of such complete temporal and geographical coverage. We take advantage of the large number of airborne lidar profiles (representing 67% of the O₃ measurements over Canada and 26% over Greenland). The measured ozone climatology established in this paper can also be used for future model evaluation at high latitudes. For example, in our work, while a good correspondence of the measured O₃ vertical and latitudinal distribution is found with model results from WRF-Chem, a negative O₃ bias of -6 to -15 ppbv between the model and the observations is found in the free troposphere over 4 km, especially over Canada. This deficiency is partly related to the WRF-Chem model stratospheric ozone initialization.

The WRF-Chem model simulation is also used to discuss the relative influence of tropospheric ozone sources at high latitude in summer. Ozone average concentrations are of the order of 65 ppbv at altitudes above 4 km both over Canada and Greenland, while they are less than 50 ppbv in the lower troposphere. For Canada, the analysis of the modeled CO distribution and the weak correlation (< 30%) of O₃ and PV suggest that stratosphere-troposphere exchange (STE) is not the major contribution to tropospheric ozone at latitudes less than 70° N, where transport of North

American biomass burning (BB) emissions took place during the 2008 summer. Conversely, significant STE is found over Greenland according to the better O₃ versus PV correlation (> 40%) and the higher value of the 75th PV percentile. This is related to the persistence of cyclonic activity during the summer over Baffin Bay.

A weak negative latitudinal summer ozone gradient of -6 to -8 ppbv is found over Canada in the mid-troposphere between 4 and 8 km because the O₃ photochemical production from BB emissions mainly takes place at latitudes less than 65° N, while STE plays a larger role at latitudes higher than 70° N. A positive ozone latitudinal gradient of 12 ppbv is observed in the same altitude range over Greenland not because of an increasing latitudinal influence of STE, but because of different long-range transport from multiple mid-latitude sources (North America, Europe and even Asia for latitudes higher than 77° N).

For the Arctic latitudes (> 80° N), free tropospheric O₃ concentrations are related to a mixture of stratospheric O₃ transport across the tropopause and Asian pollution, as already suggested by Roiger et al. (2011) using a case study of aircraft observations in the Arctic. Our study using more ozone measurements leads to the same conclusions.”

The abstract has been also changed according to the changes made in the conclusions.

- Need to quantify the differences between the backtrajectory analysis for zone 3 and 4 (p 20 line 16)

The calculation of the fraction of air masses coming from North America (-150W/-60W, 50N/70N) has been added to quantify the role of the North American surface emissions on ozone in zone 3 and 4. The fractions of air masses coming from North America are 76% and 72% for zone 3 and 4, respectively. The fraction for zone 6 and 7 over Greenland is smaller (< 30%) and support the statement about the multiple mid-latitude sources controlling ozone in zone 7. This is now included in the text (p. 21 line 21-22 and p. 22 line 2-4)

- Need to remove the sentence describing the WRF-Chem model characteristics for aerosol modelling. (P. 7 line 18)

Done

Answer to Reviewer 2

We thank reviewer 2 for his review. We apologize for the large number of English errors and the new manuscript has been carefully edited by one of the English speaking co-author. Find below the answers to the questions raised by the reviewer.

The paper provides a detailed description of tropospheric ozone measurements during the POLARCAT campaign in summer 2008, and a series of analysis of its variability and possible sources affecting the variability by using WRF-Chem chemistry-transport model. The description is intensive and the analysis is generally sound, reaching some interesting conclusions. I found the paper be a nice piece of work contributing to better understanding of sources and processes affecting the summertime ozone at high-latitudes. The paper is well within the scope of ACP, and can eventually be published. However, I found some descriptions/illustrations are redundant. There are many errors in English. Some figures look preliminary. These made me difficult to follow what the authors are trying to tell us. So, I would suggest some technical suggestions that should be addressed before publication.

Figures 5, 7, and 9: The same data are plotted in both linear and logarithmic scales. I doubt if the authors really need logarithmic plots, as they do not discuss much on the log plots. The log plots seem redundant to me.

The log plots have been removed as suggested by the reviewer because they are not specifically discussed in the text. We also agree with the reviewer that some plots needed improvment. The measured/modeled ozone scatterplot (Figure 5) has been changed to better distinguish tropospheric and stratospheric data (now in white) and to include the O3 concentration unit in the axis label. The back trajectory plots (Fig. 11 and 12) include now the name of the corresponding zone shown in Fig. 6 and 8 and unit for the altitude color scale. A new figure is created (Fig. 10) for the map of the MODIS aerosol optical depth, including also the map of the CFS fire counts distribution.

There are many errors in English:

L1: The goals of the paper are ...

L7: Ozone, CO and . . . is too much detail in Abstract, and can be removed.

L10: The average ozone concentrations are 65 ppbv ...

L14: ... modeled CO ...

L18: ... ozone gradient of -6 to -8 ppbv . . .

P7, L17: MEGAN (Model of ...)

Figure 2 caption: Intercomparison of ozone measurements ...

We thank the reviewer for his help to edit the manuscript. The English-speaking co-authors have also corrected many grammar errors.

Captions in other figures: Measured O3 (ppbv), Modeled O3 (ppbv) - need units!

Done in Fig. 5, 7, 9

Figures 3 and 4: I would make difference plots between observation and model, to illustrate where in height and latitude the model is good or bad. This is not necessary but please consider.

We understand the reviewer remark but the measurement/model comparison does not focus on small scale differences. The goal of the comparison is to check that the main latitudinal and vertical gradients are well reproduced in the WRF-Chem simulations. This is why we do not wish to produce a detailed 2-D plot of the modeled/measured ozone differences which will show mainly model/measurement spatial/temporal mismatches. We believe that the differences can be discussed using two different 2D ozone plots for the model and the data and the detailed statistical informations about model /measurement differences provided in table 5 for each region selected for the vertical and latitudinal gradient analysis .

Analysis of the latitudinal variability of tropospheric ozone in the Arctic using the large number of aircraft and ozonesonde observations in early summer 2008

Gerard Ancellet¹, Nikos Daskalakis¹, Jean Christophe Raut¹, David Tarasick², Jonathan Hair³, Boris Quennehen¹, François Ravetta¹, Hans Schlager⁴, Andrew J. Weinheimer⁵, Anne M. Thompson⁶, Bryan Johnson⁷, Jennie L. Thomas¹, and Katharine S. Law¹

¹LATMOS/IPSL, UPMC Univ. Paris 06 Sorbonne Universités, UVSQ, CNRS, Paris, France

³NASA Langley Research Center, Hampton, VA, USA

²Environment and Climate Change Canada, Downsview, ON, Canada

⁴Institut für Physik der Atmosphäre, DLR, Oberpfaffenhofen, Germany

⁵NCAR, Boulder, CO, USA

⁶NASA/GSFC, Greenbelt, MD, USA

⁷NOAA/Earth System Research Laboratory (ESRL), Boulder, CO, USA

Correspondence to: Gerard Ancellet : gerard.ancellet@upmc.fr

Abstract. ~~The goal of the paper are to: (1) present tropospheric ozone climatologies in summer. During the 2008 based on a large amount of measurements, during the International Polar Year when, the Polar Study using Aircraft, Remote Sensing, Surface Measurements, and Models of Climate Chemistry, Aerosols, and Transport (POLARCAT) campaigns were conducted (2) investigate the processes that determine concentrations in two different regions (Canada and Greenland) that were thoroughly studied using campaign, conducted in summer over Greenland and Canada, produced a large number of measurements from 3 aircraft and 7 ozonesonde stations. This paper provides an integrated analysis. Here we present an observation integrated analysis is based on three different types of O₃ measurements: airborne lidar, airborne UV absorption or chemiluminescence measurement, and intensified Electrochemical Concentration Cell (ECC) ozonesonde profiles. Discussion of these observations and the discussion of the latitudinal and vertical variability of tropospheric ozone north of 55°N during this period is performed using with the aid of a regional model (WFR-Chem). Ozone, CO and potential vorticity (PV) distributions are extracted from the simulation at the measurement locations.~~ The model is able to reproduce the O₃ latitudinal and vertical variability but with a negative O₃ bias of 6-15 ppbv is found in the free troposphere over above 4 km, especially over Canada.

Ozone average concentrations are of the order of 65 ppbv at altitudes For Canada, large average CO concentrations in the free troposphere above 4 km both over Canada and Greenland, while they are less than 50 ppbv in the lower troposphere. The relative influence of stratosphere-troposphere exchange (STE) and of ozone production related to the local biomass burning (BB) emissions is discussed using differences between average values of , CO and PV for Southern and Northern Canada or Greenland and two vertical ranges in the troposphere: 0-4 km and 4-8 km. For Canada, the model CO distribution and the (> 130 ppbv) and the weak correlation (< 30%) of O₃ and PV suggests that stratosphere-troposphere exchange (STE) is not the major contribution to average tropospheric ozone at latitudes less than 70°N, due to the fact that local biomass burning (BB) emissions were significant during the 2008 summer period. Conversely over Greenland, significant STE is found over

Greenland according to the better O₃ versus PV correlation (> 40%) and the higher values of the 75th PV percentile. It is related to the persistence of cyclonic activity during the summer over Baffin Bay.

- ~~A~~ Using differences between average concentration above Northern and Southern Canada, a weak negative latitudinal summer ozone gradient of -6 to -8 ppbv is found ~~over Canada~~ in the mid troposphere between 4 and 8 km. This is attributed to an efficient O₃ photochemical production ~~due to the~~ from BB emissions at latitudes less than 65°N, while the STE contribution is more homogeneous in the latitude range 55°N to 70°N. A positive ozone latitudinal gradient of 12 ppbv is observed in the same altitude range over Greenland not because of an increasing latitudinal influence of STE, but because of different ~~long range~~ long-range transport from multiple mid-latitude sources (North America, Europe and even Asia for latitudes higher than 77°N).
- 10 For the Arctic latitudes (> 80°N), free tropospheric O₃ concentrations during summer 2008 are related to a mixture of stratospheric O₃ transport across the tropopause and Asian pollution.

1 Introduction

- Ozone concentrations are still increasing in many locations in the Northern Hemisphere mostly due to an increase in Asian precursor emissions (Lin et al., 2015; Cooper et al., 2010). Since tropospheric ozone is an effective greenhouse gas with a relatively long lifetime its main impact on climate and air quality is within mid-latitudes regions. Several studies have shown that ozone ~~makes also~~ makes an important contribution to the Arctic surface temperature increases due to direct local warming in Arctic as well as heat transport following warming due to ozone at mid-latitudes (Shindell, 2007; Shindell et al., 2009; AMAP, 2015). The Arctic ozone budget still requires better quantification and is complicated by the interplay between the long-range transport, including the downward transport of stratospheric ozone ~~the~~ (Hess and Zbinden, 2013; Walker et al., 2012), removal of boundary layer O₃ due to halogen chemistry especially in springtime (Simpson et al., 2007; Abbatt et al., 2012), and ~~the~~ photochemical production due to local sources like boreal forest fires (Stohl et al., 2007; Thomas et al., 2013), NO_x enhancement from snowpack emissions (Honrath et al., 1999; Legrand et al., 2009), local summertime production from peroxyacetyl nitrate (PAN) decomposition (Walker et al., 2012) or ship emissions (Granier et al., 2006).
- ~~Ozone distributions~~ The O₃ distribution over North America ~~have~~ has been discussed for the spring period at high latitude using the Tropospheric Ozone Production about the Spring Equinox (TOPSE) and Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) data set in several publications (Browell et al., 2003; Wang et al., 2003; Olson et al., 2012; Koo et al., 2012) showing: (i) frequent occurrence of ozone depletion ~~event~~ events (ODE) ~~but not extending to the free troposphere, a net in the planetary boundary layer (PBL),~~ (ii) a net O₃ photochemical production rate ~~of~~ equal to zero throughout most of the troposphere, ~~while transport from mid-latitude~~ (iii) a latitudinal increase of tropospheric O₃ concentrations due to transport from mid-latitudes and Stratosphere-Troposphere Exchange (STE) ~~explain the ozone increase with latitude.~~

For the summer period ozone photochemical production is expected according to the numerous studies conducted at mid-latitudes (Crutzen et al., 1999; Parrish et al., 2012), but little attention ~~was~~ has been given to the high latitude distribution

during this season. During the ARCTAS-B (Jacob et al., 2010) and Polar Study using Aircraft, Remote Sensing, Surface Measurements, and Models of Climate Chemistry, Aerosols, and Transport (POLARCAT) campaigns (Law et al., 2014), many ozone measurements ~~have been~~ were carried out over Canada and Greenland from 15 June to 15 July 2008 ~~in addition to regular ozonesonde observation by the Canadian network~~ via aircraft and regular ozone soundings. This allows a detailed analysis of the ozone regional distribution at high latitudes between 55°N and 90°N and a discussion about the relevant ozone sources driving the summer ozone values. So far, knowledge of the relative influence of the main ozone summer sources ~~was~~ has been mainly derived from modeling studies, e.g. summer simulations of global model simulation of the ozone source attribution (Wespes et al., 2012; Walker et al., 2012; Monks et al., 2015), or regional modeling of biomass burning case studies for North American fires (Thomas et al., 2013) or Asian fires (Dupont et al., 2012).

In order to interpret the measurements, we use a hemispheric simulation performed using the regional chemical transport model WRF-Chem (Grell et al., 2005; Fast et al., 2006). WRF-Chem ~~predicts~~ produces simultaneously a meteorological forecast, including the dynamics of the ~~UTLS region, as well as emissions and upper troposphere and lower stratosphere (UTLS) region, with online~~ chemistry to predict ozone (and other trace gas and aerosol) concentrations. Here, we use the WRF-Chem model to investigate dynamics that determine ozone concentrations as a function of latitude, ~~referred to as latitudinal gradients, including stratosphere-troposphere exchanges including STE~~ processes, which can bring ~~high ozone~~ high-ozone air from the stratosphere into the upper troposphere in the Arctic. In addition, we use the model to compare directly predicted and measured ozone in summer 2008 and use ~~a pollution tracer (i.e. carbon monoxide, CO)~~ CO as a tracer to separate air influenced by ~~pollution~~ surface emissions and subsequent ozone formation via photochemistry from air influenced by stratosphere-troposphere mixing processes.

The objectives of this paper are thus twofold : (i) to establish the summer tropospheric ozone latitudinal variability over Greenland and Canada based on the large number of ozone measurements available in the free troposphere during June/July 20008, and (ii) to explore the role of photochemistry and ~~stratosphere/troposphere exchange~~ STE in the observed high latitude ozone distribution by extracting the potential vorticity (PV) and CO distribution from a 2-month WRF-Chem regional model simulation. The WRF-Chem simulation also provides the modeled ozone distribution to verify the coherence between the modeled PV and CO with the observed ozone distributions. The ozone data set and the WRF-Chem simulation are described in ~~section~~ sections 2 and 3, respectively. The model ~~and~~ versus measurement ozone comparison is discussed in section 4, while the latitudinal ~~distribution~~ distributions of ozone, CO and PV are presented in section 5.

2 Summer 2008 ozone data set

We use three ~~types of ozone measurements~~ different ozone measuring instrument to build the data set considered in this work: airborne lidar, in-situ aircraft ozone analyzer, and electrochemical concentration cell (ECC) ~~ozonesondes~~ ozonesonde. Two airborne ozone Differential Absorption Lidars (DIAL) are ~~considered~~ considered: the NASA-DC8 instrument and the ALTO lidar on the French ATR-42. In-situ ozone ~~monitors~~ analyzers have been installed onboard 3 different aircraft: the NASA DC-8, the DLR Falcon-20 and the French ATR-42. We also considered 7 ground-based stations ~~where ozonesondes are~~

~~regularly-launched~~, that participated to the ARC-IONS campaign (Tarasick et al., 2010), over Canada (latitudes $> 50^{\circ}\text{N}$ in the longitude range between -70°W and -160°W) and over Greenland (latitudes $> 55^{\circ}\text{N}$ in the longitude range between -60°W and -20°W). The times and positions of the June/July 2008 measurements included in our study are given in Table 1 for the aircraft observations and in Table 2 for the ozonesondes. The aircraft flight paths over Canada and Greenland are shown in

5 Fig.1 and the measurements are always performed during daytime.




Figure 1. Horizontal distribution of the aircraft positions (blue) and sounding locations (red) over the selected Canada (left panel) and Greenland regions (right panel).

The German DLR Falcon-20 was based in Greenland and used an UV absorption instrument (Thermo Environment Instruments, TEI49C) to measure O_3 with an uncertainty of ± 2 ppbv ($\pm 5\%$ of the signal) (Schlager et al., 1997; Roiger et al., 2011). The ATR-42 aircraft was also based in Greenland and the ozone measurements ~~are were~~ made using a similar instrument (TEI49-103) calibrated against a NIST (National Institute of Standards and Technology) referenced O_3 calibrator ~~Model49PS~~

10 Model 49PS at zero, 250, 500 and 750 ppbv (Marenco et al., 1998). A 4-ppbv negative bias related to ~~a-slight~~ O_3 loss in the ATR-42 air inlet (e.g. 10% for 40 ppbv and 5% for 80 ppbv) has been corrected in this study. On 14 July 2008, the comparison ~~with-the-of the ATR-42 and the~~ DLR-Falcon ~~at-two-ozone data at two altitude~~ levels over Greenland (near 67°N) shows an uncertainty better than 2 ppbv (Fig.2). The ~~American~~-NASA DC-8 (Weinheimer et al., 1994) O_3 measurements ~~are were~~ made with a chemiluminescence technique, with the instrument calibrated by additions of O_3 determined by UV optical absorption at

15 254 nm. Uncertainties in the DC-8 O_3 data are typically ± 2 ppbv ($\pm 5\%$ of the signal). The ~~DC8-NASA DC-8~~ flew mainly over Canadian forest fire regions in summer (Jacob et al., 2010). The comparison ~~shows-a-very-good-precision-but-a-slight-4-ppbv positive-difference-with-the-between the NASA DC-8 and the~~ DLR-Falcon ~~data-(Fig.2)~~ observed during the O_3 ~~data during~~

an aircraft intercomparison flight on July 9th over Northern Greenland ~~shows very good precision, but a 4-ppbv positive difference (Fig 2).~~ None of the DLR-Falcon and NASA ~~DC8-DC-8~~ data were corrected.

The ~~ATR-42 ozone Differential Absorption Lidar (DIAL) instrument~~ ~~ALTO lidar~~ was mounted in a zenith-viewing mode making ozone vertical profiles above the aircraft, ~~thus~~ limiting the number of ~~data available~~ ~~O₃ data available at altitudes~~ below 3 km. The lidar measurement altitude range is of the order of 6 km above the aircraft ~~altitude flight level~~ with a 300 m vertical resolution and ~~2-minute temporal resolution corresponding~~ a 10 km horizontal resolution ~~-(i.e. a 2 minute integration time).~~ The system is ~~fully~~ described in Ancellet and Ravetta (1998), ~~while performance during various airborne the applications is given and the instrument performances for different examples of daytime airborne measurements are discussed~~ in Ancellet and Ravetta (2003) ~~where several~~ ~~(urban pollution in the boundary layer, ozone in the UTLS, long-range transport in the free troposphere).~~ ~~Several~~ comparisons with in situ measurements (ECC ozonesonde ~~of and~~ airborne UV photometer) show no specific biases in clear air measurements. Measurements taken near clouds or thick aerosol layers are not included here since corrections of systematic errors related to aerosol interference ~~are unreliable. It corresponds to become very large~~ (Papayannis et al., 1990). ~~These represent~~ 20% of the lidar profiles recorded during the campaign.

The NASA-DC8 Ozone DIAL system and configuration implemented during the campaign is described by Richter et al. (1997). The instrument provides simultaneous zenith and nadir profiles to cover the troposphere and lower stratosphere. The ~~measurement resolution for the archived data~~ ~~O₃ measurement resolution~~ is 300 m in the vertical and approximately 70 km (3 min) in the horizontal ~~for daytime measurements.~~ On all field experiments, the airborne DIAL O₃ measurements are compared with in situ O₃ measurements made on the DC-8 during ascents, descents, and spirals ~~and,~~ ~~while~~ comparisons to ozonesondes during coincident overflights were ~~also~~ conducted. In the troposphere, the DIAL O₃ measurements have been shown to be accurate to better than 10% or 2 ppb, whichever is larger (Browell et al., 1983, 1985). More recently, the precision of the DIAL O₃ measurements during the high-latitude SAGE-III Ozone Loss and Validation Experiment in 2003 (SOLVE II) were found to be better than 5% from near surface to about 24 km and the accuracy was found to be better than 10% in comparison with ~~the~~ Ny-Ålesund lidar, ~~ozonesonde~~ ~~ozonesondes~~, and in situ DC-8 measurements (Lait et al., 2004). DIAL and Microwave Limb Sounder (MLS) O₃ measurements from the Polar Aura Validation Experiment (PAVE) in 2005 were found to agree within 7% across the 12-24 km altitude range (Froidevaux et al., 2008). Comparisons between DIAL and MLS were also examined in the upper troposphere and lower stratosphere (215-100 hPa region) from data obtained during the International Intercontinental Chemical Transport Experiment (INTEX-B) field experiment, and these results show good agreement in ~~lower stratosphere the lower stratosphere,~~ with decreasing performance of the MLS measurements into the troposphere (Livesey et al., 2008).

The ozonesonde monitoring was intensified in 2008 over North America in the framework of the ARCIONS initiative and the characteristics of the ozone measurements are fully described in Tarasick et al. (2010). Nearly daily sounding ~~have been~~ ~~were~~ made during the aircraft flight period at 4 stations over Canada ~~in the latitude band between 53°N and 62°N. Only weekly sounding are made in,~~ ~~and at Summit, Greenland, while weekly soundings were made at~~ the high latitude stations Alert and Resolute.

The latitudinal cross sections of the ozone data set selected in this study are shown in Fig. 3 considering two domains to produce a bidimensional latitude-altitude plot over Canada (-160°W to -70°W) and Greenland (-70°W to -20°W). The




Figure 2. ~~Ozone intercomparison~~ Intercomparison of O_3 measurements in ppbv during two wing-tip to wing-tip flights over Greenland between the ATR-42 (blue) and the DLR-Falcon (red) on 14 July (left panel) and between the NASA-DC8 (black) and the DLR-Falcon (red) on 9 July (right panel). The aircraft altitude changes are also shown in m (multiply scale by 50 on right panel and by 100 on left panel)

- horizontal distributions of the data set used for producing the latitude-altitude plots are shown in Fig. 1. For latitudes higher than 80°N , the data comes from a limited number of observations: 8 sondes from Alert and 3 DC-8 flights from 8 July to 10 July. The locations of the ozonesonde stations are shown as red vertical bars in the latitudinal cross sections. In the troposphere at altitudes less than 8 km over Canada, lidar, in-situ and ozonesonde measurements contribute 67%, 20% and 13% of the ozone data set, respectively, while they correspond to 26% 69% and 5% ~~respectively~~, respectively, over Greenland (see [Table 3 with](#) the number of observations using each technique [in Table 3](#)). The aircraft data (lidar and in-situ) therefore strongly contribute to the high measurement density even though they correspond to a limited number of flying days, i.e. 18 days from 26 June to 18 July. In this work, ozone data are ~~hourly-averaged~~ averaged hourly when they are in the same cell of a grid with a 0.5×0.5 degrees and a 1 km vertical resolution in order to avoid an oversampling of similar air masses. Only hourly averages are ~~accounted for~~ considered in Table 3.
- For both regions similar ozone vertical distributions were observed ~~with low ozone~~: low O_3 mixing ratio ≤ 40 ppbv below 3 km and ~~similar upper-troposphere-lower-stratosphere (UTLS)~~ altitude range of high O_3 concentrations typical of the UTLS in the altitude range 8-11 km between 60°N and 80°N (i.e. red and dark region with ~~ozone~~ O_3 mixing ratio > 150 ppbv). The average O_3 mixing ratio in the altitude range 4-8 km is of the order of 65 ppbv for both regions but the latitudinal O_3 gradients are more visible over Greenland than over Canada. Two mid-tropospheric ozone branches are seen in the latitude band 65°N - 73°N and 78°N - 85°N over Greenland, the ~~first one~~ former being tilted to the ~~South~~ south at 70°N and the ~~second one to the North~~ latter to the north at 80°N .

Table 1. Characteristics of the ensemble of ozone measurements made with airborne instruments during summer 2008

Instrument	Number of flights	Latitude range	Longitude range	Altitude range	Time period
ATR42 in-situ	12	59°N-71°N	-60°W/-20°W	0-7 km	30/6-14/7
ATR42 lidar	12	59°N-71°N	-60°W/-20°W	2-12 km	30/6-14/7
DC8 in-situ	11	45°N-88°N	-132°W/-38°W	0-12 km	26/6-13/7
DC8 lidar	11	45°N-88°N	-132°W/-38°W	0-15 km	26/6-10/7
F20 in-situ	18	57°N-79°N	-65°W/-20°W	0-11 km	30/6-18/7

Table 2. Characteristics of the ECC sounding stations used during summer 2008

Station	Number of ECC	Latitude	Longitude	Altitude range	Time period
Summit	22	72.6°N	-38.5°W	3.2-15 km	6/6-22/7
Alert	8	82.5°N	-62.3°W	0-15 km	4/6-24/7
Resolute	8	74.7°N	-95°W	0-15 km	4/6-30/7
Churchill	16	58.7°N	-94°W	0-15 km	4/6-30/7
Yellowknife	19	62.5°N	-114.5°W	0-15 km	23/6-12/7
Whitehorse	15	60.7°N	-135.1°W	0-15 km	27/6-12/7
Stonyplain	16	53.55°N	-114.11°W	0-15 km	26/6-12/7

3 WRF-Chem model simulation

3.1 Model description

For this study we use the regional Weather Research Forecasting model coupled with Chemistry (WRF-Chem) to study ozone during this period. WRF-Chem is a fully coupled, online meteorology and chemistry and transport mesoscale model. It has been successfully used in Arctic-focused studies in the past (Thomas et al., 2013; Marelle et al., 2015), for both gas phase and aerosol analysis. Initial meteorological conditions and boundaries are from the National Center for Environmental Prediction (NCEP) Global Forecast System (GFS) with nudging applied to temperature, wind and humidity every 6 hours. The simulation uses the Noah Land Surface model scheme with 4 soil layers, the YSU (Yonsei University-) planetary boundary layer (PBL) scheme (Hong et al., 2006), coupled with the MM5 similarity surface layer physics, the Morrison 2-moment (Morrison et al., 2009) microphysics scheme, and the Grell-3D ensemble (Grell and Dévényi, 2002) convective implicit parametrization. The radiation schemes are the Goddard (Max and Suarez, 1994) and Rapid Radiative Transfer Model (Mlawer et al., 1997) for shortwave and longwave radiation respectively. Chemical boundary conditions were taken from the Model For Ozone and Related Chemical Tracers, version 4 (Emmons et al., 2010). For gas phase chemical calculations the CBM-Z (Zaveri and




Figure 3. Latitudinal cross section of the measured ozone mixing ratio in ppbv over Canada (left panel) and over Greenland (right panel). The red bars show the location of ozonesonde stations. Black regions with $O_3 > 160$ ppbv correspond to the stratosphere.

Peters, 1999) chemical scheme is used and aerosols were calculated using the Model for Simulating Aerosol Interactions and Chemistry (Zaveri et al., 2008). The model was run from 15 March to 1 August 2008 using a polar stereographic grid (100×100 km resolution) over a domain that covers most of the Northern Hemisphere, from about 28°N . Vertically 50 hybrid layers up to 50 hPa are used with approximately 10 levels in the first two kilometers. The corresponding vertical resolution ranges from 100 m in the PBL to 500 m in the free troposphere. Anthropogenic emissions ~~used are with a $0.5^\circ \times 0.5^\circ$ spatial resolution were taken from~~ the ECLIPSE (Evaluating the CLimate and Air Quality ImPacts of Short-livEd Pollutants) version 4.0 (Klimont et al., 2013), ~~available in 0.5×0.5 spatial resolution~~. Wildfire emissions were taken from GFED 3.1 (van der Werf et al., 2010), while aircraft/shipping emissions were from the RCP 6.0 scenario (Lee et al., 2009) and (Buhaug et al., 2009), respectively.

Biogenic emissions were calculated online thanks to ~~the~~ the-MEGAN (Model of Emissions of Gases and Aerosols from Nature) (Guenther et al., 2012). ~~WRF-Chem also provides online dust and sea-salt emissions.~~

Using observations from aircraft, surface stations and satellites, atmospheric model simulations of ~~ozone~~- O_3 have been evaluated as part of POLMIP including WRF-Chem (Monks et al., 2015; AMAP, 2015). The model was run using different emissions and gas/aerosol schemes than in the POLMIP simulations, but the POLMIP results are still a good basis to choose WRF-Chem. While all models have deficiencies in reproducing trace gas concentration in the Arctic, WRF-Chem performs better than many models in re-producing tropospheric ~~ozone~~- O_3 and CO, which are used here. Given the advantages of ~~predicting also online also predicting~~ stratosphere-troposphere exchange processes online for this study, WRF-Chem is a good model for interpreting ~~the these~~ ozone climatologies constructed from measurements in summer 2008.

3.2 Potential Vorticity calculations

The WRF-Chem model does not explicitly calculate potential vorticity (PV). As a result, ~~for conducting the comparisons,~~ PV was calculated offline based on WRF meteorological fields: ~~Model:~~ potential temperature, total mass density, geopotential height and wind speed and direction ~~was used to calculate PV per model gridbox.~~

For each model grid cell, wind and temperature are interpolated from model vertical levels to the potential temperature in the center of the grid cell and the curl of the wind vector is calculated on the corresponding isentropic surfaces using the original model grid (100 km \times 100 km) and the full model vertical resolution, i.e. approximately 500 m in the free troposphere. Potential vorticity is expressed in potential vorticity units (PVu) using the definition $1 \text{ PVu} = 10^{-6} \text{ Kkg}^{-1}\text{m}^2\text{s}^{-1}$. The main uncertainty in the PV calculation is related to the representation of smaller scales (50 km) than the model resolution (e. g. narrow stratospheric streamers near the tropopause).

3.2 Model results interpolations

The measurements used in this study vary in temporal and spatial resolution (section 2), where the model results are 3-hourly on a polar stereographic grid (described in section 3). In order to avoid favoring ~~of~~ data with the highest temporal resolution, the measured data was averaged to 1 min for all the in situ and lidar measurements. The ~~ozone sondes~~ ~~ozonesondes~~ are considered to be instantaneous at the time of the balloon release. A vertical resolution of ~~1 km~~ 1 km is used for the lidars and ~~ozone sondes~~ ~~ozonesondes~~. The measured data is then gridded to $0.5^\circ \times 0.5^\circ$. If more than one ~~measurements~~ measurement is available in the same grid box from the same campaign and instrument within ~~a time frame~~ an interval of one hour, then these are considered to be the same measurement, and a mean value is calculated.

The model grid cell which is the most representative for the measurement's location in space is selected and a linear interpolation in time is done to calculate one minute increments from the 2 modelled values separated by 3 hours and surrounding the measurement time. ~~This~~ In this way the two resulting datasets (modelled and measured) both have ~~a~~ 1-min temporal resolution, and spatially only the vertical resolution is changed to set the altitude increment to 1 km. The model results, including PV, are also horizontally interpolated from ~~the~~ 100 km resolution to match the new $0.5^\circ \times 0.5^\circ$ horizontal data resolution. Each model grid cell is split into 100 mini-grid cells. Then each mini-grid cell is assigned to the appropriate $0.5^\circ \times 0.5^\circ$ final grid cell before calculating the new model values.

4 Comparison of measured and modeled ozone

The WRF-Chem ozone mixing ratios corresponding to the times and locations of the the June/July observations have been used to produce latitudinal cross sections comparable to the results shown in Fig. 3. The ~~model~~ modeled ozone vertical cross sections for Canada and Greenland are shown in Fig. 4. The altitude and latitude ozone variability of the model is comparable ~~with to~~ the observations over both regions. The vertical structure, i.e., the transition between the low ozone values below 3 km and the higher ozone concentration in the free troposphere and, the UTLS ozone variability, are well reproduced. The

latitudinal gradient of the ozone concentration in the 4-8 km altitude range over Greenland is also visible in the simulation results where the two mid-tropospheric ozone branches are also visible at 70°N and 80°N. The agreement is less good for the latitudinal gradient over Canada where the low ozone values (< 50 ppbv) seen in the model at 65°N in the ~~mid-troposphere~~ mid-troposphere are not seen in the observations, although the signs of the latitudinal gradients seem correct.




Figure 4. ~~same~~ Same as Fig. 3 but for the WRF-Chem model ozone mixing ratio corresponding to the measurement ~~sampling characteristics~~ summarized in Tables 1 and 2.

- 5 To quantify the observation/model agreement the scatter plot of modeled versus measured ozone is also presented in Fig. 5 using a PV color scale to distinguish the tropospheric and stratospheric contributions. The ~~correlation measured versus modeled~~ O₃ correlation (red line in in Fig. 5) is of the order of 0.9 in the altitude range 0-15 km over both regions because the occurrence of stratospheric ozone intrusions are very well reproduced by WRF-Chem. The correlation is however between 0.5 and 0.7 in the troposphere only using observations with PV values less than 1 PVu ~~-(green line in in Fig. 5)~~. Considering that the ozone
- 10 variability in the free troposphere is not very large (< 50 ppb), the spatial and temporal variability is still well reproduced by WRF-Chem even below the tropopause region. Even though the UTLS temporal variability is well reproduced by the model (Fig. 4), there is a significant underestimation of ozone by a factor 1.5 in the WRF-Chem simulation for the lowermost stratosphere (PV > 2 PVu) (Fig. 5). In the troposphere there is also a negative bias of the model data of the order of -6 to -15 ppbv with the largest differences over Canada (see Table 5). A fraction of this tropospheric underestimate by WRF-Chem
- 15 is likely related to the ~~ozone-O₃ underestimate in the lowermost stratosphere~~ which implies a smaller source of stratospheric ozone in the free troposphere. This, causing the stratospheric ozone source in the upper troposphere being too small. The modeled O₃ underestimate in the lower stratosphere likely originates from the ~~ozone-O₃ climatology~~ ozone-O₃ climatology used to initialize the model in the stratosphere. The other part of this 10-ppbv bias is due ~~an uncertainty to an underestimate~~ an uncertainty to an underestimate of the lightning NO_x contribution in the WRF-Chem simulation and/or an underestimate of the vertical transport of continental emissions to the

mid- and upper troposphere. Wespes et al. (2012) have shown that the lightning NO_x ozone-source and vertical transport of continental emissions both contribute to 15% of the ozone- O_3 concentration in the free troposphere at latitudes higher than 60°N in summer. Despite these flaws in the WRF-Chem simulations the spatial ozone variability is still very valuable because it the spatial O_3 variability compares rather well with the observations. The PV and CO distribution available from the WRF-Chem simulations can will therefore be used to examine the respective roles of photochemistry and STE on in the observed ozone distribution in the following sections.

5 Latitudinal ozone distribution at high latitudes

In this section, we investigate the mean latitudinal and vertical ozone gradient in the troposphere using the POLARCAT observations. To quantify these gradients, the ozone- O_3 cross sections plotted in Fig.3 have been divided into several regions using the model CO latitudinal cross sections. The vertical boundaries are defined according to the mean ozone vertical profile in the troposphere, i.e. the the depth of the low ozone values layer in the lower troposphere and the downward extent of the tropopause region. The variability of the CO distribution is only taken from the model because CO, since CO data is not available for many of the ozone observations (ozonesonde and lidar profiles).

5.1 Measurements over Canada

For the data taken over Canada, 5 regions have been considered to calculate the mean O_3 latitudinal gradient. They correspond to the blue boxes with the label-labels 1, 2, 3, 4 and 8 shown in the CO latitudinal distribution plotted in Fig. 6. The regional extent of the zones have the same order of magnitude different zones have similar regional extent, in order to make them comparable. Two are below 4 km in the altitude range where the lowest ozone values were recorded for latitudes > 60°N. The boundary between these two boxes is set according to the strong CO latitudinal difference due to the biomass burning emissions south of 65°N and lack of local emissions of ozone precursors in the region between 65°N and 80°N. In the altitude range 4-8 km, which corresponds to the largest tropospheric ozone values, two other regions were defined. The latitude boundary at 63°N is again prescribed according to the latitude where CO and O_3 concentrations are simultaneously decreasing. The last box corresponds to the tropospheric observations at high latitudes (> 80 °N) where CO concentrations are increasing, especially in the altitude range 3-8 km. These ozone observations were mainly made in northeastern Canada by the DC8 aircraft. The CO distribution derived from the WRF-Chem simulation is consistent with the analysis of CO observations by Bian et al. (2013), who show the major influence of boreal and Asian emissions on the ARCTAS-B CO observations.

The mean and median measured O_3 mixing ratio-ratios for the different boxes are shown in Table 3 including, along with the number of observations from the different measurement techniques (lidar, in-situ and sondes). The mean and median O_3 and CO mixing ratios from the model are also reported in Table 4 and the statistics relevant to the model evaluation are reported in Table 5. The -10 to -15 ppbv differences between the model and measured ozone in zones 3, 4 and 8 above 4 km are consistent with the bias of the model in the troposphere over Canada discussed in the previous section. This bias is small (0 to

-4 ppbv) in the lower troposphere, showing that the emissions used in the model simulation are good enough to calculate the O₃ photochemical production.

The negative latitudinal gradient of ozone between zones 3 and 4 ($\Delta O_3 = -6$ ppbv for the measurements and $\Delta O_3 \approx -8$ ppbv for the model) and to a lesser extent between zones 1 and 2 ($\Delta O_3 = -4$ ppbv for the measurements and $\Delta O_3 = -5$ ppbv for the model) are correlated with a significant negative latitudinal CO gradient. Here we compare median rather than mean values as the latter are biased by a few very high values (see below). For latitudes lower than 65°N there is a strong standard deviation because some of the measurements were taken very close to fresh biomass burning sources (forest fires). Sampling of the biomass burning sources during ARCTAS by the DC8 aircraft between 50°N and 63°N has been already discussed in several papers (Singh et al., 2010; Alvarado et al., 2010; Thomas et al., 2013). In the mid-troposphere the CO enhancement in zone 3 where the largest O₃ mixing ratio (70 ppbv) is recorded, is 130 ppbv, well above the CO tropospheric baseline of 60 ppbv (the strong difference between the mean and the median corresponds to the sampling of one biomass burning plume with CO > 500 ppbv which biases the mean). Zone 8 data in Tables 3 and 4 ~~includes~~ combines the aircraft sampling over both Canada and Greenland for the high latitude boxes (> 80°N) because they ~~characterized the same~~ characterize regions with similar ~~ozone and CO distribution. The ozone~~ O₃ and CO distributions. The O₃ mean for zone 8 (70 ppbv) is similar to the ~~ozone mean one~~ found in zone 3 while CO is ~~also~~ increasing again at high latitudes (40 ppbv above the CO tropospheric baseline of 60 ppbv).

The average PV latitudinal cross section is also calculated for the ozone dataset as it is a good tracer of the latitudinal variability of the stratospheric O₃ source (Fig. 7). Although frequent stratospheric air mass intrusions are seen in the 8-10 km altitude range at latitudes higher than 55°N, the role of the stratospheric source is not clearly visible in the average PV values in the free troposphere. Mean PV values larger than 0.5 PVu are only seen for latitudes higher than 75°N in the 3-6 km altitude range (the large PV values at altitudes below 2 km should not be considered as they are related to the low level cyclonic circulation due the orographic circulation around the Greenland ice cap). This suggests that the latitudinal ozone gradients between zones 3 and 4 is not related to the latitudinal distribution of stratospheric ~~intrusion. Ozone intrusions. However~~ O₃ mixing ratios larger than 70 ppbv (Fig. 3) are ~~however~~ correlated with the descending high PV tongue at latitudes between 77°-87°N (Fig. 7). Because the analysis of the ozone spatial distribution shows ~~contrasted behaviors~~ contrasting behavior of the ozone-PV relationship, ~~we also used~~ it is also necessary to look at the PV versus O₃ mixing ratio scatter plot ~~for all the~~. When only tropospheric ozone data ~~, which shows are considered (PV < 1 PVu), there is~~ a poor Pearson correlation ($r < 0.3$) between ozone and ~~stratospheric intrusion~~ PV, implying that stratospheric intrusions are not the only source of high tropospheric ozone values (Fig. 7). ~~Using the color scale of the scatter plot to separate the tropospheric air masses and the UTLS region in the 8-10 km altitude range. Indeed~~ one can see that a significant number of high tropospheric ozone mixing ratios (> 70 ppbv) are ~~indeed~~ related to PV less than 1 PVu when using the altitude color scale to separate the UTLS (8-10 km) and tropospheric data.

The difference in the slopes of the regression line, when including or not the PV values larger than 1 PVu (red versus green lines in Fig. 7), is also larger than similar variation observed at mid-latitudes in Europe where the O₃ to PV ratio decreases from 30 to 150 ppbv/PVu (Ravetta et al., 1999). This suggests that the ozone variability ~~is not from stratospheric air that during summer 2008 is not driven by the variability of stratospheric air~~ mixed into the free troposphere ~~during summer 2008, instead the ozone cycle was mainly driven by,~~ but mainly related to emissions and photochemistry over Canada.




Figure 5. Linear (left) and log-scale (right) scatter plot of WRF-Chem modeled versus measured O₃ mixing ratio in ppbv for Canada (top panel) and Greenland (bottom panel) domains. The black lines are the one to one line and the PV color scale distinguishes stratospheric (red and black-white dots) and tropospheric data (blue and green dots). The regression line parameters and the Pearson correlation coefficient with its p-value are coloured in red and green for the PV range 0-4 PVu and 0-1 PVu, respectively.




Figure 6. Latitudinal cross section of the tropospheric-modeled CO profiles-mixing ratio in ppbv over Canada for the measurement spatio-temporal distribution from the June/July 2008 WRF-Chem simulation. The blue boxes correspond to the regions where data are averaged for discussing vertical and latitudinal gradients in Table 4.

Table 3. Mean, median and standard deviation of the observed ozone mixing ratio in the different boxes shown in Fig. 6 and 8, except for the two boxes at latitudes $> 80^{\circ}\text{N}$ which have been merged in the last row of the table.

Zone N°	Latitude range	Longitude range	Altitude km	O ₃ Mean ppbv	O ₃ Median ppbv	O ₃ St.Dev. ppbv	Number lidar	Number in-situ	Number ECC
1	50°/65°N	-132°/-70°W	0-4	45.8	45.3	10.8	1046	422	243
2	65°/80°N	-132°/-70°W	0-4	42.3	40.9	11.4	401	27	28
3	50°/63°N	-132°/-70°W	4-8	69.4	68.0	18.8	1184	349	240
4	63°/75°N	-132°/-70°W	4-8	63.4	62.0	16.5	360	123	28
5	60°/80°N	-70°/-20°W	0-4	45.1	45.4	8.0	243	603	21
6	57°/65°N	-70°/-20°W	4-8	57.4	56.1	13.8	22	203	0
7	65°/75°N	-70°/-20°W	4-8	68.7	68.1	16.3	320	906	81
8	80°/87°N	-132°/-20°W	3-8	69.8	67.6	14.8	166	48	35

Table 4. same as Table 3 for the WRF-Chem model O₃, CO mixing ratio and PV 75th percentile.

Zone N°	Latitude range	Longitude range	Altitude km	O ₃ Mean ppbv	O ₃ Median ppbv	CO Mean ppbv	CO Median ppbv	PV 75th PVu
1	50°/65°N	-132°/-70°W	0-4	45.9	42.9	441.5	129.6	-
2	65°/80°N	-132°/-70°W	0-4	38.3	37.7	71.8	67.8	-
3	50°/63°N	-132°/-70°W	4-8	60.4	55.4	509.7	129.0	0.35
4	63°/75°N	-132°/-70°W	4.8	45.7	47.9	87.8	87.6	0.31
5	60°/80°N	-70°/-20°W	0-4	45.5	45.6	66.4	66.1	-
6	57°/65°N	-70°/-20°W	4-8	51.4	49.0	83.9	78.8	0.45
7	65°/75°N	-70°/-20°W	4-8	62.8	61.1	76.2	75.8	0.44
8	80°/87°N	-132°/-20°W	3-8	57.7	58.0	100.0	98.7	0.53

Table 5. Metrics of the WRF-Chem O₃ simulation performance for the regions reported in Table 4: mean bias, root mean square error (RMSE), normalized mean bias

Zone N°	Latitude range	Longitude range	Altitude km	Mean bias ppbv	RMSE ppbv	Normalized mean bias, %
1	50°/65°N	-132°/-70°W	0-4	0.1	24.0	0.3
2	65°/80°N	-132°/-70°W	0-4	-4	10.3	-9.5
3	50°/63°N	-132°/-70°W	4-8	-9	24.9	-12.9
4	63°/75°N	-132°/-70°W	4.8	-15	21.9	-28.0
5	60°/80°N	-70°/-20°W	0-4	0.4	7.5	0.8
6	57°/65°N	-70°/-20°W	4-8	-6	15.3	-10.6
7	65°/75°N	-70°/-20°W	4-8	-6	17.1	-8.6
8	80°/87°N	-132°/-20°W	3-8	-13	18.0	-17.2




Figure 7. Latitudinal cross section of the average PV profiles over Canada for the summer measurement distribution using the WRF-Chem simulation (top panel) **Linear**-(left)-and-**log-scale**-(right)-**scatterplot**-**Scatterplot** of **measured**-**modeled** O₃ in ppbv versus PV in PVu using an altitude color coded scale (bottom panel). Stratospheric data points with PV > 4 PVu are not included. The regression line parameters and the Pearson correlation coefficient with its p-value are also given for the PV range 0-4 PVu (red) and 0-1 PVu (green).

10 5.2 Measurements over Greenland

The same procedure was applied to the latitudinal distribution over Greenland. The regions considered for the latitudinal gradient analysis are slightly different because the CO latitudinal cross sections and the ozone vertical structure are different. The selected regions are shown by the blue boxes in Fig. 8. Only one zone is chosen between 60°N and 80°N for the altitude range below 4 km where there are the lowest tropospheric O₃ and CO concentrations values. The CO latitudinal gradient is indeed very weak in the lower troposphere between 60°N and 80°N (< 5 ppbv). However, ~~like-as~~ over Canada, two boxes are chosen in the altitude range 4-8 km because of their differences in term of CO and O₃ concentrations. An additional box corresponds to the tropospheric observations at high latitudes (> 80 °N) where the CO concentrations are also higher, especially in the altitude range 3-8 km. This is related to the observations mainly made in northwestern Greenland by the DC8 aircraft.

Table 5 shows that, there is no O₃ underestimate by the model in the lower troposphere and the model bias above 4 km is of the order of -6 ppbv. This suggests that the model performance over Greenland, away from the continental sources, is better than over Canada. The remaining -6 ppbv bias can be easily explained by the stratospheric O₃ climatology having too low values, while ~~lightning-lightning~~ is known to be less important over Greenland (Cecil et al., 2014; Christian et al., 2003). The CO concentration in the lower troposphere (zone 5 of Table 3) is close to the tropospheric baseline, while O₃ is not markedly different from the values seen over Canada. The positive latitudinal O₃ gradient between zones 6 and 7 in the mid-troposphere above 4 km ($\Delta O_3=12$ ppbv) is two times larger than the latitudinal gradient over Canada, but the overall O₃ mid-tropospheric concentration over Greenland is ~~not-significantly-higher-than~~ similar to its counterpart over Canada (62 ~~ppbv~~-versus 65 ppbv). The corresponding negative CO latitudinal gradient between zone 6 and 7 is weak (difference of ~~the-order-of-8~~ ppbv), but it is however anticorrelated with the ozone gradient. The CO excess above baseline is only 20 ppbv in zone 6, where CO is maximum. The anticorrelation between the O₃ and CO latitudinal gradient may ~~correspond-to-indicate~~ more occurrences of stratospheric ozone intrusion in zone 7 at latitudes higher than 65°N during the POLARCAT period. ~~Less-CO-due-to-If the lower CO were due to upwelling of~~ pristine air from the Arctic lower troposphere ~~would-lead-to-~~ it would be accompanied by smaller O₃ concentrations according to Fig. 3.

Looking at the average PV distribution extracted from the WRF-Chem simulation (Fig. 9), frequent stratospheric air mass intrusions are seen in the altitude range 8-10 km especially at 75°N (PV > 1 PVu). PV values are also larger in the ~~mid-troposphere-mid-troposphere~~ over Greenland than over Canada (75th PV percentile is 0.45 PVu over Greenland but 0.3 PVu over Canada). The PV versus O₃ mixing ratio scatter plot also shows a higher Pearson correlation ($r > 0.4$) between ozone and stratospheric intrusion than over Canada (Fig. 9). ~~Unlike-Compared to~~ the results obtained over Canada, the difference between the O₃ to PV ratio including ~~or-not~~ PV values larger than 1 and that excluding them (red versus green regression line) is smaller (200 ppbv/PVu instead of 400 ppbv/PVu for Canada). ~~It-~~ This is consistent with a larger fraction of ozone from the UTLS over Greenland. However there is not a significant latitudinal PV gradient in the mid-troposphere between zones 6 and 7 where a positive 12 ppbv ozone latitudinal gradient is detected. At high latitudes above 80°N, the PV and CO latitudinal

distribution in the troposphere are almost identical to the latitudinal cross section over Canada (Fig. 7), which ~~support~~supports merging data in a single zone 8 for the statistics reported in Table 3 and 4.




Figure 8. ~~same~~Same as Fig. 6 ~~but~~but for the Greenland region.




Figure 9. same as Fig. 7 but for the Greenland region.

10 5.3 Discussion

This section aims to discuss the latitudinal ozone distribution above Greenland and Canada using (i) existing studies of the role of biomass burning, anthropogenic emissions and STE on high latitude tropospheric ozone, (ii) satellite observations of biomass burning locations and associated plumes during summer 2008 and, (iii) backtrajectory analysis of the air masses where observations were made.

15 In the lower troposphere (0-4 km), the fact that ozone is lower than 50 ppbv everywhere (except close to the fires south west of Hudson Bay) is expected considering a near zero net ozone photochemical production (+0.3 ppbv/day near the surface and -0.3 ppbv/day in the lowermost free troposphere) and a weak influence of STE below 5 km in Northern Canada (Walker et al., 2012; Mauzerall et al., 1996).

In the free troposphere above 4 km, ~~the a~~ negative CO latitudinal gradient in the latitude range 50°N to 75°N is seen over both
20 Greenland and Canada. The large CO latitudinal gradient over Canada can only be explained when considering the biomass burning plume at the continental scale which is superimposed onto the anthropogenic emissions south of 50°N. The position of the biomass burning plume is shown by the MODIS monthly mean aerosol optical depth (~~last panel of Fig. 12~~) ~~with maxima,~~
together with the Canadian Forest Service (CFS) fire counts (Fig. 10). Maxima of aerosol optical depth are seen both south west of Hudson Bay and over the Atlantic ocean in the latitude band 50°N to 60°N. The ozone latitudinal gradient is also negative
25 and correlated with CO over Canada, which can be explained by ~~the~~ ozone photochemical production in the Canadian biomass burning plumes. Thomas et al. (2013) ~~indeed~~ showed a 3 ppbv ozone increase downwind of biomass burning emissions and an enhancement ratio $\Delta O_3 / \Delta CO$ ranging from 0.1 near the fires to 0.5 downwind. ~~It~~ This is consistent with a ΔO_3 of 6-8 ppbv between zone 3 and 4 over Canada where a 40 ppbv difference in the median CO value is found in the model simulation. The contribution of other ~~emissions~~ emission sources can be also estimated using 4-day backward trajectories calculated with the
30 FLEXTRA model and T213/L91 ECMWF analysis (Fig. 11). ~~The differences in the long-range transport between~~ According to the upper panels of Fig. 11, long-range transport into zones 3 and 4 over Canada ~~are not very large with a similar inflow from northern Pacific according to the upper panels of Fig. 11.~~ shows similar patterns with 75% of the air masses coming from North American mid-latitudes, i.e. the area between -60°W/-150°W and 50°N/70°N. Regional variability of STE has also a weak influence during this period considering the weak dependency of O_3 with PV over Canada. Local emissions from biomass burning are the most reasonable explanation for the ozone latitudinal gradient between 50°N and 75°N.

Over southern Greenland the effect of ozone photochemical production related to fire plumes and the North American anthropogenic emissions ~~apperas~~ appears to be less than 4 ppbv, considering the weak CO gradient between zones 6 and 7 (\approx -8 ppbv). The negative O_3 gradient due to increased photochemical production in zone 6 can be then easily counterbalanced by
5 another source in zone 7 to explain the positive latitudinal O_3 gradient over Greenland. The STE ozone source ~~may contribute because of~~ is a likely contributor, as suggested by the better correlation between O_3 and PV over Greenland than Canada and larger PV over Greenland. Frequent tropopause polar vortices developing over Baffin Bay and Davis Strait still occur during the summer period, because of the important role of radiation to maintain and intensify the upper troposphere cyclonic activity during the summer (Cavallo and Hakim, 2010). However, in the absence of a clear PV latitudinal gradient between

10 55°N and 75°N in the model simulation cannot explain the significant +12 ppbv ozone latitudinal gradient. Therefore, while
STE certainly contributes ~~in the ozone level to the ozone budget~~ over both southern and northern Greenland, ~~but~~ it does not
explain the largest O₃ values found in zone 7. Looking at the ~~long-range-long-range~~ transport plot for zones 6 and 7 (lower
panels of Fig. 11), ~~multiple the fraction of air masses coming from the North American mid-latitude area are now 34% and~~
15 ~~18%, respectively. Multiple mid-latitude sources including North America, Europe and even cross-Arctic transport can lead~~
~~to a more efficient photochemical production. Asia are also related to ozone observations in zone 7 compared to zone 6.~~
and may explain the differences in ozone production efficiency between zone 6 and zone 7. Wespes et al. (2012) concluded
using the Model MOZART-4 that anthropogenic pollution from Europe dominates O₃ concentrations in summer 2008 in the
Arctic, while Roiger et al. (2011) shows using aircraft measurements that Asian anthropogenic pollution can be mixed with
stratospheric air masses over Greenland in the 75°-80° latitude band.

20 For the free troposphere at high latitudes above 80 °N, the STE contribution is maximum in this region for the 4-8 km altitude
range according to the average PV distribution shown in Fig.7 or Fig.9. The 75th percentile of the PV distribution is also >
0.5 PVu in zone 8 while it is 0.3 PVu at latitudes lower than 75° over Canada and 0.4 PVu over Greenland. Zängl and Hoinka
(2001) discussed the horizontal gradient of the tropopause height over the Arctic using ECMWF analysis and radiosondes.
The region with the largest horizontal gradient is displaced to the North near 80 °N in July over North America, while vertical
25 tilting of the isentropic surfaces remains small. These conditions are favorable for isentropic motion across the tropopause
with more efficient STE. However CO ~~is also increasing again also increases~~ at high latitude because of ~~the transport-transport~~
from Asian and North Siberian pollution sources according to the air mass transport pathway for zone 8 (Fig. 12). Such a
mixture of stratospheric ~~ozone~~-O₃ and Asian pollution has been already suggested by Roiger et al. (2011) to explain the ozone
concentrations observed at very high latitudes in the Arctic. ~~This~~ Our study using more ozone measurements leads to the same
30 conclusions.

6 Conclusions

The purpose of this work is to provide a complete picture of Arctic ozone using measurements available during the summer
2008 POLARCAT campaigns over Canada and Greenland, where 3 aircraft were deployed and 7 ozonesonde stations intensi-
fied their ozone vertical profiling. This is the first case of such complete temporal and geographical coverage, ~~specifically we~~
. We take advantage of the large number of airborne lidar profiles (representing 67% of the O₃ measurements over Canada
and 26% over Greenland. ~~A~~). The measured ozone climatology established in this paper can also be used for future model
evaluation at high latitudes. For example, in our work, while a good correspondence of the measured O₃ vertical and latitudinal
5 distribution is found with model results from WRF-Chem, ~~especially over Greenland. A~~ a negative O₃ bias of -6 to -15 ppbv
between the model and the observations is found in the free troposphere over 4 km, especially over Canada. ~~This deficiency~~ is
partly related to the ~~model deficiencies in the lowermost stratosphere, because of the ozone climatology used for stratospheric~~
~~ozone. The measured ozone climatology established in this paper can serve as a benchmark for future model evaluation at high~~
~~latitudes.~~ WRF-Chem model stratospheric ozone initialization.





Figure 10. Map of the Canadian Forest Service fire counts in red for summer 2008 (left) MODIS aerosol optical depth at 0.55 μm from 15 June to 15 July 2008 (right).


- 10 The ~~analysis of the~~ WRF-Chem model simulation is also ~~useful~~ used to discuss the relative influence of tropospheric ozone sources at high latitude in summer. Ozone average concentrations are of the order of 65 ppbv at altitudes above 4 km both over Canada and Greenland, while they are less than 50 ppbv in the lower troposphere. For Canada, the analysis of the ~~model-modeled~~ CO distribution and the weak correlation ($< 30\%$) of O_3 and PV suggest that stratosphere-troposphere exchange (STE) is not the major contribution to tropospheric ozone at latitudes less than 70°N , ~~because local~~ where transport of
- 15 North American biomass burning (BB) emissions ~~were significant~~ took place during the 2008 summer ~~period~~. ~~Conversely over Greenland~~. ~~Conversely~~, significant STE is found over Greenland according to the better O_3 versus PV correlation ($> 40\%$) and the higher value of the 75th PV percentile. This is related to the persistence of cyclonic activity during the summer over Baffin Bay.

A weak negative latitudinal summer ozone gradient of -6 to -8 ppbv is found over Canada in the mid troposphere between 4 and 8 km because the O_3 photochemical production ~~due to the from~~ BB emissions mainly takes place at latitudes less than 65°N , while STE plays ~~only a significant role at latitude~~ a larger role at latitudes higher than 70°N . A positive ozone latitudinal gradient of 12 ppbv is observed in the same altitude range over Greenland not because of an increasing latitudinal influence of STE, but because of different ~~long-range~~ long-range transport from multiple mid-latitude sources (North America, Europe and even Asia for latitudes higher than 77°N).


Zone 3 Back Trajectory - Altitude (km) color scale - STE 0.037018



Zone 4 Back Trajectory - Altitude (km) color scale - STE 0.051489



Zone 6 Back Trajectory - Altitude (km) color scale - STE 0.078363



Zone 7 Back Trajectory - Altitude (km) color scale - STE 0.13453




Figure 11. Map of the air mass daily positions using 4-days FLEXTRA trajectories which correspond to the observations in zones 3 and 4 over Canada (top row) and in zones 6 and 7 over Greenland (bottom row). Black crosses show the measurement positions and the color scale is the air mass altitude in km. The fraction is the relative number of trajectory positions reaching the tropopause (PV=1.5 PVu).

For the Arctic latitudes ($> 80^{\circ}\text{N}$), free tropospheric O_3 concentrations are related to a mixture of stratospheric O_3 transport across the tropopause and Asian pollution, as already suggested by Roiger et al. (2011) using a case study of aircraft observations in the Arctic. Our study using more ozone measurements leads to the same conclusions.

Zone 8 Back Trajectory - Altitude (km) color scale - STE 0.13977




Figure 12. same as Fig.11 for the observations at latitudes $> 80^{\circ}\text{N}$ in zone 8 (left). Map of MODIS aerosol optical depth at 0.55 from 15 June to 15 July 2008 (right). 8.

Acknowledgements. We are very grateful to the support of the Meteo France/CNRS/CNES UMS SAFIRE for the ATR-42 aircraft deployment over Greenland. This work was supported by funding from ANR and LEFE INSU/CNRS (CLIMSLIP project) and from the ICE-ARC programme from the European Union 7th Framework Programme, grant number 603887, ICE-ARC contribution number XXXXXX

10 -603887. The FLEXTRA team (A. Stohl, and co-workers) is acknowledged for providing and supporting the FLEXTRA code. NASA and DLR are acknowledged for their support to the deployment of the DC8 and Falcon 20 aircraft. WOUDC is and the NASA MODIS team are acknowledged for providing the ozonesonde data and the MODIS data, respectively.

References

- Abbatt, J. P. D., Thomas, J. L., Abrahamsson, K., Boxe, C., Granfors, A., Jones, A. E., King, M. D., Saiz-Lopez, A., Shepson, P. B., Sodeau, J., Toohey, D. W., Toubin, C., von Glasow, R., Wren, S. N., and Yang, X.: Halogen activation via interactions with environmental ice and snow in the polar lower troposphere and other regions, *Atmospheric Chemistry and Physics*, 12, 6237–6271, doi:10.5194/acp-12-6237-2012, <http://www.atmos-chem-phys.net/12/6237/2012/>, 2012.
- Alvarado, M. J., Logan, J. A., Mao, J., Apel, E., Riemer, D., Blake, D., Cohen, R. C., Min, K.-E., Perring, A. E., Browne, E. C., Wooldridge, P. J., Diskin, G. S., Sachse, G. W., Fuelberg, H., Sessions, W. R., Harrigan, D. L., Huey, G., Liao, J., Case-Hanks, A., Jimenez, J. L., Cubison, M. J., Vay, S. A., Weinheimer, A. J., Knapp, D. J., Montzka, D. D., Flocke, F. M., Pollack, I. B., Wennberg, P. O., Kurten, A., Crounse, J., Clair, J. M. S., Wisthaler, A., Mikoviny, T., Yantosca, R. M., Carouge, C. C., and Le Sager, P.: Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: an integrated analysis of aircraft and satellite observations, *Atmospheric Chemistry and Physics*, 10, 9739–9760, doi:10.5194/acp-10-9739-2010, <http://www.atmos-chem-phys.net/10/9739/2010/>, 2010.
- 25 AMAP: Assessment 2015: Black carbon and ozone as Arctic climate forcers. Arctic Monitoring and Assessment Programme (AMAP), pp. 1–116, Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, <http://www.amap.no/documents/doc/AMAP-Assessment-2015-Black-carbon-and-ozone-as-Arctic-climate-forcers/1299>, 2015.
- Ancellet, G. and Ravetta, F.: Compact airborne lidar for tropospheric ozone: description and field measurements, *Appl. Opt.*, 37, 5509–5521, doi:10.1364/AO.37.005509, <http://ao.osa.org/abstract.cfm?URI=ao-37-24-5509>, 1998.
- 30 Ancellet, G. and Ravetta, F.: On the usefulness of an airborne lidar for O₃ layer analysis in the free troposphere and the planetary boundary layer, *J. Environ. Monit.*, 5, 47–56, doi:10.1039/B205727A, <http://dx.doi.org/10.1039/B205727A>, 2003.
- Bian, H., Colarco, P. R., Chin, M., Chen, G., Rodriguez, J. M., Liang, Q., Blake, D., Chu, D. A., da Silva, A., Darmenov, A. S., Diskin, G., Fuelberg, H. E., Huey, G., Kondo, Y., Nielsen, J. E., Pan, X., and Wisthaler, A.: Source attributions of pollution to the Western Arctic during the NASA ARCTAS field campaign, *Atmospheric Chemistry and Physics*, 13, 4707–4721, doi:10.5194/acp-13-4707-2013, <http://www.atmos-chem-phys.net/13/4707/2013/>, 2013.
- 35 Browell, E., Carter, A., Shipley, S., Allen, R., Butler, C., Mayo, M., J. Siviter, Jr., and Hall, W.: NASA multipurpose airborne DIAL system and measurements of ozone and aerosol profiles, *Appl. Opt.*, 22, 522–534, doi:10.1364/AO.22.000522, 1983.
- Browell, E., Ismail, S., and Shipley, S.: Ultraviolet DIAL measurements of O₃ profiles in regions of spatially inhomogeneous aerosols, *Appl. Opt.*, 24, 2827–2836, doi:10.1364/AO.24.002827, 1985.
- Browell, E. V., Hair, J. W., Butler, C. F., Grant, W. B., DeYoung, R. J., Fenn, M. A., Brackett, V. G., Clayton, M. B., Brasseur, L. A., Harper, D. B., Ridley, B. A., Klonecki, A. A., Hess, P. G., Emmons, L. K., Tie, X., Atlas, E. L., Cantrell, C. A., Wimmers, A. J., Blake, D. R., Coffey, M. T., Hannigan, J. W., Dibb, J. E., Talbot, R. W., Flocke, F., Weinheimer, A. J., Fried, A., Wert, B., Snow, J. A., and Lefer, B. L.: Ozone, aerosol, potential vorticity, and trace gas trends observed at high-latitudes over North America from February to May 2000, *Journal of Geophysical Research: Atmospheres*, 108, doi:10.1029/2001JD001390, <http://dx.doi.org/10.1029/2001JD001390>, 8369, 2003.
- Buhaus, O., Corbett, J., Endresen, O., Eyring, V., Faber, J., Hanayama, S., Lee, D.S. and Lee, D., Lindstad, H., Markowska, A., Mjelde, A., Nelissen, D., Nilsen, J., Palsson, C., Winebrake, J., Wu, W., and Yoshida, K.: Second IMO GHG study 2009, Tech. rep., 2009.
- 10 Cavallo, S. M. and Hakim, G. J.: Composite Structure of Tropopause Polar Cyclones, *Monthly Weather Review*, 138, 3840–3857, doi:10.1175/2010MWR3371.1, <http://dx.doi.org/10.1175/2010MWR3371.1>, 2010.

- Cecil, D. J., Buechler, D. E., and Blakeslee, R. J.: Gridded lightning climatology from TRMM-LIS and OTD: Dataset description, *Atmospheric Research*, 135-136, 404 – 414, doi:10.1016/j.atmosres.2012.06.028, <http://www.sciencedirect.com/science/article/pii/S0169809512002323>, 2014.
- Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Goodman, S. J., Hall, J. M., Koshak, W. J., Mach, D. M., and Stewart, M. F.: Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *Journal of Geophysical Research: Atmospheres*, 108, ACL 4–1–ACL 4–15, doi:10.1029/2002JD002347, <http://dx.doi.org/10.1029/2002JD002347>, 4005, 2003.
- 20 [Cooper, O. R., Parrish, D. D., Stohl, A., Trainer, M., Nedelec, P., Thouret, V., Cammas, J. P., Oltmans, S. J., Johnson, B. J., Tarasick, D., Leblanc, T., McDermid, I. S., Jaffe, D., Gao, R., Stith, J., Ryerson, T., Aikin, K., Campos, T., Weinheimer, A., and Avery, M. A.: Increasing springtime ozone mixing ratios in the free troposphere over western North America, *Nature*, 463, 344–348, <http://dx.doi.org/10.1038/nature08708>, 2010.](#)
- Crutzen, P., Lawrence, M., and Pöschl, U.: On the background photochemistry of tropospheric ozone, *Tellus*, 51A-B, 123–146, doi:10.3402/tellusb.v51i1.16264, <http://www.tellusb.net/index.php/tellusb/article/view/16264>, 1999.
- 25 Dupont, R., Pierce, B., Worden, J., Hair, J., Fenn, M., Hamer, P., Natarajan, M., Schaack, T., Lenzen, A., Apel, E., Dibb, J., Diskin, G., Huey, G., Weinheimer, A., Kondo, Y., and Knapp, D.: Attribution and evolution of ozone from Asian wild fires using satellite and aircraft measurements during the ARCTAS campaign, *Atmospheric Chemistry and Physics*, 12, 169–188, doi:10.5194/acp-12-169-2012, <http://www.atmos-chem-phys.net/12/169/2012/>, 2012.
- 30 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, doi:10.5194/gmd-3-43-2010, <http://www.geosci-model-dev.net/3/43/2010/>, 2010.
- Fast, J. D., Gustafson, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *Journal of Geophysical Research: Atmospheres*, 111, n/a–n/a, doi:10.1029/2005JD006721, <http://dx.doi.org/10.1029/2005JD006721>, d21305, 2006.
- 35 Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Browell, E. V., Hair, J. W., Avery, M. A., McGee, T. J., Twigg, L. W., Sumnicht, G. K., Jucks, K. W., Margitan, J. J., Sen, B., Stachnik, R. A., Toon, G. C., Bernath, P. F., Boone, C. D., Walker, K. A., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder stratospheric ozone measurements, *Journal of Geophysical Research: Atmospheres*, 113, n/a–n/a, doi:10.1029/2007JD008771, <http://dx.doi.org/10.1029/2007JD008771>, d15S20, 2008.
- Granier, C., Niemeier, U., Jungclaus, J. H., Emmons, L., Hess, P., Lamarque, J.-F., Walters, S., and Brasseur, G. P.: Ozone pollution from future ship traffic in the Arctic northern passages, *Geophysical Research Letters*, 33, doi:10.1029/2006GL026180, <http://dx.doi.org/10.1029/2006GL026180>, 113807, 2006.
- 10 Grell, G. and Dévényi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation techniques, *Geophysical Research Letters*, 29, 38–1–38–4, doi:10.1029/2002GL015311, <http://dx.doi.org/10.1029/2002GL015311>, 2002.

- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled online chemistry within the WRF model, *Atmospheric Environment*, 39, 6957–6975, doi:<http://dx.doi.org/10.1016/j.atmosenv.2005.04.027>, <http://www.sciencedirect.com/science/article/pii/S1352231005003560>, 2005.
- 15 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, <http://www.geosci-model-dev.net/5/1471/2012/>, 2012.
- Hess, P. G. and Zbinden, R.: Stratospheric impact on tropospheric ozone variability and trends: 1990-2009, *Atmospheric Chemistry and Physics*, 13, 649–674, doi:10.5194/acp-13-649-2013, <http://www.atmos-chem-phys.net/13/649/2013/>, 2013.
- 20 Hong, S.-Y., Noh, Y., and Dudhia, J.: A New Vertical Diffusion Package with an Explicit Treatment of Entrainment Processes, *Monthly Weather Review*, 134, 2318–2341, doi:10.1175/MWR3199.1, <http://dx.doi.org/10.1175/MWR3199.1>, 2006.
- Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell, B.: Evidence of NO_x production within or upon ice particles in the Greenland snowpack, *Geophysical Research Letters*, 26, 695–698, doi:10.1029/1999GL900077, <http://dx.doi.org/10.1029/1999GL900077>, 1999.
- 25 Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, *Atmospheric Chemistry and Physics*, 10, 5191–5212, doi:10.5194/acp-10-5191-2010, <http://www.atmos-chem-phys.net/10/5191/2010/>, 2010.
- 30 Klimont, Z., Smith, S. J., and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000-2011 emissions, *Environmental Research Letters*, 8, 014003, <http://stacks.iop.org/1748-9326/8/i=1/a=014003>, 2013.
- Koo, J.-H., Wang, Y., Kurosu, T. P., Chance, K., Rozanov, A., Richter, A., Oltmans, S. J., Thompson, A. M., Hair, J. W., Fenn, M. A., Weinheimer, A. J., Ryerson, T. B., Solberg, S., Huey, L. G., Liao, J., Dibb, J. E., Neuman, J. A., Nowak, J. B., Pierce, R. B., Natarajan, M., and Al-Saadi, J.: Characteristics of tropospheric ozone depletion events in the Arctic spring: analysis of the ARCTAS, ARCPAC, and
- 35 ARCIIONS measurements and satellite BrO observations, *Atmospheric Chemistry and Physics*, 12, 9909–9922, doi:10.5194/acp-12-9909-2012, <http://www.atmos-chem-phys.net/12/9909/2012/>, 2012.
- Lait, L. R., Newman, P. A., Schoeberl, M. R., McGee, T., Twigg, L., Browell, E. V., Fenn, M. A., Grant, W. B., Butler, C. F., Bevilacqua, R., Davies, J., DeBacker, H., Andersen, S. B., Kyrö, E., Kivi, E., von der Gathen, P., Claude, H., Benesova, A., Skrivankova, P., Dorokhov, V., Zaitcev, I., Braathen, G., Gil, M., Litynska, Z., Moore, D., and Gerding, M.: Non-coincident inter-instrument comparisons of ozone measurements using quasi-conservative coordinates, *Atmospheric Chemistry and Physics*, 4, 2345–2352, doi:10.5194/acp-4-2345-2004, <http://www.atmos-chem-phys.net/4/2345/2004/>, 2004.
- Law, K. S., Stohl, A., Quinn, P. K., Brock, C. A., Burkhart, J. F., Paris, J.-D., Ancellet, G., Singh, H. B., Roiger, A., Schlager, H., Dibb, J.,
- 5 Jacob, D. J., Arnold, S. R., Pelon, J., and Thomas, J. L.: Arctic Air Pollution: New Insights from POLARCAT-IPY, *Bull. Amer. Meteor. Soc.*, 95, 1873–1895, doi:10.1175/BAMS-D-13-00017.1, <http://dx.doi.org/10.1175/BAMS-D-13-00017.1>, 2014.
- Lee, D. S., Fahey, D. W., Forster, P. M., Newton, P. J., Wit, R. C., Lim, L. L., Owen, B., and Sausen, R.: Aviation and global climate change in the 21st century, *Atmospheric Environment*, 43, 3520 – 3537, doi:<http://dx.doi.org/10.1016/j.atmosenv.2009.04.024>, <http://www.sciencedirect.com/science/article/pii/S1352231009003574>, 2009.
- 10 Legrand, M., Preunkert, S., Jourdain, B., Gallée, H., Goutail, F., Weller, R., and Savarino, J.: Year-round record of surface ozone at coastal (Dumont d’Urville) and inland (Concordia) sites in East Antarctica, *Journal of Geophysical Research: Atmospheres*, 114, doi:10.1029/2008JD011667, <http://dx.doi.org/10.1029/2008JD011667>, d20306, 2009.

- [Lin, M., Horowitz, L. W., Cooper, O. R., Tarasick, D., Conley, S., Iraci, L. T., Johnson, B., Leblanc, T., Petropavlovskikh, I., and Yates, E. L.: Revisiting the evidence of increasing springtime ozone mixing ratios in the free troposphere over western North America, *Geophysical Research Letters*, 42, 8719–8728, doi:10.1002/2015GL065311, <http://dx.doi.org/10.1002/2015GL065311>, 2015GL065311, 2015.](#)
- 15 Livesey, N. J., Filipiak, M. J., Froidevaux, L., Read, W. G., Lambert, A., Santee, M. L., Jiang, J. H., Pumphrey, H. C., Waters, J. W., Cofield, R. E., Cuddy, D. T., Daffer, W. H., Drouin, B. J., Fuller, R. A., Jarnot, R. F., Jiang, Y. B., Knosp, B. W., Li, Q. B., Perun, V. S., Schwartz, M. J., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. A., Avery, M., Browell, E. V., Cammas, J.-P., Christensen, L. E., Diskin, G. S., Gao, R.-S., Jost, H.-J., Loewenstein, M., Lopez, J. D., Nedelec, P., Osterman, G. B., Sachse, G. W., and Webster, C. R.: Validation of Aura
- 20 Microwave Limb Sounder O₃ and CO observations in the upper troposphere and lower stratosphere, *Journal of Geophysical Research: Atmospheres*, 113, n/a–n/a, doi:10.1029/2007JD008805, <http://dx.doi.org/10.1029/2007JD008805>, d15S02, 2008.
- Marelle, L., Raut, J.-C., Thomas, J. L., Law, K. S., Quennehen, B., Ancellet, G., Pelon, J., Schwarzenboeck, A., and Fast, J. D.: Transport of anthropogenic and biomass burning aerosols from Europe to the Arctic during spring 2008, *Atmospheric Chemistry and Physics*, 15, 3831–3850, doi:10.5194/acp-15-3831-2015, <http://www.atmos-chem-phys.net/15/3831/2015/>, 2015.
- 25 Marengo, A., Thouret, V., Nédélec, P., Smit, H., Helten, M., Kley, D., Karcher, F., Simon, P., Law, K., Pyle, J., Poschmann, G., Von Wrede, R., Hume, C., and Cook, T.: Measurement of ozone and water vapor by Airbus in-service aircraft: The MOZAIC airborne program, an overview, *Journal of Geophysical Research: Atmospheres*, 103, 25 631–25 642, doi:10.1029/98JD00977, <http://dx.doi.org/10.1029/98JD00977>, 1998.
- Mauzerall, D., Jacob, D., Fan, S. M., Bradshaw, J., Gregory, G., Sachse, G., and Blake, D.: Origin of tropospheric ozone at remote high
- 30 northern latitudes in summer, *J. Geophys. Res.*, 101(D2), 4175–4188, 1996.
- Max, M.-D. C. and Suarez, M. J.: Volume 3: An Efficient Thermal Infrared Radiation Parameterization For Use In General Circulation Models in Technical report series on global modeling and data assimilation, Tech. rep., 1994.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *Journal of Geophysical Research: Atmospheres*, 102, 16 663–16 682,
- 35 doi:10.1029/97JD00237, <http://dx.doi.org/10.1029/97JD00237>, 1997.
- Monks, S. A., Arnold, S. R., Emmons, L. K., Law, K. S., Turquety, S., Duncan, B. N., Flemming, J., Huijnen, V., Tilmes, S., Langner, J., Mao, J., Long, Y., Thomas, J. L., Steenrod, S. D., Raut, J. C., Wilson, C., Chipperfield, M. P., Diskin, G. S., Weinheimer, A., Schlager, H., and Ancellet, G.: Multi-model study of chemical and physical controls on transport of anthropogenic and biomass burning pollution to the Arctic, *Atmospheric Chemistry and Physics*, 15, 3575–3603, doi:10.5194/acp-15-3575-2015, <http://www.atmos-chem-phys.net/15/3575/2015/>, 2015.
- Morrison, H., Thompson, G., and Tatarskii, V.: Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes, *Mon. Weather Rev.*, 137, 991–1007,
- 5 doi:10.1175/2008MWR2556.1, <http://dx.doi.org/10.1175/2008MWR2556.1>, 2009.
- Olson, J. R., Crawford, J. H., Brune, W., Mao, J., Ren, X., Fried, A., Anderson, B., Apel, E., Beaver, M., Blake, D., Chen, G., Crounse, J., Dibb, J., Diskin, G., Hall, S. R., Huey, L. G., Knapp, D., Richter, D., Riemer, D., Clair, J. S., Ullmann, K., Walega, J., Weinbringer, P., Weinheimer, A., Wennberg, P., and Wisthaler, A.: An analysis of fast photochemistry over high northern latitudes during spring and summer using in-situ observations from ARCTAS and TOPSE, *Atmospheric Chemistry and Physics*, 12, 6799–6825, doi:10.5194/acp-12-
- 10 6799-2012, <http://www.atmos-chem-phys.net/12/6799/2012/>, 2012.
- [Papayannis, A., Ancellet, G., Pelon, J., and Megie, G.: Multiwavelength lidar for ozone measurements in the troposphere and the lower stratosphere, *Appl. Opt.*, 29, 467–476, 1990.](#)

- Parrish, D. D., Law, K. S., Staehelin, J., Derwent, R., Cooper, O. R., Tanimoto, H., Volz-Thomas, A., Gilge, S., Scheel, H.-E., Steinbacher, M., and Chan, E.: Long-term changes in lower tropospheric baseline ozone concentrations at northern mid-latitudes, *Atmospheric Chemistry and Physics*, 12, 11 485–11 504, doi:10.5194/acp-12-11485-2012, <http://www.atmos-chem-phys.net/12/11485/2012/>, 2012.
- Ravetta, F., Ancellet, G., Kowol-Santen, J., Wilson, R., and Nedeljkovic, D.: Ozone, temperature and wind field measurements in a tropopause fold: comparison with a mesoscale model simulation, *Mon. Weather Rev.*, pp. 2641–2653, 1999.
- Richter, D. A., Browell, E. V., Butler, C. F., and Higdon, N. S.: Advanced airborne UV DIAL for stratospheric and tropospheric ozone and aerosol measurements, in: *Advances in Atmospheric Remote Sensing with Lidar*, edited by Ansmann, A., pp. 395–398, Springer-Verlag New York, 1997.
- Roiger, A., Schlager, H., Schäfler, A., Huntrieser, H., Scheibe, M., Aufmhoff, H., Cooper, O. R., Sodemann, H., Stohl, A., Burkhart, J., Lazzara, M., Schiller, C., Law, K. S., and Arnold, F.: In-situ observation of Asian pollution transported into the Arctic lowermost stratosphere, *Atmospheric Chemistry and Physics*, 11, 10 975–10 994, doi:10.5194/acp-11-10975-2011, <http://www.atmos-chem-phys.net/11/10975/2011/>, 2011.
- Schlager, H., Konopka, P., Schulte, P., Schumann, U., Ziereis, H., Arnold, F., Klemm, M., Hagen, D. E., Whitefield, P. D., and Ovarlez, J.: In situ observations of air traffic emission signatures in the North Atlantic flight corridor, *Journal of Geophysical Research: Atmospheres*, 102, 10 739–10 750, doi:10.1029/96JD03748, <http://dx.doi.org/10.1029/96JD03748>, 1997.
- Shindell, D.: Local and remote contributions to Arctic warming, *Geophys. Res. Lett.*, 34, L14 704, doi:10.1029/2007GL030 221, 2007.
- Shindell, D. T., Faluvegi, G., Koch, D. M., Schmidt, G. A., Unger, N., and Bauer, S. E.: Improved Attribution of Climate Forcing to Emissions, *Science*, 326, 716–718, doi:10.1126/science.1174760, <http://www.sciencemag.org/content/326/5953/716.abstract>, 2009.
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmospheric Chemistry and Physics*, 7, 4375–4418, doi:10.5194/acp-7-4375-2007, <http://www.atmos-chem-phys.net/7/4375/2007/>, 2007.
- Singh, H., Anderson, B., Brune, W., Cai, C., Cohen, R., Crawford, J., Cubison, M., Czech, E., Emmons, L., Fuelberg, H., Huey, G., Jacob, D., Jimenez, J., Kaduwela, A., Kondo, Y., Mao, J., Olson, J., Sachse, G., Vay, S., Weinheimer, A., Wennberg, P., and Wisthaler, A.: Pollution influences on atmospheric composition and chemistry at high northern latitudes: Boreal and California forest fire emissions, *Atmospheric Environment*, 44, 4553 – 4564, doi:<http://dx.doi.org/10.1016/j.atmosenv.2010.08.026>, <http://www.sciencedirect.com/science/article/pii/S1352231010007004>, 2010.
- Stohl, A., Berg, T., Burkhart, J. F., Fjaeraa, A. M., Forster, C., Herber, A., Hov, O., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Torseth, K., Treffeisen, R., Virkkunen, K., and Yttri, K. E.: Arctic smoke – record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmospheric Chemistry and Physics*, 7, 511–534, <http://www.atmos-chem-phys.net/7/511/2007/>, 2007.
- Tarasick, D. W., Jin, J. J., Fioletov, V. E., Liu, G., Thompson, A. M., Oltmans, S. J., Liu, J., Sioris, C. E., Liu, X., Cooper, O. R., Dann, T., and Thouret, V.: High-resolution tropospheric ozone fields for INTEX and ARCTAS from IONS ozonesondes, *Journal of Geophysical Research: Atmospheres*, 115, doi:10.1029/2009JD012918, <http://dx.doi.org/10.1029/2009JD012918>, d20301, 2010.
- Thomas, J. L., Raut, J.-C., Law, K. S., Marelle, L., Ancellet, G., Ravetta, F., Fast, J. D., Pfister, G., Emmons, L. K., Diskin, G. S., Weinheimer, A., Roiger, A., and Schlager, H.: Pollution transport from North America to Greenland during summer 2008, *Atmospheric Chemistry and Physics*, 13, 3825–3848, doi:10.5194/acp-13-3825-2013, <http://www.atmos-chem-phys.net/13/3825/2013/>, 2013.

- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997-2009), *Atmospheric Chemistry and Physics*, 10, 11 707–11 735, doi:10.5194/acp-10-11707-2010, <http://www.atmos-chem-phys.net/10/11707/2010/>, 2010.
- Walker, T. W., Jones, D. B. A., Parrington, M., Henze, D. K., Murray, L. T., Bottenheim, J. W., Anlauf, K., Worden, J. R., Bowman, K. W., Shim, C., Singh, K., Kopacz, M., Tarasick, D. W., Davies, J., von der Gathen, P., Thompson, A. M., and Carouge, C. C.: Impacts of mid-latitude precursor emissions and local photochemistry on ozone abundances in the Arctic, *Journal of Geophysical Research: Atmospheres*, 117, doi:10.1029/2011JD016370, <http://dx.doi.org/10.1029/2011JD016370>, d01305, 2012.
- Wang, Y., Ridley, B., Fried, A., Cantrell, C., Davis, D., Chen, G., Snow, J., Heikes, B., Talbot, R., Dibb, J., Flocke, F., Weinheimer, A., Blake, N., Blake, D., Shetter, R., Lefer, B., Atlas, E., Coffey, M., Walega, J., and Wert, B.: Springtime photochemistry at northern mid and high latitudes, *Journal of Geophysical Research: Atmospheres*, 108, doi:10.1029/2002JD002227, <http://dx.doi.org/10.1029/2002JD002227>, 8358, 2003.
- Weinheimer, A. J., Walega, J. G., Ridley, B. A., Gary, B. L., Blake, D. R., Blake, N. J., Rowland, F. S., Sachse, G. W., Anderson, B. E., and Collins, J. E.: Meridional distributions of NO_x, NO_y, and other species in the lower stratosphere and upper troposphere during AASE II, *Geophysical Research Letters*, 21, 2583–2586, doi:10.1029/94GL01897, <http://dx.doi.org/10.1029/94GL01897>, 1994.
- Wespes, C., Emmons, L., Edwards, D. P., Hannigan, J., Hurtmans, D., Saunio, M., Coheur, P.-F., Clerbaux, C., Coffey, M. T., Batchelor, R. L., Lindenmaier, R., Strong, K., Weinheimer, A. J., Nowak, J. B., Ryerson, T. B., Crounse, J. D., and Wennberg, P. O.: Analysis of ozone and nitric acid in spring and summer Arctic pollution using aircraft, ground-based, satellite observations and MOZART-4 model: source attribution and partitioning, *Atmospheric Chemistry and Physics*, 12, 237–259, doi:10.5194/acp-12-237-2012, <http://www.atmos-chem-phys.net/12/237/2012/>, 2012.
- Zängl, G. and Hoinka, K. P.: The Tropopause in the Polar Regions, *Journal of Climate*, 14, 3117–3139, doi:10.1175/1520-0442(2001)014<3117:TTITPR>2.0.CO;2, [http://dx.doi.org/10.1175/1520-0442\(2001\)014<3117:TTITPR>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2001)014<3117:TTITPR>2.0.CO;2), 2001.
- Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *Journal of Geophysical Research: Atmospheres*, 104, 30 387–30 415, doi:10.1029/1999JD900876, <http://dx.doi.org/10.1029/1999JD900876>, 1999.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *Journal of Geophysical Research: Atmospheres*, 113, n/a–n/a, doi:10.1029/2007JD008782, <http://dx.doi.org/10.1029/2007JD008782>, d13204, 2008.