

# Chemistry-climate interactions of aerosol nitrate from lightning

Tost H.<sup>1</sup>

<sup>1</sup>Institute for Atmospheric Physics, Johannes Gutenberg University Mainz, Germany

*Correspondence to:* H. Tost (tosth@uni-mainz.de)

**Abstract.** Lightning represents one of the dominant emission sources for NO<sub>x</sub> in the troposphere. The direct release of oxidised nitrogen in the upper troposphere does not only affect ozone formation, but also chemical and microphysical properties of aerosol particles in this region. This study investigates the direct impact of LNO<sub>x</sub> emissions on upper tropospheric nitrate using a global chemistry climate model. The simulation results show a substantial influence of the lightning emissions on the mixing ratios of nitrate aerosol in the upper troposphere of more than 50%. In addition to the impact on nitrate, lightning substantially affects the oxidising capacity of the atmosphere with substantial implications for gas phase sulphate formation and new particle formation in the upper troposphere. In conjunction with the condensation of nitrates, substantial differences in the aerosol size distribution occur in the upper troposphere as a consequence of lightning. This has implications for the extinction properties of the aerosol particles and for the cloud optical properties. While the extinction is generally slightly enhanced due to the LNO<sub>x</sub> emissions, the response of the clouds is ambiguous due to compensating effects in both liquid and ice clouds. Resulting shortwave flux perturbations are of  $\sim -100\text{mW/m}^2$  as determined from several sensitivity scenarios, but an uncertainty range of almost 50% has to be defined due to the large internal variability of the system and the uncertainties in the multitude of involved processes. Despite the clear statistical significance of the influence of lightning on the nitrate concentrations, the robustness of the findings gradually decreases towards the determination of the radiative flux perturbations.

## 1 Introduction

20 Lightning is one of most energetic phenomenon in the Earth atmosphere. Due to the tremendous elec-  
tricity, the associated temperatures allow for breaking up the stable molecular nitrogen compounds  
into fragments which partly recombine to nitrogen oxides (Schumann and Huntrieser, 2007). Hence  
lightning represents a natural atmospheric emission source for  $\text{NO}_x$ , in addition to the anthropogenic  
sources from industry, energy production, traffic and agriculture (e.g., Jaegle et al., 2005). Further-  
25 more, biomass burning contributes significantly to the total  $\text{NO}_x$  emissions (e.g., van der Werf et al.,  
2010). The global lightning  $\text{NO}_x$  ( $\text{LNO}_x$ ) production has been estimated to range between 2 and  
8 Tg N/yr (Schumann and Huntrieser, 2007), hence being in a similar order of magnitude as the  
soil emissions (e.g., Steinkamp and Lawrence, 2011), which represent an additional important nat-  
ural contribution to the total oxidised nitrogen in the atmosphere. In contrast to the other sources  
30 (with the exception of aircraft emissions),  $\text{LNO}_x$  represents an upper tropospheric source. Due to a  
different chemical composition and chemical reactivity compared to the boundary layer, chemical  
conversion into  $\text{HNO}_3$  is relatively efficient and most important is not subject to fast removal by  
dry deposition. Once nitric acid is formed this can condense on existing aerosol particles, mostly  
thermodynamically stabilised by ammonium ( $\text{NH}_4^+$ ), forming ammoniumnitrate ( $\text{NH}_4\text{NO}_3$ ). This  
35 aerosol species is considered semi-volatile such that in the lowermost atmosphere a substantial part  
of nitrate re-evaporates (e.g., Stelson et al., 1979). However, due to the lower temperatures in the  
upper troposphere,  $\text{NH}_4\text{NO}_3$  is thermally stabilised, and remains mostly in the aerosol phase. The  
total amount of nitrate aerosol does not only depend on the available nitric acid, but also neutralising  
cations, i.e. mostly ammonium. Furthermore, the condensation rate of nitric acid forming nitrates also  
40 depends on the available aerosol concentration, as well as competition for neutralising compounds  
with e.g. sulphate ions.

Experimental evidence for the occurrence of nitrate in convective outflows have recently been ob-  
served over the Amazon by aircraft measurements <sup>1</sup>. Furthermore,  $\text{NH}_3$  has been observed based on  
satellite retrievals in the outflow of the Asian monsoon (Höpfner et al., 2016), leading to the conclu-  
45 sion that not all soluble compounds are completely removed by scavenging during convective lifting.  
Consequently, there is observational evidence for both  $\text{NH}_3$  and  $\text{NO}_3^-$ , such that the formation of  
 $\text{NH}_4\text{NO}_3$  is likely due to the low temperatures in the upper troposphere.

Being a component of the mixture of aerosol particles, the nitrate aerosol from lightning can also  
50 influence atmospheric radiation via the direct and indirect aerosol effects by a multitude of pathways.  
First, nitrate contributes to water uptake increasing ambient aerosol size at a given relative humidity.  
Both the particulate nitrate mass as well as the additional aerosol water enhance the radiative extinc-  
tion properties of the particle and can even alter the single scattering albedo influencing the direct  
aerosol effect of nitrate particles (e.g., Adams et al., 2001).

---

<sup>1</sup>J. Schneider, pers. comm., 2016

55 Additionally, the overall chemical composition of the aerosol is modified by nitrate from light-  
ning such that the cloud condensation nuclei (CCN) or ice nuclei (IN) efficiency of the particles  
changes with implications for the indirect aerosol effects. Consequently, this indirect effect should  
be considered a competition between the formation of additional CCN (e.g., Makkonen et al., 2012)  
by the nitrate and a potential deactivation of IN for contact or deposition freezing by providing a  
60 hygroscopic coating. Immersion freezing is also affected via the concentration of the solutions, IN  
deactivation and freezing point modifications.

Currently, only a few global aerosol chemistry climate models can realistically simulate aerosol  
nitrate and its implications for the direct and partially also indirect aerosol effect due to the complex  
65 chemical interactions and the semivolatility (e.g., Jacobson, 2001; Adams et al., 2001; Bauer et al.,  
2007; Bellouin et al., 2011; Xu and Penner, 2012; Makkonen et al., 2012). However, regional scale  
air quality models often consider the formation of nitrate and also implications of the emissions from  
various sources on the  $\text{NO}_3^-$  formation, e.g. Zare et al. (2014), who investigate the role of  $\text{LNO}_x$  on  
PM<sub>2.5</sub> in Europe or Allen et al. (2012), who among other aspects analysed the influence of lightning  
70 on nitrate wet deposition in the US. The effect of  $\text{LNO}_x$  on gas phase chemistry and the oxidation  
capacity of the atmosphere has been previously studied and is continuously revisited (e.g., Labrador  
et al., 2004; Schumann and Huntrieser, 2007; Gressent et al., 2016; Finney et al., 2016a, b).

Usually, the effect of nitrates is determined from *annihilation studies*, i.e. the effect of nitrates is  
completely ignored. However, this might have implications for the overall aerosol chemical compo-  
75 sition and size distribution such that these effects are mixed with the pure impacts of nitrate. Instead  
of a total nitrate *annihilation scenario*, this study investigates the omission of only a fraction of the  
aerosol nitrate, which originates from the lightning  $\text{NO}_x$  emissions. According to our knowledge, the  
resulting climate impacts from lightning caused by particulate nitrate have not yet been quantified  
with a comprehensive chemistry climate model. Nevertheless, this scenario reveals already strong  
80 impacts due to the substantial implications of  $\text{LNO}_x$  emission on tropospheric chemistry. However,  
emission sensitivity studies on  $\text{LNO}_x$  would suffer from an even worse signal-to-noise ratio than  
found in the current study.

The results from this study are in agreement with previous studies as mentioned above with respect  
to the impact on ozone and the oxidation capacity, but explicitly analyse the impact of  $\text{LNO}_x$  on  
85 upper tropospheric nitrate concentrations and its implications for the climate system.

## 2 Model description

### 2.1 The EMAC modelling system

This study applies the ECHAM5/MESSy Atmospheric Chemistry (EMAC) model, which is a numer-  
ical chemistry and climate simulation system that includes sub-models describing tropospheric and

90 middle atmosphere processes and their interaction with oceans, land and human influences (Jöckel  
et al., 2010). It uses the second version of the Modular Earth Submodel System (MESSy2) (Jöckel  
et al., 2005) to link multi-institutional computer codes. The core atmospheric model is the 5th gen-  
eration European Centre Hamburg general circulation model (ECHAM5, Roeckner et al., 2006).  
For the present study we applied EMAC (ECHAM5 version 5.3.02, MESSy version 2.50) in the  
95 T42L31-resolution, i.e. with a spherical truncation of T42 (corresponding to a quadratic Gaussian  
grid of approx. 2.8 by 2.8 degrees in latitude and longitude) with 31 vertical hybrid pressure levels  
up to 10 hPa. The applied model setup comprised the submodels for radiation, convection, large-  
scale clouds and condensation and the budget (source, transport and loss processes) of chemical  
compounds in the gas and aerosol phase.

## 100 **2.2 Aerosol climate processes within EMAC**

To simulate the relevant processes mentioned above and sketched in Fig. 1, we employed the light-  
ning  $\text{NO}_x$  emissions scheme by Price and Rind (1992). This scheme uses convective cloud top height  
as simulated by the convection scheme (Tiedtke, 1989). Even though such a combination of two  
parameterisations is subject to large uncertainties, we have shown in the past that this special combi-  
105 nation is relatively robust and able to provide relatively realistic lightning distributions compared to  
satellite data from the LIS/OTD sensors (Tost et al., 2007). The emitted NO is subject to gas phase  
chemical transformations which are calculated with the help of the chemical model MECCA (Sander  
et al., 2011). The resulting nitric acid ( $\text{HNO}_3$ ) can subsequently partition into the aerosol phase. The  
corresponding aerosol processes are simulated with the GMXE aerosol submodel (Pringle et al.,  
110 2010; Tost and Pringle, 2012) taking both the gas-aerosol phase partitioning and the interactions  
with other chemical compounds as well as the microphysical properties of the aerosol particles into  
account. The gas-aerosol phase partitioning of inorganic semivolatile compounds is calculated by the  
ISORROPIA2 model (Fountoukis and Nenes, 2007), which is part of the GMXE aerosol scheme.  
The aerosol particles are discretised in 4 lognormal size categories, and for the larger three modes a  
115 distinction is done between internally mixed hydrophilic particles and externally mixed hydropho-  
bic particles resulting overall in 7 lognormal modes. Additional anthropogenic and natural emissions  
except for lightning are simulated with the submodels ONEMIS, OFFEMIS and TNUDGE (Kerk-  
weg et al., 2006b) submodels, providing sources for other primary and secondary aerosol particles.  
Physical loss processes (dry and wet deposition, additionally sedimentation for aerosol processes)  
120 are explicitly considered in the schemes DRYDEP, SCAV and SEDI (Kerkweg et al., 2006a; Tost  
et al., 2006).

To consider aerosol radiation interactions the prognostic aerosol mass and number concentra-  
tions (including aerosol water) are used in lookup tables from Mie calculations precalculated with  
the LIBRADTRAN (Mayer and Kylling, 2005) to determine the radiative properties of the atmo-  
125 spheric aerosol (extinction, single scattering albedo, asymmetry parameter) as described in detail

in Pozzer et al. (2012); de Meij et al. (2012) and Dietmüller et al. (2016). These parameters are explicitly considered in the radiation scheme native to ECHAM5 (Roeckner et al., 2003) replacing the aerosol climatology by Tanre et al. (1984) as described in Dietmüller et al. (2016). For the treatment of indirect aerosol effects, we have implemented a two moment cloud microphysics scheme (Lohmann and Hoose, 2009; Lohmann et al., 2010). The activation of aerosols is calculated with the scheme of Abdul-Razzak and Ghan (2000), which has been adapted to the simulated aerosol types. Furthermore, the interactions of aerosols in the homogeneous and heterogeneous freezing processes (Kärcher et al., 2006) are considered, including an adaption to the more comprehensive chemical composition of the aerosol simulated with GMXE. To investigate the sensitivity of the climate impacts of aerosol particles influenced by lightning caused by aerosol cloud interactions (ACI), a second set of simulations has been performed using a modified activation scheme based on a combination of the work of Abdul-Razzak and Ghan (2000) and Petters and Kreidenweis (2007), as described in detail by Chang et al. (2014). The set of simulations using the Abdul-Razzak and Ghan (2000) aerosol activation scheme will afterwards be abbreviated with ARG, for the other set of simulations following Chang et al. (2014) the KK (i.e. kappa-koehler) acronym will be used.

Both schemes include the effects of nitrate, but different approaches are followed to calculate the critical supersaturation: the ARG scheme uses parameters for the osmotic coefficient and dissociation of nitrate in the solution (which are more uncertain than for  $(\text{NH}_4)_2\text{SO}_4$ ), whereas the KK approach uses a volume weighted  $\kappa$  to determine the total aerosol water uptake and hence cloud formation potential. The two approaches are therefore merely a different representation of the Raoult effect in the cloud activation. Note, that the aerosol-cloud interactions are only taken into account for large-scale clouds as the treatment of cloud microphysics in convective clouds is even further simplified.

### 2.3 Simulation Setup

Decadal simulations including all feedback mechanisms with the EMAC model have been performed for present day and preindustrial conditions. Sea surface temperatures are prescribed by a climatology from the AMIP II database for all model configurations. To determine the effects of aerosols caused by lightning two simulations are performed for each scenario, one with and one without  $\text{LNO}_x$  emissions. Even though this annihilation scenario potentially cannot capture some compensation effects, we choose this approach due to the expected lower signal to noise ratio. Furthermore, as there is nitrate produced from anthropogenic, biomass burning and soil sources of  $\text{NO}_x$ , there is no complete annihilation of all nitrate. For the present day scenario we applied emissions from the AC-CMIP emission inventory (Lamarque et al., 2010, 2013) for trace gases and aerosol emissions from the AEROCOM (Dentener et al., 2006) experiment, for preindustrial conditions we followed the AEROCOM (Dentener et al., 2006) recommendation for preindustrial conditions. Biomass burning is included using the GFED data (van der Werf et al., 2010), as well as a compilation for preindus-

trial biomass burning. Note, that we applied prescribed aerosol emissions for dust and sea salt, not including a potential feedback of a changed circulation (wind speed and wind patterns) on aerosol sources. The model simulations have been initialised with results from previous experiments (see  
165 Jöckel et al. (2016)) eliminating spin-up effects. Due to the comprehensive feedback mechanisms we refrain from comparing individual years of the simulations, but focus on the decadal mean values and distributions.

### 3 Results

#### 3.1 Lightning and associated emissions

170 To estimate the effects of lightning and associated emissions, we have analysed the distribution of  $\text{LNO}_x$  emissions from lightning. Fig. 2 depicts in a 3D visualisation the temporal mean emitted  $\text{LNO}_x$  for the present day scenario.

The magnitude of the  $\text{LNO}_x$  emissions is displayed as an isosurface of  $1 \cdot 10^{-16} \text{ kg}/(\text{m}^3 \text{ s})$ . A second isosurface of  $3 \cdot 10^{-16} \text{ kg}/(\text{m}^3 \text{ s})$ , which is visible as a darker shading embedded in the first  
175 isosurface, shows that the dominant emissions take place in the upper part of the  $\text{LNO}_x$  plume. This is a consequence of the fact that the vertical emissions redistribution follows a C-shape profile according to Pickering et al. (1998), leading to enhanced emissions in the upper section of the plume. Even though more recent studies (e.g., Ott et al., 2010) suggest a different vertical emission distribution, past studies with the EMAC model have shown good agreement using the C-shape profiles  
180 (Huntrieser et al., 2007; Tost et al., 2010).

The outer emission isosurface is colour coded with the total aerosol nitrate mixing ratio (in mol/mol), depicting the amount of aerosol nitrate present in the emission peaks. Furthermore, the gray shaded isosurface visualises a level of  $0.1 \text{ ppb}_v$  of aerosol nitrate. This shows that high aerosol nitrate concentrations are located in the lower troposphere, but also enhanced mixing ratios ( $>100$   
185  $\text{ppt}_v$ ) of particulate nitrate can be found in the upper troposphere, partially concurring with the  $\text{LNO}_x$  emissions (compare the values of up to  $100 \text{ ppt}_v$  as marked by the color coding on the emission plume surfaces).

The enhanced  $\text{NO}_3^-$  mixing ratios in the lower troposphere can be explained by the anthropogenic  $\text{NO}_x$  emissions as well as sources for neutralising cations such as  $\text{NH}_4^+$  close to the surface.

190 The coloured map at the bottom of Fig. 2 depicts the mean flash frequency per  $\text{km}^2$  and minute, which is relatively well confined to the tropical continents and in reasonable agreement with observations from LIS/OTD (updated data following Christian et al. (2003)). Consequently, the  $\text{LNO}_x$  emissions are co-located over the same regions.

The total  $\text{LNO}_x$  emissions are  $5.95 \text{ Tg N/yr}$  with a  $1\sigma$ -variability of  $0.03 \text{ Tg N/yr}$  over the simulated decade in the present day scenario (ARG), for the KK scenario a total emission strength from  
195 lightning with  $6.04 \pm 0.03 \text{ Tg N/yr}$  is simulated. Note that the seasonal variability is substantially

larger. More than 80% of the emitted LNO<sub>x</sub> is placed above 500 hPa altitude in both scenarios. The difference between the two scenarios is a consequence of slightly different meteorological conditions which are caused by the feedback effects of the aerosol and cloud properties.

## 200 3.2 Tropospheric nitrate from lightning

### 3.2.1 Chemical budgets and distribution of oxidised nitrogen compounds

The emitted NO<sub>x</sub> from lightning along with other NO<sub>x</sub> molecules in the atmosphere forms HNO<sub>3</sub> mostly via the gas phase reaction of NO<sub>2</sub> + OH → HNO<sub>3</sub> on typical time scales of hours to days. Even though this reaction is much slower than the conversion between NO and NO<sub>2</sub>, it represents  
205 an important end of the circular conversion of the NO<sub>x</sub> species.

The LNO<sub>x</sub> emissions for the present day scenario result in ~ 40% higher nitric acid mixing ratios in the troposphere, and even up to ~ 61% between 500 hPa and the tropopause, compared to simulations without lightning NO<sub>x</sub> emissions (see Tab. 1). Even though the enhancement effect for N<sub>2</sub>O<sub>5</sub> is stronger in relative numbers, the overall mixing ratios for N<sub>2</sub>O<sub>5</sub> are substantially smaller  
210 (almost two orders of magnitude) such that dinitrogenpentoxide plays a minor role for the highly oxidised nitrogen compounds in the troposphere. PAN is found to see a reduction by 15% for the whole troposphere and 21% in the upper troposphere when no lightning emissions are considered, therefore being less affected by the LNO<sub>x</sub> emissions due to a potential limitation of VOCs for PAN formation.

215 The nitric acid molecules can condense on pre-existing aerosol particles forming NO<sub>3</sub><sup>-</sup> ions in the deliquesced aerosol solution or solid NH<sub>4</sub>NO<sub>3</sub> or NaNO<sub>3</sub> crystals. Tropospheric nitrate mixing ratios are typically a factor of two to three lower compared to HNO<sub>3</sub>. Nevertheless, the changes in upper tropospheric NO<sub>3</sub><sup>-</sup> mixing ratios are ~ 32% globally averaged.

Fig. 3 shows isosurfaces of tropospheric nitrate concentration differences between the simulation  
220 with LNO<sub>x</sub> emissions to the case without this source. The gray shaded isosurface depicts a 30% difference, the blue isosurface 45% differences and the red enclosed area 60% enhanced NO<sub>3</sub><sup>-</sup> mixing ratios, with only statistically significant (based on a two sided t-test of annual mean data with a significance level of 90%) data shown. Even though the maximum NO<sub>3</sub><sup>-</sup> absolute differences occur down to the surface, the largest relative differences are apparent in the UT. Most of these differ-  
225 ences are constrained to the tropics where the strongest LNO<sub>x</sub> emissions are prevalent. However, the maxima in the differences are not directly co-located with the emission maxima (i.e. Central Africa, Amazonia and the maritime continent, see Fig. 2), but generally further downwind. This is a consequence of atmospheric transport during the time required for oxidation of NO<sub>x</sub> to HNO<sub>3</sub> and subsequent partitioning into the aerosol phase.

230 To provide an estimate of the total amount of nitrate in the upper troposphere the back panel of Fig. 3 depicts the zonal mean  $\text{NO}_3^-$  mixing ratio with values between 10 and 100 ppt<sub>v</sub> for the tropical middle and upper troposphere where the impact of the  $\text{LNO}_x$  emissions is largest.

The mean upper tropospheric column burden (500 hPa to the tropopause) in  $\text{mg/m}^2$  is depicted by the coloured panel at the bottom of the plot. The turquoise isolines on this panel depict the 20%, 235 40% and 60% differences in the UT column burden between the simulations with and without  $\text{LNO}_x$  emissions. These are of course co-located with the regions of the isosurfaces.

The figure shows that the tropical upper troposphere is a region in which substantial amounts of particulate nitrate can be found. Both visualisations of the differences (UT column burden and mixing ratios) show that a large contribution of the upper tropospheric  $\text{NO}_3^-$  originates from lightning 240 emissions, but also that a sufficient amount of neutralising cations is available to stabilise the nitrate in the particulate phase. These findings are a direct consequence of the emission and conversion of  $\text{NO}_x$  to  $\text{N(V)}$ , but of course are also indirectly affected by changes in the oxidation capacity of the atmosphere and hence the  $\text{HNO}_3$  formation.

The tropospheric budget of the highly oxidised N compounds is summarised in Tab. 1, including 245 the relative importance of the  $\text{LNO}_x$  emissions. For present day conditions, the particulate nitrate contribution to the total  $\text{N(V)}$  is  $\sim 25\%$ , whereas in the upper troposphere it is only  $\sim 13\%$ .

The neglect of the  $\text{LNO}_x$  emissions leads to a shift in the contribution of the particulate phase to the total  $\text{N(V)}$ . Under these conditions 33% of the  $\text{N(V)}$  is in the form of particulate  $\text{NO}_3^-$  for the whole troposphere, whereas for the UT the fraction is higher than 21%. This is a consequence of the 250 reduced available  $\text{N(V)}$ , but a similar amount of neutralising cations, i.e. mostly  $\text{NH}_4^+$ .

The sensitivity simulations in the KK configuration result in almost identical values, (budget is shown in Tab. 1 of the supplement) depicting the fact that the description of cloud and cloud removal processes plays a minor role for the total budget of particulate nitrate in both model configurations.

The budget of the loss processes of  $\text{N(V)}$  compounds can be found in the supplement for both 255 present day and preindustrial conditions. The differences in the nitrate burden are similarly present in the loss processes, with wet deposition being mostly influenced (50% change in the loss flux), but sedimentation and dry deposition also equally contribute to the change in the total loss.

### 3.2.2 Influences on other chemical species

The reduced  $\text{NO}_x$  burden in the simulation without lightning emissions has substantial impact on 260 the concentrations of other species. As only a minor fraction of the total  $\text{LNO}_x$  emissions of  $\sim 6$  Tg  $\text{N/yr}$  are converted to  $\text{N(V+)}$  (e.g. 186 Gg  $\text{N}$  change in  $\text{N(V)}$  according to Tab. 1), the impact on ozone is comparable to previous studies (e.g., Labrador et al., 2005; Finney et al., 2016a): the calculated values for tropospheric ozone are 362 Tg in the simulation without lightning and 444 Tg if lightning is included, which corresponds to a 22% increase in the tropospheric ozone burden 265 (even though the absolute values for the tropospheric ozone burden differ substantially in each of the



studies). The effect of the LNO<sub>x</sub> emissions is again very prominent in the upper troposphere, where the O<sub>3</sub> load increases from 195 to 248 Tg by LNO<sub>x</sub> emissions, i.e. an increase of 27%. Driven by the location of most of the emissions, these effects mostly occur in the low latitudes. The resulting decrease in ozone in the zonal average resulting from neglecting lightning emissions is comparable  
270 to the findings of Grewe (2007). The change in the column burdens in the tropics also agrees with the results of Martin et al. (2002), both in the geographic patterns and the amplitude of the signal. Therefore concluding, the explicit consideration of the nitrate formation has no important impact on the tropospheric ozone distribution, such that the results of impact studies of lightning NO<sub>x</sub> (such as e.g., Banerjee et al., 2014) do not have to be revised.

275 However, the OH concentrations and therefore the oxidation capacity of the atmosphere are also substantially influenced by lightning (Labrador et al., 2004) via changes in O<sub>3</sub> and subsequent OH production via ozone photolysis and reaction of the products with water vapour. The effect on OH is displayed by a modification of the methane lifetime, as depicted in Tab. 2. The emissions of LNO<sub>x</sub> are responsible for an increase of the tropospheric methane lifetime of ~ 1.7 to 1.9 years; however  
280 in the upper troposphere (above 500 hPa up to the tropopause) an increase of the CH<sub>4</sub> lifetime of almost 10 and 13.2 years for present and preindustrial conditions, respectively, corresponding to almost a halving of the oxidation capacity of the upper troposphere.

This is especially relevant for the production of sulphate via the gas phase oxidation of SO<sub>2</sub>, which substantially influences aerosol formation and aerosol composition in the upper troposphere.  
285 The sulphate burden for present day conditions decreases from 576 Gg S(VI) to 565 Gg. This is also prominent in the UT region, where instead of 74.6 Gg S(VI) only 70.2 Gg S(VI) are simulated. As most of the oxidation in the upper troposphere takes place in the gas phase, the aqueous phase oxidation in the lower troposphere is affected to a minor degree.

Particulate ammonium mixing ratios are only affected to a minor degree by the LNO<sub>x</sub> emissions:  
290 for present day conditions the NH<sub>4</sub><sup>+</sup> burden slightly decreases without lightning emissions, but increases during preindustrial times. However, the changes in the atmospheric burden are lower than ±5%. These changes can be explained by the of lower sulphate and nitrate burdens and effects of the oxidation capacity on gaseous ammonium concentrations. A significant change due to LNO<sub>x</sub> emissions in the sulphate to bisulphate ratio is not simulated by the model despite the capabilities of  
295 the thermodynamic equilibrium model.

The KK simulations show a very similar behaviour in the burdens, the CH<sub>4</sub> lifetime and the changes induced by the omission of the LNO<sub>x</sub> emissions, such that a detailed discussion is skipped here.

### 3.3 Aerosol microphysical effects

300 The influences of the chemical composition in the upper troposphere due to lightning are also reflected in the microphysical properties of the aerosol, which are described by the size distributions

at various locations. The regions in the tropics are mostly characteristic for the tropical continents, as the oceanic aerosol distributions usually contain substantially lower particle concentrations such that the distribution of those regions is dominated by the continental grid cells. The regions have  
305 been selected following the analysis of the significant changes in Sect. 3.2.1, but also to represent various conditions of typical aerosol size distributions where lightning can play an important or only a minor role.

The temporal mean size distributions of the aerosol particles for various regions (i.e. those regions covered by the plot on the map) are shown in Fig. 4 as a 2D plot with the horizontal axis depicting the  
310 ambient aerosol diameter and the vertical axis the pressure altitude; the colour coding displays the relative difference in aerosol number concentration [in %] between the simulation with and without lightning. Furthermore, the absolute values of the size distribution [in  $\text{cm}^{-3}$ ] are depicted by the contour lines with the solid lines representing concentrations larger than 1 per  $\text{cm}^3$ . The figure shows the simulation results for the present day conditions (ARG case). The hatched regions in the 2D size  
315 distribution diagram mark those areas in which the changes do not exhibit statistical significance.

Over the tropical continents in the upper troposphere enhanced nucleation mode particle numbers are simulated if lightning emissions are considered, which is a result of the enhanced S(VI) concentrations and consequently enhanced new particle formation. At approximately 10 nm diameter the particle number is smaller in case of  $\text{LNO}_x$  emissions, due to the interaction of enhanced coagulation and condensation on the enhanced particle concentrations. Also additional condensation on  
320 existing particles and enhanced coagulation with the small particles causes the reduction of particle numbers at  $\sim 80$  nm. For larger particles a slight reduction is found as a consequence of the more efficient coagulation (due to higher particle numbers in the small mode) versus the slower condensational growth. For the lower troposphere the picture is more ambiguous and most changes are not  
325 statistically significant: in South America and Indonesia reduced nucleation takes place. However, the absolute particle concentrations are so low that this process is almost negligible, whereas in Central Africa enhanced new particle formation is caused by lightning emissions. For the larger particles the impact is rather small.

In the mid-latitudes relevant changes in the size distributions are only found in the Eastern US  
330 and China but little impact in Europe and Siberia. The first two regions are located slightly more southwards such that lightning frequencies are enhanced compared to the latter two regions (see bottom map of Fig. 2). The general pattern of the changes is comparable to the tropical continents.

In the Southern Atlantic region the changes in the size distribution profile are relatively small, as lightning emissions play a minor role for the total particle concentration as well as for the oxidation  
335 capacity of the atmosphere and therefore sulphate formation, but they are still robust. In the Central Pacific where enhanced nitrate concentrations are simulated in the upper troposphere (see Fig. 3), the changes in the aerosol size distribution are moderate with strongest signals in the middle troposphere (between 700 and 500 hPa). As lightning  $\text{NO}_x$  emissions are substantially smaller over the ocean, the

oxidation capacity is affected to a minor degree; however nitrate and gaseous N(V) are transported  
340 downwind from the source region affecting the nitrate concentrations and particle numbers, mostly  
via coagulation.

To analyse the impact of the LNO<sub>x</sub> emissions on the aerosol water uptake, the mean growth  
factor (GF = ambient diameter / dry diameter) is compared. Using this parameter has the advantage  
that, e.g. in contrast to the aerosol water content, it is independent of the aerosol size and particle  
345 number concentrations. The overall impact of lightning on the GF is relatively small: for present day  
conditions the nucleation and aiten mode depict a slight increase in the upper troposphere (up to  
2%), whereas in the accumulation and coarse mode a decrease of up to -2% is simulated caused  
by LNO<sub>x</sub> emissions. Due to the relatively dry conditions in the upper troposphere, where the nitrate  
changes are strongest, the effect on water uptake is expected to be that small. Closer to the surface  
350 the differences in the growth factor are even smaller and ambiguous in their sign, lacking statistical  
significance.

### 3.4 Impacts on climate

#### 3.4.1 Aerosol optical properties

As aerosol particles scatter and absorb solar and infrared radiation, the impact of the LNO<sub>x</sub> emis-  
355 sions is investigated with the help of Fig. 5 for the ARG present day simulation. The left panel  
shows both the mean column AOD at 550 nm wavelength (on the floor level) as well as the zonal  
mean extinction (rear panel). The ceiling panel depicts statistical significant changes in the column  
AOD and the front panel visualises statistical significant changes in the extinction. Note, that for  
determining the zonal mean of statistical significant changes, only the individual data points which  
360 exhibit significance are selected and the average over these (selected) points is taken.

The rear panel depicts the zonal mean of the average extinction per km. Substantial extinction  
is simulated in the lower troposphere (below 700 hPa) with an additional enhancement of the op-  
tical depth between 20° and 50°N. This is a consequence of both the distribution of water vapour  
and hence aerosol water supporting aerosol growth and extinction in the lower troposphere of the  
365 tropics, on the other hand driven by natural sources of dust (and associated extinction) as well as  
anthropogenic pollution (especially from East Asia). The relative changes of aerosol extinction due  
to LNO<sub>x</sub> are depicted in the front panel. The most substantial enhancement of aerosol extinction is  
simulated in the middle to upper troposphere in the tropics with zonal mean enhancements higher  
than 20%. In the uppermost troposphere reductions in extinction are also simulated as a consequence  
370 of neglecting the lightning emissions. Compared to the nitrate enhancements as analysed from Fig. 3,  
the maximum enhancement of aerosol extinction by LNO<sub>x</sub> is located further downwards in the mid-  
dle troposphere, i.e. between 400 and 600 hPa, whereas the strongest nitrate enhancement has been  
simulated between 200 and 400 hPa.

The column AOD (floor panel) shows maximum values in the dust emission regions (e.g. Northern Africa) and the anthropogenic pollution centers (e.g. Eastern China). The climatological global mean value with present day emissions is 0.122, which is close to the observations as derived from MODIS (e.g., Mao et al., 2014). The influence of the LNO<sub>x</sub> emissions can be analysed with the help of the ceiling map in Fig. 5, which shows the percentage fractions of the changes in the simulated AOD. Globally a slight reduction to 0.121 can be found ( $\sim 1\%$ ). However, in some regions even higher extinction is simulated in case of no nitrate formation from lightning, which is in contrast to the findings of Sect. 3.2.1, where no regions with enhanced nitrate have been simulated. However, only in very few regions a statistical significant signal in the column AOD can be found. This is a consequence of the fact that significant changes in extinction are mostly located in the upper troposphere which contributes less to total column AOD (cf. rear panel). The right panel of Fig. 5 depicts again the absolute values of AOD (floor panel) and extinction (rear panel), but also shows isosurfaces of the location of the regions of enhancement and reduction of extinction. The pale red depicts the +10% isosurface, whereas the embedded dark red regions mark an increase of the extinction due to LNO<sub>x</sub> of more than +20%. On the other hand the pale blue marks regions of a medium reduction ( $-10\%$ ) and dark blue regions (only at the tropopause) depict extinction reductions of more than 20%. The main enhancement of extinction by lightning emissions occur between 400 and 600 hPa over the tropical continents and slightly further downwind. This general pattern corresponds to the enhanced nitrate mixing ratios (c.f. Fig. 3), but is more restricted to the continents. In the extra-tropics an even higher extinction is simulated if no LNO<sub>x</sub> emissions are considered. This is a consequence of the secondary effects caused by the emissions, i.e. the influence on sulphur oxidation via the oxidation capacity of the atmosphere and changes in the size distribution. Comparing the pattern correlation between the changes in nitrate and extinction a relatively low value is found ( $R \approx 0.3$ ). However, the changes in sulphate and extinction show a higher correlation value of  $R \approx 0.4$ . Analysing the PDF of the differences in AOD between the simulation with and without lightning emissions (see supplement) reveals that for present day conditions larger deviations occur more often compared to the preindustrial cases. The PDFs are almost symmetric, but show a slight shift towards enhanced extinction in the simulations including LNO<sub>x</sub> emissions.

The sensitivity simulations with the alternative warm cloud activation scheme in general show a similar distribution of the aerosol extinction with the enhancement in the middle and upper tropical troposphere (see supplement). The global mean column AOD for present day conditions is slightly lower (0.120) and the reduction in case of no LNO<sub>x</sub> emissions even further reduced ( $\sim 0.5\%$ ). Strongest enhancement is found over the tropical continents, i.e. the regions with maximum lightning activity, but most of the nitrate is simulated in the upper troposphere, whereas also in this configuration the extinction is enhanced in the middle to upper troposphere. For preindustrial conditions, the enhancement of the extinction by lightning is a bit stronger, but still small ( $\sim 1\%$ ).

410 To analyse whether the contribution of scattering versus absorption is influenced by the LNO<sub>x</sub>  
emissions, the single scattering albedo is compared in both model simulations. The zonal mean value  
changes by less than  $\sim +1\%$  in the free troposphere and stratosphere, and is reduced by a maximum  
of  $\sim -4\%$  in the boundary layer when lightning emissions are considered. Consequently, a change  
in the thermodynamic structure resulting from direct aerosol-radiation interactions is expected to be  
415 small.

### 3.4.2 Cloud properties

As cloud properties can be influenced indirectly by the LNO<sub>x</sub> emissions three parameters are anal-  
ysed to provide the causes for aerosol-cloud-radiation interactions. The cloud cover may hint at  
modifications of the cloud lifetime effect (Albrecht, 1989), whereas the combination of the liquid  
420 (ice) water content and the cloud droplet (ice crystal) numbers, which can also be expressed via the  
effective radii, can shed light onto the direct influence of the lightning emissions on the radiative  
properties of the clouds via the Twomey (1977) effect.

- Cloud cover:

425 The cloud and precipitation cover changes only to a minor degree between the simulations  
with and without lightning for the present day scenario (ARG case). In the tropics the LNO<sub>x</sub>  
emissions lead to a small increase in the mid troposphere cloud coverage of the Northern  
hemisphere, whereas the dipole pattern is compensated by reduced cloudiness in the Southern  
hemisphere. Apart from this, only the polar latitudes show some larger (but not statistical  
significant) changes in cloud coverage. Consequently, the impact of the cloud lifetime effect  
430 is expected to be small.

- Liquid / ice water content:

435 The patterns for cloud water and cloud ice are characterised by local dipole changes, but there  
are no significant modifications of the distribution of cloud water and ice, respectively. Only  
the tropical middle to upper troposphere of the Northern hemisphere shows an enhanced ice  
mixing ratio in the simulation with lightning emissions, which correlates to the increase in  
cloud coverage. However, the decrease in cloud coverage in the corresponding region in the  
Southern hemisphere is not accompanied by a substantial decrease in ice water. In the zonal  
average the increase in total water is less than 10%.

- Cloud droplet / ice crystal numbers:

440 Compared to the other two parameters, some substantial changes in cloud droplet and es-  
pecially in ice crystal number are found as a consequence of the LNO<sub>x</sub> emissions. For the  
present day scenario, in the troposphere between 650 and 400 hPa an increase in cloud droplets  
is simulated in case of active lightning emissions in the tropics. Further North a decrease of  
cloud droplets between 900 and 600 hPa is calculated. Both are of approximately the same

445 magnitude both in absolute and in relative terms and are a consequence of more hydrophilic  
particles due to nitrate coating and a reduction of lower tropospheric CCN due to reduction in  
lifetime and faster conversion from hydrophobic to hydrophilic conditions. However, the sta-  
tistical significance of the changes is relatively low. For the ice crystal number the influence  
is even stronger: in the tropical upper troposphere the LNO<sub>x</sub> emissions lead to an increase in  
450 the ice crystal number in a region where the highest ice crystal numbers are simulated. In the  
mid-latitudes a reduction of the ice crystal numbers is simulated in both hemispheres above  
400 hPa. Both modifications are of the order of more than  $\pm 20\%$ . Due to the coarse tempo-  
ral resolution of the model output it is not possible to attribute the differences to a change in  
regime from homogeneous to heterogeneous freezing. Nevertheless, it is possible that locally  
455 this effect can occur and has a substantial influence on the ice crystal number concentration.

By Analysing the effective radii (Fig. 6), some of the effects on water mixing ratios and numbers  
compensate, whereas others positively amplify each other. This is both valid for liquid clouds in  
the lower and middle troposphere as well as ice clouds in the middle and upper troposphere. For  
liquid clouds, the effective radii increase in the simulation with lightning in the polewards of 40°  
460 below 800 hPa and in the tropics (0° to 20°N) up to 400 hPa. However, most of the changes are not  
statistically significant, such that Fig. 6 depicts only the changes in ice crystal size; a corresponding  
figure for the liquid clouds can be found in the supplement. The back panel depicts in colours the  
statistical significant zonal mean change in effective ice crystal size (in absolute units of  $\mu\text{m}$ ). The  
white contours represent absolute ice crystal sizes with maximum values in the upper part of the mid-  
465 latitude storm tracks and the upper tropical troposphere. Note, that the absolute values (not marked)  
are of secondary importance as mean values over cloudy and non-cloudy conditions are taken), but  
the difference is robust. For ice clouds the effective radii also increase in the same tropical latitude  
range between 600 and 200 hPa. Additionally, a decrease in effective radii of similar magnitude  
on the other side of the ITCZ is simulated. Also in the high latitudes the effective ice crystal size  
470 mostly increases in the simulation with LNO<sub>x</sub> emissions, both in the upper troposphere as well as  
in the boundary layer. The shape of the regions of ice crystal changes extends vertically over more  
than 100 hPa and is not located in the uppermost troposphere only. This indicates that the statistical  
significant changes are more associated to the process of immersion freezing in mixed phase clouds  
than to the cirrus regime with competition between homogeneous and heterogeneous freezing.

475 Using the KK model configuration, which applies a different cloud droplet activation scheme,  
results in a lower sensitivity of the liquid droplet effective radius at elevated altitude for present day  
conditions. However, in the lower troposphere a similar strong signal is simulated, which leads to  
an increase of effective droplet radius in the tropical Southern hemisphere and a decrease in the  
Northern counterpart. For ice crystals, mostly a reduction of effective crystal size is simulated which  
480 is a consequence of the liquid droplet freezing. The effects for preindustrial conditions are similar,  
with even lower sensitivity of the low clouds on the lightning emissions.

In general, the regions with positive and negative effects in the effective droplet and crystal size show ambiguous signals, such that the impact on cloud radiative properties can only be addressed in conjunction with cloud coverage and the calculation of the radiative fluxes.

### 485 3.4.3 Radiative Fluxes

The combined effects of modifications in the aerosol extinction and the cloud optical properties lead to a modification of the shortwave and longwave radiation fluxes. In Fig. 7 the mean impact on the total sky shortwave flux is depicted in the upper panel. Shown are the differences in the SW flux at the top of the atmosphere between a simulation with and one without LNO<sub>x</sub> emissions (ARG configuration). Despite the relatively large internal variability of the system, several regions show a statistical significant change in the TOA short wave flux (based on a t-test with 90% significance threshold on the annual mean data), which are marked by the hatched regions. However, the regions with increased or decreased fluxes are not directly co-located with the LNO<sub>x</sub> emissions. Downwind of Central Africa an increased TOA short wave flux (corresponding to a heating) is simulated, whereas in Northern Amazonia and the maritime continent a decrease of the fluxes (corresponding to a cooling) is calculated. Also in the mid-latitude storm tracks mostly a slight increase in the fluxes is simulated, whereas over the tropical oceans the model suggests decreased fluxes. In the global mean the overall effect is determined as an enhanced backward reflection of the shortwave radiation of  $\sim -0.1 \text{ W/m}^2$  for all sky conditions. However, the time series of the global mean differences (depicted in the lower panel of Fig. 7) is characterised by large internal variability with both positive and negative differences. The clear sky fluxes show a substantially weaker signal with only  $\sim -0.05 \text{ W/m}^2$  and hardly any regions with statistical significance. Furthermore, the patterns of substantial changes are clearly separated from the regions where lightning is dominant (i.e. mid-latitudes versus tropical continents). Consequently, the clear sky signal is interpreted as statistical noise.

505 Considering both the temporal variability of the mean and the large spatial variability, the uncertainty in the total effect has to be considered as relatively large. Using the sensitivity simulations with the alternative cloud activation scheme reveals for both present day and preindustrial conditions a shortwave flux perturbation of  $\sim -0.06 \text{ W/m}^2$  again with no significant contribution from the clear sky fluxes.

To bring these numbers into context with the total shortwave flux disturbances from anthropogenic aerosols, simulation results with both cloud activation schemes and present day and preindustrial emission scenarios are intercompared (for figures see supplement). Both model configurations show a statistical significant cooling over the regions of dominant anthropogenic aerosol pollution (Eastern US, Europe, China) for both clear sky and all sky conditions. The clear sky disturbance is  $\sim -0.57$  for the ARG and  $\sim -0.50 \text{ W/m}^2$  for the KK configurations. All sky fluxes change by  $\sim -1.62$  for the ARG and  $\sim -1.42 \text{ W/m}^2$  for the KK simulation setups, respectively. The overall anthropogenic

aerosol effect is relatively large in the simulations compared to the results published in the latest IPCC report (IPCC, 2013) and therefore an overestimated sensitivity to the aerosol disturbance by lightning  $\text{NO}_x$  nitrate cannot be ruled out completely.

The effect on the longwave radiation via both aerosol extinction and cloud effects is relatively small. Especially, the clear sky flux disturbances are almost negligible, such that the total effect is dominated by ACI.

Consequently, as the differences between both cloud activation schemes indicate an uncertainty of almost 50% the quantitative estimate of  $-100 \text{ mW/m}^2$  should also be used with a comparable uncertainty range of  $50 \text{ mW/m}^2$ .

#### 4 Conclusions

The overall impact of chemically produced aerosol nitrate from lightning  $\text{NO}_x$  emissions is analysed with a global chemistry climate model. Based on a total emission of  $6 \text{ Tg N/yr}$ , the contribution of  $\text{LNO}_x$  to the concentrations of upper tropospheric oxidised nitrogen by more than 57% for present day and 75% for preindustrial conditions. Given a sufficient amount of neutralising cations, especially  $\text{NH}_3$ , substantial amounts of aerosol nitrate are formed with the increases in the upper tropospheric burden exceeding 30% for present day and 50% for preindustrial conditions. Therefore, lightning is also a major contributor to the aerosol nitrate burden in the upper troposphere. The concentration enhancements are not uniformly distributed, but follow the regions of maximum emissions, and further downwind transport.

Besides the well known impacts of lightning on  $\text{O}_3$ , the impact on the chemical oxidation capacity of the atmosphere is highly important for the upper tropospheric aerosol loading. Because of a decrease of the oxidation potential (as represented by an increase in the methane lifetime), the sulphate formation via the gas phase reaction pathway is reduced in case of neglecting  $\text{LNO}_x$  emissions. Hence new particle formation is suppressed, efficiently modifying the small part of the aerosol size distribution. In addition to the condensation of nitrate on the aerosol particles, the combination of a reduced particle number and the coagulation of the particles, substantial effects on the particle size distribution in the upper troposphere are simulated.

The changes in aerosol size, chemical composition and number concentrations have implications on climate via both aerosol-radiation and aerosol-cloud interactions. Lightning and its consequences (nitrate formation, size distribution changes) causes mostly an increase in the total aerosol extinction with pronounced increase maxima in the upper troposphere. On the other hand, both liquid and ice phase cloud optical properties are modified by the  $\text{LNO}_x$  emissions, represented by e.g. effective droplet and ice crystal size. However, in contrast to the nitrate concentration increases the cloud effects resulting from lightning are ambiguous.



The resulting shortwave flux perturbations caused by LNO<sub>x</sub> emissions are quantified to be ~  
–100 mW/m<sup>2</sup>. The resulting effect is caused mostly by the aerosol-cloud interactions, whereas the  
direct aerosol-radiation interactions are of secondary importance. However, an uncertainty range of  
555 almost ~ 50 mW/m<sup>2</sup> has to be assumed due to large internal variability and uncertainties in the  
process description (mostly in the aerosol-cloud interactions).

Nevertheless, an increase in lightning activity in the future or impacts of a more efficient trans-  
formation of the LNO<sub>x</sub> emissions into nitrate under future climate conditions might have a non-  
negligible impact on the radiation balance of the atmosphere. On the other hand, aerosol nitrate  
560 formed from lightning offers further possibilities to address the feedback and (potentially compen-  
sating) impacts of combined chemistry-aerosol-climate interactions.

## Appendix A: Preindustrial conditions

The results from the sensitivity simulations for preindustrial conditions are summarised in this sec-  
tion.

565 – Emissions:

In the preindustrial scenario total LNO<sub>x</sub> emissions of  $6.12 \pm 0.03$  Tg N/yr are simulated for  
the ARG scenario and  $6.16 \pm 0.03$  Tg N/yr for the KK simulations.

– Nitrate enhancement:

570 During preindustrial times, an increase in the HNO<sub>3</sub> burden with ~ 67% and ~ 76% in the  
total and upper troposphere is even stronger compared to present day conditions. This stronger  
enhancement is due to the larger contribution of the LNO<sub>x</sub> emissions compared to the total  
NO<sub>x</sub> release, which is mostly from anthropogenic sources in present day conditions. Enhance-  
ments for N<sub>2</sub>O<sub>5</sub> are of ~ 80%, but also under these conditions N<sub>2</sub>O<sub>5</sub> is a minor contributor to  
the total highly oxidised nitrogen tropospheric load (less than 5%). A similar figure as Fig. 3  
575 for preindustrial conditions can be found in the supplement. Due to the lower other sources  
of NO<sub>x</sub> in the atmosphere, the importance of the LNO<sub>x</sub> emissions is substantially increased.  
Therefore, the relative differences in NO<sub>3</sub><sup>-</sup> are larger which can be seen from the extended  
areas included by the respective isosurfaces and the larger areas covered by the turquoise con-  
tour lines at the floor panel of the figure. The lower overall nitrate mixing ratios are obvious  
580 from the colour scale of the column nitrate which is almost an order of magnitude lower com-  
pared to present day conditions. A lower contribution of particulate nitrate to the total N(V)  
load with 22% for the whole and 9% for the upper troposphere is simulated, which can be  
mostly attributed to lower NH<sub>3</sub> emissions and consequently less NH<sub>4</sub><sup>+</sup> ions to neutralise and  
thermodynamically stabilise available aerosol nitrate. However, the enhancement of the par-  
585 ticulate phase due to LNO<sub>x</sub> is even stronger compared to present day conditions to ~ 34%  
for the whole troposphere and 17% for the UT.

– Methane lifetime and sulphate:

The lifetime increase is even stronger without anthropogenic emissions, such that for the upper troposphere almost a doubling of the CH<sub>4</sub> lifetime occurs (see lower part of Tab. 2). Nevertheless, this results in a similar reduction of the sulphate burden compared to present day without LNO<sub>x</sub> emissions despite the substantially lower total SO<sub>2</sub> emissions.

– Size distributions:

For preindustrial conditions (see supplement) the simulation provides similar results from the tropics. However, in the mid-latitudes where a substantial reduction of other NO<sub>x</sub> emissions is applied compared to the present day scenario the impact of lightning on the nitrogen budget as well as the size distribution becomes more important. Obviously, a similar pattern as for the tropical continents is simulated; however, due to the weaker LNO<sub>x</sub> emissions in the mid-latitudes the effects are substantially smaller compared to the tropics.

– Growth factors:

The results for the GF are similar in their distribution compared to the present day scenario with a slightly increased amplitude (up to ±5%). The sensitivity simulations *KK* are characterised by the same distribution and dependencies on the emissions by lightning. Corresponding figures can be found in the supplement.

– Extinction and AOD:

For preindustrial conditions the situation is quite comparable (see supplement), despite the intensive AOD signals in the regions with anthropogenic pollution, especially East Asia. The simulated global mean column AOD at 550 nm is 0.090. Consequently, the enhancement of the extinction between 20° and 50°N is less strong. The neglect of the lightning emissions results in a reduced column AOD in most regions, resulting in a global mean value of 0.089. A reduction in column AOD occurs only in regions of strong hydrophobic emissions (dust or BC) where the aerosol lifetime is reduced by a coating by nitrates and subsequent faster conversion from hydrophobic to hydrophilic categories. However, also in this case the statistical significance is relatively small. With respect to the zonal mean extinction the changes are slightly larger compared to present day conditions.

– Cloud response:

For preindustrial conditions the response to the LNO<sub>x</sub> emissions is comparable, but shows a slightly higher amplitude.

– Radiative fluxes:

For preindustrial conditions (for a graph see supplement) the situation is comparable with a slightly increased amplitude of  $\sim -0.14 \text{ W/m}^2$  and a similar pattern distribution. The clear sky signal is almost negligible with  $\sim -0.01 \text{ W/m}^2$  and no statistical significance.

*Acknowledgements.* Parts of this research were conducted using the supercomputer Mogon and advisory services offered by Johannes Gutenberg University Mainz ([www.hpc.uni-mainz.de](http://www.hpc.uni-mainz.de)), which is a member of the AHRP and the Gauss Alliance e.V. The authors gratefully acknowledge the computing time granted on the  
625 supercomputer Mogon at Johannes Gutenberg University Mainz ([www.hpc.uni-mainz.de](http://www.hpc.uni-mainz.de)).

The author wishes to acknowledge use of the PARAVIEW program by the Sandia National Laboratories. Sandia is a multiprogram laboratory operated by Sandia Corporation, a Lockheed Martin Company, for the United States Department of Energy's National Nuclear Security Administration under contract DE-AC04-94AL85000.

630 The author wishes to acknowledge use of the Ferret program for analysis and graphics in this paper. Ferret is a product of NOAA's Pacific Marine Environmental Laboratory; further information is available at <http://ferret.pmel.noaa.gov/Ferret/>.

Furthermore, the author thanks all MESSy developers and users for support; further information can be found at <http://www.messy-interface.org>.

635 The author thanks the two anonymous reviewers for their constructive comments, which helped to improve this manuscript.

## References

- Abdul-Razzak, H. and Ghan, S.: A parameterization of aerosol activation: 2. Multiple aerosol types, *Journal of Geophysical Research: Atmospheres*, 105, 6837–6844, 2000.
- 640 Adams, P. J., Seinfeld, J. H., Koch, D., Mickley, L., and Jacob, D.: General circulation model assessment of direct radiative forcing by the sulfate-nitrate-ammonium-water inorganic aerosol system, *Journal of Geophysical Research: Atmospheres*, 106, 1097–1111, 2001.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, 1989.
- Allen, D. J., Pickering, K. E., Pinder, R. W., Henderson, B. H., Appel, K. W., and Prados, A.: Impact of  
645 lightning-NO on eastern United States photochemistry during the summer of 2006 as determined using the CMAQ model, *Atmospheric Chemistry and Physics*, 12, 1737–1758, doi:10.5194/acp-12-1737-2012, 2012.
- Banerjee, A., Archibald, A. T., Maycock, A. C., Telford, P., Abraham, N. L., Yang, X., Braesicke, P., and Pyle, J. A.: Lightning NO<sub>x</sub>, a key chemistry–climate interaction: impacts of future climate change and consequences for tropospheric oxidising capacity, *Atmospheric Chemistry and Physics*, 14, 9871–9881,  
650 doi:10.5194/acp-14-9871-2014, 2014.
- Bauer, S. E., Koch, D., Unger, N., Metzger, S. M., Shindell, D. T., and Streets, D. G.: Nitrate aerosols today and in 2030: a global simulation including aerosols and tropospheric ozone, *Atmospheric Chemistry and Physics*, 7, 5043–5059, 2007.
- Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the Climate  
655 Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, *Journal of Geophysical Research: Atmospheres*, 116, n/a–n/a, 2011.
- Chang, D. Y., Tost, H., Steil, B., and Lelieveld, J.: Aerosol–cloud interactions studied with the chemistry-climate model EMAC, *Atmospheric Chemistry and Physics Discussions*, 14, 21 975–22 043, 2014.
- Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Good-  
660 man, S. J., Hall, J. M., Koshak, W. J., Mach, D. M., and Stewart, M. F.: Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *J. Geophys. Res.*, 108, 4005, doi:10.1029/2002JD002347, 2003.
- de Meij, A., Pozzer, A., Pringle, K. J., Tost, H., and Lelieveld, J.: EMAC model evaluation and analysis of atmospheric aerosol properties and distribution with a focus on the Mediterranean region, *Atmospheric Research*,  
665 114–115, 38 – 69, doi:10.1016/j.atmosres.2012.05.014, 2012.
- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmospheric Chemistry and Physics*, 6, 4321–4344, doi:10.5194/acp-6-4321-2006, 2006.
- 670 Dietmüller, S., Jöckel, P., Tost, H., Kunze, M., Gellhorn, C., Brinkop, S., Frömming, C., Ponater, M., Steil, B., Lauer, A., and Hendricks, J.: A new radiation infrastructure for the Modular Earth Submodel System (MESSy, based on version 2.51), *Geoscientific Model Development Discussions*, 2016, 1–21, doi:10.5194/gmd-2015-277, 2016.
- Finney, D. L., Doherty, R. M., Wild, O., and Abraham, N. L.: The impact of lightning on tropospheric ozone  
675 chemistry using a new global lightning parametrisation, *Atmospheric Chemistry and Physics*, 16, 7507–7522, doi:10.5194/acp-16-7507-2016, 2016a.


- Finney, D. L., Doherty, R. M., Wild, O., Young, P. J., and Butler, A.: Response of lightning NO<sub>x</sub> emissions and ozone production to climate change: Insights from the Atmospheric Chemistry and Climate Model Intercomparison Project, *Geophysical Research Letters*, 43, 5492–5500, doi:10.1002/2016GL068825, 2016b.
- 680 Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, NH<sub>4</sub><sup>+</sup>, Na<sup>+</sup>, SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, Cl<sup>-</sup>, H<sub>2</sub>O aerosols, *Atmospheric Chemistry and Physics*, 7, 4639–4659, doi:10.5194/acp-7-4639-2007, 2007.
- Gressent, A., Sauvage, B., Cariolle, D., Evans, M., Leriche, M., Mari, C., and Thouret, V.: Modeling lightning-NO<sub>x</sub> chemistry on a sub-grid scale in a global chemical transport model, *Atmospheric Chemistry and*  
685 *Physics*, 16, 5867–5889, doi:10.5194/acp-16-5867-2016, 2016.
- Grewe, V.: Impact of climate variability on tropospheric ozone, *Science of The Total Environment*, 374, 167–181, doi:http://dx.doi.org/10.1016/j.scitotenv.2007.01.032, 2007.
- Höpfner, M., Volkamer, R., Grabowski, U., Grutter, M., Orphal, J., Stiller, G., von Clarmann, T., and Wetzel, G.: First detection of ammonia (NH<sub>3</sub>) in the Asian summer monsoon upper troposphere, *Atmospheric Chemistry*  
690 *and Physics*, 16, 14 357–14 369, doi:10.5194/acp-16-14357-2016, 2016.
- Huntrieser, H., Schlager, H., Roiger, A., Lichtenstern, M., Schumann, U., Kurz, C., Brunner, D., Schwierz, C., Richter, A., and Stohl, A.: Lightning-produced NO<sub>x</sub> over Brazil during TROCCINOX: airborne measurements in tropical and subtropical thunderstorms and the importance of mesoscale convective systems, *Atmospheric Chemistry and Physics*, 7, 2987–3013, doi:10.5194/acp-7-2987-2007, 2007.
- 695 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/CBO9781107415324, www.climatechange2013.org, 2013.
- Jacobson, M. Z.: Global direct radiative forcing due to multicomponent anthropogenic and natural aerosols,  
700 *Journal of Geophysical Research: Atmospheres*, 106, 1551–1568, 2001.
- Jaegle, L., Steinberger, L., Martin, R. V., and Chance, K.: Global partitioning of NO<sub>x</sub> sources using satellite observations: Relative roles of fossil fuel combustion, biomass burning and soil emissions, *Faraday Discuss.*, 130, 407–423, 2005.
- Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) - a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433–444, 2005.
- 705 Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geoscientific Model Development*, 3, 717–752, doi:10.5194/gmd-3-717-2010, 2010.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geoscientific Model Development*, 9, 1153–1200, doi:10.5194/gmd-9-1153-2016, 2016.
- 715 Kärcher, B., Hendricks, J., and Lohmann, U.: Physically based parameterization of cirrus cloud formation for use in global atmospheric models, *J. Geophys. Res.*, 111, D01 206, doi:10.1029/2005JD006219, 2006.

- Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal processes DRY DEPosition and SEDimentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617–4632, 2006a.
- 720 Kerkweg, A., Sander, R., Tost, H., and Jöckel, P.: Technical Note: Implementation of prescribed (OFFLEM), calculated (ONLEM), and pseudo-emissions (TNUDGE) of chemical species in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 3603–3609, 2006b.
- Labrador, L. J., v. Kuhlmann, R., and Lawrence, M. G.: Strong sensitivity of the global mean OH concentration and the tropospheric oxidizing efficiency to the source of NO<sub>x</sub> from lightning, *Geophys. Res. Lett.*, 31, L06 102, doi:10.1029/2003GL019229, 2004.
- 725 Labrador, L. J., v. Kuhlmann, R., and Lawrence, M. G.: The effects of lightning-produced NO<sub>x</sub> and its vertical distribution on atmospheric chemistry: sensitivity simulations with MATCH-MPIC, *Atmos. Chem. Phys.*, 5, 1815–1834, 2005.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- 730 Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): overview and description of models, simulations and climate diagnostics, *Geoscientific Model Development*, 6, 179–206, doi:10.5194/gmd-6-179-2013, 2013.
- 735 Lohmann, U. and Hoose, C.: Sensitivity studies of different aerosol indirect effects in mixed-phase clouds, *Atmospheric Chemistry and Physics*, 9, 8917–8934, 2009.
- Lohmann, U., Rotstayn, L., Storelvmo, T., Jones, A., Menon, S., Quaas, J., Ekman, A. M. L., Koch, D., and Ruedy, R.: Total aerosol effect: radiative forcing or radiative flux perturbation?, *Atmospheric Chemistry and Physics*, 10, 3235–3246, 2010.
- 740 Makkonen, R., Romakkaniemi, S., Kokkola, H., Stier, P., Räisänen, P., Rast, S., Feichter, J., Kulmala, M., and Laaksonen, A.: Brightening of the global cloud field by nitric acid and the associated radiative forcing, *Atmospheric Chemistry and Physics*, 12, 7625–7633, 2012.
- Mao, K., Ma, Y., Xia, L., Chen, W. Y., Shen, X., He, T., and Xu, T.: Global aerosol change in the last decade: An analysis based on {MODIS} data, *Atmospheric Environment*, 94, 680 – 686, 2014.
- 750 Martin, R. V., Jacob, D. J., Logan, J. A., Bey, I., Yantosca, R. M., Staudt, A. C., Li, Q., Fiore, A. M., Duncan, B. N., Liu, H., Ginoux, P., and Thouret, V.: Interpretation of TOMS observations of tropical tropospheric ozone with a global model and in situ observations, *Journal of Geophysical Research: Atmospheres*, 107, ACH 4–1–ACH 4–27, doi:10.1029/2001JD001480, 4351, 2002.
- 755 Mayer, B. and Kylling, A.: Technical note: The libRadtran software package for radiative transfer calculations - description and examples of use, *Atmospheric Chemistry and Physics*, 5, 1855–1877, 2005.


- Ott, L. E., Pickering, K. E., Stenchikov, G. L., Allen, D. J., DeCaria, A. J., Ridley, B., Lin, R.-F., Lang, S., and Tao, W.-K.: Production of lightning NO<sub>x</sub> and its vertical distribution calculated from three-dimensional cloud-scale chemical transport model simulations, *Journal of Geophysical Research: Atmospheres*, 115, doi:10.1029/2009JD011880, 2010.
- 760 Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmospheric Chemistry and Physics*, 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007.
- Pickering, K. E., Wang, Y., Tao, W.-K., Price, C., and Müller, J.-F.: Vertical distribution of lightning NO<sub>x</sub> for use in regional and chemical transport models, *J. Geophys. Res.*, 103, 31 203–31 216, 1998.
- 765 Pozzer, A., de Meij, A., Pringle, K. J., Tost, H., Doering, U. M., van Aardenne, J., and Lelieveld, J.: Distributions and regional budgets of aerosols and their precursors simulated with the EMAC chemistry-climate model, *Atmospheric Chemistry and Physics*, 12, 961–987, doi:10.5194/acp-12-961-2012, 2012.
- Price, C. and Rind, D.: A simple Lightning Parametrization for Calculating Global Lightning Distributions, *J. Geophys. Res.*, 97, 9919–9933, 1992.
- 770 Pringle, K. J., Tost, H., Metzger, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P., Vignati, E., and Lelieveld, J.: Description and evaluation of GMXe: a new aerosol submodel for global simulations (v1), *Geoscientific Model Development*, 3, 391–412, doi:10.5194/gmd-3-391-2010, 2010.
- Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblue, L., Manzini, E., Rhodin, A., Schleese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circulation model ECHAM5: Part 1, Tech. Rep. 349, Max-Planck-Institut für Meteorologie, 2003.
- 775 Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L., Manzini, E., Schleese, U., and Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Clim.*, 19, 3771–3791, 2006.
- 780 Sander, R., Baumgaertner, A., Gromov, S., Harder, H., Jöckel, P., Kerkweg, A., Kubistin, D., Regelin, E., Riede, H., Sandu, A., Taraborrelli, D., Tost, H., and Xie, Z.-Q.: The atmospheric chemistry box model CAABA/MECCA-3.0, *Geoscientific Model Development*, 4, 373–380, doi:10.5194/gmd-4-373-2011, 2011.
- Schumann, U. and Huntrieser, H.: The global lightning-induced nitrogen oxides source, *Atmos. Chem. Phys.*, 7, 3823–3907, doi:10.5194/acp-7-3823-2007, 2007.
- 785 Steinkamp, J. and Lawrence, M. G.: Improvement and evaluation of simulated global biogenic soil NO emissions in an AC-GCM, *Atmospheric Chemistry and Physics*, 11, 6063–6082, 2011.
- Stelson, A., Friedlander, S., and Seinfeld, J.: A note on the equilibrium relationship between ammonia and nitric acid and particulate ammonium nitrate, *Atmospheric Environment (1967)*, 13, 369 – 371, 1979.
- Tanre, D., Geleyn, J.-F., and Slingo, J. M.: *Aerosols and Their Climatic Effects*, chap. First results of the introduction of an advanced aerosol-radiation interaction in the ecmwf low resolution global model, A. Deepak, Hampton, Va., 1984.
- 790 Tiedtke, M.: A Comprehensive Mass Flux Scheme for Cumulus Parametrization in Large-Scale Models, *Mon. Weather Rev.*, 117, 1779–1800, 1989.
- Tost, H. and Pringle, K. J.: Improvements of organic aerosol representations and their effects in large-scale atmospheric models, *Atmospheric Chemistry and Physics*, 12, 8687–8709, doi:10.5194/acp-12-8687-2012, 2012.
- 795

- Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical Note: A new comprehensive SCAV-enging submodel for global atmospheric chemistry modelling, *Atmos. Chem. Phys.*, 6, 565–574, 2006.
- 800 Tost, H., Jöckel, P., and Lelieveld, J.: Lightning and convection parameterisations - uncertainties in global modelling, *Atmos. Chem. Phys.*, 7, 4553–4568, 2007.
- Tost, H., Lawrence, M. G., Brühl, C., Jöckel, P., GABRIEL-Team, and SCOUT-O3-DARWIN/ACTIVE-Team: Uncertainties in atmospheric chemistry modelling due to convection parameterisations and subsequent scavenging, *Atmospheric Chemistry and Physics*, 10, 1931–1951, doi:10.5194/acp-10-1931-2010, 2010.
- 805 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, doi:10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2, 1977.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric Chemistry and Physics*, 10, 11 707–11 735, doi:10.5194/acp-10-11707-2010, 2010.
- 810 Xu, L. and Penner, J. E.: Global simulations of nitrate and ammonium aerosols and their radiative effects, *Atmospheric Chemistry and Physics*, 12, 9479–9504, 2012.
- Zare, A., Christensen, J. H., Gross, A., Irannejad, P., Glasius, M., and Brandt, J.: Quantifying the contributions of natural emissions to ozone and total fine PM concentrations in the Northern Hemisphere, *Atmospheric Chemistry and Physics*, 14, 2735–2756, doi:10.5194/acp-14-2735-2014, 2014.







**Figure 1.** Sketch of the simulated processes from the emission of  $\text{NO}_x$  molecules by lightning over gas phase conversion to  $\text{HNO}_3$ , gas aerosol partitioning to particulate  $\text{NO}_3^-$  and implications for the direct and indirect aerosol effects.




**Figure 2.** 3D visualisation of LNO<sub>x</sub> emissions (coloured isosurface of  $1 \cdot 10^{-16}$  kg/(m<sup>3</sup>s) and darker shaded isosurface of  $3 \cdot 10^{-16}$  kg/(m<sup>3</sup>s)) and the total aerosol nitrate mixing ratios (gray isosurface of 0.1 ppb<sub>v</sub>). Additionally, the mean flash rate in 1/(km<sup>2</sup> min) is depicted by the 2D slice at the bottom. Note the logarithmic scaling of both colour bars.



**Figure 3.** 3D visualisation of the relative differences in tropospheric aerosol nitrate mixing ratios between the simulations with and without LNO<sub>x</sub> emissions to the simulation including LNO<sub>x</sub> emissions. The gray isosurface depicts a relative difference of 30%, the blue isosurface of 45%, and the red isosurface of 60%. Additionally, the upper tropospheric aerosol nitrate column burden (in mg/m<sup>2</sup>) between 500 hPa and the tropopause is depicted by the coloured panel at the bottom of the graph. The turquoise contour lines depict relative differences of 20%, 40% and 60% difference in this column burden between the two simulations.




**Figure 4.** Map of vertically resolved percentage differences (including LNO<sub>x</sub> emissions as the reference case) in the aerosol size distributions as spatial and regional average (for the respective regions). In each small frame the y-axis depicts pressure altitude and the x-axis the aerosol diameter from 1nm to 10 μm. Overlaid are the contours of the absolute values of the size distributions, i.e. the absolute particle numbers calculated from the overlaying of the individual modes, determined from the spatial and temporal mean in particles/cm<sup>3</sup>. The solid lines depict 1, 10, 100, 1000 particles, whereas the dashed lines represent 10<sup>-1</sup>, 10<sup>-2</sup>, etc. particles. The figure depicts the present day conditions. The respective regions are: Central Pacific (150W:110W, 20S:0), Amazonia (80W:35W, 20S:10N), Eastern US (100W:75W, 30N:45N), South Atlantic (50W:10W, 80S:60S), Central Africa (5E:45E, 10S:10N), Europe (10W:25E, 40N:60N), Indonesia (95E:155E, 12S:10N), East Asia (105E:125E, 20N:42N), Siberia (60E:100E, 50N:70N). The hatched regions mark areas without statistical significance.




**Figure 5.** 3D visualisation of aerosol extinction and the influence of LNO<sub>x</sub> emissions. The left panel shows a map of the vertically integrated column AOD (at 550 nm) when lightning emissions are included (floor plane). The ceiling depicts the relative differences of the integrated column AOD between the simulation with lightning emissions minus the simulation without lightning LNO<sub>x</sub>, with the full setup serving as reference. The back panel displays the zonal average aerosol extinction (in 1/km at 550 nm) of the full simulations (Please, note the logarithmic scale.). The front panel depicts relative percentage differences due NO<sub>x</sub> emissions from lightning. White areas mark regions without statistical significance.

The right panel also shows AOD (floor) and zonal mean extinction (rear panel), and additionally 3D isosurfaces in the center of the box represent the +10% (pale red) and +20% (dark red) of the enhanced extinction due to active LNO<sub>x</sub> emissions, whereas the -10% (pale blue) and -20% (dark blue) isosurfaces mark regions, in which the emissions result in a reduction of the extinction.



**Figure 6.** Visualisation of changes in the effective radius of ice crystals. The back panel displays the statistically significant absolute changes in zonal average ice crystal size in  $\mu\text{m}$ . The white contours depict a mean ice crystal size and therefore mark regions with large and smaller crystals. Additionally, the isosurfaces represent the regions for substantial absolute changes for the effective ice crystal size (turquoise for negative and purple for positive) due the  $\text{LNO}_x$  emissions.



**Figure 7.** Absolute changes in the shortwave all-sky fluxes at the top of the atmosphere. The upper panel depicts the long term time average of the difference in the simulation with LNO<sub>x</sub> emissions minus the fluxes without lightning emissions. The hatches mark regions with a statistical significant signal (compared to the internal interannual variability). The bottom panel depicts the time series of the monthly mean global mean differences with the red line marking a 0 W/m<sup>2</sup> change.

**Table 1.** Tropospheric and upper tropospheric burden in the ARG simulation of the important highly oxidised nitrogen species, i.e. gaseous  $\text{HNO}_3$ , gaseous  $\text{N}_2\text{O}_5$ , aerosol  $\text{NO}_3^-$  and the sum of those three compounds. All values are given in Gg N (except for the relative differences which are provided in %) and are globally and vertically integrated over the whole and the upper troposphere (500hPa up to the tropopause).

	$\text{HNO}_3$	$\text{N}_2\text{O}_5$	$\text{NO}_3^-$	Total N(V+)
Present day				
Absolute values (with $\text{LNO}_x$ emissions):				
Tropospheric Column burden	403	8.0	143	554
UT Column burden	174	5.0	27.8	207
Absolute differences due to $\text{LNO}_x$ emissions:				
Tropospheric Column burden	162	3.9	20.0	186
UT Column burden	107	3.5	9.0	119
Relative differences in (%) due to $\text{LNO}_x$ emissions:				
Tropospheric Column burden	40.1	48.4	14.0	33.5
UT Column burden	61.3	70.3	32.3	57.7
Preindustrial conditions				
Absolute values (with $\text{LNO}_x$ emissions):				
Tropospheric Column burden	238	4.5	67.8	310
UT Column burden	143	4.1	15.0	161
Absolute differences due to $\text{LNO}_x$ emissions:				
Tropospheric Column burden	158	3.7	25.8	188
UT Column burden	110	3.5	8.1	121
Relative differences in (%) due to $\text{LNO}_x$ emissions:				
Tropospheric Column burden	66.6	82.6	38.1	60.6
UT Column burden	76.9	86.0	54.3	75.1



**Table 2.** CH<sub>4</sub> tropospheric and upper tropospheric (500 hPa to tropopause) lifetime and absolute change due to LNO<sub>x</sub> emissions.

	CH <sub>4</sub> lifetime [years]	Increase of the CH <sub>4</sub> lifetime [years]
<b>Present day</b>		
troposphere	7.4	1.7
UT only	14.0	9.7
<b>Preindustrial conditions</b>		
troposphere	8.9	1.9
UT only	14.1	13.2

The CH<sub>4</sub> lifetime is calculated with the help of the actual methane and OH concentrations, calculating a pseudo-first order loss rate. The corresponding temperature dependent reaction rate is identical to the one used in the online chemistry calculations, using also the temperature output of the simulations. Calculating the lifetime for each individual cell, average values are determined using grid mass weighted factors for certain regions or the whole troposphere.