

Pre-monsoon air quality over Lumbini, a world heritage site along the Himalayan foothills

Dipesh Rupakheti^{1,2*}, Bhupesh Adhikary³, Puppala S. Praveen³, Maheswar Rupakheti^{4,5},
Shichang Kang^{6,7*}, Khadak S. Mahata⁴, Manish Naja⁸, Qianggong Zhang^{1,7}, Arnico K. Panday³,
Mark G. Lawrence⁴

¹Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of
Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

²University of Chinese Academy of Sciences, Beijing 100049, China

³International Centre for Integrated Mountain Development (ICIMOD), Kathmandu, Nepal

⁴Institute for Advanced Sustainability Studies (IASS), Potsdam 14467, Germany

⁵Himalayan Sustainability Institute (HIMSI), Kathmandu, Nepal

⁶State Key Laboratory of Cryospheric Science, Cold and Arid Regions Environmental and
Engineering Research Institute (CAREERI), Lanzhou 730000, China

⁷Center for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing
100085, China

⁸Aryabhata Research Institute of Observational Sciences (ARIES), Nainital, India

**Correspondence to:*

D. Rupakheti (dipesh.rupakheti@itpcas.ac.cn), S.C. Kang (shichang.kang@lzb.ac.cn)

Abstract

Lumbini, in southern Nepal, is a UNESCO world heritage site of universal value as the birthplace of Buddha. Poor air quality in Lumbini and surrounding regions is a great concern for public health as well as for preservation, protection and promotion of Buddhist heritage and culture. We present here results from measurements of ambient concentrations of key air pollutants (PM, BC, CO, O₃) in Lumbini, first of its kind for Lumbini, conducted during an intensive measurement period of three months (April-June 2013) in the pre-monsoon season. The measurements were carried out as a part of the international air pollution measurement campaign; SusKat-ABC (Sustainable Atmosphere for the Kathmandu Valley - Atmospheric Brown Clouds). The main objective of this work was to understand and document the level of air pollution, diurnal characteristics and the influence of open biomass burning on air quality in Lumbini. The hourly average concentrations during the entire measurement campaign ranged as follows: BC: 0.3 - 30.0 $\mu\text{g m}^{-3}$, PM₁: 3.6-197.6 $\mu\text{g m}^{-3}$, PM_{2.5}: 6.1 - 272.2 $\mu\text{g m}^{-3}$, PM₁₀: 10.5 - 604.0 $\mu\text{g m}^{-3}$, O₃: 1.0 - 118.1 ppbv, and CO: 125.0 - 1430.0 ppbv. These levels are comparable to other very heavily polluted sites in South Asia. Higher fraction of coarse mode PM was found as compared to other nearby sites in the IGP-region. $\Delta\text{BC}/\Delta\text{CO}$ ratio obtained in Lumbini indicated considerable contributions of emissions from both domestic and transportation sectors. The 24-h average PM_{2.5} and PM₁₀ concentrations exceeded the WHO guideline very frequently (94% and 85% of the sampled period, respectively), which implies significant health risks for the residents and visitors in the region. These air pollutants exhibited clear diurnal cycles with high values in the morning and evening. During the study period, the worst air pollution episodes were mainly due to agro-residue burning and regional forest fires combined with meteorological conditions conducive of pollution transport to Lumbini. Fossil fuel combustion also contributed significantly, accounting for more than half of the ambient BC concentration according to aerosol spectral light absorption coefficients obtained in Lumbini. WRF-STEM, a regional chemical transport model, was used to simulate the meteorology and the concentrations of pollutants to understand the pollutant transport pathways. The model estimated values were ~ 1.5 to 5 times lower than the observed concentrations for CO and PM₁₀ respectively. Model simulated regionally tagged CO tracers showed that the majority of CO came from the upwind region of Ganges Valley. Model needs significant improvement in simulating aerosols in the

region. Given the high pollution level, there is a clear and urgent need for setting up a network of long-term air quality monitoring stations in the greater Lumbini region.

1. Introduction

The Indo-Gangetic plain (IGP) stretches over 2000 km encompassing a vast area of land in northern South Asia: the eastern parts of Pakistan, most of northern and eastern India, southern part of Nepal, and almost all of Bangladesh. The Himalayan mountains and their foothills stretch along the northern edge of IGP. The IGP region is among the most fertile and most intensely farmed region of the world. It is a heavily populated region with about 900 million residents or 12% of the world's population. Four megacities - Lahore, Delhi, Kolkata, and Dhaka are located in the IGP region, with dozens more cities with populations exceeding one million. The region has witnessed impressive economic growth in recent decades but unfortunately it has also become one of the most polluted, and an air pollution 'hot spot' of local, regional and global concern (Ramanathan et al., 2007). Main factors contributing to air pollution in the IGP and surrounding regions include emissions from vehicles, thermal power plants, industries, biomass and fossil fuel used in cooking and heating activities, agricultural activities, crop residue burning and forest fires. Air pollution gets transported long distances away from emission sources and across national borders. As a result, the IGP and adjacent regions get shrouded with a dramatic annual buildup of regional scale plumes of air pollutants, known as Atmospheric Brown Clouds (ABC), during the long and dry winter and pre-monsoon seasons each year (Ramanathan and Carmichael, 2008). **Figure 1 shows monthly synoptic wind and mean aerosol optical depth (AOD) during April-June, 2013 over South Asia.** Very high aerosol optical depth along the entire stretch of IGP reflects severity of air pollution over large areas in the region.

Poor air quality continues to pose significant threat to human health in the region. In a new study of global burden of disease released recently, Forouzanfar et al. (2015) estimated that in 2013 around 1.7 million people died prematurely in Pakistan, India, Nepal, and Bangladesh as a result of air pollution exposure, nearly 30% of global total premature deaths due to air pollution. Air pollution also affects precipitation (e.g. South Asian monsoon), agricultural productivity, ecosystems, tourism, climate, and broadly socio-economic and national development goals of the countries in the region (Burney and Ramanathan, 2014; Shindell, 2011; Ramanathan and

Carmichael, 2008). It has also been linked to intensification of cold wave and winter fog in the IGP region over recent decades (Lawrence and Lelieveld, 2010 and references therein; Safai et al., 2009; Ganguly et al., 2006). Besides high levels of aerosol loading as shown in Fig. 1, Indo-Gangetic plains also have very high levels of ground level ozone or tropospheric ozone (O_3) (e.g., Ramanathan and Carmichael (2008)) which is a toxic pollutant to plant and human health, and a major greenhouse gas (IPCC, 2013; Shindell, 2011; Mohnen et al., 1993). South Asia, in particular IGP region, has been projected to be the most ozone polluted region in world by 2030 (Stevenson et al., 2006). Majority of crop loss in different parts of the world results from effects of ozone on crop health and productivity (Shindell, 2011). Burney and Ramanathan (2014) also reported a significant loss in wheat and rice yields in India from 1980 to 2010 due to direct effects of black carbon (BC) and ozone (O_3). BC and O_3 are two key short-lived climate pollutants (SLCP). Similarly, species like fine particles and carbon monoxide (CO) are potent to health damages by posing impacts upon the respiratory and cardiovascular system and even also to the climate system (Singh et al., 2017 and references therein). Because of the IGP's close proximity to the Himalaya-Tibetan plateau region, this once relatively clean region, is now subjected to increasing air pollution transported from regions such as the IGP, which can exert additional risks to sensitive ecosystems in the mountain region (e.g., (Lüthi et al., 2015; Marinoni et al., 2013; Duchi et al., 2011). However, air pollution transport pathways to Himalayas are still not yet fully understood.

Monuments and buildings made with stones are vulnerable to air pollution damage (Brimblecombe, 2003; Gauri and Holdren, 1981). Sulfur dioxide, which forms sulfuric acid upon reaction with water, is the most harmful substance for the monuments as it can corrode and damage them (Baedecker et al., 1992; Gauri and Holdren, 1981). A recent study has reported that deposition of light absorbing aerosol particles (black carbon, brown carbon) and dust is responsible for the discoloration of Taj Mahal, a world famous monument in India (Bergin et al., 2015). Lumbini, located near the northern edge of the central Indo-Gangetic plain, is famous as the birthplace of the Lord Buddha and thus a UNESCO world heritage site of outstanding universal value to humanity. Since the study area is renowned due to its historical and archaeological significance, Lumbini is getting the worldwide attention also for poor air quality

in the region. There was no regular air quality monitoring in Lumbini at the time of our measurement campaign.

Through this study, we want to understand the level of air pollution, its diurnal characteristic, and the influence of open biomass burning on air quality in Lumbini. We carried out continuous measurements of ambient concentrations of key air pollutants (particulate matter, black carbon, carbon monoxide, ozone) and meteorological parameters during an intensive measurement period of three months (April-June) in the year 2013. These are the first reported pollutant measurements for Lumbini. A regional chemical transport model called Sulfur Transport and dEposition Model (STEM) was used to simulate the variations of meteorological parameters and air pollutants during the observation period to examine the extent to which a state-of-the-art, widely-used air quality model is able to simulate the data, as an indication for where there are still gaps in our knowledge and what further measurements and emissions dataset developments are needed. Model simulated regionally tagged CO tracers were used to identify emission source regions impacting pollutant concentration observed at Lumbini. Satellite data has also been used to understand the high pollution events during the monitoring period. These measurements were carried out as a part of the SusKat-ABC international air pollution measurement campaign (*M. Rupakheti, manuscript in preparation for ACPD*) jointly led by the International Centre for Integrated Mountain Development (ICIMOD), Kathmandu, Nepal and Institute for Advanced Sustainability Studies (IASS), Potsdam, Germany.

2. Experimental set up

2.1 Sampling site

The Lumbini measurement site (27°29.387' N, 83°16.745' E, elevation: ~100 m above sea level) is located at the premise of the Lumbini International Research Institute (LIRI), a Buddhist library in Lumbini. Lumbini lies in the Nepal's southern lowland plain or *Terai* region, termed as "bread basket of Nepal" due to the availability of very fertile land suitable for crop production, which forms the northern edge of the Indo-Gangetic Plains (IGP). About 25 km north of Lumbini the foothills begin while the main peaks of the Himalayas are 140 km to the north. The remaining three sides are surrounded by flat plain land of Nepal and India. The site is only about 8 km from the Nepal-India border in the south. A three storied 10 m tall water tower was used as

the platform for the automatic weather station (AWS) whereas remaining instruments were placed inside a room near the base of the tower. **Figure S1** shows the location of Lumbini, the Kenzo Tange Master Plan area of the Lumbini development project, and the sampling tower. An uninterrupted power back up was set up in order to assure the regular power supply even during hours with scheduled power cuts during the monitoring period. The nearby premises of the monitoring site consist of the LIRI main office and staff quarters. Further away is a museum, a local bus park for the visitors to Lumbini, the office of the Lumbini Development Trust, monasteries, and thinly forested area with grassland within the master plan area. Outside of the master plan area lie vast area of agricultural fields, village pockets, and several brick kilns and cement industries. A local road (black topped), that cuts through the master plan area, lies about 200 m north of the sampling site and experiences intermittent passing of vehicles. According to the Ministry of Culture, Tourism and Civil Aviation of Nepal over 130 thousand tourists (excluding Nepalese and Indian citizens) visited the Lumbini area in 2014 (<http://tourism.gov.np/en>).

2.2 Monitoring Instruments

The summary of instruments deployed in Lumbini is presented in Table 1. All data were collected in Nepal Standard Time (NST) which is GMT +05:45 hour. PM_1 , $PM_{2.5}$ and PM_{10} mass concentrations were monitored continuously with GRIMM EDM164 (GRIMM Aerosol Technik, Germany) which uses the light scattering at 655 nm to derive mass concentrations. Similarly, aerosol light absorptions at 7 wavelengths (370, 470, 520, 590, 660, 880, 950 nm) were measured continuously with an Aethalometer (Model AE-42, Magee Scientific, USA), averaging and reporting data every 5 min. It was operated at a flow rate of 5 l min^{-1} . No cut-off was applied for inlet; hence the reported concentration of BC is total suspended BC particles. As described by the manufacturer, ambient BC concentration is derived from light absorption at 880 nm using a specific mass absorption cross section. To obtain BC concentration in Lumbini, we used a specific mass absorption cross-section value of $8\text{ m}^2\text{ g}^{-1}$ for the 880 nm channel. A similar value has been previously used for BC measurement in the Indo-Gangetic plain (Praveen et al., 2012). To remove the filter loading effect, we used correction method suggested by Schmid et al. (2006) which was also used by Praveen et al. (2012) for BC measurements at a rural site in the Indo-Gangetic plain. Surface ozone (O_3) concentration was measured continuously with an ozone

analyzer (Model 49i, Thermo Scientific, USA) which utilizes UV (254 nm wavelength) photometric technology to measure ozone concentration in ambient air. ~~CO analyzer (Model 48i, Thermo Scientific, USA) was used to monitor ambient CO concentration which is based on the principle that CO absorbs infrared radiation at the wavelength of 4.6 microns.~~ The ambient air was drawn through 6-micron pore size SAVILLEX 47 mm filter at the inlet in order to remove the particles before sending air into the CO and O₃ analyzers using a Teflon tube. The filters were replaced every 7-10 days depending on particle loading, based on manual inspection. CO instrument was set to auto-zero at a regular interval of 6 hours. Local meteorological parameters (temperature, relative humidity, wind speed, wind direction, precipitation, and global solar radiation) were monitored with an automatic weather station (AWS) (Campbell Scientific, Loughborough, UK), recording data every minute.

2.3 Regional chemical transport model

Aerosol and trace gas distributions were simulated using a regional chemical transport model. Sulfur Transport and dEposition Model (STEM), a 3D eulerian model that has been used extensively in the past to characterize air pollutants in South Asian region ~~was used to understand~~ observations at Lumbini (Kulkarni et al., 2015; Adhikary et al., 2007). The Weather Research and Forecasting (WRF) model (Skamarock et al., 2008) version 3.5.1 was used to generate the required meteorological variables necessary for simulating pollutant transport in STEM. The model domain was centered at 24.94° N latitude and 82.55° E longitude covering a region from 3.390° N to 43.308° N latitude and 34.880° E to 130.223° E longitude. The model has 425×200 horizontal grid cells with grid resolution of 25×25 km and 41 vertical layers with top of the model set at 50 mbar. The WRF model was run from November 1, 2012 to June 30, 2013. However, for this study, modeled data only from April to June 2013 have been used. The WRF model was initialized with FNL data available from NCAR/UCAR site (<http://rda.ucar.edu/datasets/ds083.2/>).

The tracer version of the STEM model provides mass concentration of sulfate, BC (hydrophilic and hydrophobic), Organic carbon (OC), sea salt (fine and coarse mode), dust (fine PM_{2.5} and PM₁₀), CO (biomass and anthropogenic) and region tagged CO tracers. STEM model domain size, resolution and projection are those of the WRF model. Details about tracer version of the

STEM model is outlined elsewhere (Kulkarni et al., 2015; Adhikary et al., 2007). Anthropogenic emission of various pollutants (CH_4 , CO , SO_2 , NO_x , NMVOC, NH_3 , PM_{10} , $\text{PM}_{2.5}$, BC and OC) used in this analysis were taken from the EDGAR-HTAP_v2 (http://edgar.jrc.ec.europa.eu/htap_v2/index.php?SECURE=123). Open biomass burning emissions on a daily basis during the simulated period were taken from data obtained from the FINN model (Wiedinmyer et al., 2011). Both these emissions were re-gridded to the STEM model domain using four point interpolation techniques available in the STEM model emissions preprocessor. As with the WRF model, the STEM model was run from November 2, 2012 to June 30, 2013 however, data presented here are only during the intensive field campaign period.

3. Results and discussions

3.1 Meteorology

Hourly average time series of various meteorological parameters like precipitation in mm hr^{-1} (Prec), temperature in $^{\circ}\text{C}$ (T), relative humidity in % (RH), wind speed in m s^{-1} (WS) and direction in degree (WD) during the monitoring period are shown in Figure 2. Meteorological parameters were obtained with the sensors at the height of ~ 12 m from the ground. Meteorology results from WRF model simulations have been used to indicate if any significantly different air mass was present during the measurement campaign after the meteorological observations malfunctioned. Precipitation data was derived from TRMM satellite (TRMM_3B42_007 at a horizontal resolution of 0.25°) from the Giovanni platform (<http://giovanni.gsfc.nasa.gov/giovanni/>) as the rain gauge malfunctioned during the sampling period. Precipitation data from TRMM (Figure 2) show that Lumbini was relatively dry in the early portion of the measurement campaign while as the pre-monsoon edged closer to the monsoon onset, the site did experience some rainfall events. This lowered aerosol loading in the later half of the measurement campaign due to washout. Comparison of WRF model outputs with TRMM data shows that the model under-predicts rainfall through out the campaign.

Average observed temperature for the sampling period until the sensor stopped working (on 8th May, 2013, i.e., for 38 days of measurement) was 28.1°C (minimum: 16.5°C , maximum: 40°C). Average temperature from the model, during same period, was 31°C with values ranging between $19 - 40^{\circ}\text{C}$. As shown in Figure 2, the model captures the synoptic variability of

temperature and is mostly within the range of daily values. However, the model has a high bias and does not capture well daily minimum temperature values. In addition, the model does not show any large variation in temperature for the campaign period after the sensors stopped working. This insight will be useful to interpret pollution data later on. For the same period (until the sensor stopped working), the average (observed) RH was ~ 50% (ranging from 10.5 to 97.5%) whereas the model showed the average RH to be ~ 23% with values ranging between 6 to 78%. RH values are highly underestimated by the model, however as previously mentioned, the model does not show significant changes in RH during the measurement campaign after the observations stopped working.

Average observed wind speed during the study period was 2.4 m s^{-1} , with hourly values ranging between $0.03 - 7.4 \text{ m s}^{-1}$ whereas from the WRF model average wind speed was found to be 3.2 m s^{-1} (range: $0.06 - 11.1 \text{ m s}^{-1}$). Diurnal variation of observed hourly average wind speed suggested that wind speeds were lower during nights and mornings while higher wind speed prevailed during day time, with average winds $> 3 \text{ m s}^{-1}$ up to $\sim 3.3 \text{ m s}^{-1}$ between 09:00-13:00 local time (Supplementary materials, Figure S2, lower panel). High speed strong winds ($> 4 \text{ m s}^{-1}$) were from the NW direction during the month of April which later switched to almost opposite direction, i.e., SE direction from the month of May onwards. The monthly wind rose plot using the data from both observation and modeling where the difference in the pattern could be potentially due to the data resolution is shown in Figure S3. Comparing modeled wind direction prediction skills at the surface with one point measurement is not sufficient. However, in the absence of other measurements, we also show the comparison of wind direction as an indication of model performance not as model validation. Discrepancy on model results might have occurred due to various factors inherently uncertain in a weather prediction using a model. Besides, air pollution transport also occurs via elevated layers and is not limited to surface winds. We show NCEP/NCAR reanalysis plots at 850 hPa in Fig. S3 to illustrate the distinctly differing wind direction compared to the surface winds seen from observations as well as NCEP/NCAR reanalysis plot at 1000 hPa shown in Fig. 1. There are no upper wind measurement data nearby Lumbini to show model performance. Regardless, we believe that air quality model data is vital for understanding pollutant transport in an area where observation data are non-existent or are incomplete.

3.2 Air Quality

3.2.1 General overview, PM ratios and influence of meteorology on pollution concentrations

Figure 3 shows hourly averaged time series of observed BC, PM₁, PM_{2.5}, PM₁₀, O₃ and CO observed at Lumbini during the study period. Similar temporal behaviour was shown by BC, particulate matter fractions (PM₁, PM_{2.5} and PM₁₀) and CO. The gap in the figure (for PM time series) is due to the power interruption to the instrument. BC concentrations during the measurement period ranged between 0.3-29.9 $\mu\text{g m}^{-3}$ with a mean (\pm SD) value of 4.9 (\pm 3.8) $\mu\text{g m}^{-3}$. BC concentrations in Lumbini during pre-monsoon months are lower compared to BC concentrations observed in the Kathmandu Valley because of high number of vehicles plying on the street, brick kilns and other industries in Kathmandu valley (Sharma et al., 2012;Putero et al., 2015). The lowest concentration was observed during a rainy day (21-22 April) whereas the highest concentration was observed during a period of forest fire (detailed in Section 3.3). For the entire measurement period, we found average (of hourly average values) PM₁: 35.8 \pm 25.6 $\mu\text{g m}^{-3}$ (minimum-maximum range: 3.6 - 197.6 $\mu\text{g m}^{-3}$), PM_{2.5}: 53.1 \pm 35.1 $\mu\text{g m}^{-3}$ (6.1 - 272.2 $\mu\text{g m}^{-3}$), PM₁₀: 128.9 \pm 91.9 $\mu\text{g m}^{-3}$ (10.5-603.9 $\mu\text{g m}^{-3}$) and coarse-mode (PM_{10-2.5}): 75.65 \pm 61.67 $\mu\text{g m}^{-3}$ (1.98-331.80 $\mu\text{g m}^{-3}$). The coarse-mode fraction was ~ 60% of the PM₁₀. The share of coarse-mode aerosol to PM₁₀ in Lumbini was higher than that observed in other sites in the IGP; Guwahiti, India (42%) (Tiwari et al., 2017) and Dibrugarh, India (9-16%) (Pathak et al., 2013) both in eastern IGP and Delhi (38%) (Tiwari et al., 2015) in western IGP indicating the higher contribution of coarse aerosols in Lumbini, likely lifted from soils from nearby agricultural fields and construction materials by stronger winds during pre-monsoon season. Similar value of coarse-mode fraction, as in Lumbini, has been reported by Misra et al. (2014) at Kanpur for dust dominated and mixed aerosols events.

The share of BC in PM fractions were found to be ~13% in PM₁, 9% in PM_{2.5} and ~4% in PM₁₀ but the correlation coefficients of BC with three PM fractions were found to be 0.89 (PM₁), 0.88 (PM_{2.5}) and 0.69 (PM₁₀), indicating the commonality in the sources of these pollutants. The contribution of BC in PM₁ was found to be of ~12% in Kanpur during February-March (Kumar et al., 2016a) similar to Lumbini. Regarding the share of BC in PM₁₀, the share observed in

Lumbini (~4%) was similar to that observed over Varanasi (~340 km due south of our site) in central IGP (5%) (Tiwari et al., 2016) and Dibrugarh in eastern IGP (~5%) (Pathak et al., 2013). Thus our results indicate that despite our station being located at the northern edge of the IGP along the foothills of the Himalayan range, its aerosol characteristics are similar to those found in heavily polluted sites in the central and eastern IGP.

In Lumbini, the average (hourly) share of PM_1 in $PM_{2.5}$, PM_1 in PM_{10} and $PM_{2.5}$ in PM_{10} were found to be ~70%, 34% and 47% respectively. The share of average (sampling period) coarse-mode aerosols to PM_{10} (60%) was found to be higher as compared to that of average fine mode i.e., $PM_{2.5}$ (40%). Regarding other sites in IGP region, $PM_{2.5}/PM_{10}$ ratios were reported to be 56% in Kanpur (Snider et al., 2016), 60% in Varanasi (Kumar et al., 2015), 57% in Guwahati (Tiwari et al., 2017), 90% in Dribugarh (Pathak et al., 2013) and 62% in Delhi (Tiwari et al., 2015) indicating local differences within IGP as well as suggesting that influence of combustion sources at Lumbini is still lower compared to other locations in Indian section of the IGP. A recent study (Putero et al., 2015) reported the PM_1/PM_{10} during pre-monsoon of 2013 was found to be 0.39 in the Kathmandu Valley of Nepal. Lumbini has significantly lower vehicle emissions and human population than the Kathmandu Valley yet the ratios are similar, indicating the importance of regional combustion sources in Lumbini for finer aerosols (PM_1), and soil-based emissions such as road dust in the Kathmandu Valley. Future studies will need to explore the emission sources around Lumbini in much greater detail. Lower $PM_{2.5}/PM_{10}$ in Lumbini as compared to other regions mentioned earlier could be due to emissions from cement industries located within 15 km distance from the measurement site. Cement factories emit coarse sized particles but we are not able to distinguish in our measurement without having an analysis of certain marker species. Trivedi et al. (2014) reported a ratio of 0.39 (during pre-monsoon) over Delhi, which is lower than the ratio in Lumbini. The lower ratio in Delhi was due to the presence of coarse sized windblown desert dust and suspended soil materials due to strong winds.

The observed 24-hour average particulate matter concentrations ($PM_{2.5}$ and PM_{10}) were found frequently higher than the WHO prescribed guidelines for $PM_{2.5}$ ($25 \mu g m^{-3}$) and PM_{10} ($50 \mu g m^{-3}$) with $PM_{2.5}$: exceeding 94% and PM_{10} : 85% of the measurement period of 53 days in Lumbini.

Observed CO concentrations ranged between 124.9-1429.7 ppbv with an average value of 344.1±160.3 ppbv. CO concentration observed in Lumbini is lower than that of Mohali, Western India where the average concentration was 566.7 ppbv during pre-monsoon season due to intense biomass and agro-residue burning over the region (Sinha et al., 2014). Temporal variation of CO concentrations is similar to that of BC as both of these species are emitted during incomplete combustion of fuel. Moreover, a very strong correlation ($r = 0.9$) was observed between BC and CO. Past studies have shown that the ratio of BC to CO depends upon multiple factors like site location, combustion characteristics (fuel and technology) at the sources, and type of air mass (Girach et al., 2014; Pan et al., 2011; Zhou et al., 2009). Formation of the soot depends on the carbon to oxygen ratio of fuel whereas CO can also be produced naturally due to the oxidation of VOCs (Girach et al., 2014). Figure 4 shows the comparison of the average $\Delta BC/\Delta CO$ ratio (0.021) at Lumbini with that obtained from other sites. Please refer to Figure S4 in the supplementary materials for the time series of $\Delta BC/\Delta CO$ ratio observed in Lumbini. We used the method described by Pan et al. (2011) to calculate the $\Delta BC/\Delta CO$ values. The ratio was calculated using the equation $(BC-BC_0)/(CO-CO_0)$ assuming the background values (BC_0 or CO_0) as 1.25 percentile of the data. The $\Delta BC/\Delta CO$ ratio in Lumbini is similar to that obtained at a suburban site, Pantnagar in India (0.017) (Joshi et al., 2016) and in Maldives (0.017) (Dickerson et al., 2002). As compared to Lumbini, the different $\Delta BC/\Delta CO$ ratio obtained over megacities such as Beijing and Shanghai are due to the higher number of gasoline and diesel vehicles (Zhou et al., 2009). However, the ratio obtained at Lumbini were within the range of emission ratios from diesel used in transport sector (0.0013-0.055), coal (0.0019-0.0572) and biofuels (0.0087-0.0266) for domestic activities (Verma et al., 2010 and references therein). The hourly averaged observed ozone concentration ranged between 1.0 and 118.1 ppbv with a mean value of 46.6±20.3 ppbv during the sampling period. The 8-hr maximum O_3 concentration exceeded WHO guidelines of $100 \mu g m^{-3}$ (WHO, 2006) during 88% of the measurement period. Our results clearly indicate that the current pollution levels in Lumbini is of great concern to health of the people living in the region as well as over a million visitors who visit Lumbini, as well as ecosystems, particularly agro-ecosystem, especially in warm and sunny pre-monsoon months.

The relationship of wind speed (WS) with aerosol and gaseous pollutants in Lumbini is shown in Figure S5 (Supplementary information). We were interested in studying the relationship between wind speed and the pollutants since the wind governs the horizontal dilution of the pollutants (Huang et al., 2012) and also likelihood of lifting soil dust. Except ozone, all other pollutants exhibited negative correlation with wind speed. BC shows negative correlation ($r = -0.42$) with the wind speed which is similar with other pollutants as well (as can be seen from the figure). Past studies have also reported a similar negative correlation of BC with wind speed over urban and sub-urban areas (Huang et al., 2012; Cao et al., 2009; Ramachandran and Rajesh, 2007; Sharma et al., 2002; Tiwari et al., 2013) indicating that the locally generated BC can accumulate in the atmosphere during lower wind speed conditions (Cao et al., 2009). Tiwari et al. (2013) also reported similar negative correlation ($r = -0.45$) during the pre-monsoon season over Delhi. On the other hand, secondary pollutants like ozone exhibited a positive relation with the WS ($r=0.38$) indicating the WS could be one of the potential factors of high ozone in Lumbini. Solar radiation is one of the most important factors for production of ozone in the atmosphere (Naja et al., 2003). The correlation of hourly ozone concentration with solar radiation (not shown here) was found to be 0.41 whereas wind speed during the daytime only (06:00-18:00) showed very weak correlation of 0.02 with ozone, indicating the calm condition as conducive to formation and accumulation of ozone in the region. Interestingly, the highest concentrations of all measured pollutants were obtained when the wind speed was less than 1 m s^{-1} . In a separate analysis (not shown here), we considered only the WS $>1 \text{ m s}^{-1}$ and calculated the correlation coefficients to investigate the influence of regional emissions. We found the similar correlation values as previous when all WS values were considered (BC vs WS = -0.41, CO vs WS = -0.42, O₃ vs WS = 0.29, PM₁ vs WS = -0.40, PM_{2.5} vs WS = -0.38, PM₁₀ vs WS = -0.33). The correlation of WS ($>1 \text{ m/s}$) with concentration of air pollutants elucidates that air pollution over Lumbini is not only of the local origin, it is rather transported from other nearby regions as well.

Past studies near this site have been focused on the cities like Kathmandu (Sharma et al., 2012; Ram et al., 2010; Panday and Prinn, 2009; Putero et al., 2015) and Kanpur (Ram et al., 2010) and agro-residue burning dominated regions of IGP (Rastogi et al., 2016; Sinha et al., 2014; Sarkar et al., 2013) or a remote mountain location in India (Naja et al., 2014). Very high

aerosol loading is observed in South Asia during pre-monsoon, mostly over the IGP region (Supplementary materials, Figure S6). As this is the first study over an IGP site located in Nepal, pollution concentrations observed at Lumbini were compared with other sites in the region (Table 2). Different sites located at urban, semi-urban and remote locations were used for comparison to get a clear comparative picture of the situation at Lumbini amongst other locations in the region. Pre-monsoon seasonal average $PM_{2.5}$ concentration in Lumbini has been found to be lower than the megacity like Delhi (Bisht et al., 2015) and north-western IGP (Sinha et al., 2014), possibly due to higher level of emissions (from traffic and biomass burning, respectively) over those regions. In addition, average BC and CO concentrations in Lumbini were found falling in between concentrations observed at rural sites (up to 6 times higher) and cities in the region (see Table 2), indicating that Lumbini, in a way, can still be considered as semi-urban location. The hourly average O_3 concentration in Lumbini were found to be higher than the cities like Kathmandu (Putero et al., 2015) and Kanpur during pre-monsoon season (Gaur et al., 2014). However from a mesoscale perspective, the hourly average O_3 concentrations were lower at Lumbini as compared to base camp of Mt. Everest region due to the uplift of polluted air masses (Marinoni et al., 2013), stratospheric intrusion (Cristofanelli et al., 2010) and even the regional or long-range transport of the air pollutants (Bonasoni et al., 2010) to the high altitude site.

Regarding the monthly average concentration, the concentrations of all measured pollutants decreased as the pre-monsoon months advanced. The monthly average concentrations of the monitored species are shown in the Figure S7 along with the monthly fire hotspots over the region. Reduction in concentration (except PM) during the month of May (as compared to April) could be attributed to the fewer fire events during May as well as previously discussed washout by rainfall. Two peak pollution episodes were observed during the first half of April and May which is discussed in more detail in the next section.

3.2.2 Observation-model inter-comparison

Chemical transport models provide insight to observed phenomena; however, interpretation has to take into account model performance before arriving at any conclusion. This section describes pollution concentrations simulated by the WRF-STEM model. A comparison of model calculated pollutant concentration along with the minimum and maximum concentrations of various

pollutants (with observation) is shown in Table 3. The model based concentrations used here are instantaneous values for every third hour of the day. BC concentrations ranged between 0.4-3.7 $\mu\text{g m}^{-3}$ with a mean value of $1.8 \pm 0.7 \mu\text{g m}^{-3}$ for a period of 1st April-15th June 2013. The average model BC concentration was ~2.7 times lower than the observed BC. Regarding PM₁, PM_{2.5} and PM₁₀, the model simulated average concentration was 12.3 ± 5.5 (0.9-41.7) $\mu\text{g m}^{-3}$, 17.3 ± 6.7 (1.9-48.3) $\mu\text{g m}^{-3}$ and 25.4 ± 12.9 (2.1-68.8) $\mu\text{g m}^{-3}$, respectively. The model estimated values were lower by the factor of 3 and 5 respectively than the observed concentrations. The data show that model needs much improvement in its ability to adequately predict observed aerosol characteristics at Lumbini. Since pollutant concentration is a function of emissions, transport and transformation and deposition, improvements in any of these areas would improve the model. However, given observation insights by PM ratios, it seems that improvements are much needed in the emissions of primary aerosols. Current emissions do not account for trash burning, roadside dust and increasingly newer industries, especially emissions from cement factories that have propped up in recent years.

Average observed CO concentration was 255.7 ± 83.5 ppbv, ranging between 72.2-613.1 ppbv, with average model CO ~1.35 times lower than observed. Time series comparison of modeled CO versus observation is shown in Figure 3. Apart from two peak episodes the model does a better job in predicting CO concentration over Lumbini. Previous study using the STEM model over Kathmandu valley showed that the model was able to capture annual BC mean value but completely missed the concentrations during pre-monsoon and post monsoon period (Adhikary et al., 2007). Similar behavior is seen this time for CO where the model misses the peak values but reasonably captures CO concentration after mid-May when no biomass burning events are observed (model to observation ratio improves to 1.16). STEM model CO performance can be significantly improved via better constraining emissions of open biomass burning as discussed in Section 3.3. This activity is beyond the scope of this current paper although the improvements are underway for all these sectors.

3.2.3 Diurnal variations of air pollutants and boundary layer height

In the emission source region, diurnal variations of primary pollutants provide information about the time dependent emission activities (Kumar et al., 2016b). Figure 5 shows the diurnal

variation of hourly averaged concentrations of measured pollutants during the sampling period. Primary pollutants like BC, PM and CO showed typical characteristics of an urban environment, i.e., diurnal variation with a morning and an evening peak. However, Lumbini data shows higher concentrations in the evenings compared to morning hours. Elevated concentrations can be linked to morning and evening cooking hours for BC and CO where emission inventory show that residential sector has significant contribution. However, explanation for elevated evening concentration compared to morning needs further investigation. Increase in the depth of boundary layer, reduction in the traffic density on the roads, absence of open biomass burning during mid-day and increasing wind speed often contribute to the dispersion of pollutants resulting in lower concentration during afternoon. Diurnal variation of wind direction (Supplementary information, Figure S2, upper panel) shows the dominance of wind coming from south (mainly during the month of May and till mid-June). Morning and evening period experienced the winds coming from the southeast direction while the winds were predominantly from southwest direction during late afternoon. Increase in CO concentrations in the evening hours might be due to transport of CO from source regions upwind of Lumbini which along with the local emissions gets trapped under reduced Planetary Boundary Layer (PBL) heights. Ozone concentration was lowest in the morning before the sunrise and highest in late afternoon around 15:00 PM after which concentrations started declining, exhibiting a typical characteristic of a polluted urban site. Photo-dissociation of accumulated NO_x reservoirs (like HONO) provides sufficient NO concentration leading to the titration of O₃ resulting in minimum O₃ just before sunrise (Kumar et al., 2016b). The PBL height (in meters (m)) was obtained from the WRF model as observations were not available. The study period average PBL height over Lumbini was ~ 910 m (ranging between 24.28 and 3807 m observed at 06:00 and 15:00 respectively). The daily average PBL height obtained from the model is compared with published values (Wan et al., 2017) as shown in Figure 6, which indicate that the value is captured by our model during initial measurement period and overestimated in the months of mid May onwards. As the pre-monsoon month advances, PBL height also increased. The monthly average PBL height was 799 m, 956 m and 1014 m respectively during the month of April, May and (1st-15th) June. As presented in the figure, the monthly average diurnal variation also showed that the boundary

layer height was maximum during 15:00 local time during each month which coincides with the period of lowest concentration of the pollutants.

3.3 Influence of forest fires on Lumbini air quality

3.3.1 Identification of forest fire influence over large scale using in-situ observations satellite and model data

Forest fires and agricultural biomass burning (mostly agro-residue burning in large scale) are common over the South Asia and the IGP region during pre-monsoon season. North Indo-Gangetic region is characterized by fires even during the monsoon and post-monsoon season (Kumar et al., 2016b; Putero et al., 2014). These activities influence air quality not only over nearby regions but also get transported towards high elevation pristine environments like Mt. Everest (Putero et al., 2014) and Tibet (Cong et al., 2015a;2015b). So, one of the main objectives of this study was to identify the influence of open burning on Lumbini air quality. Average wind speed during the whole measurement period was 2.4 m s^{-1} . Based on this data, open fire counts within the grid size of $200 \times 200 \text{ km}$ centering over Lumbini was used for this analysis assuming that the emissions will take a maximum period of one day to reach our monitoring site. Forest fire counts were obtained from MODIS satellite data product **Fire Information for Resource Management System (FIRMS)**. More on this product has already been described by Putero et al. (2014). Figure 7 shows the daily average $\Delta\text{BC}/\Delta\text{CO}$ ratio, aerosol absorption Ångström exponent (AAE) which is derived from Aethalometer data and daily open fire count within the specified grid. The green box in the figure is used to show two peak events (presented earlier in Fig. 3) with the elevated BC and CO concentrations observed during the monitoring period. The first peak was observed during 7-9 April and second peak during 3-4 May, 2013. Two pollutants having biomass burning as the potential primary source: BC and CO were taken in consideration. High AAE values (~ 1.6) during these two events are also an indication of presence of BC of biomass burning origin. The chemical composition of TSP filter samples collected at Lumbini also showed higher concentration of Levoglucosan, a biomass burning tracer in Lumbini during the pre-monsoon season as compared to other seasons of the year (Wan et al., 2017). Wan et al. (2017) also reported that the higher correlation between K^+ with tracers of dust (Ca^{2+} and Mg^{2+}) indicated that dust is the main source of potassium in Lumbini.

489 Contrary to our expectation, we could not observe any significant influence of forest fire within
 490 the specified grid of 200x200 km (or the influence of local forest fire on the air quality over
 491 Lumbini was not observed). Therefore, a wider area, covering South and Southeast Asian
 492 regions, was selected for the forest fire count. Figure 8 (A-B) shows the active fire hotspots from
 493 MODIS, over the region, during the peak events which shows the first peak could have occurred
 494 due to the forest fire over the eastern India region whereas the second peak was influenced by the
 495 forest fire over western IGP region. Moreover, in order to strengthen our hypothesis, we have
 496 utilized satellite data products for various gaseous pollutants like CO and NO₂ (Atmospheric
 497 Infrared Sounder (AIRS) for CO and Ozone Monitoring Instrument (OMI) for NO₂ both
 498 obtained from Giovanni platform). Figure 8 (C-H) shows the daytime total column CO before,
 499 during and after occurrence of two events (peaks) as stated earlier. Atmospheric Infrared
 500 Sounder (AIRS) satellite with daily temporal resolution and 1°×1° spatial resolution have been
 501 utilized to understand the CO concentration over the area. CO concentration over Lumbini
 502 during both of the peaks confirmed the role of open fires over the IGP region for elevated
 503 concentration of CO in Lumbini. To further strengthen our finding, the aid of HYSPLIT back
 504 trajectories plots was taken. Figure 8 (I-J) represent the 6-hourly back trajectories only for these
 505 two events respectively. However, the back trajectories (during both events) indicated that the air
 506 mass passed over the fire events in the north western IGP. We note that using back trajectories to
 507 identify source regions are also uncertain as identified by Jaffe et al. (1997). Figure 8 (K) shows
 508 model biomass CO peak coincident with observed CO. Although the magnitudes are
 509 significantly different, the timing of the peaks is well captured by the model. This, we believe, is
 510 due to the fact that satellite based open fire detection also has limitation as it does not capture
 511 numerous small fires that are prevalent over south Asia which usually burn out before the next
 512 satellite overpass. More research is needed to assess the influence of these small fires on regional
 513 air quality.

514 In a separate analysis (not shown here), elevated O₃ concentration during these two events were
 515 also observed. Average O₃ concentration before, during and after the events were found to be
 516 46.2±20.3 ppbv, 53.5±31.1 ppbv and 50.3±20.9 ppbv respectively (Event-I) whereas it was
 517 found to be 54.8±23.8 ppbv, 56.7±35 ppbv and 55.6±13.4 ppbv respectively (Event-II).
 518 Increased ozone concentrations during the high peak events have been analyzed using the

satellite NO₂ concentration over the region considering the role of NO₂ as precursor for ozone formation. Daily total column NO₂ were obtained from OMI satellite (data available at the Giovanni platform; <http://giovanni.gsfc.nasa.gov/giovanni/>) at the spatial resolution of 0.25°×0.25°. Figure 9 shows the NO₂ column value before, during and after both events. Even for the NO₂, maximum concentrations were observed during these two special events. It is likely that the local as well as regional pollution (transported from NW IGP region as indicated by synoptic wind in Figure S8) contributed to the ozone peak. However, we are not able to quantify the individual contributions, even with the model simulation because ozone was not simulated in this experiment.

3.3.2 Identifying regional and local contribution

WRF-STEM model has been used to identify the anthropogenic emission source region influencing the air quality over Lumbini. As previously explained, the model is able to capture the observed CO concentration when intense open burning events were not present. A recent study (Kulkarni et al., 2015) has explored the source region contribution of various pollutants over the Central Asia using similar technique. Figure 10 (A) shows the average contribution from different regions on CO concentration over Lumbini during the whole measurement period. Major share of CO was from the Ganges valley (46%) followed by Nepal region (25%) and rest of Indian region (~17.5%). Contribution from other South Asian countries like Bangladesh and Pakistan were ~ 11% whereas China contributed for ~1% of the CO concentration in Lumbini. Regarding the monthly average contribution, the Ganges Valley and Nepal's contribution were almost equal during the month of April (~34% and ~37% respectively) but increased for the Ganges Valley region during the month of May (~44%) and got reduced for Nepal region (~25%) (Figure S9).

Figure 10 (B) is the time series of percentage contribution to total CO concentration during whole measurement period showing different air mass arriving at a 3 hourly intervals. During the whole measurement period, majority of the CO reaching Lumbini were from the Ganges valley (mainly the states of Punjab, Haryana, Uttar Pradesh, Bihar and West Bengal) region with the contribution sometimes reaching up to ~80%. Other India (central, south, east and north) regions also contributed significantly. Bangladesh's contribution in CO loading was seen only after mid-

April lasting for only about a week and after the first week of May. The contribution from Bangladesh was sporadic comparing to other regions. Highest contribution from this Bangladesh region was observed after the first week of June with the arrival of monsoonal air mass. Pakistan also contributed for the CO loading significantly. Others region as mentioned in the figure covered the regions like Afghanistan, Middle east, West Asia, East Asia, Africa and Bhutan. Contributions from these regions were less than 5%. Contribution from China was not evident till the first week of June where a specific air mass arrival shows contribution reaching up to 25% of total CO loading.

A sensitivity analysis was performed for emission uncertainty in the model grid containing Lumbini. Lumbini and surrounding regions in the recent years has seen significant rise in urban activities and industrial activity and related emissions which may not be accurately reflected in the HTAPv2 emissions inventory. A month long simulation was carried out with emissions from Lumbini and the surrounding four grids off and another simulation with Lumbini and surrounding four grid's emissions increased by 5 times the amount from HTAPv2 emissions inventory. The results are shown in Figure 10 (C) as percentage increase or decrease compared to model results using the current HTAPv2 emissions inventory. The black line shows concentration as 100% for the current HTAPv2 emissions inventory. Despite making Lumbini and the surrounding grids emissions zero, model calculation shows pollutant concentration on average is still about 78% of the original value indicating dominance of background and regional sources compared to local source in the model. Increasing emissions 5 times for the Lumbini and surrounding four grids only increases the concentration on average by 151%. Thus uncertainty in emissions are not a local uncertainty for Lumbini rather for the whole region which needs to be better understood for improving model performance against observations at Lumbini.

3.4 Does fossil fuel or biomass influence the Lumbini air?

The aerosol spectral absorption is used to gain insight into nature and potential source of black carbon. This method enables to analyze the contributions of fossil fuel combustion and biomass burning contributions to the observed BC concentration (Kirchstetter et al., 2004). Besides BC, other light absorbing (in the UV region) aerosols are also produced in course of combustion, collectively termed as organic aerosols (often also called brown carbon or BrC) (Andreae and

Gelencsér, 2006). Figure 11 shows the comparison of normalized light absorption as function of the wavelength for BC observed at Lumbini during cooking and non-cooking hours and also for the both events. Our results are compared with the published data of Kirchstetter et al. (2004) and that observed over a village center site of Project Surya in the IGP (Praveen et al., 2012) (figure not shown). We discuss light absorption data from two distinct times of the day. The main reason behind using data during 07:00-08:00 h and 16:00-17:00 h is these periods represent highest and lowest ambient concentration (Fig. 5). Also these period represent cooking and non-cooking or high and low vehicular movement hours (Praveen et al., 2012). To understand the influence of biomass and fossil fuel we plotted normalized aerosol absorption at 700 nm wavelength for complete aethalometer measured wavelengths in Fig. 11. Kirchstetter et al. (2004) reported OC absorption efficiency at 700 nm to be zero. Thus we normalized measured absorption spectrum by 700 nm wavelength absorption. Since aethalometer does not provide 700 nm wavelength absorption values, we used methodology followed by Praveen et al. (2012). Our results show that the normalized absorption for biomass burning aerosol is ~3 times higher at 370 nm compared to that at 700 nm whereas fossil fuel absorption is about 2.6 times higher at the same wavelength. In addition, the curve obtained for the both events are inclined towards the published biomass burning curve. The normalized curve obtained during both cooking and non-cooking period lies in between the standard curve of Kirchstetter et al. (2004). As shown in Fig. 11, the curve obtained for the prime cooking time is closer towards the published curve on biomass burning whereas that obtained during the non-cooking time is closer towards the published fossil fuel curve. Similar result was also observed over the Project Surya village in the IGP region (Praveen et al., 2012; Rehman et al., 2011). This clearly indicates there is contribution of both sources: biomass as well as fossil fuel on the observed BC concentration over Lumbini.

In order to identify fractional contribution of biomass burning and fossil fuel combustion to observed BC aerosol, we adopted the method described by Sandradewi et al. (2008). Wavelength dependence of aerosol absorption coefficient (b_{abs}) is proportional to $\lambda^{-\alpha}$ where λ is the wavelength and α is the absorption Ångstrom exponent. The α values ranges from 0.9-2.2 for fresh wood smoke aerosol (Day et al., 2006) and between 0.8-1.1 for traffic or diesel soot (references in Sandradewi et al. (2008)). We have taken α value of 1.86 for biomass burning and 1.1 for fossil fuel burning as suggested by previous literature (Sandradewi et al., 2008). Figure

12 shows diurnal variation of the biomass burning BC. Minimum contribution of biomass burning to total BC concentration was observed during 04:00-06:00 local time (only about 30% of the total BC). As the cooking activities start in morning, the contribution of biomass BC starts to increase and reaches about 50%. Similar pattern was repeated during evening cooking hours. Only during these two cooking periods, fossil fuel fraction BC was lower. Otherwise it remained significantly higher than biomass burning BC throughout the day. On average, ~40% of BC was from biomass burning whereas remaining 60% was contributed by fossil fuel combustion during our measurement period. Interestingly, this is the opposite of the contributions that were concluded by Lawrence and Lelieveld (2010). Lawrence and Lelieveld (2010) concluded that ~60% BC from biomass versus ~40% fossil fuel, based on a review of numerous previous studies to be likely for the outflow from Southern Asia during the winter monsoon. When we compared observed Ångström exponent with Praveen et al. (2012), we noticed that Lumbini values were lower than Project Surya Village center site. This implies Surya village center had higher biomass fraction, also it was observed absorption Ångström exponent exceeded 1.86 during cooking hours which indicates 100% biomass contribution. The difference is attributed to the fact that Lumbini sampling site is not a residential site like Surya village which can capture cooking influence efficiently. Further Lumbini sampling site is surrounded by commercial activities such as a local bus park, hotels, office buildings and industries and brick kilns slightly further away. Although the reason for this difference is not clear, it is an indication of the important role of diesel and coal emissions in the Lumbini and upwind regions.

4. Conclusions

Our measurements, a first for the Lumbini area, have shown very high air pollution at Lumbini. Black carbon (BC), carbon monoxide (CO), ozone (O₃) and particulate matter (PM₁₀, PM_{2.5} and PM₁) were measured during the pre-monsoon of 2013 as a regional site of the *SusKat-ABC campaign*. Average pollutant concentrations during the monitoring period were found to be: BC: $4.9 \pm 3.8 \mu\text{g m}^{-3}$; CO: 344.1 ± 160.3 ppbv; O₃: 46.6 ± 20.3 ppbv; PM₁₀: $128.8 \pm 91.9 \mu\text{g m}^{-3}$ PM_{2.5}: $53.14 \pm 35.1 \mu\text{g m}^{-3}$ and PM₁: $36.6 \pm 25.7 \mu\text{g m}^{-3}$ which is comparable with other urban sites like Kanpur and Delhi in the IGP region. However, our study finds higher fraction of coarse mode PM in Lumbini as compared to other sites in the IGP region. In addition, $\Delta\text{BC}/\Delta\text{CO}$ ratio obtained in Lumbini was within the range of emission from both residential and transportation

sectors, indicating them as potential key sources of BC and CO, and likely most of PM₁ in Lumbini. The diurnal variation of the pollutants is similar to that of any urban location, with peaks during morning and evening. However, our results show higher evening concentration compared to morning concentration values and needs further research to explain this behavior. During our measurement period, air quality in Lumbini was influenced by regional forest fires as shown by chemical transport model and satellite data analysis. A regional chemical transport model, WRF-STEM was used to **understand** observations. Inter-comparison of WRF-STEM model outputs with observations showed that the model underestimated the observed pollutant concentrations by a factor of ~ 1.5 to 5 but was able to capture the temporal variability. Model uncertainties are attributed mostly to uncertainties in meteorology and regional emissions as shown from sensitivity analysis with local emissions. Region-tagged CO as air-mass tracers are employed in WRF-STEM model to understand the anthropogenic emission source region influencing Lumbini. Our analysis shows that the adjacent regions; mostly the Ganges valley, other parts of India and Nepal accounted for the highest contribution to pollutant concentration in the Lumbini. The normalized light absorption curve clearly indicated the contribution to BC in Lumbini from both sources: biomass as well as fossil fuel. On average, ~40% BC was found to be from the biomass burning and ~60% from fossil fuel burning.

Various improvements and extensions would be possible in future studies. More reliable functioning of the AWS (temperature and RH sensor, rain gauge) would have allowed more in-depth analysis of the relationship between meteorological parameters and pollutants concentration. Continuous measurements of air pollutants throughout the year would allow for annual and seasonal variation study. Improvements in the **model** are much needed in its ability to simulate observed meteorology. Significant uncertainty lies with regional emissions inventory developed at national and continental scale versus local bottoms up inventory and pollutant emissions from small scale open burning not captured by satellites. There is a clear need for setting up of a continuous air quality monitoring station at Lumbini and the surrounding regions for long-term air quality monitoring.

Data availability

The data used for this manuscript can be obtained by sending an email to the corresponding authors and/or to IASS (Maheswar.Rupakheti@iass.potsdam.de) and/or to ICIMOD (arnico.panday@icimod.org). Modeling data can be obtained from B. Adhikary (Bhupesh.adhikary@icimod.org).

Authors' contributions

M.R. and M.L. conceived the Lumbini portion of the SusKat experiment. M.R. and A.K.P. coordinated the Lumbini field campaign. D.R. and K.S.M conducted the field observations at Lumbini. B.A. designed and ran the WRF-STEM model. P.S.P., B.A. and D.R. finalized the manuscript composition. D.R., P.S.P, B.A., M.R. and S.K. conducted the data analysis. D.R. and B.A. prepared the manuscript with inputs from all coauthors.

Acknowledgements

This study was partly supported by the Institute for Advanced Sustainability Studies (IASS), Germany, the International Centre for Integrated Mountain Development (ICIMOD), and the National Natural Science Foundation of China (41121001, 41225002), and the Strategic Priority Research Program (B) of the Chinese Academy of Sciences (XDB03030504). Dipesh Rupakheti is supported by CAS-TWAS President's Fellowship for International PhD Students. The IASS is grateful for its funding from the German Federal Ministry for Education and Research (BMBF) and the Brandenburg Ministry for Science, Research and Culture (MWFK). ICIMOD authors would like to acknowledge that this study was partially supported by core funds of ICIMOD contributed by the governments of Afghanistan, Australia, Austria, Bangladesh, Bhutan, China, India, Myanmar, Nepal, Norway, Pakistan, Switzerland, and the United Kingdom. The views and interpretations in this publication are those of the authors and are not necessarily attributable to the institutions they are associated with. We thank B. Kathayat, B.R. Bhatta, and Venerable Vivekananda and his colleagues (Panditarama Lumbini International Vipassana Meditation Center) for providing logistical support which was vital in setting up and running the site. We also thank C. Cüppers and M. Pahlke of the Lumbini International Research Institute (LIRI) for providing the space and power to run the instruments at the LIRI premises. Satellite data providers (MODIS, AIRS, OMI) and **HYSPLIT team** are also equally acknowledged.

694 **References**

- 695 Adhikary, B., Carmichael, G. R., Tang, Y., Leung, L. R., Qian, Y., Schauer, J. J., Stone, E. A.,
696 Ramanathan, V., and Ramana, M. V.: Characterization of the seasonal cycle of south Asian
697 aerosols: A regional-scale modeling analysis, *Journal of Geophysical Research*, 112,
698 D22S22, 1-22, 10.1029/2006jd008143, 2007.
- 699 Andreae, M. O., and Gelencsér, A.: Black carbon or brown carbon? The nature of light-
700 absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131-3148, doi: 10.5194/acp-6-
701 3131-2006, 2006.
- 702 Baedecker, P. A., Reddy, M. M., Reimann, K. J., and Sciammarella, C. A.: Effects of acidic
703 deposition on the erosion of carbonate stone — experimental results from the U.S. National
704 Acid Precipitation Assessment Program (NAPAP), *Atmos. Environ.*, 26, 147-158, doi:
705 10.1016/0957-1272(92)90018-N, 1992.
- 706 Bergin, M. H., Tripathi, S. N., Jai Devi, J., Gupta, T., McKenzie, M., Rana, K., Shafer, M. M.,
707 Villalobos, A. M., and Schauer, J. J.: The Discoloration of the Taj Mahal due to Particulate
708 Carbon and Dust Deposition, *Environ. Sci. Technol.*, 49, 808-812, doi: 10.1021/es504005q,
709 2015.
- 710 Bisht, D. S., Dumka, U. C., Kaskaoutis, D. G., Pipal, A. S., Srivastava, A. K., Soni, V. K., Attri,
711 S. D., Sateesh, M., and Tiwari, S.: Carbonaceous aerosols and pollutants over Delhi urban
712 environment: Temporal evolution, source apportionment and radiative forcing, *Sci. Total*
713 *Environ.*, 521-522C, 431-445, doi: 10.1016/j.scitotenv.2015.03.083, 2015.
- 714 Bonasoni, P., Laj, P., Marinoni, A., Sprenger, M., Angelini, F., Arduini, J., Bonafè, U., Calzolari,
715 F., Colombo, T., Decesari, S., Di Biagio, C., di Sarra, A. G., Evangelisti, F., Duchi, R.,
716 Facchini, M. C., Fuzzi, S., Gobbi, G. P., Maione, M., Panday, A., Roccato, F., Sellegri, K.,
717 Venzac, H., Verza, G. P., Villani, P., Vuillermoz, E., and Cristofanelli, P.: Atmospheric
718 Brown Clouds in the Himalayas: first two years of continuous observations at the Nepal
719 Climate Observatory-Pyramid (5079 m), *Atmospheric Chemistry and Physics*, 10, 7515-
720 7531, doi: 10.5194/acp-10-7515-2010, 2010.
- 721 Brimblecombe, P.: *The effects of air pollution on the built environment*, Imperial College Press,
722 London, 2003.
- 723 Burney, J., and Ramanathan, V.: Recent climate and air pollution impacts on Indian agriculture,
724 *Proc. Natl. Acad. Sci. USA*, 111, 16319-16324, doi: 10.1073/pnas.1317275111, 2014.
- 725 Cao, J.-J., Zhu, C.-S., Chow, J. C., Watson, J. G., Han, Y.-M., Wang, G.-h., Shen, Z.-x., and An,
726 Z.-S.: Black carbon relationships with emissions and meteorology in Xi'an, China,
727 *Atmospheric Research*, 94, 194-202, <http://dx.doi.org/10.1016/j.atmosres.2009.05.009>, 2009.
- 728 Cong, Z., Kang, S., Kawamura, K., Liu, B., Wan, X., Wang, Z., Gao, S., and Fu, P.:
729 Carbonaceous aerosols on the south edge of the Tibetan Plateau: concentrations, seasonality
730 and sources, *Atmospheric Chemistry and Physics*, 15, 1573-1584, doi: 10.5194/acp-15-
731 1573-2015, 2015a.
- 732 Cong, Z., Kawamura, K., Kang, S., and Fu, P.: Penetration of biomass-burning emissions from
733 South Asia through the Himalayas: new insights from atmospheric organic acids, *Scientific*
734 *Reports*, 5, 1-7, doi: 10.1038/srep09580, 2015b.
- 735 Cristofanelli, P., Bracci, A., Sprenger, M., Marinoni, A., Bonafè, U., Calzolari, F., Duchi, R.,
736 Laj, P., Pichon, J. M., Roccato, F., Venzac, H., Vuillermoz, E., and Bonasoni, P.:
737 Tropospheric ozone variations at the Nepal Climate Observatory-Pyramid (Himalayas, 5079

- m a.s.l.) and influence of deep stratospheric intrusion events, *Atmospheric Chemistry and Physics*, 10, 6537-6549, doi: 10.5194/acp-10-6537-2010, 2010.
- Das, S. K., and Jayaraman, A.: Role of black carbon in aerosol properties and radiative forcing over western India during premonsoon period, *Atmos. Res.*, 102, 320-334, doi: 10.1016/j.atmosres.2011.08.003, 2011.
- Day, D., Hand, J., Carrico, C., Engling, G., and Malm, W.: Humidification factors from laboratory studies of fresh smoke from biomass fuels, *J. Geophys. Res.*, 111, D22202, doi: 10.1029/2006JD007221, 2006.
- Dickerson, R. R., Andreae, M. O., Campos, T., Mayol-Bracero, O. L., Neusuess, C., and Streets, D. G.: Analysis of black carbon and carbon monoxide observed over the Indian Ocean: Implications for emissions and photochemistry, *Journal of Geophysical Research: Atmospheres*, 107, INX2 16-11-INX12 16-11, 10.1029/2001JD000501, 2002.
- Duchi, R., Cristofanelli, P., Marinoni, A., Laj, P., Marcq, S., Villani, P., Sellegri, K., Angelini, F., Calzolari, F., Gobbi, G. P., Verza, G. P., Vuillermoz, E., Sapkota, A., and Bonasoni, P.: Continuous observations of synoptic-scale dust transport at the Nepal Climate Observatory-Pyramid (5079 m a.s.l.) in the Himalayas, *Atmos. Chem. Phys. Discuss.*, 11, 4229-4261, doi: 10.5194/acpd-11-4229-2011, 2011.
- Forouzanfar, M. H., Alexander, L., Anderson, H. R., Bachman, V. F., Biryukov, S., Brauer, M., Burnett, R., Casey, D., Coates, M. M., and Cohen, A.: Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013, *Lancet*, 386, 2287-2323, doi: 10.1016/S0140-6736(15)00128-2, 2015.
- Ganguly, D., Jayaraman, A., Rajesh, T. A., and Gadhavi, H.: Wintertime aerosol properties during foggy and nonfoggy days over urban center Delhi and their implications for shortwave radiative forcing, *Journal of Geophysical Research*, 111, 1-15, 10.1029/2005jd007029, 2006.
- Gaur, A., Tripathi, S. N., Kanawade, V. P., Tare, V., and Shukla, S. P.: Four-year measurements of trace gases (SO₂, NO_x, CO, and O₃) at an urban location, Kanpur, in Northern India, *Journal of Atmospheric Chemistry*, 71, 283-301, doi: 10.1007/s10874-014-9295-8, 2014.
- Gauri, K. L., and Holdren, G.: Pollutant effects on stone monuments, *Environ. Sci. Technol.*, 15, 386-390, doi: 10.1021/es00086a001, 1981.
- Girach, I. A., Nair, V. S., Babu, S. S., and Nair, P. R.: Black carbon and carbon monoxide over Bay of Bengal during W_ICARB: Source characteristics, *Atmospheric Environment*, 94, 508-517, <http://dx.doi.org/10.1016/j.atmosenv.2014.05.054>, 2014.
- Huang, X.-F., Sun, T.-L., Zeng, L.-W., Yu, G.-H., and Luan, S.-J.: Black carbon aerosol characterization in a coastal city in South China using a single particle soot photometer, *Atmospheric Environment*, 51, 21-28, 10.1016/j.atmosenv.2012.01.056, 2012.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. , Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.
- Jaffe, D., Mahura, A., Kelley, J., Atkins, J., Novelli, P. C., and Merrill, J.: Impact of Asian emissions on the remote North Pacific atmosphere: Interpretation of CO data from Shemya,

- Guam, Midway and Mauna Loa, *Journal of Geophysical Research*, 102, 28627, 10.1029/96jd02750, 1997.
- Joshi, H., Naja, M., Singh, K., Kumar, R., Bhardwaj, P., Babu, S. S., Satheesh, S., Moorthy, K. K., and Chandola, H.: Investigations of aerosol black carbon from a semi-urban site in the Indo-Gangetic Plain region, *Atmos. Environ.*, 125, 346-359, doi: 10.1016/j.atmosenv.2015.04.007, 2016.
- Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *Journal of Geophysical Research*, 109, 1-12, doi:10.1029/2004JD004999, 2004.
- Kulkarni, S., Sobhani, N., Miller-Schulze, J. P., Shafer, M. M., Schauer, J. J., Solomon, P. A., Saide, P. E., Spak, S. N., Cheng, Y. F., Denier van der Gon, H. A. C., Lu, Z., Streets, D. G., Janssens-Maenhout, G., Wiedinmyer, C., Lantz, J., Artamonova, M., Chen, B., Imashev, S., Sverdlik, L., Deminter, J. T., Adhikary, B., D'Allura, A., Wei, C., and Carmichael, G. R.: Source sector and region contributions to BC and PM_{2.5} in Central Asia, *Atmos. Chem. Phys.*, 15, 1683-1705, doi: 10.5194/acp-15-1683-2015, 2015.
- Kumar, B., Chakraborty, A., Tripathi, S. N., and Bhattu, D.: Highly time resolved chemical characterization of submicron organic aerosols at a polluted urban location, *Environmental Science: Processes & Impacts*, 18, 1285-1296, 10.1039/C6EM00392C, 2016a.
- Kumar, M., Tiwari, S., Murari, V., Singh, A. K., and Banerjee, T.: Wintertime characteristics of aerosols at middle Indo-Gangetic Plain: Impacts of regional meteorology and long range transport, *Atmospheric Environment*, 104, 162-175, 10.1016/j.atmosenv.2015.01.014, 2015.
- Kumar, V., Sarkar, C., and Sinha, V.: Influence of post harvest crop residue fires on surface ozone mixing ratios in the NW IGP analyzed using two years of continuous in-situ trace gas measurements, *J. Geophys. Res. Atmos.*, 121, 3619-3633, doi: 10.1002/2015JD024308, 2016b.
- Lawrence, M. G., and Lelieveld, J.: Atmospheric pollutant outflow from southern Asia: a review, *Atmos. Chem. Phys.*, 10, 11017-11096, doi: 10.5194/acp-10-11017-2010, 2010.
- Lüthi, Z. L., Škerlak, B., Kim, S. W., Lauer, A., Mues, A., Rupakheti, M., and Kang, S.: Atmospheric brown clouds reach the Tibetan Plateau by crossing the Himalayas, *Atmospheric Chemistry and Physics*, 15, 6007-6021, 10.5194/acp-15-6007-2015, 2015.
- Marinoni, A., Cristofanelli, P., Laj, P., Duchi, R., Putero, D., Calzolari, F., Landi, T. C., Vuillermoz, E., Maione, M., and Bonasoni, P.: High black carbon and ozone concentrations during pollution transport in the Himalayas: Five years of continuous observations at NCO-P global GAW station, *Journal of environmental Science* 25, 1618-1625, doi: 10.1016/s1001-0742(12)60242-3, 2013.
- Misra, A., Gaur, A., Bhattu, D., Ghosh, S., Dwivedi, A. K., Dalai, R., Paul, D., Gupta, T., Tare, V., Mishra, S. K., Singh, S., and Tripathi, S. N.: An overview of the physico-chemical characteristics of dust at Kanpur in the central Indo-Gangetic basin, *Atmospheric Environment*, 97, 386-396, <http://dx.doi.org/10.1016/j.atmosenv.2014.08.043>, 2014.
- Mohnen, V. A., Goldstein, W., and Wang, W. C.: Tropospheric Ozone and Climate Change, *Air & Waste*, 43, 1332-1334, 10.1080/1073161x.1993.10467207, 1993.
- Naja, M., Lal, S., and Chand, D.: Diurnal and seasonal variabilities in surface ozone at a high altitude site Mt Abu (24.6° N, 72.7° E, 1680m asl) in India, *Atmospheric Environment*, 37, 4205-4215, doi: 10.1016/S1352-2310(03)00565-X, 2003.

- Naja, M., Mallik, C., Sarangi, T., Sheel, V., and Lal, S.: SO₂ measurements at a high altitude site in the central Himalayas: Role of regional transport, *Atmos. Environ.*, 99, 392-402, doi: 10.1016/j.atmosenv.2014.08.031, 2014.
- Pan, X. L., Kanaya, Y., Wang, Z. F., Liu, Y., Pochanart, P., Akimoto, H., Sun, Y. L., Dong, H. B., Li, J., Irie, H., and Takigawa, M.: Correlation of black carbon aerosol and carbon monoxide in the high-altitude environment of Mt. Huang in Eastern China, *Atmos. Chem. Phys.*, 11, 9735-9747, doi: 10.5194/acp-11-9735-2011, 2011.
- Panday, A. K., and Prinn, R. G.: Diurnal cycle of air pollution in the Kathmandu Valley, Nepal: Observations, *Journal of Geophysical Research*, 114, 1-19, doi: 10.1029/2008jd009777, 2009.
- Pathak, B., Bhuyan, P. K., Biswas, J., and Takemura, T.: Long term climatology of particulate matter and associated microphysical and optical properties over Dibrugarh, North-East India and inter-comparison with SPRINTARS simulations, *Atmospheric Environment*, 69, 334-344, 10.1016/j.atmosenv.2012.12.032, 2013.
- Praveen, P. S., Ahmed, T., Kar, A., Rehman, I. H., and Ramanathan, V.: Link between local scale BC emissions in the Indo-Gangetic Plains and large scale atmospheric solar absorption, *Atmospheric Chemistry and Physics*, 12, 1173-1187, 10.5194/acp-12-1173-2012, 2012.
- Putero, D., Landi, T., Cristofanelli, P., Marinoni, A., Laj, P., Duchi, R., Calzolari, F., Verza, G., and Bonasoni, P.: Influence of open vegetation fires on black carbon and ozone variability in the southern Himalayas (NCO-P, 5079 m asl), *Environmental Pollution*, 184, 597-604, 10.1016/j.envpol.2013.09.035, 2014.
- Putero, D., Cristofanelli, P., Marinoni, A., Adhikary, B., Duchi, R., Shrestha, S. D., Verza, G. P., Landi, T. C., Calzolari, F., Busetto, M., Agrillo, G., Biancofiore, F., Di Carlo, P., Panday, A. K., Rupakheti, M., and Bonasoni, P.: Seasonal variation of ozone and black carbon observed at Paknajol, an urban site in the Kathmandu Valley, Nepal, *Atmospheric Chemistry and Physics*, 15, 13957-13971, 10.5194/acp-15-13957-2015, 2015.
- Ram, K., Sarin, M., and Tripathi, S.: A 1 year record of carbonaceous aerosols from an urban site in the Indo-Gangetic Plain: Characterization, sources, and temporal variability, *J. Geophys. Res.*, 115, D24313, doi: 10.1029/2010JD014188, 2010.
- Ramachandran, S., and Rajesh, T. A.: Black carbon aerosol mass concentrations over Ahmedabad, an urban location in western India: Comparison with urban sites in Asia, Europe, Canada, and the United States, *Journal of Geophysical Research*, 112, 1-19, 10.1029/2006JD007488, 2007.
- Ramanathan, V., Li, F., Ramana, M., Praveen, P., Kim, D., Corrigan, C., Nguyen, H., Stone, E. A., Schauer, J. J., and Carmichael, G.: Atmospheric brown clouds: Hemispherical and regional variations in long-range transport, absorption, and radiative forcing, *Journal of Geophysical Research*, 112, 1-26, 10.1029/2006JD008124, 2007.
- Ramanathan, V., and Carmichael, G.: Global and regional climate changes due to black carbon, *Nature geoscience*, 1, 221-227, 2008.
- Rastogi, N., Singh, A., Sarin, M. M., and Singh, D.: Temporal variability of primary and secondary aerosols over northern India: Impact of biomass burning emissions, *Atmos. Environ.*, 125, 396-403, doi: 10.1016/j.atmosenv.2015.06.010, 2016.
- Rehman, I. H., Ahmed, T., Praveen, P. S., Kar, A., and Ramanathan, V.: Black carbon emissions from biomass and fossil fuels in rural India, *Atmospheric Chemistry and Physics*, 11, 7289-7299, 10.5194/acp-11-7289-2011, 2011.

- Safai, P. D., Kewat, S., Pandithurai, G., Praveen, P. S., Ali, K., Tiwari, S., Rao, P. S. P., Budhawant, K. B., Saha, S. K., and Devara, P. C. S.: Aerosol characteristics during winter fog at Agra, North India, *J. Atmos. Chem.*, 61, 101-118, doi: 10.1007/s10874-009-9127-4, 2009.
- Sandradewi, J., Prévôt, A. S. H., Szidat, S., Perron, N., Alfarra, M. R., Lanz, V. A., Weingartner, E., and Baltensperger, U.: Using Aerosol Light Absorption Measurements for the Quantitative Determination of Wood Burning and Traffic Emission Contributions to Particulate Matter, *Environmental Science & Technology*, 42, 3316-3323, doi: 10.1021/es702253m, 2008.
- Sarkar, C., Kumar, V., and Sinha, V.: Massive emissions of carcinogenic benzenoids from paddy residue burning in north India, *Curr. Sci. India*, 104, 1703-1709, 2013.
- Schmid, O., Artaxo, P., Arnott, W. P., Chand, D., Gatti, L. V., Frank, G. P., Hoffer, A., Schnaiter, M., and Andreae, M. O.: Spectral light absorption by ambient aerosols influenced by biomass burning in the Amazon Basin. I: Comparison and field calibration of absorption measurement techniques, *Atmospheric Chemistry and Physics*, 6, 3443-3462, doi: 10.5194/acp-6-3443-2006, 2006.
- Sharma, R. K., Bhattarai, B. K., Sapkota, B. K., Gewali, M. B., and Kjeldstad, B.: Black carbon aerosols variation in Kathmandu valley, Nepal, *Atmos. Environ.*, 63, 282-288, 10.1016/j.atmosenv.2012.09.023, 2012.
- Sharma, S., Brook, J. R., Cachier, H., Chow, J., Gaudenzi, A., and Lu, G.: Light absorption and thermal measurements of black carbon in different regions of Canada, *Journal of Geophysical Research: Atmospheres*, 107 (D24), 1-11, 10.1029/2002JD002496, 2002.
- Shindell, D.: Integrated Assessment of Black Carbon and Tropospheric Ozone: Summary for Decision Makers, United Nations Environment Programme, 303pp., 2011.
- Singh, N., Murari, V., Kumar, M., Barman, S. C., and Banerjee, T.: Fine particulates over South Asia: Review and meta-analysis of PM_{2.5} source apportionment through receptor model, *Environmental Pollution*, 223, 121-136, 10.1016/j.envpol.2016.12.071, 2017.
- Sinha, V., Kumar, V., and Sarkar, C.: Chemical composition of pre-monsoon air in the Indo-Gangetic Plain measured using a new air quality facility and PTR-MS: high surface ozone and strong influence of biomass burning, *Atmospheric Chemistry and Physics*, 14, 5921-5941, 10.5194/acp-14-5921-2014, 2014.
- Skamarock, W., Klemp, J., Dudhia, J., Gill, D., and Barker, D.: A description of the Advanced Research WRF version 3, Technical Report NCAR/TN475+STR, National Center for Atmospheric Research Technical Note, Boulder, Colorado, 2008.
- Snider, G., Weagle, C. L., Murdymootoo, K. K., Ring, A., Ritchie, Y., Stone, E., Walsh, A., Akoshile, C., Anh, N. X., Balasubramanian, R., Brook, J., Qonitan, F. D., Dong, J., Griffith, D., He, K., Holben, B. N., Kahn, R., Lagrosas, N., Lestari, P., Ma, Z., Misra, A., Norford, L. K., Quel, E. J., Salam, A., Schichtel, B., Segev, L., Tripathi, S., Wang, C., Yu, C., Zhang, Q., Zhang, Y., Brauer, M., Cohen, A., Gibson, M. D., Liu, Y., Martins, J. V., Rudich, Y., and Martin, R. V.: Variation in global chemical composition of PM_{2.5}: emerging results from SPARTAN, *Atmospheric Chemistry and Physics*, 16, 9629-9653, 10.5194/acp-16-9629-2016, 2016.
- Stevenson, D., Dentener, F., Schultz, M., Ellingsen, K., Van Noije, T., Wild, O., Zeng, G., Amann, M., Atherton, C., and Bell, N.: Multimodel ensemble simulations of present-day and

- near-future tropospheric ozone, *Journal of Geophysical Research*, 111, D08301, 1-23, 10.1029/2005JD006338, 2006.
- Tiwari, S., Srivastava, A. K., Bisht, D. S., Parmita, P., Srivastava, M. K., and Attri, S. D.: Diurnal and seasonal variations of black carbon and PM_{2.5} over New Delhi, India: Influence of meteorology, *Atmospheric Research*, 125-126, 50-62, 10.1016/j.atmosres.2013.01.011, 2013.
- Tiwari, S., Pipal, A. S., Hopke, P. K., Bisht, D. S., Srivastava, A. K., Tiwari, S., Saxena, P. N., Khan, A. H., and Pervez, S.: Study of the carbonaceous aerosol and morphological analysis of fine particles along with their mixing state in Delhi, India: a case study, *Environmental Science and Pollution Research*, 22, 10744-10757, 10.1007/s11356-015-4272-6, 2015.
- Tiwari, S., Dumka, U. C., Kaskaoutis, D. G., Ram, K., Panicker, A. S., Srivastava, M. K., Tiwari, S., Attri, S. D., Soni, V. K., and Pandey, A. K.: Aerosol chemical characterization and role of carbonaceous aerosol on radiative effect over Varanasi in central Indo-Gangetic Plain, *Atmospheric Environment*, 125, 437-449, 10.1016/j.atmosenv.2015.07.031, 2016.
- Tiwari, S., Dumka, U. C., Gautam, A. S., Kaskaoutis, D. G., Srivastava, A. K., Bisht, D. S., Chakrabarty, R. K., Sumlin, B. J., and Solmon, F.: Assessment of PM_{2.5} and PM₁₀ over Guwahati in Brahmaputra River Valley: Temporal evolution, source apportionment and meteorological dependence, *Atmospheric Pollution Research*, 8, 13-28, 10.1016/j.apr.2016.07.008, 2017.
- Trivedi, D. K., Ali, K., and Beig, G.: Impact of meteorological parameters on the development of fine and coarse particles over Delhi, *Science of The Total Environment*, 478, 175-183, 10.1016/j.scitotenv.2014.01.101, 2014.
- Verma, R. L., Sahu, L. K., Kondo, Y., Takegawa, N., Han, S., Jung, J. S., Kim, Y. J., Fan, S., Sugimoto, N., Shammaa, M. H., Zhang, Y. H., and Zhao, Y.: Temporal variations of black carbon in Guangzhou, China, in summer 2006, *Atmos. Chem. Phys.*, 10, 6471-6485, doi: 10.5194/acp-10-6471-2010, 2010.
- Wan, X., Kang, S., Li, Q., Rupakheti, D., Zhang, Q., Guo, J., Chen, P., Tripathi, L., Rupakheti, M., Panday, A. K., Wang, W., Kawamura, K., Gao, S., Wu, G., and Cong, Z.: Organic molecular tracers in the atmospheric aerosols from Lumbini, Nepal, in the northern Indo-Gangetic Plain: Influence of biomass burning, *Atmos. Chem. Phys. Discuss.*, 2017, 1-40, 10.5194/acp-2016-1176, 2017.
- WHO: Air quality guidelines: global update 2005: particulate matter, ozone, nitrogen dioxide, and sulfur dioxide, World Health Organization, Geneva, 22 pp., 2006.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625-641, doi: 10.5194/gmd-4-625-2011, 2011.
- Zhou, X., Gao, J., Wang, T., Wu, W., and Wang, W.: Measurement of black carbon aerosols near two Chinese megacities and the implications for improving emission inventories, *Atmospheric Environment*, 43, 3918-3924, 10.1016/j.atmosenv.2009.04.062, 2009.

957 **Table 1.** Summary of instruments deployed during monitoring in Lumbini

| Instrument (Model) | Manufacturer | Parameters | Inlet/sensor height (above ground) | Sampling interval | Sampled period |
|--|-----------------------------------|--|---------------------------------------|----------------------|-----------------------------|
| Environmental Dust monitor (EDM 164) | GRIMM Aerosol Technik, Germany | PM ₁ , PM _{2.5} , PM ₁₀ | 5 m | 5 min | 04/02-05/10, 06/02-06/13 |
| Aethalometer (AE42) | Magee Scientific, USA | Aerosol light absorption at seven wavelengths, and BC concentration | 3 m | 5 min | 01/04-05/06 |
| CO analyzer (48i) | Thermo Scientific, USA | CO concentration | 3 m | 1 min | 01/04-15/06 |
| O ₃ analyzer (49i) | Thermo Scientific, USA | O ₃ concentration | 3 m | 1 min | 01/04-15/06 |
| Automatic Weather Station (AWS) | Campbell Scientific, UK | T, RH, WS, WD, Global Radiation, Precipitation | 12 m | 1 min | 01/04-15/06 |

958

959

960 **Table 2.** Comparison of PM_{2.5}, BC, CO and O₃ concentrations at Lumbini with those at other sites in South Asia

| Sites | Characteristics | Measurement period | PM _{2.5} (µg m ⁻³) | BC (µg/m ³) | CO (ppbv) | O ₃ (ppbv) | References |
|--------------------|-----------------|--------------------------------|---|-------------------------|-------------|-----------------------|---|
| Lumbini, Nepal | Semi-urban | Pre-monsoon, 2013 | 53.1±35.1 | 4.9±3.8 | 344.1±160.3 | 46.6±20.3 | This study |
| Kathmandu, Nepal | Urban | Pre-monsoon, 2013 | - | 14.5±10 | - | 38.0±25.6 | (Putero et al., 2015) |
| Mt. Everest, Nepal | Remote | Pre-monsoon | - | 0.4±0.4 | - | 61.3±7.7 | (Marinoni et al., 2013) |
| Delhi, India | Urban | Pre-monsoon (night-time) | 82.3±50.5 | 7.70±7.25 | 1800±890 | - | (Bisht et al., 2015) |
| Kanpur, India | Urban | June 2009-May 2013, April-June | - | 2.1±0.9 | 721±403 | 27.9±17.8 | (Gaur et al., 2014) (Ram et al., 2010) |
| Mohali, India | Semi-urban | May, 2012 | 104±80.3 | - | 566.7±239.2 | 57.8±25.4 | (Sinha et al., 2014) |
| Mt. Abu, India | Remote | Jan 1993-Dec 2000, pre-monsoon | - | 0.7±0.14 | 131±36 | 39.9±10.8 | (Naja et al., 2003) (Das and Jayaraman, 2011) |

Table 3. Inter-comparison of observed and model simulated hourly average concentrations of air pollutants during the measurement campaign period. Unit: BC and PM in $\mu\text{g}/\text{m}^3$ and CO in ppbv.

| Pollutants | Observed (mean and range) | Modeled (mean and range) | Ratio of mean (observed/modeled) |
|-------------------------|----------------------------------|---------------------------------|---|
| BC | 4.9 (0.3-29.9) | 1.8 (0.4-3.7) | 2.7 |
| PM₁ | 36.6 (3.6-197.6) | 12.3 (0.9-41.7) | 3 |
| PM_{2.5} | 53.1 (6.1-272.2) | 17.3 (1.9-48.3) | 3 |
| PM₁₀ | 128.8 (10.5-604.0) | 25.4 (2.1-68.8) | 5 |
| CO | 344.1(124.9-1429.7) | 255.7 (72.2-613.1) | 1.35 |

Figures




Figure 1. Monthly synoptic wind (at 1000 hPa) for April, May and June 2013, based on NCEP/NCAR reanalysis data where the orientations of arrows refer to wind direction and the length of arrows represents the magnitude of wind (m/s). Red square box in the figure (left) represents the location of Lumbini. Figures on the right side represent monthly aerosol optical depth acquired with the MODIS instrument aboard TERRA satellite. High aerosol loading can be seen over the entire Indo-Gangetic Plains (IGP). Light gray color used in the figure represents the absence of data.




Figure 2. Time series of hourly average observed (red) and model estimated (blue) meteorological parameters at Lumbini, Nepal for the entire sampling period from 1 April to 15 June 2013.





Figure 3. Time series of the observed (red line) and model estimated (blue line) hourly average concentrations of BC, PM₁, PM_{2.5}, PM₁₀, O₃ and CO at Lumbini, Nepal for the entire sampling period from 1 April to 15 June 2013.



985

986 **Figure 4.** Comparison of BC concentrations to CO concentrations ($\Delta BC / \Delta CO$) ratios obtained
 987 for Lumbini with other sites. The red horizontal bar represents standard deviation.

988




Figure 5. Diurnal variations of hourly average ambient concentrations of BC, PM₁, PM_{2.5}, PM₁₀, O₃ and CO at Lumbini during the monitoring period (1 April -15 June 2013). In each box, lower and upper boundary of the box represents 25th and 75th percentile respectively, top and bottom of the whisker represents 90th and 10th percentile respectively, the mid-line represents median, and the square mark represents the mean for each hour.




Figure 6. Daily time series of PBL height obtained from the model and reported values over Lumbini (obtained from Xin et al., 2017). The lower panel shows the monthly average diurnal variation of the PBL height. The square mark in each box represents the mean PBL height, bottom and top of the box represents 25th and 75th percentile, top and bottom of the whisker represents 90th and 10th percentile respectively.




Figure 7. Time series of daily average $\Delta BC/\Delta CO$ ratio, absorption Ångström exponent (AAE), along with fire counts acquired with the MODIS instrument onboard TERRA satellite for a 200×200 km grid centered at Lumbini. Two rectangular green boxes represent time of two episodes with high peaks in CO and BC concentrations as shown in earlier figures.




Figure 8. Active fire hotspots in the region acquired with the MODIS instrument on Aqua satellite during (A) Event-I (7-9 April) and (B) Event-II (3-4 May). CO emissions, acquired with AIRS satellite, in the region two days before (3-5 April), during (7-9 April) and two days after (10-12 April) the Event-I are shown in panels (C), (E) and (G), respectively while panels (D), (F) and (H) show CO emissions two days before (1-2 May), during (3-4 May) and two days after (5-6 May) the Event-II. Panels (I) and (J) represent the 6-hr interval HYSPLIT back trajectories during Event I and II, respectively. Location of the Lumbini site is indicated by the red star in the panel (I and J). Observed CO versus Model open burning CO illustrating the contribution of forest fires during peak CO loading is shown in panel (K).




Figure 9. NO₂ total column obtained with OMI satellite over the region (a) before, (b) during, and (c) after the Event- I. The panels (d), (e), (f) show NO₂ total column before, during and after the Event- II.




Figure 10. (A) WRF-STEM model estimated contributions of various source regions to average CO concentration in Lumbini for the sampling period, (B) time series of region tagged CO tracer during the whole measurement period using HTAP emission inventory and (C) Figure showing percentage increase/decrease in CO concentration with different emissions scenario.




Figure 11. Comparison of normalized spectral light absorption coefficients obtained during the prime cooking (07:00-08:00 local time) and non cooking time (16:00-17:00 LT) at Lumbini with published data from Kirchstetter et al. (2004).




Figure 12. Diurnal variation of the fractional contribution of biomass burning to ambient BC concentration at Lumbini for the measurement period. In each box, lower and upper boundary of the box represent 25th and 75th percentile, respectively, top and bottom of the whisker represents 90th and 10th percentile, respectively. The mid-line in each box represents median while the square mark represents the mean for each hour.