

Heterogeneous ice nucleation on dust particles sourced from 9 deserts worldwide - Part 1: Immersion freezing

Yvonne Boose¹, André Welti^{1,2}, James Atkinson¹, Fabiola Ramelli¹, Anja Danielczok³, Heinz G. Bingemer³, Michael Plötze⁴, Berko Sierau¹, Zamin A. Kanji¹, and Ulrike Lohmann¹

¹Institute for Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland

²now at Leibniz Institute for Tropospheric Research, Leipzig, Germany

³Institute for Atmospheric and Environmental Sciences, J. W. Goethe-University, Frankfurt am Main, Germany

⁴Institute for Geotechnical Engineering, ETH Zürich, Zürich, Switzerland

Correspondence to: Y.Boose (yvonne.boose@env.ethz.ch) and Z. A. Kanji (zamin.kanji@env.ethz.ch)

Table 1. GPS coordinates of sample collection sites.

Collection site	type	latitude	longitude
Atacama	sieved	-22.03	-67.88
Atacama	milled	-22.03	-67.88
Australia	milled	-25.32	131.63
Crete	airborne	35.34	25.67
Dubai	sieved	24.83	55.66
Egypt	airborne	28.93	33.21
Great Basin	sieved	37.16	-116.49
Peloponnese	airborne	37.97	23.78
Israel	sieved	30.84	34.79
Israel	milled	30.84	34.79
Mojave	sieved	36.71	-117.17
Morocco	milled	31.21	-3.99
Namib	sieved	18.01	16.01
Tenerife	airborne	28.31	-16.50