

Cleaning up the air: Effectiveness of air quality policy for SO₂ and NO_x emissions in China

Ronald J. van der A^{1,2}, Bas Mijling¹, Jieying Ding^{1,3}, Maria Elissavet Koukouli⁴, Fei Liu¹, Qing Li⁵, Huiqin Mao⁵, Nicolas Theys⁶

5 ¹Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

²Nanjing University of Information Science and Technology, Nanjing, P.R.China

³Delft University of Technology, Delft, the Netherlands

⁴Laboratory of Atmospheric Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece

⁵Satellite Environment Center, Ministry of Environmental Protection, Beijing, P.R.China

10 ⁶Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

Correspondence to: Ronald J. van der A (avander@knmi.nl)

Abstract. Air quality observations by satellite instruments are global and have a regular temporal resolution, which make them very useful in studying long-term trends in atmospheric species. To monitor air quality trends in China for the period 15 2005-2015 we derive SO₂ columns and NO_x emissions on a provincial level with improved accuracy. To put these trends into perspective they are compared with public data on energy consumption and the environmental policies of China. We distinguish the effect of air quality regulations from economic growth by comparing them relatively to fossil fuel consumption. Pollutant levels, per unit of fossil fuel, are used to assess the effectiveness of air quality regulations. We note that the desulphurisation regulations enforced in 2005-2006 only had a significant effect in the years 2008-2009 when a 20 much stricter control of the actual use of the installations began. For national NO_x emissions a distinct decreasing trend is only visible since 2012, but the emission peak year differs from province to province. Unlike SO₂, emissions of NO_x are highly related to traffic. Furthermore, regulations for NO_x emissions are partly decided on a provincial level. The last three years show both a reduction in SO₂ and NO_x emissions per fossil fuel unit, since the authorities have implemented several new environmental regulations. Despite an increasing fossil fuel consumption and a growing transport sector, the effects of 25 air quality policy in China are clearly visible. Without the air quality regulations the concentration of SO₂ would be about 2.5 times higher and the NO₂ concentrations would be at least 25% higher than they are today in China.

1. Introduction

Satellite instruments can monitor air quality from space by mapping e.g. aerosols and tropospheric ozone, but are especially useful for emission estimates in observing the relatively short-living gases nitrogen dioxide (NO₂) and sulphur dioxide (SO₂). 30 For these two trace gases improved data sets recently became available, enabling analysis of air quality time series on a national or provincial level with improved accuracy. Theys et al. (2015) presented a new data set of SO₂ column densities derived from the Ozone Monitoring Instrument (OMI) satellite instrument (Levelt et al., 2006). They conclude that the SO₂ concentrations derived from OMI agree on average within 12% with ground observations. This data set strongly improves on earlier SO₂ data sets from satellites, which motivated this study. For NO₂, instead of using concentration data, we assess 35 directly the emission data of nitrogen oxides (NO_x = NO₂ + NO) that was derived from satellite observations by Mijling and Van der A (2012) and removes the meteorological influences. The precision of the derived NO_x emissions per grid cell of 0.25 x 0.25 degree is estimated as 20% (Ding et al., 2016).

China is one of the biggest emitters of SO₂ and NO₂ into the atmosphere because its large economy depends heavily on fossil fuels as an energy source. China alone is responsible for about 30 % of the global total emissions of SO₂ into the atmosphere 40 (Klimont et al., 2013), while over 90% of the SO₂ emissions are caused by coal consumption in China (Chen and Xu, 2010).

Coal is mainly used by thermal power plants and energy-intensive industry (e.g. steel, cement and glass), and to a lesser extent by residential use. SO₂ is also released by the use of oil and natural gas, but the sulphur content in these fuel types is much lower. Of these sources power plants are responsible for about 30-40 % of all emissions and industry for another 50-60 % (He K. et al., 2012, ChinaFAQs project, 2012). According to the Multi-resolution Emission Inventory for China (MEIC) (http://www.meicmodel.org/) the source of SO₂ emissions in 2010 was 29.4 % from power plants, 57.7 % from industry and 11.7 % residential and 1.2 % from transport. Figure 1 shows the location of the 600 biggest thermal power plants on the map of China including a list of provinces mentioned in this study. At a global scale, volcanic activity is another important source of atmospheric SO₂. However, plumes of active volcanoes are seldom observed over China.

NO_x is released by more or less the same anthropogenic sources, i.e. the burning of coal or oil. The main difference with SO₂ is that traffic is a much more important source for NO_x. NO_x emission factors (i.e. emissions per fossil fuel unit) in the transport sector are generally much higher than emission factors in energy and industry, which makes traffic one of the major sources of NO_x in China. According the MEIC inventory, 25% of NO₂ in 2010 was released by traffic, 32% by power plants, 4% by residential sources and 39% by industry, with the cement industry being the biggest emitter in this sector.

To reduce SO₂ in China, the authorities have implemented several environmental regulations. The most important regulation was the desulphurization of coal-fired power plants in 2005/2006 (Xu, 2011). This was later followed in the 12th five-year plan (2011-2015) by stricter control on the implementation of the regulations, additional filtering efforts, switching to low-sulphur coal and gasoline, phasing out obsolete capacity in coal-using industry, phasing out of small-scale coal mining, and gradually using more oil, gas and renewable energies instead of coal since 2011. An overview of all regulations related to SO₂ is shown in Table 1, which includes the year of the beginning of the implementation.

The regulation of NO_x was started much later than for SO₂. Although the 12th five-year plan already mentioned the intention to reduce NO₂ by 10 % (target) (ChinaFAQs project, 2012), only from 2012 onward NO_x filtering systems were installed, mainly at power plants but also for heavy industry. These regulations for NO_x were announced in 2013 in the Air Pollution Prevention and Control Action Plan (CAAC, 2013) for the coming 5 years. Gradually China has implemented several new national emission standards for cars during the time period of our study (see Table 2). Some regulations for road vehicles (e.g. ban on older polluting cars) have been introduced in China on a provincial or even city level, rather than nationwide. Strong regulations have also been enforced during specific events like the Olympic Games in 2008, Shanghai World Expo in 2010, Nanjing Youth Olympic Games in 2014, APEC meeting in 2014, etc. but those regulations are mostly of a temporary nature as shown by e.g. Mijling et al. (2012) for the Olympic Games in 2008.

To study the efficiency of the environmental policies, we analysed satellite observations of SO₂ and tropospheric NO₂ of the last 11 years. SO₂ satellite observations over China have been studied earlier by Lee et al. (2010), Li et al (2011), He (2012), Yang et al. (2013), Fioletov et al. (2015), and Krotkov et al. (2015). Satellite observations are very useful for SO₂ trend studies, as recently McLinden et al. (2016) showed that bottom-up inventories are underestimating SO₂ emissions worldwide with about 0-10%. NO₂ satellite observations over China have been evaluated by e.g. Richter et al. (2005), van der A et al. (2006), Zhang et al. (2012), and Krotkov et al. (2015). Emission estimates of NO_x over China have been analysed by Stavrou et al. (2008), Kurokawa et al. (2009), and more recently by Mijling et al. (2013) and by Liu et al. (2016a). In these studies linear trends of the concentration of air pollutants are often used. Here, however, we will relate changes derived on a provincial level for China with the energy consumption and the environmental policies of the country. This gives insight in the efficiency of the applied air quality policies and regulations. The comparison of SO₂ trends with those of NO_x emissions enables us to distinguish environmental policies specifically applied on coal-based industry and power plants with general environmental measures and trends in traffic.

2 Observational data

2.1 Satellite observations of SO₂

SO₂ is observed in the UV spectral range of satellite observations of SCIAMACHY (on Envisat), GOME-2 (on METOP-A) and OMI (on EOS-AURA). SO₂ retrieval algorithms have been earlier developed for GOME-1 by Eisinger and Burrows (1998), for SCIAMACHY (Lee et al., 2008), GOME-2 and for OMI by Krotkov et al (2006). Recently a new retrieval algorithm has been developed (Theys et al., 2015) that improves the precision of the SO₂ data for OMI with a factor 2, allowing us to derive more accurate trends based on OMI. The retrieval method is based on a Differential Optical Absorption Spectroscopy (DOAS) scheme to determine the slant columns from measured spectra in the 312-326 nm spectral range, which are then background corrected and converted to vertical columns using an Air Mass Factor (AMF). The AMF is calculated with the radiative transfer model LIDORT (LInearized Discrete Ordinate Radiative Transfer model). More details about the retrieval procedure are described in Theys et al. (2015). Also the operational algorithm of NASA for SO₂ from OMI has recently been improved. This algorithm and the algorithm of Theys et al. (2015) have a very comparable performance as shown by Fioletov et al. (2016). For this study, the algorithm of Theys et al. (2015) has been applied on the observations of the OMI instrument (Levelt et al., 2006) for its whole mission from 2004 onwards

To improve the quality of the OMI SO₂ data we exclude observations with a cloud fraction of more than 50 percent or with a fitting chi-square higher than 1. The solar zenith angle is limited to 75° and the viewing angle to 50°. Since the OMI instrument is suffering from the so-called row anomaly since 2007 (KNMI, 2012), we filter the affected rows (24-49, 54-55) in the same way for all years in the time series.

As we focus on anthropogenic SO₂, the SO₂ data for 15 June - 9 July 2011 have been removed because of its contamination with volcanic SO₂ from the eruption of the Nabro volcano in Africa and the transport of its plume to China (Brenot et al., 2014).

As a first step in our study we have made monthly means for the whole data set by averaging and gridding the data to a resolution of 1/8° by 1/8°. The gridding algorithm takes into account the area of each satellite footprint overlapping the grid cell. The resulting data set is a time series of monthly means for the time period October 2004 to December 2015.

For comparison we also use the official ESA SCIAMACHY/Envisat SO₂ product version SGP 5.02 and the standard data product from the GOME-2/METOP-A version GDP 4.7, as developed within the EUMETSAT Satellite Application Facility for Atmospheric Composition and UV radiation, O3MSAF, project and distributed by <http://atmos.caf.dlr.de/gome2/>. The data of these instruments are noisier than the OMI datasets because of the lower spatial coverage, different fit window and the lower signal-to-noise ratio of the SCIAMACHY and GOME-2 instruments. Therefore, their quality-controlled monthly mean SO₂ data have been recalculated by spatially averaging for each grid cell the data from the eight surrounding neighbouring cells, hence creating a smoothed SO₂ field. For details on the methodology and findings refer to Koukouli et al. (2016).

2.2 NO_x emission estimates from satellite observations

For NO_x emission data we use the results of an update (version 4) of the DECSO (Daily Emission estimates Constrained by Satellite Observation) algorithm developed by Mijling and van der A (2012). DECSO calculates emissions by applying a Kalman filter for the inversion of satellite data and a regional Chemical Transport Model (CTM) for the forward model calculation. It takes transport from the source into account with a semi-Lagrangian approach. The CTM we use is CHIMERE v2013 (Menut et al., 2013) with meteorological information from the European Centre for Medium-range Weather Forecasts (ECMWF) with a horizontal resolution of approximately 25x25 km². The DECSO algorithm is applied to OMI NO₂ observations derived by the DOMINO v.2 algorithm (Boersma et al., 2011). The latest improvements of the DECSO

algorithm resulting in version 4 are described by Ding et al. (2015, 2016). The monthly average emission data over China we use is available on 0.25 degree resolution for the period 2007-2015 on the web-portal www.globemission.eu.

3 Temporal analysis over China

3.1 Sources of SO₂ and NO_x in China

5 The multi-annual mean of SO₂ for 2005-2015 is shown in Figure 2. As the lifetime of SO₂ is relatively short (typically 4-48 hours) (Lee et al., 2011, Fioletov et al., 2015), the observed SO₂ concentrations are a good proxy for the location of SO₂ emissions. Regions with large SO₂ concentrations are South Hebei, the province Shandong (around the city Zibo) and the region around Chongqing. South Hebei is a region with many power plants just east of the mountainous coal-mining area in Shanxi. The hot spot in the province Shandong is related to a strongly industrialized area with lots of coal-using industry. In 10 the Chongqing region both coal mines and heavy industry are located.

Rather than located at hot spots, high NO₂ concentrations are more distributed over the East of China, mainly because traffic is an important source of NO_x emissions (see Figure 3a). The underlying NO_x emissions are shown in Figure 3b. Like for the SO₂ concentrations, NO_x emission spots can be found at the location of big power plants. Also clearly visible are the megacities of China, and ship tracks along the coast and sources along the big rivers.

15 3.2 SO₂ trends over China

To construct time series of SO₂ we have averaged the data to annual means of the vertical columns derived from OMI. From these annual mean SO₂ data we constructed time series for each province (see Table A.1). Figure 4 shows the mean normalized time series for the 10 provinces with the highest total SO₂ column densities (i.e. Tianjin, Shandong, Hebei, Shanxi, Henan, Beijing, Jiangsu, Shanghai, Anhui and Liaoning), together responsible for 60% of all ambient SO₂ in China. 20 The individual time series are drawn as thin black lines. The minimum and maximum of these time series for each year are shown in the grey shaded area to indicate the variability. The time series of Shanghai is the lowest black line of the 10 series, thus the reductions have been strongest in this province since 2005. Apart from Ningxia province, all provincial time series show very similar patterns. In general, the SO₂ concentrations were at a maximum in the year 2007, when the start of decreasing trend is visible in China. Despite some fluctuations the SO₂ concentrations remain relatively constant from 2010 25 till 2013, where after they are decreasing again.

A different trend is observed for Ningxia, a province in the mid-north of the country with a relative low population density and large coal resources. Here an increasing trend emerges for the years starting from 2010 when several new coal power plants were put into operation. A list of largest power plants (with a capacity of more than 600 MW) and the start year of their operation is shown in Table 3. From 2012 onward, the more stringent SO₂ emission regulations also started to have 30 effect in Ningxia.

3.3 NO_x emission trends over China

National NO_x emission trends show a different pattern than those of SO₂. We observe an increasing trend till about 2012, with an exception of the year 2009 which is related to regulations started at the Olympic Games in 2008 (Mijling et al., 2008) and the global economic crisis which shortly slowed down the Chinese economic growth. Total NO_x emissions in East 35 China reached their peak levels in 2012, and have been decreasing since. While the economy kept growing after 2012, the emission of NO_x slowly decreases again as a result of the air quality regulations described in Section 1. According to the DECSO emission inversion, in 2015 the NO_x emissions were 4.9 Tg N/yr, which is 22.8% lower than in the peak year 2012. However, the 2015 emissions were still 14.1% higher than in the reference year 2007. The trends per province (see Table A.2) show very similar patterns with only the starting year (the year with maximum NO_x emissions) of the decrease in

emissions varying over the provinces. In Figure 5a, the normalized (to the year 2007) time series of annual NO_x emissions for East China (102-132°E, 18-50°N) is shown in similar way as for SO_2 in Figure 4. The mean, minimum and maximum of the 10 provinces with highest NO_x emission are shown (Shandong, Hebei, Henan, Jiangsu, Guangdong, Shanxi, Zhejiang, Anhui, Sichuan and Hubei), together responsible for 65% of all Chinese NO_x emissions. The thin black lines show the time series for the individual 10 provinces, where the lower line represents Guangdong. Figure 5b shows the peak year for each province. Provinces where air pollution regulations, for e.g. traffic, got a lot of attention at an early stage, like Beijing and Shanghai, have reached their maximum before 2011. Most industrialised regions show their peak in the years 2011-2013. Some of the less developed and populated provinces show a maximum in 2014, which means that their decrease in NO_x emissions is very recent. Regional variations are mainly due to the fact that regulations for the NO_x emission reductions, for instance in traffic or power plants, are determined and implemented on a provincial level (Liu et al., 2016b). For the province Ningxia we see a very similar pattern occurring as for SO_2 , which shows that for this low-densely populated province traffic plays a small role and the trend is determined by the operation of newly-built power plants.

3.4 Air pollution in relation to fossil fuel consumption

To relate the observed SO_2 and NO_x reduction to environmental regulations we have to take into account the coal and oil consumption in the same time period. The total coal consumption in Standard Coal Equivalent (SCE) units per year for China and the total oil consumption (also in SCE units) are shown in Figure 6, based on data of NBSC (2015). According to Guan et al. (2012) and Hong et al. (2016) the sum of the coal consumption of all provinces is more accurate than the number provided for the whole of China, thus we use the provincial totals for coal consumption. For NO_x emissions the transport sector plays an important role, especially ships are one of the largest NO_x emitters per fuel unit in the transport sector. The total freight transport almost doubles every 6 years in China.

Since the burning of coal and oil are the dominant sources of SO_2 and NO_x emissions, we can consider the total emissions of these air pollutants as the product of the national use of coal and oil (activity) and the average emission factor of one unit coal/oil. The effectiveness of environmental regulation will be reflected in a decrease of this emission factor. Therefore, we divide the annual SO_2 column measured from satellites and the annual NO_x emissions by the annual coal and oil consumption in China. In this way we get a measure of the emitted SO_2 or NO_x per unit (SCE) of fossil fuel consumption reflecting the Chinese environmental policy. The results are shown in Figure 7. One might argue that SO_2 is more related to coal than oil, but division by only coal yields the same results. In our analysis we omit gas consumption since this is very limited in China and hence does not affect the results significantly.

We focus here mainly on the results for OMI, because of the instrument's high spatial resolution and lack of instrumental degradation. However, SO_2 data of the SCIAMACHY and GOME-2 instrument are also added in Figure 7 to be able to further look into the past (starting in 2003) and to verify the results of OMI. The SO_2 data of SCIAMACHY and GOME-2 are averaged over the summer months (April-September). The remaining monthly means are excluded from the analysis due to a lower accuracy at higher latitudes and a large part of the higher latitudes is missing due to snow cover. For OMI each data point is averaged over 12 months and the total area of China, which reduces the root-mean-square error to a negligible level. Biases among all instruments are removed by normalizing the values to those in reference year 2007. Up to 2009, the results agree fairly well. After 2009, we see the results of GOME-2 and OMI for SO_2 slowly diverge in time, which might be result of the instrument degradation of the UV spectra of GOME-2 after 2009 (Munro et al., 2016).

Changing weather conditions from year-to-year can affect the results for SO_2 concentrations and when these weather conditions are different during the overpass of SCIAMACHY and GOME-2 (around 9:30 local time) and overpass of OMI (around 13:30 local time), this can lead to differences between the instruments. The global coverage of SCIAMACHY is once every 6 days and for GOME-2 and OMI almost daily. The limited number of samples for SCIAMACHY makes this

data more sensitive to weather conditions. Note that due to the nature of the inversion algorithm the NO_x emission data is in general not sensitive to meteorological variability.

For SO_2 we see a big decrease in the years 2008 and 2009, while the desulphurization program of the 11th five-year plan started already in 2005/2006, when the authorities begin to reduce SO_2 emissions by installing desulphurization devices in many power plants (Lu et al., 2010). In 2006 SO_2 monitoring devices were also installed in the chimneys of the power plants. This resulted in a decrease in SO_2 emissions from 2006, while the much bigger decrease of SO_2 in 2008-2009 reflects the stronger government control at that time on the actual use of the equipment (Xu et al., 2011). After 2009, the SO_2 content per consumed coal unit only slowly decrease until 2011. From 2012 onwards we see a stronger annual decrease in SO_2 . This coincides with the 12th five-year program when new measures were taken to upgrade the coal quality, to modernize the industry and to put more effort on law enforcement. Especially the law enforcement in the last years concerning the prohibition of flue gas bypass and the use of desulphurization devices in the steel industry played an important role.

For the NO_x emissions the total annual emissions are used and divided in the same way as for SO_2 by the total coal and oil consumption. Here, however we should keep in mind that the transport sector (especially by shipping) emits much more NO_x per fuel unit than the power and industrial sectors (see e.g. Zhao et al., 2013). Thus the percentage of the total fuel used by transport is relevant for the graph of NO_x . In the early years we see in general a small increase in NO_x emissions per fuel unit due to the increasing fraction of the transport sector in the fuel use. Exceptions are the year 2009 and the recent year 2015. The year 2009 coincides with the global economic crisis when there was less export of goods from China. This affected especially the transport sector, mostly transport over water, as shown in De Ruyter de Wildt et al. (2012). Faber et al. (2012) and Boersma et al. (2015) showed that the economic crisis also resulted in a significant reduction of the average vessel speed to save fuel used by ship transport. This caused not only a shift in source sectors but in general led to lower NO_x per fuel values. This explains the dip in pollution per fuel unit in 2009. After 2012 the gradual increase of NO_x per fuel (as a result of the strongly growing transport sector) slowly stops, and the year 2015 shows a sharp decline in NO_x per fossil fuels unit. This can be directly related to the rapidly growing installation of Selective Catalytic Reduction (SCR) equipment at power plants since 2012 and the introduction of new emission standards for cars, as shown by Liu et al. (2016). This strong reduction in NO_x for 2015 and the equally strong reduction for SO_2 in 2014 and 2015 are a result of very effective recent environmental regulations in the last years in China. By comparing the efficiency level in 2015 with earlier levels we can conclude from Figure 7 that without these air quality regulations SO_2 concentrations would nowadays be about 2.5 times higher and the NO_2 concentrations would be at least 25% higher in China today.

4 Discussion

The current developments in data products derived from satellite observations provide high quality time series of the air pollutants NO_x and SO_2 . Although the mean of observed SO_2 columns are not linearly related to the SO_2 emissions because of the influence of the weather, it can still be argued that these satellite data products, whether concentrations or emissions, provide a fair comparison over the various regions from year to year. By comparing these time series with fossil fuel energy consumption the economic growth is removed from the equation and we can monitor the effectiveness of air quality policies. We foresee that this method will become a valuable tool for policy makers concerning air quality regulations.

For China we see similar patterns in the trends of SO_2 per province. In 2006 a nation-wide implementation of desulphurisation installations started. However, the effects are only visible in 2008 and 2009 when a strict control by the Chinese authorities on the actual use of the desulphurisation installations started. In 2009, we see the effect of the air quality regulations for SO_2 and NO_x resulting from the global economic recession at the end of 2008. The increasing relative contribution of the transport sector to the NO_x emission slowly increases the amount of NO_x per fossil fuel unit after 2009. After 2011 we see a steadily decreasing SO_2 pollution per fossil fuel unit caused by various Chinese environmental regulations. In the last year of our time series, 2015, a clear effect becomes visible of very recent regulations for NO_x

emissions from power plants and heavy industry. The fit of linear trends often used in earlier studies is therefore no longer applicable to the Chinese situation.

The availability of high quality satellite data for the last ten years is especially interesting for China where the situation is rapidly changing. For instance in Europe and Japan desulphurisation started much earlier when these satellite data were not yet available. On the other hand, in India SO₂ and NO_x emissions are still growing and possible new regulations can be monitored in the years to come with an even better quality using forthcoming sensors as e.g. TROPOMI on-board Sentinel-5 Precursor.

Despite the growing use of coal and oil in the last ten years in China we see reduced emissions per fuel unit in the past few years. This decreasing trend in both SO₂ and NO_x for China is likely to continue in the coming years for which the Chinese national government has announced less use of coal, more environmental regulations for SO₂ and NO_x and stricter reinforcement of control of environmental policies.

Acknowledgements

This research has been funded by the MarcoPolo project of the European Union Seventh Framework Programme (FP7/2007-2013) under Grant Agreement n° 606953 and by the GlobEmission project (Contract No 4000104001/11/I-NB) of the Data User Element programme of the European Space Agency.

5 References

- Boersma, K.F., H.J. Eskes, R. J. Dirksen, R. J. van der A, J. P. Veefkind, P. Stammes, V. Huijnen, Q. L. Kleipool, M. Sneep, J. Claas, J. Leitao, A. Richter, Y. Zhou, and D. Brunner, An improved retrieval of tropospheric NO₂ columns from the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905-1928, 2011
- Boersma, K.F., G.C.M. Vinken, J. Tournadre, Ships going slow in reducing their NO_x emissions: changes in 2005–2012 ship exhaust inferred from satellite measurements over Europe, *Environ. Res. Letters*, 10, 7, doi:10.1088/1748-9326/10/7/074007, 2015
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: mission objectives and measurement modes, *J. Atmos. Sci.*, 56, 127–150, doi:10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2, 1999.
- Brenot, H., N. Theys, L. Clarisse, J. van Geffen, J. van Gent, M. Van Roozendael, R.J. van der A, D. Hurtmans, P.F. Coheur, C. Clerbaux, P. Valks, P. Hedelt, F. Prata, O. Rasson and C. Zehner, Support to Aviation Control Service (SACS): an online service for near-real-time satellite monitoring of volcanic plumes, *Natural Hazards and Earth System Sciences*, 2014, 14, 1099-1123, doi:10.5194/nhess-14-1099-2014.
- CAAC 2013, Clean Air Alliance of China, State Council air pollution prevention and control action plan, issue II, October 2013, <http://en.cleairchina.org/product/6346.html> (English translation). Last accessed: 8 October 2015
- ChinaFAQs Project, <http://www.chinafaqs.org/library/chinafaqs-china-adopts-world-class-pollutant-emissions-standards-coal-power-plants>, 2012. Last accessed: 8 October 2015
- De Ruyter de Wildt, M, H. Eskes, K. F. Boersma, The global economic cycle and satellite-derived NO₂ trends over shipping lanes, *Geophys. Res. Letters*, doi:10.1029/2011GL049541, 2012
- Ding, J., van der A, R. J., Mijling, B., Levelt, P. F., and Hao, N.: NO_x emission estimates during the 2014 Youth Olympic Games in Nanjing, *Atmos. Chem. Phys.*, 15, 9399-9412, doi:10.5194/acp-15-9399-2015, 2015.
- Ding, J., van der A, R. J., Mijling, B., Levelt, P. F., New NO_x emission inversion algorithm improves estimates over low-emission regions., *J. Geophys. Res.*, submitted, 2016

- Eisinger, M., and J. P. Burrows (1998), Tropospheric sulfur dioxide observed by the ERS-2 GOME instrument, *Geophys. Res. Lett.*, 25, 4177–4180.
- Faber J, Nelissen D, Hon G, Wang H and Tsimplis M, Regulated slow steaming in maritime transport—an assessment of options, costs and benefits CE Delft (The Netherlands: Delft)
- 5 (www.cedelft.eu/publicatie/regulated_slow_steaming_in_maritime_transport/1224) p 117 , 2012
- Fioletov, V. E., C. A. McLinden, N. Krotkov, and C. Li (2015), Lifetimes and emissions of SO₂ from point sources estimated from OMI. *Geophys. Res. Lett.*, 42, 1969–1976. doi: 10.1002/2015GL063148.
- Fioletov, V. E., McLinden, C. A., Krotkov, N., Li, C., Joiner, J., Theys, N., Carn, S., and Moran, M. D.: A global catalogue of large SO₂ sources and emissions derived from the Ozone Monitoring Instrument, *Atmos. Chem. Phys. Discuss.*,
- 10 doi:10.5194/acp-2016-417, in review, 2016.
- Guan, D., Z. Liu, Y. Geng, S. Lindner and K. Hubacek, The gigatonne gap in China's carbon dioxide inventories, *Nature Climate Change*, 2, 672–675, (2012), doi:10.1038/nclimate1560
- Hong, C., Zhang, Q., He, K., Guan, D., Li, M., Liu, F., and Zheng, B.: Variations of China's emission estimates response to uncertainties in energy statistics, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-459, in review, 2016.
- 15 He, Kebin: Multi-resolution Emission Inventory for China (MEIC): model framework and 1990-2010 anthropogenic emissions, in International Global Atmospheric Chemistry Conference, 17-21 September, Beijing, China. [online] Available from: <http://adsabs.harvard.edu/abs/2012AGUFM.A32B..05H> (Accessed 4 February 2015), 2012.
- He, H., et al. (2012), SO₂ over central China: Measurements, numerical simulations and the tropospheric sulfur budget, *J. Geophys. Res.*, 117, D00K37, doi:10.1029/2011JD016473.
- 20 Klimont, Z., Smith, S. J., and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions, *Environ. Res. Lett.*, 8, 014003, doi:10.1088/1748-9326/8/1/014003, 2013.
- KNMI: Background information about the Row Anomaly in OMI, Retrieved from: <http://www.knmi.nl/omi/research/product/rowanomaly-background.php>, 2012, Last accessed: 8 October 2015.
- Koukouli, M.E., Balis, D.S., van der A, R., Theys, N., Hedelt, P., Lichtenberg, G., Richter, A., Krotkov, N., Li, C., Taylor,
- 25 M., Characteristics of the anthropogenic SO₂ load over China observed by SCIAMACHY/Envisat, OMI/Aura and GOME2/MetopA, paper in preparation for Atmospheric Environment, 2016.
- Krotkov, N.A., S.A. Carn, A.J. Krueger, P.K. Bhartia, and K. Yang, Band residual difference algorithm for retrieval of SO₂ from the aura ozone monitoring instrument (OMI), *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 44, No. 5, 2006.
- 30 Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z., and Streets, D. G.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2014, *Atmos. Chem. Phys. Discuss.*, 15, 26555-26607, doi:10.5194/acpd-15-26555-2015, 2015.
- Kurokawa, J. I., Yumimoto, K., Uno, I., & Ohara, T. (2009). Adjoint inverse modeling of NO_x emissions over eastern China using satellite observations of NO₂ vertical column densities. *Atmospheric Environment*, 43(11), 1878-1887.
- 35 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Greet, J.-M., Fukui, T., Kawashima, K. and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058.
- Lee, C., Richter, A., Weber, M., and Burrows, J. P.: SO₂ Retrieval from SCIAMACHY using the Weighting Function DOAS (WFDOAS) technique: comparison with Standard DOAS retrieval, *Atmos. Chem. Phys.*, 8, 6137-6145, doi:10.5194/acp-8-6137-2008, 2008.
- 40

- Lee, C., R. V. Martin, A. van Donkelaar, H. Lee, R. R. Dickerson, J. C. Hains, N. Krotkov, A. Richter, K. Vinnikov, and J. J. Schwab (2011), SO₂ emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations, *J. Geophys. Res.*, 116, D06304, doi:10.1029/2010JD014758.
- Levelt, P. F., G. H. J. van den Oord, M. R. Dobber, A. Mälkki, H. Visser, J. de Vries, P. Stammes, J. Lundell, and H. Saari (2006), The ozone monitoring instrument, *IEEE Trans. Geo. Rem. Sens.*, 44(5), 1093–1101, doi:10.1109/TGRS.2006.872333.
- Li, C., Zhang, Q., Krotkov, N. A., Streets, D. G., He, K., Tsay, S.-C., and Gleason, J. F.: Recent large reduction in sulfur dioxide emissions from Chinese power plants observed by the Ozone Monitoring Instrument, *Geophys. Res. Lett.*, 37, 1–6, doi:10.1029/2010GL042594, 2010.
- 10 Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., and He, K. B.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, *Atmos. Chem. Phys.*, 15, 13299–13317, doi:10.5194/acp-15-13299-2015, 2015.
- Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K., and Wagner, T.: NO_x lifetimes and emissions of cities and power plants in polluted background estimated by satellite observations, *Atmos. Chem. Phys.*, 16, 5283–5298, doi:10.5194/acp-16-5283-2016, 2016a.
- 15 Liu, F., Q. Zhang, R. J. van der A, B. Zheng, D. Tong, L. Yan and K. B. He, Recent reduction in NO_x emissions over China from satellite observations and emission inventories, submitted to *Env. Res. Letters*, 2016b.
- Lu, Z., Streets, D. G., Zhang, Q., Wang, S., Carmichael, G. R., Cheng, Y. F., Wei, C., Chin, M., Diehl, T., and Tan, Q.: Sulphur dioxide emissions in China and sulphur trends in East Asia since 2000, *Atmos. Chem. Phys.*, 10, 6311–6331, doi:10.5194/acp-10-6311-2010, 2010.
- 20 McLinden, C. A., V. Fioletov, M. W. Shephard, N. Krotkov, C. Li, R. V. Martin, M. D. Moran, and J. Joiner, Space-based detection of missing sulfur dioxide sources of global air pollution, *Nature Geoscience*, DOI: 10.1038/ngeo2724, 2016
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G., Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R. and Vivanco, M. G.: CHIMERE 2013: a model for regional atmospheric composition modelling, *Geosci. Model Dev.*, 6(4), 981–1028, doi:10.5194/gmd-6-981-2013, 2013
- MEP, Ministry of Environmental Protection, http://english.mep.gov.cn/Plans_Reports/, last access: 8 October 2015.
- Mijling, B. and R.J. van der A, Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale, *J. Geophys. Res.*, 117, 2012, doi:10.1029/2012JD017817.
- 30 Mijling, B., R.J. van der A and Q. Zhang, Regional nitrogen oxides emission trends in East Asia observed from space, *Atm. Chem. Phys.*, 2013, 13, 12003–12012, doi:10.5194/acp-13-12003-2013.
- Munro, R., Lang, R., Klaes, D., Poli, G., Retscher, C., Lindstrot, R., Huckle, R., Lacan, A., Grzegorski, M., Holdak, A., Kokhanovsky, A., Livschitz, J., and Eisinger, M.: The GOME-2 instrument on the Metop series of satellites: instrument design, calibration, and level 1 data processing – an overview, *Atmos. Meas. Tech.*, 9, 1279–1301, doi:10.5194/amt-9-1279-2016, 2016.
- 35 NBSC, National Bureau of Statistics China, <http://data.stats.gov.cn>, last accessed: 29 December 2015
- NDRC, 2013, National Development and Reform Commission (reported on 16 August 2013), http://www.ndrc.gov.cn/zcfb/zcfbtz/201308/t20130827_555124.html (last accessed on 01-11-2015)
- Richter, A., J. P. Burrows, H. Nuß, C. Granier, and U. Niemeier (2005), Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129 – 132, doi:10.1038/nature04092.
- 40 State Council, 2014, State Council announcement on 15-5-2014 of environmental regulations), http://www.gov.cn/zhengce/content/2014-05/26/content_8824.htm (last accessed on 01-11-2015)

- Stavrakou, T., Müller, J. F., Boersma, K. F., De Smedt, I., & van der A, R. J. (2008). Assessing the distribution and growth rates of NO_x emission sources by inverting a 10-year record of NO₂ satellite columns. *Geophysical Research Letters*, 35(10).
- They, N., De Smedt, I., van Gent, J., Danckaert, T., Wang, T., Hendrick, F., Stavrakou, T., Bauduin, S., Clarisse, L., Li, C., Krotkov, N., Yu, H., Brenot, H., and Van Roozendaal, M.: Sulfur dioxide vertical column DOAS retrievals from the Ozone Monitoring Instrument: global observations and comparison to ground-based and satellite data, *J. Geophys. Res.-Atmos.*, 120, 2470–2491, doi:10.1002/2014JD022657, 2015.
- van der A, R. J., Peters, D. H. M. U., Eskes, H., Boersma, K. F., Van Roozendaal, M., De Smedt, I., and Kelder, H. M.: Detection of the trend and seasonal variation in tropospheric NO₂ over China, *J. Geophys. Res.*, 111, D12317, doi:10.1029/2005JD006594.
- Xu, Y., Improvements in the Operation of SO₂ Scrubbers in China's Coal Power Plants, *Environ. Sci. Technol.*, 2011, 45 (2), pp 380–385, DOI: 10.1021/es1025678.
- Yang, K., R. R. Dickerson, S. A. Carn, C. Ge, and J. Wang (2013), First observations of SO₂ from the satellite Suomi NPP OMPS: Widespread air pollution events over China, *Geophys. Res. Lett.*, 40, 4957–4962, doi:10.1002/grl.50952.
- Zhang, Q., Geng, G., Wang, S., Richter, A., & He, K. (2012). Satellite remote sensing of changes in NO_x emissions over China during 1996–2010. *Chinese Science Bulletin*, 57(22), 2857-2864.
- Zhang R. (2013, August 11): Heating causes pollution spike in Northern China, *CCTV.com*, Retrieved from <http://english.cntv.cn/program/china24/20131108/101015.shtml>
- Zhu Ningzhu (2013, October 12): China to shut coal mines in safety overhaul, *Xinhua News Agency*, Retrieved from http://news.xinhuanet.com/english/china/2013-10/12/c_132793667.htm
- Zhao, B., Wang, S. X., Liu, H., Xu, J. Y., Fu, K., Klimont, Z., Hao, J. M., He, K. B., Cofala, J., and Amann, M.: NO_x emissions in China: historical trends and future perspectives, *Atmos. Chem. Phys.*, 13, 9869-9897, doi:10.5194/acp-13-9869-2013, 2013.



Figure 1: Location of power plants in China according to REAS v.2 (Kurokawa et al., 2013). The size of each dot indicates the emission of the power plants (Power plants in close proximity are combined in a single dot). In addition, a list is given of the provinces mentioned in this study.





Figure 2: Average SO₂ concentrations for the period 2005 to 2015 as observed by the OMI satellite instrument. Data below 0.1 DU is masked (grey colour).



5

(a)

(b)

Figure 3: (a) The averaged tropospheric NO₂ concentrations over China measured by OMI in the period 2005-2014. (b) The NO_x emissions in the year 2010 derived from the OMI satellite observations.





Figure 4: Time series (red line) of the annual mean of the 10 provinces with the highest SO_2 concentrations derived from the OMI satellite observations. The time series are normalized to their value in 2005. The grey area indicates the maximum range of the individual values of the times series of each of the 10 provinces. The thin black lines show the individual time series of those provinces. The province of Ningxia has a distinct deviating trend, here shown in blue.

5



(a)

(b)

Figure 5: (a) Shown are the annual total NO_x emission estimates for the last 9 years for the top 10 of highest NO_x emitting provinces in East China. Emissions are derived with DECSO V4 using OMI observations. The thin black lines show the individual time series of those provinces. (b) Peak year of the NO_x emissions per province.

10




Figure 6: In black the annual coal consumption and in red the annual oil consumption for China is shown.




Figure 7: Time series of the ratio of the mean SO₂ columns and the fossil fuel consumption in China based on observations of OMI (black), SCIAMACHY (green), and GOME-2 (blue). The ratios of the annual NO_x emissions and the fossil fuel consumption is based on observations of OMI (red). All time series are normalized to the year 2007.

Table 1: Environmental regulations of the Chinese national government to reduce SO₂ in the air.

Start year of implementation	Regulation	Reference
2005-2006	Desulphurization techniques in power plants.	Li et al., 2010
2005-2012	Closure of several of the most polluting power plants	Liu et al., 2015
2008	Stricter control of implementation of desulphurization in power plants	Xu et al., 2011 Liu et al., 2015
2011	Use of more gas and renewable energies instead of coal	NBSC, 2015
January 2012	New emission standard of air pollutants for thermal power plants	MEP, 2015
2013	Mandatory SO ₂ filtering of small-scale coal-fired industry	Zhang, 2013, NDRC, 2013
End of 2013	Stricter control of environmental policy	CAAC, 2013, State Council, 2014
End of 2013	Further desulphurization in industry	CAAC, 2013, NDRC, 2013
2014	Phasing out small-scale coal-fires boilers	CAAC, 2013, State Council, 2014
2014	Closure of 2000 small-scale coal mines	Zhu, 2013
End of 2014	Use of low-sulphur coal	State Council, 2014
End of 2014	Cap on coal consumption	State Council, 2014

Table 2: Environmental regulations of the Chinese national government to reduce NO_x emissions.

Year of implementation	Regulation	Reference
2011-2015	Installation of Selective Catalytic Reduction (SCR) equipment at power plants	Liu et al., 2016, CAAC, 2013
2007	China 3 (Euro 3) emissions standards for cars, nationwide	www.dieselnat.com
2011	China 4 (Euro 4) emissions standards for gasoline cars, nationwide.	www.dieselnat.com
2015	China 4 (Euro 4) emissions standards for diesel cars, nationwide.	www.dieselnat.com

Table 3: Main power plants in Ningxia province (> 600 MW). Data collected from www.sourcewatch.org.

Power plant	Capacity (MW)	In operation since	Remark
CPI Linhezhen	700	unknown	
Daba-1	1200	< 2000	
Daba-2	1100	unknown	An extension of Daba-1
Ningxia Zhongning-2	660	2005-2006	
Guodian Shizuishan-2	1980	2006	
Ningdong Maliantai	660	2006	
Huadian Ningxia Lingwu units 1&2	1200	2007	
Guodian-Dawukou	1100	2010	Extension of the original 440 MW plant
Guohua Ningdong	660	2010	
Ningxia Liupanshan	660	2010	
Huadian Ningxia Lingwu units 3&4	2120	2010-2011	
Shenhua Yuanyang Lake	1320	2010-2011	
Shuidonggou	1200	2011	
Ningdong Younglight	660	2013	

Appendix A

Table A1 Annual SO₂ column densities (DU/grid cell) per province observed by OMI

Province	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Anhui	0.625	0.586	0.953	0.637	0.499	0.567	0.671	0.553	0.602	0.379	0.280
Beijing	0.753	0.829	0.989	0.711	0.778	0.749	0.850	0.673	0.634	0.640	0.491
Chongqing	0.514	0.509	0.530	0.567	0.580	0.580	0.492	0.370	0.469	0.269	0.136
Fujian	0.099	0.123	0.196	0.135	0.104	0.113	0.112	0.080	0.107	0.076	0.064
Gansu	0.144	0.136	0.150	0.130	0.135	0.123	0.134	0.127	0.131	0.105	0.103
Guangdong	0.251	0.257	0.280	0.239	0.171	0.177	0.138	0.095	0.118	0.086	0.080
Guangxi	0.199	0.203	0.270	0.236	0.127	0.190	0.179	0.092	0.134	0.091	0.072
Guizhou	0.424	0.478	0.532	0.516	0.418	0.424	0.357	0.261	0.345	0.167	0.100
Hainan	0.098	0.086	0.091	0.092	0.060	0.090	0.106	0.027	0.087	0.055	0.000
Hebei	0.903	0.931	0.996	0.908	0.874	0.881	0.922	0.863	0.844	0.716	0.540
Heilongjiang	0.134	0.141	0.144	0.142	0.124	0.138	0.154	0.162	0.125	0.134	0.135
Henan	1.036	0.920	1.222	0.938	0.709	0.778	0.992	0.827	0.762	0.585	0.439
Hubei	0.487	0.477	0.603	0.490	0.342	0.386	0.479	0.365	0.378	0.288	0.176
Hunan	0.364	0.330	0.448	0.371	0.270	0.281	0.320	0.240	0.259	0.180	0.112
Jiangsu	0.847	0.782	1.054	0.917	0.678	0.716	0.871	0.687	0.735	0.524	0.326
Jiangxi	0.272	0.278	0.373	0.267	0.202	0.222	0.244	0.197	0.230	0.184	0.136
Jilin	0.205	0.233	0.260	0.259	0.191	0.207	0.201	0.187	0.203	0.192	0.156
Liaoning	0.512	0.576	0.602	0.568	0.504	0.478	0.475	0.479	0.515	0.480	0.321
NeiMongol	0.154	0.170	0.200	0.174	0.175	0.180	0.191	0.178	0.177	0.186	0.152
Ningxia	0.234	0.208	0.273	0.225	0.258	0.241	0.350	0.349	0.306	0.242	0.209
Qinghai	0.079	0.085	0.077	0.080	0.079	0.088	0.083	0.091	0.096	0.082	0.086
Shaanxi	0.357	0.301	0.401	0.324	0.261	0.269	0.338	0.304	0.315	0.246	0.224
Shandong	1.197	1.309	1.531	1.315	1.113	1.188	1.323	1.232	1.191	0.870	0.592
Shanghai	0.874	0.744	0.883	0.828	0.588	0.656	0.544	0.460	0.507	0.325	0.202
Shanxi	0.748	0.806	0.928	0.779	0.593	0.612	0.789	0.703	0.661	0.614	0.493
Sichuan	0.429	0.429	0.513	0.376	0.415	0.427	0.394	0.293	0.350	0.198	0.123
Taiwan	0.089	0.071	0.081	0.085	0.074	0.090	0.074	0.055	0.086	0.052	0.051
Tianjin	1.197	1.344	1.577	1.132	1.217	1.289	1.176	1.140	1.150	1.005	0.708
XinjiangU.	0.073	0.074	0.087	0.093	0.088	0.094	0.090	0.090	0.111	0.101	0.084
Xizang/Tibet	0.080	0.097	0.086	0.091	0.094	0.111	0.096	0.097	0.083	0.087	0.087
Yunnan	0.140	0.159	0.182	0.147	0.144	0.153	0.149	0.131	0.127	0.100	0.085
Zhejiang	0.383	0.337	0.452	0.395	0.297	0.316	0.403	0.258	0.321	0.212	0.158
P.R.China	0.397	0.392	0.444	0.373	0.330	0.342	0.358	0.335	0.332	0.280	0.225

Table A2 Annual NO_x emissions (Gg N/year) per province derived from OMI observations

Province	2007	2008	2009	2010	2011	2012	2013	2014	2015
Anhui	167	169	187	224	242	292	288	282	215
Beijing	91	62	90	107	88	80	89	74	64
Chongqing	54	57	70	75	87	95	96	100	70
Fujian	96	114	100	114	161	162	153	167	137
Gansu	31	38	37	42	61	73	60	78	52
Guangdong	383	383	331	341	371	413	360	374	331
Guangxi	118	148	118	145	152	224	224	200	157
Guizhou	107	130	142	154	131	194	191	180	122
Hainan	8	13	11	17	23	22	25	37	30
Hebei	427	423	403	515	563	543	544	511	436
Heilongjiang	33	36	30	25	43	54	40	49	31
Henan	334	347	370	445	470	481	491	433	315
Hubei	135	140	144	186	248	263	233	270	199
Hunan	109	112	124	162	163	216	184	218	187
Jiangsu	374	344	344	421	470	433	428	445	365
Jiangxi	51	58	65	73	85	105	111	150	112
Jilin	30	22	18	20	45	50	43	48	43
Liaoning	122	128	124	169	205	225	178	199	173
NeiMongol	98	116	117	156	215	215	169	142	111
Ningxia	30	34	32	41	75	72	66	61	36
Qinghai	6	6	6	10	14	13	15	13	13
Shaanxi	118	118	113	181	216	222	196	208	158
Shandong	464	510	493	629	689	677	731	712	580
Shanghai	96	103	101	95	109	75	83	93	84
Shanxi	292	284	253	328	397	395	373	313	260
Sichuan	155	158	179	204	232	254	271	280	205
Taiwan	100	106	98	106	113	118	106	114	111
Tianjin	77	86	99	136	152	114	102	97	88
Yunnan	83	109	105	97	118	159	144	176	126
Zhejiang	243	253	247	270	327	281	294	292	240
EastChina	4332	4502	4454	5382	6150	6402	6179	6201	4941