

# Cleaning up the air: Effectiveness of air quality policy for SO<sub>2</sub> and NO<sub>x</sub> emissions in China

Ronald J. van der A<sup>1,2</sup>, Bas Mijling<sup>1</sup>, Jieying Ding<sup>1,3</sup>, Maria Elissavet Koukouli<sup>4</sup>, Fei Liu<sup>1</sup>, Qing Li<sup>5</sup>, Huiqin Mao<sup>5</sup>, Nicolas Theys<sup>6</sup>

5 <sup>1</sup>Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

<sup>2</sup>Nanjing University of Information Science and Technology, Nanjing, P.R.China

<sup>3</sup>Delft University of Technology, Delft, the Netherlands

<sup>4</sup>Laboratory of Atmospheric Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece

<sup>5</sup>Satellite Environment Center, Ministry of Environmental Protection, Beijing, P.R.China

10 <sup>6</sup>Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

*Correspondence to:* Ronald J. van der A (avander@knmi.nl)

**Abstract.** Air quality observations by satellite instruments are global and have a regular temporal resolution, which make them very useful in studying long-term trends in atmospheric species. To monitor air quality trends in China for the period 15 2005-2015 we derive SO<sub>2</sub> columns and NO<sub>x</sub> emissions on a provincial level with improved accuracy. To put these trends into perspective they are compared with public data on energy consumption and the environmental policies of China. We distinguish the effect of air quality regulations from economic growth by comparing them relatively to fossil fuel consumption. Pollutant levels, per unit of fossil fuel, are used to assess the effectiveness of air quality regulations. We note that the desulphurisation regulations enforced in 2005-2006 only had a significant effect in the years 2008-2009 when a 20 much stricter control of the actual use of the installations began. For national NO<sub>x</sub> emissions a distinct decreasing trend is only visible since 2012, but the emission peak year differs from province to province. Unlike SO<sub>2</sub>, emissions of NO<sub>x</sub> are highly related to traffic. Furthermore, regulations for NO<sub>x</sub> emissions are partly decided on a provincial level. The last three years show both a reduction in SO<sub>2</sub> and NO<sub>x</sub> emissions per fossil fuel unit, since the authorities have implemented several new environmental regulations. Despite an increasing fossil fuel consumption and a growing transport sector, the effects of 25 air quality policy in China are clearly visible. Without the air quality regulations the concentration of SO<sub>2</sub> would be about 2.5 times higher and the NO<sub>2</sub> concentrations would be at least 25% higher than they are today in China.

## 1. Introduction

Satellite instruments can monitor air quality from space by mapping e.g. aerosols and tropospheric ozone, but are especially useful for emission estimates in observing the relatively short-living gases nitrogen dioxide (NO<sub>2</sub>) and sulphur dioxide (SO<sub>2</sub>). 30 For these two trace gases improved data sets recently became available, enabling analysis of air quality time series on a national or provincial level with improved accuracy. Theys et al. (2015) presented a new data set of SO<sub>2</sub> column densities derived from the Ozone Monitoring Instrument (OMI) satellite instrument (Levelt et al., 2006). They conclude that the SO<sub>2</sub> concentrations derived from OMI agree on average within 12% with ground observations. This data set strongly improves on earlier SO<sub>2</sub> data sets from satellites, which motivated this study. For NO<sub>2</sub>, instead of using concentration data, we assess 35 directly the emission data of nitrogen oxides (NO<sub>x</sub> = NO<sub>2</sub> + NO) that was derived from satellite observations by Mijling and Van der A (2012) and removes the meteorological influences. The precision of the derived NO<sub>x</sub> emissions per grid cell of 0.25 x 0.25 degree is estimated as 20% (Ding et al., 2016).

China is one of the biggest emitters of SO<sub>2</sub> and NO<sub>2</sub> into the atmosphere because its large economy depends heavily on fossil fuels as an energy source. China alone is responsible for about 30 % of the global total emissions of SO<sub>2</sub> into the atmosphere 40 (Klimont et al., 2013), while over 90% of the SO<sub>2</sub> emissions are caused by coal consumption in China (Chen and Xu, 2010).

Coal is mainly used by thermal power plants and energy-intensive industry (e.g. steel, cement and glass), and to a lesser extent by residential use. SO<sub>2</sub> is also released by the use of oil and natural gas, but the sulphur content in these fuel types is much lower. Of these sources power plants are responsible for about 30-40 % of all emissions and industry for another 50-60 % (He K. et al., 2012, ChinaFAQs project, 2012). According to the Multi-resolution Emission Inventory for China (MEIC) (http://www.meicmodel.org/) the source of SO<sub>2</sub> emissions in 2010 was 29.4 % from power plants, 57.7 % from industry and 11.7 % residential and 1.2 % from transport. Figure 1 shows the location of the 600 biggest thermal power plants on the map of China including a list of provinces mentioned in this study. At a global scale, volcanic activity is another important source of atmospheric SO<sub>2</sub>. However, plumes of active volcanoes are seldom observed over China.

NO<sub>x</sub> is released by more or less the same anthropogenic sources, i.e. the burning of coal or oil. The main difference with SO<sub>2</sub> is that traffic is a much more important source for NO<sub>x</sub>. NO<sub>x</sub> emission factors (i.e. emissions per fossil fuel unit) in the transport sector are generally much higher than emission factors in energy and industry, which makes traffic one of the major sources of NO<sub>x</sub> in China. According the MEIC inventory, 25% of NO<sub>2</sub> in 2010 was released by traffic, 32% by power plants, 4% by residential sources and 39% by industry, with the cement industry being the biggest emitter in this sector.

To reduce SO<sub>2</sub> in China, the authorities have implemented several environmental regulations. The most important regulation was the desulphurization of coal-fired power plants in 2005/2006 (Xu, 2011). This was later followed in the 12<sup>th</sup> five-year plan (2011-2015) by stricter control on the implementation of the regulations, additional filtering efforts, switching to low-sulphur coal and gasoline, phasing out obsolete capacity in coal-using industry, phasing out of small-scale coal mining, and gradually using more oil, gas and renewable energies instead of coal since 2011. An overview of all regulations related to SO<sub>2</sub> is shown in Table 1, which includes the year of the beginning of the implementation.

The regulation of NO<sub>x</sub> was started much later than for SO<sub>2</sub>. The 12<sup>th</sup> five-year plan mentioned the intention to reduce NO<sub>2</sub> by 10 % (target) (ChinaFAQs project, 2012), from 2011 onward NO<sub>x</sub> filtering systems were installed, mainly at power plants but also for heavy industry. These regulations for NO<sub>x</sub> were announced in 2013 in the Air Pollution Prevention and Control Action Plan (CAAC, 2013). According to Liu et al. (2016) Selective Catalytic Reduction (SCR) equipment was installed in this period and growing from a penetration of about 18% in 2011 until 86% in 2015. SCR equipment in power plants are expected to reduce the emissions of the power plant with at least 70% (ICAC, 2009). The SCR installation is the most significant measure taken to reduce the NO<sub>x</sub> emissions and it largely coincides with the peak year of observed NO<sub>2</sub> concentrations (Liu et al., 2016). At the same time China has implemented several new national emission standards for cars during the time period of our study (see Table 2). The change from China 3 to China 4 standard for cars in the period 2011-2015 reduces the maximum allowed amount of NO<sub>x</sub> emissions for on-road vehicles with 50% (Wu et al., 2017). More strict regulations for on-road vehicles (e.g. ban on older polluting cars) have been introduced on a city level, e.g. in Beijing, rather than nationwide. To our knowledge no regulations for ship emissions have been announced. Strong regulations have also been enforced during specific events like the Olympic Games in 2008, Shanghai World Expo in 2010, Nanjing Youth Olympic Games in 2014, and the APEC meeting in 2014, but those regulations were mostly of a temporary nature as shown by e.g. Mijling et al. (2012) for the Olympic Games in 2008.

To study the efficiency of the environmental policies, we analysed satellite observations of SO<sub>2</sub> and tropospheric NO<sub>2</sub> of the last 11 years. SO<sub>2</sub> satellite observations over China have been studied earlier by Lee et al. (2010), Li et al (2011), He (2012), Yang et al. (2013), Fioletov et al. (2015), and Krotkov et al. (2015). Satellite observations are very useful for SO<sub>2</sub> trend studies, as recently McLinden et al. (2016) showed that bottom-up inventories are underestimating SO<sub>2</sub> emissions worldwide with about 0-10%. NO<sub>2</sub> satellite observations over China have been evaluated by e.g. Richter et al. (2005), van der A et al. (2006), Zhang et al. (2012), and Krotkov et al. (2015). All these studies showed a strong increase in NO<sub>2</sub> over East China. On a city-scale or regional level trends are analysed and reported by Gu et al. (2013), Schneider et al. (2015), and Duncan et al. (2016). Although some cities showed already a decreasing trend, notably in the Pearl river delta, an overall decrease in NO<sub>2</sub> concentrations in China is only recently observed by Irie et al. (2016) Liu et al., (2016) and De Foy et al. (2016). To

exclude meteorology as a factor for variability in NO<sub>2</sub> several authors evaluated NO<sub>x</sub> emissions instead. Emission estimates of NO<sub>x</sub> over China have been analysed by Stavrou et al. (2008), Kurokawa et al. (2009), and more recently by Mijling et al. (2013) and by Liu et al. (2016a).

In these studies, whether of concentrations or emissions, linear trends of the air pollutants are often used. Here, however, we will relate changes derived on a provincial level for China with the energy consumption and the environmental policies of the country. This gives insight in the efficiency of the applied air quality policies and regulations. We apply this to NO<sub>x</sub> emissions instead of concentrations for the period 2007 until 2015. The comparison of SO<sub>2</sub> trends with those of NO<sub>x</sub> emissions enables us to distinguish environmental policies specifically applied on coal-based industry and power plants with general environmental measures and trends in traffic.

## 2 Observational data

### 2.1 Satellite observations of SO<sub>2</sub>

SO<sub>2</sub> is observed in the UV spectral range of satellite observations of SCIAMACHY (on Envisat), GOME-2 (on METOP-A) and OMI (on EOS-AURA). SO<sub>2</sub> retrieval algorithms have been earlier developed for GOME-1 by Eisinger and Burrows (1998), for SCIAMACHY (Lee et al., 2008), GOME-2 and for OMI by Krotkov et al (2006). Recently a new retrieval algorithm has been developed (Theys et al., 2015) that improves the precision of the SO<sub>2</sub> data for OMI with a factor 2, allowing us to derive more accurate trends based on OMI. The retrieval method is based on a Differential Optical Absorption Spectroscopy (DOAS) scheme to determine the slant columns from measured spectra in the 312-326 nm spectral range, which are then background corrected and converted to vertical columns using an Air Mass Factor (AMF). The AMF is calculated with the radiative transfer model LIDORT (LInearized Discrete Ordinate Radiative Transfer model). More details about the retrieval procedure are described in Theys et al. (2015). Also the operational algorithm of NASA for SO<sub>2</sub> from OMI has recently been improved. This algorithm and the algorithm of Theys et al. (2015) have a very comparable performance as shown by Fioletov et al. (2016). For this study, the algorithm of Theys et al. (2015) has been applied on the observations of the OMI instrument (Levelt et al., 2006) for its whole mission from 2004 onwards

To improve the quality of the OMI SO<sub>2</sub> data we exclude observations with a cloud fraction of more than 50 percent or with a fitting chi-square higher than 1. The solar zenith angle is limited to 75° and the viewing angle to 50°. Since the OMI instrument is suffering from the so-called row anomaly since 2007 (KNMI, 2012), we filter the affected rows (24-49, 54-55) in the same way for all years in the time series.

As we focus on anthropogenic SO<sub>2</sub>, the SO<sub>2</sub> data for 15 June - 9 July 2011 have been removed because of its contamination with volcanic SO<sub>2</sub> from the eruption of the Nabro volcano in Africa and the transport of its plume to China (Brenot et al., 2014).

As a first step in our study we have made monthly means for the whole data set by averaging and gridding the data to a resolution of 1/8° by 1/8°. The gridding algorithm takes into account the area of each satellite footprint overlapping the grid cell. The resulting data set is a time series of monthly means for the time period October 2004 to December 2015.

For comparison we also use the official ESA SCIAMACHY/Envisat SO<sub>2</sub> product version SGP 5.02 and the standard data product from the GOME-2/METOP-A version GDP 4.7, as developed within the EUMETSAT Satellite Application Facility for Atmospheric Composition and UV radiation, O3MSAF, project and distributed by <http://atmos.caf.dlr.de/gome2/>. The data of these instruments are noisier than the OMI datasets because of the lower spatial coverage, different fit window and the lower signal-to-noise ratio of the SCIAMACHY and GOME-2 instruments. Therefore, their quality-controlled monthly mean SO<sub>2</sub> data have been recalculated by spatially averaging for each grid cell the data from the eight surrounding neighbouring cells, hence creating a smoothed SO<sub>2</sub> field. For details on the methodology and findings refer to Koukouli et al. (2016).

## 2.2 NO<sub>x</sub> emission estimates from satellite observations

For NO<sub>x</sub> emission data we use the results of an update (version 4) of the DECSO (Daily Emission estimates Constrained by Satellite Observation) algorithm developed by Mijling and van der A (2012). DECSO calculates emissions by applying a Kalman filter for the inversion of satellite data and a regional Chemical Transport Model (CTM) for the forward model calculation. It takes transport from the source into account with a semi-Lagrangian approach. The CTM we use is CHIMERE v2013 (Menut et al., 2013) with meteorological information from the European Centre for Medium-range Weather Forecasts (ECMWF) with a horizontal resolution of approximately 25x25 km<sup>2</sup>. The DECSO algorithm is applied to OMI NO<sub>2</sub> observations derived by the DOMINO v.2 algorithm (Boersma et al., 2011). The latest improvements of the DECSO algorithm resulting in version 4 are described by Ding et al. (2015, 2016). The monthly average emission data over China we use is available on 0.25 degree resolution for the period 2007-2015 on the web-portal [www.globemission.eu](http://www.globemission.eu).

## 3 Temporal analysis over China

### 3.1 Sources of SO<sub>2</sub> and NO<sub>x</sub> in China

The multi-annual mean of SO<sub>2</sub> for 2005-2015 is shown in Figure 2. As the lifetime of SO<sub>2</sub> is relatively short (typically 4-48 hours) (Lee et al., 2011, Fioletov et al., 2015), the observed SO<sub>2</sub> concentrations are a good proxy for the location of SO<sub>2</sub> emissions. Regions with large SO<sub>2</sub> concentrations are South Hebei, the province Shandong (around the city Zibo) and the region around Chongqing. South Hebei is a region with many power plants just east of the mountainous coal-mining area in Shanxi. The hot spot in the province Shandong is related to a strongly industrialized area with lots of coal-using industry. In the Chongqing region both coal mines and heavy industry are located.

Rather than located at hot spots, high NO<sub>2</sub> concentrations are more distributed over the East of China, mainly because traffic is an important source of NO<sub>x</sub> emissions (see Figure 3a). The underlying NO<sub>x</sub> emissions are shown in Figure 3b. Like for the SO<sub>2</sub> concentrations, NO<sub>x</sub> emission spots can be found at the location of big power plants. Also clearly visible are the megacities of China, and ship tracks along the coast and sources along the big rivers.

### 3.2 SO<sub>2</sub> trends over China

To construct time series of SO<sub>2</sub> we have averaged the data to annual means of the vertical columns derived from OMI. From these annual mean SO<sub>2</sub> data we constructed time series for each province (see Table A.1). Figure 4 shows the mean normalized time series for the 10 provinces with the highest total SO<sub>2</sub> column densities (i.e. Tianjin, Shandong, Hebei, Shanxi, Henan, Beijing, Jiangsu, Shanghai, Anhui and Liaoning), together responsible for 60% of all ambient SO<sub>2</sub> in China. The individual time series are drawn as thin black lines. The minimum and maximum of these time series for each year are shown in the grey shaded area to indicate the variability. The time series of Shanghai is the lowest black line of the 10 series, thus the reductions have been strongest in this province since 2005. Apart from Ningxia province, all provincial time series show very similar patterns. In general, the SO<sub>2</sub> concentrations were at a maximum in the year 2007, when the start of decreasing trend is visible in China. Despite some fluctuations the SO<sub>2</sub> concentrations remain relatively constant from 2010 till 2013, where after they are decreasing again.

A different trend is observed for Ningxia, a province in the mid-north of the country with a relative low population density and large coal resources. Here an increasing trend emerges for the years starting from 2010 when several new coal power plants were put into operation. A list of largest power plants (with a capacity of more than 600 MW) and the start year of their operation is shown in Table 3. From 2012 onward, the more stringent SO<sub>2</sub> emission regulations also started to have effect in Ningxia.

### 3.3 NO<sub>x</sub> emission trends over China

National NO<sub>x</sub> emission trends show a different pattern than those of SO<sub>2</sub>. We observe an increasing trend till about 2012, with an exception of the year 2009 which is related to regulations started at the Olympic Games in 2008 (Mijling et al., 2008) and the global economic crisis which shortly slowed down the Chinese economic growth. Total NO<sub>x</sub> emissions in East China reached their peak levels in 2012, and have stopped increasing since this year. While the economy kept growing after 2012, the emission of NO<sub>x</sub> slowly decreases again as a result of the air quality regulations described in Section 1. According to the DECSO emission inversion, in 2015 the NO<sub>x</sub> emissions were 4.9 Tg N/yr, which is 22.8% lower than in the peak year 2012. However, the 2015 emissions were still 14.1% higher than in the reference year 2007. The trends per province (see Table A.2) show very similar patterns with only the starting year (the year with maximum NO<sub>x</sub> emissions) of the decrease in emissions varying over the provinces. Events like the Olympic Games in Beijing in 2008 and the World Expo in Shanghai in 2010, when temporary strict air quality regulations have been enforced, can be recognised in this Table as years with significant lower emissions for these provinces. In Figure 5a, the normalized (to the year 2007) time series of annual NO<sub>x</sub> emissions for East China (102-132°E, 18-50°N) is shown in similar way as for SO<sub>2</sub> in Figure 4. The mean, minimum and maximum of the 10 provinces with highest NO<sub>x</sub> emission are shown (Shandong, Hebei, Henan, Jiangsu, Guangdong, Shanxi, Zhejiang, Anhui, Sichuan and Hubei), together responsible for 65% of all Chinese NO<sub>x</sub> emissions. The thin black lines show the times series for the individual 10 provinces, where the lower line represents Guangdong. Figure 5b shows the peak year for each province. Provinces where air pollution regulations, for e.g. traffic, got a lot of attention at an early stage, like Beijing and Shanghai, have reached their maximum before 2011. Most industrialised regions show their peak in the years 2011-2013. Some of the less developed and populated provinces show a maximum in 2014, which means that their decrease in NO<sub>x</sub> emissions is very recent. Regional variations are mainly due to the fact that regulations for the NO<sub>x</sub> emission reductions, for instance in traffic or power plants, are determined and implemented on a provincial level (Liu et al., 2016b). For the province Ningxia we see a very similar pattern occurring as for SO<sub>2</sub>, which shows that for this low-densely populated province traffic plays a small role and the trend is determined by the operation of newly-built power plants.

### 3.4 Air pollution in relation to fossil fuel consumption

To relate the observed SO<sub>2</sub> and NO<sub>x</sub> reduction to environmental regulations we have to take into account the coal and oil consumption in the same time period. The total coal consumption in Standard Coal Equivalent (SCE) units per year for China and the total oil consumption (also in SCE units) are shown in Figure 6, based on data of NBSC (2015). According to Guan et al. (2012) and Hong et al. (2016) the sum of the coal consumption of all provinces is more accurate than the number provided for the whole of China, thus we use the provincial totals for coal consumption. For NO<sub>x</sub> emissions the transport sector plays an important role, especially ships are one of the largest NO<sub>x</sub> emitters per fuel unit in the transport sector. The total freight transport almost doubles every 6 years in China.

Since the burning of coal and oil are the dominant sources of SO<sub>2</sub> and NO<sub>x</sub> emissions, we can consider the total emissions of these air pollutants as the product of the national use of coal and oil (activity) and the average emission factor of one unit coal/oil. The effectiveness of environmental regulation will be reflected in a decrease of this emission factor. Therefore, we divide the annual SO<sub>2</sub> column measured from satellites and the annual NO<sub>x</sub> emissions by the annual coal and oil consumption in China. In this way we get a measure of the emitted SO<sub>2</sub> or NO<sub>x</sub> per unit (SCE) of fossil fuel consumption reflecting the Chinese environmental policy. The results are shown in Figure 7. One might argue that SO<sub>2</sub> is more related to coal than oil, but division by only coal yields the same results. In our analysis we omit gas consumption since this is very limited in China and hence does not affect the results significantly.

We focus here mainly on the results for OMI, because of the instrument's high spatial resolution and lack of instrumental degradation. However, SO<sub>2</sub> data of the SCIAMACHY and GOME-2 instrument are also added in Figure 7 to be able to further look into the past (starting in 2003) and to verify the results of OMI. The SO<sub>2</sub> data of SCIAMACHY and GOME-2

are averaged over the summer months (April-September). The remaining monthly means are excluded from the analysis due to a lower accuracy at higher latitudes and a large part of the higher latitudes is missing due to snow cover. For OMI each data point is averaged over 12 months and the total area of China, which reduces the root-mean-square error to a negligible level. Biases among all instruments are removed by normalizing the values to those in reference year 2007. Up to 2009, the results agree fairly well. After 2009, we see the results of GOME-2 and OMI for SO<sub>2</sub> slowly diverge in time, which might be result of the instrument degradation of the UV spectra of GOME-2 after 2009 (Munro et al., 2016).

Changing weather conditions from year-to-year can affect the results for SO<sub>2</sub> concentrations and when these weather conditions are different during the overpass of SCIAMACHY and GOME-2 (around 9:30 local time) and overpass of OMI (around 13:30 local time), this can lead to differences between the instruments. The global coverage of SCIAMACHY is once every 6 days and for GOME-2 and OMI almost daily. The limited number of samples for SCIAMACHY makes this data more sensitive to weather conditions. Note that due to the nature of the inversion algorithm the NO<sub>x</sub> emission data is in general not sensitive to meteorological variability.

For SO<sub>2</sub> we see a big decrease in the years 2008 and 2009, while the desulphurization program of the 11<sup>th</sup> five-year plan started already in 2005/2006, when the authorities begin to reduce SO<sub>2</sub> emissions by installing desulphurization devices in many power plants (Lu et al., 2010). In 2006 SO<sub>2</sub> monitoring devices were also installed in the chimneys of the power plants. This resulted in a decrease in SO<sub>2</sub> emissions from 2006, while the much bigger decrease of SO<sub>2</sub> in 2008-2009 reflects the stronger government control at that time on the actual use of the equipment (Xu et al., 2011). After 2009, the SO<sub>2</sub> content per consumed coal unit only slowly decrease until 2011. From 2012 onwards we see a stronger annual decrease in SO<sub>2</sub>. This coincides with the 12<sup>th</sup> five-year program when new measures were taken to upgrade the coal quality, to modernize the industry and to put more effort on law enforcement. Especially the law enforcement in the last years concerning the prohibition of flue gas bypass and the use of desulphurization devices in the steel industry played an important role.

For the NO<sub>x</sub> emissions the total annual emissions are used and divided in the same way as for SO<sub>2</sub> by the total coal and oil consumption. Here, however we should keep in mind that the transport sector (especially by shipping) emits much more NO<sub>x</sub> per fuel unit than the power and industrial sectors (see e.g. Zhao et al., 2013). Thus the percentage of the total fuel used by transport is relevant for the graph of NO<sub>x</sub>. In the early years we see in general a small increase in NO<sub>x</sub> emissions per fuel unit due to the increasing fraction of the transport sector in the fuel use. Exceptions are the year 2009 and the recent year 2015. The year 2009 coincides with the global economic crisis (Lin and McElroy, 2011) when there was less export of goods from China. This affected especially the transport sector, mostly transport over water, as shown in De Ruyter de Wildt et al. (2012). Faber et al. (2012) and Boersma et al. (2015) showed that the economic crisis also resulted in a significant reduction of the average vessel speed to save fuel used by ship transport. This caused not only a shift in source sectors but in general led to lower NO<sub>x</sub> per fuel values. This explains the dip in pollution per fuel unit in 2009. After 2009 the NO<sub>x</sub> per fossil fuel is slowly increasing because the transport sector is growing faster than the energy sector and has a higher emission factor. Statistics of the NBSC (2015) show that the transport is growing with a factor two every 5-6 years (Wu et al., 2017). After 2012 the gradual increase of NO<sub>x</sub> per fuel slowly stops, and the year 2015 shows a sharp decline in NO<sub>x</sub> per fossil fuels unit. This can be directly related to the rapidly growing installation of SCR equipment at power plants since 2012 and to a lesser account to the introduction of new emission standards for cars, as shown by Liu et al. (2016). This strong reduction in NO<sub>x</sub> for 2015 and the equally strong reduction for SO<sub>2</sub> in 2014 and 2015 are a result of very effective recent environmental regulations in the last years in China. By comparing the efficiency level in 2015 with earlier levels we can conclude from Figure 7 that without these air quality regulations SO<sub>2</sub> concentrations would nowadays be about 2.5 times higher. For NO<sub>x</sub> per fossil fuel we were expecting a gradual growth after 2012 because of the continuing relative growth of the transport sector. Keeping this in mind we compare the years 2015 with 2012 and conclude that without air quality regulations the NO<sub>2</sub> concentrations would be at least 25% higher in China today.

On a provincial scale we can in principle do similar analyses, but unfortunately the provincial energy consumption related to coal and oil have a very high uncertainty due to inconsistencies in interprovincial imports and exports (Hong et al., 2016). We see this reflected in a high variability in the annual provincial data and sometimes missing data. Especially for the oil consumption the data have high uncertainties (Guan et al., 2011, Hong et al., 2016). Therefore, we have only analysed the 5 provinces with dominating coal consumption as shown in Figure 8. In this graph we excluded Guizhou province because of its difficult to interpret coal consumption in 2011 as a result of large power shortages (NBSC, 2015, Xinhua, 2011). For SO<sub>2</sub> per fossil fuel unit we see that all provinces follow the national trend. For NO<sub>x</sub> per fossil fuel we see more variation per province depending on the role of transport. Most of these by coal consumption dominated provinces start their decreasing trend from 2011 reflecting the national program on SCR installations starting that same year. It is interesting to see that the commissioning of new power plants in 2011 causes a strong increase in both SO<sub>2</sub> and NO<sub>x</sub> in the province Ningxia (see Figure 4 and 5a). However, when compensating for fossil fuel usage, one can see that the same national air quality regulations are applied here, as the trend in Figure 8 shows the same pattern as for other provinces. This show the strength of the presented method to assess the efficiency of air pollution regulations.

## 15 4 Discussion

The current developments in data products derived from satellite observations provide high quality time series of the air pollutants NO<sub>x</sub> and SO<sub>2</sub>. Although the mean of observed SO<sub>2</sub> columns are not linearly related to the SO<sub>2</sub> emissions because of the influence of the weather, it can still be argued that these satellite data products, whether concentrations or emissions, provide a fair comparison over the various regions from year to year. By comparing these time series with fossil fuel energy consumption the economic growth is removed from the equation and we can monitor the effectiveness of air quality policies. We foresee that this method will become a valuable tool for policy makers concerning air quality regulations.

For China we see similar patterns in the trends of SO<sub>2</sub> per province. In 2006 a nation-wide implementation of desulphurisation installations started. However, the effects are only visible in 2008 and 2009 when a strict control by the Chinese authorities on the actual use of the desulphurisation installations started. In 2009, we see the effect of the air quality regulations for SO<sub>2</sub> and NO<sub>x</sub> resulting from the global economic recession at the end of 2008. The increasing relative contribution of the transport sector to the NO<sub>x</sub> emission slowly increases the amount of NO<sub>x</sub> per fossil fuel unit after 2009. After 2011 we see a steadily decreasing SO<sub>2</sub> pollution per fossil fuel unit caused by various Chinese environmental regulations. In the last year of our time series, 2015, a clear effect becomes visible of very recent regulations for NO<sub>x</sub> emissions from power plants and heavy industry. The fit of linear trends often used in earlier studies is therefore no longer applicable to the Chinese situation.

The availability of high quality satellite data for the last ten years is especially interesting for China where the situation is rapidly changing. For instance in Europe and Japan desulphurisation started much earlier when these satellite data were not yet available. On the other hand, in India SO<sub>2</sub> and NO<sub>x</sub> emissions are still growing and possible new regulations can be monitored in the years to come with an even better quality using forthcoming sensors as e.g. TROPOMI on-board Sentinel-5 Precursor.

Despite the growing use of coal and oil in the last ten years in China we see reduced emissions per fuel unit in the past few years. This decreasing trend in both SO<sub>2</sub> and NO<sub>x</sub> for China is likely to continue in the coming years for which the Chinese national government has announced less use of coal, more environmental regulations for SO<sub>2</sub> and NO<sub>x</sub> and stricter reinforcement of control of environmental policies.

40

## Acknowledgements

This research has been funded by the MarcoPolo project of the European Union Seventh Framework Programme (FP7/2007-2013) under Grant Agreement n° 606953 and by the GlobEmission project (Contract No 4000104001/11/I-NB) of the Data User Element programme of the European Space Agency.

## 5 5 References

- Boersma, K.F., H.J. Eskes, R. J. Dirksen, R. J. van der A, J. P. Veefkind, P. Stammes, V. Huijnen, Q. L. Kleipool, M. Sneep, J. Claas, J. Leitao, A. Richter, Y. Zhou, and D. Brunner, An improved retrieval of tropospheric NO<sub>2</sub> columns from the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905-1928, 2011
- Boersma, K.F., G.C.M. Vinken, J. Tournadre, Ships going slow in reducing their NO<sub>x</sub> emissions: changes in 2005–2012 ship exhaust inferred from satellite measurements over Europe, *Environ. Res. Letters*, **10**, 7, doi:10.1088/1748-9326/10/7/074007, 2015
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: mission objectives and measurement modes, *J. Atmos. Sci.*, 56, 127–150, doi:10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2, 1999.
- 15 Brenot, H., N. Theys, L. Clarisse, J. van Geffen, J. van Gent, M. Van Roozendaal, R.J. van der A, D. Hurtmans, P.F. Coheur, C. Clerbaux, P. Valks, P. Hedelt, F. Prata, O. Rasson and C. Zehner, Support to Aviation Control Service (SACS): an online service for near-real-time satellite monitoring of volcanic plumes, *Natural Hazards and Earth System Sciences*, 2014, 14, 1099-1123, doi:10.5194/nhess-14-1099-2014.
- CAAC 2013, Clean Air Alliance of China, State Council air pollution prevention and control action plan, issue II, October 2013, <http://en.cleairchina.org/product/6346.html> (English translation). Last accessed: 8 October 2015
- 20 ChinaFAQs Project, <http://www.chinafaqs.org/library/chinafaqs-china-adopts-world-class-pollutant-emissions-standards-coal-power-plants>, 2012. Last accessed: 8 October 2015
- de Foy, B., Z. Lu and D. G. Streets, Satellite NO<sub>2</sub> retrievals suggest China has exceeded its NO<sub>x</sub> reduction goals from the twelfth Five-Year Plan, *Scientific Reports* 6, 35912 (2016), doi:10.1038/srep35912
- 25 De Ruyter de Wildt, M., H. Eskes, K. F. Boersma, The global economic cycle and satellite-derived NO<sub>2</sub> trends over shipping lanes, *Geophys. Res. Letters*, doi:10.1029/2011GL049541, 2012
- Ding, J., van der A, R. J., Mijling, B., Levelt, P. F., and Hao, N.: NO<sub>x</sub> emission estimates during the 2014 Youth Olympic Games in Nanjing, *Atmos. Chem. Phys.*, 15, 9399-9412, doi:10.5194/acp-15-9399-2015, 2015.
- Ding, J, van der A, R. J., Mijling, B., Levelt, P. F., New NO<sub>x</sub> emission inversion algorithm improves estimates over low-  
30 emission regions., *J. Geophys. Res.*, submitted, 2016
- Duncan B N, Lamsal L N, Thompson A M, Yoshida Y, Lu Z, Streets D G, Hurwitz M M and Pickering K E, A space-based, high-resolution view of notable changes in urban NO<sub>x</sub> pollution around the world (2005–2014), *J. Geophys. Res.* 121, 976–96, 2016
- Eisinger, M., and J. P. Burrows (1998), Tropospheric sulfur dioxide observed by the ERS-2 GOME instrument, *Geophys. Res. Lett.*, 25, 4177–4180.
- 35 Faber J, Nelissen D, Hon G, Wang H and Tsimplis M, Regulated slow steaming in maritime transport—an assessment of options, costs and benefits CE Delft (The Netherlands: Delft) ([www.cedelft.eu/publicatie/regulated\\_slow\\_steaming\\_in\\_maritime\\_transport/1224](http://www.cedelft.eu/publicatie/regulated_slow_steaming_in_maritime_transport/1224)) p 117, 2012
- Fioletov, V. E., C. A. McLinden, N. Krotkov, and C. Li (2015), Lifetimes and emissions of SO<sub>2</sub> from point sources  
40 estimated from OMI. *Geophys. Res. Lett.*, 42, 1969–1976. doi: 10.1002/2015GL063148.



- Fioletov, V. E., McLinden, C. A., Krotkov, N., Li, C., Joiner, J., Theys, N., Carn, S., and Moran, M. D.: A global catalogue of large SO<sub>2</sub> sources and emissions derived from the Ozone Monitoring Instrument, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-417, in review, 2016.
- Gu D, Wang Y, Smeltzer C and Liu Z 2013 Reduction in NO<sub>x</sub> emission trends over China: regional and seasonal variations, *Environ. Sci. Technol.*, 47, 12912–9
- Guan, D., Z. Liu, Y. Geng, S. Lindner and K. Hubacek, The gigatonne gap in China's carbon dioxide inventories, *Nature Climate Change*, 2, 672–675, (2012), doi:10.1038/nclimate1560
- Hong, C., Zhang, Q., He, K., Guan, D., Li, M., Liu, F., and Zheng, B.: Variations of China's emission estimates response to uncertainties in energy statistics, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-459, in review, 2016.
- 10 He, Kebin: Multi-resolution Emission Inventory for China (MEIC): model framework and 1990-2010 anthropogenic emissions, in International Global Atmospheric Chemistry Conference, 17-21 September, Beijing, China. [online] Available from: <http://adsabs.harvard.edu/abs/2012AGUFM.A32B..05H> (Accessed 4 February 2015), 2012.
- He, H., et al. (2012), SO<sub>2</sub> over central China: Measurements, numerical simulations and the tropospheric sulfur budget, *J. Geophys. Res.*, 117, D00K37, doi:10.1029/2011JD016473.
- 15 ICAC. Institute of Clean Air Companies, Selective Catalytic Reduction (SCR) control of NO<sub>x</sub> emissions from fossil fuel-fired electric power plants, [https://c.yimcdn.com/sites/icac.site-ym.com/resource/resmgr/Standards\\_WhitePapers/SCR\\_WhitePaper\\_final\\_2009.pdf](https://c.yimcdn.com/sites/icac.site-ym.com/resource/resmgr/Standards_WhitePapers/SCR_WhitePaper_final_2009.pdf), 2009, Last accessed: 22 November 2016.
- Klimont, Z., Smith, S. J., and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions, *Environ. Res. Lett.*, 8, 014003, doi:10.1088/1748-9326/8/1/014003, 2013.
- 20 KNMI: Background information about the Row Anomaly in OMI, Retrieved from: <http://www.knmi.nl/omi/research/product/rowanomaly-background.php>, 2012, Last accessed: 8 October 2015.
- Koukoulis, M.E., Balis, D.S., van der A, R., Theys, N., Hedelt, P., Lichtenberg, G., Richter, A., Krotkov, N., Li, C., Taylor, M., Characteristics of the anthropogenic SO<sub>2</sub> load over China observed by SCIAMACHY/Envisat, OMI/Aura and
- 25 GOME2/MetopA, paper in preparation for Atmospheric Environment, 2016.
- Krotkov, N.A., S.A. Carn, A.J. Krueger, P.K. Bhartia, and K. Yang, Band residual difference algorithm for retrieval of SO<sub>2</sub> from the aura ozone monitoring instrument (OMI), *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 44, No. 5, 2006.
- Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J.,
- 30 Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z., and Streets, D. G.: Aura OMI observations of regional SO<sub>2</sub> and NO<sub>2</sub> pollution changes from 2005 to 2014, *Atmos. Chem. Phys. Discuss.*, 15, 26555-26607, doi:10.5194/acpd-15-26555-2015, 2015.
- Kurokawa, J. I., Yumimoto, K., Uno, I., & Ohara, T. (2009). Adjoint inverse modeling of NO<sub>x</sub> emissions over eastern China using satellite observations of NO<sub>2</sub> vertical column densities. *Atmospheric Environment*, 43(11), 1878-1887.
- 35 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Greet, J.-M., Fukui, T., Kawashima, K. and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058.
- Lee, C., Richter, A., Weber, M., and Burrows, J. P.: SO<sub>2</sub> Retrieval from SCIAMACHY using the Weighting Function DOAS (WFDOAS) technique: comparison with Standard DOAS retrieval, *Atmos. Chem. Phys.*, 8, 6137-6145, doi:10.5194/acp-8-6137-2008, 2008.
- 40 Lee, C., R. V. Martin, A. van Donkelaar, H. Lee, R. R. Dickerson, J. C. Hains, N. Krotkov, A. Richter, K. Vinnikov, and J. J. Schwab (2011), SO<sub>2</sub> emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations, *J. Geophys. Res.*, 116, D06304, doi:10.1029/2010JD014758.

- Levelt, P. F., G. H. J. van den Oord, M. R. Dobber, A. Mälkki, H. Visser, J. de Vries, P. Stammes, J. Lundell, and H. Saari (2006), The ozone monitoring instrument, *IEEE Trans. Geo. Rem. Sens.*, 44(5), 1093–1101, doi:10.1109/TGRS.2006.872333.
- Li, C., Zhang, Q., Krotkov, N. A., Streets, D. G., He, K., Tsay, S.-C., and Gleason, J. F.: Recent large reduction in sulfur dioxide emissions from Chinese power plants observed by the Ozone Monitoring Instrument, *Geophys. Res. Lett.*, 37, 1–6, doi:10.1029/2010GL042594, 2010.
- Lin J T and McElroy M B 2011 Detection from space of a reduction in anthropogenic emissions of nitrogen oxides during the Chinese economic downturn *Atmos. Chem. Phys.* 11, 8171–88
- Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., and He, K. B.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, *Atmos. Chem. Phys.*, 15, 13299–13317, doi:10.5194/acp-15-13299-2015, 2015.
- Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K., and Wagner, T.: NO<sub>x</sub> lifetimes and emissions of cities and power plants in polluted background estimated by satellite observations, *Atmos. Chem. Phys.*, 16, 5283–5298, doi:10.5194/acp-16-5283-2016, 2016a.
- Liu, F., Q. Zhang, R. J. van der A, B. Zheng, D. Tong, L. Yan and K. B. He, Recent reduction in NO<sub>x</sub> emissions over China from satellite observations and emission inventories, submitted to *Env. Res. Letters*, 2016b.
- Lu, Z., Streets, D. G., Zhang, Q., Wang, S., Carmichael, G. R., Cheng, Y. F., Wei, C., Chin, M., Diehl, T., and Tan, Q.: Sulphur dioxide emissions in China and sulphur trends in East Asia since 2000, *Atmos. Chem. Phys.*, 10, 6311–6331, doi:10.5194/acp-10-6311-2010, 2010.
- McLinden, C. A., V. Fioletov, M. W. Shephard, N. Krotkov, C. Li, R. V. Martin, M. D. Moran, and J. Joiner, Space-based detection of missing sulfur dioxide sources of global air pollution, *Nature Geoscience*, DOI: 10.1038/ngeo2724, 2016
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G., Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R. and Vivanco, M. G.: CHIMERE 2013: a model for regional atmospheric composition modelling, *Geosci. Model Dev.*, 6(4), 981–1028, doi:10.5194/gmd-6-981-2013, 2013
- MEP, Ministry of Environmental Protection, [http://english.mep.gov.cn/Plans\\_Reports/](http://english.mep.gov.cn/Plans_Reports/), last access: 8 October 2015.
- Mijling, B. and R.J. van der A, Using daily satellite observations to estimate emissions of short-lived air pollutants on a mesoscopic scale, *J. Geophys. Res.*, 117, 2012, doi:10.1029/2012JD017817.
- Mijling, B., R.J. van der A and Q. Zhang, Regional nitrogen oxides emission trends in East Asia observed from space, *Atm. Chem. Phys.*, 2013, 13, 12003–12012, doi:10.5194/acp-13-12003-2013.
- Munro, R., Lang, R., Klaes, D., Poli, G., Retscher, C., Lindstrot, R., Huckle, R., Lacan, A., Grzegorski, M., Holdak, A., Kokhanovsky, A., Livschitz, J., and Eisinger, M.: The GOME-2 instrument on the Metop series of satellites: instrument design, calibration, and level 1 data processing – an overview, *Atmos. Meas. Tech.*, 9, 1279–1301, doi:10.5194/amt-9-1279-2016, 2016.
- NBSC, National Bureau of Statistics China, <http://data.stats.gov.cn>, last accessed: 29 December 2015
- NDRC, 2013, National Development and Reform Commission (reported on 16 August 2013), [http://www.ndrc.gov.cn/zcfb/zcfbtz/201308/t20130827\\_555124.html](http://www.ndrc.gov.cn/zcfb/zcfbtz/201308/t20130827_555124.html) (last accessed on 01-11-2015)
- Richter, A., J. P. Burrows, H. Nuß, C. Granier, and U. Niemeier (2005), Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, 437, 129 – 132, doi:10.1038/nature04092.
- Schneider, P., W.A. Lahoz and R.J. van der A, Recent Satellite-based Trends of Tropospheric Nitrogen Dioxide over Large Urban Agglomerations Worldwide, *Atm. Chem. Phys.*, 2015, 15, 1205–1220, doi:10.5194/acp-15-1205-2015.
- State Council, 2014, State Council announcement on 15-5-2014 of environmental regulations), [http://www.gov.cn/zhengce/content/2014-05/26/content\\_8824.htm](http://www.gov.cn/zhengce/content/2014-05/26/content_8824.htm) (last accessed on 01-11-2015)

- Stavrakou, T., Müller, J. F., Boersma, K. F., De Smedt, I., & van der A, R. J. (2008). Assessing the distribution and growth rates of NO<sub>x</sub> emission sources by inverting a 10-year record of NO<sub>2</sub> satellite columns. *Geophysical Research Letters*, 35(10).
- 5 Theys, N., De Smedt, I., van Gent, J., Danckaert, T., Wang, T., Hendrick, F., Stavrakou, T., Bauduin, S., Clarisse, L., Li, C., Krotkov, N., Yu, H., Brenot, H., and Van Roozendael, M.: Sulfur dioxide vertical column DOAS retrievals from the Ozone Monitoring Instrument: global observations and comparison to ground-based and satellite data, *J. Geophys. Res.-Atmos.*, 120, 2470–2491, doi:10.1002/2014JD022657, 2015.
- 10 van der A, R. J., Peters, D. H. M. U., Eskes, H., Boersma, K. F., Van Roozendael, M., De Smedt, I., and Kelder, H. M.: Detection of the trend and seasonal variation in tropospheric NO<sub>2</sub> over China, *J. Geophys. Res.*, 111, D12317, doi:10.1029/2005JD006594, 2005.
- Wu, Y., S. Zhang, J. Hao, H. Liu, X. Wu, J. Hu, M. P. Walsh, T. J. Wallington, K. M. Zhang, S. Stevanovic, On-road vehicle emissions and their control in China: A review and outlook, *Science of The Total Environment*, 574, 332-349, doi:10.1016/j.scitotenv.2016.09.040, 2017
- 15 Sun, L. (ed.), N. Zhou (2011, October 22):, Guizhou power supply tight, *Xinhua News Agency*, Retrieved from [http://news.xinhuanet.com/fortune/2011-10/22/c\\_111115420.htm](http://news.xinhuanet.com/fortune/2011-10/22/c_111115420.htm)
- Xu, Y., Improvements in the Operation of SO<sub>2</sub> Scrubbers in China's Coal Power Plants, *Environ. Sci. Technol.*, 2011, 45 (2), pp 380–385, DOI: 10.1021/es1025678.
- 20 Yang, K., R. R. Dickerson, S. A. Carn, C. Ge, and J. Wang (2013), First observations of SO<sub>2</sub> from the satellite Suomi NPP OMPS: Widespread air pollution events over China, *Geophys. Res. Lett.*, 40, 4957–4962, doi:10.1002/grl.50952.
- Zhang, Q., Geng, G., Wang, S., Richter, A., & He, K. (2012). Satellite remote sensing of changes in NO<sub>x</sub> emissions over China during 1996–2010. *Chinese Science Bulletin*, 57(22), 2857-2864.
- Zhang R. (2013, August 11): Heating causes pollution spike in Northern China, *CCTV.com*, Retrieved from <http://english.cntv.cn/program/china24/20131108/101015.shtml>
- 25 Zhu N. (2013, October 12): China to shut coal mines in safety overhaul, *Xinhua News Agency*, Retrieved from [http://news.xinhuanet.com/english/china/2013-10/12/c\\_132793667.htm](http://news.xinhuanet.com/english/china/2013-10/12/c_132793667.htm)
- Zhao, B., Wang, S. X., Liu, H., Xu, J. Y., Fu, K., Klimont, Z., Hao, J. M., He, K. B., Cofala, J., and Amann, M.: NO<sub>x</sub> emissions in China: historical trends and future perspectives, *Atmos. Chem. Phys.*, 13, 9869-9897, doi:10.5194/acp-13-9869-2013, 2013.



**Figure 1: Location of power plants in China according to REAS v.2 (Kurokawa et al., 2013). The size of each dot indicates the emission of the power plants (Power plants in close proximity are combined in a single dot). In addition, a list is given of the provinces mentioned in this study.**





Figure 2: Average SO<sub>2</sub> concentrations for the period 2005 to 2015 as observed by the OMI satellite instrument. Data below 0.1 DU is masked (grey colour).



5

Figure 3: (a) The averaged tropospheric NO<sub>2</sub> concentrations over China measured by OMI in the period 2005-2014. (b) The NO<sub>x</sub> emissions in the year 2010 derived from the OMI satellite observations.





Figure 4: Time series (red line) of the annual mean of the 10 provinces with the highest  $\text{SO}_2$  concentrations derived from the OMI satellite observations. The time series are normalized to their value in 2005. The grey area indicates the maximum range of the individual values of the times series of each of the 10 provinces. The thin black lines show the individual time series of those provinces. The province of Ningxia has a distinct deviating trend, here shown in blue.

5



(a)

(b)

Figure 5: (a) Shown are the annual total  $\text{NO}_x$  emission estimates for the last 9 years for the top 10 of highest  $\text{NO}_x$  emitting provinces in East China. Emissions are derived with DECSO V4 using OMI observations. The thin black lines show the individual time series of those provinces. (b) Peak year of the  $\text{NO}_x$  emissions per province.

10




Figure 6: In black the annual coal consumption and in red the annual oil consumption for China is shown.





Figure 7: Time series of the ratio of the mean SO<sub>2</sub> columns and the fossil fuel consumption in China based on observations of OMI (black), SCIAMACHY (green), and GOME-2 (blue). The ratios of the annual NO<sub>x</sub> emissions and the fossil fuel consumption is based on observations of OMI (red). All time series are normalized to the year 2007.



5 Figure 8: Same graph as Figure7 but with time series for the provinces Hebei, Henan, Nei Mongol, Ningxia and Shanxi included. Time series per province of the ratio of the mean SO<sub>2</sub> columns and the fossil fuel consumption are drawn in blue. The ratios of the annual NO<sub>x</sub> emissions and the fossil fuel consumption per province are shown in red. All time series are normalized to the year 2007 and based on OMI observations.



**Table 1: Environmental regulations of the Chinese national government to reduce SO<sub>2</sub> in the air.**

<b>Start year of implementation</b>	<b>Regulation</b>	<b>Reference</b>
2005-2006	Desulphurization techniques in power plants.	Li et al., 2010
2005-2012	Closure of several of the most polluting power plants	Liu et al., 2015
2008	Stricter control of implementation of desulphurization in power plants	Xu et al., 2011 Liu et al., 2015
2011	Use of more gas and renewable energies instead of coal	NBSC, 2015
January 2012	New emission standard of air pollutants for thermal power plants	MEP, 2015
2013	Mandatory SO <sub>2</sub> filtering of small-scale coal-fired industry	Zhang, 2013, NDRC, 2013
End of 2013	Stricter control of environmental policy	CAAC, 2013, State Council, 2014
End of 2013	Further desulphurization in industry	CAAC, 2013, NDRC, 2013
2014	Phasing out small-scale coal-fires boilers	CAAC, 2013, State Council, 2014
2014	Closure of 2000 small-scale coal mines	Zhu, 2013
End of 2014	Use of low-sulphur coal	State Council, 2014
End of 2014	Cap on coal consumption	State Council, 2014

5

**Table 2: Environmental regulations of the Chinese national government to reduce NO<sub>x</sub> emissions.**

<b>Year of implementation</b>	<b>Regulation</b>	<b>Reference</b>
2011-2015	Installation of Selective Catalytic Reduction (SCR) equipment at power plants. In 2013 the SCR equipment was installed in about 50% of all power plants.	Liu et al., 2016, CAAC, 2013
2007	China 3 (Euro 3) emissions standards for cars, nationwide	Wu et al., 2017
2011	China 4 (Euro 4) emissions standards for gasoline cars, nationwide.	Wu et al., 2017
2015	China 4 (Euro 4) emissions standards for diesel cars, nationwide.	Wu et al., 2017

**Table 3: Main power plants in Ningxia province (> 600 MW). Data collected from [www.sourcewatch.org](http://www.sourcewatch.org).**

<b>Power plant</b>	<b>Capacity (MW)</b>	<b>In operation since</b>	<b>Remark</b>
CPI Linhezhen	700	unknown	
Daba-1	1200	< 2000	
Daba-2	1100	unknown	An extension of Daba-1
Ningxia Zhongning-2	660	2005-2006	
Guodian Shizuishan-2	1980	2006	
Ningdong Maliantai	660	2006	
Huadian Ningxia Lingwu units 1&2	1200	2007	
Guodian-Dawukou	1100	2010	Extension of the original 440 MW plant
Guohua Ningdong	660	2010	
Ningxia Liupanshan	660	2010	
Huadian Ningxia Lingwu units 3&4	2120	2010-2011	
Shenhua Yuanyang Lake	1320	2010-2011	
Shuidonggou	1200	2011	
Ningdong Younglight	660	2013	

## Appendix A

**Table A1 Annual SO<sub>2</sub> column densities (DU/grid cell) per province observed by OMI**

Province	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015
Anhui	0.625	0.586	0.953	0.637	0.499	0.567	0.671	0.553	0.602	0.379	0.280
Beijing	0.753	0.829	0.989	0.711	0.778	0.749	0.850	0.673	0.634	0.640	0.491
Chongqing	0.514	0.509	0.530	0.567	0.580	0.580	0.492	0.370	0.469	0.269	0.136
Fujian	0.099	0.123	0.196	0.135	0.104	0.113	0.112	0.080	0.107	0.076	0.064
Gansu	0.144	0.136	0.150	0.130	0.135	0.123	0.134	0.127	0.131	0.105	0.103
Guangdong	0.251	0.257	0.280	0.239	0.171	0.177	0.138	0.095	0.118	0.086	0.080
Guangxi	0.199	0.203	0.270	0.236	0.127	0.190	0.179	0.092	0.134	0.091	0.072
Guizhou	0.424	0.478	0.532	0.516	0.418	0.424	0.357	0.261	0.345	0.167	0.100
Hainan	0.098	0.086	0.091	0.092	0.060	0.090	0.106	0.027	0.087	0.055	0.000
Hebei	0.903	0.931	0.996	0.908	0.874	0.881	0.922	0.863	0.844	0.716	0.540
Heilongjiang	0.134	0.141	0.144	0.142	0.124	0.138	0.154	0.162	0.125	0.134	0.135
Henan	1.036	0.920	1.222	0.938	0.709	0.778	0.992	0.827	0.762	0.585	0.439
Hubei	0.487	0.477	0.603	0.490	0.342	0.386	0.479	0.365	0.378	0.288	0.176
Hunan	0.364	0.330	0.448	0.371	0.270	0.281	0.320	0.240	0.259	0.180	0.112
Jiangsu	0.847	0.782	1.054	0.917	0.678	0.716	0.871	0.687	0.735	0.524	0.326
Jiangxi	0.272	0.278	0.373	0.267	0.202	0.222	0.244	0.197	0.230	0.184	0.136
Jilin	0.205	0.233	0.260	0.259	0.191	0.207	0.201	0.187	0.203	0.192	0.156
Liaoning	0.512	0.576	0.602	0.568	0.504	0.478	0.475	0.479	0.515	0.480	0.321
NeiMongol	0.154	0.170	0.200	0.174	0.175	0.180	0.191	0.178	0.177	0.186	0.152
Ningxia	0.234	0.208	0.273	0.225	0.258	0.241	0.350	0.349	0.306	0.242	0.209
Qinghai	0.079	0.085	0.077	0.080	0.079	0.088	0.083	0.091	0.096	0.082	0.086
Shaanxi	0.357	0.301	0.401	0.324	0.261	0.269	0.338	0.304	0.315	0.246	0.224
Shandong	1.197	1.309	1.531	1.315	1.113	1.188	1.323	1.232	1.191	0.870	0.592
Shanghai	0.874	0.744	0.883	0.828	0.588	0.656	0.544	0.460	0.507	0.325	0.202
Shanxi	0.748	0.806	0.928	0.779	0.593	0.612	0.789	0.703	0.661	0.614	0.493
Sichuan	0.429	0.429	0.513	0.376	0.415	0.427	0.394	0.293	0.350	0.198	0.123
Taiwan	0.089	0.071	0.081	0.085	0.074	0.090	0.074	0.055	0.086	0.052	0.051
Tianjin	1.197	1.344	1.577	1.132	1.217	1.289	1.176	1.140	1.150	1.005	0.708
XinjiangU.	0.073	0.074	0.087	0.093	0.088	0.094	0.090	0.090	0.111	0.101	0.084
Xizang/Tibet	0.080	0.097	0.086	0.091	0.094	0.111	0.096	0.097	0.083	0.087	0.087
Yunnan	0.140	0.159	0.182	0.147	0.144	0.153	0.149	0.131	0.127	0.100	0.085
Zhejiang	0.383	0.337	0.452	0.395	0.297	0.316	0.403	0.258	0.321	0.212	0.158
P.R.China	0.397	0.392	0.444	0.373	0.330	0.342	0.358	0.335	0.332	0.280	0.225

**Table A2 Annual NO<sub>x</sub> emissions (Gg N/year) per province in the domain of DECSO (in parenthesis the fraction of provincial area considered) derived from OMI observations**

Province	2007	2008	2009	2010	2011	2012	2013	2014	2015
Anhui	167	169	187	224	242	292	288	282	215
Beijing	91	62	90	107	88	80	89	74	64
Chongqing	54	57	70	75	87	95	96	100	70
Fujian	96	114	100	114	161	162	153	167	137
Gansu (61%)	31	38	37	42	61	73	60	78	52
Guangdong	383	383	331	341	371	413	360	374	331
Guangxi	118	148	118	145	152	224	224	200	157
Guizhou	107	130	142	154	131	194	191	180	122
Hainan	8	13	11	17	23	22	25	37	30
Hebei	427	423	403	515	563	543	544	511	436
Heilongjiang(74%)	33	36	30	25	43	54	40	49	31
Henan	334	347	370	445	470	481	491	433	315
Hubei	135	140	144	186	248	263	233	270	199
Hunan	109	112	124	162	163	216	184	218	187
Jiangsu	374	344	344	421	470	433	428	445	365
Jiangxi	51	58	65	73	85	105	111	150	112
Jilin	30	22	18	20	45	50	43	48	43
Liaoning	122	128	124	169	205	225	178	199	173
NeiMongol (83%)	98	116	117	156	215	215	169	142	111
Ningxia	30	34	32	41	75	72	66	61	36
Shaanxi	118	118	113	181	216	222	196	208	158
Shandong	464	510	493	629	689	677	731	712	580
Shanghai	96	103	101	95	109	75	83	93	84
Shanxi	292	284	253	328	397	395	373	313	260
Sichuan (51%)	155	158	179	204	232	254	271	280	205
Taiwan	100	106	98	106	113	118	106	114	111
Tianjin	77	86	99	136	152	114	102	97	88
Yunnan (36%)	83	109	105	97	118	159	144	176	126
Zhejiang	243	253	247	270	327	281	294	292	240
EastChina	4332	4502	4454	5382	6150	6402	6179	6201	4941