

Manuscript acp-2016-447

Screening of cloud microorganisms isolated at the puy de Dôme (France) station for the production of biosurfactants
P. Renard, I. Canet, M. Sancelme, N. Wirgot, L. Deguillaume, and A.-M. Delort

Answer to Referee #1

First we would like to thank to the reviewers for their work and interest in our work. We have taken into account their comments to improve the manuscript and answered point by point to their questions. Changes in the manuscript are underlined in yellow.

Anonymous Referee #1

The authors report equilibrium surface tension values for every microorganism in their study. However, it is unclear how they determine when equilibrium has been reached. All of the surface tension time profiles in Figure 2 appear to be decreasing when the measurements were stopped. That is, the reported equilibrium surface tension values are the minimum values for the time profiles given, but may not be if the time profile was extended. In section 2.3, the authors state a 30-minute maximum for surface tension measurements but give no justification for this time frame.

Author's response

You are absolutely right, the minimum of the equilibrium region (σ_{eq}), is difficult to determine experimentally (small variations of surface tension over long timescales) (Nozière et al., 2014). The surface tension decreases asymptotically and the logarithmic scale of the figure 2 is probably misleading, it looks clearer when presented in a non-logarithmic form (see enclosed Figure 2. Time profile of surface tension measurements in a non-logarithmic form.). The overall equilibration time, t_{eq} , is of the order of $2 \times t_m$ (time of meso-equilibrium) (Nozière et al., 2014). After this period, surface tension decreases marginally. According to the measurements we did on longer period (few hours), our overestimation is comprised between 0.1 and 0.2 mN m⁻¹. That is why, above 30 mN m⁻¹, we only give nominal surface tension (Table S2 in the supplementary). Below 30 mN m⁻¹ and above the CMC, surface tension decreases quickly and it is easier to be more accurate.

Finally, we decided to keep our original Figure 2 presented in the logarithmic mode as it is the usual way in publications and allows to show the initial surface tension (σ_0), the surface tensions in the meso-equilibrium (σ_m) and in the equilibrium (σ_{eq}) phase (See Nozière et al 2014 for instance).

Anonymous Referee #1

In Figure 2, the authors show the surface tension time profile for the R2A broth, which is the medium used for all cultures of their isolated microorganisms. However, this may not be a good baseline because incubation period lasts between six to ten days. We accept that the microorganisms are altering the composition of the broth by producing biosurfactants, but they are consuming nutrients in the broth as well. It remains to address how the removal of nutrients would impact the surface tension of the crude extracts.

Author's response

We agree that broth, i.e. carbon sources, influences biosurfactant production. This is true for industrial production of biosurfactants performed in aqueous media with a rich carbon source feedstock, such as carbohydrates, hydrocarbons, fats, and oils. Such enriched broths increase the production. However in our case R2A broth is very poor so when the microorganisms consume the carbon sources, it does not make a great change. We confirmed this point thanks to the following experiments: We did purification of few biosurfactants from microbial R2A cultures, and then we measured the surface tension of these pure compounds both in water and in R2A medium, the differences were marginal (confidential, to be published).

Author's changes

We modified the text as follows:

P 9 line 9: In this collection, we observed 34 strains (7%) that reduce the surface tension of the R2A broth below 30 mN m⁻¹.

Anonymous Referee #1

Furthermore, the authors present a large amount of data regarding the surface tension of crude extracts but do not make a connection to the surface tension of cloud water, which is arguably the basis for this work. Since the authors have already collected the cloud water in order to isolate the microorganisms, it would be useful to also report surface tension values for the cloud water samples as well extend the crude extract results to cloud water.

Author's response

We share your viewpoint; the surface tension of cloud water could have been relevant. Unfortunately, the cloud samplings have been performed before the acquisition of the tensiometer. However, according to the Köhler theory, the surface tension, as well as, the saturation vapor pressure and the CCN diameter, drive the activation of particles into cloud droplets. The activation occurs when the radius of the cloud droplet is minimal (few nm, i.e., wet aerosol) and the concentration of organic compounds, such as biosurfactant, is maximal (Nozière et al., 2014). The effect of surface tension is maximal during the activation. In cloud droplet (few μm), organic compounds are diluted, and biosurfactants are likely under the CMC. Nevertheless, measuring surface tension in concentrated cloud water could be a complementary work, especially since we observed in 300 fold-concentrated rain, a strong decrease of the surface tension (30 mN m^{-1}) (unpublished data).

Here, we demonstrate that bacteria sampled in clouds are able to produce biosurfactants under lab conditions. We are currently isolating and characterizing these biosurfactants. We have identified 11 different structures by mass spectrometry. In the future we want to collect cloud and rain samples and also aerosols and look for these structures in these atmospheric samples (this is what is proposed in the conclusion). This is a long term research plan.

Author's changes

In order to emphasize this point, we have modified the following sentences:

P 13 lines 4: "In conclusion, the results of the present study showed that the microbial strains isolated from cloud waters produce strong biosurfactants under laboratory conditions. The major and most active producers belong to the *Pseudomonas* genus, which is prevalent in cloud water and typically originates from the phyllosphere. Although the presence of surfactants has been shown on aerosols (Nozière et al., 2014), it has not yet been demonstrated in clouds, and the structure of these compounds has not been established. The biosurfactants overproduced by the best producers in the present study will be isolated to analyze their chemical structure. In parallel, the biosurfactants from cloud aerosols and rain samples will also be extracted, and their structural fingerprints will be analyzed and compared with the signatures of microbial surfactants isolated from clouds."

Anonymous Referee #1

Finally, the statistical analysis section did not seem to add much to the paper. The main takeaway was that α -Proteobacteria are efficient biosurfactant producers, which reinforces conclusions from section 3.2. However, the entire analysis seems unsubstantiated. The distinction between air mass origins seems arbitrary. The distinction between chemical compositions is more logical, but the conclusions for that analysis are weaker.

Author's response

Our statistics are based on 480 strains but these strains are grouped into 39 cloud events, thus partially dependent. This sampling was spread over 10 years, and represented with related analyzes, a considerable work and a more than correct observation of Puy de Dôme clouds.

However it is still difficult to make statistics on samples with such intra- and inter-sample variations. For example, in marine clouds, we identified only one strain in few events (e.g., event 29) compared to the 62 strains in the event 54 (see Table S1 in supplementary). This makes our Mann-Whitney and Kruskal-Wallis tests a bit weak.

We could use mixed model. Nevertheless, you are right, these statistics would not add much new, i.e., the correlation of *Pseudomonas* / surface tension. We therefore concluded it would be better to be limited to a high-quality observation.

In conclusion we decided to keep the paragraph “Impact of the origin and chemical composition of clouds on biosurfactant production” to give some general tendency. The obtained results are interesting as they suggest a link between the vegetation origin and the biosurfactant production. This should be studied in more details in the future.

Author's changes

In the abstract, we replaced:

Statistical analyses showed some positive correlations between the origin of air masses and chemical composition of cloud waters with the presence of biosurfactant-producing microorganisms, suggesting a “biogeography” of this production.

by:

We observed some correlations between the chemical composition of cloud water and the presence of biosurfactant-producing microorganisms, suggesting the “biogeography” of this production.

Page 4 line 5: we replaced:

“In order to evaluate the potential correlation between the origin of air masses and composition of cloud waters and the presence of biosurfactant-producing microorganisms, statistical analyses are performed.”

by:

“We observed a potential correlation between the composition of cloud waters and the presence of biosurfactant-producing microorganisms.”

P5 line 14: This text has been deleted:

2.4 Statistical analyses

Herein, we investigate the differences, in terms of impact on the non-normally distributed surface tension, due to the origin of air mass and the chemical composition of clouds using the PAST software version 3.09 (Hammer et al., 2001).

Using a non-parametric method, the Kruskal-Wallis one-way analysis of variance (Siegel, 1956), we compare the distributions of surface tensions between 4 air mass origin sectors: west (W), north-west/north (NW/N), north-east (NE) and south-west/south (SW/S) and between 4 chemical composition groups (Marine, Highly marine, Continental and Polluted). P-value < 0.05 is considered statistically significant.

Mann-Whitney test (Mann and Whitney, 1947), which is a measure of how different two populations are, allows specifying which group dominates, with two-by-two comparison.

Page 10 line 1: we totally rewrote the section 3.3 and replaced by:

3.3 Potential impact of the chemical composition of the clouds on biosurfactant production

In the present study, the screened microbial strains were isolated from 39 cloud events presenting different profiles. Information on the cloud chemical composition and the physicochemical parameters measured at the puy de Dôme station and described in (Deguillaume et al., 2014) is provided on the website of the Observatory of Earth Physics in Clermont-Ferrand (<http://www.obs.univ-bpclermont.fr/SO/beam/data.php>). The main parameters, including pH, SO_4^{2-} , NO_3^- , Cl^- , acetate, formate, oxalate, succinate, malonate, Na^+ , NH_4^+ , Mg^{2+} , K^+ , and Ca^{2+} , are summarized in the Supplemental materials (Table S1). These physico-chemical parameters were used for the ACP analysis as described in Deguillaume et al. (2014). The ACP generated 4 different types of clouds, classified as “highly marine”, “marine”, “continental” and “polluted”. Typically, the more “polluted” clouds have a lower pH and higher concentrations of NH_4^+ , NO_3^- , and SO_4^{2-} . The more “marine” clouds have a higher concentration of NaCl. The 39 cloud events were divided into 2 “highly marine”, 26 “marine”, 8 “continental” and 3 “polluted” clouds (Table S1).




Figure 4 (a). Surface tension (σ) distribution of the 480 strains examined for biosurfactant production according to the physicochemical characteristics of cloud waters (marine, highly marine, continental and polluted). Highlighted in blue, the number of tested strains. Box and whisker plots are shown with the minimal (red) and maximal (green) surface tensions. The orange boxes represent the 25th and 75th percentiles of the measurements (b). Phyla distribution according to the physicochemical characteristics of the cloud waters (marine, highly marine, continental and polluted).

Figure 4a shows the distribution of the surface tensions values (σ) measured from the 480 strains examined for biosurfactant production according to the cloud water chemical composition (marine, highly marine, continental or polluted). A comparison of the distribution of the phyla of the strains in the same cloud events is presented in Figure 4b. The samples from marine clouds constitute the majority of this collection (323/480 strains). We observed a difference between the surface tension values from continental and highly marine strains (medians: 56 and 61 mN m⁻¹, respectively). Highly marine clouds are characterized by the highest minimal surface tension (45 mN m⁻¹, Figure 4a), consistent with the almost complete absence of γ -Proteobacteria, which are the most efficient biosurfactant-producing microorganisms ($\sigma \leq 45$ mN m⁻¹) (1/57 isolates, see Figure 4b). These observations were based on 39 cloud events with 480 different strains, representing, to our knowledge, the largest cloud sample data set studied; this data set is representative of cloud sampling over more than 10 years at the puy de Dôme station. Although it remains difficult to generate statistics on samples with such intra- and inter-sample variations, these results provide a general tendency that could be reinforced and confirmed with more data in the future.

Figure 4, Figure 5a, Figure 6a, Table S3 and Figure S1 have been deleted.

We have kept only Figure 5b and Figure 6b which are now Figure 4(a) and (b) in the revised manuscript, note that the presentation of the data has been modified as suggested by referee 2.

Anonymous Referee #1

The grammatical errors are too numerous to list individually. This paper would greatly benefit from editing by a native English speaker.

Authors: The manuscript has been proof read by ACS services

Page 5, line 2. Might be helpful to keep units consistent with Page 4, line 33. Either g (preferred) or rpm.

Authors: We put these values: 10,480 g / 3 min

Page 5, line 11. Change section number from Roman to Arabic numerals.

Authors: done

Page 9, lines 9-10. This is a misrepresentation because the biosurfactants are reducing the surface tension of the R2A broth, not pure water.

Authors: Changed

Page 9, line 11. I think you mean surface tension values between 30 and 45 mN m⁻¹ not up to 45 mN m⁻¹.

Authors: Yes, we agree, we have changed it to "between 30 and 45 mN m⁻¹"

Page 9, line 18. Third and fourth is clearer than third and last.

Authors: Changed

Page 11, lines 12-14. There is not a significant difference between all four sectors, just between NW/N and the others, according to your supplementary information.

Authors: Actually we have deleted all the data linked to the back-trajectories of the air masses (see answer to referee 2)

Page 14, lines 3-5. Citation for this sentence?

Authors: _ we have added "(Joly et al., 2015 and references hereinafter)".

Manuscript acp-2016-447

Screening of cloud microorganisms isolated at the puy de Dôme (France) station for the production of biosurfactants
P. Renard, I. Canet, M. Sancelme, N. Wirgot, L. Deguillaume, and A.-M. Delort

Answer to Referee #2

First we would like to thank to the reviewers for their work and interest in our work. We have taken into account their comments to improve the manuscript and answered point by point to their questions. Changes in the manuscript are underlined in yellow.

Anonymous Referee #2

English grammar: This paper must be completely proofread by a native English speaker before publication should be considered.

Author's changes

The manuscript has been proof read by ACS services

Anonymous Referee #2

-Clouds: the largest hole in this manuscript is the lack of cloud-water analysis. The authors acceptably demonstrate that biosurfactant producing bacteria exist in their cloud water samples but fail to demonstrate if the bacteria actually have a measurable effect on their collected cloud droplets. Even if the authors are unable to measure surface tension depression in their cloud samples, this should still be noted and contextualized in the manuscript. The paragraph in the discussion section, starting P14-L33, would greatly benefit from this analysis.

Author's response

We deeply agree with this comment. To demonstrate if the bacteria have a measurable effect on the formation of cloud droplets is the critical issue, and the point of further studies.

Nozière et al. (2014) observed strong decreases of surface tension on aerosols: "The first results have shown that these fractions are much more surface-active than expected (σ : 30 mN m⁻¹) and display properties similar to those of biosurfactants such as surfactin or rhamnolipids".

Here, we demonstrate that bacteria sampled in clouds are able to produce biosurfactants under lab conditions. We are currently isolating and characterizing these biosurfactants. We have identified 11 different structures by mass spectrometry. In the future we want to collect atmospheric water samples (rain and cloud) and also aerosols and look for these structures in these atmospheric samples (this is what is proposed in the conclusion). This is a long term research plan.

We would like to add that, although we did not measure surface tension in cloud waters, we recently measured it in concentrated rain (x300) and found a value of 30 mN/m⁻¹.

Author's changes

In order to emphasize this point, we have modified the following sentences:

P 11 Line 7: "In the present study, we showed that under laboratory conditions, the most efficient biosurfactant-producing microorganisms ($\sigma < 45$ mN m⁻¹) belonged to a limited number of bacterial genera (*Pseudomonas* and *Xanthomonas*) from the γ -Proteobacteria class (78%) and a yeast genus (*Udeniomyces*) from the Basidiomycota phylum (11%)."

P 13 lines 4: "In conclusion, the results of the present study showed that the microbial strains isolated from cloud waters produce strong biosurfactants under laboratory conditions. The major and most active producers belong to the *Pseudomonas* genus, which is prevalent in cloud water and typically originates from the phyllosphere. Although the presence of surfactants has been shown on aerosols (Nozière et al., 2014), it has not yet been demonstrated in clouds, and the structure of these compounds has not been established. The biosurfactants overproduced by the best producers in the present study will be isolated to analyze their chemical structure. In parallel, the biosurfactants from cloud aerosols and rain samples will also be extracted, and their structural fingerprints will be analyzed and compared with the signatures of microbial surfactants isolated from clouds."

Anonymous Referee #2

Arbitrary choices: While not debilitating to the paper itself, two arbitrary divisions are made in this manuscript: 1) the surface tension division of Fig.3 ... For the surface tension divisions, the authors cite Baudel et al. (2012) and Ekstrom et al. (2010) saying that their divisions are chosen in a similar way. However, neither Baudel nor Ekstrom divide their data in the same way that the authors here are trying to do. It would be more correct to say that the authors are choosing bins for their samples that match 1-2 bins of previously published works. The other bins are completely arbitrary and the authors do not provide any reasons for why they chose >55, 45-55, 30-45, and <30 mN/m. Some reasoning behind these bins needs to be present in the manuscript.

Author's response

You are right, Ekström et al. (2010) and Baduel et al. (2012) have not, strictly speaking, defined categories. Our categories rather correspond to the values observed by them. As cited thereafter the value of 30 mN m⁻¹ refers to strong biosurfactants, the value of 45 mN m⁻¹ to HULIS. We agree that the value of 55 mN m⁻¹ is not so well defined and is more arbitrary. Note that our comments in the text greatly refer to values < 45 mN m⁻¹.

Furthermore, we agree, to categorize quantitative variables is always somewhat arbitrary but it helps to present the results.

According to Ekström et al. (2010), 30 mN m⁻¹ is the distinct signature of microbial surfactants: *"The very low surface tension values obtained with the aerosol samples were attributed to the presence of biosurfactants, because these compounds are the only natural substances able to have such strong effects on the surface tension"*

"Comparing the curves for the standard compounds with those obtained with the aerosol extracts on Fig. 3 (curves with open circles) clearly show that the latter have the distinct signatures of microbial surfactants: a surface tension below 30 mN m⁻¹ at high concentrations, and a sharp transition characteristic of micelle-forming surfactants".

Baduel et al. (2012) also observed strong decrease of surface tension on their atmospheric aerosol samples between 30 and 45 mN m⁻¹: *"The minimum surface tension obtained from the summer samples was systematically lower (30 mN m⁻¹) than that of the winter samples (35-45 mN m⁻¹)."*

According to Ekström et al. (2010), humic-like substances (HULIS) would only lower the surface tension to 45 mN m⁻¹: *"This implies that only a few tens of μM of biosurfactants would lower the surface tension of water to about 30 mN m⁻¹. By contrast, 20mM (20 g L⁻¹) of HULIS would only lower the surface tension to 45 mN m⁻¹ (Taraniuk et al., 2007), and 10M of malonic acid would lower it to 50 mN/m."*

55 mN m⁻¹ is probably the most arbitrary limit; it approximates the first surface tension values measured on aerosols filter samples: *"So far, the only way to perform such measurements is with aerosol filter samples, which are extracted, and the surface tension of the extracts measured with a tensiometer. The first studies using these methods reported surface tensions between 52 mN m⁻¹ (Mircea et al., 2005) and 60 mN m⁻¹ (Capel et al., 1990; Facchini et al., 1999, 2000; Hitzenberger et al., 2002; Decesari et al., 2005; Mircea et al., 2005). The small amounts of material on the filters made these methods challenging. An additional drawback was that the extraction, usually in water, was not specific to surfactants, leading to mixtures where the contribution of the surfactants was underestimated."* (Baduel et al., 2012).

Author s' changes

According to the reviewer remarks we have modified the text as follows:

P 9 line 7: we deleted this sentence: "These 4 categories are chosen in a similar way to Baduel et al. (2012) and Ekström et al. (2010)."

P 9 line 8: we modified the text as follow:

"The first category ($\sigma \leq 30$ mN m⁻¹) is rare among man-made surfactants and is typical of surfactants of biological origin (Christofi and Ivshina, 2002). In this collection, we observed 34 strains (7%) that reduce the surface tension of the R2A broth below 30 mN m⁻¹. These strains exclusively belonged to the genera *Pseudomonas* and *Xanthomonas* (γ -Proteobacteria, Fig. 1b). The second category corresponded to surface tension values between 30 and 45 mN m⁻¹. The 55 mN m⁻¹ limit is often considered the threshold in terms of the surface tension decrease originating from HULIS (humic like substances) (Kiss et al., 2005; Taraniuk et al., 2007). We observed only 30 strains (6%) in this second category. In summary, from the first two categories ($\sigma \leq 45$ mN m⁻¹), although new phyla were observed in the second category, the phylum distribution of the most efficient biosurfactant-producing microorganisms remains largely dominated by γ -Proteobacteria (78% of all strains) and more moderately by Basidiomycota (11%) (Fig. 3). Notably, the two other major taxa of all studied strains, Actinobacteria and α -Proteobacteria, almost completely disappear in these categories. The third and fourth categories ($45 < \sigma \leq 55$ and $\sigma > 55$ mN m⁻¹) represented 28 and 59% of the collection, respectively. The 55 mN m⁻¹ limit is relatively arbitrary but approximates the first surface tension values measured on the aerosol filter samples (Baduel et al., 2012; Capel et al., 1990; Decesari et al., 2005; Facchini et al., 1999, 2000; Hitzenberger et al., 2002; Mircea et al., 2005)."

Anonymous Referee #2

-Arbitrary choices: While not debilitating to the paper itself, two arbitrary divisions are made in this manuscript: [...] 2) the source region division of Fig. 4 For the source region divisions, the authors present no reasoning for dividing the air masses into marine/highly marine/etc. From Table S1, it is impossible to tell why marine and highly marine are split. Fig. S1 suggests the split is because of time over open ocean but the authors need to be explicit here.

I would also argue that Fig. 4 adds nothing to the manuscript and should be replaced with the HYSPLIT trajectories and some additional meteorological statistics (e.g. wind direction histogram for the sampling site for both cloud-sampling days and non-sampling days). There is also no reason to not show the air mass height results from HYSPLIT.

Author's response

We understand from the reviewer's remarks that our text was not clear and did not give enough details to be understandable.

The different categories as described in Figure 4 are based on the paper of Deguillaume et al. (2014). In this paper the air masses are defined according two different types of criteria:

1) Their back trajectories : They are calculated using the HYSPLIT (Hybrid Single-Particle Lagrangian Integrated Trajectory) model with the GDAS1 meteorological data archive and default settings (Draxler and Rolph, 2010). Considering 10 years of monitoring at the puy de Dôme station, Deguillaume et al (2014) divided France in four sectors to classify these back trajectories crossing France (West, North East, NorthWest/North, SouthWest/West).

2) The chemical content of cloud water: The physicochemical parameters presented in **Table S1** (pH, SO_4^{2-} , NO_3^- , Cl^- , Acetate, Formate, Oxalate, Succinate, Malonate, Na^+ , NH_4^+ , Mg^{2+} , K^+ , Ca^{2+}) are used to make an ACP analysis as described in Deguillaume et al. (2014). This ACP gives 4 different groups which have been named “highly marine”, “marine”, “continental” and “polluted”. Typically, the more “polluted” are the clouds, the lower is the pH and the higher are the concentrations of NH_4^+ , NO_3^- , SO_4^{2-} . The more “marine” are the clouds, the higher is the concentration of NaCl.

In our opinion the second type of categories based of chemical measurements is more accurate as it reflects the whole history of the clouds integrating their eventual complex trajectories.

Author's changes

Therefore, in the revised manuscript we deleted all the data linked to the back trajectories and only kept the “highly marine”, “marine”, “continental” and “polluted” categories. This includes deletion of “W, NE, NW, NW/N and SW/W” in Tables S1 and S2, of Table S3, of Figure S1, of Figure 4, of Figure 5(a) and Figure 6 (a).

Figures 5b and Figure 6b are now Figures 4(a) and (b) in the revised manuscript.

P 10, line 11, this text was deleted: “Cloud events are first classified according to the air mass origin - *i.e.*, west (W), north-west/north (NW/N), north-east (NE) and south-west/south (SW/S) - determined from backward trajectories (see Fig. 4 and Fig. S1 in the supplement). Second, cloud events are classified according to the physicochemical characteristics of cloud waters (Marine, Highly marine, Continental and Polluted, see Fig S1 and Table S1 in the supplement) as described by Deguillaume et al. (2014).

Figure 4 shows that 18 events come from W consisting of 2 Highly marine, 14 Marine and 2 Continental cloud events, 13 events from NW/N with 10 marine, 2 continental and 1 polluted cloud events, 3 events from NE with 2 polluted cloud events and the other continental and, from the SW/S, 5 events of which 2 marine and 3 continental cloud events.”

P 11 line 10, this text was deleted: Figure 5a shows the distribution of surface tensions values (σ) measured from the 480 strains tested for biosurfactant production, according to the air mass origins (4 sectors: W, NW/N, NE and SW/S). Samples from west sector constitute the great majority of our collection (318/480 strains). From statistical analysis, we observe a significant difference (Kruskal-Wallis p-value: 0.0049 << 0.05) in the distribution of biosurfactant-producing microorganisms between the NW/N sectors and the others (W, NE and SW/S). The Mann-Whitney test (see details in the Supplement, Table S3) allows us to attribute this difference to the NW/N sector with a surface tension median (53 mN m^{-1}) significantly lower (Mann-Whitney p-values < 0.05) than the other three sectors (medians: 59, 60 and 61 mN m^{-1} , for W, NE and SW/S sectors, respectively). This difference cannot be completely attributed to the differences in the phyla distribution within the different air masses (Fig. 6). Indeed, as shown before, the most efficient biosurfactant-producing microorganisms belong to γ -Proteobacteria class, which represents 23% of all strains, but its distribution regarding the air mass origin sectors remains unselective (26, 28, 3 and 14% for W, NW/N, NE and SW/S sectors, respectively). The difference in the distribution of biosurfactant-producing microorganisms between the four sectors is rather due to the proportion of the most efficient biosurfactant-producing microorganisms ($\sigma \leq 45 \text{ mN m}^{-1}$) amongst the γ -Proteobacteria class (Fig. 6). In the NW/N sector, most

efficient biosurfactant-producing microorganisms account for 68% of γ -Proteobacteria (13/19 isolates), against 40% in the other three sectors (37/93 isolates). No such difference amongst the γ -Proteobacteria class is observed in the chemical composition groups.

P 10 line 6, We have added these sentences in the modified manuscript:

"The main parameters, including pH, SO_4^{2-} , NO_3^- , Cl^- , acetate, formate, oxalate, succinate, malonate, Na^+ , NH_4^+ , Mg^{2+} , K^+ , and Ca^{2+} , are summarized in the Supplemental materials (Table S1). These physico-chemical parameters were used for the ACP analysis as described in Deguillaume et al. (2014). The ACP generated 4 different types of clouds, classified as "highly marine", "marine", "continental" and "polluted". Typically, the more "polluted" clouds have a lower pH and higher concentrations of NH_4^+ , NO_3^- , and SO_4^{2-} . The more "marine" clouds have a higher concentration of NaCl. The 39 cloud events were divided into 2 "highly marine", 26 "marine", 8 "continental" and 3 "polluted" clouds (Table S1)."

Anonymous Referee #2

Furthermore, the analysis starting on page 11, which attributes statistical difference between the air mass divisions (one of which has 3 samples!), seems weak. I would suggest that there are stronger ways to segregate the air masses given the Table S1. I believe the whole analysis should be rerun using the chemical speciation data.

Shorter Comments:

-Mann-Whitney and KW ANOVA – can the authors comment on how appropriate it is to assume statistical independence for air masses that, though they are measured at the sampling site as from 2 different directions, shared the same path as few as 5 days prior to sampling?

-Figure 5 needs to be replaced with a proper box and whiskers plot or some other way to judge what the real spread in the data looks like (possibly note the std. dev.?)

Author's response

As explained above, Deguillaume et al (2014) classifies clouds according to two different criteria: back trajectories and physico-chemical parameters. In our opinion the second type of categories based of chemical measurements is more accurate as it reflects the whole history of the clouds integrating their eventual complex trajectories. Therefore, in the revised manuscript we have deleted all the data and text linked to the back trajectories and only kept the "highly marine", "marine", "continental" and "polluted" categories.

Our statistical analyses are based on 39 cloud events with 480 different strains. This represents, to our knowledge, the largest data set on cloud samples ever studied; it is representative of cloud sampling over more than 10 years at the puy de Dôme station. These cloud events when classified according to ACP analysis (Deguillaume et al. 2014) are independent and can be compared. The only problem is that it is still difficult to make statistics on samples with such intra- and inter-sample variations (not because there is only 3 samples for example). For example, in marine clouds, we identified only one strain in few events (e.g., event 29) compared to the 62 strains in the event 54 (see Table S1 in supplementary). This makes our Mann-Whitney and Kruskal-Wallis tests a bit weak.

We have therefore concluded it would be better to be limited to a high-quality observation rather than making questionable statistics, which would not adding much new, i.e., the correlation of *Pseudomonas* / surface tension.

In conclusion we decided to keep the paragraph "Impact of the origin and chemical composition of clouds on biosurfactant production" to give some general tendency. The obtained results are interesting as they suggest a link between the vegetation origin and the biosurfactant production. This should be studied in more details in the future.

Author's changes

In the abstract, we replaced:

Statistical analyses showed some positive correlations between the origin of air masses and chemical composition of cloud waters with the presence of biosurfactant-producing microorganisms, suggesting a “biogeography” of this production.

by:

We observed some correlations between the chemical composition of cloud water and the presence of biosurfactant-producing microorganisms, suggesting the “biogeography” of this production.

Page 4 line 5: we replaced:

“In order to evaluate the potential correlation between the origin of air masses and composition of cloud waters and the presence of biosurfactant-producing microorganisms, statistical analyses are performed.”

by:

"We observed a potential correlation between the composition of cloud waters and the presence of biosurfactant-producing microorganisms."

P5 line 14: This text has been deleted:

2.4 Statistical analyses

Herein, we investigate the differences, in terms of impact on the non-normally distributed surface tension, due to the origin of air mass and the chemical composition of clouds using the PAST software version 3.09 (Hammer et al., 2001).

Using a non-parametric method, the Kruskal-Wallis one-way analysis of variance (Siegel, 1956), we compare the distributions of surface tensions between 4 air mass origin sectors: west (W), north-west/north (NW/N), north-east (NE) and south-west/south (SW/S) and between 4 chemical composition groups (Marine, Highly marine, Continental and Polluted). P-value < 0.05 is considered statistically significant.

Mann-Whitney test (Mann and Whitney, 1947), which is a measure of how different two populations are, allows specifying which group dominates, with two-by-two comparison.

Page 10 line 1: we totally rewrote the section 3.3 and replaced by:

3.3 Potential impact of the chemical composition of the clouds on biosurfactant production

In the present study, the screened microbial strains were isolated from 39 cloud events presenting different profiles. Information on the cloud chemical composition and the physicochemical parameters measured at the puy de Dôme station and described in (Deguillaume et al., 2014) is provided on the website of the Observatory of Earth Physics in Clermont-Ferrand (<http://www.obs.univ-bpclermont.fr/SO/beam/data.php>). The main parameters, including pH, SO_4^{2-} , NO_3^- , Cl^- , acetate, formate, oxalate, succinate, malonate, Na^+ , NH_4^+ , Mg^{2+} , K^+ , and Ca^{2+} , are summarized in the Supplemental materials (Table S1). These physico-chemical parameters were used for the ACP analysis as described in Deguillaume et al. (2014). The ACP generated 4 different types of clouds, classified as “highly marine”, “marine”, “continental” and “polluted”. Typically, the more “polluted” clouds have a lower pH and higher concentrations of NH_4^+ , NO_3^- , and SO_4^{2-} . The more “marine” clouds have a higher concentration of NaCl. The 39 cloud events were divided into 2 “highly marine”, 26 “marine”, 8 “continental” and 3 “polluted” clouds (Table S1).




Figure 4 (a). Surface tension (σ) distribution of the 480 strains examined for biosurfactant production according to the physicochemical characteristics of cloud waters (marine, highly marine, continental and polluted). Highlighted in blue, the number of tested strains. Box and whisker plots are shown with the minimal (red) and maximal (green) surface tensions. The orange boxes represent the 25th and 75th percentiles of the measurements (b). Phyla distribution according to the physicochemical characteristics of the cloud waters (marine, highly marine, continental and polluted).

Figure 4a shows the distribution of the surface tensions values (σ) measured from the 480 strains examined for biosurfactant production according to the cloud water chemical composition (marine, highly marine, continental or polluted). A comparison of the distribution of the phyla of the strains in the same cloud events is presented in Figure 4b. The samples from marine clouds constitute the majority of this collection (323/480 strains). We observed a difference between the surface tension values from continental and highly marine strains (medians: 56 and 61 mN m⁻¹, respectively). Highly marine clouds are characterized by the highest minimal surface tension (45 mN m⁻¹, Figure 4a), consistent with the almost complete absence of γ -Proteobacteria, which are the most efficient biosurfactant-producing microorganisms ($\sigma \leq 45$ mN m⁻¹) (1/57 isolates, see Figure 4b). These observations were based on 39 cloud events with 480 different strains, representing, to our knowledge, the largest cloud sample data set studied; this data set is representative of cloud sampling over more than 10 years at the puy de Dôme station. Although it remains difficult to generate statistics on samples with such intra- and inter-sample variations, these results provide a general tendency that could be reinforced and confirmed with more data in the future.

Figure 4, Figure 5a, Figure 6a, Table S3 and Figure S1 have been deleted.

We have kept only Figure 5b and Figure 6b which are now Figure 4(a) and (b) in the revised manuscript, note that the presentation of the data has been modified as suggested by the referee (whiskers plots).

Anonymous Referee #2

Paragraph starting P15, L26 – I am generally a proponent of contextualizing your work but I am not convinced this paragraph adds anything to the manuscript. It should either be removed or condensed and moved to the introduction.

Author's response

We agree with the referee that this paragraph is a bit out of scope of this paper, therefore it has been deleted.

Author's changes

Page 15 line 26, this paragraph has been deleted:

To our knowledge, research on the potential impact of biosurfactants on human health due to their presence in atmospheric waters remains marginal, compared to terrestrial or aquatic ecosystem studies (Olkowska et al., 2014). Aerosols are now well-known to represent a major concern for the populations as shown by epidemiology studies (Bernstein et al., 2004; Dominici et al., 2014; Pope, 2000). The toxicological impact is inversely proportional to the particle size (Nel, 2005; Novák et al., 2014). For example, fine particles ($PM_{1-2.5}$) reach lung alveoli and ultrafine particles ($PM_{0.1}$), once in the lung, would readily pass into the bloodstream and cause a direct insult to the cardiovascular system and other organs (Moshhammer and Neuberger, 2003; Polichetti et al., 2009). The composition of the aerosols is also of major importance and should not be underestimated (Brimblecombe and Latif, 2004; Škarek et al., 2007). Regarding more precisely surface-active organic aerosols, few reports are devoted to synthetic molecules. Thus, Poulsen et al. (2000) suggested that molecules with surfactant properties could interfere in the immunological pathways, which could explain the increase of allergic diseases in industrialized societies. At high concentration in the atmosphere, surfactants can lead to asthma, can disrupt the stability of human respiratory systems, as well as can cause dry eyes allergies (Ahmad et al., 2009; Xinxin et al., 2016). Rhamnolipids inhibit ciliary function and produce damage to the human bronchial epithelium (Abdel-Mawgoud et al., 2011). Surfactants, by lowering the surface tension of tear films of the eyes could also be at the origin of dry eyes sensation (Vejrup and Wolkoff, 2002). Hence, biosurfactants present on aerosols could have a double impact on human cells, first because they could destroy cell membranes of the host, second because they could concentrate and dissolve toxic pollutants and help their penetration within the host cells. Further studies are needed to better evaluate the impact of surfactant on human health.

Anonymous Referee #2

Minor comments:

- S2.2, P4, L23-25 : Given that most of the people reading ACP are not biologists, a one sentence explanation on why those three specific agars were chosen should be added to the text. Also, TSA should be defined.

-P5, L11: Should read “3.2” not “III.2”

Author's changes

- Page 4 line 18, the text has been modified as follows:

"Triplicate volumes of 0.1 mL of cloud water were plated onto R2A agar growth medium (Reasoner and Geldreich, 1985; DIFCO™), and eventually onto R2A medium supplemented with NaCl 20 g L⁻¹ and King's B (King et al., 1954), Sabouraud (DIFCO™) and TSA (Trycase Soy Agar, DIFCO™) media. The plates were incubated at 17°C or 5°C under aerobic-dark conditions until the appearance of colonies (typically 6 days

at 17°C or 10 days at 5°C) (Vařtilingom et al., 2012). R2A medium is a poor medium initially developed to isolate microorganisms from tap water and is well adapted to cloud samples, which are also poor. The addition of NaCl to R2A favors the selection of marine microorganisms; King's B medium is selective for *Pseudomonas* strains, while Sabouraud medium is selective for yeast strains."

- Page 5, Line 1, Should read "3.2" not "III.2":

Done

Screening of cloud microorganisms isolated at the puy de Dôme (France) station for the production of biosurfactants

Pascal Renard^{1,2}, Isabelle Canet¹, Martine Sancelme¹, Nolwenn Wirgot^{1,2}, Laurent Deguillaume³, Anne-Marie Delort^{1,2}

¹ Institut de Chimie de Clermont-Ferrand, Université Clermont Auvergne, Université Blaise Pascal, BP 10448, F-63000 Clermont-Ferrand, France

² CNRS, UMR 6296, ICCF, F-63171 Aubière, France

³ Laboratoire de Météorologie Physique/OPGC, Université Clermont Auvergne, Université Blaise Pascal, BP 10448, F-63000 Clermont-Ferrand, France

Correspondence to: Anne-Marie Delort (A-Marie.Delort@univ-bpclermont.fr)

Abstract. A total of 480 microorganisms collected from 39 clouds sampled at the puy de Dôme station (alt. 1465 m, 45°46'19" N, 2°57'52" E, Massif Central, France) were isolated and identified. This unique collection was screened for biosurfactant (surfactants of microbial origin) production by measuring the surface tension (σ) of the crude extracts, comprising the supernatants of the pure cultures, using the pendant drop technique. The results showed that 41% of the tested strains were active producers ($\sigma < 55 \text{ mN m}^{-1}$), 7% being extremely active ($\sigma < 30 \text{ mN m}^{-1}$). The most efficient biosurfactant producers ($\sigma < 45 \text{ mN m}^{-1}$) belong to a few bacterial genera (*Pseudomonas* and *Xanthomonas*) from the γ -Proteobacteria class (78%) and a yeast genus (*Udeniomyces*) from the Basidiomycota phylum (11%). Some *Bacillus* strains from the Firmicutes phylum were also active but represented a small fraction of the collected population. Strains from the Actinobacteria phylum in the collection examined in the present study showed moderate biosurfactant production ($45 < \sigma < 55 \text{ mN m}^{-1}$). *Pseudomonas* (γ -Proteobacteria), the most frequently detected genus in clouds, with some species issued from the phyllosphere, was the dominant group for the production of biosurfactants. We observed some correlations between the chemical composition of cloud water and the presence of biosurfactant-producing microorganisms, suggesting the “biogeography” of this production. Moreover, the potential impact of the production of biosurfactants by cloud microorganisms on atmospheric processes and human health is discussed.

Keywords: Biosurfactants, cloud condensation nuclei, microorganisms

1 Introduction

Atmospheric aerosol particles act as cloud condensation nuclei (CCN) upon which liquid droplets can form. More aerosols increase the concentration of smaller droplets, leading to a brighter cloud (Twomey effect). However, owing to their complexity, all aerosol–cloud–interactions (ACI) can amplify or dampen this effect. Therefore, ACI, particularly CCN activation, still account for major uncertainties in global climate and future climate change predictions (Boucher et al., 2013).

Among organic aerosols, water soluble organic compounds (WSOC) represent a significant fraction of the tropospheric aerosol mass (Kanakidou et al., 2005; Murphy et al., 2006; Saxena and Hildemann, 1996; Zenchelsky and Youssefi, 1979; Zhang et al., 2007) and constitute a complex mixture of neutral and acidic polar organic compounds (Decesari et al., 2001). As some WSOC are amphipathic compounds, these aerosols can act as surfactants (surface-active agents) by creating a partition between the droplet gas-liquid interface and the bulk volume. Seidl and Hänel (1983) were among the first to estimate concentrations of surface-active soluble substances on rainwater and wet aerosol by measuring the lowering of the surface tension. Capel et al. (1990) correlated the surface tension of fog samples with the dissolved organic carbon content. Hitzenberger et al. (2002) observed slight reductions in the surface tension for most of the 23 cloud water samples collected in mountainous and sparsely populated areas.

Some groups (Decesari et al., 2005; Facchini et al., 1999) have renewed research on surface-active compounds in the atmosphere with a great deal of speculation about the potential impact of these compounds on the climate (Brimblecombe and Latif, 2004). Indeed, surfactants affect cloud droplet growth in two main ways according to the Köhler equation (Köhler, 1936): by increasing soluble mass (“Raoult term”) and by decreasing cloud droplet surface tension (“Kelvin term”) (Decesari et al., 2003; Facchini et al., 1999; Lance et al., 2004; Mircea et al., 2002; Rodhe, 1999; Shulman et al., 1996). Thus, considering both the solute concentration increase and the surface tension decrease, Mircea et al. (2002) calculated a substantial attenuation of the aerosol critical supersaturation, revealing a significant increase in the CCN number concentration. By adding surfactant to the gas-aerosol interface, Sareen et al. (2013) assessed significant enhancements in CCN activity (up to 7.5% reduction in critical dry diameter for activation), which for ambient aerosol would lead to a 10% increase in the cloud droplet number concentration. Facchini et al. (1999) estimated that a population rise resulting from surfactants might result, in all stratus clouds, in a 1% increase in albedo and, subsequently, in a calculated global radiative forcing of -1 W m^{-2} .

In addition to the impact of surfactants on CCN activity *via* both Raoult and Kelvin terms, a third effect should be considered. Surfactants can either enhance or slow down the transfer of water across the surface according to the hydrophilic or hydrophobic nature of this aerosol organic coating (Aumann and Tabazadeh, 2008; Chakraborty and Zachariah, 2011; Feingold and Chuang, 2002; Rudich, 2003). These organic coats are common on aerosol particles and might retard the evaporation of molecules present in the water phase, reduce gas transfer, influence chemical reactions, and alter absorption or reflection properties of aerosols (Clifford et al., 2007; Decesari et al., 2003; Gill et al., 1983; Gilman and Vaida, 2006). Nenes et al. (2002) examined the sensitivity of the cloud droplet number concentration to different chemical factors as the dissolution of soluble gases and solutes or formation of organic films at the droplet surface, demonstrating that these chemical effects on droplet activation could be significant. Similarly, the $\cdot\text{OH}$ heterogeneous reactions that occur on organics in the troposphere can significantly modify the hygroscopic properties and CCN ability of these organic surfaces (Bertram et al., 2001; Ellison et al., 1999), thus potentially playing an important role in the Earth’s radiative balance by affecting the properties of clouds, *e.g.*, Twomey effect and cloud life time (Aumann et al., 2010; Nenes et al., 2002; Rodhe, 1999).

Recent experiments have focused on CCN enhancement resulting from biogenic influence (Facchini et al., 2000; O’Dowd et al., 2002; Svenningsson et al., 2006). Thus, some atmospheric organic compounds, such as pinic and

pinonic acids (produced by the oxidation of terpenes in organic vapors released from the canopy) have been implicated in the reduction of the surface tension of water, even at extremely low concentrations (Li et al., 2010; O'Dowd et al., 2002). Another important class of hydrophobic WSOC is humic-like substances (HULIS). These complex mixtures of high molecular weight compounds can depress surface tension in fog water samples by 15-20% (Decesari et al., 2003; Dinar et al., 2006; Facchini et al., 2000).

Nevertheless, although a number of organic surface-active compounds have been detected in aerosol particles and cloud droplets, a large fraction of WSOC remains poorly characterized (Herckes et al., 2013). Moreover, due to their limited concentrations in aerosols, it remains unknown whether atmospheric organic surfactants decrease the surface tension of atmospheric water or contribute to the CCN properties of atmospheric particles (McFiggans et al., 2006).

In the last decade, a few studies have identified strong organic surfactants in atmospheric aerosols (Baduel et al., 2012). Their exceptional tension-active properties suggested that these compounds could be "biosurfactants" of microbial origin that could affect cloud formation (Ekström et al., 2010; Nozière et al., 2014). Biosurfactants are secondary metabolites produced by microorganisms, including low molecular biosurfactants (mainly glycolipids and lipopeptides) and high molecular mass biosurfactants (polysaccharides, proteins, lipopolysaccharides, lipoproteins or complex mixtures of these polymers) (Gautam and Tyagi, 2006; Rosenberg and Ron, 1999). These amphiphilic compounds reduce the surface tension of atmospheric water below 30 mN m^{-1} (*i.e.*, $> -40\%$ compared with pure water) and at concentrations 5 or 6 orders of magnitude lower than organic acids (Ekström et al., 2010). In comparison, HULIS decreased the surface tension in fog water samples by 20% at 100 mg C L^{-1} (Facchini et al., 2000).

Baduel et al. (2012) measured the surface tensions in summer samples, which would be consistent with the high biogenic activity observed during this season. Ahern et al. (2007) showed that fluorescent pseudomonads isolated from clouds and rainwater produce biosurfactants. This study was the first and sole report on the exploration of the potential production of biosurfactants by microorganisms isolated from the cloud environment. Biosurfactants could be both directly issued from the Earth's surface during aerosolization or directly produced in cloud waters. However, a multitude of bacteria, fungi and yeasts display metabolic activities in clouds (Amato et al., 2005, 2007a; Hill et al., 2007; Sattler et al., 2001; Vaitilingom et al., 2012, 2013). These microorganisms survive and resist atmospheric stresses (Delort et al., 2010; Joly et al., 2015).

Poorly considered in the atmosphere, biosurfactants have been extensively studied in soil- and plant-associated environments. Indeed, biosurfactant-producing bacteria in both undisturbed and contaminated soils have been well characterized (Bodour et al., 2003; Raaijmakers et al., 2010). Biosurfactants have been investigated for their capacity to remove heavy metals, and bioremediation is one of the main industrial applications of biosurfactants (Banat et al., 2010; Mulligan, 2009). Moreover, in terms of microbial life and activity, biosurfactants play a key role in bacterial cell motility, organic compound solubilization, microbial biofilm formation and disruption or anti-microbial activity (Chrzanowski et al., 2012; D'aes et al., 2010; Mann and Wozniak, 2012; Raaijmakers et al., 2010; Ron and Rosenberg, 2001).

Within a project aimed at examining atmospheric surfactants and characterizing their effects on cloud droplet formation, we focused on the biosurfactant-producing microorganisms present in atmospheric waters. Cloud

water samples are collected at the puy de Dôme station (France) belonging to the GAW (Global Atmosphere Watch) stations network. A total of 480 bacterial and yeast strains were isolated and identified. This unique collection of microorganisms was screened to identify biosurfactant-producing microorganisms. The surface tension of crude extracts, comprising supernatants of the pure culture, was determined using the pendant drop technique (Hansen and Rødsrud, 1991). We observed a potential correlation between the composition of cloud waters and the presence of biosurfactant-producing microorganisms. Finally, we discuss the potential impact of the production of biosurfactants by cloud microorganisms on atmospheric physicochemical processes.

2 Materials and Methods

2.1 Cloud sampling and physicochemical characterization of the cloud water samples

Cloud water samples were collected using a cloud droplet impactor sterilized by autoclaving and installed on the summit of the puy de Dôme Mountain (1465 m above sea level, 45°46'19"N, 2°57'52"E, Massif Central). Non-precipitating and non-convective cloud samples were collected. The experiments were conducted using the 480 microbial strains collected during 39 cloud events from 2004 to 2014. The physicochemical content of the aqueous cloud samples was characterized (concentrations of organic acids, inorganic ions and pH, see Table S1 in the supplement). Details about the sampling site, instrumentation and procedures for cloud sampling as well as the methods for the chemical analysis of cloud water samples, are provided in Deguillaume et al. (2014).

2.2 Isolation and identification of microorganisms from cloud waters

Triplicate volumes of 0.1 mL of cloud water were plated onto R2A agar growth medium (Reasoner and Geldreich, 1985; DIFCO™), and eventually onto R2A medium supplemented with NaCl 20 g L⁻¹ and King's B (King et al., 1954), Sabouraud (DIFCO™) and TSA (Trycase Soy Agar, DIFCO™) media. The plates were incubated at 17°C or 5°C under aerobic-dark conditions until the appearance of colonies (typically 6 days at 17°C or 10 days at 5°C) (Vaïtilingom et al., 2012). R2A medium is a poor medium initially developed to isolate microorganisms from tap water and is well adapted to cloud samples, which are also poor. The addition of NaCl to R2A favors the selection of marine microorganisms; King's B medium is selective for *Pseudomonas* strains, while Sabouraud medium is selective for yeast strains.

Representative colonies were selected based on colony morphology and pigment production. The isolates obtained in pure cultures (R2A, 17°C) were stored in 10% (v/v) glycerol at -80°C. The strains were identified by ribosomal RNA gene sequencing (16S or 26S rRNA gene sequences for bacteria and yeasts, respectively). A complete description of the methods of identification is available in Vaïtilingom et al. (2012).

2.3 Surface tension measurements

Strains from the glycerol stocks were used to inoculate R2A broth in 96 deep-well plates (500 µL/well). The plates were incubated at 17°C under agitation for 5 days, followed by centrifugation (3000 g/20 min). The supernatants were transferred into 1-mL microtubes and stored at -30°C until subsequent surface tension measurements. The thawed samples were centrifuged (10,480 g/3 min) just prior to surface tension measurements.

All surface tension measurements were performed using the pendant drop method with an OCA 15 Pro tensiometer (Data Physics, Germany). The camera analyzes the pendant drop profile of the crude extract. A dosing needle with a 1.65-mm outside diameter was used, producing drops of 12 μL . The software fits this latter measurement to the Young-Laplace equation and averages out surface tension from all measurements (Hansen and Rødsrud, 1991). The measurements were obtained at 295 K every second. The tensiometer was calibrated using Milli-Q water. The uncertainty of the instrument was $\pm 0.01 \text{ mN m}^{-1}$. Each dynamic surface tension curve was measured three times for the most efficient biosurfactant-producing microorganisms, and the measurements displayed $\pm 10\%$ variation. These dynamic surface tension measurements lasted until the equilibrium region is reached (maximum 30 min, see below [section 3.2](#)). Along with the surface tension, each measurement also provided real-time monitoring of the droplet volume, facilitating an assessment of the evaporation. No significant evaporation ($< 5\%$) was observed during the experiments (Fig. 2).

3 Results

3.1 Identification of cloud microbial isolates




Figure 1. (a) Phyla distribution of the 480 strains examined for biosurfactant production. (b) Genera distribution of the most representative strains (85%: α - and γ -Proteobacteria, Actinobacteria and Basidiomycota). The 4 categories of surface tensions ($\sigma \leq 30$, $30 < \sigma \leq 45$, $45 < \sigma \leq 55$ and $\sigma > 55 \text{ mN m}^{-1}$) are indicated in red, orange, yellow and green, respectively.

The identification of the 480 strains (bacteria and yeasts) collected during the 39 cloud events at the puy de Dôme station, together with the values of surface tension obtained from their crude supernatants after 5 days of culture in R2A broth, is described in the Supplemental materials (Table S2).

Figure 1(a) shows the distribution of the different phyla or classes of these microbial isolates. Three phyla of microorganisms were dominant: Proteobacteria (α , β and γ - Proteobacteria), Actinobacteria and Basidiomycota, accounting for 89.6% of the collection, while 2.5% of the latter remain unidentified.

In detail (Fig. 1b), the phylum of Proteobacteria was predominant (220 isolates, 45.8%), particularly α - and γ - Proteobacteria (95 and 112 isolates, 19.8% and 23.3%, respectively). In these latter two classes, the most recurrent strains belonged to the genera *Sphingomonas* (83 isolates) and *Pseudomonas* (78 isolates), respectively. β -Proteobacteria represent only 2.7% of the total. The phylum Actinobacteria (116 isolates, 24.2%) primarily contained strains from the genera *Clavibacter* (25 isolates), *Curtobacterium* (14 isolates), *Frigoribacterium* (18 isolates) and *Rhodococcus* (21 isolates). Notably, the phylum Actinobacteria presents a much greater diversity of genera compared with the other phyla, centered on one dominant genus. Among the bacterial strains, the phyla Bacteroidetes and Firmicutes were also represented, but to a lesser extent (20 and 12 isolates, respectively, 4.2% and 2.5%).

Concerning yeasts, the major group belonged to the phylum Basidiomycota (94 isolates, 19.6%), primarily containing strains from the genera *Cryptococcus* (14 isolates), *Dioszegia* (39 isolates) and *Udeniomyces* (25 isolates). The phylum Ascomycota was also present but with only 6 isolates (1.2%).

Globally, the phylogeny of the isolated strains examined in the present study is consistent with the previously published phylogeny of the strains isolated from the same sampling sites, except for the genus *Bacillus*, which was much less abundant in the selected clouds events (Vaithilingom et al., 2012). Notably, many strains originated from the phyllosphere, consistent with the predominance of the phylum Proteobacteria. These 480 strains therefore constitute a unique collection of cloud microorganisms, representative of a cloud community that can be tested for biosurfactant production.

3.2 Screening for biosurfactant-producing microorganisms




Figure 2. Time profile of the surface tension measurements. In black, R2A broth medium. In purple, strain 14b13 (*Frigoribacterium* sp., Actinobacteria). In blue, strain 26b18 (*Cryptococcus* sp., Basidiomycota). In green, strain 14b02 (*Pseudomonas* sp., Y-Proteobacteria). In orange, strain 59b03 (*Pseudomonas syringae*, Y-Proteobacteria). σ_0 : initial surface tension. Surface tensions in the meso-equilibrium (σ_m) and equilibrium (σ_{eq}) phases.

Figure 2 shows the time profile of the surface tension measurements performed on the culture supernatants of 4 selected microbial strains. As a reference, the surface tension obtained from R2A medium, which served as the culture medium, is also presented. In this case, the observed surface tension (63 mN m^{-1}) remained close to the value of pure water (72.8 mN m^{-1}). As expected, a more or less strong reduction of the surface tension values was observed when the microorganisms produced biosurfactants in the culture medium. This phenomenon is time-dependent, and the time profiles were dependent on the studied strain. These profiles were consistent with those obtained from atmospheric aerosol by Nozière et al. (2014) and are typical of a surface tension dynamic (Hua and Rosen, 1991). Indeed, three distinct kinetic regimes follow each other during the equilibration process: first, a rapid decline of the R2A value to σ_0 occurs, which happens too rapidly to record ($< 0.1 \text{ s}$). This is followed by the meso-equilibrium phase, during which the surface tension decreases to σ_m . Then, the appearance of the equilibrium region occurs, where the minimum, σ_{eq} , is reached. This region corresponds to the saturation of the surface (Γ_∞) with surfactant molecules. Hereafter, the surface tension measurements (σ) are referred to as σ_{eq} .

Monitoring the surface tension over time revealed that the equilibration time (apparent diffusion coefficient) in the pendant drops varied from a few seconds to 30 min, likely depending on the concentration and chemical structure of the expressed biosurfactants, which affect molecular interactions and/or diffusion. For example, from the supernatant of the strains 59b03 and 14b02, two *Pseudomonas* strains (Y-Proteobacteria), the measured

equilibrium surface tensions (σ_{eq}) were close (below 28 mN m^{-1}), while the time profiles were much different, and the equilibration stage occurred at approximately 2 and 10 minutes, respectively.




Figure 3. (a) Surface tension (σ) distribution of the 480 strains examined for biosurfactant production and **(b)** the phylum distribution for the most efficient biosurfactant-producing microorganisms ($\sigma \leq 45 \text{ mN m}^{-1}$).

The 480 strains tested for biosurfactant production (see Supplement Table S2) were differentiated into four main categories according to the measured surface tension ($\sigma \leq 30$, $30 < \sigma \leq 45$, $45 < \sigma \leq 55$ and $\sigma > 55 \text{ mN m}^{-1}$, Fig. 1b and Fig. 3). The first category ($\sigma \leq 30 \text{ mN m}^{-1}$) is rare among man-made surfactants and is typical of surfactants of biological origin (Christofi and Ivshina, 2002). In this collection, we observed 34 strains (7%) that reduce the surface tension of the R2A broth below 30 mN m^{-1} . These strains exclusively belonged to the genera *Pseudomonas* and *Xanthomonas* (γ -Proteobacteria, Fig. 1b). The second category corresponded to surface tension values between 30 and 45 mN m^{-1} . The 55 mN m^{-1} limit is often considered the threshold in terms of the surface tension decrease originating from HULIS (humic like substances) (Kiss et al., 2005; Taraniuk et al., 2007). We observed only 30 strains (6%) in this second category. In summary, from the first two categories ($\sigma \leq 45 \text{ mN m}^{-1}$), although new phyla were observed in the second category, the phylum distribution of the most efficient biosurfactant-producing microorganisms remains largely dominated by γ -Proteobacteria (78% of all strains) and more moderately by Basidiomycota (11%) (Fig. 3). Notably, the two other major taxa of all studied strains, Actinobacteria and α -Proteobacteria, almost completely disappear in these categories. The third and fourth categories ($45 < \sigma \leq 55$ and $\sigma > 55 \text{ mN m}^{-1}$) represented 28 and 59% of the collection, respectively. The 55 mN m^{-1} limit is relatively arbitrary but approximates the first surface tension values measured on the aerosol filter samples (Baduel et al., 2012; Capel et al., 1990; Decesari et al., 2005; Facchini et al., 1999, 2000; Hitzenberger et al., 2002; Mircea et al., 2005). Remarkably, *Pseudomonas* (γ -Proteobacteria) and *Sphingomonas* (α -Proteobacteria), the most frequently observed genera in the clouds (Väitilingom et al., 2012), showed completely different behaviors: *Pseudomonas* provide the most active biosurfactant-producing microorganisms, while almost all *Sphingomonas* are not efficient for the production of biosurfactants.

3.3 Potential impact of the chemical composition of the clouds on biosurfactant production

In the present study, the screened microbial strains were isolated from 39 cloud events presenting different profiles. Information on the cloud chemical composition and the physicochemical parameters measured at the puy de Dôme station and described in (Deguillaume et al., 2014) is provided on the website of the Observatory of Earth Physics in Clermont-Ferrand (<http://www.obs.univ-bpclermont.fr/SO/beam/data.php>). The main parameters, including pH, SO_4^{2-} , NO_3^- , Cl^- , acetate, formate, oxalate, succinate, malonate, Na^+ , NH_4^+ , Mg^{2+} , K^+ , and Ca^{2+} , are summarized in the Supplemental materials (Table S1). These physico-chemical parameters were used for the ACP analysis as described in Deguillaume et al. (2014). The ACP generated 4 different types of clouds, classified as “highly marine”, “marine”, “continental” and “polluted”. Typically, the more “polluted” clouds have a lower pH and higher concentrations of NH_4^+ , NO_3^- , and SO_4^{2-} . The more “marine” clouds have a higher concentration of NaCl. The 39 cloud events were divided into 2 “highly marine”, 26 “marine”, 8 “continental” and 3 “polluted” clouds (Table S1).




Figure 4 (a). Surface tension (σ) distribution of the 480 strains examined for biosurfactant production according to the physicochemical characteristics of cloud waters (marine, highly marine, continental and polluted). Highlighted in blue, the number of tested strains. Box and whisker plots are shown with the minimal (red) and maximal (green) surface tensions. The orange boxes represent the 25th and 75th percentiles of the measurements (b). Phyla distribution according to the physicochemical characteristics of the cloud waters (marine, highly marine, continental and polluted).

Figure 4a shows the distribution of the surface tensions values (σ) measured from the 480 strains examined for biosurfactant production according to the cloud water chemical composition (marine, highly marine, continental or polluted). A comparison of the distribution of the phyla of the strains in the same cloud events is presented in Figure 4b. The samples from marine clouds constitute the majority of this collection (323/480 strains). We observed a difference between the surface tension values from continental and highly marine strains (medians: 56 and 61 mN m^{-1} , respectively). Highly marine clouds are characterized by the highest minimal surface tension

(45 mN m⁻¹, Figure 4a), consistent with the almost complete absence of γ -Proteobacteria, which are the most efficient biosurfactant-producing microorganisms ($\sigma \leq 45$ mN m⁻¹) (1/57 isolates, see Figure 4b). These observations were based on 39 cloud events with 480 different strains, representing, to our knowledge, the largest cloud sample data set studied; this data set is representative of cloud sampling over more than 10 years at the puy de Dôme station. Although it remains difficult to generate statistics on samples with such intra- and inter-sample variations, these results provide a general tendency that could be reinforced and confirmed with more data in the future.

4 Discussion and conclusion

The study of biosurfactant production in the environment has primarily focused on microorganisms isolated from soils, rhizospheres and the phyllosphere or from the marine environment (Bodour et al., 2003; Jackson et al., 2015; Raaijmakers et al., 2010; Satpute et al., 2010). Concerning atmospheric waters, Ahern et al. (2007) reported the presence of biosurfactant-producing bacteria in the air, studying 100 strains isolated from 4 rain and cloud samples. Here, we investigated 480 strains isolated from 39 different cloud events. When we consider that these strains typically produce biosurfactants when the measured surface tension (σ_{eq}) is less than 55 mN m⁻¹, 41% of the tested strains were active, 7% being extremely active ($\sigma < 30$ mN m⁻¹). Although the methods used to evaluate biosurfactant production were different, this result is consistent with that of Ahern et al. (2007), who reported 55% active strains in rain and cloud samples. In the present study, we showed that under laboratory conditions, the most efficient biosurfactant-producing microorganisms ($\sigma < 45$ mN m⁻¹) belonged to a limited number of bacterial genera (*Pseudomonas* and *Xanthomonas*) from the γ -Proteobacteria class (78%) and a yeast genus (*Udeniomyces*) from the Basidiomycota phylum (11%). Some *Bacillus* strains from the Firmicutes phylum were also active but represented a small fraction (3%) of the total population of the cloud collection examined in the present study. Strains from the Actinobacteria phylum were primarily present in the group with moderate biosurfactant production ($45 < \sigma < 55$ mN m⁻¹). In previous studies, *Pseudomonas* (γ -Proteobacteria) and *Bacillus* (Firmicutes) have been reported as high biosurfactant-producing microorganisms; *Acinetobacter* (α -Proteobacteria) has also been frequently reported (Desai and Banat, 1997; Rosenberg and Ron, 1999). For yeasts, the major high biosurfactant-producing genera include *Candida* and *Torulopsis* from phylum Ascomycota (Desai and Banat, 1997; Karanth et al., 1999; Rosenberg and Ron, 1999). In the present study, *Pseudomonas* strains were clearly the dominant group and the most active biosurfactant-producing microorganisms, whereas the *Acinetobacter* genus was absent. This result is highly consistent with studies performed on environmental samples, such as soils (Bodour et al., 2003), plants (D'aes et al., 2010; Raaijmakers et al., 2010), seawater (Cai et al., 2015) and in atmospheric water (Ahern et al., 2007).

Notably, the *Pseudomonas* genus is commonly detected in the phyllosphere, the main source of primary bioaerosols (Amato et al., 2007b; Morris et al., 2014; Våtilingom et al., 2012). Interestingly, biosurfactants play a specific role in the interactions between plants and *Pseudomonas* (D'aes et al., 2010; Raaijmakers et al., 2010). Biosurfactants present versatile functions, including interactions with other organisms (such as antibiotic activity) and modifications of the leaf-surface properties. These surface modifications enable cell mobility, biofilm formation and the colonization of the leaves by these bacteria. Particularly, rhamnolipids are involved in different processes of biofilm formation; the final step involves the release of the planktonic daughter population

(Mann and Wozniak, 2012). This production of biosurfactants could therefore be important for the formation of biofilms on leaf surfaces, facilitating the aerosolization and dispersion of *Pseudomonas* strains in the air.

This aerosolization of *Pseudomonas* strains could explain the correlation observed between the clouds composition and the distribution of biosurfactant-producing microorganisms observed in the present study. It is clear that microbial isolates from highly marine clouds, significantly impacted by the ocean source (an almost total absence of γ -Proteobacteria, see Fig. 4b), are lower biosurfactant producers than microorganisms isolated from continental clouds. Continental clouds, travelling over vegetated zones, thus contain more *Pseudomonas* strains. More generally, the correlation between the different origins of the air masses and the production of biosurfactants by cloud microorganisms could be explained by the significant differences in the vegetation of France. For example, the predominance of the most efficient biosurfactant-producing microorganisms in clouds originating from the Northwest-North sector could reflect agricultural practices. These French regions are characterized by uniform monocultures (Ministère de l'Agriculture, de l'Agroalimentaire et de la Forêt, 2014) extremely favorable to plant pathogens, such as *Pseudomonas syringae* (McDonald and Linde, 2002).

The occurrence of biosurfactants has recently been shown in atmospheric aerosols (Baduel et al., 2012; Ekström et al., 2010; Nozière et al., 2014). Indeed, the presence of biosurfactants in the atmosphere could result from different mechanisms. First, these aerosols could be directly produced by microorganisms in the clouds. This idea is relevant considering that microorganisms are metabolically active in clouds and can survive under atmospheric conditions (Joly et al., 2015 and references hereinafter). Indeed, microbial metabolic activity has been demonstrated by measurements of the ATP (adenosine 5'triphosphate) content (Amato et al., 2005, 2007a, Vaïtilingom et al., 2012, 2013) and by the uptake of the dye CTC (5-cyano-2,3-ditolyl tetrazolium chloride) directly *in situ* in cloud water samples (Hill et al., 2007). This activity has even been demonstrated at low temperatures (Sattler et al., 2001). In addition, these microorganisms also survive and resist atmospheric stresses, including evaporation-condensation cycles, freeze-thaw cycles, and exposure to oxidants and solar light and cold temperature (Delort et al., 2010; Joly et al., 2015). Biosurfactants can also be produced in extreme environments, such as deep sea or Arctic soil (Jackson et al., 2015; Janek et al., 2013). Under laboratory conditions, the microorganisms produced biosurfactants within the first 24 hours, even in poor nutritive medium (R2A). As the residence time of microorganisms in the atmosphere was modeled between 2 to 10 days (Burrows et al., 2009), these metabolically active microorganisms could thus synthesize biosurfactants in the clouds. A second and obvious pathway for the incorporation of biosurfactants in the atmosphere is associated with the presence of these molecules at the surface of the microorganisms. Biosurfactants can therefore be carried together with the microorganism when aerosolized or in a biofilm existing on dust or leaf particles. Moreover, biosurfactants could be directly emitted into the atmosphere as biogenic aerosols, particularly biosurfactants of marine origin, which can be emitted in sea sprays during bubbling and wave braking processes (Blanchard, 1989; Elliott et al., 2014).

The presence of biosurfactants might have implications for atmospheric processes. First, these molecules could impact atmospheric chemistry. For example, the *in situ* biosynthesis of biosurfactants by microorganisms in clouds could be considered as the production of secondary organic aerosols, thus modifying the global carbon balance in the water phase. The presence of biosurfactants (aerosolized from the earth surface or produced *in situ*) at the surface of cloud droplets could also change the precise picture of the mass transfer between the gas and water phases of clouds. Organic surface films can provide barriers against transportation across the air-

particle interface, inhibiting the uptake of water and gas phase species. These organic films can be an auspicious medium for solubilizing gas phase organic species, perhaps reflecting the observed non-Henry's law concentration of organics in field samples (Barnes, 1986; Davies et al., 2013; Lo and Lee, 1996; Park et al., 2009; Renard et al., 2014). The chemical characterization and the study of the reactivity of these organic layers will be of special interest to further understand cloud chemistry.

Second, biosurfactants could affect atmospheric microphysics by modifying CCN activation. Owing to their exceptional scope in reducing surface tension, biosurfactants *per se*, whether present on aerosols or associated at the microorganisms surface, are thus likely to enhance the propensity of these aerosols to form clouds, as the activation of particles into cloud droplets depends on surface tension according to Köhler's theory (Köhler, 1936). This topic has been controversial, but recently, Nozière et al. (2014) showed that the total surfactant fraction of atmospheric particles is much more surface-active ($\sigma \leq 30 \text{ mN m}^{-1}$) than HULIS. These authors demonstrated that the equilibration time of biosurfactants might hinder the measurement of such an effect when using classical instruments, such as a hygroscopic tandem differential mobility analyzer or a CCN counter. Upon further examination of the results obtained in the present study, it is reasonable to consider that biosurfactant partitioning in macroscopic pendant droplets might decrease the surface tension values relative to atmospheric conditions. Indeed, in a microscopic ($D_{\text{wet}} \approx 1 \text{ }\mu\text{m}$) droplet, the partitioning impact could be insignificant owing to a surface area to volume ratio several orders of magnitude higher (Prisle et al., 2012; Sorjamaa et al., 2004). Nevertheless, the lower the radius of the droplet or the higher the surface to volume ratio, the higher the bacteria concentration (Aller et al., 2005; Hejkal et al., 1980) and the higher the WSOC concentration (Ervens and Volkamer, 2010). By dividing the surfactant concentration in the atmosphere ($\sim 10^{-12} - 10^{-9} \text{ mol m}^{-3}$ in Olkowska et al., 2014) by the liquid water content of wet aerosol ($\sim 10^{-6} - 10^{-5} \text{ g m}^{-3}$ in Ervens and Volkamer, 2010), we obtained a significant concentration of $\sim \text{mM}$. A recent study (Gérard et al., 2016) reported concentrations above the typical CMC. Thus, Ruehl et al. (2012) presented strong evidence that surface tension reduction can occur in microscopic droplets, and even more in wet aerosol, provided that the particles predominantly (*i.e.*, $\gtrsim 80\%$) comprise surfactants.

The influence of biological surfactants on the prediction of particle cloud activation and indirect aerosol climate effects should be implemented in models. Indeed, if the effects of organic surfactants (particularly carboxylic acids) on the surface tension of activating droplets is considered in parameterizations (Abdul-Razzak and Ghan, 2004), then recent studies have shown that the surface partitioning of organic molecules to a microscopic aqueous droplet interface should also be considered in models (Nozière, 2016; Prisle et al., 2012; Ruehl et al., 2016).

Moreover, because *Pseudomonas* strains are the most efficient biosurfactant-producing bacteria and are dominant in clouds and rain (the present study and Ahern et al., 2007), the question arises about the potential role of biosurfactants in the cycle of *Pseudomonas* in the atmosphere. Indeed, biosurfactants facilitate the aerosolization of these strains from leaf surfaces and favor the formation of cloud droplets through the modification of the surface properties of the cells. Moreover, most of these strains belong to *Pseudomonas syringae* species, which are ice nuclei active and can induce precipitations back to the earth. Thus, biosurfactants

should be integrated into the life history of *Pseudomonas syringae* and its relationship to the water cycle (Morris et al., 2008, 2014).

In conclusion, the results of the present study showed that the microbial strains isolated from cloud waters produce strong biosurfactants under laboratory conditions. The major and most active producers belong to the *Pseudomonas* genus, which is prevalent in cloud water and typically originates from the phyllosphere. Although the presence of surfactants has been shown on aerosols (Nozière et al., 2014), it has not yet been demonstrated in clouds, and the structure of these compounds has not been established. The biosurfactants overproduced by the best producers in the present study will be isolated to analyze their chemical structure. In parallel, the biosurfactants from cloud aerosols and rain samples will also be extracted, and their structural fingerprints will be analyzed and compared with the signatures of microbial surfactants isolated from clouds. These comparisons should provide evidence of the microbial origin of the surfactants present on aerosols. Studying such biosurfactants in the atmosphere is of special interest for the chemical characterization and reactivity of organic layers and the characterization of their impact on mass transfer and water uptake. The activation of aerosols containing organic matter is a major topic directly associated with the climatic effects of aerosol-cloud-interactions. A small change in the droplet population could affect cloud albedo and the formation of precipitation (Li et al., 2011; Rodhe, 1999). Hence, there is a need to enhance our knowledge about biosurfactants, focusing on the extent of their impact on human health and the global climate (Brimblecombe and Latif, 2004).

Acknowledgments: This work was funded by the French-USA program ANR-NSF SONATA and the French-Slovak Program Stefanik. Cloud sampling was possible thanks to the staff of the Observatoire de Physique du Globe de Clermont-Ferrand. The authors would like to thank Magali Abrantes for her technical assistance.

References

- Abdul-Razzak, H. and Ghan, S. J.: Parameterization of the influence of organic surfactants on aerosol activation, *J. Geophys. Res. Atmospheres*, 109(D3), D03205, doi:10.1029/2003JD004043, 2004.
- Ahern, H. E., Walsh, K. A., Hill, T. C. J. and Moffett, B. F.: Fluorescent pseudomonads isolated from Hebridean
 5 cloud and rain water produce biosurfactants but do not cause ice nucleation, *Biogeosciences*, 4(1), 115–124, doi:10.5194/bg-4-115-2007, 2007.
- Aller, J. Y., Kuznetsova, M. R., Jahns, C. J. and Kemp, P. F.: The sea surface microlayer as a source of viral and bacterial enrichment in marine aerosols, *J. Aerosol Sci.*, 36(5–6), 801–812, doi:10.1016/j.jaerosci.2004.10.012, 2005.
- 10 Amato, P., Ménager, M., Sancelme, M., Laj, P., Mailhot, G. and Delort, A.-M.: Microbial population in cloud water at the Puy de Dôme: Implications for the chemistry of clouds, *Atmos. Environ.*, 39(22), 4143–4153, doi:10.1016/j.atmosenv.2005.04.002, 2005.
- Amato, P., Parazols, M., Sancelme, M., Mailhot, G., Laj, P. and Delort, A.-M.: An important oceanic source of micro-organisms for cloud water at the Puy de Dôme (France), *Atmos. Environ.*, 41(37), 8253–8263,
 15 doi:10.1016/j.atmosenv.2007.06.022, 2007a.
- Amato, P., Hennebelle, R., Magand, O., Sancelme, M., Delort, A.-M., Barbante, C., Boutron, C. and Ferrari, C.: Bacterial characterization of the snow cover at Spitzberg, Svalbard, *FEMS Microbiol. Ecol.*, 59(2), 255–264, doi:10.1111/j.1574-6941.2006.00198.x, 2007b.
- Aumann, E. and Tabazadeh, A.: Rate of organic film formation and oxidation on aqueous drops, *J. Geophys.*
 20 *Res.*, 113(D23), doi:10.1029/2007JD009738, 2008.
- Aumann, E., Hildemann, L. M. and Tabazadeh, A.: Measuring and modeling the composition and temperature-dependence of surface tension for organic solutions, *Atmos. Environ.*, 44(3), 329–337, doi:10.1016/j.atmosenv.2009.10.033, 2010.
- Baduel, C., Nozière, B. and Jaffrezo, J.-L.: Summer/winter variability of the surfactants in aerosols from
 25 Grenoble, France, *Atmos. Environ.*, 47, 413–420, doi:10.1016/j.atmosenv.2011.10.040, 2012.
- Banat, I. M., Franzetti, A., Gandolfi, I., Bestetti, G., Martinotti, M. G., Fracchia, L., Smyth, T. J. and Marchant, R.: Microbial biosurfactants production, applications and future potential, *Appl. Microbiol. Biotechnol.*, 87(2), 427–444, doi:10.1007/s00253-010-2589-0, 2010.
- Barnes, G. T.: The effects of monolayers on the evaporation of liquids, *Adv. Colloid Interface Sci.*, 25, 89–200,
 30 doi:10.1016/0001-8686(86)80004-5, 1986.
- Bertram, A. K., Ivanov, A. V., Hunter, M., Molina, L. T. and Molina, M. J.: The Reaction Probability of OH on Organic Surfaces of Tropospheric Interest, *J. Phys. Chem. A*, 105(41), 9415–9421, doi:10.1021/jp0114034, 2001.

- Blanchard, D. C.: The ejection of drops from the sea and their enrichment with bacteria and other materials: a review, *Estuaries Coasts*, 12(3), 127–137, 1989.
- Bodour, A. A., Drees, K. P. and Maier, R. M.: Distribution of Biosurfactant-Producing Bacteria in Undisturbed and Contaminated Arid Southwestern Soils, *Appl. Environ. Microbiol.*, 69(6), 3280–3287, doi:10.1128/AEM.69.6.3280-3287.2003, 2003.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U. and others: Clouds and aerosols, in *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 571–657, Cambridge University Press., 2013.
- 10 Brimblecombe, P. and Latif, M. T.: Rediscovering Atmospheric Surfactants, *Environ. Chem. - Env. CHEM*, 1(1), doi:10.1071/EN04044, 2004.
- Burrows, S. M., Elbert, W., Lawrence, M. G. and Pöschl, U.: Bacteria in the global atmosphere – Part 1: Review and synthesis of literature data for different ecosystems, *Atmos Chem Phys*, 9(23), 9263–9280, doi:10.5194/acp-9-9263-2009, 2009.
- 15 Cai, Q., Zhang, B., Chen, B., Song, X., Zhu, Z. and Cao, T.: Screening of biosurfactant-producing bacteria from offshore oil and gas platforms in North Atlantic Canada, *Environ. Monit. Assess.*, 187(5), 1–12, doi:10.1007/s10661-015-4490-x, 2015.
- Capel, P. D., Gunde, R., Zuercher, F. and Giger, W.: Carbon speciation and surface tension of fog, *Environ. Sci. Technol.*, 24(5), 722–727, doi:10.1021/es00075a017, 1990.
- 20 Chakraborty, P. and Zachariah, M. R.: On the structure of organic-coated water droplets: From “net water attractors” to “oily” drops, *J. Geophys. Res.*, 116(D21), doi:10.1029/2011JD015961, 2011.
- Christofi, N. and Ivshina, I. b.: Microbial surfactants and their use in field studies of soil remediation, *J. Appl. Microbiol.*, 93(6), 915–929, doi:10.1046/j.1365-2672.2002.01774.x, 2002.
- Chrzanowski, Ł., Ławniczak, Ł. and Czaczyk, K.: Why do microorganisms produce rhamnolipids?, *World J. Microbiol. Biotechnol.*, 28(2), 401–419, doi:10.1007/s11274-011-0854-8, 2012.
- 25 Clifford, D., Bartels-Rausch, T. and Donaldson, D. J.: Suppression of aqueous surface hydrolysis by monolayers of short chain organic amphiphiles, *Phys. Chem. Chem. Phys.*, doi:10.1039/b617079j, 2007.
- D’aes, J., De Maeyer, K., Pauwelyn, E. and Höfte, M.: Biosurfactants in plant–*Pseudomonas* interactions and their importance to biocontrol, *Environ. Microbiol. Rep.*, 2(3), 359–372, doi:10.1111/j.1758-2229.2009.00104.x, 2010.
- 30 Davies, J. F., Miles, R. E. H., Haddrell, A. E. and Reid, J. P.: Influence of organic films on the evaporation and condensation of water in aerosol, *Proc. Natl. Acad. Sci.*, 110(22), 8807–8812, doi:10.1073/pnas.1305277110, 2013.

- Decesari, S., Facchini, M. C., Matta, E., Lettini, F., Mircea, M., Fuzzi, S., Tagliavini, E. and Putaud, J.-P.: Chemical features and seasonal variation of fine aerosol water-soluble organic compounds in the Po Valley, Italy, *Atmos. Environ.*, 35(21), 3691–3699, doi:10.1016/S1352-2310(00)00509-4, 2001.
- Decesari, S., Facchini, M. C., Mircea, M., Cavalli, F. and Fuzzi, S.: Solubility properties of surfactants in atmospheric aerosol and cloud/fog water samples, *J. Geophys. Res. Atmospheres*, 108(D21), 4685, doi:10.1029/2003JD003566, 2003.
- Decesari, S., Facchini, M. C., Fuzzi, S., McFiggans, G. B., Coe, H. and Bower, K. N.: The water-soluble organic component of size-segregated aerosol, cloud water and wet depositions from Jeju Island during ACE-Asia, *Atmos. Environ.*, 39(2), 211–222, doi:10.1016/j.atmosenv.2004.09.049, 2005.
- 10 Deguillaume, L., Charbouillot, T., Joly, M., Vaïtilingom, M., Parazols, M., Marinoni, A., Amato, P., Delort, A.-M., Vinatier, V., Flossmann, A., Chaumerliac, N., Pichon, J. M., Houdier, S., Laj, P., Sellegri, K., Colomb, A., Brigante, M. and Mailhot, G.: Classification of clouds sampled at the puy de Dôme (France) based on 10 yr of monitoring of their physicochemical properties, *Atmospheric Chem. Phys.*, 14(3), 1485–1506, doi:10.5194/acp-14-1485-2014, 2014.
- 15 Delort, A.-M., Vaïtilingom, M., Amato, P., Sancelme, M., Parazols, M., Mailhot, G., Laj, P. and Deguillaume, L.: A short overview of the microbial population in clouds: Potential roles in atmospheric chemistry and nucleation processes, *Atmospheric Res.*, 98(2–4), 249–260, doi:10.1016/j.atmosres.2010.07.004, 2010.
- Desai, J. D. and Banat, I. M.: Microbial production of surfactants and their commercial potential., *Microbiol. Mol. Biol. Rev.*, 61(1), 47–64, 1997.
- 20 Dinar, E., Taraniuk, I., Graber, E. R., Katsman, S., Moise, T., Anttila, T., Mentel, T. F. and Rudich, Y.: Cloud Condensation Nuclei properties of model and atmospheric HULIS, *Atmos Chem Phys*, 6(9), 2465–2482, doi:10.5194/acp-6-2465-2006, 2006.
- Ekström, S., Nozière, B., Hultberg, M., Alsberg, T., Magnér, J., Nilsson, E. D. and Artaxo, P.: A possible role of ground-based microorganisms on cloud formation in the atmosphere, *Biogeosciences*, 7(1), 387–394, doi:10.5194/bg-7-387-2010, 2010.
- 25 Elliott, S., Burrows, S. M., Deal, C., Liu, X., Long, M., Ogunro, O., Russell, L. M. and Wingenter, O.: Prospects for simulating macromolecular surfactant chemistry at the ocean–atmosphere boundary, *Environ. Res. Lett.*, 9(6), 64012, doi:10.1088/1748-9326/9/6/064012, 2014.
- Ellison, G. B., Tuck, A. F. and Vaida, V.: Atmospheric processing of organic aerosols, *J. Geophys. Res. Atmospheres*, 104(D9), 11633–11641, doi:10.1029/1999JD900073, 1999.
- 30 Ervens, B. and Volkamer, R.: Glyoxal processing by aerosol multiphase chemistry: towards a kinetic modeling framework of secondary organic aerosol formation in aqueous particles, *Atmospheric Chem. Phys.*, 10(17), 8219–8244, doi:10.5194/acp-10-8219-2010, 2010.
- Facchini, M. C., Mircea, M., Fuzzi, S. and Charlson, R. J.: Cloud albedo enhancement by surface-active organic solutes in growing droplets, *Nature*, 401(6750), 257–259, doi:10.1038/45758, 1999.
- 35

- Facchini, M. C., Decesari, S., Mircea, M., Fuzzi, S. and Loglio, G.: Surface tension of atmospheric wet aerosol and cloud/fog droplets in relation to their organic carbon content and chemical composition, *Atmos. Environ.*, 34(28), 4853–4857, doi:10.1016/S1352-2310(00)00237-5, 2000.
- Feingold, G. and Chuang, P. Y.: Analysis of the Influence of Film-Forming Compounds on Droplet Growth: Implications for Cloud Microphysical Processes and Climate, *J. Atmospheric Sci.*, 59(12), 2006–2018, doi:10.1175/1520-0469(2002)059<2006:AOTIOF>2.0.CO;2, 2002.
- Gautam, K. K. and Tyagi, V. K.: Microbial Surfactants: A Review, *J. Oleo Sci.*, 55(4), 155–166, doi:10.5650/jos.55.155, 2006.
- Gérard, V., Nozière, B., Baduel, C., Fine, L., Frossard, A. A. and Cohen, R. C.: Anionic, Cationic, and Nonionic Surfactants in Atmospheric Aerosols from the Baltic Coast at Askö, Sweden: Implications for Cloud Droplet Activation, *Environ. Sci. Technol.*, 50(6), 2974–2982, doi:10.1021/acs.est.5b05809, 2016.
- Gill, P. S., Graedel, T. E. and Weschler, C. J.: Organic films on atmospheric aerosol particles, fog droplets, cloud droplets, raindrops, and snowflakes, *Rev. Geophys.*, 21(4), 903–920, doi:10.1029/RG021i004p00903, 1983.
- Gilman, J. B. and Vaida, V.: Permeability of Acetic Acid through Organic Films at the Air–Aqueous Interface, *J. Phys. Chem. A*, 110(24), 7581–7587, doi:10.1021/jp061220n, 2006.
- Hansen, F. K. and Rødsrud, G.: Surface tension by pendant drop, *J. Colloid Interface Sci.*, 141(1), 1–9, doi:10.1016/0021-9797(91)90296-K, 1991.
- Hejkal, T. W., LaRock, P. A. and Winchester, J. W.: Water-to-Air Fractionation of Bacteria, *Appl. Environ. Microbiol.*, 39(2), 335–338, 1980.
- Herckes, P., Valsaraj, K. T. and Collett, J. L.: A review of observations of organic matter in fogs and clouds: Origin, processing and fate, *Atmospheric Res.*, 132–133, 434–449, doi:10.1016/j.atmosres.2013.06.005, 2013.
- Hill, K. A., Shepson, P. B., Galbavy, E. S., Anastasio, C., Kourtev, P. S., Konopka, A. and Stirm, B. H.: Processing of atmospheric nitrogen by clouds above a forest environment, *J. Geophys. Res. Atmospheres*, 112(D11), D11301, doi:10.1029/2006jd008002, 2007.
- Hitzenberger, R., Berner, A., Kasper-Giebl, A., Löflund, M. and Puxbaum, H.: Surface tension of Rax cloud water and its relation to the concentration of organic material, *J. Geophys. Res. Atmospheres*, 107(D24), 4752, doi:10.1029/2002JD002506, 2002.
- Hua, X. Y. and Rosen, M. J.: Dynamic surface tension of aqueous surfactant solutions, *J. Colloid Interface Sci.*, 141(1), 180–190, doi:10.1016/0021-9797(91)90313-W, 1991.
- Jackson, S. A., Borchert, E., O’Gara, F. and Dobson, A. D. W.: Metagenomics for the discovery of novel biosurfactants of environmental interest from marine ecosystems, *Curr. Opin. Biotechnol.*, 33, 176–182, doi:10.1016/j.copbio.2015.03.004, 2015.
- Janek, T., Łukaszewicz, M. and Krasowska, A.: Identification and characterization of biosurfactants produced by the Arctic bacterium *Pseudomonas putida* BD2, *Colloids Surf. B Biointerfaces*, 110, 379–386, doi:10.1016/j.colsurfb.2013.05.008, 2013.

- Joly, M., Amato, P., Sancelme, M., Vinatier, V., Abrantes, M., Deguillaume, L. and Delort, A.-M.: Survival of microbial isolates from clouds toward simulated atmospheric stress factors, *Atmos. Environ.*, 117, 92–98, doi:10.1016/j.atmosenv.2015.07.009, 2015.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R.,
 5 Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G. and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos Chem Phys*, 5(4), 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- Karanth, N. G. K., Deo, P. G. and Veenanadig, N. K.: Microbial production of biosurfactants and their
 10 importance, *Curr. Sci.*, 77(1), 116–126, 1999.
- King, E. O., Ward, M. K. and Raney, D. E.: Two simple media for the demonstration of pyocyanin and fluorescein, *J. Lab. Clin. Med.*, 44(2), 301–307, 1954.
- Kiss, G., Tombácz, E. and Hansson, H.-C.: Surface Tension Effects of Humic-Like Substances in the Aqueous Extract of Tropospheric Fine Aerosol, *J. Atmospheric Chem.*, 50(3), 279–294, doi:10.1007/s10874-005-5079-5,
 15 2005.
- Köhler, H.: The nucleus in and the growth of hygroscopic droplets, *Trans. Faraday Soc.*, 32, 1152, doi:10.1039/tf9363201152, 1936.
- Lance, S., Nenes, A. and Rissman, T. A.: Chemical and dynamical effects on cloud droplet number: Implications for estimates of the aerosol indirect effect, *J. Geophys. Res. Atmospheres*, 109(D22), D22208, doi:10.1029/2004JD004596, 2004.
 20
- Li, X., Hede, T., Tu, Y., Leck, C. and Ågren, H.: Surface-Active *cis* -Pinonic Acid in Atmospheric Droplets: A Molecular Dynamics Study, *J. Phys. Chem. Lett.*, 1(4), 769–773, doi:10.1021/jz9004784, 2010.
- Li, Y., Ezell, M. J. and Finlayson-Pitts, B. J.: The impact of organic coatings on light scattering by sodium chloride particles, *Atmos. Environ.*, 45(25), 4123–4132, doi:10.1016/j.atmosenv.2011.05.031, 2011.
- Lo, J.-H. A. and Lee, W.-M. G.: Effect of surfactant film on solubility of hydrophobic organic compounds in fog droplets, *Chemosphere*, 33(7), 1391–1408, doi:10.1016/0045-6535(96)00273-1, 1996.
 25
- Mann, E. E. and Wozniak, D. J.: *Pseudomonas* biofilm matrix composition and niche biology, *FEMS Microbiol. Rev.*, 36(4), 893–916, doi:10.1111/j.1574-6976.2011.00322.x, 2012.
- McDonald, B. A. and Linde, C.: The population genetics of plant pathogens and breeding strategies for durable resistance, *Euphytica*, 124(2), 163–180, doi:10.1023/A:1015678432355, 2002.
 30
- McFiggans, G., Artaxo, P., Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S., Gysel, M., Laaksonen, A., Lohmann, U., Mentel, T. F., Murphy, D. M., O'Dowd, C. D., Snider, J. R. and Weingartner, E.: The effect of physical and chemical aerosol properties on warm cloud droplet activation, *Atmos Chem Phys*, 6(9), 2593–2649, doi:10.5194/acp-6-2593-2006, 2006.

Ministère de l'Agriculture, de l'Agroalimentaire et de la Forêt: Agreste La statistique, l'évaluation et la prospective agricole, Territoire et climat., [online] Available from: <http://agreste.agriculture.gouv.fr/IMG/pdf/Gaf15p013-017.pdf> (Accessed 20 May 2016), 2014.

- 5 Mircea, M., Facchini, M. C., Decesari, S., Fuzzi, S. and Charlson, R. J.: The influence of the organic aerosol component on CCN supersaturation spectra for different aerosol types, *Tellus B*, 54(1), 74–81, doi:10.1034/j.1600-0889.2002.00256.x, 2002.
- Mircea, M., Facchini, M. C., Decesari, S., Cavalli, F., Emblico, L., Fuzzi, S., Vestin, A., Rissler, J., Swietlicki, E., Frank, G. and others: Importance of the organic aerosol fraction for modeling aerosol hygroscopic growth and activation: a case study in the Amazon Basin, *Atmospheric Chem. Phys.*, 5(11), 3111–3126, 2005.
- 10 Morris, C. E., Sands, D. C., Vinatzer, B. A., Glaux, C., Guilbaud, C., Buffière, A., Yan, S., Dominguez, H. and Thompson, B. M.: The life history of the plant pathogen *Pseudomonas syringae* is linked to the water cycle, *ISME J.*, 2(3), 321–334, 2008.
- Morris, C. E., Conen, F., Alex Huffman, J., Phillips, V., Pöschl, U. and Sands, D. C.: Bioprecipitation: a feedback cycle linking Earth history, ecosystem dynamics and land use through biological ice nucleators in the atmosphere, *Glob. Change Biol.*, 20(2), 341–351, doi:10.1111/gcb.12447, 2014.
- 15 Mulligan, C. N.: Recent advances in the environmental applications of biosurfactants, *Curr. Opin. Colloid Interface Sci.*, 14(5), 372–378, doi:10.1016/j.cocis.2009.06.005, 2009.
- Murphy, D. M., Cziczo, D. J., Froyd, K. D., Hudson, P. K., Matthew, B. M., Middlebrook, A. M., Peltier, R. E., Sullivan, A., Thomson, D. S. and Weber, R. J.: Single-particle mass spectrometry of tropospheric aerosol particles, *J. Geophys. Res.*, 111(D23), doi:10.1029/2006JD007340, 2006.
- 20 Nenes, A., Charlson, R. J., Facchini, M. C., Kulmala, M., Laaksonen, A. and Seinfeld, J. H.: Can chemical effects on cloud droplet number rival the first indirect effect?, *Geophys. Res. Lett.*, 29(17), 1848, doi:10.1029/2002GL015295, 2002.
- Nozière, B.: Don't forget the surface, *Science*, 351(6280), 1396–1397, doi:10.1126/science.aaf3253, 2016.
- 25 Nozière, B., Baduel, C. and Jaffrezo, J.-L.: The dynamic surface tension of atmospheric aerosol surfactants reveals new aspects of cloud activation, *Nat. Commun.*, 5, doi:10.1038/ncomms4335, 2014.
- O'Dowd, C. D., Aalto, P., Hmeri, K., Kulmala, M. and Hoffmann, T.: Aerosol formation: Atmospheric particles from organic vapours, *Nature*, 416(6880), 497–498, doi:10.1038/416497a, 2002.
- Olkowska, E., Ruman, M. and Polkowska, Ż.: Occurrence of Surface Active Agents in the Environment, *J. Anal. Methods Chem.*, 2014, e769708, doi:10.1155/2014/769708, 2014.
- 30 Park, S.-C., Burden, D. K. and Nathanson, G. M.: Surfactant Control of Gas Transport and Reactions at the Surface of Sulfuric Acid, *Acc. Chem. Res.*, 42(2), 379–387, doi:10.1021/ar800172m, 2009.
- Prisle, N. L., Ottosson, N., Öhrwall, G., Söderström, J., Dal Maso, M. and Björneholm, O.: Surface/bulk partitioning and acid/base speciation of aqueous decanoate: direct observations and atmospheric implications, *Atmospheric Chem. Phys.*, 12(24), 12227–12242, doi:10.5194/acp-12-12227-2012, 2012.
- 35

- Raaijmakers, J. M., Bruijn, I. D., Nybroe, O. and Ongena, M.: Natural functions of lipopeptides from *Bacillus* and *Pseudomonas*: more than surfactants and antibiotics, *FEMS Microbiol. Rev.*, 34(6), 1037–1062, doi:10.1111/j.1574-6976.2010.00221.x, 2010.
- Reasoner, D. J. and Geldreich, E. E.: A new medium for the enumeration and subculture of bacteria from potable water, *Appl. Environ. Microbiol.*, 49(1), 1–7, 1985.
- Renard, P., Reed Harris, A. E., Rapf, R. J., Ravier, S., Demelas, C., Coulomb, B., Quivet, E., Vaida, V. and Monod, A.: Aqueous Phase Oligomerization of Methyl Vinyl Ketone by Atmospheric Radical Reactions, *J. Phys. Chem. C*, 118(50), 29421–29430, doi:10.1021/jp5065598, 2014.
- Rodhe, H.: Atmospheric chemistry: Clouds and climate, *Nature*, 401(6750), 223–225, doi:10.1038/45701, 1999.
- Ron, E. Z. and Rosenberg, E.: Natural roles of biosurfactants, *Environ. Microbiol.*, 3(4), 229–236, doi:10.1046/j.1462-2920.2001.00190.x, 2001.
- Rosenberg, E. and Ron, E. Z.: High- and low-molecular-mass microbial surfactants, *Appl. Microbiol. Biotechnol.*, 52(2), 154–162, doi:10.1007/s002530051502, 1999.
- Rudich, Y.: Laboratory Perspectives on the Chemical Transformations of Organic Matter in Atmospheric Particles, *Chem. Rev.*, 103(12), 5097–5124, doi:10.1021/cr020508f, 2003.
- Ruehl, C. R., Chuang, P. Y., Nenes, A., Cappa, C. D., Kolesar, K. R. and Goldstein, A. H.: Strong evidence of surface tension reduction in microscopic aqueous droplets, *Geophys. Res. Lett.*, 39(23), L23801, doi:10.1029/2012GL053706, 2012.
- Ruehl, C. R., Davies, J. F. and Wilson, K. R.: An interfacial mechanism for cloud droplet formation on organic aerosols, *Science*, 351(6280), 1447–1450, doi:10.1126/science.aad4889, 2016.
- Sareen, N., Schwier, A. N., Lathem, T. L., Nenes, A. and McNeill, V. F.: Surfactants from the gas phase may promote cloud droplet formation, *Proc. Natl. Acad. Sci.*, 110(8), 2723–2728, doi:10.1073/pnas.1204838110, 2013.
- Satpute, S. K., Banat, I. M., Dhakephalkar, P. K., Banpurkar, A. G. and Chopade, B. A.: Biosurfactants, bioemulsifiers and exopolysaccharides from marine microorganisms, *Biotechnol. Adv.*, 28(4), 436–450, doi:10.1016/j.biotechadv.2010.02.006, 2010.
- Sattler, B., Puxbaum, H. and Psenner, R.: Bacterial growth in supercooled cloud droplets, *Geophys. Res. Lett.*, 28(2), 239–242, doi:10.1029/2000GL011684, 2001.
- Saxena, P. and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical review of the literature and application of thermodynamics to identify candidate compounds, *J. Atmospheric Chem.*, 24(1), 57–109, doi:10.1007/BF00053823, 1996.
- Seidl, W. and Hänel, G.: Surface-active substances on rainwater and atmospheric particles, *Pure Appl. Geophys.*, 121(5–6), 1077–1093, 1983.

Shulman, M. L., Jacobson, M. C., Carlson, R. J., Synovec, R. E. and Young, T. E.: Dissolution behavior and surface tension effects of organic compounds in nucleating cloud droplets, *Geophys. Res. Lett.*, 23(3), 277–280, doi:10.1029/95GL03810, 1996.

Sorjamaa, R., Svenningsson, B., Raatikainen, T., Henning, S., Bilde, M. and Laaksonen, A.: The role of surfactants in Köhler theory reconsidered, *Atmos Chem Phys*, 4(8), 2107–2117, doi:10.5194/acp-4-2107-2004, 2004.

Svenningsson, B., Rissler, J., Swietlicki, E., Mircea, M., Bilde, M., Facchini, M. C., Decesari, S., Fuzzi, S., Zhou, J., Mønster, J. and Rosenørn, T.: Hygroscopic growth and critical supersaturations for mixed aerosol particles of inorganic and organic compounds of atmospheric relevance, *Atmos Chem Phys*, 6(7), 1937–1952, doi:10.5194/acp-6-1937-2006, 2006.

Taraniuk, I., Graber, E. R., Kostinski, A. and Rudich, Y.: Surfactant properties of atmospheric and model humic-like substances (HULIS), *Geophys. Res. Lett.*, 34(16), L16807, doi:10.1029/2007GL029576, 2007.

Vařtilingom, M., Attard, E., Gaiani, N., Sancelme, M., Deguillaume, L., Flossmann, A. I., Amato, P. and Delort, A.-M.: Long-term features of cloud microbiology at the puy de Dôme (France), *Atmos. Environ.*, 56, 88–100, doi:10.1016/j.atmosenv.2012.03.072, 2012.

Vařtilingom, M., Deguillaume, L., Vinatier, V., Sancelme, M., Amato, P., Chaumerliac, N. and Delort, A.-M.: Potential impact of microbial activity on the oxidant capacity and organic carbon budget in clouds, *Proc. Natl. Acad. Sci.*, 110(2), 559–564, doi:10.1073/pnas.1205743110, 2013.

Zenchelsky, S. and Youssefi, M.: Natural organic atmospheric aerosols of terrestrial origin, *Rev. Geophys.*, 17(3), 459–464, doi:10.1029/RG017i003p00459, 1979.

Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M. and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34(13), L13801, doi:10.1029/2007GL029979, 2007.