

Replies to reviewer 2

by E. Christner, M. Kohler, M. Schneider

January 4, 2017

We like to thank reviewer 2 for the review and the constructive comments, which helped to improve the manuscript.

According to the comments of reviewer 1, we implemented (1) kinetic fractionation during surface evaporation at skin temperatures above 0°C and (2) reduced fractionation under conditions of supersaturation in clouds into the model. Because of this, most numbers and figures throughout the manuscript are slightly changed compared to the initial version. A marked-up manuscript version is attached at the end of this document.

General comments:

The manuscript by Christner et al. presents 17 months of new, continuous, low-level atmospheric water vapor δD measurements at a site near Karlsruhe, Germany in conjunction with a Lagrangian isotope model to inform understanding of the along-trajectory controls on measured δD variability. Integration of HYSPLIT-calculated, low-level back-trajectories with the isotope model shows that much of the observed variability in measured δD values is the result of identifiable along-transport processes most generally associated with moistening and dehydration of the air parcel through evapotranspiration and precipitation processes, respectively. A subset of backtrajectories associated with 'cold snaps' result from a systemic shift to continental source regions and easterly trajectories. For this subset, the authors investigate an additional controlling mechanism for isotopic evolution of these easterly trajectories, that of isotopic modification via moistening from surface snow sublimation. In the closing section of the manuscript, the authors investigate a range of possible conditions for isotopic modification of regional snowpacks that best explains measured δD , namely skin temperature controls on fractionating versus non-fractionating sublimation processes. From this, the authors determine the relevant skin temperature window for which post-depositional isotopic modification of snowpacks and associated impacts on low-level atmospheric moisture δD is most relevant.

This is a detailed and comprehensive manuscript that presents a new and robust long-term δD dataset that proves valuable for investigating controls on the isotopic evolution of low-level atmospheric moisture. The methods applied are appropriate. I particularly like the integration of HYSPLIT-derived trajectories into the new Lagrangian model presented and find the observed model δD congruence (Figs. 6 and 7) impressive and supporting of the Lagrangian-model approach. I find the discussion of the isotope effects of sublimation both nuanced and convincing, which is important given that this impactful process is generally neglected or assumed to be negligible in similarly-focused studies. I expect the findings and research design of this study will be of interest to a broad audience, particularly in light of the expansion of laser-based isotope analyzers that are sure to increase the number of similar isotope records in coming years. Accordingly, I strongly support publication of the manuscript in ACP pending some minor/moderate revisions to the text and some figures in order to (1) reduce redundancies in the text, (2) provide additional clarification for components of the HYSPLIT-Lagrangian isotope model integration and (3) ensure the more complex aspects of the manuscript are understandable to the broader audience likely to be interested in this work (e.g., earth scientists interested in proxy-based investigations of paleoaltimetry and paleoclimate).

Specific comments:

(1) *Reorganization and reducing redundancies* – much of the text in Section 3.3 is redundant with Section 2.3.2. It seems much of 3.3 could be moved and combined with 2.3.2. Similarly, section 3.4 falls under the heading 'Measurements'; however, this moisture source data seems more relevant to the model description in Section 2.3.

Our reply: We removed the redundancies by moving the information from sections 3.3 and 3.4 into the sections 2.3.1 and 2.3.2.

The opening paragraph of Section 2.3.3 presents some basic back-trajectory statistics but explanation is limited here. Important clarifying information is not provided until Section 4, specifically Section 4.3. I suggest saving back-trajectory statistics for Section 4 when more details needed for clarification are presented.

Our reply: We slightly shortened the trajectory statistics and moved it to Sect. 4.1.

Discussion of 24-hour smoothing window on page 4 could be removed there and saved for the same discussion on p. 14 (lines 19-28).

Our reply: We shifted information regarding model uncertainty from the smoothing to Sect. 5.1 (p.15). Since already the moisture source diagnostics in Sect. 4.1 and at the beginning of Sect. 5.1 refer to the smoothing of q , we added a clarifying motivation for the smoothing to Sect. 2.1.: *"In addition, using GDAS wind fields with three-hourly time resolution for the calculation of back trajectories – which is a coarse time resolution compared to the GDAS time steps on the order of minutes – may cause deviations between the HYSPLIT back trajectories and the exact trajectories that air parcels followed in the GDAS. This, in turn, may result in artificial changes of q along the HYSPLIT back trajectories. For instance HYSPLIT trajectories not fully capturing diurnal vertical movement in consequence of thermal expansion of the atmospheric boundary layer in the GDAS may show artificial diurnal changes of q . Like Stohl and James (2004) and Sodemann et al. (2008b), we assume P and E to be the dominant processes and ignore the other effects. To avoid an overestimation of the formation of precipitation and moisture uptake in consequence of potentially artificial diurnal and sub-diurnal variations of q along the HYSPLIT trajectories, we smoothed q along the back trajectories with a 24 h rectangle function."*

(2) *Model clarification and limitations* – Given the uncertainty in vapor δD at high altitudes (> 2 km) and low proportion of trajectories (2%) encountering these altitudes, it seems these trajectories could simply be removed from the analysis.

Our reply: We removed the 2% air masses during cold snap which originated from altitudes higher than 2km above ground level from the analysis and deleted the discussion of model uncertainty from model initialization at high altitudes from Sect. 5.1. We updated slightly changed numbers and figures resulting from this change and added two sentences to the second paragraph in Section 5.1: *"Only 2% of the corresponding air masses originated from altitudes higher than 2000m above ground level. Since uncertainty of model initialization is especially high for these air masses, we finally excluded the 2% of air masses originating from high altitudes."*

Beyond the vapor measurements at the Karlsruhe study site, are there other published datasets that would ground truth the isotope model presented? Specifically, are there any regional records of soil water δD ($\delta^{18}O$) that can be presented to constrain the RCWIP values used?

Our reply: We expanded the motivation for assuming the δD of soil water to be close to the GNIP based monthly RCWIP data (Sect. 2.3.2): *"Because soil water in Central Europe is generally frequently recharged by precipitation, we ignore systematic enrichment of $HD^{16}O$ in the uppermost soil layer caused by continuous fractionating evaporation from the soil. For instance measurements of precipitation amount at the measurement site in Karlsruhe indicate recharge of soil water around the site by more than 1mm precipitation per day on average every 2.9 days. The assumption regarding enrichment is supported by findings of (Risi et al., 2016), who observe insignificant systematic deviations of isotope ratios of water within the upper 15 cm of the soil from isotope ratios in precipitation at several sites in France, Germany, and the Czech Republic."*

Additionally, are there any regional snowpack δD ($\delta^{18}O$) records that would give a better understanding of the degree of snowpack δD variability? It is likely that snowpack δD varies both spatially and temporally throughout the accumulation and melt season in the study region, thus some discussion on how this variability limits the model presented is important.

Our reply: In general the GNIP based RCWIP data reflects the spatial variability of the δD of snowfall (Fig. 1b) and also accounts for changes of δD in different months of a year. We expanded the discussion regarding the δD of surface layer snow during the accumulation and melt season in Sect. 5.1 and added the following paragraph: *“In contrast to this, post-depositional fractionation processes may increase the δD values of surface layer snow on the order of +10 to +20‰ during periods with small accumulation rates (Gurney & Lawrence, 2004; Moser & Stichler, 1974), suggesting a scenario where the δD values of surface layer snow are higher than the δD from the RCWIP. Please note that this scenario is not likely for a seasonal snowpack during melt season since ablation may uncover old snow from colder winter months, causing changes of the δD of surface layer snow of -50‰ (Dahlke & Lyon, 2013). Given the relatively small snow depths in the investigated moisture source region (Fig. 8b), we ignore the uncovering of older snow layers and only consider a potential post-depositional increase of the δD of surface layer snow.”*

Another limitation that might be more explicitly discussed or tied into the previous point about spatial δD variability of the snowpack is that of the $1^\circ \times 1^\circ$ resolution of the GDAS data set. How might this spatial smoothing impact ability to model δD variation?

Our reply: We added two paragraphs clarifying the meaning of $T_{\text{subl,max}}$ and the impact of horizontal resolution to Sect. 6: *“The determined $T_{\text{subl,max}}$ refers to a GDAS skin temperature that was weighted with positive latent heat flux at ground level. For this reason $T_{\text{subl,max}}$ is representative for GDAS skin temperatures during the day, when evaporation is strongest. However it should be kept in mind that due to the coarse resolution of $1 \times 1^\circ$ of the GDAS data, much spatial variability of the skin temperature is smoothed out. The meaning of the $T_{\text{subl,max}}$ derived in this study is therefore an average temperature in a $1 \times 1^\circ$ grid cell above which the evaporation of meltwater exceeds the amount of moisture from sublimation. A way to derive a more physical temperature separating the regimes of sublimation and meltwater evaporation could be using data with a higher horizontal resolution. This wouldn't necessarily improve accuracy with respect to the back trajectories' positions but locations with enhanced skin temperatures and especially high amounts of surface evaporation would be more realistically represented, presumably resulting in higher evaporation weighted skin temperatures and a higher $T_{\text{subl,max}}$. In addition, higher horizontal resolution would allow to better account for spatial heterogeneity of the δD of surface layer snow in mountainous regions, for instance by also weighting the δD of the snow with the amount of surface evaporation. In this context please note that systematic uncertainty regarding the δD of surface layer snow turned out to be the main limitation for determining $T_{\text{subl,max}}$ with the approach presented here. For this reason regular analysis of the δD of surface layer snow samples for instance at selected GNIP stations would be a very desirable and efficient measure to reduce uncertainty of $T_{\text{subl,max}}$.”*

(3) *Clarifying complexity* – I find Section 5.2 and corresponding Figure 10 difficult to comprehend. I understand the general idea that multiple model runs were used to identify the cutoff temperature between fractionation and non-fractionating sublimation, but is not clear to me how the 16 scenarios of ‘side constraint’ variability and associated 128 total model runs correspond to the lines shown in Fig. 10. How do 128 total model runs translate to 9 distinct lines in Fig. 10? Please clarify in the text and figure caption and reconsider what Figure 10 should show to more clearly communicate the information in this section.

Our reply: Figure 10 shows the results from 144 model scenarios with 16 different $T_{\text{subl,max}}$ and the 9 configurations M_{S_MW} , $M_{S_MW,-,snow}$, $M_{S_MW,-,ini}$, $M_{S_MW,-,upt.12h}$, $M_{S_MW,-,-}$, $M_{S_MW,+ ,snow}$, $M_{S_MW,+ ,ini}$, $M_{S_MW,+ ,upt.36h}$, $M_{S_MW,+++}$. For clarification we adapted Figure 10 and the respective caption and added some further explanation to Sect. 5.2 : “To assess this uncertainty of the optimal $T_{\text{subl,max}}$, we performed $16 \cdot 8 = 128$ additional model runs (Fig. 10, thin lines) corresponding to 16 different $T_{\text{subl,max}}$ from -15 to 0°C and 8 different model configurations with the same assumptions about a changed δD of snow, a changed δD at the model initialization, a different amount of moisture uptake, and superposition of the three effects as in the above uncertainty assessment for the Ms and the Mmw. Analogous to the uncertainty assessment for the Ms and the Mmw, we refer to the model runs as $M_{S_MW,+ ,snow, T_{\text{subl,max}}}$, $M_{S_MW,+ ,ini, T_{\text{subl,max}}}$, $M_{S_MW,+ ,upt.36h, T_{\text{subl,max}}}$, $M_{S_MW,+++ , T_{\text{subl,max}}}$, $M_{S_MW, snow, T_{\text{subl,max}}}$, $M_{S_MW,- ,ini, T_{\text{subl,max}}}$, $M_{S_MW,- ,upt.12h, T_{\text{subl,max}}}$, and $M_{S_MW,- , T_{\text{subl,max}}}$.”




Figure 10. Mean difference between modeled and observed δD of water vapor at Karlsruhe for selected cold snap events ($\Delta\delta D$). Each cross represents the $\Delta\delta D$ obtained from one model configuration with specific model assumptions regarding the $T_{\text{subl,max}}$, the δD of snow, the δD at the model initialization, and the amount of moisture uptake. The thick black line connects the $\Delta\delta D$ from 16 model configurations with $T_{\text{subl,max}}$ between -15 and 0°C and the model assumptions of $M_{S_MW, T_{\text{subl,max}}}$ and illustrates the increase of the modeled δD values with increasing maximum skin temperature allowing non-fractionating sublimation. The gray shaded area depicts possible systematic changes of the $\Delta\delta D$ in the case of a systematic deviation of the δD of snow from the RCWIP climatology (light blue), a systematically changed δD at the model initialization (yellow), a systematically changed amount of moisture uptake (green), and superposition of the different assumptions (thin black lines). The thin dashed black line corresponds to the model configuration $M_{S_MW,+++ , T_{\text{subl,max}}}$, which doesn't allow reproducing the observations in group “warm” and is therefore not considered in the gray shaded area of uncertainty. Magenta: optimal $T_{\text{subl,max}}$ for best agreement of model and observation (dot) and uncertainty of the optimal $T_{\text{subl,max}}$ (error bar) due to uncertainty of model assumptions.

Given the focus on easterly trajectories, a figure more clearly showing association of easterly trajectory pathways with corresponding snow cover in that region would be helpful. This might be accomplished by adding a panel to Figure 8 showing snow cover.

Our reply: We added a second panel to Figure 8 showing the mean GDAS snow height during the cold snaps.




Figure 8. (a) Lagrangian source region analysis of low-level water vapor in Karlsruhe for the observed cold snaps. The mean identified fraction of moisture along the five-day back trajectories is 48%. (b) Mean snow depth during the cold snaps based on GDAS data.

(4) The title, introduction, and conclusions sections place focus exclusively on the sublimation aspects of this work. I think this undersells and undervalues the importance of the broader Lagrangian isotope model approach and its applications. Sublimation appears to only relate to the easterly trajectory subset (< 50% of trajectories investigated). I am not sure if there is a companion paper planned/submitted/published for the (north)westerly trajectories but these trajectories seem important to discuss in more detail as well, even with a single summary paragraph somewhere in Section 4 of what was learned from modeling these trajectories. If manuscript focus is to be exclusively on sublimation effects, as the title implies, I don't think all trajectory data should be included (i.e., Fig. 5) and the δD record should focus on 'cold snap' trajectories. Currently, there seems to be too much data shown in Figures 5-7 that receive too cursory of a discussion.

Our reply: We added the following paragraph to Sect. 4.2: *"With respect to surface evaporation the air masses from the West and from the East contain very different types of information. The air masses from the West are exposed to moisture uptake from the ocean and to continental evapotranspiration at relatively warm temperatures. Therefore they contain information about isotope fractionation during evaporation from the ocean, evaporation from warm land surfaces, and plant transpiration. This makes the modeled δD of westerly air masses especially sensitive to simplifying model assumptions regarding the δD of moisture from continental evapotranspiration at warm skin temperatures. Increasing for instance the fraction of plant transpiration on total evapotranspiration from 0.7 to 0.8 in the model increases the mean modeled δD from summer by 2.0‰. Assuming δD values of soil water which are systematically increased by +10‰ increases the mean modeled δD from summer by +2.4‰. In contrast to the westerly air masses, the easterly air masses during cold snaps are sensitive to isotope fractionation during surface evaporation at temperatures where snow and meltwater exist. For these easterly air masses assumptions regarding evapotranspiration at warm skin temperatures only have a small impact."*

We replaced the seasonal panels in Fig. 5 by one source region map for the whole time period covered by the isotope observations in Karlsruhe and shortened the respective caption. In addition, we shortened Section 4.1 and combined it with Section 4.2. In Fig. 6 we kept the whole data for demonstrating the exceptional temperature and δD during the cold snaps. In Fig. 7 we also kept all data, because it underlines the capability of the model of reproducing observations over a wide range of temperatures and for different synoptic conditions.




Figure 5. Source regions of moisture (q) 30m above ground level in Karlsruhe (black star) based on five-day back trajectories for the time period January 2012 to May 2013. The color code indicates the contribution of different source regions to q in Karlsruhe in % per km^2 . Integration over the whole map gives the total identified humidity of 47%.

Technical comments

p. 1, Line 16: 'isotopologues' of water more appropriate since molecules are listed

Our reply: changed

p. 2, Lin4 4: somewhat misleading to say snowpacks are most sensitive to isotopic modification. Lakes, for example, generally provide longer-duration integration of post-depositional modification.

Our reply: We changed this sentence to "*Most problematic in that context are systematic post-depositional changes of isotope ratios into one direction, which most likely affect water reservoirs with long exposure to the atmosphere such as water intercepted in the snowpack.*"

p. 2, Lines 7-8: Here and throughout I think 'fractioning' should be replaced with 'fractionating'. I am unfamiliar with the term 'fractioning' with regards to isotopes.

Our reply: changed

p. 2, Line 8: clarify what is meant by 'other part'

Our reply: We changed the sentence to "*To explain the observations, the authors suggested non-fractionating sublimation with part of the sublimated vapor fractionating recondensing and part of the sublimated vapor escaping the snow layer.*"

p. 2, Line 13: It might be helpful to introduce the delta notation earlier in the manuscript so that it can be used here, i.e., 'increase of δ values'

Our reply: We moved the definition of δD into the introduction: "*In consequence, not only specific humidity and dew point temperature but also the isotope concentration ratio $R_D = [HD^{16}O]/[H_2^{16}O]$ – commonly referred to as $\delta D = R_D/R_{D,VSMOW} - 1$ with $R_{D,VSMOW} = 0.00031152$ – decreases in a cooling and raining air mass.*"

p. 2, Line 21: What is meant by 'resublimation'?

Our reply: Corrected. What we meant is "*deposition*".

p. 4, Lines 12-13: Quantify what is meant by 'fast' – sub-diurnal?

Our reply: We replaced 'Fast and random variations' (now on p.15) by "*Potentially artificial diurnal variations [...]*" and 'may smooth out real diurnal variations' (now on p.15) by "*[...] may smooth out real sub-diurnal and diurnal variations [...]*"

p. 4, Line 15: What is meant by 'displacement' of moisture uptake amount?

Our reply: We mean by 'displacement' that pronounced moisture uptake at a certain location is interpreted as continuous moisture uptake over a longer time interval in consequence of smoothing q . As we added more expanded clarifying information regarding the smoothing, this seems to be clear enough. So we removed 'displacement' from the sentence (now on p.15): "*Arbitrarily choosing a width of 24 h may smooth out real sub-diurnal and diurnal variations of q and thereby, may result in an underestimation of the amount of moisture uptake.*"

p. 5, Line 3: I think the phrasing 'more likely' should be changed – 'preferential fractionation of D into the liquid phase' is more precise language.

Our reply: We changed the respective sentence to "*Because of preferential fractionation of D into the liquid phase, the formation of precipitation results in a decreasing isotope concentration ratio ($R_D = [HD^{16}O]/[H_2^{16}O]$) in an air mass.*"

p. 5, Line 25: Clarify what is meant by 'not affected by dilution'

Our reply: For clarification we changed the paragraph to "*If the entrained moisture from evaporation was transported via small-scale turbulence to the trajectory altitude, mixing with air from below the trajectory level is likely. However, applying the above equations, changes of R_D and q in consequence of mixing with low-level air masses are ignored. We therefore implicitly assume that the air masses below the trajectory level experienced a similar transport and precipitation history as the tracked air parcel. R_D and q of the tracked air parcel are not affected by mixing with that air from below, only the by freshly evaporated moisture.*"

p. 6, Line 15: Where does the 2.6 day value come from?

Our reply: Ten-minutely measurements of precipitation amount at the measurement site indicate precipitation with more than 0.1mm per ten minutes on average every 2.6 days. As precipitation per day might be a more intuitive unit, we changed the sentence to: *“For instance measurements of precipitation amount at the measurement site in Karlsruhe indicate recharge of soil water around the site by more than 1mm precipitation per day on average every 2.9 days.”*

p. 11, Line 1: Here and elsewhere, should use ‘value’ instead of ‘ratio’ to describe δD

Our reply: replaced

p. 11, Line 4: I think the statement ‘caused by the relation between δD and condensation temperature’ is misleading. The more direct control on the continental effect is degree of rainout.

Our reply: We replaced ‘condensation temperature’ by *“degree of rainout”*

p. 12, Lines 18-19: Redundant sentences: ‘Second...’ And ‘To this end...’

Our reply: We removed the two sentences.

p. 12, Lines 20-21: Rationale for only considering trajectories above 28% median level needs to be more clearly communicated.

Our reply: We clarified rationale for only considering these trajectories: *“The air masses with the smallest moisture uptake within the last five days are least sensitive to the δD of moisture from surface evaporation and therefore don’t allow to robustly evaluate the model description of isotope fractionation during the sublimation of snow sublimation and the evaporation of meltwater. For a meaningful interpretation, we therefore only used the half of data with a contribution from surface evaporation at $T_{skin} < 0$ higher than 28%.”*

p. 13, Lines 4-5: The term ‘side constraints’ here and following is unclear to me. Can you clarify what is meant by ‘side’?

Our reply: We replaced the term “side constraints” by *“model assumptions”*.

p. 15, Line 29: Clarify what ‘both groups’ refer to.

Our reply: We changed the sentence to: *“To simultaneously bring into agreement modeled and observed δD of data corresponding to group “cold” and group “warm”, we suggest the existence of two regimes of T_{skin} with predominant non-fractionating sublimation in the colder regime and predominant fractionating evaporation of meltwater in the warmer regime.”*

- Dahlke, H. E., & Lyon, S. W. (2013). Early melt season snowpack isotopic evolution in the Tarfala valley, northern Sweden. *Annals of Glaciology*, *54*(62), 149–156. <http://doi.org/10.3189/2013AoG62A232>
- Gurney, S., & Lawrence, D. (2004). Seasonal trends in the stable isotopic composition of snow and meltwater runoff in a subarctic catchment at Okstindan, Norway. *Nordic Hydrology*, *35*(2), 119–137.
- Moser, H., & Stichler, W. (1974). Deuterium and oxygen-18 contents as an index of the properties of snow covers. *International Association of Hydrological Sciences Publication*, *114*, 122–135.
- Risi, C., Ogée, J., Bony, S., Bariac, T., Raz-yaseef, N., Wingate, L., ... Guglielmo, F. (2016). The Water Isotopic Version of the Land-Surface Model ORCHIDEE: Implementation, Evaluation, Sensitivity to Hydrological Parameters. *Hydrology: Current Research*, *7*(4), 2157–7587. <http://doi.org/10.4172/2157-7587.1000258>

The influence of snow sublimation and meltwater evaporation on ~~stable isotopes~~ δD of water vapor in the atmospheric boundary layer of Central Europe

Emanuel Christner^{1,2}, Martin Kohler¹, and Matthias Schneider²

¹Institute of Meteorology and Climate Research – Department Troposphere (IMK-TRO), Karlsruhe Institute of Technology (KIT), Germany

²Institute of Meteorology and Climate Research – Department Atmospheric Trace Gases and Remote Sensing (IMK-ASF), Karlsruhe Institute of Technology (KIT), Germany

Correspondence to: E. Christner (emanuel.christner@kit.edu)

Abstract. Post-depositional fractionation of stable water isotopes due to fractioning-fractionating surface evaporation introduces uncertainty to various isotope applications such as the reconstruction of paleotemperatures, paleoaltimetry, and the investigation of groundwater formation. In this study, we investigate isotope fractionation ~~during the evaporation of snow at ground level~~ at snow covered moisture sources by combining 17 months of observations of isotope concentration ratios [HD¹⁶O]/[H₂¹⁶O] in low-level water vapor in Central Europe with a new Lagrangian isotope model. The isotope model is capable of reproducing variations of the observed isotope ratios with a correlation coefficient R of 0.82. Observations from 38 days were associated with cold snaps and moisture uptake in snow covered regions. Deviations between modeled and measured isotope ratios during the cold snaps were related to differences in skin temperatures (T_{skin}). Analysis of T_{skin} provided by the Global Data Assimilation System (GDAS) of the NCEP implies the existence of two regimes of T_{skin} with different types of isotope fractionation during evaporation: A cold regime with $T_{\text{skin}} < T_{\text{subl,max}} = -7.17.7^\circ\text{C}$, which is dominated by non-fractioning-non-fractionating sublimation of snow and a warmer regime with $T_{\text{subl,max}} < T_{\text{skin}} < 0^\circ\text{C}$, which is dominated by fractioning-fractionating evaporation of meltwater. Based on a sensitivity study, we assess an uncertainty range of the determined $T_{\text{subl,max}}$ of ~~$-11.5-11.9$~~ to ~~$-2.52.9$~~ $^\circ\text{C}$. The existence of the two fractionation regimes has important implications for the interpretation of isotope records from snow covered regions as well as for a more realistic modeling of isotope fractionation at snow covered moisture sources. For these reasons, more detailed experimental studies at snow covered sites are needed to better constrain the $T_{\text{subl,max}}$ and to further investigate isotope fractionation in the two regimes.

1 Introduction

The hydrological cycle of the atmosphere is usually investigated by water vapor concentration measurements. A new dimension is opened by the analysis of H₂O stable isotope ratios, which are modified by H₂O phase changes during evaporation, cloud formation, and in-cloud physics. The main reason for this fractionation is the difference between vapor pressures of ~~the stable isotopes~~ stable isotopologues such as H₂¹⁶O and HD¹⁶O, which results in a preferential condensation of HD¹⁶O. In consequence, not only specific humidity and dew point temperature but also the isotope concentration ratio $R_D = [\text{HD}^{16}\text{O}]/[\text{H}_2^{16}\text{O}]$ ~

commonly referred to as $\delta D = R_D / R_{D,VSMOW} - 1$ with $R_{D,VSMOW} = 0.00031152$ – decreases in a cooling and raining air mass. The resulting relation between condensation temperature and isotope ratios of water vapor or precipitation is the basis for a variety of applications. Water isotopes in ice cores are used for the high-resolution reconstruction of paleotemperatures (Dansgaard, 1964; Masson-Delmotte et al., 2005). The temperature-induced gradient of isotope ratios with altitude makes water isotopes in precipitation a proxy for paleotopography (Poage, 2001; Rowley et al., 2001; Blisniuk, 2005; Rowley and Garzzone, 2007). Hydrological studies exploit the altitude-dependence and a seasonality of isotope ratios in precipitation to investigate groundwater formation (de Vries and Simmers, 2002).

Isotope fractionation during evaporation at the ground level post-depositionally modifies the isotope ratio of water at the surface and in the upper soil layers (Barnes and Allison, 1983). Because the various isotope applications rely on a close relation between the isotope ratio of precipitation and the isotope ratio of water at the surface, post-depositional effects increase the uncertainty of the isotope applications. Most problematic in that context are systematic post-depositional changes of isotope ratios into one direction, which most likely affect water intercepted in the snowpack because of reservoirs with long exposure to the atmosphere such as water intercepted in the snowpack.

Several studies report post-depositional enrichment of heavy isotopes in the snowpack. Such enrichment was observed over a wide range of temperatures and even at skin temperatures far below the freezing point. Epstein et al. (1965) observed enhanced isotope ratios δD values in depth-hoar layers in South Pole firn. To explain the observations, the authors suggested non-fractionating sublimation and subsequent fractionating recondensation of a non-fractionating sublimation with part of the vapor combined with the escape of the other sublimated vapor fractionating recondensing and part of the vapor sublimated vapor escaping the snow layer. Moser and Stichler (1974) analyzed surface layer snow from Switzerland (2450 m a.s.l.) during an eight-day fair weather period with air temperatures between about -5 and 0°C . They observed a continuous increase of isotope ratios δD values in the aging surface layer snow, which they attributed to fractionating evaporation or sublimation of snow. Stichler et al. (2001) report a similar experiment from the Chilean Andes (5536 m a.s.l.) with air temperatures between about -12 and -5°C . They observed an increase of isotope ratios δD values in the aging surface layer snow as well. They attributed this increase to kinetic fractionation during sublimation, caused by smaller coefficients of diffusion of the heavier isotopes. Based on a seasonal increase of isotope ratios δD values in a snowpack in northern Norway (~ 900 m a.s.l.), Gurney and Lawrence (2004) suggested fractionating evaporation of meltwater and subsequent recrystallization of residual meltwater. Consistent with this, Lechler and Niemi (2011) report an altitude-dependent seasonal modification of isotope ratios δD values in an alpine snowpack, which they attributed to fractionating evaporation of meltwater during ablation.

A clear assignment of such observations to a specific type of interaction between snowpack and atmosphere or to certain meteorological conditions is difficult, as continuous time series of high-resolution isotope profiles of the snowpack are hard to obtain. Furthermore, processes such as meltwater percolation, diffusion, sublimation, and resublimation-deposition within the snowpack additionally modify the profiles of isotope ratios, making an interpretation even more challenging. The current understanding of fractionation during the evaporation of snow is therefore highly uncertain and stretches from non-fractionating non-fractionating layer-by-layer sublimation of snow without any modification of isotope ratios in the snow-

pack (Ambach et al., 1968; Dansgaard, 1973; Friedman et al., 1991) to systematic enrichment of heavy isotopes in the snow in consequence of ~~fractioning-fractionating~~ evaporation of meltwater (Gurney and Lawrence, 2004; Lechler and Niemi, 2011). This uncertainty complicates interpretation of isotope records from snow covered regions, as the contribution of the various potential post-depositional effects may be different during climatologically different time periods. In addition, this uncertainty
5 limits the isotope modeling of atmospheric moisture sources in snow covered regions, as isotope-enabled models in general only consider one of the types of isotope fractionation for the sublimation of snow (e.g. Yoshimura et al., 2006; Werner et al., 2016).

In this context, continuous observations of isotope ratios of low-level water vapor may provide new insights by offering the opportunity to investigate fractionation during evaporation from the snowpack from a complementary point of view. A case
10 study by Noone et al. (2013) investigates local variations of ~~isotope ratios- δD values~~ in water vapor subsequent to a winter storm. Based on observations on a research tower the study partitioned the different surface fluxes assuming local evapotranspiration to consist of ~~non-fractioning sublimation and fractioning~~ non-fractionating sublimation and fractionating evaporation of meltwater.

A promising way to extend the investigation of isotope fractionation during evaporation of snow to the remote moisture
15 source regions is the combination of isotope observations with Lagrangian isotope modeling. A number of earlier studies applied Lagrangian isotope modeling along idealized, climatological trajectories (Jouzel and Merlivat, 1984; Johnsen et al., 1989) or along individual back trajectories (Schlosser et al., 2004; Helsen et al., 2004, 2005, 2007; Sodemann et al., 2008a; Pfahl and Wernli, 2009). As these studies mainly focused on polar or marine regions, the applied Lagrangian isotope models are not optimized to simulate continental evapotranspiration in the middle latitudes or evaporation of snow or meltwater at
20 temperatures close to the freezing point.

In this paper, we present a time series with 17 months of continuous measurements of ~~R_D - δD~~ in low-level water vapor in Central Europe. The measurements cover two winters, which were marked by a number of cold snaps and related snowfall. By combining ~~isotope the δD~~ observations during the cold snaps with a new Lagrangian isotope model, we investigate isotope fractionation ~~during sublimation or evaporation of surface layer snow~~ at snow covered moisture sources.

25 In the following, we present the Lagrangian isotope model (Sect. 2) and characterize the uncertainty of our ~~isotope- δD~~ measurements (Sect. 3). In Sect. 4, we relate variations of the ~~isotope ratios- δD~~ to air temperature and specific moisture source regions during cold snaps. In Sect. 5, ~~isotope ratios- δD values~~ during cold snaps are analyzed with respect to ~~fractionation during evaporation or sublimation of snow at the surface~~ isotope fractionation during surface evaporation in snow covered regions.

2 Lagrangian moisture diagnostics

30 2.1 Back trajectories

Isotope ratios to be analyzed in this paper were measured at a site near Karlsruhe in Central Europe (49.10° N, 8.44° E, 110.4 m a.s.l.). Kinematic five-day back trajectories from the site were calculated with the Hybrid Single Particle Lagrangian Integrated Trajectory Model 4.0 (Draxler and Hess, 1998) with a time resolution of 1 h. Three-dimensional wind fields for trajectory cal-

5 culation were derived from the Global Data Assimilation System (GDAS) of the National Centers for Environmental Prediction (NCEP) (Kanamitsu, 1989; Derber et al., 1991; Parrish and Derber, 1992). The GDAS data (V1.5) is available for every 3 h at a $1 \times 1^\circ$ horizontal resolution and on 23 sigma pressure levels between 1000 and 20 hPa. To account for uncertainty of the back trajectories, we used trajectory ensembles. Each ensemble consists of nine trajectories, starting 30 m above ground level at the measurement site as well as 50 km N, NE, E, SE, S, SW, W, and NW from the site. Back trajectories were calculated for every three hours of measurement time and were initialized at 0, 3, 6, 9, 12, 15, 18, and 21 UTC.

2.2 Lagrangian diagnostic of moisture sources

In order to identify major source regions of low-level water vapor in Karlsruhe, we applied a Lagrangian source region analysis similar to the method described by Sodemann et al. (2008b). The method traces air parcels along kinematic five-day back trajectories and analyzes changes of specific humidity (q) in time intervals of one hour. Such changes are possible due to the formation of precipitation (P), evaporation from the ground (E), evaporation of falling rain, air mass mixing due to convection or small-scale turbulence, diffusion, and numerical errors. In addition, using GDAS wind fields with three-hourly time resolution for the calculation of back trajectories – which is a coarse time resolution compared to the GDAS time steps on the order of minutes – may cause deviations between the HYSPLIT back trajectories and the exact trajectories that air parcels followed in the GDAS. This, in turn, may result in artificial changes of q along the HYSPLIT back trajectories. For instance HYSPLIT trajectories not fully capturing diurnal vertical movement in consequence of thermal expansion of the atmospheric boundary layer in the GDAS may show artificial diurnal changes of q . Like Stohl and James (2004) and Sodemann et al. (2008b), we assume P and E to be the dominant processes and ignore the other effects. Using this simplification to avoid an overestimation of the formation of precipitation and moisture uptake in consequence of potentially artificial diurnal variations of q along the HYSPLIT trajectories, we smoothed q along the back trajectories with a 24 h rectangle function. Using these simplifications, the change of specific humidity (Δq) per time step (Δt) is:

$$\Delta q / \Delta t = (E - P) / \Delta t. \quad (1)$$

Assuming further that either E or P are dominating (James et al., 2004), the net change $\Delta q / \Delta t$ per time step is attributed to only E or to P . Corresponding to these assumptions, an identified decrease of specific humidity is attributed to the formation of precipitation. In cases of a positive increment of specific humidity, the method assumes moisture uptake from evaporation at ground level. In this case, the contribution of surface evaporation (f_m) in a time interval m to total specific humidity at the end of this time interval (q_m) is

$$f_m = \frac{\Delta q_m}{q_m}. \quad (2)$$

The formation of precipitation in a later time interval does not affect the f_m calculated for the earlier time interval. In contrast to that, further moisture uptake in a later time interval n reduces the relative contribution of surface evaporation in the earlier time interval m to the moisture at the end of n . In this case, the contribution from earlier moisture uptake to q_n is recalculated

according to:

$$f'_m = f_m \cdot \frac{q_n - \Delta q_n}{q_n}. \quad (3)$$

Fast and random variations of q along the back trajectories could result in an overestimation of the formation of precipitation and moisture uptake. To avoid such an overestimation, fast variations of q were removed by smoothing q along the back trajectories with a 24h rectangle function. Arbitrarily choosing a width of 24h may smooth out real diurnal variations of q and thereby, may result in a displacement or underestimation of the amount of moisture uptake. Wherever necessary for the interpretation of our results, we therefore assess the related uncertainty of findings by changing the width of the rectangle function to 12 and 36h. In cases of moisture uptake above the atmospheric boundary layer an attribution to surface evaporation is not directly evident. A maximum altitude for consideration of moisture uptake might therefore be appropriate. However, Aemisegger et al. (2014) relate moisture uptake at higher levels for trajectories starting from Rietholzbach in northern Switzerland to the outflow of shallow convection. As moisture in Karlsruhe and Rietholzbach originates from similar source regions, we also do not assume a maximum altitude for the consideration of surface evaporation.

2.3 Lagrangian isotope model

A Lagrangian isotope model was developed to serve as a benchmark for our δD - δD measurements. Like the Lagrangian moisture source diagnostic, the model runs along kinematic five-day back trajectories and attributes changes of specific humidity (q) to the formation of precipitation or moisture uptake from surface evaporation. Analogous to the moisture source diagnostic, the model does not apply a maximum altitude for moisture uptake, smoothes q along the trajectories with a 24 h rectangle function, and uses the same trajectory ensembles. From each trajectory ensemble, nine modeled δD -ratios- δD values for Karlsruhe are obtained, which are combined to one value by calculating the average value by weighting the nine values with q -weighted average at the arrival.

2.3.1 Dehydration

A decrease of specific humidity in a time interval indicates the formation of precipitation. During the formation of precipitation, condensation of $HD^{16}O$ is more likely than condensation of $H_2^{16}O$, because of $HD^{16}O$'s lower vapor pressure. The Because of preferential fractionation of D into the liquid phase, the formation of precipitation therefore results in a decreasing isotope concentration ratio ($R_D = [HD^{16}O]/[H_2^{16}O]$) in an air mass. Assuming equilibrium conditions during condensation and subsequent immediate rainout of the condensate, we calculate-simulate the change of isotope ratios of the residual water vapor according to the Rayleigh distillation model (Rayleigh and Ramsay, 1894). Based on a fractionation factor $\alpha_D(T)$, this Rayleigh model calculates changes of R_D for infinitesimal changes of specific humidity q :

$$d \ln R_D = [\alpha_D(T) - 1] \cdot d \ln q \quad (4)$$

Specific humidity and air temperatures along the back trajectories were derived from the identical GDAS data set used for the calculation of the back trajectories. Under equilibrium conditions α_D only depends on the temperature (T) of an air parcel and increases from about 1.082 to ~~1.209~~ 1.240 between $+20^\circ\text{C}$ and -30 ~~40~~40 $^\circ\text{C}$. For $T \geq 0^\circ\text{C}$ we use a parametrization of α_D over liquid water (Horita and Wesolowski, 1994). For $T < 0^\circ\text{C}$, we assume enhanced fractionation over ice and use the parametrization of Jancso and Van Hook (1974). ~~Additional non-equilibrium ("kinetic") fractionation~~ A reduction of the fractionation factor $\alpha_D(T)$ in the case of supersaturation in ice clouds (Merlivat and Jouzel, 1979; Jouzel and Merlivat, 1984) is ignored by the model because of moderate temperatures and correspondingly low degrees of supersaturation in the examined source regions by a factor on the order of 0.964 to 1 between -40°C and 0°C was considered following Jouzel and Merlivat (1984), with a supersaturation parameter λ of 0.004 according to Risi et al. (2010).

10

2.3.2 Moistening

In the case that specific humidity increases in a time interval $[t_1, t_2]$, we assume a permeable air parcel which takes up moisture Δq by turbulent mixing. We attribute that moisture to evaporation at ground level with the isotope ratio $R_{D,E}$, which was lofted via small-scale turbulence or convection to the trajectory level. To calculate the corresponding change of R_D of moisture in the tracked air parcel, we apply the following mixing equations:

$$q(t_2) = q(t_1) + \Delta q, \quad (5)$$

$$R_D(t_2) = \frac{R_D(t_1) \cdot q(t_1) + \Delta q \cdot R_{D,E}(t_1)}{q(t_1) + \Delta q}. \quad (6)$$

If the entrained moisture from evaporation was transported via small-scale turbulence to the trajectory altitude, ~~dilution mixing~~ with air from below the trajectory level is likely. ~~Applying However, applying~~ the above equations, ~~we changes of R_D and q in consequence of mixing with low-level air masses are ignored. We~~ therefore implicitly assume that the air masses below the trajectory level experienced a similar transport and precipitation history as the tracked air parcel. ~~In this case,~~ R_D and q of the tracked air parcel are not affected by ~~dilution mixing~~ with that air from below, only the by freshly evaporated moisture.

Depending on the type of ground and skin temperature (T_{skin}), we calculate $R_{D,E}$ assuming evaporation from the ocean ($R_{D,E_{\text{ocean}}}$), continental evapotranspiration ($R_{D,E_{\text{ET}}}$), evaporation of melted snow ($R_{D,E_{\text{snowevap.}}}$), or sublimation of snow ($R_{D,E_{\text{snowsubl.}}}$).

For evaporation from the ocean, we assume ~~equilibrium vapor pressure~~ fractionation over liquid water with $\alpha_D(T_{\text{skin}})$ according to the parameterization of Horita and Wesolowski (1994). ~~Kinetic and T_{skin} at the trajectory position. To account for additional kinetic~~ fractionation during evaporation on the order of $\alpha_{D,\text{kin}} = 1.002 - 1.007$ (Merlivat and Jouzel, 1979; Pfahl and Wernli, 2009) due to different coefficients of diffusion of the different water ~~isotopes is ignored by the model isotopologues, we increased the α_D by the factor $\alpha_{D,\text{kin}} = 1.005$.~~ Please note that $\alpha_{D,\text{kin}}$ is much smaller than the vapor pressure fractionation factor α_D . Therefore, ~~consideration of kinetic fractionation considering a dependence of $\alpha_{D,\text{kin}}$ on environmental conditions~~ is less important for R_D than for other frequently used isotope ratios such as $[\text{H}_2^{18}\text{O}]/[\text{H}_2^{16}\text{O}]$. Given the above assumptions, the isotope ratio of evaporated moisture $R_{D,E_{\text{ocean}}}$ only depends on the ~~sea surface temperature (SST) surface temperature~~

and the isotope ratio of sea surface water ($R_{D,ocean}$, ~~which we derive from~~) at the trajectory position:

$$R_{D,E_ocean} = \frac{R_{D,ocean}}{\alpha_D(SST)} \frac{R_{D,ocean}}{\alpha_D(T_{skin})}. \quad (7)$$

We derived R_{D,E_ocean} from observations of δD and $\delta^{18}O$ of ocean surface water collected in the the Global Seawater Oxygen-18 Database (Schmidt et al., 1999). Since little data with the δD of ocean surface water exists, we calculated a median $\delta D/\delta^{18}O$ ratio of 6.56 from Fröhlich et al. (1988), Duplessy (1970), Delaygue et al. (2001), Gat et al. (1996), Ostlund et al. (1987), Aharon and Chappell (1986), Yobbi (1992), and Weiss et al. (1979) and used this ratio to calculate δD from $\delta^{18}O$ data. The $\delta^{18}O$ of ocean surface water along the back trajectories was derived from the spatial $1 \times 1^\circ$ interpolation (V1.1) of the Global Seawater Oxygen-18 Database by LeGrande and Schmidt (2006) (Fig. 1a), which we linearly interpolated to the locations along the trajectories. To derive skin temperatures representative of conditions during maximum evaporation (T_{skin}), we weighted skin temperatures along the trajectories ($T_{skin,unweighted}$) with positive surface latent heat flux in time intervals of ± 12 h (Fig. 2). If there were less than 12 trajectory points (time resolution of 1 h) with significant latent heat flux above 2 W m^{-2} in an interval, it was extended for ± 1 h until it contained 12 data points. $T_{skin,unweighted}$ and accumulated surface latent heat fluxes along the back trajectories were derived from a reduced GDAS data set with the same horizontal resolution of $1 \times 1^\circ$ as it was used for the calculation of the back trajectories but with data only every 6 h. The data was interpolated linearly in space and time to the locations along the trajectories. The accumulated surface latent heat flux from the GDAS was divided by six to account for the hourly resolution of the trajectories.

Over the continent, evapotranspiration consists of evaporation from the bare soil, transpiration of plants, and evaporation from canopy interception. As a first simplification we ignore canopy interception, i.e. we consider condensation with subsequent complete re-evaporation as a neutral process with respect to q and R_D . Moisture from the two other sources strongly differs in isotopic composition.

Evaporation from the bare soil is accompanied by isotope fractionation (Zhang et al., 2010). To calculate the isotope ratio of moisture evaporated from the bare soil ($R_{D,E_soilevap.}$), we assume ~~equilibrium fractionation~~. ~~Kinetic vapor pressure fractionation over liquid with $\alpha_D(T_{skin})$ according to the parameterization of Horita and Wesolowski (1994)~~. To account for kinetic fractionation during evaporation from the soil on the order of $\alpha_{D,kin}=1.017-1.025$ (Mathieu and Bariac, 1996) ~~is ignored~~. ~~For the calculation of $\alpha_D(T)$, we apply the parameterization of for equilibrium fractionation over liquid and use the skin temperature at the trajectory position~~. Since time of moisture uptake and evaporation might differ, we weighted skin temperatures along the back trajectories with surface evaporation within 24h (T_{skin}), we increased α_D by the factor 1.021. We further assume the isotope ratios of soil water to be the same as the isotope ratios of precipitation ($R_{D,prec.}$):

$$R_{D,E_soilevap.} = \frac{R_{D,prec.}}{\alpha_D(T_{skin})}. \quad (8)$$

Observations of $R_{D,prec.}$ have been collected in the Global Network of Isotopes in Precipitation (GNIP) (Araguas et al., 1996) of the IAEA since the 1960s. We used climatological monthly means of $R_{D,prec.}$ of the Regional Cluster-based Water Isotope Prediction (RCWIP) (Terzer et al., 2013), which provides a spatial interpolation of the GNIP data. RCWIP data is available with

a horizontal resolution of $0.17 \times 0.17^\circ$ (Fig. 1b/c) and was linearly interpolated to the locations along the trajectories. Because soil water around the measurement site is in Central Europe is generally frequently recharged by precipitation on average every 2.6 days, we ignore the systematic enrichment of HD^{16}O in the soil. We further assume the isotope ratio uppermost soil layer caused by continuous fractionating evaporation from the soil. For instance measurements of precipitation amount at the measurement site in Karlsruhe indicate recharge of soil water to be the same as the isotope ratio of precipitation ($R_{D,\text{prec.}}$); which we derive from the climatological monthly means of the Regional Cluster-based Water Isotope Prediction (RCWIP):

$$R_{D,E_{\text{soilevap.}}} = \frac{R_{D,\text{prec.}}}{\alpha_D(T_{\text{skin}})}.$$

around the site by more than 1 mm precipitation per day on average every 2.9 days. The assumption regarding enrichment is supported by findings of Risi et al. (2016), who observe insignificant systematic deviations of isotope ratios of water within the upper 15 cm of the soil from isotope ratios in precipitation at several sites in France, Germany, and the Czech Republic.

In contrast to bare soil evaporation, plants take up soil water from the soil and transpire that water completely and therefore with no fractionation into the atmosphere. Still some fractionation is possible on short timescales, due to asynchronous accumulation and the release of HD^{16}O and H_2^{16}O in leaves (Zhang et al., 2010). However, this process averages out over a day (Harwood et al., 1999; Farquhar et al., 2007) and is therefore ignored for modeling isotope ratios along the five-day back trajectories. To calculate the isotope ratio of moisture originating from plant transpiration, we assume:

$$R_{D,E_{\text{transp.}}} = R_{D,\text{prec.}}. \quad (9)$$

The isotope ratio of total evapotranspiration ($R_{D,E_{\text{ET}}}$) depends on the fraction of plant transpiration (FT) on total evapotranspiration:

$$R_{D,E_{\text{ET}}} = R_{D,E_{\text{soilevap.}}} \cdot (1 - \text{FT}) + R_{D,E_{\text{transp.}}} \cdot \text{FT} \\ = \frac{R_{D,\text{prec.}}}{\alpha_D(T_{\text{skin}})} \cdot (1 - \text{FT}) + R_{D,\text{prec.}} \cdot \text{FT}. \quad (10)$$

FT varies with region and on seasonal, synoptic, and diurnal timescales. For modeling we ignore these variations and use a constant fraction of transpiration. Based on Choudhury et al. (1998), Lawrence et al. (2007), and Aemisegger et al. (2014) we assume an average FT in Europe of 0.7.

Whenever we observe moisture uptake at continental skin temperatures below 0°C , we ignore transpiration of plants (FT=0) and attribute the moisture to the evaporation of melted snow or ice. In this case, we again assume equilibrium fractionation over liquid. We further assume the isotope ratio of snow to be the same as the climatological monthly means of the RCWIP:

$$R_{D,E_{\text{snowevap.}}} = \frac{R_{D,\text{prec.}}}{\alpha_D(T_{\text{skin}})}. \quad (11)$$

In Sect. 5 we investigate a possible role of snow sublimation. In this case we define a skin temperature $T_{\text{subl,max}}$. Below that temperature we assume complete layer-by-layer sublimation of snow without isotope fractionation:

$$R_{D,E_{\text{snowsubl.}}} = R_{D,\text{prec.}}. \quad (12)$$

2.3.3 Initialization

~~Forty-three percent of the analyzed air masses originated from oceanic regions and~~ For initializing of R_D of air masses originating from altitudes below 2 km ~~Forty-two percent originated from the continent and also from altitudes below 2km~~ above ground level. ~~For the initialization of R_D of these air masses,~~ we assume isotope ratios ($R_{D,ini}$) in a convectively well-
 5 mixed atmospheric boundary layer (ABL), where water vapor and ocean surface water or soil water are in isotopic equilibrium:

$$R_{D,ini,<2km} = \frac{R_{D,ocean|prec.}}{\alpha_D(SST|T_{skin})} \frac{R_{D,ocean|prec.}}{\alpha_D(T_{skin})}. \quad (13)$$

For initialization at surface temperatures above 0°C, we calculate fractionation factors according to the parametrization of Horita and Wesolowski (1994). At skin temperatures below 0°C we assume fractionation over ice and apply the parametrization
 10 of Jancso and Van Hook (1974).

~~Fifteen percent of air masses originated~~ For air masses originating from an altitude (h) higher than 2 km above ground level ~~For these air masses~~ we assume a linear decrease of $R_{D,ini}$ from boundary layer ratios at an altitude of 2 km to $R_{D,10 km}$ of 0.45 (Hanisco et al., 2007; Sayres et al., 2010) at an altitude of 10 km.

$$R_{D,ini,>2km} = R_{D,ini,<2km} - (R_{D,ini,<2km} - R_{D,10 km}) \frac{h - 2 \text{ km}}{10 \text{ km} - 2 \text{ km}}. \quad (14)$$

15 The dependence of modeled isotope ratios from initialization decreases with moisture uptake along the back trajectories (Fig. 43). The uncertainty of $R_{D,ini}$ is especially strong at high altitudes, where air masses are strongly dehydrated and have a long history of isotope fractionation. Because air masses from high altitudes take up a lot of humidity during descent and transport to Karlsruhe, the uncertainty of R_D from initialization is strongly reduced in Karlsruhe. Back trajectories corresponding to smaller moisture uptake typically originate from the ABL. R_D in Karlsruhe of these back trajectories depends more strongly on $R_{D,ini}$,
 20 which on the other hand is much better defined within the ABL. Considering typical atmospheric moisture residence times in the range of 4–8 days (Läderach and Sodemann, 2016; Trenberth, 1998), using ten-day back trajectories instead of five-day back trajectories should almost eliminate model uncertainty of the δD at Karlsruhe from uncertainty of $R_{D,ini}$. However, such long trajectories easily cover distances of 10,000 km, which makes the modeled δD sensitive to potentially very different conditions in far-distant regions. This, in turn, makes uncertainty assessment of the modeled δD more complex. Using five-day
 25 back trajectories is therefore a trade-off between a reasonably small sensitivity of the modeled δD at Karlsruhe on $R_{D,ini}$ and concentrating the analysis to the North Atlantic and Eurasia. Whenever it is necessary for the interpretation of our results, we assess ~~related~~ uncertainty of the modeled δD δD from the model initialization by changing $R_{D,ini}$ in different model runs.

3 Measurements

3.1 Isotope water vapor measurements

The concentrations of H_2^{16}O and HD^{16}O in low-level water vapor were measured for 17 months on a research campus 12 km north of Karlsruhe in Southwest Germany (49.10° N, 8.44° E, 110.4 m a.s.l.).

- 5 For the continuous measurements we used a Picarro water isotopologue analyzer L2120-i, which analyzed the ambient water vapor with a sampling rate of 0.6 Hz. The measurement technique is based on cavity ring-down spectroscopy, where the beam of a tunable diode laser is directed through a cavity, filled with the air to be analyzed. Based on the ring-down time of the laser light intensity, absorption spectra are measured between 7183.5 and 7184 cm^{-1} . A characterization of two similar analyzers (L1115-i and L2130-i) can be found in Aemisegger et al. (2012). Please note that the isotopologue analyzer also measures
- 10 concentrations of the water isotope H_2^{18}O . As the isotope concentration ratio $R_{18\text{O}} = [\text{H}_2^{18}\text{O}]/[\text{H}_2^{16}\text{O}]$ is more sensitive to kinetic fractionation than R_D , it would be essential to [more detailed](#) consider kinetic fractionation during evaporation for modeling $R_{18\text{O}}$ with the Lagrangian isotope model. Uncertainty of the modeled $R_{18\text{O}}$ related to uncertainty of the kinetic fractionation factor does not allow deeper insight from analysis of R_D and $R_{18\text{O}}$ than from analysis of R_D alone. Thus, we do not use the H_2^{18}O measurements in this paper.
- 15 The Picarro water isotopologue analyzer was located on the sixth floor of the Institute of Meteorology and Climate Research – Atmospheric Trace Gases and Remote Sensing of the Karlsruhe Institute of Technology. A downward-facing inlet funnel was installed one meter above the edge of the roof, which corresponds to an altitude above ground level of 28 m. The connection to the inlet was established with a 6 m long tube with a diameter of 6.4 mm. To reduce wall effects, we permanently flushed the inlet line with 30 slpm of air and used tubing made of electropolished stainless steel. To avoid condensation, the wall temperature
- 20 of the 5 m of tubing inside the building was regulated to 22°C. Saturation humidity corresponding to this temperature is above the analyzer limit of 14.9 $\text{g} \cdot \text{kg}^{-1}$. On five days in August 2012, humidity slightly exceeded that analyzer limit. Corresponding measurements were removed from the time series.

We report isotope measurements in [the](#) δ -notation, which normalizes isotope ratios to a standard scale, defined by the Vienna standard mean ocean water (VSMOW: $\delta D \delta D = 0\text{‰}$) and standard light Antarctic precipitation (SLAP: $\delta D \delta D = -428.0\text{‰}$)

- 25 (IAEA, 2009):

$$\delta D = \frac{R_D}{R_{D,\text{standard}}} - 1.$$

- For the automated calibration of the analyzer we applied a Picarro standard delivery module (A0101), which allows the alternating injection of two different water standards into a Picarro vaporizer (A0211). In this vaporizer the liquid standards immediately evaporate in a constant flow of dry synthetic air (140°C, 0.3 slpm dry air flow with 1.2 $\text{mg} \cdot \text{kg}^{-1}$ residual humidity).
- 30 A two-point calibration was done for 2 h every 10 h at $\delta D \delta D$ of -62.1‰ and -142.2‰ .

Instrumental drift during the 17 months was below 3‰ (Fig. 24a). The total accuracy of our $\delta D \delta D$ measurements due to uncertainty of calibrations and instrumental drift between two calibrations was 0.98‰. The 0.6 Hz precision of the measurements was below 1‰ and can be ignored for 10-minute averages shown in this paper.

Based on the two-point calibrations, we applied linear stretching to the measurements. Sixty-three percent of our observations are within the range of isotope ratios covered by the two standards. To approve linearity of the applied correction for isotope ratios below that range, we performed repeated calibrations with a third standard ($\delta D/\delta D = -245.3\%$) in the two years subsequent to the campaign. We found additional uncertainty of $\delta D/\delta D$ at -245.3% due to a slight non-linearity of the applied correction to be smaller than 0.3 ‰, which is in agreement with the more detailed characterization of Aemisegger et al. (2012).

To identify a potential humidity dependence of the isotope ratio measurements, we generated three humidity levels between $1.8 \text{ g} \cdot \text{kg}^{-1}$ and $13.7 \text{ g} \cdot \text{kg}^{-1}$ during each calibration. We found the humidity dependence of the instrument to be smaller than the uncertainty of individual calibrations. Therefore, we only applied the average humidity dependence of the whole found using all calibrations of $-0.021 \text{ ‰}/(\text{g} \cdot \text{kg}^{-1})$ to the data set (Fig. 24b).

3.2 Meteorological data at the measurement site

Observations of specific humidity at the measurement site were derived from the the Picarro isotope analyzer. For calibration of the Picarro humidity measurements we used observations of a VPT6 Thygan dew point mirror hygrometer (Meteolabor, Switzerland), which was mounted on a meteorological tower 30 m above ground level 900 m WSW in the west-southwest. Since the topography at the measurement site is flat for some kilometers in all directions, we assume the tower observations to be representative for the measurement site. The dew point hygrometer performed a measurement of one minute every ten minutes and has an uncertainty of $\pm 0.1 \text{ K}$. Ten-minute averages of specific humidity derived from the Picarro Analyzer and observations of the dew point hygrometer show a correlation coefficient R of 0.9913. For the calibration of the Picarro humidity observations we applied the mean linear regression between hygrometer data and Picarro measurements of 1.13, which varied by 1% between the first and second half of the measurement period.

The amount of precipitation was measured at the meteorological tower at ground level with a time resolution of ten minutes.

3.3 Isotope ratios in the moisture source regions

Observations of δD and $\delta^{18}\text{O}$ of ocean surface water are collected in the the Global Seawater Oxygen-18 Database. Since little data with δD of ocean surface water exists, we calculated a median $\delta D/\delta^{18}\text{O}$ ratio of 6.56 from , , , , , and and used this ratio to calculate δD from $\delta^{18}\text{O}$ data. The $\delta^{18}\text{O}$ of ocean surface water along the back trajectories was derived from the spatial 1×1 interpolation (V1.1) of the Global Seawater Oxygen-18 Database by (Fig. 3a), which we linearly interpolated to the locations along the trajectories. For the estimation of δD of soil water along the back trajectories, we assumed soil water to have the same isotopic composition as precipitation. Observations of isotope ratios in precipitation have been collected in the Global Network of Isotopes in Precipitation (GNIP) of the IAEA since the 1960s. We use climatological monthly means of δD of the Regional Cluster-based Water Isotope Prediction (RCWIP), which provides a spatial interpolation of the GNIP data. RCWIP data is available with a horizontal resolution of 0.17×0.17 (Fig. 3b/e) and was linearly interpolated to the locations along the trajectories.

3.3 Meteorological data in the moisture source regions

Specific humidity and air temperatures along back trajectories were derived from the identical GDAS data set used for the calculation of the back trajectories. Skin temperatures ($T_{\text{skin,unweighted}}$), accumulated surface latent heat flux, and a flag for snow cover along the back trajectories were derived from a reduced GDAS data set with the same horizontal resolution of 1×1 but with data only every 6h. The data was interpolated linearly in space and time to the locations along the trajectories. The accumulated surface latent heat flux from the GDAS was divided by six to account for the hourly resolution of the trajectories. To derive skin temperatures representative of conditions during maximum evaporation (T_{skin}), we weighted the $T_{\text{skin,unweighted}}$ along the trajectories with positive surface latent heat flux in time intervals of 12h (Fig. 4). If there were less than 12 trajectory points (time resolution of 1h) with significant latent heat flux above 2 in an interval, it was extended for 1h until it contained 12 data points.

4 Analysis of seasonal and synoptic variations

~~In this section,~~ In this section, we present the measurements from Karlsruhe, covering the time period from January 2012 to May 2013. For this time period, we identify specific circulation regimes related to cold snaps in Karlsruhe. Subsequent to this, we examine the capability of the Lagrangian isotope model of reproducing corresponding variations of δD .

4.1 Source regions of moisture

~~δD .~~ For the

4.1 Continental temperatures and zonal circulation

According to five-day back trajectories, 85% of the sampled low-level air masses at Karlsruhe originated from altitudes below 2 km above ground level. Consequently, most of the air masses were exposed to continuous moisture uptake from surface evaporation during transport to Karlsruhe. For identification of major source regions of low-level the water vapor in Karlsruhe we applied the Lagrangian moisture source diagnostic described in Sect. 2.2. ~~During all seasons~~ Based on this analysis, we found the North Atlantic was to be the most important moisture source (Fig. 5), from where westerlies transported the tracked air masses to the measurement site. ~~Due to a weakening~~ In addition to the predominantly westerly moisture transport, inversions of zonal circulation in summer, the source region was smallest in this season and concentrated on the eastern part of the North Atlantic. In winter, the source region was more extended, in consequence of a more pronounced zonal circulation, and expanded as far as the East Coast of winter occasionally led to easterly moisture transport. Because of the finite length of the five-day back trajectories, the total identified humidity is lower than 100%. If smoothing specific humidity along the back trajectories for 24 h, the total identified humidity accounts for 47%. If smoothing specific humidity for 12 h, diurnal and sub-diurnal variations in humidity are interpreted as the United States. In addition, occasional inversions of zonal circulation in winter led to easterly moisture transport.

4.2 Continental temperatures and zonal circulation

~~On a seasonal timescale air formation of precipitation and moisture uptake, which increases the total identified humidity to 63%. Both numbers are in reasonable agreement with more extended studies implying global atmospheric moisture residence times in the range of 4–8 days~~ (Läderach and Sodemann, 2016; Trenberth, 1998).

5 Air temperatures, specific humidity, and $\delta D/\delta D$ in Karlsruhe followed ~~a similar pattern~~ similar seasonal patterns (Fig. 6). In winter (~~DJF~~ December, January, February), air temperatures 30 m above ground level (T) were on average 2°C . Towards summer (~~JJA~~ June, July, August), T increased to on average 20°C . Higher T in summer corresponds to a higher saturation vapor pressure. That allows the transport of marine air to Karlsruhe with less condensation in summer than in winter. Consequently specific humidity (q) in Karlsruhe rose from $6\text{ g}\cdot\text{kg}^{-1}$ (DJF) to $14.8\text{ g}\cdot\text{kg}^{-1}$ (JJA). $\delta D/\delta D$ changed from -162% (DJF) to
10 -109% (JJA) and thereby consistently with T and q implies a lower degree of condensation and rainout in summer than in winter.

The gray color in Fig. 6 identifies circulation regimes with easterly moisture transport (25% of data). Such regimes predominantly occurred in winter and resulted in the transport of continental air masses to Karlsruhe. The corresponding air masses usually were marked by especially low T and q , which led to pronounced cold snaps in Karlsruhe. Consistent with the low T
15 and q , the air masses during cold snaps showed an especially low $\delta D/\delta D$ value.

Both findings - the seasonality of $\delta D/\delta D$ and the especially low $\delta D/\delta D$ in cold, continental air masses from the East - are in good agreement with the well-known "continental effect". That effect describes a decrease of $\delta D/\delta D$ in precipitation over continents with distance to the coast (Fig. ~~3b/e1b~~), caused by the relation between ~~δD and condensation temperature~~ δD and
20 degree of rainout. Since the $\delta D/\delta D$ of rain depends on the $\delta D/\delta D$ of the water vapor it is formed from, it is reasonable to find a similar continental effect imprinted to water vapor as well.

4.2 Comparison of measured and modeled $\delta D/\delta D$

The Lagrangian isotope model described in Sect. 2.3 is able to reproduce the observed slow seasonal variation of $\delta D/\delta D$, as well as the strong and relatively fast variations of $\delta D/\delta D$ due to circulation regimes with predominantly easterly moisture
25 transport in winter (Fig. 7a/b). The mean difference between modeled and observed $\delta D/\delta D$ is $+1.0\%$ ($\Delta\delta D$) is -4.1% . The correlation coefficient R of modeled and observed $\delta D/\delta D$ is 0.82. Thereby, the correlation calculated for different seasons strongly differs from the overall correlation. For summer R is only 0.06 -0.03 due to the small variability of $\delta D/\delta D$ in this season. For winter R is 0.87.

Figure 7c shows the scatter plot of measured and modeled $\delta D/\delta D$. Furthermore, the figure illustrates the impact of the formation
30 of precipitation and surface evaporation on the modeled $\delta D/\delta D$. If the formation of precipitation is ignored in the model, the overall correlation with the observations is still 0.84 0.79 . The main reason for this is the relation of observations with low $\delta D/\delta D$ ratios δD values to easterly moisture transport. Corresponding back trajectories are initialized with relatively low $\delta D/\delta D$ values according to GNIP observations in the respective continental source regions. This means that we only ignore the forma-

tion of precipitation in the five days covered by the back trajectories, but we implicitly consider the formation of precipitation which determined the isotope ratios in precipitation in the moisture source regions. However, due to surface evaporation with relatively high δD ratios, the $\Delta\delta D$ values, the $\Delta\delta D$ in this scenario is +29.423.9‰. If one considers the formation of precipitation but ignores the surface evaporation the overall R is reduced to 0.62. The corresponding $\Delta\delta D$ is -35.634.8‰.

5 So consideration of both processes, surface evaporation and the formation of precipitation ~~and surface evaporation~~ is essential for reproducing the observed δD .

With respect to surface evaporation the air masses from the West and from the East contain very different types of information. The air masses from the West are exposed to moisture uptake from the ocean and to continental evapotranspiration at relatively warm temperatures. Therefore they contain information about isotope fractionation during evaporation from the ocean, evaporation
10 from warm land surfaces, and plant transpiration. This makes the modeled δD of westerly air masses especially sensitive to simplifying model assumptions regarding the δD of moisture from continental evapotranspiration at warm skin temperatures. Increasing for instance the fraction of plant transpiration on total evapotranspiration from 0.7 to 0.8 in the model increases the mean modeled δD from summer by 2.0‰. Assuming δD values of soil water which are systematically increased by +10‰ increases the mean modeled δD from summer by +2.4‰. In contrast to the westerly air masses, the easterly air masses during
15 cold snaps are sensitive to isotope fractionation during surface evaporation at temperatures where snow and meltwater exist. For these easterly air masses assumptions regarding evapotranspiration at warm skin temperatures only have a small impact.

An interesting feature in Fig. 7a are spikes of low δD values (blue), which are not reproduced by the model (red). We frequently observed such spikes from spring to autumn. Potential processes causing the spikes are the evaporation of rain below the cloud base or isotope exchange between falling raindrops and the low-level water vapor. An impact of these sub-
20 cloud processes on the δD of low-level water vapor is demonstrated for individual weather fronts by Wen et al. (2008) and Aemisegger et al. (2015) and is complementarily supported by observations of isotope ratios in precipitation (Friedman et al., 1962; Stewart, 1975; Gedzelman and Arnold, 1994). As isotope processes below clouds are not represented in the Lagrangian isotope model, the observations related to sub-cloud processes can't be further investigated by means of this model. However, a relevant role of sub-cloud processes for our observations of δD spikes is supported by the strongly increased probability of
25 precipitation during the spikes. For respective observations with δD values smaller than expected from the standard deviation between observed and modeled δD (23.5‰) the probability to observe precipitation in Karlsruhe within ± 3 h was 44%, whereas for the other observations this probability was only 24%. For the investigation of δD during cold snaps in winter sub-cloud processes can be ignored, because the interaction between falling precipitation and water vapor is strongly suppressed in cases of solid precipitation.

30 5 δD during cold snaps

The good agreement of modeled and measured δD in winter (R=0.87) underlines the strong potential of the Lagrangian isotope model for analyzing isotope processes in the remote moisture source regions during cold snaps. In this section, we select respective observations and investigate isotope fractionation during sublimation or evaporation at ground level.

5.1 Sublimation of snow or snowmelt evaporation?

Evaporation below 0°C was historically often considered as ~~non-fractioning non-fractionating~~ layer-by-layer sublimation of snow and ice (Ambach et al., 1968; Dansgaard, 1973; Friedman et al., 1991), because of a low coefficient of self-diffusion of water molecules in ice. However, this assumption ignores snow melt and fractionation during evaporation from the liquid phase, as implied by Gurney and Lawrence (2004), Lechler and Niemi (2011), and Noone et al. (2013). Both assumptions result in very different δD - δD of moisture from evaporation. In the case of ~~non-fractioning non-fractionating~~ sublimation, this δD equals the δD - δD equals the δD of the snow. In the case of ~~fractioning fractionating~~ evaporation, the δD -ratio- δD value of moisture from evaporation is about 90% lower. Our trajectory model provides an opportunity to test both formulations and to assess which one is more reasonable for Central Europe.

10 ~~For the investigation of isotope fractionation during evaporation at low skin temperatures~~ According to the Lagrangian moisture source diagnostic, only 5% of the humidity analyzed in Karlsruhe from January 2012 to May 2013 originated from surface evaporation at locations with a GDAS based snow depth greater than 0.5 cm. For this reason, we used data fulfilling ~~two selection criteria~~ three selection criteria for the further investigation of isotope fractionation during evaporation from snow covered surfaces. First, we excluded observations which are strongly affected by evapotranspiration at skin temperatures (T_{skin}) above the freezing point. For this purpose, we identified moisture uptake at $T_{\text{skin}} > 0^{\circ}\text{C}$ by means of the Lagrangian-moisture source diagnostic and excluded air masses with a respective contribution above 2%. ~~Second, we ensured relevant moisture uptake at $T_{\text{skin}} < 0$. To this end, we identified moisture uptake at $T_{\text{skin}} < 0$.~~ For air masses meeting the first criterion the median contribution of moisture evaporated at $T_{\text{skin}} < 0^{\circ}\text{C}$ is 28%. The air masses with the smallest moisture uptake within the last five days are least sensitive to the δD of moisture from surface evaporation and therefore don't allow to robustly evaluate the model description of isotope fractionation during the sublimation of snow and the evaporation of meltwater. For a meaningful interpretation, we therefore only used the half of data with a contribution from surface evaporation at $T_{\text{skin}} < 0$ higher than 28%. Only 2% of the corresponding air masses originated from altitudes higher than 2000 m above ground level. Since uncertainty of model initialization is especially high for these air masses, we finally excluded the 2% of air masses originating from high altitudes.

25 Some ~~178-174~~ of the three-hourly modeled data points meet both the three selection criteria. They belong to 38 different days and were used for further interpretation. Respective air masses mainly originated from the East (Fig. 8) ~~and from altitudes below 2000 a~~. The mean fraction of moisture identified for these air masses by the moisture source diagnostic is 48%. If smoothing humidity along the back trajectories for 12 m above ground level instead of 24 h the mean identified fraction is 68%. The GDAS data indicates the existence of snow on the ground at 96% of locations along the selected back trajectories ~~(Fig. 8b) with a median snow depth of 1.8 ± 0.9 cm (\pm gives the interquartile range).~~ The median T_{skin} at the trajectory positions five days back was $-11.7 \pm 5.25.1^{\circ}\text{C}$ (~~\pm gives the interquartile range~~). During transport to Karlsruhe the T_{skin} rose on average by $6.0 \pm 3.43.3$ K. A decrease of relative humidity due to warming of the air masses was partially compensated by moisture uptake and a corresponding increase of specific humidity by on average $52 \pm 35\%$.

Assuming equilibrium fractionation during evaporation of meltwater at $T_{\text{skin}} < 0^{\circ}\text{C}$ in the reference run (M_{MW} , Table 1), the

model underestimates the selected δD -ratios- δD values by on average $\Delta\delta D/\Delta\delta D = -19.718.6 \pm 1.61.5\%$ (\pm gives the statistical uncertainty of the mean). Assuming ~~non-fractioning~~ non-fractionating sublimation at $T_{\text{skin}} < 0^\circ\text{C}$ in a further model run (M_S , Table 1) results in δD -ratios- δD values that are on average $\Delta\delta D/\Delta\delta D = +25.726.9 \pm 1.71.6\%$ above the observations.

Considering the relatively high δD -ratios- δD values of moisture from ~~non-fractioning~~ non-fractionating sublimation and the about 90% lower δD -ratios- δD values in the case of ~~fractioning~~ fractionating evaporation, the difference of mean δD - δD between both model runs is qualitatively reasonable. To judge if one run provides more realistic results, we tested if one of the two scenarios can be brought into agreement with the observations when considering uncertainty of ~~side constraints in the model~~ model assumptions.

As the most important ~~side constraints~~ source of uncertainty for modeling the δD in Karlsruhe, ~~we consider interannual variability of δD - δD in Karlsruhe during cold snaps~~ we consider variability of the δD of surface layer snow in the moisture source regions. ~~Such variability is not that is not fully~~ captured by the model ~~and therefore~~. This variability may systematically affect (1) the assumed ~~δD of snow~~ δD of moisture from surface evaporation as well as (2) ~~δD at the~~ the δD at the model initialization. In addition, (3) the amount of identified moisture uptake needs to be accurate to reliably simulate the impact of ~~sublimation or evaporation of snow on the δD~~ surface evaporation on the δD of water vapor.

In the following, we estimate uncertainty of ~~these side constraints~~ respective model assumptions. Subsequent to this, we vary the ~~side constraints~~ model assumptions in different model runs, to assess corresponding systematic uncertainty of the modeled δD - δD .

(1) The δD - δD of moisture from sublimation or evaporation of ~~snow-meltwater~~ depends on the δD of the snowpack δD of the surface layer snow, which we assume to be equal to the δD - δD of precipitation. The δD - δD of precipitation in an individual year may systematically differ from the used climatological monthly means of the Regional Cluster-based Water Isotope Prediction (RCWIP). To assess typical interannual variability of δD - δD of precipitation in winter, we used data from 134 European and Russian GNIP stations from the mid-latitudes between 8.4 and 50° E with observations from at least three years. For each of the stations we calculated the mean δD - δD in the different winters (November, December, January, and February) and the standard deviation of the winter averages. The mean of standard deviations of the different stations was 11.5%, which we assume to reflect the mean interannual variability of δD - δD in precipitation in the moisture source regions related to cold snaps.

~~In addition, the average monthly δD of snow may differ from the average δD of the monthly total precipitation recorded by the GNIP.~~ Because there is a general relation between surface air temperatures and δD - δD of precipitation in Central Europe (Schoch-Fischer et al., 1983; Jacob and Sonntag, 1991), winter months from years with especially high air temperatures and a potentially strong contribution from liquid precipitation are related to relatively high δD -ratios- δD values. Since we want to estimate ~~δD of the snowpack~~ the δD of snow, data from winters with especially low temperatures and a strong contribution from solid precipitation with low δD -ratios ~~is more suitable~~ δD values is likely to be most representative. During these winters the δD - δD of precipitation is probably closest to δD - δD of the RCWIP minus the 11.5%.

In contrast to this, post-depositional fractionation processes may increase the δD values of surface layer snow on the order of +10 to +20% during periods with small accumulation rates (Gurney and Lawrence, 2004; Moser and Stichler, 1974).

suggesting a scenario where the δD values of surface layer snow are higher than the δD from the RCWIP. Please note that this scenario is not likely for a seasonal snowpack during melt season since ablation may uncover old snow from colder winter months, causing changes of the δD of surface layer snow of -50% (Dahlke and Lyon, 2013). Given the relatively small snow depths in the investigated moisture source region (Fig. 8b), we ignore the uncovering of older snow layers and only consider a potential post-depositional increase of the δD of surface layer snow.

To test whether the too high modeled δD -ratios- δD values from the scenario of sublimation (M_S) can be significantly reduced when considering the systematic uncertainty of δD -of-the snowpack- δD of surface layer snow regarding interannual variability of the δD of snowfall, we performed one model run ($M_{S-,snow}$), in which we shifted δD -the δD of snow by -11.5% . To test whether the too low modeled δD -ratios- δD values from the scenario of evaporation of meltwater (M_{MW}) can be increased if considering a post-depositional increase of the δD of surface layer snow, we performed one further model run ($M_{MW+,snow}$), in which we shifted δD -of-snow-by-the δD of surface layer snow by the same absolute value of $+11.5\%$, although a positive shift is not as likely as a negative shift. The mean difference between modeled and observed δD ($\Delta\delta D$) ($\Delta\delta D$) of the different model runs is listed in Table 1.

(2) During cold snaps in Karlsruhe, on average 48% of humidity could be attributed to moisture uptake along the five-day back trajectories. The other side of this argument is that the δD - δD of 52% of humidity in Karlsruhe is determined by the initialization of isotope ratios.

For initialization below 2km above ground level, we assume δD - δD in a well-mixed atmospheric boundary layer and isotopic equilibrium between water vapor and the climatological monthly δD - δD of precipitation. This assumption is in agreement with one of the rare extended, simultaneous time series of δD - δD in water vapor and precipitation, conducted 45 km NNE from our site (Jacob and Sonntag, 1991). This study shows monthly averages of δD - δD in precipitation and water vapor at ground level for the years 1981 to 1988. The average deviation of the δD - δD of water vapor to isotopic equilibrium with precipitation in November, December, January, and February was $+4.1 \pm 7.7\%$ (\pm states the standard deviation calculated from the winter averages of the different years). Ignoring the interannually varying deviation between $4.1 - 7.7 = -3.6\%$ and $4.1 + 7.7 = +11.8\%$ from the isotopic equilibrium may systematically bias δD - δD at the model initialization.

For air masses originating from altitudes higher than 2km above ground level, the uncertainty of δD at initialization might be higher. Our model assumes a linear decrease of δD -ratios to -550 at an altitude of 10km. Since in situ observations of δD from the free troposphere are rare, uncertainty of this assumption is hard to assess. We assume that a systematic deviation of δD to the estimated profile is smaller than ± 100 . During cold snaps only 2 of the back trajectories originate from altitudes higher than 2km above ground level, so that uncertainty of the average modeled δD in Karlsruhe is only slightly affected by the uncertainty of an initialization in high altitudes. To test how much the values of $\Delta\delta D$ - $\Delta\delta D$ for M_S and M_{MW} can be reduced if considering the uncertainty of δD - δD at the model initialization, we performed two further model runs. For the scenario of sublimation we performed a model run in which we shifted δD - δD at the initialization for -3.6% ± 100 in cases of initialization below / above 2km above ground level ($M_{S-,ini}$). For the scenario of evaporation of meltwater we shifted δD - δD at the initialization for $+11.8\%$ ± 100 in cases of initialization below / above 2km above ground level ($M_{MW+,ini}$).

(3) To avoid misinterpretation of fast and random variations of specific humidity. Potentially artificial diurnal variations of

q along the back trajectories ~~as could result in an overestimation of~~ the formation of precipitation and moisture uptake, ~~we smoothed humidity~~. To avoid such an overestimation, diurnal variations of q were suppressed by smoothing q along the back trajectories with a 24 h broad rectangle kernel. ~~Arbitrarily assuming function. Arbitrarily choosing~~ a width of 24 h may ~~smooth out real sub-diurnal and diurnal variations of q and thereby, may~~ result in an ~~overestimation or~~ underestimation of the ~~formation of precipitation and amount of~~ moisture uptake. To assess the potential impact of the smoothing on the modeled $\delta D/\delta D$, we changed the width of the applied rectangle kernel to 12 h in $M_{S-,upt.12h}$ and to 36 h in $M_{MW+,upt.36h}$.

To finally assess the minimum possible values of $\Delta\delta D/\Delta\delta D$ in the case of superposition of the three sources of uncertainty discussed above, we combined the assumptions of $M_{S-,snow}$, $M_{S-,ini}$, and $M_{S-,upt.12h}$ in the model run M_{S-} and $M_{MW+,snow}$, $M_{MW+,ini}$, and $M_{MW+,upt.36h}$ in the model run M_{MW+} .

Table 1 summarizes $\Delta\delta D/\Delta\delta D$ for the different model runs. None of the model runs considering only one source of uncertainty is able to reduce $\Delta\delta D/\Delta\delta D$ for the scenarios of sublimation or evaporation of meltwater to values close to 0. Even for M_{S-} , which simultaneously assumes all the uncertainties of ~~side constraints model assumptions~~ in the scenario of sublimation, $\Delta\delta D/\Delta\delta D$ is ~~+13.5~~ 16.0%. The discussed uncertainty terms are therefore not able to bring model and observations into agreement with each other, if only considering ~~non-fractioning non-fractionating~~ sublimation. This implies that ~~fractioning fractionating~~ evaporation of meltwater played a significant role during our observations.

For M_{MW+} , which simultaneously assumes all the uncertainties of ~~side constraints model assumptions~~ in the scenario of evaporation of meltwater, the ~~value of $\Delta\delta D$ absolute value of $\Delta\delta D$~~ is reduced to ~~1.30~~ 0.9%. Considering the statistical uncertainty of $\Delta\delta D/\Delta\delta D$ of 1.6%, the average modeled and measured ~~δD of the 178~~ δD of the 174 selected air masses may therefore be brought into rough agreement with each other, if assuming ~~fractioning fractionating~~ evaporation of meltwater.

However, this requires superposition of the different uncertainty terms.

In order to refine this result, we split the selected observations into two groups of equal size according to the predominant skin temperature during moisture uptake ($T_{skin,predom.}$). For this purpose, we weighted skin temperatures along the individual ensembles of back trajectories with moisture uptake identified by the Lagrangian moisture source diagnostic. The median $T_{skin,predom.}$ of the ~~178-174~~ selected trajectory ensembles is -6.92°C . We attributed data points to a group "cold", if $T_{skin,predom.}$ of the respective trajectory ensemble is below -6.92°C . For $T_{skin,predom.}$ above -6.92°C we attributed data points to a group "warm". Please note that also points of group "warm" have a $T_{skin,predom.}$ below 0°C according to our selection criteria. Due to interannual variability of $T_{skin,predom.}$, data of the two groups is not randomly distributed in time. ~~Seventy-six~~ Seventy-seven percent of group "cold" correspond to an especially pronounced cold snap in February/March 2012, whereas 85% of group "warm" belong to cold snaps between October 2012 and February 2013.

Figure 9 shows two-dimensional probability distributions of the selected modeled and measured $\delta D/\delta D$. Blue denotes data from group "cold" and red denotes data from group "warm". Under the assumption of ~~non-fractioning non-fractionating~~ sublimation (M_S), modeled ~~δD ratios δD values~~ of both groups are significantly higher than the observed values (Fig. 9a). The overestimation of modeled ~~δD ratios δD values~~ is especially strong in the regime with higher T_{skin} , where snow melt and ~~fractioning fractionating~~ evaporation would be expected. Figure 9b shows the respective probability distributions under the assumption of snow melt and ~~fractioning fractionating~~ evaporation of meltwater (M_{MW}). Under this assumption, the modeled $\delta D/\delta D$ of

group "warm" is close to the observations. However, modeled δD -ratios- δD values of group "cold" are now far too low. Table 1 lists the mean differences between modeled and measured δD - δD for the two groups ($\Delta\delta D_{\text{cold}}$, $\Delta\delta D_{\text{warm}}$, $\Delta\delta D_{\text{cold}}$, $\Delta\delta D_{\text{warm}}$). As δD - δD of the individual groups may deviate more from the observations than the mean δD -of-all-178- δD of all 174 selected air masses, analyzing $\Delta\delta D_{\text{cold}}$ and $\Delta\delta D_{\text{warm}}$, $\Delta\delta D_{\text{cold}}$ and $\Delta\delta D_{\text{warm}}$ allows drawing less ambiguous conclusions than analysis of $\Delta\delta D$ - $\Delta\delta D$. For $M_{S,---$ the value of $\Delta\delta D_{\text{warm}}$ (16.7% $\Delta\delta D_{\text{warm}}$ (18.2%)) is larger than the value of $\Delta\delta D$ - $\Delta\delta D$ of all selected data (13.5% 16.0%) which underlines the importance of fractioning-fractionating evaporation for reproducing the observations. For $M_{MW,+++}$ the value of $\Delta\delta D_{\text{cold}}$ (11.9% absolute value of $\Delta\delta D_{\text{cold}}$ (10.5%)) is larger than the value of $\Delta\delta D$ (1.3% absolute value of $\Delta\delta D$ (0.9%)). So even $M_{MW,+++}$, in which we simultaneously assumed all the uncertainties of side constraints-model assumptions in the scenario of fractioning-fractionating evaporation of meltwater doesn't allow reproducing the observations of group "cold". This, in turn, implies significant non-fractioning-non-fractionating sublimation during our observations.

Comparison of δD -observations with δD - δD observations with δD of the Lagrangian isotope model therefore implies a relevant role of both types of isotope fractionation in Central Europe.

5.2 Temperature-dependent types of fractionation

To simultaneously bring into agreement modeled and observed δD -of-both-groups δD of data corresponding to group "cold" and group "warm", we suggest the existence of two regimes of T_{skin} with predominant non-fractioning-non-fractionating sublimation in the colder regime and predominant fractioning-fractionating evaporation of meltwater in the warmer regime. For the characterization of these two regimes we assume a maximum temperature for non-fractioning-non-fractionating sublimation in the model: $T_{\text{subl,max}}$. For $T_{\text{skin}} < T_{\text{subl,max}}$, we assume non-fractioning-non-fractionating sublimation. In the case of $T_{\text{subl,max}} < T_{\text{skin}} < 0^\circ\text{C}$, we assume equilibrium fractionation during the evaporation of meltwater. To assess $T_{\text{subl,max}}$ for optimal agreement between modeled and observed δD - δD , we performed 16 model runs with a different $T_{\text{subl,max}}$ in each run (-15 to 0°C in steps of 1 K). We refer to these runs as $M_{S_MW, T_{\text{subl,max}}}$. The thick black line in Fig. 10 shows the mean differences between modeled and observed δD -of-the-178- δD of the 174 selected data points ($\Delta\delta D$ - $\Delta\delta D$) from the 16 different $M_{S_MW, T_{\text{subl,max}}}$. Modeled δD -ratios- δD values are highest in $M_{S_MW, 0^\circ\text{C}}$, which assumes non-fractioning non-fractionating sublimation of snow for all $T_{\text{skin}} < 0^\circ\text{C}$ and is therefore identical with M_S . In-Out of the 16 model runs $M_{S_MW, -15^\circ\text{C}}$ the modeled δD -ratios are lowest- δD -ratios gives the lowest δD values. The δD values from $M_{S_MW, -15^\circ\text{C}}$ are close to δD - δD from M_{MW} , as most T_{skin} along the back trajectories were above -15°C , which means that almost no sublimation below $T_{\text{subl,max}}$ is considered. $M_{S_MW, -29^\circ\text{C}}$ would give-gives the same results as M_{MW} , because there was no moisture uptake identified for $T_{\text{skin}} \leq -29^\circ\text{C}$.

The agreement between observed and modeled δD - δD from the $M_{S_MW, T_{\text{subl,max}}}$ is best for a $T_{\text{subl,max}}$ of $-7.17.7^\circ\text{C}$. For this $T_{\text{subl,max}}$ the mean $\Delta\delta D$ - $\Delta\delta D$ of all selected data points is 0 and also δD - δD of group "cold" as well as the δD - δD of group "warm" is approximately reproduced by the model. Figure 9c shows the respective two-dimensional probability distributions. The statistical uncertainty of this optimal $T_{\text{subl,max}}$ due to scatter between modeled and observed δD - δD is 0.7°C . Further uncertainty is introduced to the determined optimal $T_{\text{subl,max}}$ by the assumptions of the Lagrangian isotope model, which can

systematically change the mean modeled δD and the optimal $T_{\text{subl,max}}$. To assess this uncertainty of the optimal $T_{\text{subl,max}}$, we varied side constraints (~~type_of_variation~~) in the $M_{S_MW,T_{\text{subl,max}}}$ analogous to the uncertainty assessment for the average modeled δD . For this purpose we performed 16·8=128 additional model runs (Fig. 10, thin lines) corresponding to 16 different $T_{\text{subl,max}}$ from -15 to 0°C and 8 different model configurations with the same assumptions about a changed δD of snow, a changed δD at the model initialization, a different amount of moisture uptake, and superposition of the three effects as in the above uncertainty assessment for the M_S and the M_{MW} . Analogous to the uncertainty assessment for the M_S and the M_{MW} , which we refer to as $M_{S_MW,\text{type_of_variation},T_{\text{subl,max}}}$. Model the model runs as $M_{S_MW,+,\text{snow},T_{\text{subl,max}}}$, $M_{S_MW,+,\text{ini},T_{\text{subl,max}}}$, $M_{S_MW,+,\text{upt.36h},T_{\text{subl,max}}}$, $M_{S_MW,+++,T_{\text{subl,max}}}$, $M_{S_MW,-,\text{snow},T_{\text{subl,max}}}$, $M_{S_MW,-,\text{ini},T_{\text{subl,max}}}$, $M_{S_MW,-,\text{upt.12h},T_{\text{subl,max}}}$ and $M_{S_MW,---,T_{\text{subl,max}}}$. The model runs with assumptions related to higher modeled δD -ratios- δD values (dashed thin lines) result in a lower optimal $T_{\text{subl,max}}$ and model runs with assumptions related to lower modeled δD -ratios- δD values (solid thin lines) result in a higher optimal $T_{\text{subl,max}}$.

Thin black lines in Fig. 10 depict the maximum possible shift of the average modeled δD - δD in Karlsruhe in the case of superposition of the ~~three examined side constraints~~ examined model assumptions. The solid thin black line reflects a maximum unfavorable superposition of assumptions related to lower modeled δD -ratios- δD values ($M_{S_MW,---,T_{\text{subl,max}}}$). The $M_{S_MW,---,T_{\text{subl,max}}}$ therefore allow to assess the upper bound of $-3.23.6^\circ\text{C}$ of the $T_{\text{subl,max}}$ for optimal agreement between model and observation. The lower bound of the optimal $T_{\text{subl,max}}$, which is derived from the $M_{S_MW,+++,T_{\text{subl,max}}}$, is -15°C (thin dashed line). Here again, individual analysis of data from the groups "cold" and "warm" allows us to refine the result. For $M_{MW,+++}$ $\Delta\delta D_{\text{warm}}/\Delta\delta D_{\text{warm}}$ is $+9.38.6\%$ (Table 1). Since $M_{MW,+++}$ assumes fractionating evaporation of meltwater for all $T_{\text{skin}} < 0^\circ\text{C}$, it marks the lower boundary of δD -ratios- δD values from the $M_{S_MW,+++,T_{\text{subl,max}}}$, ~~the~~. The corresponding set of assumptions ~~doesn't~~ does therefore not allow reproducing the observations in group "warm" even if assuming a very low $T_{\text{subl,max}}$. Assuming maximum unfavorable superposition of the uncertainties of ~~side constraints~~ model assumptions in the $M_{S_MW,+++,T_{\text{subl,max}}}$ is therefore too conservative for the uncertainty assessment of the optimal $T_{\text{subl,max}}$. For this reason, we assess the lower bound of the optimal $T_{\text{subl,max}}$ by means of the $M_{S_MW,+,\text{snow},T_{\text{subl,max}}}$, which only consider one uncertainty term and just allow the reproduction of the observed δD - δD in group "warm". From the $M_{S_MW,+,\text{snow},T_{\text{subl,max}}}$ we derive a better confined lower bound of the optimal $T_{\text{subl,max}}$ of $-10.811.2^\circ\text{C}$.

So the uncertainty of model assumptions translates into an uncertainty range of $T_{\text{subl,max}}$ for optimal agreement of model and observation from $-10.8-11.2$ to $-3.23.6^\circ\text{C}$. Together with the statistical uncertainty of 0.7°C , the total uncertainty range of $T_{\text{subl,max}}$ sums up to $-11.5-11.9$ to $-2.52.9^\circ\text{C}$.

6 Conclusions

In this paper, we investigated isotope fractionation during ~~the sublimation or evaporation of snow at ground level~~ surface evaporation in snow covered regions. For this purpose, we combined 17 months of measurements of δD - δD in low-level water vapor in Central Europe with a new Lagrangian isotope model.

By means of this approach, we identified two regimes of GDAS skin temperatures (T_{skin}) below the freezing point with signif-

icantly different deviation between modeled and observed $\delta D/\delta D$. To resolve this difference, we suggest two regimes of T_{skin} with different types of predominant isotope fractionation. Based on sensitivity tests with the Lagrangian isotope model, we found that the colder regime is described best by ~~non-fractioning~~ non-fractionating sublimation of snow. The warmer regime is described best by fractioning-fractionating evaporation of meltwater.

- 5 We determined a $T_{subl,max}$ separating both regimes of T_{skin} by optimizing the agreement between modeled and observed δD . ~~For δD . For a $T_{subl,max}$ of $-7.17.7^\circ C$~~ this agreement is best. Uncertainty related to assumptions of the isotope model corresponds to a range of uncertainty of $T_{subl,max}$ from ~~$-11.5-11.9$~~ to $-2.52.9^\circ C$.

The finding of a cold temperature regime with a small impact of fractionation during sublimation at ground level does not contradict earlier studies on snow which indicate fractioning-fractionating interaction between the ~~snowpack~~ surface layer snow and atmospheric water vapor, even in cases of temperatures far below the freezing point (Epstein et al., 1965; Stichler et al., 2001). These studies document systematic post-depositional increases of isotope ratios in the snowpack, which imply processes such as slight kinetic fractionation during sublimation in consequence of different coefficients of diffusion of the different water isotopes or fractioning-fractionating vapor deposition. Given the uncertainty of the Lagrangian isotope model, these small effects would not be detectable by our approach. Nevertheless, these effects might result in significant post-depositional modifications of isotope ratios in the snowpack on timescales longer than the five days covered by the trajectories.

For GDAS skin temperatures between $T_{subl,max}$ and $0^\circ C$ our results imply significant fractioning-fractionating evaporation of meltwater. ~~Despite skin temperatures below the freezing point, the formation of meltwater is likely to exist within this temperature regime. Since we weighted skin temperatures with positive latent heat flux at ground level, $T_{subl,max}$ already refers to skin temperatures during the day, when evaporation is strongest. However, due to the coarse resolution of the six-hourly 1×1 GDAS data, locations of locally or temporally enhanced skin temperatures are smoothed out. The evaporation of meltwater at these locations may exceed the amount of moisture from sublimation for skin temperatures above $T_{subl,max}$.~~ ~~The~~ identification of a fractioning-fractionating "meltwater regime" is consistent with earlier observations of isotope ratios in snow, which point to fractioning-fractionating evaporation during ablation at temperatures close below the freezing point (Moser and Stichler, 1974; Gurney and Lawrence, 2004; Lechler and Niemi, 2011). Since snow samples give an integrated signal over long time periods, a detailed attribution of these observations to certain meteorological conditions is difficult. Complementary to the studies on snow, the method presented here allows the post-depositional isotope fractionation to be attributed to meteorological conditions with GDAS skin temperatures between $T_{subl,max}$ and $0^\circ C$.

The determined $T_{subl,max}$ refers to a GDAS skin temperature that was weighted with positive latent heat flux at ground level. For this reason $T_{subl,max}$ is representative for GDAS skin temperatures during the day, when evaporation is strongest. However it should be kept in mind that due to the coarse resolution of $1\times 1^\circ$ of the GDAS data, much spatial variability of the skin temperature is smoothed out. The meaning of the $T_{subl,max}$ derived in this study is therefore an average temperature in a $1\times 1^\circ$ grid cell above which the evaporation of meltwater exceeds the amount of moisture from sublimation. A way to derive a more physical temperature separating the regimes of sublimation and meltwater evaporation could be using data with a higher horizontal resolution. This wouldn't necessarily improve accuracy with respect to the back trajectories' positions but locations with enhanced skin temperatures and especially high amounts of surface evaporation would be more realistically represented,

presumably resulting in higher evaporation weighted skin temperatures and a higher $T_{\text{subl,max}}$.

In addition, higher horizontal resolution would allow to better account for spatial heterogeneity of the δD of surface layer snow in mountainous regions, for instance by also weighting the δD of the snow with the amount of surface evaporation.

In this context please note that systematic uncertainty regarding the δD of surface layer snow turned out to be the main

5 limitation for determining $T_{\text{subl,max}}$ with the approach presented here. For this reason regular analysis of the δD of surface layer snow samples for instance at selected GNIP stations would be a very desirable and efficient measure to reduce uncertainty of $T_{\text{subl,max}}$.

Our results show that surface evaporation in the two identified regimes of skin temperature has a strong impact on the δD - δD of low-level water vapor in Central Europe. For isotope applications based on relations between δD - δD and temperature, the

10 consideration of the different types of isotope fractionation in both regimes is therefore of great interest. For instance, seasonal ablation in coastal regions of Greenland might systematically affect the relation between the δD - δD of water vapor and dew point temperature over Greenland. Because such a seasonality may be different in climatologically different time periods, it may introduce uncertainty to temperature reconstructions from Greenland ice cores.

Furthermore, ~~fractioning-fractionating~~ evaporation of meltwater will increase the δD -ratio- δD value of the residual meltwater

15 and in the case of recrystallization, the δD -ratio- δD value of the snowpack. Ignoring ~~fractioning-fractionating~~ evaporation therefore introduces uncertainty to a variety of isotope applications from reconstructions of paleotemperatures and paleogeography to studies on the formation of groundwater. The specification of a temperature regime with enhanced fractionation during evaporation may therefore help to identify, investigate, and reduce biases inherent to these applications.

Acknowledgements. This study was funded in part by the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no. 256961 and by the German Climate Modeling Initiative (PalMod). We acknowledge support by Deutsche Forschungsgemeinschaft and Open Access Publishing Fund of Karlsruhe Institute of Technology.

References

- Aemisegger, F., Sturm, P., Graf, P., Sodemann, H., Pfahl, S., Knohl, A., and Wernli, H.: Measuring variations of $\delta^{18}\text{O}$ and $\delta^2\text{H}$ in atmospheric water vapour using two commercial laser-based spectrometers: an instrument characterisation study, *Atmospheric Measurement Techniques*, 5, 1491–1511, doi:10.5194/amt-5-1491-2012, 2012.
- 5 Aemisegger, F., Pfahl, S., Sodemann, H., Lehner, I., Seneviratne, S. I., and Wernli, H.: Deuterium excess as a proxy for continental moisture recycling and plant transpiration, *Atmospheric Chemistry and Physics*, 14, 4029–4054, doi:10.5194/acp-14-4029-2014, 2014.
- Aemisegger, F., Spiegel, J. K., Pfahl, S., Sodemann, H., Eugster, W., and Wernli, H.: Isotope meteorology of cold front passages: A case study combining observations and modeling, *Geophysical Research Letters*, 42, 5652–5660, doi:10.1002/2015GL063988, 2015.
- Aharon, P. and Chappell, J.: Oxygen isotopes, sea level changes and the temperature history of a coral reef environment in New Guinea over the last 105 years, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 56, 337–379, doi:10.1016/0031-0182(86)90101-X, 1986.
- 10 Ambach, W., Dansgaard, W., Eisner, H., and Moller, J.: The altitude effect on the isotopic composition of precipitation and glacier ice in the Alps, *Tellus*, 20, 595–600, doi:10.1111/j.2153-3490.1968.tb00402.x, 1968.
- Araguas, L. A., Danesi, P., Froehlich, K., and Rozanski, K.: Global monitoring of the isotopic composition of precipitation, *Journal of Radioanalytical and Nuclear Chemistry Articles*, 205, 189–200, doi:10.1007/BF02039404, 1996.
- 15 Barnes, C. and Allison, G.: The distribution of deuterium and ^{18}O in dry soils, *Journal of Hydrology*, 60, 141–156, doi:10.1016/0022-1694(83)90018-5, 1983.
- Blisniuk, P. M.: Stable isotope paleoaltimetry: A critical review, *American Journal of Science*, 305, 1033–1074, doi:10.2475/ajs.305.10.1033, 2005.
- Choudhury, B. J., DiGirolamo, N. E., Susskind, J., Darnell, W. L., Gupta, S. K., and Asrar, G.: A biophysical process-based estimate of global land surface evaporation using satellite and ancillary data II. Regional and global patterns of seasonal and annual variations, *Journal of Hydrology*, 205, 186–204, doi:10.1016/S0022-1694(97)00149-2, 1998.
- 20 Dahlke, H. E. and Lyon, S. W.: Early melt season snowpack isotopic evolution in the Tarfala valley, northern Sweden, *Annals of Glaciology*, 54, 149–156, doi:10.3189/2013AoG62A232, 2013.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436–468, doi:10.1111/j.2153-3490.1964.tb00181.x, 1964.
- 25 Dansgaard, W.: Stable isotope glaciology, Reitzel, 1973.
- de Vries, J. J. and Simmers, I.: Groundwater recharge: an overview of processes and challenges, *Hydrogeology Journal*, 10, 5–17, doi:10.1007/s10040-001-0171-7, 2002.
- Delaygue, G., Bard, E., Rollion, C., Jouzel, J., Stiévenard, M., Duplessy, J.-C., and Ganssen, G.: Oxygen isotope/salinity relationship in the northern Indian Ocean, *Journal of Geophysical Research*, 106, 4565, doi:10.1029/1999JC000061, 2001.
- 30 Derber, J. C., Parrish, D. F., and Lord, S. J.: The New Global Operational Analysis System at the National Meteorological Center, *Weather and Forecasting*, 6, 538–547, doi:10.1175/1520-0434(1991)006<0538:TNGOAS>2.0.CO;2, 1991.
- Draxler, R. R. and Hess, G. D.: An Overview of the HYSPLIT_4 Modelling System for Trajectories, Dispersion, and Deposition, *Australian Meteorological Magazine*, 47, 295–308, 1998.
- Duplessy, J.-C.: Note préliminaire sur les variations de la composition isotopique des eaux superficielles de l’Océan Indien: La relation ^{18}O -salinité, *CR Acad. Sci. Paris*, 271, 1075—1078, 1970.
- 35 Epstein, S., Sharp, R. P., and Gow, A. J.: Six-year record of oxygen and hydrogen isotope variations in South Pole firn, *Journal of Geophysical Research*, 70, 1809–1814, doi:10.1029/JZ070i008p01809, 1965.

- Farquhar, G. D., Cernusak, L. a., and Barnes, B.: Heavy water fractionation during transpiration., *Plant physiology*, 143, 11–8, doi:10.1104/pp.106.093278, 2007.
- Friedman, I., Machta, L., and Soller, R.: Water-vapor exchange between a water droplet and its environment, *Journal of Geophysical Research*, 67, 2761–2766, doi:10.1029/JZ067i007p02761, 1962.
- 5 Friedman, I., Benson, C., and Gleason, J.: Isotopic changes during snow metamorphism, *Stable Isotope Geochemistry: A Tribute to Samuel Epstein*, pp. 211–221, 1991.
- Fröhlich, K., Grabczak, J., and Rozanski, K.: Deuterium and oxygen-18 in the baltic sea, *Chemical Geology: Isotope Geoscience section*, 72, 77–83, doi:10.1016/0168-9622(88)90038-3, 1988.
- Gat, J. R., Shemesh, A., Tziperman, E., Hecht, A., Georgopoulos, D., and Basturk, O.: The stable isotope composition of waters of the eastern
10 Mediterranean Sea, *Journal of Geophysical Research*, 101, 6441, doi:10.1029/95JC02829, 1996.
- Gedzelman, S. D. and Arnold, R.: Modeling the isotopic composition of precipitation, *Journal of Geophysical Research*, 99, 10455, doi:10.1029/93JD03518, 1994.
- Gurney, S. and Lawrence, D.: Seasonal trends in the stable isotopic composition of snow and meltwater runoff in a subarctic catchment at Okstindan, Norway., *Nordic hydrology*, 35, 119–137, 2004.
- 15 Hanisco, T. F., Moyer, E. J., Weinstock, E. M., St. Clair, J. M., Sayres, D. S., Smith, J. B., Lockwood, R., Anderson, J. G., Dessler, a. E., Keutsch, F. N., Spackman, J. R., Read, W. G., and Bui, T. P.: Observations of deep convective influence on stratospheric water vapor and its isotopic composition, *Geophysical Research Letters*, 34, L04814, doi:10.1029/2006GL027899, 2007.
- Harwood, K. G., Gillon, J. S., Roberts, A., and Griffiths, H.: Determinants of isotopic coupling of CO₂ and water vapour within a Quercus petraea forest canopy, *Oecologia*, 119, 109–119, doi:10.1007/s004420050766, 1999.
- 20 Helsen, M. M., van de Wal, R. S. W., van den Broeke, M. R., Kerstel, E. R. T., Masson-Delmotte, V., Meijer, H. a. J., Reijmer, C. H., and Scheele, M. P.: Modelling the isotopic composition of snow using backward trajectories: a particular precipitation event in Dronning Maud Land, Antarctica, *Annals of Glaciology*, 39, 293–299, doi:10.3189/172756404781814230, 2004.
- Helsen, M. M., Van De Wal, R. S. W., Van Den Broeke, M. R., Van As, D., Meijer, H. A. J., and Reijmer, C. H.: Oxygen isotope variability in snow from western Dronning Maud Land, Antarctica and its relation to temperature, *Tellus B*, 57, 423–435,
25 doi:10.1111/j.1600-0889.2005.00162.x, 2005.
- Helsen, M. M., Van de Wal, R. S. W., and Van den Broeke, M. R.: The Isotopic Composition of Present-Day Antarctic Snow in a Lagrangian Atmospheric Simulation*, *Journal of Climate*, 20, 739–756, doi:10.1175/JCLI4027.1, 2007.
- Horita, J. and Wesolowski, D. J.: Liquid-vapor fractionation of oxygen and hydrogen isotopes of water from the freezing to the critical temperature, *Geochimica et Cosmochimica Acta*, 58, 3425–3437, doi:10.1016/0016-7037(94)90096-5, 1994.
- 30 IAEA: Reference Sheet for VSMOW2 and SLAP2 international measurement standards. Issued 2009-02-13, International Atomic Energy Agency, 2009.
- Jacob, H. and Sonntag, C.: An 8-year record of the seasonal variation of 2H and 18O in atmospheric water vapour and precipitation at Heidelberg, Germany, *Tellus B*, 43, 291–300, doi:10.1034/j.1600-0889.1991.t01-2-00003.x, 1991.
- James, P., Stohl, A., Spichtinger, N., Eckhardt, S., and Forster, C.: Climatological aspects of the extreme European rainfall of August 2002
35 and a trajectory method for estimating the associated evaporative source regions, *Natural Hazards and Earth System Science*, 4, 733–746, doi:10.5194/nhess-4-733-2004, 2004.
- Jancso, G. and Van Hook, W. A.: Condensed phase isotope effects, *Chemical Reviews*, 74, 689–750, doi:10.1021/cr60292a004, 1974.

- Johnsen, S. J., Dansgaard, W., and White, J. W. C.: The origin of Arctic precipitation under present and glacial conditions, *Tellus B*, 41B, 452–468, doi:10.1111/j.1600-0889.1989.tb00321.x, 1989.
- Jouzel, J. and Merlivat, L.: Deuterium and oxygen 18 in precipitation: Modeling of the isotopic effects during snow formation, *Journal of Geophysical Research*, 89, 11 749, doi:10.1029/JD089iD07p11749, 1984.
- 5 Kanamitsu, M.: Description of the NMC Global Data Assimilation and Forecast System, *Weather and Forecasting*, 4, 335–342, doi:10.1175/1520-0434(1989)004<0335:DOTNGD>2.0.CO;2, 1989.
- Läderach, A. and Sodemann, H.: A revised picture of the atmospheric moisture residence time, *Geophysical Research Letters*, 43, 924–933, doi:10.1002/2015GL067449, 2016.
- Lawrence, D. M., Thornton, P. E., Oleson, K. W., and Bonan, G. B.: The Partitioning of Evapotranspiration into Transpiration, Soil Evaporation, and Canopy Evaporation in a GCM: Impacts on Land-Atmosphere Interaction, *Journal of Hydrometeorology*, 8, 862–880, doi:10.1175/JHM596.1, 2007.
- 10 Lechler, A. R. and Niemi, N. A.: The influence of snow sublimation on the isotopic composition of spring and surface waters in the southwestern United States: Implications for stable isotope-based paleoaltimetry and hydrologic studies, *Geological Society of America Bulletin*, 124, 318–334, doi:10.1130/B30467.1, 2011.
- 15 LeGrande, A. N. and Schmidt, G. a.: Global gridded data set of the oxygen isotopic composition in seawater, *Geophysical Research Letters*, 33, L12 604, doi:10.1029/2006GL026011, 2006.
- Masson-Delmotte, V., Jouzel, J., Landais, A., Stievenard, M., Johnsen, S. J., White, J. W. C., Werner, M., Sveinbjornsdottir, A., and Fuhrer, K.: GRIP deuterium excess reveals rapid and orbital-scale changes in Greenland moisture origin., *Science (New York, N.Y.)*, 309, 118–21, doi:10.1126/science.1108575, 2005.
- 20 Mathieu, R. and Bariac, T.: A numerical model for the simulation of stable isotope profiles in drying soils, *Journal of Geophysical Research*, 101, 12 685, doi:10.1029/96JD00223, 1996.
- Merlivat, L. and Jouzel, J.: Global climatic interpretation of the deuterium-oxygen 18 relationship for precipitation, *Journal of Geophysical Research*, 84, 5029, doi:10.1029/JC084iC08p05029, 1979.
- Moser, H. and Stichler, W.: Deuterium and oxygen-18 contents as an index of the properties of snow covers, *International Association of Hydrological Sciences Publication*, 114, 122–135, 1974.
- 25 Noone, D., Risi, C., Bailey, A., Berkelhammer, M., Brown, D. P., Buening, N., Gregory, S., Nusbaumer, J., Schneider, D., Sykes, J., Vanderwende, B., Wong, J., Meillier, Y., and Wolfe, D.: Determining water sources in the boundary layer from tall tower profiles of water vapor and surface water isotope ratios after a snowstorm in Colorado, *Atmospheric Chemistry and Physics*, 13, 1607–1623, doi:10.5194/acp-13-1607-2013, 2013.
- 30 Ostlund, H. G., Craig, H., Broecker, W. S., and Spenser, D.: Shorebased data and graphics, *GEOSECS Atlantic, Pacific and Indian Ocean Expeditions*, 7, 1987.
- Parrish, D. F. and Derber, J. C.: The National Meteorological Center's Spectral Statistical-Interpolation Analysis System, *Monthly Weather Review*, 120, 1747–1763, doi:10.1175/1520-0493(1992)120<1747:TSMCSS>2.0.CO;2, 1992.
- Pfahl, S. and Wernli, H.: Lagrangian simulations of stable isotopes in water vapor: An evaluation of nonequilibrium fractionation in the Craig-Gordon model, *Journal of Geophysical Research*, 114, D20 108, doi:10.1029/2009JD012054, 2009.
- 35 Poage, M. A.: Empirical relationships between elevation and the stable isotope composition of precipitation and surface waters: considerations for studies of paleoelevation change, *American Journal of Science*, 301, 1–15, doi:10.2475/ajs.301.1.1, 2001.

- Rayleigh, L. and Ramsay, W.: Argon, a New Constituent of the Atmosphere., *Proceedings of the Royal Society of London*, 57, 265–287, doi:10.1098/rspl.1894.0149, 1894.
- Risi, C., Bony, S., Vimeux, F., and Jouzel, J.: Water-stable isotopes in the LMDZ4 general circulation model: Model evaluation for present-day and past climates and applications to climatic interpretations of tropical isotopic records, *Journal of Geophysical Research*, 115, D12 118, doi:10.1029/2009JD013255, 2010.
- Risi, C., Ogée, J., Bony, S., Bariac, T., Raz-yaseef, N., Wingate, L., Welker, J., Knohl, A., Kurz-Besson, C., Leclerc, M., Zhang, G., Buchmann, N., Santrucek, J., Hronkova, M., David, T., Peylin, P., and Guglielmo, F.: The Water Isotopic Version of the Land-Surface Model ORCHIDEE: Implementation, Evaluation, Sensitivity to Hydrological Parameters, *Hydrology: Current Research*, 7, 2157–7587, doi:10.4172/2157-7587.1000258, 2016.
- 10 Rowley, D. B. and Garzione, C. N.: Stable Isotope-Based Palealtimetry, *Annual Review of Earth and Planetary Sciences*, 35, 463–508, doi:10.1146/annurev.earth.35.031306.140155, 2007.
- Rowley, D. B., Pierrehumbert, R. T., and Currie, B. S.: A new approach to stable isotope-based palealtimetry: implications for paleoaltimetry and paleohypsometry of the High Himalaya since the Late Miocene, *Earth and Planetary Science Letters*, 188, 253–268, doi:10.1016/S0012-821X(01)00324-7, 2001.
- 15 Sayres, D. S., Pfister, L., Hanisco, T. F., Moyer, E. J., Smith, J. B., St. Clair, J. M., O'Brien, a. S., Witinski, M. F., Legg, M., and Anderson, J. G.: Influence of convection on the water isotopic composition of the tropical tropopause layer and tropical stratosphere, *Journal of Geophysical Research*, 115, D00J20, doi:10.1029/2009JD013100, 2010.
- Schlosser, E., Reijmer, C., Oerter, H., and Graf, W.: The influence of precipitation origin on the d18O-T relationship at Neumayer station, Ekströmisen, Antarctica, *Annals of Glaciology*, 39(1), 41–48, 2004.
- 20 Schmidt, G., Bigg, G. R., and Rohling, E. J.: "Global Seawater Oxygen-18 Database - v1.21, 1999.
- Schoch-Fischer, H., Rozanski, K., Jacob, H., Sonntag, C., Jouzel, I., Östlund, G., and Geyh, M. A.: Hydrometeorological factors controlling the time variation of D, 18O and 3H in atmospheric water vapour and precipitation in the northern westwind belt. IAEA,(ed.), in: *Isotope Hydrology*, pp. 3–30, IAEA, Vienna, 1983.
- Sodemann, H., Masson-Delmotte, V., Schwierz, C., Vinther, B. M., and Wernli, H.: Interannual variability of Greenland winter precipitation sources: 2. Effects of North Atlantic Oscillation variability on stable isotopes in precipitation, *Journal of Geophysical Research*, 113, D12 111, doi:10.1029/2007JD009416, 2008a.
- Sodemann, H., Schwierz, C., and Wernli, H.: Interannual variability of Greenland winter precipitation sources: Lagrangian moisture diagnostic and North Atlantic Oscillation influence, *Journal of Geophysical Research*, 113, D03 107, doi:10.1029/2007JD008503, 2008b.
- Stewart, M. K.: Stable isotope fractionation due to evaporation and isotopic exchange of falling waterdrops: Applications to atmospheric processes and evaporation of lakes, *Journal of Geophysical Research*, 80, 1133–1146, doi:10.1029/JC080i009p01133, 1975.
- 30 Stichler, W., Schotterer, U., Fröhlich, K., Ginot, P., Kull, C., Gäggeler, H., and Pouyaud, B.: Influence of sublimation on stable isotope records recovered from high-altitude glaciers in the tropical Andes, *Journal of Geophysical Research*, 106, 22 613, doi:10.1029/2001JD900179, 2001.
- Stohl, A. and James, P.: A Lagrangian Analysis of the Atmospheric Branch of the Global Water Cycle. Part I: Method Description, Validation, and Demonstration for the August 2002 Flooding in Central Europe, *Journal of Hydrometeorology*, 5, 656–678, doi:10.1175/1525-7541(2004)005<0656:ALAOTA>2.0.CO;2, 2004.

- Terzer, S., Wassenaar, L. I., Araguás-Araguás, L. J., and Aggarwal, P. K.: Global isoscapes for $\delta^{18}\text{O}$ and $\delta^2\text{H}$ in precipitation: improved prediction using regionalized climatic regression models, *Hydrology and Earth System Sciences*, 17, 4713–4728, doi:10.5194/hess-17-4713-2013, 2013.
- Trenberth, K. E.: Atmospheric Moisture Residence Times and Cycling: Implications for Rainfall Rates and Climate Change, *Climatic Change*, 39, 667–694, doi:10.1023/A:1005319109110, 1998.
- Weiss, R., Östlund, H., and Craig, H.: Geochemical studies of the Weddell sea, *Deep Sea Research Part A. Oceanographic Research Papers*, 26, 1093–1120, doi:10.1016/0198-0149(79)90059-1, 1979.
- Wen, X.-F., Sun, X.-M., Zhang, S.-C., Yu, G.-R., Sargent, S. D., and Lee, X.: Continuous measurement of water vapor D/H and $^{18}\text{O}/^{16}\text{O}$ isotope ratios in the atmosphere, *Journal of Hydrology*, 349, 489–500, doi:10.1016/j.jhydrol.2007.11.021, 2008.
- 10 Werner, M., Haese, B., Xu, X., Zhang, X., Butzin, M., and Lohmann, G.: Glacial–interglacial changes of H_2^{18}O , HDO and deuterium excess – results from the fully coupled Earth System Model ECHAM5/MPI-OM, *Geoscientific Model Development Discussions*, 8, 8835–8894, doi:10.5194/gmdd-8-8835-2015, 2016.
- Yobbi, D. K.: Effects of Tidal Stage and Ground-Water Levels on the Discharge and Water Quality of Springs in Coastal Citrus and Hernando Counties, Florida, *Water Resources Investigations Report 92-4096*, USGS, 1992.
- 15 Yoshimura, K., Miyazaki, S., Kanae, S., and Oki, T.: Iso-MATSIRO, a land surface model that incorporates stable water isotopes, *Global and Planetary Change*, 51, 90–107, doi:10.1016/j.gloplacha.2005.12.007, 2006.
- Zhang, S., Wen, X., Wang, J., Yu, G., and Sun, X.: The use of stable isotopes to partition evapotranspiration fluxes into evaporation and transpiration, *Acta Ecologica Sinica*, 30, 201–209, doi:10.1016/j.chnaes.2010.06.003, 2010.

1)

Table 1. Average differences between modeled and measured δD of the 178-174 selected data points ($\Delta\delta D$), data points of group "cold" ($\Delta\delta D_{\text{cold}}$), and data points of group "warm" ($\Delta\delta D_{\text{warm}}$) from different model runs (M). \pm states the statistical uncertainty of the averages (root mean-square error divided by the square root of the number of observations). Values of particular interest are printed in bold type.

Name of model run	Description of model run
M_{MW}	<u>fractioning-fractionating</u> evaporation of meltwater; (reference run)
$M_{\text{MW+},\text{snow}}$	<u>fractioning-fractionating</u> evaporation of meltwater; δD ratios of snow increased by 11.5% <u>snow increased by 11.5%</u>
$M_{\text{MW+},\text{ini}}$	<u>fractioning-fractionating</u> evaporation of meltwater; δD ratios δD values at initialization increased by 11.8%/100for air masses originating from altitudes-
lower / higher than 2000m above ground level $M_{\text{MW+},\text{upt.36h}}$	<u>fractioning-fractionating</u> evaporation of meltwater; reduced moisture uptake in consequence of smoothing q along the trajectories with a 36 h broad rectangle kernel (instead of 24 h)
$M_{\text{MW+++}}$	<u>fractioning-fractionating</u> evaporation of meltwater; simultaneous occurrence of the of the three assumptions above
M_{S}	<u>non-fractioning-non-fractionating</u> sublimation of snow
$M_{\text{S-},\text{snow}}$	<u>non-fractioning-non-fractionating</u> sublimation of snow; δD ratios of snow decreased by 11.5% <u>snow decreased by 11.5%</u>
$M_{\text{S-},\text{ini}}$	<u>non-fractioning-non-fractionating</u> sublimation of snow; δD ratios δD values at initialization decreased by 3.6%/100for air masses originating from altitudes-
lower / higher than 2000m above ground level $M_{\text{S-},\text{upt.12h}}$	<u>non-fractioning-non-fractionating</u> sublimation of snow; increased moisture uptake in consequence of smoothing q along the trajectories with a 12 h broad rectangle kernel (instead of 24 h)
$M_{\text{S---}}$	<u>non-fractioning-non-fractionating</u> sublimation of snow; simultaneous occurrence of the of the three assumptions above




Figure 1. (a) δD of ocean surface water derived from the interpolation (V1.1) of the Global Seawater Oxygen-18 Database by LeGrande and Schmidt (2006), assuming a constant factor of 6.56 between δD and $\delta^{18}O$ in ocean surface water. (b) Climatological δD in precipitation in winter (DJF) from the Regionalized Cluster-based Water Isotope Prediction (RCWIP), which in turn is based on observations of the Global Network of Isotopes in Precipitation (GNIP). (c) same as in (b) but for summer (JJA).




Figure 2. (a) Probability distributions of continental GDAS skin temperatures ($T_{\text{skin,unweighted}}$) (blue) and of skin temperatures weighted with the accumulated hourly latent heat flux at ground level within ± 12 h (T_{skin}) (green). The occurrence of low temperatures is reduced as a consequence of the weighting. A peak around 0°C becomes more clearly visible. (b) Illustration of the weighting algorithm for one exemplary back trajectory (arrival in Karlsruhe May 4, 2012, 21 UTC).




Figure 3. Illustration of the isotope modeling for one exemplary back trajectory (arrival in Karlsruhe March 18, 2012, 0 UTC). **(a)** Altitude of the back trajectory (black) and terrain height (gray/blue). The isotope model was initialized at 80° N in the marine boundary layer (MBL). In a low-pressure system near Iceland the tracked air parcel ascended to an altitude of 3200 m. During the last three days of transport to Karlsruhe the air parcel was sinking to the sampling altitude. **(b)** After initialization in the MBL, δD of the tracked air parcel specific humidity q (light blue and green colored line) and δD values (thick black line) was of the tracked air parcel were slightly decreasing, due to the formation of precipitation (dashed light blue lines) within the first day. More pronounced formation of precipitation, in consequence of lofting in a low-pressure system near Iceland, resulted in a second pronounced decrease of specific humidity q (gray) and the modeled δD -ratio- δD value dropped accordingly. Due to moisture uptake (dashed green lines), related to a descent of the air parcel in the subsequent days, q and the δD -ratio- δD value were increasing, until the air parcel reached Karlsruhe. Black thin curves illustrate the modeled δD - δD for different initializations of δD - δD (section Sect. 2.3.3). The dependence on the initialization decreases with the amount of moisture uptake along the trajectories and is only low in Karlsruhe.




Figure 4. (a) Deviation of individual calibration measurements with the Picarro water isotopologue analyzer from isotope ratios of the liquid standards. Each point represents a calibration of one hour. Dots: Standard 1; crosses: Standard 2. (b) Humidity dependence of the δD measurements. Long-term drift depicted in (a) was removed in (b) by subtracting the five-week running average of calibrations. Red regression lines were calculated for both standards simultaneously by subtracting the mean difference between both standards. \pm gives the difference between the slopes calculated for the first and second half of the measurement period.

(a) δD of ocean surface water derived from the interpolation (V1.1) of the Global Seawater Oxygen-18 Database by , assuming a constant factor of 6.56 between δD and $\delta^{18}O$ in ocean surface water. (b) Climatological δD in precipitation in winter (DJF) from the Regionalized Cluster-based Water Isotope Prediction (RCWIP), which in turn is based on observations of the Global Network of Isotopes in Precipitation (GNIP). (c) same as in (b) but for summer (JJA):

(a) Probability distributions of continental GDAS skin temperatures ($T_{\text{skin,unweighted}}$) (blue) and of skin temperatures weighted with the accumulated hourly latent heat flux at ground level within 12h (T_{skin}) (green). The occurrence of low temperatures is reduced as a consequence of the weighting. A peak around 0 becomes more clearly visible. (b) Illustration of the weighting algorithm for one exemplary back trajectory (arrival in Karlsruhe May 4, 2012, 21UTC):




Figure 5. Source regions of moisture (q) 30 m above ground level in Karlsruhe (black star) ~~in~~ based on five-day back trajectories for the different seasons: ~~December, time period~~ January, February (DJF), March, April, 2012 to May (MAM), June, July, August (JJA), September, October, November (SON), 2013. The color code indicates the contribution of different source regions to q in Karlsruhe in % per km². Integration over the whole map gives the identified humidity per season. Because of the finite length of the back trajectories, the total identified humidity is lower than 100 of 47% and accounts for 52 (DJF), 51 (MAM), 39 (JJA) and 50 (SON).




Figure 6. Measurements in Karlsruhe from January 2012 ~~until~~to May 2013 30 m above ground level (ten-minute averages). Black: five-day back trajectories originate from the West; gray: five-day back trajectories originate from the East. **(a)** δD - δD of water vapor. Gaps in the time series are caused by instrumental issues with analyzer and calibration device. **(b)** Air temperature (T). **(c)** Specific humidity (q).




Figure 7. (a) Time series of measured (blue) and modeled (red) $\delta D - \delta D$ of water vapor in Karlsruhe. (b) Enlarged section of (a), which demonstrates the capability of the model of capturing the high variability of $\delta D - \delta D$ in winter. (c) Measured versus modeled $\delta D - \delta D$. The three-hourly available modeled $\delta D - \delta D$ is compared to the temporally closest ten-minute average of the $\delta D - \delta D$ observations. Gray: reference run, surface evaporation (E) and the formation of precipitation (P) are considered ($R=0.82$, $\Delta\delta D \Delta\delta D = +1.0 - 4.1\%$); green: only E is considered, P is ignored ($R=0.81$, $\Delta\delta D \Delta\delta D = +29.423.9\%$); magenta: only P is considered, E is ignored ($R=0.62$, $\Delta\delta D \Delta\delta D = -35.634.8\%$); black: 1:1 line.




Figure 8. (a) Lagrangian source region analysis of low-level water vapor in Karlsruhe for the observed cold snaps. The mean identified fraction of moisture along the five-day back trajectories is 48%. (b) Mean snow depth during the cold snaps based on GDAS data.




Figure 9. Two-dimensional probability distributions of measured and modeled δD of low-level water vapor in Karlsruhe for selected cold snap events. Red: group "warm"; blue: group "cold". Probabilities were calculated for a $20\% \times 20\%$ δD grid, smoothed with a 20% broad rectangle kernel, and finally interpolated to a $1\% \times 1\%$ grid. Probabilities are normalized to 1 at the maximum, contours show probability levels of **0.8**, 0.7, 0.6, 0.45, 0.35. **(a)** The model assumes sublimation of snow (no isotope fractionation) in the case of moisture uptake and skin temperature (T_{skin}) below 0°C . **(b)** The model assumes evaporation of melted snow (equilibrium isotope fractionation) in the case of moisture uptake and $T_{\text{skin}} < 0^\circ\text{C}$. **(c)** The model assumes sublimation of snow in the case of moisture uptake and $T_{\text{skin}} < -7.17^\circ\text{C}$. In the case of moisture uptake and $-7.17^\circ\text{C} < T_{\text{skin}} < 0^\circ\text{C}$ the model assumes evaporation of melted snow.




Figure 10. Mean difference between modeled and observed δD of water vapor at Karlsruhe for selected cold snap events ($\Delta\delta D$). Each cross represents the $\Delta\delta D$ obtained from one model configuration with specific model assumptions regarding the $T_{\text{subl,max}}$, the δD of snow, the δD at the model initialization, and the amount of moisture uptake. The thick black line connects the $\Delta\delta D$ from 16 model configurations with $T_{\text{subl,max}}$ between -15 and 0°C and the model assumptions of $M_{S,MW,T_{\text{subl,max}}}$ and illustrates the increase of the modeled δD values with increasing maximum skin temperature allowing non-fractionating sublimation ($T_{\text{subl,max}}$). Magenta: optimal $T_{\text{subl,max}}$ for best agreement. The gray shaded area depicts possible systematic changes of model and observation (dot) and uncertainty of the optimal $T_{\text{subl,max}}$ (error bar) due to potential $\Delta\delta D$ in the case of a systematic deviation of δD the δD of snow from the RCWIP climatology (light blue), a systematically changed δD at the model initialization of δD (yellow), and uncertainty of the a systematically changed amount of moisture uptake (green). Thin, and superposition of the different assumptions (thin black lines: modeled δD). The thin dashed black line corresponds to the model configuration $M_{S,MW,+,+,T_{\text{subl,max}}}$, which doesn't allow reproducing the observations in group "warm" and is therefore not considered in the case-gray shaded area of superposition uncertainty. Magenta: optimal $T_{\text{subl,max}}$ for best agreement of the different sources model and observation (dot) and uncertainty of the optimal $T_{\text{subl,max}}$ (error bar) due to uncertainty of model assumptions.