Letter to the Editor

ACP Discussions doi: 10.5194/acp-2016-463 (Editor - Peter Haynes) 'Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012'

Dear Peter Haynes,

many thanks for handling our manuscript. We prepared and submitted a revised version of our manuscript and are confident that we have satisfactorily addressed all comments of referee s#1 to #3. A detailed point-by-point response to all referee comments (#1 to #3) is attached. Further, a document specifying all changes in the revised manuscript compared to the ACPD version is added.

Best wishes

Bärbel Vogel

Author Comment to Referee #1

ACP Discussions doi: 10.5194/acp-2016-463 (Editor - Peter Haynes) 'Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower strato-

sphere during the Asian monsoon season 2012'

We thank Referee #1 for further guidance on how to to revise our paper. Following the reviewers advice we have elaborated some minor points, which strengthen our findings. Our reply to the reviewer comments is listed in detail below. Questions and comments of the referee are shown in italics.

This paper presents an analysis of the evolution and transport of air with emission sources in the Asian monsoon region to the northern hemisphere extratropical lower stratosphere using primarily output from the Chemical Lagrangian Model of the Stratosphere (CLaMS) driven with winds from the ERA-Interim reanalysis. Aircraft observations are used to establish confidence in the model simulations and demonstrate linkages between transport signatures and air masses observed in the extratopical lower stratosphere downstream. The main conclusions of the study are that the air lofted into the Asian monsoon anticyclone is confined to the tropical upper troposphere during the monsoon season, but is transported poleward into the extratropical lower stratosphere during the breakup of the Asian monsoon anticyclone in early fall via mostly Rossby wave breaking events over the Pacific and Atlantic ocean basins. The paper is well-detailed, well-structured, and well-reasoned. I do not find any fundamental errors in the analysis or questionable claims in the attendant discussion, but I believe the paper will benefit from a bit more cohesion of arguments throughout and a bit more analysis of the seasonality of transport. I recommend that the paper be considered for publication after mostly minor revisions. My general and specific comments to help quide the revisions are provided below.

General Comments

1. The linkages between transport of the monsoon upper troposphere air to the extratropical lower stratosphere and poleward Rossby wave breaking downstream of the anticyclone are clearly established in the manuscript. However, I feel the authors miss an opportunity to strengthen these points as their analysis expands in the latter part of the paper. Namely, the recent climatological studies of Rossby wave breaking cited in the paper provide support for the observed seasonality and timing of "flooding" of moist monsoon air into the extratropical lower stratosphere and for the dominant locations of poleward wave breaking events pointed out in Figure 16. Such additional detail on these linkages will provide better continuity in the Results section and strengthen the presentation and discussion of the conceptual model given in Figure 16.

Many thanks for this suggestion. We agree that such details would strengthen our results and added therefore the following paragraph to the introduction:

"Exchange of air masses from the troposphere to the stratosphere occurs preferably in poleward flow structures such as tropospheric intrusions (e. g. Sprenger et al., 2007; Pan et al., 2009; Vogel et al., 2011). These intrusions develop into elongated potential vorticity (PV) streamers and are manifestation of Rossby wave breaking. Rossby wave breaking is identified as an important mechanism for exchange of air masses between the tropical upper troposphere and the extratropical lower stratosphere with a pronounced peak during summer in each hemisphere controlled by the presence of monsoon anticyclones, in particular of the Asian monsoon (e. g. Homeyer and Bowman, 2013; Kunz et al., 2015). High frequency of PV streamers are found over the eastern North Pacific and over the Atlantic in summer demonstrated in a recently published climatology of PV streamers (Kunz et al., 2015)."

2. While the authors do a good job of including observational support for the model results with analysis of a few flights during the TACTS/ESMVal campaign, a more thorough test of the CLaMS model (particularly related to the influence of Asian monsoon air on extratropical lower stratosphere water vapor – i.e., Figure 15) using the same chemical dataset that was used for initialization – Aura MLS – is requested. Is the enhancement in extratropical lower stratosphere water vapor at 380 K following breakup of the Asian monsoon anticyclone an observed characteristic? I expect this is a straight-forward test of the model and would go a long way in strengthening the perceived impact of this paper. If not an observed characteristic, this is a questionable result.

We followed the reviewers advice and performed a comparison between CLaMS and MLS water vapor in the extra-tropical northern lower stratosphere. Overall, MLS water vapor measurements confirm our model results. Both MLS and CLaMS water vapor clearly show an increase of water vapor during summer and autumn 2012 in the northern extra-tropical lower stratosphere. However some differences between CLaMS and MLS remains. This differences are discusses within the revised version of our paper as follows.

"Because water vapor is an important greenhouse gas and even small perturbations of water vapor mixing ratios in the ExUTLS have a significant impact on surface climate, we are interested to estimate the impact of the Asian monsoon on moistening the lower stratosphere. From our simulations, we roughly estimate the fraction of H_2O originating in India/China, Southeast Asia, and tropical Pacific Ocean contributing to the water budget in the lower northern hemisphere stratosphere. Fig. 1 shows the mean water vapor content in the northern lower stratosphere for PV values larger than 7.2 PVU (10 PVU) and northward of 30° N at 380 K (400 K) calculated with CLaMS (black line). In CLaMS at this altitude, contributions of cirrus clouds to the total water content are of minor importance. An increase of H_2O in the northern lower stratosphere is found in our simulation during summer and autumn as reported in previous studies (e.g. Ploeger et al., 2013; Zahn et al., 2014; Müller et al., 2015). The fraction of H₂O from different boundary tracers is indicated by different colors. End of October 2012, a contribution of approximately 1.5 ppmv (1.0 ppmv) H₂O originates from source regions in Asia and the tropical Pacific compared to a mean water vapor content of \approx 5 ppm (4.5 ppm) at 380 K (400 K). Total H₂O without contributions of all boundary tracers i. e. the contribution of aged air (see Fig. 1, green line) shows a decrease during summer and autumn. Mean H₂O from AURA-MLS (version 3.3 and version 4) in the northern stratosphere calculated similar as mean CLaMS H_2O values for PV values larger than 7.2 PVU (10 PVU) and northward of 30° N at 380 K (400 K) also show an increase of water vapor within the lower northern stratosphere during summer and autumn 2012 (see Fig. 1, gray and purple line) and therefore support our findings from CLaMS simulations. Differences between CLaMS and MLS mean values in

the lower northern stratosphere in particular at 380 K could by explained by sampling issues (different spacial resolution of MLS and CLaMS), vertical resolution of MLS limiting measurements of steep tracer gradient around the tropopause and the initialization of CLaMS H_2O at 1 May 2012 (above 400 K: AURA-MLS; below 350 K: CLaMS multi-annual simulation based on ERA-Interim water vapor with a linear transition between 350 K and 400 K, more details see (Vogel et al., 2015)). Previous comparisons between CLaMS and MLS (v3.3) water vapor demonstrate that differences in H_2O found in the lower stratosphere at high latitudes are likely an artifact of the MLS averaging kernels (Ploeger et al., 2013). Further, a comparison of water vapor climatologies from international limb sounder by Hegglin et al. (2013) demonstrate that Aura-MLS (v3.3) H_2O measurements tend to be low at high latitudes in the lowermost stratosphere. Because of the good agreement between CLaMS H₂O and in situ water vapor measurements by the FISH instrument during TACTS/ESMVal in the lower northern stratosphere (see Figs. 9-11) we are confident that CLaMS water vapor simulations within the extra-tropical lower stratosphere are reliable."

3. The naming convention for emission sources outside of the Asian monsoon region is inconsistent in the text and Figures of the manuscript. While the use of "residual" is common and seems to be the primary intention of the authors, "residue" appears in other places (e.g., Figures 5, 6, 14, and 15; Table 2; and Page 12, line 23; Page 12, line 29). Please update the text and figures to refer to this as "residual" throughout.

We agree that within the paper the naming convention should be consistent and would like to thank the reviewer for carefully reading the manuscript. We changed "residue" to "residual" overall in the revised version of the paper (both within text and figures).

4. While the figures are (for the most part) visually appealing, an effort should be made to have the spacing, scaling, and text sizes consistent throughout. For example, the color bar labels run into the latitude axes in Figures 3 and 4, the cross-sections are misaligned in Figures 5 and 6, text sizes of the two panels in Figure 7 are different, the bottom panel is unnecessarily displaced from the top three in Figure 12, and the text sizes in Figures 14 and 15 are not legible at normal zoom.



Figure 1: (see Fig. 15 in ACPD paper) The increase of H_2O mixing ratios in the lower northern hemisphere stratosphere at 380 K (top) and at 400 K (bottom) during summer 2012 is shown (black line). The green line indicated H_2O mixing ratios without fractions from the Earth's boundary layer. Mean H_2O mixing ratios derived from MLS version v3.3 (gray) and version v4 (purple) in the northern extra-tropical lower stratosphere show also an increase in water vapor during summer and autumn. A rough estimation of the fraction of H_2O mixing ratios originating in India/China (red), Southeast Asia (yellow) and the tropical Pacific Ocean (blue) in the northern lower stratosphere is also given.

We revised the figures according the reviewers advice.

Specific Comments

- 1. Page 1, line 7: "...jet such as the..." should be "jet such that the" done
- 2. Page 1, lines 16-17: This statement is confusing here. This should be clarified to say that sources from Asia and the tropical Pacific account for ~ 1.5 ppmv of the ~ 5 ppmv mean in the extratropical lower stratosphere.

We revised the sentence as follows: "End of October 2012, approximately 1.5 ppmv H₂O is found in the lower northern hemisphere stratosphere (at 380 K) from source regions both in Asia and in the tropical Pacific compared to a mean water vapor content of ≈ 5 ppmv."

- 3. Page 2, line 3: "...is acting..." should be "...acts..." done
- 4. Page 3, lines 16-23: This discussion is limited to large-scale transport processes, correct? For example, we know that moist convection (apart from a large organized system like a typhoon) is capable of transporting air across the tropopause but such small-scale processes (though possibly represented to an extent) are not resolved in these models. A bit more detail and context should be given to clarify these points here, which you do reflect on near the end of the paper.

The following sentence is added: "In addition to this large-scale transport process of water vapor by air mass exchange between the tropics and the extratropics, convection induced injections of water vapor in mid-latitudes can also occur in large storm systems such as tropical cyclones and by deep continental convection (e. g. Anderson et al., 2012; Homeyer et al., 2014; Vogel et al., 2014)."

- 5. Page 3, line 27: "...are used..." should be "...is used..." done
- 6. Page 4, line 14: "Measurements of..." should be "Measurements from..." done
- 7. Page 6, line 30: "...occurs like for all..." should be "...occurs in an equivalent fashion to all..."

on page 5, line 30: "...occurs in an equivalent way to all chemical species" is added

- 8. Page 7, line 20: "...as for..." should be "...to..." done
- 9. Page 8, line 12: "...into stratosphere." should be "...into the stratosphere." done
- 10. Page 8, lines 15-18: The Homeyer et al 2011 paper you cite can be referenced here as well. done
- 11. Page 10, lines 11-12: What do you mean by this statement? It takes 5 weeks for the parcels to be transported from their surface emission locations to the lower stratosphere over Europe? This statement needs to be clarified a bit.

The following sentence is added: "These air masses uplifted by typhoon Bolaven are transported from the Earth's surface over the West Pacific within 5 weeks to the lower stratosphere over Europe [Vogel et al. 2014]." 12. Page 10, lines 19-21: Based on the time series, No. 3 is an aircraftonly signature (i.e., no apparent plume in the CLaMS simulation – at least not to me!).

That is correct, therefore we revise the text as follows: "During the second part of the flight on 26 September 2012 (see Fig. 9, top), further signatures of tropospheric air are measured during the flight (No. 3-6). Also here, enhanced percentages of the emission tracers for India/China and Southeast Asia / tropical Pacific Ocean up to 18 % are simulated, except for region No. 3."

13. Page 10, lines 26-18: But is this really dynamics of the AMA or of a downstream RWB event? You have already demonstrated that the latter is the reason this particular air mass crossed the tropopause, correct?

We revise the sentence (page 10, line 26-28) as follows: "Our simulations in agreement with measurements show that the amount of water vapor and pollution in the lower stratosphere is enhanced in the Northern Hemisphere in September 2012 associated with both the dynamic of the Asian monsoon anticyclone and the transport of air masses from both Asia and the tropical Pacific along the subtropical jet."

- 14. Page 10, line 35: "Northern Hemisphere" should be "stratosphere" revised to "... into the northern extratropical lower stratosphere..."
- 15. Page 11, line 22: Remove "these" done
- 16. Page 12, line 25: "...masses on the..." should be "...masses to the...", and "End October..." should be "End of October..." done
- 17. Page 12, line 27: "Here, highest contributions are from tropical..." should be "Here, the highest contributions are from the tropical..." done
- Page 12, line 31: "This is in particular..." should be "This is particularly..." done

References

- Anderson, J., Wilmouth, D., Smith, J., and Sayres, D.: UV Dosage Levels in Summer: Increased Risk of Ozone Loss from Convectively Injected Water Vapor, Science, 337, 835–839, doi:10.1126/science.1222978, 2012.
- Hegglin, M. I., Tegtmeier, S., Anderson, J., Froidevaux, L., Fuller, R., Funke, B., Jones, A., Lingenfelser, G., Lumpe, J., Pendlebury, D., Remsberg, E., Rozanov, A., Toohey, M., Urban, J., von Clarmann, T., Walker, K. A., Wang, R., and Weigel, K.: SPARC Data Initiative: Comparison of water vapor climatologies from international satellite limb sounders, J. Geophys. Res., 118, 11824–11846, doi:doi:10.1002/jgrd.50752, 2013.
- Homeyer, C. R. and Bowman, K. P.: Rossby Wave Breaking and Transport between the Tropics and Extratropics above the Subtropical Jet, J. Atmos. Sci., 70, 607–626, 2013.
- Homeyer, C. R., Pan, L. L., Dorsi, S. W., Avallone, L. M., Weinheimer, A. J., O'Brien, A. S., DiGangi, J. P., Zondlo, M. A., Ryerson, T. B., Diskin, G. S., and Campos, T. L.: Convective transport of water vapor into the lower stratosphere observed during double-tropopause events, J. Geophys. Res., 119, 10941–10958, doi:10.1002/2014JD021485, 2014.
- Kunz, A., Sprenger, M., and Wernli, H.: Climatology of potential vorticity streamers and associated isentropic transport pathways across PV gradient barriers, J. Geophys. Res., 120, 3802–3821, doi:10.1002/2014JD022615, URL http://dx.doi.org/10.1002/2014JD022615, 2014JD022615, 2015.
- Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and Engel, A.: Impact of the Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe 2012, Atmos. Chem. Phys. Discuss., 15, 34765–34812, doi:10.5194/acpd-15-34765-2015, 2015.
- Pan, L. L., Randel, W. J., Gille, J. C., Hall, W. D., Nardi, B., Massie, S., Yudin, V., Khosravi, R., Konopka, P., and Tarasick, D.: Tropospheric intrusions associated with the secondary tropopause, J. Geophys. Res., 114, D10302, doi:10.1029/2008JD011374, 2009.
- Ploeger, F., Günther, G., Konopka, P., Fueglistaler, S., Müller, R., Hoppe, C., Kunz, A., Spang, R., Grooß, J.-U., and Riese, M.: Horizontal water

vapor transport in the lower stratosphere from subtropics to high latitudes during boreal summer, J. Geophys. Res., 118, 8111–8127, doi:10.1002/jgrd. 50636, 2013.

- Sprenger, M., Wernli, H., and Bourqui, M.: Stratosphere-Troposphere Exchange and Its Relation to Potential Vorticity Streamers and Cutoffs near the Extratropical Tropopause, J. Atmos. Sci., 64, 1587–1602, doi:10.1175/JAS3911.1, URL http://dx.doi.org/10.1175/JAS3911.1, 2007.
- Vogel, B., Pan, L. L., Konopka, P., Günther, G., Müller, R., Hall, W., Campos, T., Pollack, I., Weinheimer, A., Wei, J., Atlas, E. L., and Bowman, K. P.: Transport pathways and signatures of mixing in the extratropical tropopause region derived from Lagrangian model simulations, J. Geophys. Res., 116, D05306, doi:10.1029/2010JD014876, 2011.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Hoor, P., Krämer, M., Müller, S., Zahn, A., and Riese, M.: Fast transport from Southeast Asia boundary layer sources to northern Europe: rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon anticyclone, Atmos. Chem. Phys., 14, 12745–12762, doi:10.5194/acp-14-12745-2014, URL http:// www.atmos-chem-phys.net/14/12745/2014/, 2014.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, Atmos. Chem. Phys., 15, 13699-13716, doi:10.5194/acp-15-13699-2015, URL http:// www.atmos-chem-phys.net/15/13699/2015/, 2015.
- Zahn, A., Christner, E., van Velthoven, P. F. J., Rauthe-Schöch, A., and Brenninkmeijer, C. A. M.: Processes controlling water vapor in the upper troposphere/lowermost stratosphere: An analysis of 8 years of monthly measurements by the IAGOS-CARIBIC observatory, J. Geophys. Res., 119, 11,505–11,525, doi:10.1002/2014JD021687, 2014.

Author Comment to Referee #2

ACP Discussions doi: 10.5194/acp-2016-463 (Editor - Peter Haynes) 'Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012'

We thank Referee #2 for the very helpful review. Our reply to the reviewer comments is listed in detail below. Questions and comments of the referee are shown in italics.

Summary: In this study the authors examine the transport characteristics associated with the Asian summer monsoon during September-October 2012, using both measurements of trace gases (e.g. ozone, water vapor, methane) and idealized tracer simulations that provide information about the relative contributions of different boundary layer regions to upper tropospheric/lower stratospheric air masses. The study provides strong evidence that the eastward shedding of eddies from the monsoon anticyclone provides an important mechanism for transporting boundary layer air from India and South Asia to the lower stratosphere over northern midlatitudes. While this study provides a comprehensive analysis and important contribution to the field that will make it suitable for publication, I have a few major comments that need addressing before I recommend its publication. In particular I have one major concern about the authors' interpretation of the air-mass origin tracers that needs addressing, as it may potentially affect the interpretation of the main results. I also have smaller comments that are, by comparison, less important.

We thank the reviewer for the opportunity to remove unclarities about the emission tracer approach used in our paper. A detailed discussion follows below.

Major Comment

1. I am concerned about the interpretation of the air-mass tracers as a fraction. It is definitely constructive to look at the relative contributions of

different source regions and I commend the authors' use of the diagnostic. However, more care should be taken in the interpretation of the tracer concentration as giving the fraction of air that was last at the earth's surface in a given source region. In particular, the simulation only covers 1 May 2012 -31 October 2012. If the tracers are to be interpreted as fractions (as the authors intend them to be) then the sum of the air-mass tracers corresponding to all of the source regions must equal 1 (since the union of the source regions is the entire planetary boundary layer (PBL)). This is not the case, however, as shown in Figure 9. The sum of the red, blue and orange lines should. in principle, equal 1 (but does not). What this tells me is that the tracers have not been integrated to equilibrium so that there is a large amount of air that is not accounted for by the source fractions. This is a known issue when dealing with air-mass origin tracers (Orbe et al. (2015)) and I am concerned about what this means for the main conclusions in the study. Please either start the simulation much earlier (to ensure tracer equilibration by September 2012) or remove all references to "fraction" because this interpretation is not correct. Alternatively, it is possible that I am missing something important in the authors' definition of "residual" (by which I interpret the rest of the *PBL*) - *if this is the case, please clarify in the text.*

Orbe, Clara, Paul A. Newman, Darryn W. Waugh, Mark Holzer, Luke D. Oman, Feng Li, and Lorenzo M. Polvani. "Airmass Origin in the Arctic. Part I: Seasonality." Journal of Climate 28, no. 12 (2015): 4997-5014. (Figure 3b)

The reviewers interpretation of the residual is correct, however we believe that there is a misunderstanding regarding an important point in our approach: The composition of air in the UTLS is always a combination of aged air masses from the stratosphere and free troposphere and young freshly injected air masses from the boundary layer. Only for the young air masses, our characterization of air mass origin is conducted.

In our approach we cannot run the atmosphere to equilibrium as we are interested in working out the contribution of young, freshly injected air to the composition of both the Asian monsoon anticyclone and the extratropical lower stratosphere. In so far our approach is different but complementary to the approach by Orbe et al..

We are aware that we had not brought this point across very well in our original paper (ACPD Version). We have now improved the presentation of our modeling concept and its impact on the results throughout the paper (see in particular Section 3 in the revised version).

To go in more detail, the idea of using artificial tracers to identify transport has been used for a long time in both Eulerian (e.g. Mahlman and Moxim, 1978; Stone et al., 1999) and Lagrangian models (e.g. Bowman and Cohen, 1997; Bowman and Carrie, 2002; Günther et al., 2008; Vogel et al., 2011). In our approach, the artificial emission tracers in CLaMS are designed to identify possible boundary source regions in Asia that could contribute to the composition of the Asian monsoon anticyclone and the extratropical lower stratosphere during the course of the monsoon season 2012 considering advection and mixing processes.

The CLaMS transport scheme consists of advection, which is the reversible part of transport (the trajectory), and mixing, the irreversible part of transport. At beginning of the CLaMS model simulation each air particle is marked by a boundary layer tracer (Ω). That means, for example, that the boundary layer tracer (Ω) for air parcels in the boundary layer (BL) is set equal to one and everywhere else (free troposphere and stratosphere) equal to zero. Mixing in CLaMS between an air parcel from the boundary layer (Ω $(\Omega = 1)$ and an air parcel from the free troposphere or stratosphere ($\Omega = 0$) is implemented by insertion of a new particle. The boundary layer tracer (Ω) of the new particle is then equal to 0.5. Successive mixing processes between boundary layer air ('young air masses') and air masses from the free troposphere or stratosphere ('aged air masses') during the course of the simulation yield a tracer distribution differing from the initial distribution ($\Omega = 1$ or Ω = 0). In the CLaMS simulation used here, air masses in the model boundary layer are marked by boundary emission tracers every 24 h (the time step for mixing in CLaMS) during the simulation period from 1 May 2012 until end of October 2012. Thus in our simulations, the age spectrum is relatively simply constructed from the response to a single pulse during $(t_i, t_i + \Delta t_i)$ for times $t > t_i$ (see Holzer et al., 2009), with t_i equal to 1 May 2012 and Δt_i equal to 6 months. This pulse marks 'young air masses'. E.g. a value of Ω equal to 0.4 means that 40% of the air parcel is younger than Δt_i (= 6 months; 'young air masses') and 60 % are older than Δt_i released from the model boundary layer before 1 May 2012 (emission time $t_e < t_i$; 'aged air masses'). We used this single pulse approach because the focus of the paper is to analyze the influence of fresh emissions from different boundary regions using the meteorological conditions of the year 2012. Here, the air-mass fraction $f(r,t \mid \Omega)$ is defined to be the fraction of air at a location r and time t

(with $t_i \leq t \leq t_i + \Delta t_i$) that was emitted since t_i within Δt_i from the model boundary layer (BL).

To get the information about the origin of young air masses within the model boundary layer, we divided the boundary layer (BL) in different regions, i.e. different artificial tracers of air mass origin, referred to as "emission tracers" (Ω_i of the number n = 17) are introduced in CLaMS (see paper table 1). Within the boundary layer, the sum of all different emission tracers (Ω_i) is equal to 1 ($\Omega = \sum_{i=1}^{n} \Omega_i = 1$). During the course of the simulation, the air-mass fraction $f_i(r, t \mid \Omega_i)$ is defined to be the fraction of air at a location r and time t (with $t_i \leq t \leq t_i + \Delta t_i$) that was emitted since t_i within Δt_i from the model boundary layer (BL) in region Ω_i . Note however that emission tracers only describe the contribution of the young air masses.

The fraction of air from different boundary regions $\sum_{i=1}^{n} f_i(r, t \mid \Omega_i) = f(r, t \mid \Omega)$ shown in Fig. 9 and also Figs. 10 and 11 (red, blue, orange) is less then 100% because only "young air masses" emitted since 1 May 2012 are counted. Air masses originating in the free troposphere or stratosphere ('aged air masses') also contribute to the composition of the extratropical lower stratosphere. Thus in Fig. 9 region No.2 (gray area), the sum of air masses originating in India/China, in Southeast Asia/tropical Pacific and in the residual surface $(\sum_{i=1}^{n} f_i(r, t \mid \Omega_i))$ is roughly 45%. The remaining 55% of the composition of the lower stratosphere in this region is from the free troposphere and the stratosphere. Fig. 14 illustrate how the contribution of the model boundary layer $(f(r, t \mid \Omega))$ to the extratropical stratosphere rises to 44% at 360 K, to 35% at 380 K, and to 23% at 400 K (see Sect. 4.4.2) from May until the End of October 2012. The remaining percentages to 100% are contributions from "aged air masses" originating in the free troposphere and stratosphere.

We added the following sentence: 'On 26 September 2012 (see Fig. 9, top), a very pronounced signature of tropospheric air in the lower stratosphere is found between 09:05 UTC to 10:17 UTC (No. 2). Here, the contributions of the emission tracer for India/China and Southeast Asia /tropical Pacific Ocean are up to 20 % and 23 %, respectively (up to 5 % from the residual surface). Thus, the sum of all emission tracers for model boundary layer is roughly 48 %. The remaining 52 % of the composition of the lower stratosphere in this region is from aged air masses originating in the free troposphere and the stratosphere at the beginning our the CLaMS simulation on 1 May 2012.' The technique used in our paper is different compared to the approach used in Orbe et al. (2015). A detailed comparison of our approach and the approach used in Orbe et al. (2015) is published in (Vogel et al., 2015, Sect. 4 Discussion). For clarification we repeat the corresponding paragraphs published in Vogel et al. (2015) here in our authors' comment:

"In contrast to our study, Orbe et al. (2015) used tracers of air mass origin in model simulations to infer the impact of boundary regions in Asia to the tropical lower stratosphere with an approach that the different tracers have equilibrated so that the sum of all tracers of air mass origin is equal to unity within the entire atmosphere (using a spin-up of 20 years). This approach provides for each air parcel the information about the origin within the boundary layer. However, this approach provides no information on the transport times from the boundary layer in Asia to the tropical lower stratosphere. To infer the transport times in the approach by Orbe et al. (2015), they introduce a boundary impulse response that marks air that left the boundary layer on a certain day (1 July). They infer transport times of 1 month from the boundary layer in Asia into the tropical lower stratosphere in July when the Asian monsoon is active. In this respect, their pulse serves a similar purpose as our seasonal tracer set-up. However, our approach considers all air masses that left the boundary layer over the course of the 2012 monsoon season since May and thus reflects the meteorological conditions of the entire 2012 monsoon season."

Specific Comments

1. Line 29, Page 3: I am wary of the use of the term "transport pathways." The air mass origin tracers only tell you where air was last in contact with the boundary layer. They do not provide a sense for how the air arrived at the receptor location, so please remove all references to pathways. To infer pathways you would need to use idealized tracers similar to those used for inferring the age spectrum or, most appropriately, the path density tracers examined in Holzer (2009)

Holzer, Mark. "The path density of interhemispheric surface-to-surface transport. Part I: Development of the diagnostic and illustration with an analytic model." Journal of the Atmospheric Sciences 66, no. 8 (2009): 2159-2171

Our approach allows the relative contribution of air masses of varying origin to be quantified for every individual CLaMS air parcel on a certain location (r_j) and for each time (t_l) during the course of the simulation. Thus, transport pathways from the boundary layer into the UTLS (occurring since 1 May 2012) can be inferred analyzing the spatial-temporal evolution of artificial emission tracers during the simulation period. The change of the fraction of air originating in region Ω_i on different locations (r_j) for different times $(t_l) f_{ijl}(r_l, t_j | \Omega_i)$ define in our approach the transport pathways. However, we agree with the reviewer that the term "pathway" might have been used a bit to excessively and have removed the word on various locations in the paper.

2. Line 8, page 6: Again, reservation about the word "pathway."

okay, we dropped "pathway"

3. Lines 33-35, page 6: The sum of all of the air-mass fractions does not equal 1, leaving a large fraction of air unaccounted for. Therefore, I am not confident in the statement that "that air masses originating in India/China and Southeast Asia/Pacific Ocean almost exclusively contribute to the chemical composition of the separated anticyclone." Please either start your simulation earlier (to ensure equilibration of the airmass tracers) or do not use the word "fraction".

We agree the air masses originating in India/China and Southeast Asia/Pacific Ocean do not exclusively contribute to the chemical composition of the separated anticyclone. For clarification we added the following sentence to the paper:

"Further, the emission tracer for Southeast Asia / tropical Pacific Ocean contributes to the composition of the Asian monsoon anticyclone, however to a smaller extent compared to the emission tracer for India/China as shown in Fig. 4 (right). Contributions from the residual surface are of minor importance (see Sect. 4.2). Note that contribution of air masses originating in the free troposphere and stratosphere ('aged air masses') als contribute to the composition of the Asian monsoon anticyclone of about 25% at the end of September 2012 (see in Vogel et al., 2015, Fig. 9)."

References

- Bowman, K. P. and Carrie, G. D.: The mean-meridional transport circulation of the troposphere in an idealized GCM, J. Atmos. Sci., 59, 2002.
- Bowman, K. P. and Cohen, P. J.: Interhemispheric exchange by seasonal modulation of the Hadley circulation, J. Atmos. Sci., 54, 1997.
- Günther, G., Müller, R., von Hobe, M., Stroh, F., Konopka, P., and Volk, C. M.: Quantification of transport across the boundary of the lower stratospheric vortex during Arctic winter 2002/2003, Atmos. Chem. Phys., 8, 3655-3670, URL http://www.atmos-chem-phys.net/8/3655/2008/, 2008.
- Mahlman, J. and Moxim, W.: Tracer simulation using a global general circulation model - results from a mid-latitude instantaneous source experiment, J. Atmos. Sci., 35, 1340-1374, 1978.
- Orbe, C., Waugh, D. W., and Newman, P. A.: Air-mass origin in the tropical lower stratosphere: The influence of Asian boundary layer air, Geophys. Res. Lett., 42, 2015GL063937, doi:10.1002/2015GL063937, URL http: //dx.doi.org/10.1002/2015GL063937, 2015.
- Stone, E. M., Randel, W. J., and I. Stanford, J.: Transport of passive tracers in baroclinic wave life cycles, J. Atmos. Sci., 56, 1364–1381, 1999.
- Vogel, B., Pan, L. L., Konopka, P., Günther, G., Müller, R., Hall, W., Campos, T., Pollack, I., Weinheimer, A., Wei, J., Atlas, E. L., and Bowman, K. P.: Transport pathways and signatures of mixing in the extratropical tropopause region derived from Lagrangian model simulations, J. Geophys. Res., 116, D05306, doi:10.1029/2010JD014876, 2011.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, Atmos. Chem.

Phys., 15, 13699-13716, doi:10.5194/acp-15-13699-2015, URL http://www.atmos-chem-phys.net/15/13699/2015/, 2015.

Author Comment to Referee #3

ACP Discussions doi: 10.5194/acp-2016-463 (Editor - Peter Haynes) 'Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012'

We thank Referee #3 for the good evaluation of our paper. Following the reviewers advice we revised some parts of the paper for the purpose of clarification. Our reply to the reviewer comments is listed in detail below. Questions and comments of the referee are shown in italics.

The authors use the global Lagrangian CLaMS model, with artificial and chemical constituent tracers to quantify the contributions of different boundary Layer (BL) source regions in Asia to the Asian Summer Monsoon (ASM) anticyclone, and from there to extra-tropical lower stratosphere (ExLS), for the 2012 ASM season. Further, they illustrate the transport pathways for BL source air, accumulated in the ASM anticyclone, to reach the ExLS, via eddy shedding in the upper troposphere, subsequent filamentation and penetration into the stratosphere, associated with Rossby wave breaking along the subtropical jet. They also consider the westward shedding of air from the ASM anticyclone into the tropical upper troposphere.

The authors use the simulated artificial tracers, and simulated ozone, CO, and water vapor, to interpret small-scale structures observed along aircraft flight tracks as filamentary intrusions of BL source air associated with the ASM anticyclone into the lower stratosphere over Northern Europe. Further, they use the artificial tracers to quantify the contribution of different BL source regions to the ExLS over the 2012 ASM season, and use CLaMS simulated water vapor to estimate the contribution of Asian source regions to water vapor in the ExLS.

The study builds upon earlier studies that have illustrated tropospherestratosphere exchange (STE) mechanisms associated with the ASM, going back at least as far as Dethof et al. (1999), a paper which the authors cite. At the same time the use of the artificial and constituent tracers with the CLaMS model to quantify estimates of the contributions from Asian (and other) BL source regions to the ExLS is I believe a step beyond these earlier studies. Quantification of water vapor contribution to the NH lower stratosphere (p.13) is particularly interesting. The cross-section of the filamentary structure (Fig. 6) provides an illuminating illustration of intermediate (mixed) constituent and stability conditions between the tropospheric and stratospheric air mass characteristics.

The comparison of aircraft observations of the low stratosphere over Northern Europe with CLaMS tracer maps interpolated to the aircraft flight tracks illustrates effectively that the simulation of tropospheric filaments in the low stratosphere represents real-world conditions, and supports the quantitative estimates of BL source influence in the ExLS presented later. The analysis and interpretation is reminiscent of that conducted by Fairlie et al. (2007) for INTEX-NA aircraft observations; the authors may wish to add correlation scatter plots of the observed O3, CO or CH4, water vapor to further illuminate air mass origin and characteristics of mixed troposphere-stratosphere air masses.

I think the paper could use an editorial review for the English and sentence structure. There is occasional awkwardness in the sentence structure, and some choice of wording that I find confusing, and may be a translation issue (see some examples below). Nevertheless, I think the paper is suitable for publication in ACP given consideration to these issues and the points listed below, most of which are minor and for the purpose of clarification.

We agree with the reviewer that tracer-tracer correlations are a powerful technique to study mixing processes between the troposphere and stratosphere. Tracer-tracer correlations would integrate the information from the aircraft observations and the CLaMS model to quantify and characterize signatures of mixing between stratospheric and tropospheric air (e.g. monsoon air) and therefore give insights in single mixing processes. However, in this paper we didn't show tracer-correlations because that would be material for an additional new study that we are planing. Therefore, to analyze tracer-tracer correlations of different chemical species from different TACTS/ESMVal measurements and tracer-tracer correlations from the CLaMS model using in addition tracers of air mass origin (e.g. similar as in Vogel et al. (2011)) would go beyond the scope of the paper discussed here. For the TACTS flight on 30 August 2012 an analysis of tracer-tracer correlations (O_3 - CO) is already published in Müller et al. (2016).

We would like to thank the reviewer for carefully reading our paper and identifying some parts of the paper that need revisions for the purpose of clarification (details see below 'Minor Comments'). For the most comments identifying ambiguities, the reviewers interpretation was correct (see below). Following the reviewers advice, we revise those parts of the text. In general, some English corrections are already made in the revised version of the paper. After the acceptance of our paper by ACP a language revision of the paper will be made by the Language Services of Forschungszentrum Jülich.

Minor Comments

1. p.6, line 31, What is meant by "Maritime Continent"?

The expression "The Maritime Continent" describes the region between the Indian and Pacific Oceans including the archipelagos of Indonesia, Borneo, New Guinea, the Philippine Islands, the Malay Peninsula, and the surrounding seas.

We added in the revised version of the paper: '...Maritime Continent (the region between Indian and Pacific Oceans)...'

2. p.7, line 12-13: Comment: The authors will recognize that the transport is only irreversible if the tropospheric intrusion is mixed into the stratospheric surroundings. It is conceivable that an intrusion across the PV=7.2 PVU could return to the troposphere downstream.

We agree that the sentence can be misinterpreted, therefore we remove 'irreversible' and write in the revised version: 'Isentropic transport of air masses across the 7.2 PVU isoline indicates exchange between the tropics and extratropics due to wave breaking.'

3. p.7, lines 14-15: Comment: I am unable to see the PV=7.2 PVU isopleth enveloping a "region of enhanced tracers"

We agree that the sentence is somewhat unclear. We revised the sentence as follows: 'On 20 September 2012, at the northern flank of the separated anticyclone the 7.2 PVU isoline is in the region where strongest gradients of the emission tracers occur indicating the transport barrier at 380 K (see Fig. 3 and 4).'

4. p.7, line 31, instead of "surface that is" do you mean "surface, i.e.,"? I.e., are the authors stating the definition of "residual" here?

For clarification we introduced a definition for the 'residual surface' in Sect. 3 (CLaMS simulations using artificial tracers of air mass origin) as follows:

'The most important regions for our study are India/China (= Northern India (NIN) + Southern India (SIN) + Eastern China (ECH)), Southeast Asia (SEA) and the tropical Pacific Ocean (TPO). The sum of all model boundary layer tracers ($\Omega = \sum_{i=1}^{n} \Omega_i$) without contributions from India/China, Southeast Asia, and the tropical Pacific Ocean is summarized in one emission tracer referred to as "the residual surface" (= Ω - NIN - SIN - ECH - SEA - TPO).'

and revised the sentence on page 7, line 31, Sect. 4.1, as follows:

'Fig. 5 (bottom left) shows the emission tracer for the residual surface (the entire model boundary layer without contributions from India/China, Southeast Asia, and the tropical Pacific Ocean, see Sect. 3).'

5. p.7, line 32-33, reference "no signature." Would the authors be more quantitative here? Looks like up to 10-15% is due to "residual" sources in the anticyclone.

We revised the sentence as follows: 'The contribution of the emission tracer for the residual surface is approximately 10-15 % within the separated anticyclone indicating that air masses originating in India/China

and Southeast Asia / tropical Pacific Ocean almost exclusively contribute to the chemical composition of the separated anticyclone.'

6. p.8, line 12, reference "indicating transport from the troposphere into the stratosphere." This is according to the definition of the authors, based on the work of Kunz et al.

Yes, that is correct. Therefore we revised the sentence to be more precise as follows: 'At 380 K enhanced contributions of the emission tracer from India/China are also found north of the 7.2 PVU barrier indicating transport from the troposphere into the stratosphere according to the definition by Kunz et al. (2015).'

7. p.9, reference discussion of Fig. 9 emission tracer plots, here and elsewhere. Please confirm for the reader if "residual" includes all BL surfaces other than those identified (China/India, SEAsia/ tropical Pacific). How should the reader interpret the sum of these percentages being much less than 100%, e.g. does the remainder comprise background lower stratospheric air, unconnected to any BL surface in past 5 months?

As mentioned above we added a definition for 'residual surface' in Sect. 3 (CLaMS simulations using artificial tracers of air mass origin) in the revised version of the paper. The interpretation that the sum of the percentages of all boundary layer tracers is less than 100% because of contributions of aged air masses from the stratosphere is correct. For further clarification we added the following sentence in the revised version:

'On 26 September 2012 (see Fig. 9, top), a very pronounced signature of tropospheric air in the lower stratosphere is found between 09:05 UTC to 10:17 UTC (No. 2). Here, the contributions of the emission tracer for India/China and Southeast Asia /tropical Pacific Ocean are up to 20 % and 23 %, respectively (up to 5 % from the residual surface). Thus, the sum of all emission tracers for model boundary layer is roughly 48 %. The remaining 52 % of the composition of the lower stratosphere in this region is from aged air masses originating in the free troposphere and the stratosphere at the beginning our the CLaMS simulation on 1 May

8. p. 10, discussion of Figs. 9-11. It would be helpful if the authors labeled the locations of flight segments "1", "2", "3", etc. on the maps in Figs. 7-8, to help the reader identify the features highlighted in the flight data to features on the CLAMS maps. Additionally, the flight data appears to be higher temporal resolution than the CLAMS profiles (e.g. the profile of FISH H2O). It may be helpful to add a time-averaged data profile at the same resolution as the CLAMS for better comparison. Tracer-tracer correlation plots may also be a useful addition (see above).

We agree that numbers in Fig. 7-8 can help the reader to identify the regions that are impacted by young tropospheric air masses. We revised figures 7-8 and also Fig. 10 and 11 as shown below.

Within Fig. 9-11, the measurements of CO, CH_4 , H_2O and O_3 are shown in the highest resolution that was available for each species to demonstrate the variability of the measurements. For the interpolation of CLaMS results as described in Appendix B, backward trajectories are calculated every second along the flight path.

As discussed above an analysis of tracer-tracer correlations would be beyond the scope of this paper.

9. p. 11, lines 13-15. Suggestion: I think the authors mean to emphasize the locations (Atlantic and Pacific Oceans) here, rather than the mechanism (Rossby wave breaking). They may want to leave out "Rossby wave breaking" in this sentence to keep the stress on the locations.

That is a good point. We changed the sentence as follow: 'Finally, Fig. 11 shows that transport from air masses originating in the Asian monsoon anticyclone into the lower stratosphere take place most frequently over the Pacific and Atlantic Ocean'

10. p.11, lines 18-20. This sentence seems a bit out of context here. Perhaps reference to SE Asia/ Tropical Pacific contribution (the appendix) would sit better after the introduction to Fig.12 (p.11, line 2, after "September 2012").



Figure 1: (= Fig. 7 within the ACPD paper) Horizontal (top) cross-section of the fraction of air originating in India/China over Europe om 26 September 2012. The flight path transferred to noontime of the TACTS/ESMVal flight is shown as yellow line. Segments of the flight in the lower stratosphere with enhanced measured CO, CH_4 , and H_2O and reduced O_3 compared to the stratospheric background are highlighted in red and numbered for clarification (the same numbers are used in Fig. 9 The climatological isentropic transport barrier of 7.2 PVU at 380 K is shown as thick black line. The white thick line marks the position of the vertical (bottom) cross-section at 8°W longitude which is similar to the cross-sections shown in Fig. 5 and 6.

We agree. The paragraph is shifted as proposed.

11. p.11, reference discussion of "transport pathways": The title of 4.4.1 is "transport pathways into the lower stratosphere." The authors have illustrated that "eastward eddy shedding" on the NE side of the anticyclone, and subsequent transport and filamentation of material can be a pathway to reach the stratosphere (pathway 1, line 10). But, what about the "westward eddy shedding" (pathway 2, line 11) from the anticyclone to the TTL? I find no discussion of this as a potential pathway to the stratosphere, via e.g. diabatic ascent (Garny and Randel, 2015).

We agree that pathway 2 needs some further discussion and added therefore the following paragraph: 'Transport of air masses from the Asian monsoon anticyclone southeastwards into the tropics by e. g. westward eddy shedding causes increase of contributions of air masses originating in the boundary layer in India/China within the TTL. These air masses could penetrate into the upwelling in the deep branch of the Brewer-Dobson circulation and thus could be transported further up into the stratosphere (e. g. Garny and Randel, 2016).'

12. p.12, discussion of Fig. 14 and Table 2. Please clarify how these metrics are computed. Are they achieved by area weighting daily isentropic "fraction of air" maps for areas north of 30 o N and for PV greater than the "transport barrier" PV? Are you saying for example that by end October 2012, almost 20% of the air in the NH at 360K north of these delimiters originates in the India/China BL within the previous 5 months?

The reviewers interpretation of the percentages in Fig. 14. is correct. For clarification we revised the text as follows:

'The accumulation of young air masses from Asia since 1 May 2012 in the extratropical lower stratosphere is calculated using the isentropic transport barrier at different levels of potential temperature derived by Kunz et al. (2015) as shown in Fig. 14 and Tab. 2. To calculate the percentages of different emission tracers within the lower extratropical stratosphere at a certain level of potential temperature, we use the following approach: For each day between 1 May and 31 October 2012 a mean value for each emission tracers of all CLaMS air parcels is calculated for PV values larger (lower) than those at the transport barrier (see Tab. 2) and for air masses poleward of 30° N (30° S) at a specific isentropic level ($\Theta \pm 0.5$ K). End of October 2012, the contributions of all boundary emission tracers on the composition of the extratropical northern lower stratosphere are at 360 K \approx 44%, at 380 K \approx 35%, and at 400 K \approx 23%, with the remaining fraction of air consisting of aged air. The highest contributions of the boundary emission tracers are from India/China uplifted within the Asian monsoon anticyclone, from Southeast Asia, and from the tropical Pacific Ocean. The contribution of all other regions of the Earth's surface (residual surface) are of minor importance (Tab. 2).'

13. p.14, lines 12-14, reference "A mixing layer (see Fig. 5)" It is unclear to me what feature is being identified in Fig. 5. I see no discussion of such a mixing layer in earlier discussion of Fig. 5. Indeed 2 lines earlier (p.14, line 11) the thermal tropopause is described as a "strong transport barrier above the separated anticyclone." Do you really mean Fig. 5? Are you referring to the thin layer of strong vertical gradients in fractions of air and in simulated CO at the thermal tropopause south of 40 o N, 370-400 K What is the evidence for mixing, and what is the mechanism? Or do you mean Fig. 6 instead where mixed tropospherestratosphere characteristics of BL source fractions, CO, and buoyancy frequency are evident between the double thermal tropopauses? The discussion of PAN (p.14, lines 16-23) suggest you are discussing Fig. 5, but what I see is a strong vertical gradient at the tropopause, not a zone of mixed tropospheric and stratospheric characteristics. I read reference to a "small mixing layer around the tropopause" (p.14, line 33) which seems to minimize the significance of mixing here; if it's not significant (strong transport barrier), why spend a whole paragraph (lines 10-23) describing it?

We agree the discussion about the mixing layer is redundant. Therefore, we remove this paragraph within the discussion section as well as the statement "small mixing layer around the tropopause" (p.14, line 33) within the conclusions.

14. There are some places where the English is a bit obscure to me, e.g.

on p.15, line 15, I don't know what "yield predominantly" means in this context. I wonder if the words "yield to" (e.g. on p.15, line 23) is intended to mean "serves to" or "results in", i.e. "serves to increase," or "results in increasing."

The sentence is revised as follows: 'This second transport pathway is mainly caused by westward eddy shedding, however this transport pathway enhances predominantly the contribution of fresh emission from India/China to the composition of TTL air at 380 K in summer and autumn 2012.'

References

- Garny, H. and Randel, W. J.: Transport pathways from the Asian monsoon anticyclone to the stratosphere, Atmos. Chem. Phys., 16, 2703–2718, doi: 10.5194/acp-16-2703-2016, 2016.
- Kunz, A., Sprenger, M., and Wernli, H.: Climatology of potential vorticity streamers and associated isentropic transport pathways across PV gradient barriers, J. Geophys. Res., 120, 3802–3821, doi:10.1002/2014JD022615, URL http://dx.doi.org/10.1002/2014JD022615, 2014JD022615, 2015.
- Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and Engel, A.: Impact of the Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe 2012, Atmos. Chem. Phys., 16, 10573–10589, doi:10.5194/acp-16-10573-2016, 2016.
- Vogel, B., Pan, L. L., Konopka, P., Günther, G., Müller, R., Hall, W., Campos, T., Pollack, I., Weinheimer, A., Wei, J., Atlas, E. L., and Bowman, K. P.: Transport pathways and signatures of mixing in the extratropical tropopause region derived from Lagrangian model simulations, J. Geophys. Res., 116, D05306, doi:10.1029/2010JD014876, 2011.



Figure 2: (= Fig. 8 within the ACPD paper) Horizontal cross-section of the fraction of air originating in India/China over Europe on 25 September at 370 K (top) and on 23 September 2012 at 380 K (bottom). The flight paths transferred to noontime for the TACTS/EsmVal flights 2012 over northern Europe are marked as yellow lines. Segments of the flight in the lower stratosphere with enhanced measured CO, CH_4 , and H_2O and reduced O_3 compared to the stratospheric background are highlighted in red and numbered for clarification (the same numbers are used in Fig. 10 and 11). The climatological isentropic transport barrier of 6.0 PVU (at 370 K) and 7.2 PVU (at 380 K) are shown as thick black lines.



Figure 3: (= Fig. 10 within the ACPD paper) As Fig. 9, but for the flight on 25 September 2012.



Figure 4: (= Fig. 11 within the ACPD paper) As Fig. 9, but for the flight on 23 September 2012.

Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012

Bärbel Vogel¹, Gebhard Günther¹, Rolf Müller¹, Jens-Uwe Grooβ¹, Armin Afchine¹, Heiko Bozem², Peter Hoor², Martina Krämer¹, Stefan Müller², Martin Riese¹, Christian Rolf¹, Nicole Spelten¹, Gabriele P. Stiller³, Jörn Ungermann¹, and Andreas Zahn³

¹Forschungszentrum Jülich, Institute of Energy and Climate Research - Stratosphere (IEK-7), Jülich, Germany
 ²Institute for Atmospheric Physics, University of Mainz, Mainz, Germany
 ³Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

Correspondence to: Bärbel Vogel (b.vogel@fz-juelich.de)

Abstract. Global simulations with the Chemical Lagrangian Model of the Stratosphere (CLaMS) using artificial tracers of air mass origin are used to analyze transport pathways mechanisms from the Asian monsoon region into the lower stratosphere. In a case study, the transport of air masses from the Asian monsoon anticyclone originating in India/China by an eastward migrating anticyclone breaking off from the main anticyclone on 20 September 2012 and filaments separated at the north-

- 5 eastern flank of the anticyclone are analyzed. Enhanced contributions of young air masses (younger than 5 months) are found within the separated anticyclone confined at the top by the thermal tropopause. Further, these air masses are confined by the anticyclonic circulation and at the polar side by the subtropical jet such as that the vertical structure looks appears like a bubble within the upper troposphere. Subsequently, these air masses are transported eastwards along the subtropical jet and enter the lower stratosphere by quasi-horizontal transport in a region of double tropopauses most likely associated with Rossby wave
- 10 breaking events. As a result, thin filaments with enhanced signatures of tropospheric trace gases are measured in the lower stratosphere over Europe during the TACTS/ESMVal campaign in September 2012 in very good agreement with CLaMS simulations. Our simulations demonstrate that source regions in Asia and in the Pacific Ocean have a significant impact on the chemical composition of the lower stratosphere of the Northern Hemisphereby flooding the extratropical lower stratosphere with young. Young moist air masses, in particular at the end of the monsoon season in September/October 2012(, are flooding)
- 15 the extratropical lower stratosphere in the Northern Hemisphere with contributions up to $\approx 30\%$ at 380 K) in contrast (with the remaining fraction being aged air). In contrast, the contribution of young air masses to the southern hemisphere is much lower. End of October 2012, approximately 1.5 ppmv H₂O is found in the lower northern hemisphere stratosphere (at 380 K) from source regions both in Asia and in the tropical Pacific compared to a mean water vapor content of ≈ 5 ppmv. In addition to this main transport pathway from the Asian monsoon anticyclone to the east along the subtropical jet and subsequent transport into
- 20 the northern lower stratosphere, a second horizontal transport pathway out of the anticyclone to the west into the tropics (TTL) is found in agreement with MIPAS HCFC-22 measurements.

1 Introduction

15

The Asian summer monsoon is associated with strong upward transport of tropospheric source gases by deep convection. Uplifted tropospheric air masses are confined by a strong anticyclonic circulation in the upper troposphere which is acting acts as a transport barrier (e. g. Li et al., 2005; Randel and Park, 2006; Park et al., 2007, 2008, 2009; Ploeger et al., 2015). In summer

- 5 2012, enhanced tropospheric trace gases are found in the upper troposphere over Asia from mid-June until late October both during the existence of the Asian monsoon anticyclone and after its breakup (Vogel et al., 2015). Studies for the summer 2012 show that air masses from the Asian monsoon impact the chemical composition of the lower stratosphere over northern Europe, in particular at the end of the monsoon season in September and October 2012 (Vogel et al., 2014, 2015) (Müller et al., 2015). (Müller et al., 2016). An increase in the concentrations of tropospheric source gases such as CO, H₂O, and N₂O and a decrease
- 10 of the stratospheric tracer O_3 was identified in in-situ aircraft measurements in the lower stratosphere over Europe from summer to autumn 2012 associated with transport from the Asian monsoon anticyclone (Müller et al., 2015) (Müller et al., 2016).

The identification and the relative importance of different transport pathways of tropospheric source gases found within the Asian monsoon anticyclone into the lower stratosphere are subject of a longstanding debate (Dethof et al., 1999; Park et al., 2004; Randel et al., 2010; Bian et al., 2012; Bourassa et al., 2012; Ploeger et al., 2013; Fairlie et al., 2014; Fromm

- et al., 2014; Uma et al., 2014; Vogel et al., 2014, 2015; Garny and Randel, 2016; Tissier and Legras, 2016; Orbe et al., 2015). The influence of these different transport pathways on the chemical composition of the extratropical upper troposphere and lower stratosphere (ExUTLS) is important because changes in ozone and water vapor in the ExUTLS have a significant impact on surface climate, even if the perturbation is relatively small. Thus changes in the chemical composition in that part of the
- Earth's atmosphere play a crucial role for climate change (e. g. Solomon et al., 2010; Riese et al., 2012; Hossaini et al., 2015). Moreover, increasing stratospheric water vapor has the potential to affect stratospheric chemistry (e. g. Kirk-Davidoff et al., 1999; Dvortsov and Solomon, 2001; Vogel et al., 2011a; Anderson et al., 2012) and can influence the formation of cirrus clouds in the lower stratosphere (Spang et al., 2015). Further, enhanced water vapor in combination with pollution in the Asian monsoon region is discussed to play an important role in the formation of the Asian Tropopause Aerosol Layer (ATAL) which
 causes a regional radiative forcing (Vernier et al., 2015).

One important mechanism for long-range transport of air masses from the Asian monsoon anticyclone to the extratropical lower stratosphere is the separation of air masses at the northeastern flank of the anticyclone caused by disturbances of the subtropical jet by strong Rossby waves and subsequent eastward transport of these air masses within the tropics along the subtropical jet (Dethof et al., 1999; Vogel et al., 2014, 2015). In particular, eastward migrating anticyclones breaking off

30 from the main anticyclone a few times each summer (Dethof et al., 1999; Hsu and Plumb, 2001; Popovic and Plumb, 2001; Garny and Randel, 2013; Vogel et al., 2014; Ungermann et al., 2016) have the potential to carry air with high amounts of tropospheric trace gases from the Asian monsoon anticyclone to mid- and high latitudes of the Northern Hemisphere. These young air masses can be transported into the extratropical lower stratosphere where they are eventually mixed irreversibly with the surrounding aged stratospheric air (Dethof et al., 1999; Garny and Randel, 2013; Vogel et al., 2014) thus affecting the

chemical and radiative balance of the extratropical lower stratosphere. However, the exact transport mechanisms of young air masses from the troposphere including air originating within the Asian monsoon anticyclone into the lower stratosphere are yet to be identified and quantified.

- Trajectory calculations show that tropospheric air masses at the southeast edge of the anticyclonic circulation over Southeast 5 Asia may be uplifted to the outer anticyclonic circulation by a typhoon to levels of potential temperature of \approx 370 K (Vogel et al., 2014). Subsequent upward transport of these air masses occurs in a clockwise upward spiral around the core of the Asian monsoon anticyclone to levels of potential temperature around 380 K. Moreover, Vogel et al. (2014) demonstrate that the combination of very rapid uplift by a typhoon and eastward eddy shedding from the Asian monsoon anticyclone is an additional rapid transport pathway (\approx 5 weeks) connecting surface air with enhanced signatures of tropospheric trace gases
- 10 measured in the lower stratosphere over northern Europe.

Here, we use in-situ measurements obtained during two aircraft campaigns TACTS and ESMVal, jointly conducted in August and September 2012, using the German High Altitude and LOng Range Research Aircraft (HALO). Backward trajectory calculations (Vogel et al., 2014) (Müller et al., 2015) (Müller et al., 2016) show that enhanced tropospheric trace gases that were measured over northern Europe in the extratropical lower stratosphere during the TACTS/ESMVal campaign during

15 August and September 2012 are affected by air masses from the circulation of the Asian monsoon anticyclone or originating in the Asian monsoon anticyclone itself.

Further model simulations with the Chemical Lagrangian model of the stratosphere (Vogel et al., 2015) Model of the Stratosphere using artificial emission tracers for different regions on the Earth's surface demonstrate that emissions of young (Vogel et al., 2015) demonstrate that the chemical composition of the lower stratosphere of the Northern Hemisphere is influenced

- 20 by a combination of young, freshly injected air and aged air. The composition of the young air masses is dominated by air masses primarily from originating in India, China, and Southeast Asiainfluence the chemical composition of the lower stratosphere of the Northern Hemisphere, in particular at the end of the monsoon season in September/October 2012. Thus the Asian monsoon anticyclone and boundary emissions from Asia contribute to the maximum of tropospheric signatures found in the northern extratropical lower stratosphere in boreal summer and autumn (e. g. Hoor et al., 2005; Hegglin and Shep-
- 25 herd, 2007; Bönisch et al., 2009; Zahn et al., 2014) (Müller et al., 2015) (Müller et al., 2016), sometimes also referred to as 'flushing' of the lower stratosphere (Hegglin and Shepherd, 2007; Bönisch et al., 2009).

Exchange of air masses from the troposphere to the stratosphere occurs preferably in poleward flow structures such as tropospheric intrusions (e. g. Fairlie et al., 2007; Sprenger et al., 2007; Pan et al., 2009; Vogel et al., 2011b). These intrusions develop into elongated potential vorticity (PV) streamers and are manifestation of Rossby wave breaking. Rossby wave

30 breaking is identified as an important mechanism for exchange of air masses between the tropical upper troposphere and the extratropical lower stratosphere with a pronounced peak during summer in each hemisphere controlled by the presence of monsoon anticyclones, in particular of the Asian monsoon (e. g. Homeyer and Bowman, 2013; Kunz et al., 2015). High frequency of PV streamers are found over the eastern North Pacific and over the Atlantic in summer demonstrated in a recently published climatology of PV streamers (Kunz et al., 2015). In addition to this large-scale transport process of water vapor by air mass exchange between the tropics and the extratropics, convection induced injections of water vapor in mid-latitudes can also occur in large storm systems such as tropical cyclones and by deep continental convection (e. g. Anderson et al., 2012; Homeyer et al., 2014; Vogel et al., 2014).

In this paper, we identify in detail the transport mechanisms of air masses originating in boundary source regions in Asia and separated from the Asian monsoon anticyclone by eastward eddy shedding into the lower stratosphere over northern Europe including mixing processes. The same model simulation as in Vogel et al. (2015) with the three-dimensional Lagrangian chemistry transport model (CLaMS) (e. g. Pommrich et al., 2014, and references therein) are is used including artificial emission tracers that mark source regions in the boundary layer of the Earth's atmosphere. This allows the origin of air masses and the detailed transport pathways from Asian source regions into the northern lower stratosphere to be quantified. Further,

- 10 the irreversible part of transport, i.e. mixing, in CLaMS is controlled by the local horizontal strain and vertical shear rates with mixing parameters deduced from observations (Konopka et al., 2012, and references therein) and therefore allows mixing processes in the lower stratosphere as shown in several previous studies to be characterized (e.g. Pan et al., 2006; Vogel et al., 2011b; Konopka and Pan, 2012). Small scale structures indicating the impact of young tropospheric air masses within the lower stratosphere measured during the TACTS/ESMVal aircraft campaign in September 2012 are compared with results of CLaMS
- 15 model simulations to identify their origin and the associated transport pathways. Finally, the impact of boundary source regions in Asia on the composition of the extratropical Northern Hemisphere is calculated.

The paper is organized as follows: Section 2 describes the in situ measurements and Section 3 the setup for used CLaMS simulations. In Section 4, CLaMS results are presented and compared to measurements. A short discussion and conclusions are given in Sections 5 and 6.5.

20

2 TACTS/ESMVal Measurements over Europe

Here we use in-situ measurements performed during two aircraft campaigns TACTS and ESMVal, jointly conducted in August and September 2012. TACTS was designed to study 'Transport and Composition in the Upper Troposphere and Lowermost Stratosphere'. The objective of the ESMVal (Earth System Model Validation) measurement campaign was to validate model
simulations with measurements ranging from the Southern to the Northern Hemisphere (here from 65°S to 80°N). Both campaigns were performed using the German High Altitude and LOng Range Research Aircraft (HALO), a Gulfstream V. During the TACTS/ESMVal campaign 13 research flights were performed, the most of them over northern Europe. Here, we discuss the last three flights conducted on 23, 25, and 26 September 2012. Measurements of from the following in-situ instruments on board the HALO aircraft are used:

- Carbon monoxide (CO) and methane (CH₄) measurements are used from TRIHOP a three channel Quantum Cascade Laser Infrared Absorption spectrometer (for more details see Müller et al., 2015).
 (for more details see Müller et al., 2016) which is an updated version of the three-channel tunable diode laser instrument for atmospheric research (TRISTAR) used in previous aircraft campaigns (e.g. Hoor et al., 2004).
- Water vapor (H₂O) measurements were obtained from the Fast In-situ Stratospheric Hygrometer (FISH) which is based on the Lyman- α photofragment fluorescence technique (Zöger et al., 1999). The FISH inlet was mounted forward-facing to measure total water which is the sum of gas-phase water and water in ice particles. To calculate gas-phase water from FISH measurements in clouds a correction procedure is applied (for more details see Meyer et al., 2015).
- 5
- Ozone (O₃) was measured with FAIRO a light-weight (14.5 kg) instrument with high accuracy (2%) and high time resolution (10 Hz) developed for the HALO aircraft. FAIRO combines dual-beam UV photometer with an UV-LED as light source and a dry chemiluminescence detector (Zahn et al., 2012).
- Potential temperature (Θ) was deduced from the Basic HALO Measurement and Sensor System (BAHAMAS) that yields
- 10

15

basic meteorological and avionic data of all TACTS/ESMVal flights.

3 CLaMS simulations using artificial tracers of air mass origin

We use the same model simulation with the Chemical Lagrangian model of the stratosphere Model of the Stratosphere (CLaMS) (McKenna et al., 2002b, a; Pommrich et al., 2014, and references therein) as in Vogel et al. (2015) covering the Asian monsoon season 2012. The CLaMS model has the capability to reproduce strong gradients of chemical species found in regions with strong transport barriers such as the edge of the Asian monsoon anticyclone (e.g. Konopka et al., 2010; Ploeger et al., 2015; Vogel et al., 2015), the extratropical tropopause (e.g. Pan et al., 2006; Vogel et al., 2011b; Konopka and Pan, 2012) and the polar vortex (e.g. Günther et al., 2008; Vogel et al., 2008).

In brief, the global CLaMS simulation employed here covers an altitude range from the surface up to 900 K potential temperature ($\approx 37 \text{ km}$ altitude) with a horizontal resolution of 100 km and a maximum vertical resolution of approximately

- 20 400 m near the tropopause. The model simulation is driven by horizontal winds from ERA-Interim reanalysis (Dee et al., 2011) provided by the European Centre for Medium-Range Weather Forecasts (ECMWF). For the vertical velocities, the diabatic approach was applied using the diabatic heating rate as vertical velocity including latent heat release (for more details see Ploeger et al., 2010). Further, CLaMS employs a hybrid coordinate (ζ), which transforms from a strictly isentropic coordinate Θ to a pressure-based coordinate system below a certain reference level (in this paper study 300 hPa) (for more details see Nonenka et al. 2012; Pommrich et al. 2014).
- 25 Konopka et al., 2012; Pommrich et al., 2014).

The CLaMS simulation covers the time period from 1 May 2012 until 31 October 2012 including the Asian monsoon season 2012. It includes full stratospheric chemistry (Grooß et al., 2014; Sander et al., 2011). Chemical species are initialized on 1 May 2012 mainly based on satellite data from AURA-MLS (version 3.3) (Livesey et al., 2011) and ACE-FTS (version 3.0) (Waymark et al., 2013) measurements. At the upper boundary (900 K potential temperature), mainly AURA-MLS and

30 ACE-FTS measurements and tracer-tracer correlations were used (for more details see Vogel et al., 2015).

At the lower boundary (surface), O_3 is set to a constant tropospheric volume mixing ratio of 4.8×10^{-8} volume mixing ratio representing the ozone mixing ratio at 5 km (Brasseur and Solomon, 2005, p. 619). ECMWF water vapor is prescribed

at lower model levels. Lower boundary conditions for CO and CH_4 are derived from AIRS (Atmospheric Infrared Sounder) version 6 satellite measurements following the approach described by Pommrich et al. (2014).

In the CLaMS simulation used here, artificial tracers of air mass origin, referred to as "emission tracers", are implemented that mark globally all regions in the Earth's boundary layer ($\approx 2-3$ km above the surface following orography corresponding to $\zeta < 120$ K) as shown in Fig. 1 and Tab. 1. Within the boundary layer, the sum of all different emission tracers (Ω_i) is equal to 1 ($\Omega = \sum_{i=1}^{n} \Omega_i = 1$). Air masses in the model boundary layer are marked by different emission tracers every 24 h (the time step for mixing in CLaMS). Thus in our simulations, the transport of air masses from the model boundary layer up to the UTLS is considered from the response to a single pulse during ($t_i, t_i + \Delta t_i$) for times $t > t_i$, with t_i equal to 1 May 2012 and Δt_i equal

- 10 to 6 months marking young, freshly emitted air masses. We used this single pulse approach because the focus of the paper is to analyze the influence of fresh emissions from different boundary regions using the meteorological conditions of the year 2012. Transport and mixing of the emission tracers to other regions of the free troposphere or stratosphere occurs like for in an equivalent way to all chemical species included in the CLaMS simulation. The percentage of an individual emission tracer counts the contribution of the corresponding boundary layer region to the composition of an air parcel considering advection
- 15 and mixing since 1 May 2012. Thus, during the course of the simulation, the air-mass fraction $f_i(r, t \mid \Omega_i)$ is defined to be the fraction of air at a location r and time t (with $t_i \le t \le t_i + \Delta t_i$) that was emitted since t_i within Δt_i from the model boundary layer in region Ω_i . Note however that our emission tracers only describe the contribution of young freshly injected air masses from the boundary layer. Therefore, in our simulations presented here, the composition of an air mass in the free atmosphere will be a combination of 'young air masses' and 'aged air masses' originating in the free troposphere or stratosphere.
- 20 Emission tracers in CLaMS are designed to identify surface regions of the Earth's atmosphere that contribute to the composition of the Asian monsoon anticyclone and of the lower stratosphere during the course of the 2012 Asian monsoon season with a focus on the influence of fresh emissions. Therefore, the starting point for the CLaMS simulation is chosen a few weeks before the formation of the Asian monsoon anticyclone occurred. Thus, transport processes associated with the Asian monsoon anticyclone and horizontal transport from the TTL into the lower stratosphere are considered.
- Vogel et al. (2015) show that at the end of the monsoon season in September up to 75 % of the air masses within the Asian monsoon anticyclone are younger than 5 months. In addition, they show that the emission tracer for India/China (Northern India + Southern India + Eastern China) is a good proxy for the location and shape of Asian monsoon anticyclone using pattern correlations with potential vorticity (PV), and MLS O₃ and CO measurements. Therefore, the emission tracer for India/China is very well suited to analyze transport pathways of air masses from the Asian monsoon anticyclone into the northern lower
- 30 stratosphere. The emission tracer for Southeast Asia contributes also to the composition of the Asian monsoon but is in addition found in air masses circulating around the outer edge of the Asian monsoon anticyclone at levels of potential temperature of around 380 K.

The most important regions for our study are India/China (= Northern India (NIN) + Southern India (SIN) + Eastern China (ECH)), Southeast Asia (SEA) and the tropical Pacific Ocean (TPO). The sum of all model boundary layer tracers

 $(\Omega = \sum_{i=1}^{n} \Omega_i)$ without contributions from India/China, Southeast Asia, and the tropical Pacific Ocean is summarized in one emission tracer referred to as "the residual surface" (= Ω - NIN - SIN - ECH - SEA - TPO).

4 Results

4.1 Eastward eddy shedding and separation of filaments

5 We will analyze transport mechanisms and pathways of tropospheric air masses separated by eastward eddy shedding or filaments at the northeastern flank of the Asian monsoon anticyclone into the lower stratosphere over Europe for the 2012 monsoon season.

At the northeastern flank of the Asian monsoon anticyclone filaments with low PV and enhanced percentages of the emission tracer for India/China are frequently separated from the main anticyclone in 2012, e.g. on 4 and 5 September 2012 as shown

- 10 in Fig. 2. In addition, eastward migrating anticyclones break off from the main anticyclone several times during summer. A pronounced eddy shedding event occurred on 20 September 2012 (Vogel et al., 2014). A second anticyclone (130°E-210°E) characterized by low PV separated from the main Asian monsoon anticyclone (10°E-110°E) on 20 September 2012 as shown in Fig. 3 (top left). Thereafter, transport of these air masses characterized by low PV to the Pacific Ocean occurred (see Fig. 3, left) at 380 K.
- Fig. 3 (right column) shows that air masses with enhanced contributions of the emission tracers for India/China and originating from the Asian monsoon anticyclone are transported eastwards within the separated anticyclone. After 20 September 2012, a long filament of air characterized by low PV and enhanced percentages of emission tracers for India/China evolves from the separated anticyclone. On 23 September 2012, this filament is located over North America along a poleward excursion of the subtropical westerly jet. Between 24 and 26 September 2012, these air masses are transported further eastwards over North
- 20 America and afterwards to the Atlantic Ocean.

A very similar horizontal distribution as for the emission tracer for India/China is also found for simulated CO (see Fig. 4, left). However, enhanced CO values are in addition found at 380 K in the tropics associated with deep uplift in the tropics outside of the monsoon region to the upper troposphere in particular over the Maritime Continent (the region between Indian and Pacific Oceans) and the western Pacific (see emission tracer for Southeast Asia and tropical Pacific Ocean in Appendix A

- 25 to this paper). The Further, the emission tracer for Southeast Asia / tropical Pacific Ocean contributes also to the composition of the Asian monsoon anticyclone, however to a smaller extent compared to the emission tracer for India/China as shown in Fig. 4 (right). Contributions from the residual surface are of minor importance (see Sect. 4.2). Note that contribution of air masses originating in the free troposphere and stratosphere ('aged air masses') also contribute to the composition of the Asian monsoon anticyclone about 25 % at the end of September 2012 (see in Vogel et al., 2015, Fig. 9).
- 30 Maximum percentages for the emission tracer for Southeast Asia / tropical Pacific Ocean are found at the edge of the Asian monsoon anticyclone indicating the transport of these air masses around the outer edge of the Asian monsoon anticyclone outside of a PV-based transport barrier (Ploeger et al., 2015) as discussed by Vogel et al. (2015). Thus air masses from Southeast Asia / tropical Pacific Ocean are found within a widespread area around the anticyclone caused by the large-scale anticyclonic

flow in this region acting as a large-scale stirrer. Moreover, high contributions of these emission tracers are found in the tropics associated with deep uplift out side of the monsoon region.

Kunz et al. (2015) derived a climatology of PV streamers (e.g. polward moving filaments with low PV originating in the tropics) as indicators of Rossby wave breaking on isentropes between 320 and 500 K using ERA-Interim reanalyses for the

- 5 time period from 1979 to 2011. The 7.2 PVU isoline shown in black in Fig. 3 and Fig. 4 represents the isentropic transport barrier between the tropics and mid-latitudes based on the climatology of the dynamically relevant PV contours derived by Kunz et al. (2015) at 380 K for the Northern Hemisphere for September until November. Isentropic transport of air masses across the 7.2 PVU isoline indicates irreversible exchange between the tropics and extratropics due to wave breaking. On 20 September 2012, at the northern flank of the separated anticyclone the 7.2 PVU isoline envelops the region of enhanced tracers
- 10 is in the region where strongest gradients of the emission tracers occur indicating the transport barrier at 380 K (see Fig. 3 and 4).

Considering altitude-height cross-sections from the separated anticyclone caused by eddy shedding at $180^{\circ}E$ (Fig. 5) shows that air masses within the separated anticyclone ($20^{\circ}N - 40^{\circ}N$) throughout the troposphere are characterized by high percentages of the emission tracers for India/China (top left). The vertical tracer distribution of the separated anticyclone looks like a

- 15 bubble that is confined by the subtropical westerly jet in the north (35°N-55°N) and by the thermal tropopause at the top. The thermal tropopause is slightly elevated similar as for to the Asian monsoon anticyclone itself. Here, the thermal tropopause acts as a transport barrier for further upward transport from the separated anticyclone into the lower stratosphere. The terminology 'tropospheric bubble' is also used to describe the trace gas distribution within the Asian monsoon itself (Pan et al., 2016).
- In addition, the isentropic transport barriers of 4 and 10 PVU at 350 K and 400 K (Kunz et al., 2015), respectively, are shown in Fig. 5 (white thick lines) for the Northern Hemisphere for September until November. No significant transport across these PV isolines at the polar edge of the separated anticyclone was found indicating that here no air mass exchange between stratosphere and troposphere occurred due to Rossby wave breaking.

The altitude-height cross-section for the emission tracer for Southeast Asia / tropical Pacific Ocean (Fig. 5, top right) has also the structure of a bubble with enhanced percentages. These enhanced contributions of the emission tracer for Southeast

Asia / tropical Pacific Ocean are clearly separated from the tropics, outside of the monsoon circulation, which shows even higher contributions (up to 40%) of these tracers.

Fig. 5 (bottom left) shows the emission tracer for the residual surface that is the sum of all other emission tracers (all tracers without (the entire model boundary layer without contributions from India/Chinaand Southeast Asia/, Southeast Asia, and the tropical Pacific Ocean, see Sect. 3). The contribution of the emission tracer for the residual surface shows no signature is

30 approximately 10-15 % within the separated anticyclone indicating that air masses originating in India/China and Southeast

Asia / tropical Pacific Ocean almost exclusively contribute to the chemical composition of the separated anticyclone.

In general, the troposphere is characterized by weak static stability (buoyancy frequency squared, N^2) in contrast to stratospheric air masses. The altitude-height cross-section of N^2 (Fig. 5, bottom right) shows that air masses within the separated anticyclone are characterized by tropospheric N^2 values. Our model simulation demonstrates that air masses enclosed in the

35 second anticyclone are still located in the troposphere. The question arises where exactly air masses from the Asian monsoon

anticyclone separated by eddy shedding or large filaments will enter the lower stratosphere i. e. where they exactly cross the extratropical tropopause.

4.2 Isentropic transport pathways into the lower stratosphere

- 5 As shown in Figs. 3 and 4 (middle) a long filament of air characterized by low PV and high percentages of emission tracers for India/China and Southeast Asia / tropical Pacific Ocean evolves from the separated anticyclone. On 23 September 2012, this filament is located over North America along the subtropical westerly jet. At 380 K enhanced contributions of the emission tracer from India/China are also found north of the 7.2 PVU barrier indicating transport from the troposphere into stratospherethe stratosphere according to the definition by Kunz et al. (2015).
- 10 An altitude-height cross-sections at 120°W on 23 September 2012 cutting the filament over northern America is shown in Fig. 6. A double thermal tropopause (30°N-65°N) is found along the filament (see black dots in Fig. 6) which encloses a tropospheric intrusion centered around 370 K with a vertical extension of up to 50 K. Air masses associated with a tropospheric intrusion, namely tropospheric air masses intruding polward into the lower stratosphere, are characterized by lower values of ozone, PV, and N² and higher values of water vapor and CO compared to the stratospheric background (Pan et al., 2009)
- 15 (Homeyer et al., 2011) (Vogel et al., 2011b; Ploeger et al., 2013). Indeed, the air mass in the tropospheric intrusion shows low N² value (Fig. 6, middle right) and enhanced simulated CO (Fig. 6, bottom). For illustration, the 4 PVU and 10 PVU isolines are shown as thick white line in Fig. 6 indicating the climatological isentropic transport barrier at 350 K and 400 K, respectively, for the Northern Hemisphere for September (Kunz et al., 2015).

Enhanced percentages of the emission tracers for India/China (up to $\approx 30\%$) and Southeast Asia / tropical Pacific Ocean

- 20 (up to $\approx 33\%$) are found between the double tropopauses (see Fig. 6, top) compared to the stratospheric background. This demonstrates the horizontal isentropic transport of young tropospheric air masses from the filament into the lower stratosphere. The impact of the emission tracer of the residual surface is less than 10% indicating a minor influence from young air masses originating in other regions of the Earth's surface on the tropospheric intrusion (see Fig. 6 middle left). Thus the transport of tropospheric air masses from the troposphere into the lower stratosphere occurs in the region between the double tropopauses.
- 25 Previous studies also found intrusions of tropospheric air into the lower stratosphere associated with extratropical double tropopauses (e. g. Pan et al., 2009; Homeyer et al., 2011; Vogel et al., 2011b; Schwartz et al., 2015; Wu and Lü, 2015). Further, it was shown that double tropopauses are frequently associated with Rossby wave breaking events along the subtropical jet (e. g. Vaughan and Timmis, 1998; Castanheira and Gimeno, 2011; Ungermann et al., 2013)

Homeyer and Bowman 2012 (Homeyer and Bowman, 2013) (Homeyer et al., 2014).

30

Consequently thin filaments with enhanced contributions of emission tracers for India/China (also for Southeast Asia / tropical Pacific Ocean but not shown here) are found at the polar side of the 7.2 PVU isoline representing the climatological isentropic transport barrier at 380 K in September and October 2012 (see Fig. 7, top, thick black line). These small filaments in the lower stratosphere with enhanced percentages of emission tracers for India/China reach the flight path (red line) of the

TACTS/ESMVal flight on 26 September 2012 over northern Europe. An altitude-height cross-sections at 8°W on 26 September 2012 (marked as thick white line in Fig. 7, top) cross a filament with enhanced contributions of the emission tracers for India/China at 380 K between 60°N and 70°N (Fig. 7, bottom). This filament is clearly found above the first thermal tropopause (black dots) within the lower stratosphere associated with a double tropopause.

5

Similar filaments within the lower stratosphere were also measured at the flight path of the TACTS/ESMVal flights on 23 and 25 September 2012 (Fig. 8). Subsequently, these filaments are mixed with the stratospheric background and dissipate in the lower stratosphere over time.

4.3 Comparison to TACTS/ESMVal measurements

Above, in Sec. 4.1 and 4.2 we considered CLaMS simulations using artificial emission tracers to show that air masses orig inating in Asia and in the tropical Pacific affect the chemical composition of the lower stratosphere over northern Europe in September 2012. In this section, we compare results of the three-dimensional CLaMS simulation of CO, O₃, CH₄, and H₂O with in-situ measurements of three flights conducted on 23, 25, and 26 September 2012 over Europe during the TACTS/ESMVal campaign. The CLaMS results are compared with the in-situ measurements along the flight path by interpolating in time and space. The interpolation method is described in detail in Appendix B.

- In Fig. 9 11, CLaMS results interpolated along the flight path for each of the three flights are presented. Also shown are the contributions of CLaMS emission tracers for India/China, Southeast Asia / tropical Pacific Ocean, and the residual surface interpolated along the flight path (see Figs. 9 11, top). Further, measured CO, O₃, CH₄, and H₂O mixing ratios are compared with the CLaMS results. The parts of the flights conducted in the lower stratosphere and characterized by enhanced values of measured CO, CH₄, and H₂O and simultaneously reduced O₃ compared to the stratospheric background are highlighted in
- 20 gray and are numbered for each flight. The same flight segments are shown in red in Figs. 7 and 8 and are also labeled by numbers.

In general, during all three flights, air masses in the lower stratosphere over Europe are affected by boundary emissions from India/China and Southeast Asia / tropical Pacific Ocean. Contributions up to $\approx \frac{2320\%}{2320\%}$ for emission tracers for India/China and Southeast Asia / tropical Pacific Ocean are found. In contrast, the contribution of the emission tracers for the

25 residual surface are below 7% in the lower stratosphere. Higher contributions are only found in flight parts conducted in the troposphere, e. g. during take off, landing, or flight pattern with a steep decent down into the troposphere followed by a steep ascent back into the lower stratosphere, referred to as "dive" (see Fig. 9 (top) shortly after 13.00 UTC and Fig. 11 (top) around 10:00 UTC).

In regions with measured enhanced values of measured CO, CH_4 , and H_2O and reduced O_3 , generally also the contributions of the emission tracers for India/China and Southeast Asia / tropical Pacific Ocean simulated with CLaMS are enhanced (marked in gray). Further, a good overall agreement between measurements of CO, H_2O , O_3 and CLaMS simulations is found. The simulated values of CH_4 are in general lower than the measurements (up to ≈ 50 ppbv), however the simulated CH_4 signatures correspond to the measurements. Differences between model and measurements are most likely caused by uncertainties in the lower boundary conditions of CLaMS (see Sec. 3) which do not include individual emission events of CO or CH_4 (see Pommrich et al., 2014).

In the following, we discuss individual signatures of tropospheric air notable in the measurements. These signatures are marked in gray and for clarification, are numbered for each flight (see Fig. 9 - 11, top).

- 5 On 26 September 2012 (see Fig. 9, top), a very pronounced signature of tropospheric air in the lower stratosphere is found between 09:05 UTC to 10:17 UTC (No. 2). Here, the contributions of the emission tracer for India/China and Southeast Asia /tropical pacific Pacific Ocean are up to 20 % and 23 %, respectively -(up to 5 % from the residual surface). Thus, the sum of all emission tracers for model boundary layer is roughly 48 %. The remaining 52 % of the composition of the lower stratosphere in this region is from aged air masses originating in the free troposphere and the stratosphere at the beginning our the CLaMS
- 10 <u>simulation on 1 May 2012.</u>

This is consistent with backward trajectories calculations reported by Vogel et al. (2014) showing that these air masses are affected by the Asian monsoon anticyclone. Some of the trajectories (2%) originate in the West Pacific and experienced very rapid uplift in typhoon Bolaven on 24/25 August 2012. These air masses reach uplifted by typhoon Bolaven are transported from the Earth's surface over the West Pacific within 5 weeks to the lower stratosphere over Europe (Vogel et al., 2014).

- 15 Further, on 26 September 2012 (see Fig. 9, top), a second pronounced signature of tropospheric air (measured CO, O₃, CH₄, and H₂O values are of similar magnitude as for No. 2) is found in the lower stratosphere between 08:05 UTC to 10:23 UTC (No. 1). Also here, the contribution of the emission tracers for India/China and Southeast Asia / Tropical Pacific Ocean are enhanced by up to 10 % and 15 %, respectively. 60-day backward trajectories calculated by Vogel et al. (2014) show that some of these trajectories originate also in the West Pacific region in typhoon Haikui on 2/3 August 2012, however these trajectories
- 20 need a longer time (\approx 8 weeks) from their origin in the West Pacific to reach northern Europe.
 - During the second part of the flight on 26 September 2012 (see Fig. 9, top), further signatures of tropospheric air are found measured during the flight (No. 3-6), with . Also here, enhanced percentages of the emission tracers for India/China and Southeast Asia / tropical Pacific Ocean up to 18 % are simulated, except for region No. 3, that marks a tropospheric signature only found in the measurements.
- 25 During the flights on 25 and 23 September 2012 (see Figs. 10 and 11) also several lower tropospheric signatures are found in the lower stratosphere. Also here, measured tropospheric signals agree in general with results of the CLaMS model. The results of all these flights confirm that air masses with enhanced amounts of tropospheric trace gases measured in the lower stratosphere over northern Europe originating in boundary source regions in India/China and Southeast Asia / tropical Pacific Ocean. Our simulations in agreement with measurements show that the amount of water vapor and pollution in the lower strato-
- 30 sphere is enhanced in the Northern Hemisphere in September 2012 associated with both the dynamic of the Asian monsoon anticyclone and the transport of air masses from both Asia and the tropical Pacific along the subtropical jet.

4.4 Impact from Asian boundary source regions on extratropical lower stratosphere

4.4.1 Transport pathways into the lower stratosphere

In the previous sections, the long-range transport pathway from the Asian monsoon anticyclone into the extratropical lower Northern Hemisphere northern extratropical lower stratosphere was discussed using a case study. Here, the impact of this

- 5 horizontal transport on the composition of the extratropical lower stratosphere is calculated. Fig. 12 shows the horizontal distribution at 380 K of mean values of the emission tracer for India/China for July, August, and September 2012. The Figs. 10 and 11 show that also air masses originating in Southeast Asia and in the tropical Pacific contribute to the composition of the lower stratosphere over Northern Europe. The horizontal distribution of the emission tracers of Southeast Asia and the tropical Pacific at 380 K from July until September 2012 are shown and discussed in Appendix A to this paper.
- 10 In Fig. 12, the temporal evolution of the long-range transport pathway from the region of the Asian monsoon anticvclone into the extratropical lower stratosphere is evident. The frequent separation of air masses at the northeast flank of the Asian monsoon anticyclone and subsequent eastward transport along the subtropical jet is the most important mechanism for longrange transport into the lower northern stratosphere. This long-range transport pathway is most pronounced in September 2012 when the strong eddy shedding event took place as discussed in Sec. 4.1. In Summer 2012, two main horizontal transport
- pathways of air masses from the Asian monsoon anticyclone with high amounts of the emission tracer for India/China into the 15 TTL evolve (see Fig. 12d):
 - 1. northeastwards along the subtropical jet (eastward eddy shedding and separations of filaments) and the subsequent transport most likely by Rossby wave breaking events into the lower northern hemispheric stratosphere and
 - 2. southwestwards into the tropics (westward eddy shedding) and subsequent mixing within the TTL.
- 20 Transport from the Asian monsoon anticyclone both to the east and to the west yields an increase of fresh emissions from India/China within the TTL at 380 K in summer and autumn 2012. Finally, Fig. 12 shows that transport from air masses originating in the Asian monsoon anticyclone into the lower stratosphere take place most likely by Rossby wave breaking frequently over the Pacific and Atlantic Ocean (see Fig. 12c) which is in agreement with findings by Kunz et al. (2015). In our simulation, the horizontal long-range transport pathway along the subtropical jet is furthermore found at levels of potential 25
- temperature between 340 K and 420 K (not shown here).

Figs. 10 and 11 show that also Transport of air masses from the Asian monsoon anticyclone southeastwards into the tropics by e. g. westward eddy shedding causes increase of contributions of air masses originating in Southeast Asia and in the tropical Pacific contribute to the composition of the lower stratosphere over Northern Europe. The horizontal distribution of the emission tracers of Southeast Asia and the tropical Pacific at 380K from July until September 2012 are shown and discussed

30 in Appendix A to this paper, the boundary layer in India/China within the TTL. These air masses could penetrate into the upwelling in the deep branch of the Brewer-Dobson circulation and thus could be transported further up into the stratosphere (e. g. Garny and Randel, 2016).

To demonstrate that these both transport pathways are also evident in observations, we analyze global HCFC-22 measurements of the MIPAS instrument onboard the ENVISAT satellite (Chirkov et al., 2016). The production of the ozone-depleting and greenhouse gas HCFC-22 ($CHClF_2$) is regulated by the Montreal Protocol and its amendments and adjustments. In accordance with these regulations, in the last decades in some regions e.g. in Eastern Asia and in the Near East, HCFC-22 has been

- 5 used as interim replacement gas for more potent ozone-depleting substances such as CFCs, although it has been phased out in the developed countries (Fortems-Cheiney et al., 2013). Because HCFC-22 is emitted in locally restricted regions, in particular in Eastern Asia and therefore in the Asian monsoon region, this trace gas is very well suited to study transport processes in the Asian monsoon anticyclone. The region where HCFC-22 emissions occur in Eastern Asia overlap in parts with the emission tracer for India/China in our CLaMS model simulation for 2012.
- 10 For summer 2012 no MIPAS measurements are available. Therefore, we compare our results with MIPAS measurements for 2008, because MIPAS HCFC-22 measurements have a very good coverage in summer 2008 over Asia. For this comparison, we perform a CLaMS model simulation for summer 2008 with the same setup for emission tracers as for the model simulation of 2012 described in Sect. 3. Fig. 13a shows mean values of HCFC-22 (Chirkov et al., 2016) measured by the MIPAS for July, August and September (JAS) 2008 at 380 K. To improve the measurement density of HCFC-22, synoptically interpolating of
- 15 multiple days of measurements is used through CLaMS trajectory calculations. For each day in JAS 2008, trajectories were computed from the time of measurements in a time window of 5 days (i.e. -2 and +2 days) to 12:00 UTC of the selected day. Subsequently, the mean HCFC-22 values are calculated on a 3° × 3° latitude-longitude grid between 370 and 390 K.

Similar patterns of HCFC-22 and the emission tracer for India/China are found (see Figs. 13) in the region of the Asian monsoon anticyclone in 2008. This indicates that the two horizontal transport pathways from the Asian monsoon anticyclone to the northeast and to the southwest found in our model results for the India/China tracer in 2012 are also apparent in the

20 to the northeast and to the southwest found in our model results for the India/China tracer in 2012 are also apparent in the MIPAS HCFC-22 measurements and CLaMS simulations for 2008.

The detailed position of the highest tracer values are somewhat different between 2008 and 2012, probably caused by the interannual variability of the monsoon. Further, enhanced HCFC-22 values are also found in the Southern Hemisphere at the southern edge of the TTL most likely caused by upward transport in the tropics (see patterns of emission tracer for Southeast

25 Asia and the tropical Pacific in Appendix A). Nevertheless, MIPAS HCFC-22 measurements demonstrate that the large-scale transport pathways from the Asian monsoon anticyclone at its northeastern flank and at its western flank found by artificial tracers of air mass origin in the CLaMS model are also evident in global satellite measurements.

4.4.2 Flooding of the extratropical lower stratosphere

The accumulation of young air masses from Asia since 1 May 2012 in the extratropical lower stratosphere is calculated using
the isentropic transport barrier at different levels of potential temperature derived by Kunz et al. (2015) as shown in Fig. 14 and Tab. 2. Mean values for To calculate the percentages of different emission tracers are within the lower extratropical stratosphere at a certain level of potential temperature, we use the following approach: For each day between 1 May and 31 October 2012 a mean value for each emission tracers of all CLaMS air parcels is calculated for PV values larger (lower) than those at the transport barrier (see Tab. 2) and for air masses poleward of 30° N (30° S) at a specific isentropic level (Θ± 0.5 K). End of

October 2012, the contributions of all boundary emission tracers on the the composition of the extratropical northern lower stratosphere are at 360 K \approx 44%, at 380 K \approx 35%, and at 400 K \approx 23%, with highest contributions the remaining fraction of air consisting of aged air. The highest contributions of the boundary emission tracers are from India/China uplifted within the Asian monsoon anticyclone, from Southeast Asia, and from the tropical Pacific Ocean. The contribution of all other regions of the Earth's surface (residue residual surface) are of minor importance (Tab. 2).

An equivalent analysis for the Southern Hemisphere (see Fig. 14c and Tab. 2) shows that the contribution of young air masses on to the composition of southern extratropical lower stratosphere is much lower. End of October 2012, the contributions of all boundary emission tracers on to the composition of the extratropical southern hemisphere lower stratosphere are at 340 K \approx 27%, at 360 K \approx 21%, and at 380 K \approx 2%. Here, highest contributions are from the tropical Pacific, Southeast Asia, and a

5

10 minor fraction from India/China representing air masses from the Asian monsoon anticyclone. Also here, the contribution of all other regions of the Earth's surface (residual surface) are of minor importance (see Tab. 2).

Our findings demonstrate the importance of the large-scale Asian monsoon system for the transport of young air masses from Asia and the tropical Pacific into the lower stratosphere of the Northern Hemisphere. This is in particular particularly important for chemical species with lifetimes longer than a few months in the stratosphere such as CO. In agreement with our simulations,

- 15 during TACTS/ESMVal an increase of the concentrations of long-lived tropospheric source gases (lifetimes > 3 months) such as CO, H₂O, and N₂O was found by in-situ aircraft measurements in the lower stratosphere over Europe from August until September associated with transport from the Asian monsoon anticyclone (Müller et al., 2015) (Müller et al., 2016). These transport processes result in an accumulation of young air masses originating in boundary regions from Asia and the tropical Pacific Ocean within the lower stratosphere in the Northern Hemisphere during autumn 2012.
- 20 Because water vapor is an important greenhouse gas and even small perturbations of water vapor mixing ratios in the ExUTLS have a significant impact on surface climate, we are interested to estimate the impact of the Asian monsoon on moistening the lower stratosphere. From our simulations, we roughly estimate the fraction of H₂O originating in India/China, Southeast Asia, and tropical Pacific Ocean contributing to the water budget in the lower northern hemisphere stratosphere. Fig. 15 shows the mean water vapor content in the northern lower stratosphere for PV values larger than 7.2 PVU (10 PVU)
- 25 and northward of 30° N at 380 K (400 K) calculated with CLaMS (black line). In CLaMS at this altitude, contributions of cirrus clouds to the total water content are of minor importance. An increase of H₂O in the northern lower stratosphere is found in our simulation during summer and autumn as reported in previous studies (e.g. Ploeger et al., 2013; Zahn et al., 2014) (Müller et al., 2015) (Müller et al., 2016). The fraction of H₂O from different boundary tracers is indicated by different colors. End of October 2012, a contribution of approximately 1.5 ppmv (1.0 ppmv) H₂O are originates from source regions in
- 30 Asia and the tropical Pacific compared to a mean water vapor content of $\approx 5 \text{ ppmvppm}(4.5 \text{ ppm})$ at 380 K (400 K). Total H₂O without contributions of all boundary tracers i. e. the contribution of aged air (see Fig. 15, green line) shows a decrease during summer and autumn. Mean H₂O from AURA-MLS (version 3.3 and version 4) in the northern stratosphere calculated similar as mean CLaMS H₂O values for PV values larger than 7.2 PVU (10 PVU) and northward of 30° N at 380 K (400 K) also show an increase of water vapor within the lower northern stratosphere during summer and autumn 2012 (see Fig. 15, gray and
- 35 purple line) and therefore support our findings from CLaMS simulations. Differences between CLaMS and MLS mean values

in the lower northern stratosphere in particular at 380 K could by explained by sampling issues (different spacial resolution of MLS and CLaMS), vertical resolution of MLS limiting measurements of steep tracer gradient around the tropopause and the initialization of CLaMS H_2O at 1 May 2012 (above 400 K: AURA-MLS; below 350 K: CLaMS multi-annual simulation based on ERA-Interim water vapor with a linear transition between 350 K and 400 K, more details see (Vogel et al., 2015)).

- 5 Previous comparisons between CLaMS and MLS (v3.3) water vapor demonstrate that differences in H₂O found in the lower stratosphere at high latitudes are likely an artifact of the MLS averaging kernels (Ploeger et al., 2013). Further, a comparison of water vapor climatologies from international limb sounder by Hegglin et al. (2013) demonstrate that Aura-MLS (v3.3) H₂O measurements tend to be low at high latitudes in the lowermost stratosphere. Because of the good agreement between CLaMS H₂O and in situ water vapor measurements by the FISH instrument during TACTS/ESMVal in the lower northern stratosphere
- 10 (see Figs. 9-11) we are confident that CLaMS water vapor simulations within the extra-tropical lower stratosphere are reliable.

Thus, the increase of water vapor in the lower northern stratosphere during summer and autumn can be explained by transport of young tropospheric air masses from Asia and the Pacific ocean affected by the Asian monsoon anticyclone.

We refer this process as 'flooding' of the lower stratosphere with young tropospheric air masses (although a contribution of
 aged stratospheric air remains). In previous papers (e.g. Hegglin and Shepherd, 2007) (Müller et al., 2015) (Müller et al., 2016)
 also the expression 'flushing' as been used. Using the word 'flooding' we emphasize that the northern lower stratosphere is flooded with wet tropospheric young air masses from Asia and the tropical Pacific.

5 Discussion and conclusions

- In this paper, the transport mechanisms are analyzed of air masses from inside the Asian monsoon anticyclone to the northern hemisphere lower stratosphere. The combination of separation of anticyclones or filaments at the northeastern flank of the Asian monsoon anticyclone and subsequent horizontal transport along the subtropical jet associated with Rossby wave breaking is put forward here as a long-range transport mechanism from the Asian monsoon anticyclone to the northern hemisphere lower stratosphere. Our model simulations show that air masses originating in Southeast Asia and in the tropical Pacific Ocean uplifted outside of the core of the Asian monsoon anticyclone also contribute to air masses with tropospheric character found
- 25 in the northern lower stratosphere. This is consistent with 40-day backward trajectory calculations for the TACTS flight on 26 September 2012 (region No. 2 in Fig. 9) showing that about 39% of the trajectories originating in Southeast Asia, Western Pacific and the Asian monsoon region (Vogel et al., 2014). The residual 61% of the trajectories are from the background lower stratosphere. However in the 3-dimensional CLaMS simulations used here, irreversible mixing processes are additionally considered and a time period of 5 months (from 1 May 2012 until September 2012) is simulated which is longer than the
- 30 40-day backward trajectory calculations.

Our model simulation is driven by ERA-Interim reanalysis data. Convection in CLaMS is represented by vertical velocities in ERA-Interim reanalysis data. Thus, small-scale rapid uplift in convective cores is not included in CLaMS simulations. However, previous studies using 3-dimensional CLaMS simulations or trajectory calculations (e.g. Ploeger et al., 2010, 2015;

Pommrich et al., 2014; Vogel et al., 2014, 2015) (Müller et al., 2015) (Müller et al., 2016) (Ungermann et al., 2016; Konopka et al., 2016) in comparison with satellite or in-situ measurements show that ERA-Interim data are well suited to study transport processes in the vicinity of the Asian monsoon anticyclone and in the tropical tropopause layer.

For the time period of the eddy shedding event discussed in Sect. 4.1, our model study shows that the thermal tropopause

- 5 is a strong transport barrier above the separated anticyclone for young air masses (younger than 5 months) originating in Asia and the tropical Pacific Ocean. A mixing layer around the thermal tropopause up to ≈450K is found for fresh emissions (see Fig.5). The simulations also show that the vertical extension of the mixing layer above the thermal tropopause reaches ≈ 450K (see Fig.5) assuming typical mixing ratios found in the extra-tropical UTLS (≈ 30ppbv < < 60ppbv) (e.g. Hoor et al., 2004; Pan et al., 2004; Vogel et al., 2011b). Ungermann et al. (2016) analyzed an eastward eddy shedding event</p>
- 10 in August 1997 observed by the CRISTA space-shuttle mission using PAN (peroxyacetyl nitrate) measurements as a tracer of tropospheric air. They demonstrate that high-resolution CRISTA PAN measurements are very well suited to separate air masses of both the main Asian monsoon anticyclone and of the eastwards separated anticyclone from the stratospheric background. Above the separated eddy, CRISTA measurements also indicate a mixing layer 1-2 km above the thermal tropopause (up to ≈ 420K) with enhanced PAN mixing ratio a few days after the separation of the eddy from the main Asian monsoon anticyclone.
- 15 Thus, both our CLaMS simulations for 2012 presented in this paper and PAN CRISTA measurements from August 1997 byUngermann et al. (2016) indicate the existence of a mixing layer above the thermal tropopause of a separated eddy a few days after the eddy shedding event.

Our findings show that the most important pathways-mechanism for long-range transport of air masses from the Asian monsoon anticyclone to the extratropical lower stratosphere are eastward migrating anticyclones breaking off a few times each

- 20 summer from the main anticyclone (Dethof et al., 1999; Popovic and Plumb, 2001; Garny and Randel, 2013; Vogel et al., 2014; Ploeger et al., 2015) and filaments separated on the northeastern flank of the anticyclone. In our case study for the eddy shedding event on 20 September 2012, enhanced contributions of young air masses (younger than 5 months) are found within the separated anticyclone and filaments within the upper troposphere. Within the separated anticyclone, air masses with enhanced tropospheric tracers such as CO and enhanced values of artificial tracers of air mass origin are confined at the
- 25 top by the thermal tropopause ; only a small mixing layer around the tropopause is found in our model simulations. At the polar side, the subtropical jet acts as horizontal transport barrier. Therefore, the spatial structure of enhanced young air masses characterized by enhanced values of emission tracers from India/China looks like bubble within the upper troposphere. Those air masses are transported eastwards along the subtropical jet and can enter the extratropical lower stratosphere most likely driven by Rossby wave breaking events associated with double tropopauses (e. g. Vaughan and Timmis, 1998; Castanheira and
- Gimeno, 2011; Ungermann et al., 2013) Homeyer and Bowman 2012 (Homeyer and Bowman, 2013) (Homeyer et al., 2014). Our simulations show that after entering the lower stratosphere, these air masses are mixed irreversibly with the surrounding stratospheric air. During the TACTS/ESMVal campaign in August and September 2012 conducted with the German Research Aircraft HALO, filaments with enhanced amounts of tropospheric trace gases such as CO, CH_4 , and H_2O and reduced amounts of the stratospheric background were measured during three flights on 23, 25,
- and 26 September 2012 over northern Europe within the extratropical lower stratosphere. Our simulations confirm that these

signatures are remnants from of an eastwards migrating anticyclone and filaments released at the northeast flank of the Asian monsoon anticyclone that are transported within $\approx 8-14$ days to northern Europe.

In addition to this main transport pathway from the Asian monsoon anticyclone to the east along the subtropical jet and subsequent transport into the northern lower stratosphere, a second horizontal transport pathway out off of the anticyclone

5 to the west into the tropics (TTL) is found in agreement with MIPAS HCFC-22 measurements. This second transport pathway is mainly caused by westward eddy shedding, however this transport pathway vield predominantly to increase enhances predominantly the contribution of fresh emission from India/China within the TTL to the composition of TTL air at 380 K in summer and autumn 2012 (see Fig. 16).

Moreover, our simulation shows that young air masses originating in Southeast Asia and the tropical Pacific Ocean also contribute substantially to the composition of the lower stratosphere over Northern Europe in summer and autumn 2012. These 10 air masses are spread out distributed during summer and autumn globally on the tropical side of the subtropical jet stream of the Northern Hemisphere (see Appendix A). A large scale movement of those air masses around the Asian monsoon anticyclone is found in our model simulations indicating that the Asian monsoon anticyclone acts as a large stirrer within the TTL causing mixing of young air masses from Southeast Asia and the tropical Pacific Ocean into the TTL. Subsequent transport into the

northern lower stratosphere most likely by Rossby wave breaking events vield to results in an increase of young air masses 15 within the lower northern stratosphere.

Our findings demonstrate that emissions from young air masses originating in India/China and Southeast Asia, and tropical Pacific Ocean affected by the circulation of the Asian monsoon anticyclone have a significant impact on the chemical compositions of the lower stratosphere of the Northern Hemisphereby flooding the ExUTLS with young wet airmasses in particular.

20 Young moist air masses are flooding the extratropical lower stratosphere in the Northern Hemisphere with contributions up to $\approx 30\%$ at 380 K (with the remaining fraction being aged air) at the end of the monsoon season in September/October 2012 in contrast 2012. In contrast, the contribution of young air masses to the southern hemisphere is much lower. Our model simulations show that transport from boundary sources in Asia and the tropical Pacific during summer and autumn yield moistening of the lower northern stratosphere (of by roughly ≈ 1.5 ppmv/1.0 ppmv at 380 K/400 K end of October 2012) which could 25

have a potential an impact on surface climate.

Acknowledgements. We sincerely thank Andreas Engel and Harald Bönisch (University of Frankfurt) for coordinating the TACTS campaign, Hans Schlager (DLR Oberpfaffenhofen) for coordinating the ESMVal campaign, the HALO crew, and the HALO pilots. The authors gratefully acknowledge Laura Pan (NCAR, Boulder) and Eric Jensen (NASA Ames Research Center, Moffett Field) for helpful discussions 30 on transport pathways in the Asian monsoon circulation and their impact on moistening the lower stratosphere. In particular, we thank Laura Pan for comments on the concept of the Asian Monsoon anticyclone as a "tropospheric bubble". We thank the European Centre for Medium-Range Weather Forecasts (ECMWF) for providing meteorological analyses and the ERA-Interim reanalysis data. The authors gratefully acknowledge the computing time granted on the supercomputer JUROPA at the Jülich Supercomputing Centre (JSC) under the VSR project ID JICG11. Our activities were partly funded by the German Science Foundation (Deutsche Forschungsgemeinschaft) under the project LASSO (HALO-SPP 1294/GR 3786) and by the European Community's Seventh Framework Programme (FP7/2007-2013) under the project StratoClim (grant agreement no. 603557). The authors thank three anonymous reviewers for their very helpful comments.

References

5

- Anderson, J., Wilmouth, D., Smith, J., and Sayres, D.: UV Dosage Levels in Summer: Increased Risk of Ozone Loss from Convectively Injected Water Vapor, Science, 337, 835–839, doi:10.1126/science.1222978, 2012.
- Bian, J., Pan, L. L., Paulik, L., Vömel, H., and Chen, H.: In situ water vapor and ozone measurements in Lhasa and Kunmin during the Asian summer monsoon, Geophys, Res. Lett., 39, L19808. doi:10.1029/2012GL052996, 2012.
- Bönisch, H., Engel, A., Curtius, J., Birner, T., and Hoor, P.: Quantifying transport into the lowermost stratosphere using simultaneous in-situ measurements of SF₆ and CO₂, Atmos. Chem. Phys., 9, 5905–5919, 2009.
 - Bourassa, A. E., Robock, A., Randel, W. J., Deshler, T., Rieger, L. A., Lloyd, N. D., Llewellyn, E. J. T., and Degenstein, D. A.: Large Volcanic Aerosol Load in the Stratosphere Linked to Asian Monsoon Transport, Science, 337, 78–81, doi:10.1126/science.1219371, 2012.
- 10 Brasseur, G. and Solomon, S.: Aeronomy of the Middle Atmosphere: Chemistry and Physics of the Stratosphere and Mesosphere, Springer, Heidelberg, Germany, third edn., 2005.
 - Castanheira, J. M. and Gimeno, L.: Association of double tropopause events with baroclinic waves, J. Geophys. Res., 116, doi:10.1029/2011JD016163, 2011.
 - Chirkov, M., Stiller, G. P., Laeng, A., Kellmann, S., von Clarmann, T., Boone, C. D., Elkins, J. W., Engel, A., Glatthor, N., Grabowski, U.,
- Harth, C. M., Kiefer, M., Kolonjari, F., Krummel, P. B., Linden, A., Lunder, C. R., Miller, B. R., Montzka, S. A., Mühle, J., O'Doherty, S., Orphal, J., Prinn, R. G., Toon, G., Vollmer, M. K., Walker, K. A., Weiss, R. F., Wiegele, A., and Young, D.: Global HCFC-22 measurements with MIPAS: retrieval, validation, global distribution and its evolution over 2005-2012, Atmos. Chem. Phys., 16, 3345–3368, doi:10.5194/acp-16-3345-2016, www.atmos-chem-phys.net/16/3345/2016/, 2016.

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P.,

20 Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Koehler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, Q. J. R. Meteorol. Soc., 137, 553–597, doi:10.1002/qj.828, 2011.

Delaunay, B.: Sur la sphère vide. A la mémoire de Georges Voronoï, Bulletin de l'Académie des Sciences de l'URSS, 6, 793-800, 1934.

- 25 Dethof, A., O'Neill, A., Slingo, J. M., and Smit, H. G. J.: A mechanism for moistening the lower stratosphere involving the Asian summer monsoon, Q. J. R. Meteorol. Soc., 556, 1079–1106, 1999.
 - Dvortsov, V. L. and Solomon, S.: Response of the stratospheric temperatures and ozone to past and future increases in stratospheric humidity, J. Geophys. Res., 106, 7505–7514, 2001.

Fairlie, T. D., Avery, M. A., Pierce, R. B., Al-Saadi, J., Dibb, J., and Sachse, G.: Impact of multiscale dynamical processes and mixing on the

- 30 chemical composition of the upper troposphere and lower stratosphere during the Intercontinental Chemical Transport Experiment-North America, J. Geophys. Res., 112(D16S90), doi:10.1029/2006JD007923, 2007.
 - Fairlie, D., Vernier, J.-P., Natarajan, M., and Bedka, K. M.: Dispersion of the Nabro volcanic plume and its relation to the Asian summer monsoon, Atmos. Chem. Phys., 14, 7045–7057, doi:10.5194/acp-14-7045-2014, 2014.
- Fortems-Cheiney, A., Chevallier, F., Saunois, M., Pison, I., Bousquet, Cressot, C., Wang, H. J., Yokouchi, Y., and Artuso, F.: HCFC-
- 35 22 emissions at global and regional scales between 1995 and 2010: Trends and variability, J. Geophys. Res., 118, 7379–7388, doi:10.1002/jgrd.50544, 2013.

- Fromm, M., Kablick III, G., Nedoluha, G., Carboni, E., Grainger, R., Campbell, J., and Lewis, J.: Correcting the record of volcanic stratospheric aerosol impact: Nabro and Sarychev Peak, J. Geophys. Res., 119, 10343–10364, doi:10.1002/2014JD021507, http://dx.doi.org/ 10.1002/2014JD021507, 2014.
- Garny, H. and Randel, W. J.: Dynamic variability of the Asian monsoon anticyclone observed in potential vorticity and correlations with tracer distributions, J. Geophys. Res., 118, 13421–13433, doi:10.1002/2013JD020908, 2013.
- Garny, H. and Randel, W. J.: Transport pathways from the Asian monsoon anticyclone to the stratosphere, Atmos. Chem. Phys., 16, 2703–2718, doi:10.5194/acp-16-2703-2016, 2016.
 - Grooß, J.-U., Engel, I., Borrmann, S., Frey, W., Günther, G., Hoyle, C. R., Kivi, R., Luo, B. P., Molleker, S., Peter, T., Pitts, M. C., Schlager, H., Stiller, G., Vömel, H., Walker, K. A., and Müller, R.: Nitric acid trihydrate nucleation and denitrification in the Arctic stratosphere, Atmos. Chem. Phys., 14, 1055–1073. doi:10.5194/acp-14-1055-2014. 2014.
- Günther, G., Müller, R., von Hobe, M., Stroh, F., Konopka, P., and Volk, C. M.: Quantification of transport across the boundary of the lower stratospheric vortex during Arctic winter 2002/2003, Atmos. Chem. Phys., 8, 3655–3670, http://www.atmos-chem-phys.net/8/3655/2008/, 2008.
 - Hegglin, M. I. and Shepherd, T. G.: O₃-N₂O correlations from the Atmospheric Chemistry Experiment: Revisiting a diagnostic of transport
- and chemistry in the stratosphere, J. Geophys. Res., D19301, doi:10.1029/2006JD008281, 2007.
- Hegglin, M. I., Tegtmeier, S., Anderson, J., Froidevaux, L., Fuller, R., Funke, B., Jones, A., Lingenfelser, G., Lumpe, J., Pendlebury, D., Remsberg, E., Rozanov, A., Toohey, M., Urban, J., von Clarmann, T., Walker, K. A., Wang, R., and Weigel, K.: SPARC Data Initiative: Comparison of water vapor climatologies from international satellite limb sounders, J. Geophys. Res., 118, 11824–11846, doi:10.1002/jgrd.50752, 2013.
- 20 Homeyer, C. R. and Bowman, K. P.: Rossby wave breaking and transport between the tropics and extratropics above the subtropical jet, J. Atmos. Sci., 70, 607–626, 2012. 2013.
 - Homeyer, C. R., Bowman, K. P., Pan, L. L., Atlas, E. L., Gao, R.-S., and Campos, T. L.: Dynamical and chemical characteristics of tropospheric intrusions observed during START08, J. Geophys. Res., 116, D06111, doi:10.1029/2010JD015098, 2011.

Homeyer, C. R., Pan, L. L., Dorsi, S. W., Avallone, L. M., Weinheimer, A. J., O'Brien, A. S., DiGangi, J. P., Zondlo, M. A., Ryerson, T. B.,

- 25 Diskin, G. S., and Campos, T. L.: Convective transport of water vapor into the lower stratosphere observed during double-tropopause events, J. Geophys. Res., 119, 10941–10958, doi:10.1002/2014JD021485, 2014.
 - Hoor, P., Gurk, C., Brunner, D., Hegglin, M. I., Wernli, H., and Fischer, H.: Seasonality and extent of extratropical TST derived from in-situ CO measurements during SPURT, Atmos. Chem. Phys., 4, 1427–1442, 2004.

Hoor, P., Fischer, H., and Lelieveld, J.: Tropical and extratropical tropospheric air in the lowermost stratosphere over Europe: A CO-based

30 budget, Geophys. Res. Lett., 32, doi:10.1029/2004GL022018, 2005.

5

10

Hossaini, R., Chipperfield, M., Montzka, M. P., Rap, S. A., Dhomse, S., and Feng, W.: Efficiency of short-lived halogens at influencing climate through depletion of stratopsheric ozone, Nature Geoscience, 8, 186–190, doi:10.1038/ngeo2363, 2015.

Hsu, C. J. and Plumb, R. A.: Nonaxisymmetric Thermally Driven Circulations and Upper-Tropospheric Monsoon Dynamics, J. Atmos. Sci., 57, 1255–1276, 2001.

- 35 Kirk-Davidoff, D. B., Hintsa, E. J., Anderson, J. G., and Keith, D. W.: The effect of climate change on ozone depletion through changes in stratospheric water vapour, Nature, 402, 399–401, doi:10.1038/46521, 1999.
 - Konopka, P. and Pan, L. L.: On the mixing-driven formation of the Extratropical Transition Layer (ExTL), J. Geophys. Res., 117, D18301, doi:10.1029/2012JD017876, 2012.

Konopka, P., Grooß, J.-U., Günther, G., Ploeger, F., Pommrich, R., Müller, R., and Livesey, N.: Annual cycle of ozone at and above the tropical tropopause: observations versus simulations with the Chemical Lagrangian Model of the Stratosphere (CLaMS), Atmos. Chem. Phys., 10, 121–132, doi:10.5194/acp-10-121-2010, http://www.atmos-chem-phys.net/10/121/2010/, 2010.

Konopka, P., Ploeger, F., and Müller, R.: Entropy- and static stability-based Lagrangian model grids, in: Geophysical Mono-

- 5 graph Series: Lagrangian Modeling of the Atmosphere, edited by Lin, J., vol. 200, pp. 99–109, American Geophysical Union, doi:10.1029/2012GM001253, 2012.
 - Konopka, P., Ploeger, F., Tao, M., and Riese, M.: Regionally resolved impact of ENSO on the variability of the Brewer-Dobson circulation and of water vapor entering the tropical lower stratosphere, J. Geophys. Res., submitted, 2016.
 - Kunz, A., Sprenger, M., and Wernli, H.: Climatology of potential vorticity streamers and associated isentropic transport pathways
- 10 across PV gradient barriers, J. Geophys. Res., 120, 3802–3821, doi:10.1002/2014JD022615, http://dx.doi.org/10.1002/2014JD022615, 2014JD022615, 2015.
 - Li, Q., Jiang, J. H., Wu, D. L., Read, W. G., Livesey, N. J., Waters, J. W., Zhang, Y., Wang, B., Filipiak, M. J., Davis, C. P., Turquety, S., Wu, S., Park, R. J., Yantosca, R. M., and Jacob, D. J.: Convective outflow of South Asian pollution: A global CTM simulation compared with EOS MLS observations, Geophys. Res. Lett., 32, L14 826, doi:10.1029/2005GL022762, 2005.
- 15 Livesey, N. J., Read, W. G., Froidevaux, L., Lambert, A., Manney, G. L., Pumphrey, H. C., Santee, M. L., Schwartz, M. J., Wang, S., Cofield, R. E., Cuddy, D. T., Fuller, R. A., Jarnot, R. F., Jiang, J. H., Knosp, B. W., Stek, P. C., Wagner, P. A., and Wu, D. L.: EOS MLS Version 3.3 Level 2 data quality and description document, Tech. Rep. JPL D33509, Jet Propulsion Laboratory, available from http://mls.jpl.nasa.gov/, 2011.
 - McKenna, D. S., Grooß, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the
- 20 Stratosphere (CLaMS): 2. Formulation of chemistry scheme and initialization, J. Geophys. Res., 107, 4256, doi:10.1029/2000JD000113, 2002a.
 - McKenna, D. S., Konopka, P., Grooß, J.-U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 1. Formulation of advection and mixing, J. Geophys. Res., 107, 4309, doi:10.1029/2000JD000114, 2002b.
- 25 Meyer, J., Rolf, C., Schiller, C., Rohs, S., Spelten, N., Afchine, A., Zöger, M., Sitnikov, N., Thornberry, T. D., Rollins, A. W., Bozóki, Z., Tátrai, D., Ebert, V., Kühnreich, B., Mackrodt, P., Möhler, O., Saathoff, H., Rosenlof, K. H., and Krämer, M.: Two decades of water vapor measurements with the FISH fluorescence hygrometer: a review, Atmos. Chem. Phys., 15, 8521–8538, doi:10.5194/acp-15-8521-2015, http://www.atmos-chem-phys.net/15/8521/2015/, 2015.
 - Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and
- 30 Engel, A.: Impact of the Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe 2012, Atmos. Chem. Phys. Discuss., 15, 34765–34812, doi:10.5194/acpd-15-34765-2015, 2015.
 note Müller et al.(2015) is deleted and replaced by Müller et al.(2016) !!
 - Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and Engel, A.: Impact of the Asian Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe
- 35 2012, Atmos. Chem. Phys. Discuss., 15, 34, <u>16</u>, <u>10</u> 765–34573–<u>10</u> 812, <u>2015</u>. <u>589</u>, doi:10.5194/acp-16-10573-2016, <u>2016</u>.
 - Orbe, C., Waugh, D. W., and Newman, P. A.: Air-mass origin in the tropical lower stratosphere: The influence of Asian boundary layer air, Geophys. Res. Lett., 42, 2015GL063937, doi:10.1002/2015GL063937, http://dx.doi.org/10.1002/2015GL063937, 2015.

- Pan, L. L., Randel, W. J., Gary, B. L., Mahoney, M. J., and Hintsa, E. J.: Definitions and sharpness of the extratropical troppause: A trace gas perspective, J. Geophys. Res., 109, doi:10.1029/2004JD004982, 2004.
- Pan, L. L., Konopka, P., and Browell, E. V.: Observations and model simulations of mixing near the extratropical tropopause, J. Geophys. Res., 111, D05106, doi:10.1029/2005JD006480, 2006.
- 5 Pan, L. L., Randel, W. J., Gille, J. C., Hall, W. D., Nardi, B., Massie, S., Yudin, V., Khosravi, R., Konopka, P., and Tarasick, D.: Tropospheric intrusions associated with the secondary tropopause, J. Geophys. Res., 114, D10302, doi:10.1029/2008JD011374, 2009.
 - Pan, L. L., Honomichl, S. B., Kinnison, D. E., Abalos, M., Randel, W. J., Bergman, J. W., and Bian, J.: Transport of chemical tracers from the boundary layer to the stratosphere associated with the dynamics of the Asian summer monsoon, J. Geophys. Res., to be submitted, 2016.
- 10 Park, M., Randel, W. J., Kinnison, D. E., Garcia, R. R., and Choi, W.: Seasonal variation of methane, water vapor, and nitrogen oxides near the tropopause: Satellite observations and model simulations, J. Geophys. Res., 109, D03302, doi:10.1029/2003JD003706, 2004. note Pan et al.(2004) is deleted !!
 - Park, M., Randel, W. J., Gettleman, A., Massie, S. T., and Jiang, J. H.: Transport above the Asian summer monsoon anticyclone inferred from Aura Microwave Limb Sounder tracers, J. Geophys. Res., 112, D16309, doi:10.1029/2006JD008294, 2007.
- 15 Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A., and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, Atmos. Chem. Phys., 8, 757-764, 2008.
 - Park, M., Randel, W. J., Emmons, L. K., and Livesey, N. J.: Transport pathways of carbon monoxide in the Asian summer monsoon diagnosed from Model of Ozone and Related Tracers (MOZART), J. Geophys. Res., 114, D08303, doi:10.1029/2008JD010621, 2009.
 - Pion, S. and Teillaud, M.: 3D Triangulations, in: CGAL User and Reference Manual, CGAL Editorial Board, 4.2 edn., http://doc.cgal.org/4.
- 20 2/CGAL.CGAL/html/packages.html#PkgTriangulation3Summary, 2013. Ploeger, F., Konopka, P., Günther, G., Grooß, J.-U., and Müller, R.: Impact of the vertical velocity scheme on modeling transport across the tropical tropopause layer, J. Geophys. Res., 115, D03301, doi:10.1029/2009JD012023, 2010.
 - Ploeger, F., Günther, G., Konopka, P., Fueglistaler, S., Müller, R., Hoppe, C., Kunz, A., Spang, R., Grooß, J.-U., and Riese, M.: Horizontal water vapor transport in the lower stratosphere from subtropics to high latitudes during boreal summer, J. Geophys. Res., 118, 8111–8127,
- 25 doi:10.1002/jgrd.50636, 2013.
 - Ploeger, F., Gottschling, C., Grießbach, S., Grooß, J.-U., Günther, G., Konopka, P., Müller, R., Riese, M., Stroh, F., Tao, M., Ungermann, J., Vogel, B., and von Hobe, M.: A potential vorticity-based determination of the transport barrier in the Asian summer monsoon anticyclone, Atmos. Chem. Phys., 15, 13145-13159, doi:doi:10.5194/acp-15-13145-2015, 2015.

Pommrich, R., Müller, R., Grooß, J.-U., Konopka, P., Ploeger, F., Vogel, B., Tao, M., Hoppe, C. M., Günther, G., Spelten, N., Hoffmann,

- 30 L., Pumphrey, H.-C., Viciani, S., D'Amato, F., Volk, C. M., Hoor, P., Schlager, H., and Riese, M.: Tropical troposphere to stratosphere transport of carbon monoxide and long-lived trace species in the Chemical Lagrangian Model of the Stratosphere (CLaMS), Geosci. Model Dev., 7, 2895–2916, doi:10.5194/gmd-7-2895-2014, http://www.geosci-model-dev.net/7/2895/2014/, 2014.
 - Popovic, J. M. and Plumb, R. A.: Eddy Shedding from the Upper-Tropospheric Asian Monsoon Anticyclone, J. Atmos. Sci., 58, 93–104, 2001.
- 35 Randel, W. J. and Park, M.: Deep convective influence on the Asian summer monsoon anticyclone and associated tracer variability observed with Atmospheric Infrared Sounder (AIRS), J. Geophys. Res., 111, D12314, doi:10.1029/2005JD006490, 2006.
 - Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C., and Pumphrey, H.: Asian Monsoon Transport of Pollution to the Stratosphere, Science, 328, 611–613, doi:10.1126/science.1182274, 2010.

- Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative effects, J. Geophys. Res., 117, D16305, doi:10.1029/2012JD017751, 2012.
- Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Wine, P. H., Abbatt, J. P. D., Burkholder, J. B., Kolb, C. E., Moortgat, G. K., Huie, R. E., and Orkin, V. L.: Chemical kinetics and photochemical data for use in atmospheric studies, JPL Publication 10-6, 2011.
- 5 Schwartz, M. J., Manney, G. L., Hegglin, M. I., Livesey, N. J., Santee, M. L., and Daffer, W. H.: Climatology and variability of trace gases in extratropical double-tropopause regions from MLS, HIRDLS and ACE-FTS measurements, J. Geophys. Res., 120, 843–867, doi:10.1002/2014JD021964, 2015.
 - Sibson, R.: A brief description of natural neighbor interpolation, in: Interpreting Multivariate Data. Chichester, edited by Barnett, V., pp. 21–36, John Wiley, 1981.
- 10 Solomon, S., Rosenlof, K., Portmann, R., Daniel, J., Davis, S., Sanford, T., and Plattner, G.-K.: Contributions of stratospheric water vapor to decadal changes in the rate of global warming, Science, 327, 1219–1223, doi:10.1126/science.1182488, 2010.
 - Spang, R., Günther, G., Riese, M., Hoffmann, L., Müller, R., and Griessbach, S.: Satellite observations of cirrus clouds in the Northern Hemisphere lowermost stratosphere, Atmospheric Chemistry and Physics, 15, 927–950, doi:10.5194/acp-15-927-2015, http://www. atmos-chem-phys.net/15/927/2015/, 2015.
- 15 Sprenger, M., Wernli, H., and Bourqui, M.: Stratosphere-Troposphere Exchange and Its Relation to Potential Vorticity Streamers and Cutoffs near the Extratropical Tropopause, J. Atmos. Sci., 64(5), 1587–1602, doi:10.1175/JAS3911.1, http://dx.doi.org/10.1175/JAS3911.1, 2007.

Tissier, A.-S. and Legras, B.: Convective sources of trajectories traversing the tropical tropopause layer, Atmos. Chem. Phys., 16, 3383–3398, doi:10.5194/acp-16-3383-2016, 2016.

- 20 Uma, K. N., Das, S. K., and Das, S. S.: A climatological perspective of water vapor at the UTLS region over different global monsoon regions: observations inferred from the Aura-MLS and reanalysis data, Clim. Dyn., 43, 407–420, doi:10.1007/s00382-014-2085-9, 2014. Ungermann, J., Pan, L. L., Kalicinsky, C., Olschewski, F., Knieling, P., Blank, J., Weigel, K., Guggenmoser, T., Stroh, F., Hoffmann, L., and Riese, M.: Filamentary structure in chemical tracer distributions near the subtropical jet following a wave breaking event, Atmos. Chem. Phys., 13, 10517–10534, doi:10.5194/acp-13-10517-2013, 2013.
- 25 Ungermann, J., Ern, M., Kaufmann, M., Müller, R., Spang, R., Ploeger, F., Vogel, B., and Riese, M.: Observations of PAN and its confinement in the Asian Summer Monsoon Anticyclone summer monsoon anticyclone in high spatial resolution, Atmos. Chem. Phys. Discuss., 16, 8389–8403, doi:10.5194/acp-16-8389-2016, 2016.
 - Vaughan, G. and Timmis, C.: Transport of near-tropopause air into the lower midlatitude stratosphere, Q. J. R. Meteorol. Soc., 124, 1559–1578, 1998.
- 30 Vernier, J. P., Fairlie, T. D., Natarajan, M., Wienhold, F. G., Bian, J., Martinsson, B. G., Crumeyrolle, S., Thomason, L. W., and Bedka, K. M.: Increase in upper tropospheric and lower stratospheric aerosol levels and its potential connection with Asian pollution, J. Geophys. Res., doi:10.1002/2014JD022372, 2015.

Vogel, B., Konopka, P., Grooß, J.-U., Müller, R., Funke, B., López-Puertas, M., Reddmann, T., Stiller, G., von Clarmann, T., and Riese, M.: Model simulations of stratospheric ozone loss caused by enhanced mesospheric NO_x during Arctic Winter 2003/2004, Atmos. Chem.

Vogel, B., Feck, T., and Grooß, J.-U.: Impact of stratospheric water vapor enhancements caused by CH₄ and H₂ increase on polar ozone loss, J. Geophys. Res., 116, D05301, doi:10.1029/2010JD014234, 2011a.

³⁵ Phys., 8, 5279–5293, 2008.

- Vogel, B., Pan, L. L., Konopka, P., Günther, G., Müller, R., Hall, W., Campos, T., Pollack, I., Weinheimer, A., Wei, J., Atlas, E. L., and Bowman, K. P.: Transport pathways and signatures of mixing in the extratropical tropopause region derived from Lagrangian model simulations, J. Geophys. Res., 116, D05306, doi:10.1029/2010JD014876, 2011b.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., Hoor, P., Krämer, M., Müller, S., Zahn, A., and Riese, M.: Fast transport from Southeast Asia boundary layer sources to northern Europe: rapid uplift in typhoons and eastward eddy shedding of the Asian monsoon anticyclone, Atmos. Chem. Phys., 14, 12745–12762, doi:10.5194/acp-14-12745-2014, http://www.atmos-chem-phys.net/14/12745/2014, 2014.
- Vogel, B., Günther, G., Müller, R., Grooß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, Atmos. Chem. Phys., 15, 13699–13716, doi:10.5194/acp-15-13699-2015, http://www.atmos-chem-phys.net/15/13699/2015/, 2015.
- 10 Waymark, C., Walker, K. A., Boone, C. D., and Bernath, P. F.: ACE-FTS versions 3.0 data set: validation and data processing update, Ann. Geophys., 56, doi:10.4401/ag-6339, 2013.
 - Wu, X. and Lü, D.: Ten-year climatological features and air origin of midlatitude double tropopauses, Adv. Atmos. Sci., 32, 1592–1602, doi:10.1007/s00376-015-5036-4, 2015.
 - Zahn, A., Weppner, J., Widmann, H., Schlote-Holubek, K., Burger, B., Kühner, T., and Franke, H.: A fast and precise chemiluminescence
- 15 ozone detector for eddy flux and airborne application, Atmos. Meas. Tech., 5, 363–375, doi:10.5194/amt-5-363-2012, 2012.
 - Zahn, A., Christner, E., van Velthoven, P. F. J., Rauthe-Schöch, A., and Brenninkmeijer, C. A. M.: Processes controlling water vapor in the upper troposphere/lowermost stratosphere: An analysis of 8 years of monthly measurements by the IAGOS-CARIBIC observatory, J. Geophys. Res., 119, 11,505–11,525, doi:10.1002/2014JD021687, 2014.
- Zöger, M., Schiller, C., and Eicke, N.: Fast in situ hygrometers: A new family of balloonborne and airborne Lyman- α photofragment fluo-
- 20 rescence hygrometers, J. Geophys. Res., 104, 1807 1816, 1999.

5



Figure 1. Global geographic location of artificial boundary layer sources regions in the CLaMS model, also referred to as 'emission tracers' (Vogel et al., 2015). The latitude and longitude range for each emission tracer is listed in Table 1.

 Table 1. Latitude and longitude range of artificial boundary layer sources in the CLaMS model, also referred to as "emission tracers". The geographic position of each emission tracer is shown in Fig. 1.

Emission tracer	Latitude	Longitude
Northern India (NIN)	$20-40^{\circ}$ N	55–90° E
Southern India (SIN)	$0-20^{\circ}$ N	55–90° E
Eastern China (ECH)	$20-40^{\circ}$ N	90–125° E
Southeast Asia (SEA)	$12^{\circ} \text{ S}-20^{\circ} \text{ N}$	90–155° E
Siberia (SIB)	40–75° N	55–180° E
Europe (EUR)	45–75° N	20° W–55 $^{\circ}$ E
Mediterranean (MED)	$35-45^{\circ}$ N	20° W– 55° E
Northern Africa (NAF)	0–35° N	20° W–55 $^{\circ}$ E
Southern Africa (SAF)	$36^{\circ} \text{ S-}0^{\circ} \text{ N}$	7–42° E
Madagascar (MDG)	27–12° S	42–52° E
Australia (AUS)	40–12° S	110–155° E
North America (NAM)	15–75° N	$160-50^{\circ} \mathrm{W}$
South America (SAM)	55° S-15° N	80–35° W
Tropical Pacific Ocean (TPO)	$20^{\circ} \text{ S}-20^{\circ} \text{ N}$	see Fig. 1
Tropical Atlantic Ocean (TAO)	$20^{\circ} \text{ S}-20^{\circ} \text{ N}$	see Fig. 1
Tropical Indian Ocean (TIO)	$20^{\circ} \text{ S}-20^{\circ} \text{ N}$	see Fig. 1
Background	remaining surface	



Figure 2. Horizontal distribution of the fraction of air originating in India/China (here the sum of emission from North India, South India, and East China) at 380 K potential temperature on 4 (top) and 5 (bottom) September 2012. The horizontal winds are indicated by white arrows. The 7.2 PVU surface is shown as thick black line indicating the climatological isentropic transport barrier at 380 K in September (Kunz et al., 2015).



Figure 3. Horizontal distribution of PV (left) and the fraction of air originating in India/China (right) (here the sum of emission from North India, South India, and East China) at 380 K potential temperature on 20, 23, and 26 September 2012. The horizontal winds are indicated by white arrows. The 7.2 PVU surface is shown as thick black line indicating the climatological isentropic transport barrier at 380 K in September (Kunz et al., 2015). The the thick white lines at 180°E and at 120°W mark the position of vertical curtains shown in Figs. 5 and 6.



Figure 4. As Fig. 3, but for simulated CO (left) and the fraction of air originating in Southeast Asia / tropical Pacific Ocean.



Figure 5. Latitude-height cross-sections from 40°S to 90°N at 180°E longitude on 20 September 2012 including the separated anticyclone ($\approx 25-45^{\circ}$ N) caused by an eddy shedding event for the fraction of air originating in India/China (top left), the fraction of air originating in Southeast Asia / tropical Pacific Ocean (top right), the fraction of air originating in the residual surface (bottom left), and buoyancy frequency squared N² (bottom right). The climatological isentropic transport barrier of 4 and 10 PVU at 350 K and 400 K, respectively, for September (thick white lines) and thermal tropopause (black dots) are shown. The corresponding levels of potential temperature are marked by thin white lines.



Figure 6. Latitude-height cross-sections of the Northern Hemisphere at 120° W longitude on 23 September 2012 including an northward directed filament of low PV ($\approx 35-80^{\circ}$ N) moving along the subtropical jet for the fraction of air originating in India/China (top left), the fraction of air originating in Southeast Asia / tropical Pacific Ocean (top right), the fraction of air originating in the residual surface (bottom left), and buoyancy frequency squared N² (bottom right). The climatological isentropic transport barrier of 4 and 10 PVU at 350 K and 400 K, respectively, for September (thick white lines) and thermal tropopause (black dots) are shown. The corresponding levels of potential temperature are marked by thin white lines.



Figure 7. Horizontal (top) cross-section of the fraction of air originating in India/China over Europe om 26 September 2012. The flight path transferred to noontime of the TACTS/ESMVal flight is shown as red-yellow line. Segments of the flight in the lower stratosphere with enhanced measured CO, CH_4 , and H_2O and reduced O_3 compared to the stratospheric background are highlighted in red and numbered for clarification (the same numbers are used in Fig. 9) The climatological isentropic transport barrier of 7.2 PVU at 380 K is shown as thick black line. The white thick line marks the position of the vertical (bottom) cross-section at 8°W longitude which is similar to the cross-sections shown in Fig. 5 and 6.



Figure 8. Horizontal cross-section of the fraction of air originating in India/China over Europe on 25 September at 370 K (top) and on 23 September 2012 at 380 K (bottom). The flight paths transferred to noontime for the TACTS/EsmVal flights 2012 over northern Europe are marked as red-yellow lines. Segments of the flight in the lower stratosphere with enhanced measured CO, $CH_{4,1}$ and H_2O and reduced O_3 compared to the stratospheric background are highlighted in red and numbered for clarification (the same numbers are used in Fig. 10 and 11). The climatological isentropic transport barrier of 6.0 PVU (at 370 K) and 7.2 PVU (at 380 K) are shown as thick black lines.



Figure 9. Top Panel: Time evolution (given in UT time) of potential temperature, the fraction of air originating in India/China, Southeast Asia/tropical Pacific and of the residual surface (sum of all other emission tracers covering the entire Earth's surface) simulated with CLaMS for the flight on 26 September 2012. Middle and Lower panel: CO, O_3 , CH₄, and H₂O simulations and measurements. Segments of the flight in the lower stratosphere with enhanced measured CO, CH₄, and H₂O and reduced O₃ compared to the stratospheric background are highlighted in gray and numbered for clarification.



Figure 10. As Fig. 9, but for the flight on 25 September 2012.



Figure 11. As Fig. 9, but for the flight on 23 September 2012.



Figure 12. Mean values of the contribution of the emission tracer for India/China at 380 K in July (a), August (b) and September (c) 2012. Plate (d) shows mean values for India/China for July, August and September 2012. The climatological isentropic transport barriers of 7.2 PVU (northern hemisphere) and -11.5 PVU (southern hemisphere) at 380 K are shown as thick white dots.



Figure 13. Mean values of HCFC-22 (a) measured by MIPAS for July, August, and September 2008 at 380 K synoptically interpolated using CLaMS trajectory calculations (details see Sect. 4.4.1). Plate (b) is similar as Fig. 12d, but for the year 2008.

Table 2. The contributions of all boundary emission tracers on the the composition of the extratropical lower stratosphere end of October 2012 at different levels of potential temperature using the climatological isentropic transport barrier derived by Kunz et al. (2015) for the northern and southern hemisphere. Highest contributions are from India/China, Southeast Asia, and the tropical Pacific Ocean. The contribution of all other regions of the Earth's surface are of minor importance (residue residual surface).

Θ	transport barrier	contribution of	contribution of
		all boundary tracer	residual surface
Northern Hemisphere			
340 K	3.8 PVU	48 %	7 %
360 K	5.5 PVU	44 %	5%
380 K	7.2 PVU	35 %	4 %
400 K	10.0 PVU	23 %	3%
420 K	13.5 PVU	14%	1 %
Southern Hemisphere			
340 K	-3.0 PVU	27 %	3 %
360 K	-5.6 PVU	21 %	2%
380 K	-11.5 PVU	2 %	<1%



Figure 14. Contribution of different emission tracers from India/China, Southeast Asia, tropical Pacific Ocean, and residual surface to the northern lower stratosphere at different levels of potential temperature 360 K (a) and 380 K (b) from May until October 2012. The same for the southern lower stratosphere at 360 K (c).



Figure 15. The increase of H_2O mixing ratios in the lower northern hemisphere stratosphere at 380 K (top) and at 400 K (bottom) during summer 2012 is shown (black line). The green line indicated H_2O mixing ratios without fractions from the Earth's boundary layer. Mean H_2O mixing ratios derived from MLS version v3.3 (gray) and version v4 (purple) in the northern extra-tropical lower stratosphere show also an increase in water vapor during summer and autumn. A rough estimation of the fraction of H_2O mixing ratios originating in India/China (red), Southeast Asia (yellow) and the tropical Pacific Ocean (blue) in the northern lower stratosphere at 380K-is also given.



Figure 16. Horizontal Transport pathways from the Asian monsoon monsoon anticyclone and the surrounding air masses (yellow) westward into the tropical troppause layer (blue arrow) and eastward along the subtropical (red arrows). Cross-troppause transport into the extra-tropical lower stratosphere occurs mainly above the Atlantic and Pacific Ocean most likely driven by Rossby wave breaking events (green arrows).
Appendix A: Additional emission tracers

Horizontal distributions of emission tracers for Southeast Asia and the tropical Pacific Ocean

Figs. 10 and 11 show that also air masses originating in Southeast Asia and in the tropical Pacific contribute to the composition

- 5 of the lower stratosphere over Northern Europe. The horizontal distribution of the emission tracers of Southeast Asia and the tropical Pacific at 380 K from July until September 2012 are shown in Fig. 17. Air masses with enhanced percentages of the emission tracer for tropical Pacific Ocean are found over the Maritime Continent and the western Pacific in July and August 2012 indicating upward transport outside of the Asian monsoon anticyclone (see Fig. 12 a and b). In September 2012, enhanced values are also found westward and eastward of the Maritime Continent indicating horizontal transport within the TTL (see
- 10 Fig. 17c and d (left)). Low values are found in the regions of the Asian monsoon anticyclone indicating the transport barrier around the anticyclone which inhibit horizontal exchange of air masses. Further transport into the northern lower stratosphere also occurs most likely by Rossby wave breaking events along the subtropical jet similar as for air masses from the Asian monsoon anticyclone (enhanced contributions from India/China; see Fig. 12).
- Air masses with enhanced contributions from Southeast Asia are found both within the Asian monsoon anticyclone and over
 the Maritime Continent and the western Pacific caused by the geographic position of the emission tracer of Southeast Asia as shown in Fig. 17 (right). Thus, enhanced contributions from Southeast Asia are found as a superposition of transport within the Asian monsoon anticyclone and outside in the western Pacific region is evident in the horizontal distribution of the emission tracer for Southeast Asia.

20 Appendix B: Description of interpolation method

Full 3-D Delaunay triangulation

25

In this study, the used interpolation method to determine mixing ratios of chemical species and percentages of artificial tracers of air mass origin along the flight path of the TACTS/ESMVal flights (see Figs. 9–11) from CLaMS data is upgraded from the previously used semi-2-D scheme to a full 3-D Delaunay triangulation.

As a first step, CLaMS backward trajectories are calculated from time and space of the measurements to the CLaMS model output at noon one day before the measurement. Subsequently, the CLaMS data calculated on an irregular grid are interpolated on the endpoints of the backward trajectories as described below.

The complexity of creating the Delaunay triangulation increases in the worst-case with the square of the inserted points. To reduce the number of points and to get rid of the spherical shape of the points, it is advantageous to partition the surface of the Earth in six latitudinal bands (0° to 36°, 36° to 72°, and 72° to 90° for each hemisphere). The four latitudinal bands between 72°S and 72°N are in addition split into longitudinal areas (every 36° for the two bands between 0° and 36°; every 45° for the two bands between 36° and 72°). This partitioning was selected to to give roughly evenly sized areas with a sufficient amount



Figure 17. Mean values of the contribution of the emission tracer for tropical Pacific Ocean (left) and Southeast Asia (right) at 380 K in July (a), August (b) and September (c) 2012. The climatological isentropic transport barriers of 7.2 PVU (northern hemisphere) and -11.5 PVU (southern hemisphere) at 380 K are shown as thick white dots.

of air parcels. Within each of these areas, a Cartesian 3-D Delaunay triangulation including all contained parcels and a certain amount of neighboring parcels (here extending each area by 25% in longitude and latitude) is employed (Delaunay, 1934). The chosen amount of overlap assures that all relevant parcels surrounding the area are included in the triangulation for the typical air parcel density of the examined simulations. The coordinate system employed for each area is locally projected using

5 a stereographic projection on the horizontal center of each area.

In the CLaMS simulation used here, a horizontal resolution of 100 km is used with a maximum vertical resolution (thickness of the model layer) of about 400 m around the tropopause and thicker layers above and below the tropopause (in general the geometric thickness of a layer Δz is described by $\Delta z = \alpha \Delta r$ with the aspect ratio α and the horizontal resolution Δr ; α is controlled by the assumption that the entropy of the system is uniformly distributed over all air parcels (more details see

Konopka et al., 2012; Pommrich et al., 2014)). The previously used semi-2-D scheme interpolation is effectively restricted to 10 occur within one model layer.

The thickness of each layer in hybrid coordinates is defined corresponding to a fixed amount of a virtual altitude in kilometer δ within the coordinate system of each area. The vertical position of each air parcel within a layer is assigned to the virtual coordinate by its relative position in the layer: If the boundaries of the layers are defined as ζ_i with $\zeta_{i+1} > \zeta_i$, $i \in \mathbb{N}_0$, then the virtual vertical coordinate z' of an air parcel with vertical hybrid coordinate ζ within one area is defined as

5

$$z' = \left(i + \frac{\zeta - \zeta_i}{\zeta_{i+1} - \zeta_i}\right) \cdot \delta \tag{B1}$$

For the CLaMS simulation used here, a factor of 100 km was chosen for δ , corresponding to the typical horizontal distance of parcels within one layer according to the horizontal resolution.

Not used here, but useful for other applications is a simple linear relationship between the potential temperature of an air 10 parcel and its virtual vertical altitude of, e.g., $z' = \theta \cdot 10 \text{ km/K}$.

Using the coordinate transformation with the virtual vertical coordinate as defined in equation B1 and using the segmentation in different latitude-longitude areas, the triangulation can be calculated in Cartesian coordinates involving a strongly reduced amount of air parcels compared to all air parcels of the model simulation. Using the 3-D triangulation, several fast and useful interpolation options are readily available, such as nearest neighbor or barycentric interpolation. We choose to employ natural

15 neighbor (or Sibson) interpolation (Sibson, 1981), which computes the weighted means of all neighboring points in contrast to methods such as the barycentric interpolation, that use only neighboring points spanning a tetrahedron around the enclosed air parcel.

The Computational Geometry Algorithms Library (CGAL) offers all routines necessary to efficiently implement the described algorithm (Pion and Teillaud, 2013). The algorithm and its variants were implemented as a python module called

20 juregrid3d (Jülich Regridding 3D) using C++ and cython for the performance critical parts.