

Organic Aerosol source apportionment in London 2013 with ME-2: exploring the solution space with annual and seasonal analysis

Ernesto Reyes-Villegas¹, David C. Green², Max Priestman², Francesco Canonaco³, Hugh Coe¹, André S.H. Prévôt³, James D. Allan^{1,4}

5 ¹School of Earth, Atmospheric and Environmental Sciences, The University of Manchester, Manchester, M13 9PL, UK

² School of Biomedical and Health Sciences, King's College London, London, UK

³ Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen PSI, Switzerland

⁴ National Centre for Atmospheric Science, The University of Manchester, Manchester, M13 9PL, UK

Correspondence to: James D. Allan (james.allan@manchester.ac.uk)

10 **Abstract.** The Multilinear Engine (ME-2) factorisation tool is being widely used following the recent development of the Source Finder (SoFi) interface at the Paul Scherrer Institute. However, the success of this tool, when using the a-value approach, largely depends on the inputs (i.e. target profiles) applied as well as the experience of the user. A strategy to explore the solution space is proposed, where the solution that best describes the organic aerosol (OA) sources is determined according to the systematic application of predefined statistical tests. This includes trilinear regression, which proves to be a
15 useful tool to compare different ME-2 solutions. Aerosol Chemical Speciation Monitor (ACSM) measurements were carried out at the urban-background site of North Kensington, London from March to December 2013, where for the first time the behaviour of OA sources and their possible environmental implications are studied using an ACSM. Five OA sources were identified: biomass burning OA (BBOA), hydrocarbon-like OA (HOA), cooking OA (COA), semivolatile oxygenated OA (SVOOA) and low-volatility oxygenated OA (LVOOA). ME-2 analysis of the seasonal datasets (spring, summer and
20 autumn) showed a higher variability in the OA sources that was not detected in the combined March-December dataset; this variability was explored with the triangle plots f44:f43 f44:f60, where a high variation of SVOOA relative to LVOOA was observed in the f44:f43 analysis. Hence, it was possible to conclude that, when performing source apportionment to long-term measurements important information may be lost and this analysis should be done to short periods of time such as seasonally. Further analysis on the atmospheric implications of these OA sources was carried out, identifying evidence of the
25 possible contribution of heavy-duty diesel vehicles to air pollution during weekdays compared to those fuelled by petrol.

Keywords: Aerosol sources, matrix factorization, air quality, PM₁, ACSM, SoFi.

1. Introduction

Developed countries have made great improvements in air quality. However, air pollution still represents a significant air quality issue, mainly in urban cities, due to the sheer number of inhabitants and the associated anthropogenic emissions
30 resulting from the inhabitants' daily activities (transportation, energy production and industrial activities). In particular, aerosols have significant effects on air quality (Watson, 2002; Pope III and Dockery, 2006; Keywood et al., 2015).

Organic aerosols (OA) are one of the main constituents of submicron particulate matter, composing 20–90% of the total submicron particle mass (Zhang et al., 2007). OA are classified according to their origin as primary OA (POA) or secondary OA (SOA). POA are directly emitted from a range of sources while SOA are produced from gaseous precursors (VOCs) by
35 chemical reactions in the atmosphere. POA sources range from traffic emissions (hydrocarbon-like OA, HOA), biomass burning OA (BBOA), to OA emissions from cooking (COA) among others. Kupiainen and Klimont (2007) determined that the main sources of POA in Europe were emissions from traffic and the residential combustion of solid fuels. Allan et al. (2010) identified three POA sources: transport, burning of solid fuels and cooking in Manchester and London. SOA are the

40 main constituents of OA ranging from 64% in urban areas to 95% in rural sites (Zhang et al., 2007). Previous source apportionment studies (Zhang et al., 2011) often identified a highly oxygenated fraction with low volatility (LVOOA) and a less oxygenated and more volatile species (SVOOA). In general, SVOOA represent fresh SOA, which, after photochemical processing, evolve into LVOOA (Jimenez et al., 2009). POA and SOA concentrations vary over seasons and years, thus in order to study the OA sources and processes as well as their impacts on air quality, it is necessary to carry out long-term measurements and subsequent source apportionment data analysis.

45 Aerosol mass spectrometry has been widely used for measuring aerosol concentrations in a wide range of ground based measurements (Hildebrandt et al., 2011; Mohr et al., 2012; Saarikoski et al., 2012; Young et al., 2015b). In particular, the Aerosol Chemical Speciation Monitor (ACSM), which has been recently developed (Ng et al., 2011), has been used to carry out long-term measurements of non-refractory submicron aerosols around the world, for instance an industrial-residential area in Atlanta, Georgia (Budisulistiorini et al., 2013), a high-elevation mountain in Canada (Takahama et al., 2011), at
50 background locations in South Africa, (Vakkari et al., 2014) and Spain (Minguillón et al., 2015a; Ripoll et al., 2015), a semi-rural site in Paris (Petit et al., 2015) and at an urban background site in Switzerland (Canonaco et al., 2015).

Source apportionment techniques have been widely used to quantitatively determine aerosol sources. The main source apportionment models include: chemical mass balance (CMB) and positive matrix factorization (PMF).

55 CMB uses prior knowledge of source profiles and assuming that the composition of all sources is well defined and known (Henry et al., 1984). This technique is ideal when changes between the source and the receptor are minimal, although this barely happens in real atmospheric conditions and the constraints may add a high level of uncertainty.

PMF is a least-squares approach based on a receptor-only multivariate factor analytic model (Paatero and Tapper, 1994). The main difference between PMF and CMB is that PMF does not require any information as input to the model and the profiles and contributions are uniquely modelled by the solver (Paatero et al., 2002). PMF was applied to OA data measured
60 with an AMS for the first time by Lanz et al. (2007), using measurements taken at an urban background site in Zurich in the summer of 2005, where six OA sources were determined: LVOOA, SVOOA, HOA, Charbroiling-like OA, BBOA and COA. Subsequently, PMF was successfully applied to other datasets, acquired from a wide range of sampling sites and with different techniques, Ng et al. (2010) compiled and analysed 43 studies carried out at different sites around the world. This study provided a broad overview of aerosol composition and the importance of SOA as well as BBOA and HOA sources. In
65 other PMF studies, it was possible to find other relevant sources such as COA (Allan et al., 2010; Huang et al., 2010; Liu et al., 2012; Mohr et al., 2012; Sun et al., 2013; Crippa et al., 2013a).

ME-2 is a multivariate solver that determines solutions using the same equations as PMF (Paatero, 1999), with the possibility of using previous knowledge (factor time series and / or factor profiles) as inputs to the model, to partially constrain the solution, thereby reducing the rotational ambiguity (Paatero et al., 2002) and leading to more interpretable PMF solution(s)
70 as shown in Lanz et al. (2008) where three sources of OA were successfully determined (traffic related, solid fuel and secondary OA) during winter in an urban-background site in Zurich. Here unconstrained PMF runs failed to identify the environmental solution. This was most probably due to a high degree of temporal co-variation in the OA sources driven by low temperatures and periods of strong inversion.

The development of the Source Finder (SoFi) interface (Canonaco et al., 2013) written on the software package Igor Pro
75 (WaveMetrics, Inc.) together with a further standardised approach developed by Crippa et al. (2014), allowed different OA source apportionment studies to be undertaken such as a study at a suburban background site in Paris, France during January-March 2012 (Petit et al., 2014); laboratory studies analysing atmospheric ageing from the photo-oxidation of α -pinene and of wood combustion emissions in smog chambers and flow reactors (Bruns et al., 2015); long-term measurements (February

2011-February 2012) carried out at an urban background site in Zurich, Switzerland studying differences in oxygenated OA during summer and winter periods (Canonaco et al., 2015). As part of the ACTRIS project (Aerosols, Clouds, and Trace gases Research InfraStructure network) (Fröhlich et al., 2015) an intercomparison between 14 ACSM and one high resolution time of flight aerosol mass spectrometer (HR-ToF-AMS) was carried out at the SIRTA site in Gif-sur-Yvette near Paris, being able to identify 4 sources: hydrocarbon-like OA (HOA), OA related to cooking activities (COA), biomass burning related OA (BBOA) and oxygenated organic aerosol (OOA). These four sources were successfully identified from HR-ToF-AMS measurements with unconstrained PMF analysis. However, in the case of the ACSM datasets, it was necessary to partially constrain solutions via ME-2 analysis; probably due to the low signal to noise ratio of ACSM data compared to the AMS and the rural site type. Furthermore, new ME-2 source apportionment studies have been published this year (Bozzetti et al., 2016; Fountoukis et al., 2016; Milic et al., 2016; Elser et al., 2016), and even more are expected to come due to the successful application of SoFi. Thus, new strategies to systematically explore the solutions are needed.

This study includes data analysis of the first ACSM instrument deployed in the UK at the North Kensington site from March to December 2013, using the recently developed graphical interface SoFi, to perform non-refractory OA source apportionment analysis with the ME-2 factorization tool, implementing a strategy to determine the solution that best identifies OA sources, according to the statistical tests applied, and with further subsequent discussion of the various identified OA sources.

2. Methodology

The data used in this analysis (5th March – 30th December 2013) were obtained using an Aerosol Chemical Speciation Monitor (ACSM), deployed at the urban-background site in North Kensington, London. This instrument is owned by The Department for Environment, Food and Rural Affairs (DEFRA) and is part of the Aerosols, Clouds, and Trace gases Research InfraStructure Network (ACTRIS).

Source apportionment of OA was carried out using the PMF model implemented through the Multi-linear Engine tool (ME-2) and controlled via the Source Finder (SoFi) graphical user interface version 4.8, developed at the Paul Scherrer Institute (PSI), Switzerland (Canonaco et al., 2013).

2.1 Site and instrumentation

North Kensington (51.5215°, -0.2129°) is an urban background site located adjacent to a school, 7 Km to the west of central London. There is a residential road 30 metres to the east with an average traffic flow of 8,000 vehicles per day (Bigi and Harrison, 2010). This monitoring site is part of the DEFRA Automatic Urban and Rural Network (<http://uk-air.defra.gov.uk/networks/network-info?view=aur>).

As an urban background site, North Kensington is not significantly influenced by a single source or street, and concentrations may be analysed as an integrated contribution from all sources upwind of the site in London. This site is widely accepted as representative site of background air quality in central London and has a large set of long-term measurements for various pollutants (Bigi and Harrison, 2010). Different studies have been carried out at this site such as analysis of elemental and organic carbon concentrations in offline measurements of particulate matter with a diameter less than 10 micrometres (PM₁₀) (Jones and Harrison, 2005), PM₁₀ and NO_x association with wind speed (Jones et al., 2010), properties of nanoparticles (Dall'Osto et al., 2011), PM₁₀ and PM_{2.5} (Liu and Harrison, 2011) and in aerosol chemical composition (Beccaceci et al., 2015) in the atmosphere. The first long-term study of the behaviour of non-refractory inorganic and organic aerosols (PM_i) at the North Kensington site was carried out analysing cToF-AMS data collected from

January 2012 to January 2013 (Young et al., 2015a) where source apportionment analysis was carried out applying unconstrained PMF runs, with five sources identified: HOA, COA, solid fuel OA (SFOA), SVOOA and LVOOA.

120 The Aerosol Chemical Speciation Monitor (ACSM) measures, in real time, the mass and chemical composition of particulate organics, nitrate (NO₃), sulphate (SO₄), ammonium (NH₄) and chloride (Cl) ions, with a detection limit of 0.2 µgm⁻³ for an average sampling time of 30 min (Ng et al., 2011). These chemical species measured by the ACSM are determined according to the same methodology used in the AMS as defined by Allan et al. (2004). In principle, the ACSM is designed and built under the same sampling and detection technology as state-of-the-art Aerosol Mass Spectrometer (AMS) instruments. However, the ACSM is better suited for air quality monitoring applications due to its lower size, weight, cost, and power requirements; it is also more affordable to operate and is capable of measuring over long periods of time without supervision (Ng et al., 2011).

125 Time series of pollutants such as BC, CO, NO_x, OC, EC were downloaded from the DEFRA website for the North Kensington monitoring site. Wind speed and direction data were obtained from the meteorological station at Heathrow airport (located 17 Km from the sampling site). Wind data from this site was used due to its representativeness of regional winds without being affected by surrounding buildings.

2.2 Source apportionment (ME-2)

The multilinear engine algorithm (Paatero, 1999) is a multivariate solver that is typically used to solve the PMF model, which is based on a receptor-only factor analytic model (Paatero and Tapper, 1994). The bilinear representation of PMF solves Eq. (1), written in matrix notation, which represents the mass balance between the factor profiles and the concentrations.

$$X = G * F + E \quad (1)$$

The elements g_{ik} of matrix G represent the time series and the elements f_{kj} of matrix F represent the j elements of the profile (for example, mass spectrum) and E is the model residual.

140 The parameters f and g are fitted using a least squares approach that iteratively minimizes the variable Q (Paatero et al., 2002).

$$Q(f, g) = \sum_{i=1}^m \sum_{j=1}^n \left(\frac{e_{ij}}{\sigma_{ij}} \right)^2 \quad (2)$$

Where, e_{ij} represent the residuals and σ_{ij} the estimated uncertainty for the points i and j .

145 The variable Q depends on the number of selected factors and the size of the data matrix, hence it is necessary to normalize Q by the degree of freedom of the model solution (Q_{exp}) (Paatero et al., 2002) to monitor solutions.

$$Q_{exp} \cong n * m - p * (m + n) \quad (3)$$

150 Where p is the number of factors chosen, n number of samples and m the mass spectra. Ideally, if the model accurately captured the variability of the measured data, it would be expected to have a value of $Q/Q_{exp} = 1$, still this value depends on fluctuations in the source profiles, over- or underestimation of input data errors and from the model error.

Solutions using a least squares approach to solve a factor analysis problem may have linear transformations, also known as rotations (Paatero and Hopke, 2009). One advantage of ME2 over PMF is that the rotational ambiguity can be reduced by

using previous knowledge of profiles (for example mass spectra) or time series of different pollutants using the a -value approach. The equation 4 was applied using different target profiles (g_i) and a range of a -values (a) to constrain OA sources in different runs ($g_{i,run}$)

$$g_{i,run} = g_i \pm a * g_i \quad (4)$$

The a -value is a parameter that represents the degree of variability of the target profile, which typically ranges from zero to one, the closer to zero the more constrained the solution is (Lanz et al., 2008). The user should keep in mind that partially constrained solutions are carried out by compromising the Q/Q_{exp} value, which should be monitored to determine the feasibility of the solutions.

2.2.1 Target profiles and levels of constraint

In this study, solutions obtained with ME-2 were constrained using the a -value approach, by using four different sets of mass spectra from previous studies as target profiles (TP) (Table 1). The “a” set of TP represent BBOA and HOA average factor profiles obtained from an analysis carried out on different mass spectra from a variety of monitoring sites across Europe (Crippa et al., 2014) and COA obtained from a study in Paris (Crippa et al., 2013a). “c”, “s” and “w” sets of TP were provided by Young et al. (2015a) from a PMF analysis carried out on AMS measurements at the North Kensington site in London, 2012. “c” TP were obtained from an analysis performed on annual OA measured with a cToF-AMS (11 January 2012 – 23 January 2013). “s” and “w” TP were obtained from summer and winter measurements were taken with an HR-AMS (January–February 2012 and July–August 2012, respectively). The ACSM was specifically designed to deliver mass spectra that were equivalent to the AMS; with the AMS having a higher signal to noise ratio, it is expected the use of its mass spectra as TP to be appropriate. Moreover, we consider AMS-generated TP to be convenient to use especially considering there are more of these available, including the ones obtained from the same site. In this study, the suitability of different TP will be systematically assessed in the determination of OA sources using a wide range of a -values.

A wide range of combinations of TP and a -values were used during this analysis, all of them being run with three random initial values (seeds), to determine the stability of the solutions. Constraints were applied using one, two and three TP; in all the solutions, there were at least two unconstrained factors. Figure 1 shows the coding used to identify the different solutions, for example when constraining three factor profiles: wB5_H2_C3_S1.

2.3 Strategy to explore the solution space

The success of ME-2 relies on the additional use of *a priori* information in form of constraints. However, without a well-defined strategy or a limited analysis of the solution space, it may lead to a subjectively and inaccurately selected solution. Moreover, where possible, target profiles from different studies should be tested in order to determine which set of target profiles are the most adequate. Therefore, the following sections show the results of the analysis carried out on the dataset March-December 2013, where the considerations provided by Crippa et al. (2014) were applied. Moreover, new analysis techniques were developed to explore the solution space.

PMF solutions are run to determine the number of factors (sources) in the solution; this is carried out by running PMF for a different number of factors. Once the number of possible sources has been chosen, different combinations of a -values and constrained factors are tested to determine the solution that better identifies the OA sources. The residual of the solution provides important information; it is possible to determine if the solution is over estimated (negative residual) or under estimated (positive residual). When a structure on the diurnal residual is observed, it allows the factor which is affecting the residual to be determined (Crippa et al., 2014) and a decision to be made if the a -value should be modified or even if the target profile is the appropriate or not for this dataset. Together with the residual, it is recommended to look at the total

Q/Q_{exp} , which is a parameter used to monitor solutions, the best solution, according to the statistical tests applied, will be the one with values closest to one.

195 Trilinear regression is used as a new technique to explore the solution space in ME-2 analysis. Multilinear regression has been previously applied to analyse the relationship between POA and combustion tracers (Allan et al., 2010;Liu et al., 2011;Young et al., 2015b) as well as polycyclic aromatic hydrocarbons (Elser et al., 2016). This is used instead of simple linear regression because many of the combustion related variables will have multiple sources such as biomass burning and traffic. Eq. (4) shows the trilinear regression equation used to analyse the relationship between POA and combustion tracers.

$$Y = A + B[BBOA] + C[HOA] + D[COA] \quad (5)$$

200 Where “Y” is NO_x , BC, or CO.

B, C and D slopes represent the contribution of BBOA, HOA and COA to “Y” and the intercept A is representative of the “Y” background concentration. The following considerations should be taken into account: The slopes and intercepts should be positive as they represent air pollutant concentrations and the slope D is used as a validation parameter which should be close to zero, due to its low contribution to BC, NO_x and CO, owing to the fact that most cooking in the UK uses electricity or natural gas as a source of heat (DECC, 2015;NAEI, 2016). A nonzero value would indicate correlation with combustion tracers and thus the possibility that it is receiving interference from HOA, which has a similar mass spectrum. Chi-square is used as a “goodness of fit” where the lower the value the better fit between the analysed pollutants.

3. Results

3.1 Exploring the solution space for March-December dataset

210 This section shows the results from the analysis applied to determine the solution that best represents the OA sources for the complete dataset March-December 2013, according to the statistical tests applied, where a total of eight unconstrained and 25 constrained solutions were analysed.

3.1.1 Solutions, a -values and stability

215 Unconstrained runs with $f_{peak} = 0$ and 3 different seeds were performed in order to determine the number of OA sources, being five (BBOA, HOA, COA, SVOOA, LVOOA) the most adequate number of sources (Fig. S1.b) as it was possible to split the SOA into SVOOA and LVOOA. Further unconstrained analysis was performed by running five factor solutions with different f -peaks, from -1 to 1 with steps of 0.1 (Fig. S4) in order to select the PMF solution to be compared with the ME-2 analysis. ME-2 is run using a range of a -values, these a -values were selected after trial and error and according to the literature (Lanz et al., 2008;Crippa et al., 2014;Petit et al., 2014), which suggests that a -values depend on the similarity of the target profile and the factor profile being analysed: HOA mass spectra do not show high variability when compared with different sites, thus it is possible to restrict the constraint with a -values of 0.1-0.2. On the other hand, COA and BBOA mass spectra from different sites show high variability and a looser constraint should be applied (for example a -values 0.3-0.5 or higher).

225 Constraining only one or two factors of the five-factor solutions gave the least favourable results with high residuals and mixing factor profiles. When analysing the different seeds, these solutions also showed high variability between seeds; greater stability was found when three of the five factor solutions were constrained (Fig. S2), behaviour also observed by Crippa et al. (2014). As a result, in this analysis, five-factor solutions constraining three factors will be analysed for the first

seed. One PMF solution and two solutions constraining two factors were also used during the exploration (Fig. 2) for three sets of TP.

230 3.1.2 Q/Q_{exp} , diurnal residual and trilinear regression

For an ideal solution a Q/Q_{exp} value of 1.0 would be expected. However, there is not a standard criterion to define a satisfactory Q/Q_{exp} value as a certain amount of ‘model error’ will cause it to be systematically higher than unity (Ulbrich et al., 2009). When comparing different solutions from the same dataset (Fig. 2.b), it is possible to observe that there is not a significant variation on the Q/Q_{exp} (ranging between 1.88-2.2) when using different a -values, suggesting that all the solutions are mathematically acceptable. The unconstrained solution is the one with the lowest total Q/Q_{exp} with a value of 1.88, which is expected, as PMF calculates the solution by minimising this value; however, the PMF solution has a high Chi square and negative slope for COA (Fig. 2.a), implying that this solution is not environmentally acceptable, thus it is necessary to analyse all the different parameters in fig. 2 in order to select the solution that best identifies the OA sources.

235 Figure 2.a shows the diurnal residual analysis where solutions constrained with “c” target profiles present high positive residual around 14:00-19:00 hrs. Solutions constrained with “w” target profiles have a negative residual during early morning with a positive residual at 21:00 hrs. Hence, the solution with a better diurnal residual is within the solutions constrained with “a” target profiles.

240 Figure 2.b shows the trilinear regression outputs between NO_x and POA for the different solutions (see supplement S3 for BC and CO trilinear regressions). All the solutions properly identified the background NO_x concentrations (grey line). Solutions with “c” and “w” target profiles showed similar undesirable results as in the diurnal residual analysis, with “c” target profiles presenting negative COA slopes and “w” presenting high COA slopes and Chi square values, consistent with the diurnal residual analysis that the best solution, according to the statistical tests applied, is with the solutions constrained with “a” target profiles. Additionally, trilinear regression outputs show variations between different solutions constrained with “a” target profiles with changes mainly in the Chi square and the BBOA.

250 3.1.3 Diurnal concentrations and mass spectra

OA sources have characteristic diurnal trends, and they may be used, together with their respective mass spectra, to analyse the solutions and determine if all the factors in the solution are suitable, environmentally speaking. BBOA showed low concentrations during the day, with high concentrations at night, mainly related to domestic heating (Alfarra et al., 2007); HOA presents two peaks during the day related to commuting, one in the morning and another one in the evening (Zhang et al., 2005); COA has two peaks related to OA emissions from cooking activities: one peak at noon and one peak in the evening (Allan et al., 2010). SVOOA is temperature dependent with low concentrations during the day increasing in the evening due to the condensation of gas phase pollutants. LVOOA, due to its regional origin, does not show high variations in its diurnal trend.

255 Diurnal concentrations for all the solutions (Supplement S3) were analysed to determine the main sources. Here, it was possible to observe that solutions with undesirable outputs in the residual, total Q/Q_{exp} and/or trilinear regression were likely to have mixed diurnal concentrations between two sources. For example in the case of “c” TP solutions, CO and BC trilinear regressions (Fig. S5.a and S5.b) show better COA slopes with values close to zero, however due to the high diurnal residual (Fig. 2.a) and HOA with high concentrations during the evening (Fig. S5.c), suggesting mixing with BBOA, “c” TP solutions are not considered acceptable solutions.

265 These undesirable outputs previously observed were also detected when analysing the mass spectra of the different solutions; Fig. S3 shows examples of diverse situations that were found: in the solution wB7_H5_C7_S1 it is possible to observe

mixed factors where SVOOA has peaks of BBOA (m/z 60) and COA (m/z 55 and 57); one source with only one strong peak in its mass spectrum (SVOOA in solution cB3_H1_S1); The PMF solution was not able to properly identify a BBOA factor with low peaks at m/z 60 and 73 and a peak at m/z 60 for COA implying mixing with BBOA.

270 Finally, from this analysis, aB3_H2_C3_S1 was determined to be the solution that best represents the OA sources for March-December analysis, according to the statistical tests applied.

3.2 Seasonal analysis

275 When applying source apportionment, ME-2 considers that both target profiles and factor profiles remain constant over time, which may not be the case for long periods of time where meteorological conditions and pollutant emissions related to human activities vary greatly (Canonaco et al., 2015; Ripoll et al., 2015), thus the same analysis that was carried out on March-December data set was applied to data divided into seasons of the year: spring (March, April and May), summer (June, July and August) and autumn (September, October and November), see supplement S.3 for detailed information of the seasonal analysis.

280 Analysing the spring dataset (Fig. S7), solutions constrained with “a” and “c” TP were found to present the least favourable results with high Chi-square values and negative COA ratios in the trilinear analysis, as well as a higher negative diurnal residual; the solution wB3_H1_C3_S1 was deemed to be the best solution for spring analysis. Solutions constrained with “s” and “c” TP were the least favourable results for the summer analysis (Fig. S9) with low chi-square values in “s” target profiles, which show high negative residuals in the morning and at night. “c” target profiles show a high positive residual around 15:00-18:00 hrs; the solution aB5_H1_C3_S1 was found to be the best solution for the summer analysis. In the autumn analysis (Fig. S11), solutions constrained with “a” and “w” TP were found to be the least favourable results with high positive residuals in the morning, also “a” target profiles show high chi-square values. The solution cB3_H1_S1 was deemed the best solution for the autumn analysis, according to the statistical tests applied. It is worth to mention that all plausible solutions deconvolved a high percentage of the total OA mass (Fig. S12), with summer being the period with less OA mass estimated (90%) and the other periods with more than 95% of mass estimated from the total OA concentrations.

290 4. Discussion and atmospheric implications

4.1 Annual and seasonal solutions

In the following subsections, the outputs of annual and seasonal solutions are compared in order to further explore the variability of the different OA sources.

4.1.1 Total Q/Q_{exp} and diurnal residual

295 Analysing the total Q/Q_{exp} , all the solutions obtained were mathematically acceptable and with small variations between their different values: 1.95 for March-December, 2.01 for spring, 1.95 for summer and 1.96 for autumn (Fig. 3.a).

300 Q/Q_{exp} values obtained in this study are compared to values obtained in different ME-2 studies, for example Petit et al. (2014), in a study using an ACSM, obtained a Q/Q_{exp} value of 6; studies carried out in Spain during winter and summer obtained 1.15 and 0.38 respectively (Minguillón et al., 2015b). Q/Q_{exp} values obtained with PMF are also comparable with values obtained in this study, for example (Young et al., 2015a) obtained a value of 1.35 from annual measurements carried out with a cToF-AMS at this site; (Allan et al., 2010) obtained different Q/Q_{exp} values for the analysis carried out to three different data sets: a value of 3.9 from measurements obtained using a HR-ToF-AMS and values of 10.5 and 16.7 using a cToF-AMS; also Crippa et al. (2013b) identified a Q/Q_{exp} value of 4.59, on HR-ToF-AMS measurements during July 2009

305 at the urban background site in Paris. Due to all this variability of Q/Q_{exp} values found in the literature, this parameter alone cannot be used as a criterion to determine the solution that best identifies the OA sources.

It is in the diurnal residual where we can observe a high variation (Fig. 3.b), with autumn proving to be the most overestimated with negative residuals of $-0.033 \mu\text{g}\cdot\text{m}^{-3}$ mainly in the morning and at night. On the other hand, summer shows to be the most underestimated solution with values of $0.018 \mu\text{g}\cdot\text{m}^{-3}$ particularly between midday and 17:00 UTC. The fact that summer is underestimated from 12:00 to 17:00 UTC is probably related to the increase on photochemical activity, a situation that ME-2 is not able to capture as considers the mass spectra to remain constant over the period analysed. It is important to notice that these diurnal residuals of $0.03 \mu\text{g}\cdot\text{m}^{-3}$ or less are low compared with diurnal concentrations of the OA sources, which concentrations ranged $0.1\text{-}0.6 \mu\text{g}\cdot\text{m}^{-3}$.

4.1.2 Trilinear regression analysis

315 Looking at the trilinear outputs for the different periods analysed (Figure 3.a), HOA slopes present higher variability with values of 50.0 for March-Dec, 81.0 for spring, 41.0 for summer and 85.5 for autumn. The different BBOA and HOA slopes for spring, summer and autumn suggest that when looking at March-Dec solution only, there are seasonal variations, perhaps affected by changes on the inhabitants' daily activities (i.e. domestic heating) and meteorological conditions, that the March-Dec solution does not completely capture. With regard to COA slopes and background concentrations, they are well identified and relatively constant over the different periods analysed.

320 The analysis presented in section 4 shows that seasonal analysis more accurately deconvolves OA sources, being possible to obtain more detailed information that will be lost when running ME-2 for long periods of time.

4.1.3 Target profiles (TP) and their impact on the solutions

325 As previously mentioned, the chosen solutions were: aB3_H2_C3_S1 for March-December, wB3_H1_C3_S1 for spring, aB5_H1_C3_S1 for summer and wB3_H1_S1 for autumn. The fact that March-December and summer solutions were obtained with "a" TP is possibly due to the fact that these TP represent an average from different mass spectra, becoming robust TP able to deal with the variations of these two datasets; one large dataset (March-Dec) and one dataset with concentrations affected by the different photochemical processes due to the high temperatures (Summer). On the other hand, spring and autumn do not show these variations and their OA sources may be apportioned using winter TP which were obtained under similar temperatures.

330 Looking at the "c" and "s" TP, these were the ones with the least favourable results in all analyses carried out. This may be attributed to "c" being the only TP obtained with a cToF-AMS while the rest were obtained using a HR-AMS. In the case of "s" TP the unfavourable outputs are again related to the high variability present during this period of time. This analysis shows the importance of using the adequate TP when doing source apportionment as well as to explore solutions with different types of TP in order to determine the OA sources.

335 4.2 Variability of factor profiles

The variability of the different solutions previously obtained may be explored further with the triangle plots f_{44} vs f_{43} (Ng et al., 2010; Morgan et al., 2010) and f_{44} vs f_{60} (Cubison et al., 2011). The parameters f_{43} , f_{44} and f_{60} represent the ratio of the integrated signal at m/z 43, 44 and 60, respectively, to the total signal in the organic component mass spectrum. Figure 4.a shows that LVOOA, while having different values between solutions, is found in distinct areas of the plot (connecting lines are used to make the SVOOA variability clearer), whereas SVOOA shows values of f_{44} vs f_{43} with high variability. This analysis shows that the factors derived for SOA do not always conform to the model of LVOOA and SVOOA

proposed by Jimenez et al. (2009). Furthermore, the fact that the lines are going in different directions with the seasons of year means that the factorisation is identifying different aspects of the chemical complexity, as LVOOA and SVOOA rather than being originated from primary emissions are part of continuous physicochemical processes involving gases, aerosols and meteorological parameters among others. This serves to highlight that a 2-component model (LVOOA and SVOOA) is an oversimplification of a complex chemical system as concluded by Canonaco et al. (2015) who found significant f44 vs f43 difference for summer and winter analyses.

By analysing Fig. 4.b, it is possible to observe the variability in f60, with the lowest value obtained in summer (0.013) followed by spring, autumn and March-Dec (0.022, 0.024 and 0.034, respectively). Variability in biomass burning OA depends on the fuel type, burning conditions and level of processing (Weimer et al., 2008;Hennigan et al., 2011;Ortega et al., 2013;Young et al., 2015b). A study carried out by (Young et al., 2015b) in London, 2012, identified two types of solid fuel OA factors, attributed to differences in burning efficiency. BBOA evolution has been frequently observed with high f44 and low f60 values due to due to aging, oxidation and cloud processing (Huffman et al., 2009;Cubison et al., 2011). Thus, it was possible to obtain a variety of BBOA for the different seasons of the year, ranging from a fresh BBOA with a high f60 during autumn to a more oxidised BBOA with a low f60 during summer.

For all the solutions, COA presents an f60 value of approximately 0.01, which has been previously identified by Mohr et al. (2009) who obtained f60 values of 0.015 - 0.03 for different types of meat cooking. The fact that all the COA mass spectra present similar f44:f60 ratios suggest the COA footprint is relatively constant over the different seasons, being together with HOA the more adequate sources to constrain when applying the *a*-value approach.

4.3 Petrol and diesel contribution to traffic emissions

Traffic emissions contribute significantly to air pollution (Beevers et al., 2012;Carslaw et al., 2013;May et al., 2014). In order to better analyse traffic emissions and their impact on air quality, it is necessary to understand the fuel type and pollutant contribution from different vehicles. In particular, the United Kingdom has a considerable percentage of diesel-fuelled vehicles; according to the vehicle licencing statistics, the percentage of diesel-fuelled vehicles licenced has been increasing over the last few years from 22% in 2006 to 36.2% in 2014 respectively, while petrol-fuelled vehicles decreased from 77.7% to 62.9% (GOV.UK, 2015).

Diesel emits higher NO_x and HOA concentrations compared to petrol, while petrol emits higher concentrations of CO, according to the National Atmospheric Emissions inventory (NAEI, 2016), during 2014 the emission factors (units in kilotonnes of pollutant per Megatonne of fuel used) were: 11-12 for diesel and 1.9-4.3 for petrol in the case of NO_x and 2.4-5.6 for diesel and 11-50 for petrol in the case of CO. Moreover, there are variations between Light Duty Diesel (LDD) and Heavy Duty Diesel (HDD) emissions (LAEI, 2013), with LDD emitting higher NO_x concentrations and HDD emitting higher HOA concentrations.

It is possible to qualitatively analyse the impact of different fuels on air pollution by looking at weekday/weekend ratios (WD/WE), as previously done in several studies (Bahreini et al., 2012;Tao and Harley, 2014;DeWitt et al., 2015) and stating the hypothesis that different fuels will have different pollutant contribution during the week. This analysis is performed considering WD as Monday to Friday and WE only Sunday to eliminate the mixed traffic on Saturday. Another consideration is that heavy duty/light duty emissions fleet ratio is higher during the week (Lough et al., 2006;Bahreini et al., 2012;Heo et al., 2015). It is also important to state that heavy duty vehicles are exclusively diesel fuelled whereas light duty vehicles are a mixture of diesel and petrol.

380 Trilinear regression explained in section 2.3, was used with data divided into WD (Monday to Friday) and WE (Sunday) to
analyse the WD/WE contributions. Subsequently, it was possible to determine WD/WE ratios for the slopes: NO_x/HOA and
CO/HOA.

In order to compare these trilinear outputs with the WD/WE ratios between NO_x and CO, $\text{NO}_x/\Delta\text{CO}$ was calculated from
average concentrations: there is a difference in lifetime between CO (lifetime of months) and NO_x (lifetime of hours), thus it
385 is important to consider the background CO concentrations to be able to compare NO_x and CO concentrations. It is necessary
to perform a linear regression between CO and NO_x and calculate ΔCO , which is the average CO concentration minus the
intercept from the CO: NO_x linear regression.

Figure 5 shows the WD/WE ratios, where it is possible to observe $\text{NO}_x/\Delta\text{CO}$ ratios of 1.25, 1.35 and 1.36 for March-Dec,
summer and autumn, respectively, suggesting diesel with a higher contribution during WD compared to petrol. These
390 findings are confirmed by the CO/HOA ratios, which for the same periods of time, are lower than one (0.8, 0.45 and 0.9)
suggesting a lower contribution of petrol during weekdays compared to diesel. In spring, there are no considerable changes
to the WD/WE ratios, although a higher contribution of petrol is shown during WD with values of 1.28 for CO/HOA and
low diesel contribution. Analysing the NO_x/HOA ratios, the seasonal ratios show values of 1.07, 1.06 and 1.05 suggesting a
slightly higher contribution of LDD during WD than HDD.

395 **4.4 $\text{PM}_{2.5}$ daily concentrations and PM_1 composition**

$\text{PM}_{2.5}$ has been widely studied due to its potential to cause negative effects on health (Pope III and Dockery, 2006; Harrison
et al., 2012; Bohnenstengel et al., 2014). This adverse impact is directly connected to the size of the particles, making PM_1
more detrimental to health than $\text{PM}_{2.5}$ (Ramgolam et al., 2009). Moreover, analysing the aerosol contribution to PM_1 and its
association with $\text{PM}_{2.5}$ concentrations allows the possible influence of PM_1 on $\text{PM}_{2.5}$ levels to be determined. According to
400 the Daily Air Quality Index (DAQI), $\text{PM}_{2.5}$ concentrations are considered moderate when daily concentrations are between
35 and 52 $\mu\text{g m}^{-3}$ and high when levels are between 53 and 69 $\mu\text{g m}^{-3}$. Daily $\text{PM}_{2.5}$ concentrations during the sampling period
show that the majority of daily concentrations were considered to be low episodes (Fig. 6.a), with 10 episodes of moderate
concentrations and only two episodes of high $\text{PM}_{2.5}$ concentrations (55.2 and 61.5 $\mu\text{g m}^{-3}$).

Considering that PM_1 is composed mainly of OA, SO_4 , NO_3 , NH_4 and BC, it is possible to analyse the PM_1 composition
405 during $\text{PM}_{2.5}$ high concentrations (Fig. 6.b). Episodes with moderate and high $\text{PM}_{2.5}$ concentrations were observed with low
wind speeds (Fig. S13), being NO_3 and LVOOA the main PM_1 contributors. High NO_3 concentrations were observed during
spring as found in a previous study performed by Young et al. (2015a) who determined that NO_3 concentrations in spring
depend on air mass trajectory, precursors and meteorology. Different contributions from OA sources were identified: in the
episode in March, high BBOA concentrations were observed, whereas during the episodes in April and September, higher
410 concentrations of LVOOA were detected.

Defining BBOA, HOA COA and BC as primary and SVOOA, LVOOA, NO_3 , NH_4 and SO_4 as secondary aerosols, the main
 PM_1 contributors to $\text{PM}_{2.5}$ concentrations are secondary aerosols with a total contribution of 61% (Fig. 6.c). These findings
agree with a previous study carried out in this same monitoring site carried out by (Young et al., 2015a) who found
secondary aerosols to be the predominant source of PM_1 over the year, with different secondary inorganic and organic
415 aerosol contributions between winter and summer.

5. Conclusions

This study presents the source apportionment carried out using ME-2 within SoFi 4.8 of OA concentrations, measured with an ACSM from March to December 2013 at the urban-background site in North Kensington, London; the first time it was deployed in the UK.

420 ME-2 proved to be a robust tool to deconvolve OA sources. This study highlighted the importance of using appropriate mass spectra as target profiles and a -values when exploring the solution space. With the implementation of new techniques to compare different solutions, it was possible to systematically determine the solution with the best separation of OA sources, mathematically and environmentally speaking. The comparison carried out between the solution for the March-December dataset and the seasonal solutions showed high variations mainly in the SVOOA and the BBOA sources, with wide range of
425 $f_{44}:f_{43}$ values for SVOOA (Fig. 4.a) and f_{60} values ranging from 13×10^{-3} for summer to 24×10^{-3} for autumn (Fig. 4.b). These variations support the importance of running ME-2 during periods of time where weather conditions and emissions from human activities are less variable, such as seasonal analyses.

SVOOA presented a high variability in the oxidation state during the different seasons. This is due to the nature of SVOOA being affected mainly with high temperatures and ME-2 not being able to completely determine SVOOA concentrations.
430 These results support the indication that is not an accurate practice to use SVOOA as a target profile when analysing solutions. Trilinear regressions deliver quantitative information about the ratios between combustion tracers and POA. These ratios may be used as a proxy for other urban background sites to estimate POA concentrations.

From analysing heavy and light duty diesel emissions, the main contributor on weekdays was found to be from diesel emissions, particularly LDD emissions. Thus, in order to reduce traffic emissions on weekdays, LDD vehicles should be
435 targeted. For the $PM_{2.5}$ analysis (March-December 2013), the main PM_1 contributors to these concentrations were secondary aerosols and BC, which means that PM_1 contributors to $PM_{2.5}$ concentrations are related to emissions from combustion activities and also from secondary pollutants produced in the atmosphere.

This study delivers mass spectra and time series of OA sources for a long-term period as well as seasons of the year that maybe used in future studies as TP. Furthermore, the scientific findings provide significant information to strengthen
440 legislation as well as to support health studies in order to improve air quality in the UK.

Author contributions: D.C. Green, E. Reyes-Villegas and J.D. Allan designed the project; D.C. Green and M. Priestman operated, calibrated and performed QA of ACSM data; E. Reyes-Villegas performed the data analysis; E. Reyes-Villegas, F. Canonaco, D.C. Green, H. Coe, A.S.H. Prévôt, and J.D. Allan wrote the paper.

Data Availability: Monitoring data are available through the Department of the Environment, Food and Rural Affairs
445 (<https://uk-air.defra.gov.uk/networks/>). Factorisation data are available on request.

Acknowledgements: E. Reyes-Villegas is supported with a studentship by the National Council of Science and Technology–Mexico (CONACYT) under registry number 217687.

References

Alfarra, M. R., Prevot, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M., and
450 Baltensperger, U.: Identification of the mass spectral signature of organic aerosols from wood burning emissions, *Environmental Science & Technology*, 41, 5770-5777, Doi 10.1021/Es062289b, 2007.
Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A. M., Drewnick, F., Onasch, T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A generalised method for the extraction of

chemically resolved mass spectra from aerodyne aerosol mass spectrometer data, *J Aerosol Sci*, 35, 909-922, DOI 10.1016/j.jaerosci.2004.02.007, 2004.

455 Allan, J. D., Williams, P. I., Morgan, W. T., Martin, C. L., Flynn, M. J., Lee, J., Nemitz, E., Phillips, G. J., Gallagher, M. W., and Coe, H.: Contributions from transport, solid fuel burning and cooking to primary organic aerosols in two uk cities, *Atmos Chem Phys*, 10, 647-668, 2010.

460 Bahreini, R., Middlebrook, A. M., de Gouw, J. A., Warneke, C., Trainer, M., Brock, C. A., Stark, H., Brown, S. S., Dube, W. P., Gilman, J. B., Hall, K., Holloway, J. S., Kuster, W. C., Perring, A. E., Prevot, A. S. H., Schwarz, J. P., Spackman, J. R., Szidat, S., Wagner, N. L., Weber, R. J., Zotter, P., and Parrish, D. D.: Gasoline emissions dominate over diesel in formation of secondary organic aerosol mass, *Geophys Res Lett*, 39, Artn L06805 Doi 10.1029/2011gl050718, 2012.

465 Beccaceci, S., McGhee, E. A., Brown, R. J. C., and Green, D. C.: A comparison between a semi-continuous analyzer and filter-based method for measuring anion and cation concentrations in pm10 at an urban background site in london, *Aerosol Science and Technology*, 49, 793-801, 10.1080/02786826.2015.1073848, 2015.

470 Beever, S. D., Westmoreland, E., de Jong, M. C., Williams, M. L., and Carslaw, D. C.: Trends in nox and no2 emissions from road traffic in great britain, *Atmos Environ*, 54, 107-116, 10.1016/j.atmosenv.2012.02.028, 2012.

Bigi, A., and Harrison, R. M.: Analysis of the air pollution climate at a central urban background site, *Atmos Environ*, 44, 2004-2012, DOI 10.1016/j.atmosenv.2010.02.028, 2010.

475 Bohnenstengel, S. I., Belcher, S. E., Aiken, A., Allan, J. D., Allen, G., Bacak, A., Bannan, T. J., Barlow, J. F., Beddows, D. C. S., Bloss, W. J., Booth, A. M., Chemel, C., Coceal, O., Di Marco, C. F., Dubey, M. K., Faloon, K. H., Fleming, Z. L., Furger, M., Gietl, J. K., Graves, R. R., Green, D. C., Grimmond, C. S. B., Halios, C. H., Hamilton, J. F., Harrison, R. M., Heal, M. R., Heard, D. E., Helfter, C., Herndon, S. C., Holmes, R. E., Hopkins, J. R., Jones, A. M., Kelly, F. J., Kotthaus, S., Langford, B., Lee, J. D., Leigh, R. J., Lewis, A. C., Lidster, R. T., Lopez-Hilfiker, F. D., McQuaid, J. B., Mohr, C., Monks, P. S., Nemitz, E., Ng, N. L., Percival, C. J., Prévôt, A. S. H., Ricketts, H. M. A., Sokhi, R., Stone, D., Thornton, J. A., Tremper, A. H., Valach, A. C., Visser, S., Whalley, L. K., Williams, L. R., Xu, L., Young, D. E., and Zotter, P.: Meteorology, air quality, and health in london: The clearflo project, *Bulletin of the American Meteorological Society*, 10.1175/bams-d-12-00245.1, 2014.

480 Bozzetti, C., Daellenbach, K. R., Hueglin, C., Fermo, P., Sciare, J., Kasper-Giebl, A., Mazar, Y., Abbaszade, G., El Kazzi, M., Gonzalez, R., Shuster-Meiseles, T., Flasch, M., Wolf, R., Křepelová, A., Canonaco, F., Schnelle-Kreis, J., Slowik, J. G., Zimmermann, R., Rudich, Y., Baltensperger, U., El Haddad, I., and Prévôt, A. S. H.: Size-resolved identification, characterization, and quantification of primary biological organic aerosol at a european rural site, *Environmental Science and Technology*, 50, 3425-3434, 10.1021/acs.est.5b05960, 2016.

485 Bruns, E. A., El Haddad, I., Keller, A., Klein, F., Kumar, N. K., Pieber, S. M., Corbin, J. C., Slowik, J. G., Brune, W. H., Baltensperger, U., and Prévôt, A. S. H.: Inter-comparison of laboratory smog chamber and flow reactor systems on organic aerosol yield and composition, *Atmos Meas Tech*, 8, 2315-2332, 10.5194/amt-8-2315-2015, 2015.

490 Budisulistiorini, S. H., Canagaratna, M. R., Croteau, P. L., Baumann, K., Edgerton, E. S., Kollman, M. S., Ng, N. L., Verma, V., Shaw, S. L., Knipping, E. M., Worsnop, D. R., Jayne, J. T., Weber, R. J., and Surratt, J. D.: Intercomparison of an aerosol chemical speciation monitor (acsm) with ambient fine aerosol measurements in downtown atlanta, georgia, *Atmos. Meas. Tech. Discuss.*, 6, 11181-11213, 10.5194/amtd-6-11181-2013, 2013.

495 Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prevot, A. S. H.: Sofi, an igor-based interface for the efficient use of the generalized multilinear engine (me-2) for the source apportionment: Me-2 application to aerosol mass spectrometer data, *Atmos Meas Tech*, 6, 3649-3661, DOI 10.5194/amt-6-3649-2013, 2013.

Canonaco, F., Slowik, J. G., Baltensperger, U., and Prévôt, A. S. H.: Seasonal differences in oxygenated organic aerosol composition: Implications for emissions sources and factor analysis, *Atmos Chem Phys*, 15, 6993-7002, 10.5194/acp-15-6993-2015, 2015.

500 Carslaw, D. C., Williams, M. L., Tate, J. E., and Beever, S. D.: The importance of high vehicle power for passenger car emissions, *Atmos Environ*, 68, 8-16, DOI 10.1016/j.atmosenv.2012.11.033, 2013.

505 Crippa, M., DeCarlo, P. F., Slowik, J. G., Mohr, C., Heringa, M. F., Chirico, R., Poulain, L., Freutel, F., Sciare, J., Cozic, J., Di Marco, C. F., Elsasser, M., Nicolas, J. B., Marchand, N., Abidi, E., Wiedensohler, A., Drewnick, F., Schneider, J., Borrmann, S., Nemitz, E., Zimmermann, R., Jaffrezo, J. L., Prevot, A. S. H., and Baltensperger, U.: Wintertime aerosol chemical composition and source apportionment of the organic fraction in the metropolitan area of paris, *Atmos Chem Phys*, 13, 961-981, DOI 10.5194/acp-13-961-2013, 2013a.

Crippa, M., El Haddad, I., Slowik, J. G., DeCarlo, P. F., Mohr, C., Heringa, M. F., Chirico, R., Marchand, N., Sciare, J., Baltensperger, U., and Prevot, A. S. H.: Identification of marine and continental aerosol sources in paris using

high resolution aerosol mass spectrometry, *J Geophys Res-Atmos*, 118, 1950-1963, Doi 10.1002/Jgrd.50151, 2013b.

510 Crippa, M., Canonaco, F., Lanz, V. A., Aijala, M., Allan, J. D., Carbone, S., Capes, G., Ceburnis, D., Dall'Osto, M., Day, D. A., DeCarlo, P. F., Ehn, M., Eriksson, A., Freney, E., Ruiz, L. H., Hillamo, R., Jimenez, J. L., Junninen, H., Kiendler-Scharr, A., Kortelainen, A. M., Kulmala, M., Laaksonen, A., Mensah, A., Mohr, C., Nemitz, E., O'Dowd, C., Ovadnevaite, J., Pandis, S. N., Petaja, T., Poulain, L., Saarikoski, S., Sellegri, K., Swietlicki, E., Tiitta, P., Worsnop, D. R., Baltensperger, U., and Prevot, A. S. H.: Organic aerosol components derived from 25 ams data sets across europe using a consistent me-2 based source apportionment approach, *Atmos Chem Phys*, 14, 6159-6176, DOI 10.5194/acp-14-6159-2014, 2014.

515 Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., Diskin, G. S., Fisher, J. A., Fuelberg, H. E., Hecobian, A., Knapp, D. J., Mikoviny, T., Riemer, D., Sachse, G. W., Sessions, W., Weber, R. J., Weinheimer, A. J., Wisthaler, A., and Jimenez, J. L.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos Chem Phys*, 11, 12049-12064, DOI 10.5194/acp-11-12049-2011, 2011.

520 Dall'Osto, M., Thorpe, A., Beddows, D. C. S., Harrison, R. M., Barlow, J. F., Dunbar, T., Williams, P. I., and Coe, H.: Remarkable dynamics of nanoparticles in the urban atmosphere, *Atmos Chem Phys*, 11, 6623-6637, DOI 10.5194/acp-11-6623-2011, 2011.

525 Digest of united kingdom energy statistics 2015 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/450302/DUKES_2015.pdf, access: 12/08/2016, 2015.

530 DeWitt, H. L., Hellebust, S., Temime-Roussel, B., Ravier, S., Polo, L., Jacob, V., Buisson, C., Charron, A., André, M., Pasquier, A., Besombes, J. L., Jaffrezo, J. L., Wortham, H., and Marchand, N.: Near-highway aerosol and gas-phase measurements in a high-diesel environment, *Atmos. Chem. Phys.*, 15, 4373-4387, 10.5194/acp-15-4373-2015, 2015.

535 Elser, M., Huang, R. J., Wolf, R., Slowik, J. G., Wang, Q., Canonaco, F., Li, G., Bozzetti, C., Daellenbach, K. R., Huang, Y., Zhang, R., Li, Z., Cao, J., Baltensperger, U., El-Haddad, I., and André, P.: New insights into pm2.5 chemical composition and sources in two major cities in china during extreme haze events using aerosol mass spectrometry, *Atmos Chem Phys*, 16, 3207-3225, 10.5194/acp-16-3207-2016, 2016.

540 Fountoukis, C., Megaritis, A. G., Skyllakou, K., Charalampidis, P. E., Van Der Gon, H. A. C. D., Crippa, M., Prévôt, A. S. H., Fachinger, F., Wiedensohler, A., Pilinis, C., and Pandis, S. N.: Simulating the formation of carbonaceous aerosol in a european megacity (paris) during the megapoli summer and winter campaigns, *Atmos Chem Phys*, 16, 3727-3741, 10.5194/acp-16-3727-2016, 2016.

545 Fröhlich, R., Crenn, V., Setyan, A., Belis, C. A., Canonaco, F., Favez, O., Riffault, V., Slowik, J. G., Aas, W., Aijälä, M., Alastuey, A., Artiñano, B., Bonnaire, N., Bozzetti, C., Bressi, M., Carbone, C., Coz, E., Croteau, P. L., Cubison, M. J., Esser-Gietl, J. K., Green, D. C., Gros, V., Heikkinen, L., Herrmann, H., Jayne, J. T., Lunder, C. R., Minguillón, M. C., Močnik, G., O'Dowd, C. D., Ovadnevaite, J., Petralia, E., Poulain, L., Priestman, M., Ripoll, A., Sarda-Estève, R., Wiedensohler, A., Baltensperger, U., Sciare, J., and Prévôt, A. S. H.: Actris acsm intercomparison – part 2: Intercomparison of me-2 organic source apportionment results from 15 individual, co-located aerosol mass spectrometers, *Atmos. Meas. Tech. Discuss.*, 8, 1559-1613, 10.5194/amtd-8-1559-2015, 2015.

550 Vehicles statistics: <https://www.gov.uk/government/collections/vehicles-statistics>, access: 04/08/2015, 2015.

555 Harrison, R. M., Laxen, D., Moorcroft, S., and Laxen, K.: Processes affecting concentrations of fine particulate matter (pm2.5) in the uk atmosphere, *Atmos Environ*, 46, 115-124, DOI 10.1016/j.atmosenv.2011.10.028, 2012.

560 Hennigan, C. J., Miracolo, M. A., Engelhart, G. J., May, A. A., Presto, A. A., Lee, T., Sullivan, A. P., McMeeking, G. R., Coe, H., Wold, C. E., Hao, W. M., Gilman, J. B., Kuster, W. C., de Gouw, J., Schichtel, B. A., Collett Jr, J. L., Kreidenweis, S. M., and Robinson, A. L.: Chemical and physical transformations of organic aerosol from the photo-oxidation of open biomass burning emissions in an environmental chamber, *Atmos. Chem. Phys.*, 11, 7669-7686, 10.5194/acp-11-7669-2011, 2011.

Henry, R. C., Lewis, C. W., Hopke, P. K., and Williamson, H. J.: Review of receptor model fundamentals, *Atmospheric Environment (1967)*, 18, 1507-1515, [http://dx.doi.org/10.1016/0004-6981\(84\)90375-5](http://dx.doi.org/10.1016/0004-6981(84)90375-5), 1984.

Heo, J., de Foy, B., Olson, M. R., Pakbin, P., Sioutas, C., and Schauer, J. J.: Impact of regional transport on the anthropogenic and biogenic secondary organic aerosols in the los angeles basin, *Atmos Environ*, 103, 171-179, DOI 10.1016/j.atmosenv.2014.12.041, 2015.

Hildebrandt, L., Kostenidou, E., Lanz, V. A., Prevot, A. S. H., Baltensperger, U., Mihalopoulos, N., Laaksonen, A., Donahue, N. M., and Pandis, S. N.: Sources and atmospheric processing of organic aerosol in the mediterranean:

- Insights from aerosol mass spectrometer factor analysis, *Atmos Chem Phys*, 11, 12499-12515, DOI 10.5194/acp-11-12499-2011, 2011.
- 565 Huang, X. F., He, L. Y., Hu, M., Canagaratna, M. R., Sun, Y., Zhang, Q., Zhu, T., Xue, L., Zeng, L. W., Liu, X. G., Zhang, Y. H., Jayne, J. T., Ng, N. L., and Worsnop, D. R.: Highly time-resolved chemical characterization of atmospheric submicron particles during 2008 beijing olympic games using an aerodyne high-resolution aerosol mass spectrometer, *Atmos Chem Phys*, 10, 8933-8945, DOI 10.5194/acp-10-8933-2010, 2010.
- 570 Huffman, J. A., Docherty, K. S., Mohr, C., Cubison, M. J., Ulbrich, I. M., Ziemann, P. J., Onasch, T. B., and Jimenez, J. L.: Chemically-resolved volatility measurements of organic aerosol from different sources, *Environmental Science & Technology*, 43, 5351-5357, 10.1021/es803539d, 2009.
- Jimenez, J., Canagaratna, M., Donahue, N., Prevot, A., Zhang, Q., Kroll, J., DeCarlo, P., Allan, J., Coe, H., and Ng, N.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525-1529, 2009.
- 575 Jones, A. M., and Harrison, R. M.: Interpretation of particulate elemental and organic carbon concentrations at rural, urban and kerbside sites, *Atmos Environ*, 39, 7114-7126, 10.1016/j.atmosenv.2005.08.017, 2005.
- Jones, A. M., Harrison, R. M., and Baker, J.: The wind speed dependence of the concentrations of airborne particulate matter and nox, *Atmos Environ*, 44, 1682-1690, DOI 10.1016/j.atmosenv.2010.01.007, 2010.
- Keyword, M., Cope, M., Meyer, C. P. M., Iinuma, Y., and Emmerson, K.: When smoke comes to town: The impact of biomass burning smoke on air quality, *Atmos Environ*,
580 <http://dx.doi.org/10.1016/j.atmosenv.2015.03.050>, 2015.
- Kupiainen, K., and Klimont, Z.: Primary emissions of fine carbonaceous particles in europe, *Atmos Environ*, 41, 2156-2170, <http://dx.doi.org/10.1016/j.atmosenv.2006.10.066>, 2007.
- London atmospheric emissions inventory 2013: <http://data.london.gov.uk/dataset/london-atmospheric-emissions-inventory-2013>, access: 18/08/2016, 2013.
- 585 Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prevot, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, *Atmos Chem Phys*, 7, 1503-1522, 2007.
- Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., Szidat, S., Wehrli, M. N., Wacker, L., Weimer, S., Caseiro, A., Puxbaum, H., and Prevot, A. S. H.: Source attribution of submicron organic aerosols
590 during wintertime inversions by advanced factor analysis of aerosol mass spectra, *Environmental Science & Technology*, 42, 214-220, Doi 10.1021/Es0707207, 2008.
- Liu, D., Allan, J., Corris, B., Flynn, M., Andrews, E., Ogren, J., Beswick, K., Bower, K., Burgess, R., Choularton, T., Dorsey, J., Morgan, W., Williams, P. I., and Coe, H.: Carbonaceous aerosols contributed by traffic and solid fuel
595 burning at a polluted rural site in northwestern england, *Atmos Chem Phys*, 11, 1603-1619, DOI 10.5194/acp-11-1603-2011, 2011.
- Liu, Q., Sun, Y., Hu, B., Liu, Z. R., Akio, S., and Wang, Y. S.: In situ measurement of pm1 organic aerosol in beijing winter using a high-resolution aerosol mass spectrometer, *Chinese Sci Bull*, 57, 819-826, DOI 10.1007/s11434-011-4886-0, 2012.
- 600 Liu, Y. J., and Harrison, R. M.: Properties of coarse particles in the atmosphere of the united kingdom, *Atmos Environ*, 45, 3267-3276, DOI 10.1016/j.atmosenv.2011.03.039, 2011.
- Lough, G. C., Schauer, J. J., and Lawson, D. R.: Day-of-week trends in carbonaceous aerosol composition in the urban atmosphere, *Atmos Environ*, 40, 4137-4149, <http://dx.doi.org/10.1016/j.atmosenv.2006.03.009>, 2006.
- 605 May, A. A., Nguyen, N. T., Presto, A. A., Gordon, T. D., Lipsky, E. M., Karve, M., Gutierrez, A., Robertson, W. H., Zhang, M., Brandow, C., Chang, O., Chen, S., Cicero-Fernandez, P., Dinkins, L., Fuentes, M., Huang, S.-M., Ling, R., Long, J., Maddox, C., Massetti, J., McCauley, E., Miguel, A., Na, K., Ong, R., Pang, Y., Rieger, P., Sax, T., Truong, T., Vo, T., Chattopadhyay, S., Maldonado, H., Maricq, M. M., and Robinson, A. L.: Gas- and particle-phase primary emissions from in-use, on-road gasoline and diesel vehicles, *Atmos Environ*, 88, 247-260, <http://dx.doi.org/10.1016/j.atmosenv.2014.01.046>, 2014.
- 610 Milic, A., Miljevic, B., Alroe, J., Mallet, M., Canonaco, F., Prevot, A. S. H., and Ristovski, Z. D.: The ambient aerosol characterization during the prescribed bushfire season in brisbane 2013, *Science of The Total Environment*, 560-561, 225-232, <http://dx.doi.org/10.1016/j.scitotenv.2016.04.036>, 2016.
- Minguillón, M. C., Brines, M., Pérez, N., Reche, C., Pandolfi, M., Fonseca, A. S., Amato, F., Alastuey, A., Lyasota, A., Codina, B., Lee, H. K., Eun, H. R., Ahn, K. H., and Querol, X.: New particle formation at ground level and in the vertical column over the barcelona area, *Atmos Res*, 164-165, 118-130, 10.1016/j.atmosres.2015.05.003, 2015a.
- 615 Minguillón, M. C., Ripoll, A., Pérez, N., Prévôt, A. S. H., Canonaco, F., Querol, X., and Alastuey, A.: Chemical characterization of submicron regional background aerosols in the western mediterranean using an aerosol chemical speciation monitor, *Atmos. Chem. Phys.*, 15, 6379-6391, 10.5194/acp-15-6379-2015, 2015b.

620 Mohr, C., Huffman, J. A., Cubison, M. J., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of primary organic aerosol emissions from meat cooking, trash burning, and motor vehicles with high-resolution aerosol mass spectrometry and comparison with ambient and chamber observations, *Environmental Science & Technology*, 43, 2443-2449, Doi 10.1021/Es8011518, 2009.

625 Mohr, C., DeCarlo, P. F., Heringa, M. F., Chirico, R., Slowik, J. G., Richter, R., Reche, C., Alastuey, A., Querol, X., Seco, R., Penuelas, J., Jimenez, J. L., Crippa, M., Zimmermann, R., Baltensperger, U., and Prevot, A. S. H.: Identification and quantification of organic aerosol from cooking and other sources in barcelona using aerosol mass spectrometer data, *Atmos Chem Phys*, 12, 1649-1665, DOI 10.5194/acp-12-1649-2012, 2012.

Morgan, W. T., Allan, J. D., Bower, K. N., Highwood, E. J., Liu, D., McMeeking, G. R., Northway, M. J., Williams, P. I., Krejci, R., and Coe, H.: Airborne measurements of the spatial distribution of aerosol chemical composition across europe and evolution of the organic fraction, *Atmos Chem Phys*, 10, 4065-4083, DOI 10.5194/acp-10-4065-2010, 2010.

630 National atmospheric emissions inventory: <http://naei.defra.gov.uk/data/>, access: 15/08/2016, 2016.

Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prevot, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in northern hemispheric datasets from aerosol mass spectrometry, *Atmos Chem Phys*, 10, 4625-4641, DOI 10.5194/acp-10-4625-2010, 2010.

635 Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., and Sun, Y.: An aerosol chemical speciation monitor (acsm) for routine monitoring of the composition and mass concentrations of ambient aerosol, *Aerosol Science and Technology*, 45, 780-794, 2011.

640 Ortega, A. M., Day, D. A., Cubison, M. J., Brune, W. H., Bon, D., de Gouw, J. A., and Jimenez, J. L.: Secondary organic aerosol formation and primary organic aerosol oxidation from biomass-burning smoke in a flow reactor during flame-3, *Atmos. Chem. Phys.*, 13, 11551-11571, 10.5194/acp-13-11551-2013, 2013.

Paatero, P., and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics*, 5, 111-126, 1994.

645 Paatero, P.: The multilinear engine: A table-driven, least squares program for solving multilinear problems, including the n-way parallel factor analysis model, *J Comput Graph Stat*, 8, 854-888, 10.2307/1390831, 1999.

Paatero, P., Hopke, P. K., Song, X. H., and Ramadan, Z.: Understanding and controlling rotations in factor analytic models, *Chemometr Intell Lab*, 60, 253-264, Doi 10.1016/S0169-7439(01)00200-3, 2002.

Paatero, P., and Hopke, P. K.: Rotational tools for factor analytic models, *Journal of Chemometrics*, 23, 91-100, 2009.

650 Petit, J. E., Favez, O., Sciare, J., Canonaco, F., Croteau, P., Močnik, G., Jayne, J., Worsnop, D., and Leoz-Garziandia, E.: Submicron aerosol source apportionment of wintertime pollution in paris, france by double positive matrix factorization (pmf2) using an aerosol chemical speciation monitor (acsm) and a multi-wavelength aethalometer, *Atmos. Chem. Phys.*, 14, 13773-13787, 10.5194/acp-14-13773-2014, 2014.

655 Petit, J. E., Favez, O., Sciare, J., Crenn, V., Sarda-Estève, R., Bonnaire, N., Močnik, G., Dupont, J. C., Haeffelin, M., and Leoz-Garziandia, E.: Two years of near real-time chemical composition of submicron aerosols in the region of paris using an aerosol chemical speciation monitor (acsm) and a multi-wavelength aethalometer, *Atmos Chem Phys*, 15, 2985-3005, 10.5194/acp-15-2985-2015, 2015.

Pope III, C. A., and Dockery, D. W.: Health effects of fine particulate air pollution: Lines that connect, *J Air Waste Manage*, 56, 709-742, 2006.

660 Ramgolam, K., Favez, O., Cachier, H., Gaudichet, A., Marano, F., Martinon, L., and Baeza-Squiban, A.: Size-partitioning of an urban aerosol to identify particle determinants involved in the proinflammatory response induced in airway epithelial cells, *Part Fibre Toxicol*, 6, 2009.

Ripoll, A., Minguillón, M. C., Pey, J., Jimenez, J. L., Day, D. A., Sosedova, Y., Canonaco, F., Prévôt, A. S. H., Querol, X., and Alastuey, A.: Long-term real-time chemical characterization of submicron aerosols at montsec (southern pyrenees, 1570 m a.S.L.), *Atmos Chem Phys*, 15, 2935-2951, 10.5194/acp-15-2935-2015, 2015.


665 Saarikoski, S., Carbone, S., Decesari, S., Giulianelli, L., Angelini, F., Canagaratna, M., Ng, N. L., Trimborn, A., Facchini, M. C., Fuzzi, S., Hillamo, R., and Worsnop, D.: Chemical characterization of springtime submicrometer aerosol in po valley, italy, *Atmos Chem Phys*, 12, 8401-8421, DOI 10.5194/acp-12-8401-2012, 2012.

670 Sun, Y. L., Wang, Z. F., Fu, P. Q., Yang, T., Jiang, Q., Dong, H. B., Li, J., and Jia, J. J.: Aerosol composition, sources and processes during wintertime in beijing, china, *Atmos Chem Phys*, 13, 4577-4592, DOI 10.5194/acp-13-4577-2013, 2013.

- Takahama, S., Schwartz, R. E., Russell, L. M., Macdonald, A. M., Sharma, S., and Leitch, W. R.: Organic functional groups in aerosol particles from burning and non-burning forest emissions at a high-elevation mountain site, *Atmos. Chem. Phys.*, 11, 6367-6386, 10.5194/acp-11-6367-2011, 2011.
- 675 Tao, L., and Harley, R. A.: Changes in fine particulate matter measurement methods and ambient concentrations in california, *Atmos Environ*, 98, 676-684, <http://dx.doi.org/10.1016/j.atmosenv.2014.09.044>, 2014.
- Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from positive matrix factorization of aerosol mass spectrometric data, *Atmos Chem Phys*, 9, 2891-2918, 2009.
- 680 Vakkari, V., Kerminen, V. M., Beukes, J. P., Tiitta, P., Van Zyl, P. G., Josipovic, M., Venter, A. D., Jaars, K., Worsnop, D. R., Kulmala, M., and Laakso, L.: Rapid changes in biomass burning aerosols by atmospheric oxidation, *Geophys Res Lett*, 41, 2644-2651, 10.1002/2014gl059396, 2014.
- Watson, J. G.: Visibility: Science and regulation, *J Air Waste Manage*, 52, 628-713, 10.1080/10473289.2002.10470813, 2002.
- 685 Weimer, S., Alfarra, M. R., Schreiber, D., Mohr, M., Prévôt, A. S. H., and Baltensperger, U.: Organic aerosol mass spectral signatures from wood-burning emissions: Influence of burning conditions and wood type, *Journal of Geophysical Research: Atmospheres*, 113, n/a-n/a, 10.1029/2007jd009309, 2008.
- Young, D. E., Allan, J. D., Williams, P. I., Green, D. C., Flynn, M. J., Harrison, R. M., Yin, J., Gallagher, M. W., and Coe, H.: Investigating the annual behaviour of submicron secondary inorganic and organic aerosols in london, *Atmos. Chem. Phys.*, 15, 6351-6366, 10.5194/acp-15-6351-2015, 2015a.
- 690 Young, D. E., Allan, J. D., Williams, P. I., Green, D. C., Harrison, R. M., Yin, J., Flynn, M. J., Gallagher, M. W., and Coe, H.: Investigating a two-component model of solid fuel organic aerosol in london: Processes, pm1 contributions, and seasonality, *Atmos Chem Phys*, 15, 2429-2443, 10.5194/acp-15-2429-2015, 2015b.
- Zhang, Q., Worsnop, D. R., Canagaratna, M. R., and Jimenez, J. L.: Hydrocarbon-like and oxygenated organic aerosols in pittsburgh: Insights into sources and processes of organic aerosols, *Atmos Chem Phys*, 5, 3289-3311, 2005.
- 695 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced northern hemisphere midlatitudes, *Geophys Res Lett*, 34, Artn L13801 Doi 10.1029/2007gl029979, 2007.
- 700 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: A review, *Analytical and bioanalytical chemistry*, 401, 3045-3067, 2011.
- 705

Table 1: Sets of target profiles used in the study.

a	c	s	w
BBOA	SFOA	HOA	SFOA
HOA	HOA	COA	HOA
COA	COA	SVOOA	COA



720 **Figure 4:** f44 vs f43 (a) and f44 vs f60 (b) plots for different periods of time.




Figure 5: WD/WE ratios to analyse petrol and diesel contributions.




Figure 6: Daily $PM_{2.5}$ concentrations (a) daily (b) and total PM_1 composition (c), purple line in Fig. 6.c separates secondary and primary aerosols.

725