

Overview of mercury dry deposition, litterfall, and throughfall studies

L. Paige Wright¹, Leiming Zhang², Frank J. Marsik³

¹Independent Researcher, Stratford, Prince Edward Island, Canada

²Air Quality Research Division, Science and Technology Branch, Environment and Climate Change Canada

³Department of Climate and Space Sciences and Engineering, University of Michigan, U.S.

Correspondence to: leiming.zhang@canada.ca

A manuscript prepared for the special issue of *Atmospheric Chemistry Physics*

“Data collection, analysis and application for speciated atmospheric mercury”

Abstract. The current knowledge concerning mercury dry deposition is reviewed, including dry deposition algorithms used in chemical transport models (CTMs) and at monitoring sites and related deposition calculations, measurement methods and studies for quantifying dry deposition of gaseous oxidized mercury (GOM) and particulate bound mercury (PBM), and measurement studies of litterfall and throughfall mercury. Measured median GOM plus PBM dry deposition in Asia ($10.7 \mu\text{g m}^{-2} \text{yr}^{-1}$) is almost double that in North America ($6.1 \mu\text{g m}^{-2} \text{yr}^{-1}$) due to the higher anthropogenic emissions in Asia. The measured mean GOM plus PBM dry deposition in Asia ($22.7 \mu\text{g m}^{-2} \text{yr}^{-1}$), however, is less than that in North America ($30.8 \mu\text{g m}^{-2} \text{yr}^{-1}$). The variations between the median and mean values reflect the influences that single extreme measurements can have on the mean of a dataset. Measured median litterfall and throughfall mercury are 34.8 and $49.0 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively, in Asia, 12.8 and $16.3 \mu\text{g m}^{-2} \text{yr}^{-1}$ in Europe, and 11.9 and $7.0 \mu\text{g m}^{-2} \text{yr}^{-1}$ in North America. The corresponding measured mean litterfall and throughfall mercury are 42.8 and $43.5 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively, in Asia, 14.2 and $19.0 \mu\text{g m}^{-2} \text{yr}^{-1}$ in Europe, and 12.9 and $9.3 \mu\text{g m}^{-2} \text{yr}^{-1}$ in North America. The much higher litterfall mercury than GOM plus PBM dry deposition suggests the important contribution of gaseous elemental mercury (GEM) to mercury dry deposition to vegetated canopies. Over all the regions, including the Amazon, dry deposition, estimated as the sum of litterfall and throughfall minus open-field wet deposition, is more dominant than wet deposition for Hg deposition. Regardless of the measurement or modelling method used, a factor of two or larger uncertainties in GOM plus PBM dry deposition need to be kept in mind when using these numbers for mercury impact studies.

1. Introduction

Atmospheric mercury (Hg) exists in three operationally defined forms - gaseous elemental mercury (GEM), gaseous oxidized mercury (GOM), and particulate bound mercury (PBM). Quantification of atmospheric deposition of the three forms of Hg to various underlying surfaces is needed to assess Hg effects on sensitive ecosystems and subsequent human health impact. Wet deposition of Hg can be quantified through analysis of precipitation chemistry (Prestbo and Gay, 2009). While the direct measurement of Hg dry deposition is technically challenging, a number of methods have been developed for approximate Hg dry deposition, such as micrometeorological, surrogate surface, litterfall, and throughfall measurements (Cobbett and Van Heyst, 2007; Marsik et al., 2007; Graydon et al., 2008; Huang and Gustin, 2015; Risch et al., 2012; Ma et al., 2016).

The approaches used by the above methods to measure mercury dry deposition vary considerably. Efforts are now underway to obtain a better understanding of the variability of estimates from these methods, in an effort to provide a constraint of measurement uncertainties (Marsik et al., 2009; Gustin et al., 2015). The relationships between some of these methods were briefly discussed in Zhang et al. (2012b) and are further illustrated below. Mercury dry deposited to a forest canopy-soil system includes uptake of Hg by leaf stomata and cuticle, tree bark, and underlying soil. Some of the deposited Hg may emit back into the atmosphere while some may be translocated to the branches, stems, and roots. Mercury in litterfall includes a portion of the dry deposited Hg as well as capture of Hg emitted from the soil, although it may not account for the mercury that has been translocated. Mercury in throughfall includes wet deposited Hg above the canopy and a portion of dry deposited Hg washed off from the canopy. Thus, litterfall Hg

likely represents the low-end of Hg dry deposition if Hg emission from the underlying soil is small, although it can be higher than the actual dry deposition above the canopy due to the interception of emitted Hg by forest leaves if soil Hg emissions are high and the ambient Hg concentrations above the forests are low. The difference between throughfall and open-area wet deposition, with the former collected inside and the latter outside a forest, should also represent a portion of the dry deposition. Thus, on annual basis, dry deposition can be approximated as the sum of litterfall and throughfall Hg minus wet deposited Hg.

In chemical transport models (CTMs) and at monitoring sites, Hg dry deposition is commonly calculated using the inferential method, where it is the product of the dry deposition velocity and the atmospheric Hg concentration (Fulkerson, 2006; Lyman et al., 2007; Marsik et al., 2007; Engle et al., 2010; Huang et al., 2012; Zhang et al., 2012b), or using the bi-directional exchange scheme for GEM (Xu et al., 1999; Bash, 2010; Wang et al., 2014; Wright and Zhang, 2015). Development and validation of these schemes need field-measured Hg dry deposition data, which unfortunately have larger uncertainties. The magnitudes of uncertainties are expected to be similar between measurement and model estimates (Zhang et al., 2009; Fang et al., 2012b).

The present study provides a detailed review of current knowledge concerning Hg dry deposition including numerical schemes used in CTMs and at monitoring sites, available field measurements on Hg dry deposition, and litterfall and throughfall Hg measurements. Knowledge gained from this study provides guidance for future research directions.

2. Dry deposition algorithms and estimation

2.1 In chemical transport models

There are several global and regional chemical transport models from Canada, the United States,

66 Europe, Russia, and Asia that include algorithms for the dry deposition of mercury. A summary
67 of the various models' schemes and their algorithms is provided in Table S1. In chemical
68 transport models, the approach for the calculation of the dry deposition velocities of GEM,
69 GOM, and PBM varies between models. For example, in GRAHM (Dastoor et al., 2015), GEOS-
70 Chem (Song et al., 2015), ECHMERIT (De Simone et al., 2014), and GNAQPMS-Hg (Chen et
71 al., 2015), the dry deposition velocity for GEM is calculated using the multiple resistance
72 analogy approach described in Wesely (1989), Zhang et al. (2003) or Kerkweg et al. (2006). In
73 the case of many regional models, GEM is still not considered in the simulations under the
74 assumption that the dry deposition of GEM is not important considering it is also emitted from
75 the surface, for example WRF/Chem-Hg (Gencarelli et al., 2015), TEAM (Seigneur et al., 2006),
76 REMSAD (Bullock et al., 2008), and CAMx (De Foy et al., 2014). CMAQ has implemented the
77 bi-directional scheme of Bash (2010) for its treatment of GEM (Bash et al., 2014). Wang et al.
78 (2014) have recently completed a sensitivity analysis of an updated bi-directional air-surface
79 exchange scheme for GEM. A similar resistance analogy for GEM is used to calculate GOM, for
80 example in GRAHM, GEOS-Chem, and many regional models. In other models, such as
81 WRF/Chem-Hg and CTM-Hg, the dry deposition velocity of GOM is assumed equal to HNO_3
82 due to their similar solubilities. A number of modelling studies have calculated the dry
83 deposition velocity of PBM using the size-segregated particle dry deposition model described in
84 Zhang et al. (2001b) or an earlier dry deposition and gravitational approach by Slinn and Slinn
85 (1980), including Han et al. (2008); Zhang et al. (2012a); De Simone et al. (2014); and Dastoor
86 et al. (2015). Two other approaches are to set the deposition velocity equal to the deposition
87 velocity of the relatively non-reactive sulphate (Bullock and Brehme, 2002; Christensen et al.,

2004; Chen et al., 2015) or to assume specific values over land (0.1 cm s^{-1}) and water (0.01 cm s^{-1}) (Seigneur et al., 2001; 2004).

The largest major inter-model comparison study, the Intercomparison Study of Numerical Models for Long-Range Atmospheric Transport of Mercury, involved eight regional, hemispheric, and global models that compared both short and long term simulations over Europe. This inter-comparison took place between 2000 and 2005 and was conducted in three phases. The first was an examination of the physical and chemical schemes of Hg (Ryaboshapko et al., 2002); the second phase compared short-term simulations (Ryaboshapko et al., 2007a); and the third phase included simulations over an entire year (Ryaboshapko et al., 2007b). In this latter study, the annual model results conducted over the full year (1999) of the dry deposition of Hg were within $\pm 50\%$ of the observations; however, the range of the monthly modelled dry deposition results varied by up to $\pm 100\%$ from observations. The large discrepancies observed were attributed to two factors noted earlier, that not all of the models included the dry deposition of GEM and that some models used a fixed deposition velocity over forests for GOM (Ryabshapko et al., 2007b).

The North American Mercury Model Intercomparison Study involved three regional-scale models (CMAQ, REMSAD, and TEAM) running simulations over North America using the same initial/boundary conditions (Bullock et al., 2008; 2009). In this study, the dry deposition results varied between the models by close to a factor of ten, with REMSAD producing the smallest dry deposition fluxes. Only TEAM simulated dry deposition of GEM, however the GEM results were not reported due to the assumption that most of the GEM dry deposited was re-emitted back to the atmosphere (Bullock et al., 2008).

More recently, the Hemispheric Transport of Air Pollution (HTAP) modelling experiment, included inter-comparisons between GRAHM, GEOS-Chem, GLEMOS, CMAQ-Hg, and ECHMERIT (Travnikov et al., 2010). Hg dry deposition varied between the models, with GEOS-Chem simulating dry deposition fluxes between $\sim 5 \mu\text{g m}^{-2} \text{yr}^{-1}$ over the Arctic to almost $30 \mu\text{g m}^{-2} \text{yr}^{-1}$ over East Asia. For comparison, CTM-Hg fluxes ranged between $\sim 2 \mu\text{g m}^{-2} \text{yr}^{-1}$ over the Pacific and $\sim 15 \mu\text{g m}^{-2} \text{yr}^{-1}$ over Europe and North Africa while GRAHM dry deposition fluxes ranged between $\sim 2 \mu\text{g m}^{-2} \text{yr}^{-1}$ over the Pacific and $\sim 12 \mu\text{g m}^{-2} \text{yr}^{-1}$ over East Asia. Over all of the regions, GLEMOS produced the smallest dry deposition fluxes ranging between $\sim 1 \mu\text{g m}^{-2} \text{yr}^{-1}$ over the Pacific to $\sim 8 \mu\text{g m}^{-2} \text{yr}^{-1}$ over East Asia. In this study, mercury deposition results reflected the different model parameterizations. For example, GRAHM simulated higher mercury deposition in the Polar Regions due to the inclusion of AMDE events, whereas the incorporation of sea salt aerosol scavenging resulted in higher deposition by GEOS-Chem over oceans (Travnikov et al., 2010). The dry deposition varied by more than an order of magnitude over Europe/North Africa, North America, East Asia, and South Asia. Therefore, Travnikov et al. (2010) recommends the need for more measurements of dry deposition in these regions.

In a regional model study by Zhang et al. (2012a), speciated mercury was modeled over the Great Lakes using CMAQ2002, CMAQ2005, and GRAHM2005. The differences between the estimations of dry deposition of GOM and PBM in the CMAQ models and GRAHM at the regional scale were significant (up to a factor of two). For all three models, the differences between modelled and measured GOM and PBM concentrations were between factors of two and ten, whereas the GEM concentrations were within 30%.

As part of the Global Mercury Assessment (GMA), in 2013, three models (GLEMOS, GEOS-Chem, and GMHG) were used to explore global mercury deposition (Travnikov et al., 2015). These models simulated higher deposition fluxes over industrial sites and regions with higher emissions, similar to trends observed in measurements over China (Fu et al., 2010a). The largest contributions to dry deposition fluxes of GOM and PBM were over South Asia, East Asia, and Europe. All three models included AMDE events where the highest dry deposition fluxes in the Arctic were produced in the spring months. The focus of Travnikov et al. (2015)'s study was more to use the three models as an ensemble to examine the global circulation of mercury so the discussion was not on the differences between the models. However, individual results from each model were included in the report. Examples of differences between the models include the relative contribution of domestic and foreign anthropogenic sources to mercury deposition over different regions for 2013. All three models produced the highest contributions in East Asia from domestic anthropogenic sources and in the Arctic from foreign anthropogenic sources. Differences between the models were observed for predicted deposition in Europe, where GLEMOS and GEOS-Chem attributed the largest contribution to domestic sources (>30% and 20%, respectively), whereas GMHG had foreign sources as the major contributor. The other region with differences was South Asia, where again GLEMOS and GEOS-Chem had domestic sources as the larger contributor and GMHG had foreign sources as the major contributor.

Challenges still exist in modelling speciated mercury deposition in chemical transport models. These uncertainties come from uncertainties in model-simulated GOM and PBM concentrations (Baker and Bash, 2012; Holloway et al., 2012; Zhang et al., 2012a, Kos et al.,

2013), uncertainties in existing dry deposition algorithms for GOM and PBM, and the lack of proper dry deposition algorithms for GEM. A better treatment of some of the processes simulated in these models, based on new knowledge generated from analysis of large data sets of recently collected speciated atmospheric data (Cheng et al., 2014; 2015), may improve model performance in simulating GOM and PBM concentrations. For example, the fixed gas-particle partitioning of oxidized Hg in some models (e.g., GRAHM) can be replaced with temperature-dependant parameterizations described in Cheng et al. (2014). The coarse PBM fraction neglected in many models (Zhang et al., 2012a) should also be addressed since coarse PBM contributes significantly to Hg dry and wet deposition budget (Fang et al., 2012b; Cheng et al., 2015). Validation and implementation of updated bi-directional air-surface exchange schemes into chemistry transport models, such as the one for GEM by Wang et al. (2014), will be beneficial in reducing these uncertainties.

The limited amount of measurement data inhibits the development and improvement of chemical transport models. While speciated mercury data has been increasing rapidly in the recent decade through enhanced monitoring networks (Gay et al., 2013) and increases in field studies (Lindberg and Stratton, 1998; Kim et al., 2009; Maestas et al., 2011; Fu et al., 2015; Lee et al., 2016), there is no practical way for direct dry deposition measurement as further illustrated in Section 3. Besides, the monitored oxidized Hg concentration data, especially GOM, likely have large uncertainties (Jaffe et al., 2014; Gustin et al., 2015) and caution should be made when applying this data for model evaluation (Kos et al., 2013).

2.2 At monitoring networks/sites

An increase in the monitoring of speciated atmospheric Hg in the past decade has led to

an increase in the number of dry deposition estimations using the inferential method. There are several monitoring networks that have been established to monitor speciated atmospheric Hg. The largest network to date is the National Atmospheric Deposition (NADP)/Atmospheric Mercury Network (AMNet), which includes more than 30 sites mostly in the United States (Gay et al., 2013). Other networks include the European Monitoring and Evaluation Programme (EMEP) (Travnikov and Ilyin, 2009), the Arctic Monitoring and Assessment Program (AMAP) (Travnikov et al., 2015), and the Global Mercury Observation System (GMOS) (Sprovieri et al., 2013). There have been several monitoring sites which have performed year-long measurement campaigns of Hg dry deposition of GOM and PBM, such as over ten sites in Japan (Sakata and Marumoto, 2004; 2005; Sakata and Asakura, 2008) and a number in Taiwan (Fang et al., 2012a, b; 2013; 2014; 2016). A comprehensive review of atmospheric GEM and TGM mercury measurements, sites, and trends across the globe can be found in the literature (e.g. Sprovieri et al., 2010). The majority of the Hg dry deposition studies using the inferential approach and speciated ambient concentrations (referred to as modelled deposition below) were conducted for GOM and PBM (Lindberg and Stratton, 1998; Fulkerson, 2006; Lyman et al., 2007, 2009; Marsik et al., 2007; Han et al., 2008; Engle et al., 2010; Lombard et al., 2011; Castro et al., 2012; Huang et al., 2012; Petersen et al., 2012; Sather et al., 2013; Huang and Gustin, 2015; Fang et al., 2016), while a few studies also included GEM (Rea et al., 2001; Miller et al., 2005; Caldwell et al., 2006; Holmes et al., 2011; Zhang et al., 2012b; Bieser et al., 2014; Enrico et al., 2016). A summary of estimated dry deposition fluxes for GOM and PBM estimated using the data from the various monitoring networks and other field campaigns is provided in Table S2.

Dry deposition flux estimations of GOM in North America range from $0.02 \text{ ng m}^{-2} \text{ hr}^{-1}$ to

5.91 ng m⁻² hr⁻¹ with one exception in Augusta, Georgia, where the maximum estimated GOM dry deposition flux was 36.3 ng m⁻² hr⁻¹ over vents of a mercury chlor-alkali plant (Landis et al., 2004). In China, the estimation is higher at 20.4 ng m⁻² hr⁻¹ (Huang et al., 2012). Dry deposition flux estimations of PBM range from 0.003 ng m⁻² hr⁻¹ to 4.54 ng m⁻² hr⁻¹ in North America and from 0.43 ng m⁻² hr⁻¹ to 46.46 ng m⁻² hr⁻¹ in East Asia. In Europe, the median of estimated GOM+PBM dry deposition fluxes is 0.34 ng m⁻² hr⁻¹. In most studies, dry deposition of GOM has been estimated to be much higher than PBM due to the much higher deposition velocities of GOM than PBM (Han et al., 2008; Engle et al., 2010; Huang et al., 2012; Zhang et al., 2012b). Another reason that needs to be emphasized is the exclusion of coarse PBM concentration in the monitoring process in many studies (e.g., Zhang et al., 2012b). Differences between the modelled and measured fluxes are typically at a factor of two or larger due to measurement uncertainties and choice of model parameters (Lyman et al., 2007; Marsik et al., 2007; Huang and Gustin, 2015), although some studies obtained very close model-measurement values (Miller et al., 2005; Lyman et al., 2009; Zhang et al., 2012b).

Dry deposition is often assumed to be less important than wet deposition since earlier studies only estimated GOM and PBM dry deposition. Using speciated Hg data at 19 monitoring sites in North America, Zhang et al. (2012b) estimated speciated and total Hg dry deposition and also used litterfall Hg measurements at the regional scale as a constraint for model estimation. In the study of Zhang et al. (2012b), GEM was demonstrated to contribute much more than GOM+PBM to the total dry deposition over vegetated surface. Thus, total dry deposition, when including GEM, was much higher than wet deposition at the majority of the sites, a conclusion that can be supported from litterfall (Risch et al., 2012) and somewhat by throughfall

measurements (see more discussion in Sections 5 and 6).

3. Deposition (flux), litterfall, and throughfall measurement methodologies

3.1. Micrometeorological approaches

The most common micrometeorological approach employed for the measurement of the air-surface exchange of mercury is the flux-gradient approach. In a common application, time-averaged fluxes of speciated mercury (typically reported in either nanograms (or picograms) per meter squared per hour) are obtained using the relation:

$$F_{Hg,i} = -K_z \frac{\Delta C_i}{\Delta Z}$$

where $F_{Hg,i}$ is the flux of a given mercury species, i , K_z is the vertical eddy-transfer coefficient and $\frac{\Delta C_i}{\Delta Z}$ is the time-averaged vertical gradient of the mercury species, in question. Several approaches have been used to determine the values of the stability-dependent eddy-transfer coefficient K_z , including the aerodynamic approach (e.g., Edwards, 2005; Marsik et al., 2005; Fritsche et al., 2008) and the modified Bowen-ratio approach (e.g., Kim et al., 1995; Lindberg et al., 1995; Meyers et al., 1996; Poissant et al. 2004; Fritsche et al., 2008). In each case, the assumption is made that all scalar quantities are mixed in a similar fashion within the atmospheric surface layer, and thus the vertical eddy-transfer coefficient for heat (or other tracers, such as CO₂ or H₂O_v) can be used as a surrogate for the vertical eddy-transfer coefficient for mercury.

The vertical gradient of mercury has been measured using either: (a) manual approaches (such as gold-bead traps for GEM, KCl-coated denuders for GOM and quartz filters for PBM) or (b) automated sampling systems (such as a Tekran® Model 2537A Mercury Vapor Analyzer combined with a Tekran® Model 1110 Synchronized Two-Port Sampling Unit for GEM, a

Tekran® Model 1130 Mercury Speciation Unit for GOM, combined with a Tekran® Model 1135 Particulate Mercury Unit for PBM).

3.2. Dynamic gas flux chambers

Dynamic gas chambers, which have been used to study the flux of a number of gaseous species from a variety of emitting surfaces, including leaves, soil and water, have been successfully employed for use in the measurement of the air-surface exchange of GEM (e.g., Poissant and Casimir, 1998; Zhang et al., 2001a; Wallschläger et al., 2002; Marsik et al., 2005; Eckley et al., 2011; Edwards and Howard, 2013). In a typical configuration for the measurement of the air-surface exchange of GEM from soil or water surfaces, a polycarbonate flux chamber (DFC) is placed on the emitting surface to be measured. Ambient air is drawn by vacuum through the DFC at a constant flow rate, nominally 1.5 LPM. The concentration of GEM is measured at both the inlet and outlet of the DFC, typically using an automated system, such as a Tekran® Model 2537A Mercury Vapor Analyzer combined with a Tekran® Model 1110 Synchronized Two-Port Sampling Unit, though initial approaches used manual gold bead traps. The resulting GEM flux (typically reported in nanograms per meter squared per hour) is calculated using the relation:

$$F_{GEM} = \left[\frac{(C_o - C_i) \cdot Q}{A} \right]$$

where C_o is the concentration of GEM at the outlet, C_i is the concentration of GEM at the inlet, Q is the volume flow rate through the DFC and A is the area of the footprint of the DFC on the surface in question. As with other approaches, the final flux is determined using the above equation, subtracting a sample blank value from measured value. Typically, DFC blanks are obtained at the start and completion of a given sampling period by placing the DFC over a clean,

inert surface (e.g., Aluminum foil).

3.3. Surrogate surface approaches

A number of studies considered in this manuscript have utilized surrogate surface measurement approaches to provide a direct measurement of mercury to the Earth's surface. The designs of the various surrogate surfaces tend to favor the quantification of one form of mercury over the others, while not being able to totally rule out a contribution from one of the other forms. Intercomparison studies have suggested that the different approaches provide relatively consistent results (Marsik, 2009; Gustin et al., 2015), though care must be taken when interpreting surrogate surface results due to the differences in surrogate surface designs.

There are a variety of surrogate surface approaches that have been used to directly measure the dry deposition of mercury. The general classes of approaches can be categorized in the following manner: (a) water/solution-based surfaces, (b) filter-based surfaces, (c) membrane-based surfaces and (d) turf-based surrogate surfaces. A brief discussion of each of these approaches is presented below.

a. Water/solution-based surfaces

Water- or solution-based surrogate surfaces have been used by a number of groups (Sakata and Marumoto, 2005; Marsik et al., 2007; Lai et al., 2011) for the quantification of the dry deposition of mercury. In general, these approaches deploy a given volume of ultra-pure water in a reservoir that is part of an aerodynamic sampler design. Following a specified period of deployment, the sample solutions are retrieved and the amount of mercury within the sample solution is quantified. In most cases, the final results are "bias-corrected", meaning that the average amount of mercury found in a subset of unexposed solutions is subtracted from the

exposed solution to obtain a net amount of mercury which is assumed to be due to that dry deposited to the exposed solution. Some approaches utilize a static amount of solution which is deployed in a fixed reservoir within the sample holder (Marsik et al., 2007; Lai et al., 2011). Other water/solution-based systems approaches use a more dynamic approach for the measurement of the dry deposition of mercury and other species, in which the sample solution is continuously-circulated through the sampling apparatus, with the exposed surface area remaining constant through the sample period (e.g., Sakata and Marumoto, 2004).

In general, surrogate water surface approaches using “ultra-pure” water are considered to collect both GOM and PBM, with limited uptake expected from GEM due to the relatively insolubility of this latter species. In most cases, the resultant deposition flux of mercury (in picograms per square meter per hour) is determined by

$$Deposition\ flux = \left[\frac{(Hg_{Sample} - Hg_{Blank})}{A \cdot T} \right]$$

where Hg_{Sample} is the amount of mercury in the sample solution, Hg_{Blank} is the amount of mercury in the sample blank, A is the exposed surface area of the solution and T is the length of time over which the sample was exposed to the ambient air.

It should be noted that some approaches use slightly acidified solutions for the purpose of “fixing” deposited mercury to avoid volatilization after deposition to the exposed solution. It is possible that such dilute-acid solutions may lead to an enhanced uptake of GEM due to the fixation of any GEM which may even temporarily become dissolved in the solution.

b. Filter-based surfaces

A number of filter-based approaches have been used in the collection of mercury dry-deposition (Lai et al., 2011). The general concept of this approach is that the three-dimensional structure of the filters will lead to a capture of GOM and PBM. Like the water/solution-based approaches, the filter surfaces are exposed for a given amount of time and then returned to the analytical laboratory at the end of the sampler period, with subsequent quantification of the amount of mercury deposited to the surface, followed by a bias-correction for the average amount of mercury found in a subset of unexposed filters. As before, the mercury dry-deposition flux is quantified as:

$$Deposition\ flux = \left[\frac{(Hg_{Sample} - Hg_{Blank})}{A \cdot T} \right]$$

where Hg_{Sample} is the amount of mercury on the exposed sample filter, Hg_{Blank} is the amount of mercury on the sample filter blank, A is the exposed surface area of the filter and T is the length of time over which the sample was exposed to the ambient air.

While some of the filter-based approaches utilize the filter surfaces as received from the manufacturer, some approaches coat the filter surfaces either with a solution (that is subsequently allowed to dry) or, in some cases, with a fine film of gold. In these latter cases, the goal is to enhance the collection of GEM mercury that would otherwise be missed by the other filter-based approaches.

c. Membrane-based surfaces

One of the more recently developed approaches uses cation exchange membranes as the exposed deposition surfaces (e.g., Lyman et al., 2007; Castro et al., 2012; Sather et al., 2013; Huang and Gustin, 2015). These membranes have been designed to specifically favor the uptake of GOM over either GEM or PBM, using the assumption that the deposition of mercury is dominated by the former species. This class of filters is typically deployed facing downward, which serves to reduce contamination by large particles and by precipitation. An example of one such filter (Pall Corporation, P/N ICD45S3R) was employed by Lyman *et al.* (2007) and is constructed out of negatively-charged polysulfane on one side, with a non-woven polyester backing on the other. As before, the resultant deposition flux of mercury (in picograms per square meter per hour) is determined by

$$Deposition\ flux = \left[\frac{(Hg_{Sample} - Hg_{Blank})}{A \cdot T} \right]$$

where Hg_{Sample} is the amount of mercury on the exposed sample filter, Hg_{Blank} is the amount of mercury on the sample filter blank, A is the exposed surface area of the filter and T is the length of time over which the sample was exposed to the ambient air. The inverted sampler design employed with cation exchange membrane samplers protects the membranes from contamination by rainfall, thus allowing these samplers to be deployed for long periods of time and thus providing a long-term integrated measure of Hg dry deposition.

d. Turf Surrogate Samplers

Much in the way that cation-exchange membrane surfaces were developed in a way that protects them from contamination by rainfall, the turf surrogate samples (TSS) approach was

developed to allow for dry deposition measurements to be performed in such a way that the samplers could remain deployed during both dry- and precipitating-periods (Marsik et al., 2009; Lynam et al., 2015), thus allowing its application for long-term (days to week) quantification of Hg dry deposition. In addition to reducing the amount of person-hours required to cover surrogate surfaces in the advent of precipitation, this approach also sought to characterize dry-deposition to wetted surfaces which can occur following instances of dewfall or rainfall. TSS utilize circular disks of artificial turf (e.g., polyethylene Astro Turf®, GrassWorx, LLC, St. Louis, MO), which serve as three-dimensional deposition surfaces. The turf has approximately 2-cm high polyethylene “blades of grass” which are placed into a Teflon well-style insert (such as that used in the surrogate water-based samplers noted above) that is part of an aerodynamic airfoil. The backing of the turf has a lattice-like construction which allows precipitation to flow through the turf surface into a “throughfall” sample bottle which is included as part of the sample train.

For the period over which the TSS was exposed to ambient air, the estimated dry deposition of mercury is calculated as:

$$\text{Dry Deposition Flux} = Hg \text{ flux}_{TSS} + Hg \text{ flux}_{TF} - Hg \text{ flux}_{WO}$$

where $Hg \text{ flux}_{TSS}$ is the flux of mercury to the TSS over the period of exposure, $Hg \text{ flux}_{TF}$ is the flux of mercury captured within the throughfall bottle and $Hg \text{ flux}_{WO}$ is the flux of mercury captured with a collocated wet-only precipitation sampler. In this case, the wet-only measurement is used to remove the wet-deposited mercury flux from the throughfall

measurement, which would contain a combination of both wet- and dry-deposited mercury. All components of the TSS system are acid-cleaned prior to deployment, with field blanks collected for the circular pieces of turf, the throughfall samples and the wet-only precipitation samples. The appropriate corrections are then made to the data in the determination of the final estimation of the measured flux.

3.4. Litterfall and throughfall-based approaches

There are several methods for collecting litterfall: manual sampling of leaves using gloves (Roulet et al., 1998; 1999; Kalicin et al., 2008; Michelazzo et al., 2010; Ding et al., 2011; Juillerat et al., 2012; Obrist et al., 2011; 2012; Melendez-Perez et al., 2014; Jiskra et al., 2015); nylon nets attached to aluminium frames placed a certain distance from the ground (Iverfeldt et al., 1991; Munthe et al., 1995; Lee et al., 2000; Nóvoa-Muñoz et al., 2008; da Silva et al., 2009; Wang et al., 2009; Fisher and Wolfe, 2012); litterfall traps/collectors (Rea et al., 1996; Grigal et al., 2000; St. Louis et al., 2001; Johnson, 2002; Sheehan et al., 2006; Silva-Filho et al., 2006; Bushey et al., 2008; Graydon et al., 2008; Larssen et al., 2008; Selvendiran et al., 2008; Fu et al., 2010a; Risch et al., 2012; Zhou et al., 2013); funnel and net combinations (Schwesig and Matzner 2000; 2001); and the use of litterbags (Hall and St. Louis, 2004; Demers et al., 2007; Pokharel and Obrist, 2011). Fresh foliage was cut directly from trees and shrubs manually from inside branches and outside branches (McLaughlin et al., 2008); from the crowns of the trees (Mélières et al., 2003; Michelazzo et al., 2010; Blackwell and Driscoll, 2015b; Fostier et al., 2015; Luo et al., 2015); middle branches (Rasmussen et al., 1991; 1995; Rea et al., 2002; Richardson and Friedland, 2015); lower tree branches (Michelazzo et al., 2010; Obrist et al., 2012; Gong et al., 2014; Hutnik et al., 2014; Fostier et al., 2015); or shot down with a pellet gun

(Bushey et al., 2008; Blackwell and Driscoll, 2015a).

Mercury that is dry deposited to leaves, needle surfaces, branches, and trunks gets washed off during precipitation events. This Hg-enriched rain then lands on the surface and becomes throughfall (Rea et al., 2000; 2001; Mélières et al., 2003; da Silva et al., 2009; Risch et al., 2012). The Hg in the throughfall is then either adsorbed by the litter or forest floor or re-volatized (Demers et al., 2007). In addition to dry deposition to the leaf/needle surface, foliar leaching was thought to be a possible source of Hg in throughfall but was found to be negligible (Rea et al., 2001). Throughfall Hg is calculated by multiplying the volume-mean-weighted Hg concentration in throughfall by the total throughfall volume. In throughfall field studies, samplers are collected weekly (Choi et al., 2008; Wang et al., 2009; Fu et al., 2010a; 2010b; Luo et al., 2015), bi-weekly (Kolka et al., 1999; Grigal et al., 2000; Schwesig and Matzner, 2000; Witt et al., 2009; Åkerblom et al., 2015); monthly (Munthe et al., 1995; Fostier et al., 2000; Lee et al., 2000; Larssen et al., 2008); on an event basis (Rea et al., 2001; Demers et al., 2007; Kalicin et al., 2008); or the funnels are covered between precipitation events to avoid contamination by dry deposition of Hg on the funnels (Rea et al., 1996). Lengths of campaign measurements vary between a single growing season (Munthe et al., 1995; Rea et al., 1996; 2001; Fostier et al., 2000) to up to a five-year period (Lee et al., 2000; Porvari and Verta, 2003), with the current majority of throughfall studies having been for a one-year length (Larssen et al., 2008; Wang et al., 2009; Fu et al., 2010a; 2010b; Fisher and Wolfe, 2012; Åkerblom et al., 2015; Blackwell and Driscoll, 2015b; Luo et al., 2015). The time schedule for throughfall collection is important as there is the opportunity for coniferous forests to scavenge atmospheric Hg outside of the growing season if the field studies last for longer than just the growing season (Fisher et al., 2012). The

sampling frequency can affect the measured throughfall Hg as there is the possibility of contamination by Hg dry deposition occurring if the samplers are not covered during extensive dry periods (Rea et al., 1996). Over long-term studies however, sample-to-sample throughfall Hg concentrations may vary on a smaller timescale (i.e. one month) using different sampling frequencies but the cumulative Hg throughfall deposition will be statistically insignificant (Choi et al., 2008).

4. Oxidized mercury flux measurements

Measurements of GEM flux have been reviewed in Zhu et al. (2016) so this section only focuses on GOM and PBM. The majority of the dry deposition flux measurement campaigns for GOM and PBM (88%) have used surrogate surface techniques, with 67% of them being at sites in North America and 33% of them at sites in Asia. These measurement studies have included several long-term campaigns of up to three years in length in both North America (Lyman et al., 2009; Castro et al., 2012; Peterson et al., 2012; Weiss-Penzias et al., 2012; Sather et al., 2013; 2014; Wright et al., 2014; Huang and Gustin, 2015) and Asia (Sakata and Asakura, 2008; Fang et al., 2016; Han et al., 2016). Current knowledge on particle size distribution of PBM, although limited, has started to be studied in more depth in recent years (Keeler et al., 1995; Ames et al., 1998; Feddersen et al., 2012; Kim et al., 2012). Although PBM mass distributes more in fine ($PM_{2.5}$) than coarse particles ($>2.5 \mu m$), dry deposition from fine and coarse PBM are equally important due to the faster deposition velocity of coarse particles (Fang et al., 2012a). An increase in North American PBM studies would be beneficial in further exploration of these trends. For a detailed summary of the measured dry deposition fluxes of GOM and PBM ($ng\ m^{-2}\ hr^{-1}$), as well as the measured concentrations of GOM and PBM ($pg\ m^{-3}$), see Table S3.

In North America, the median and mean values are 6.1 and $30.8 \mu\text{g m}^{-2} \text{yr}^{-1}$ with a range between 0.26 and $520 \mu\text{g m}^{-2} \text{yr}^{-1}$. Long term studies have shown a stronger dependence on GOM dry deposition (40-57%) to the total Hg deposition over arid regions, such as the Four Corners and Nevada (Caldwell et al., 2006; Lyman et al., 2007; Sather et al., 2014) than in more humid regions such as Texas and Oklahoma (5-27%) and Florida (Lyman et al., 2007; Marsik et al., 2007; Peterson and Gustin, 2008; Weiss-Penzias et al., 2011; 2012; Sather et al., 2014). Higher GOM and PBM concentrations and dry deposition fluxes have been observed in the close proximity to, or downwind from, point source emissions (Huang et al., 2012; Sather et al., 2014) than at rural sites (Castro et al., 2012; Sather et al., 2014). GOM concentration and dry deposition have shown positive correlations to wind speed (Lyman et al., 2010; Gustin et al., 2012; Sather et al., 2014; Huang and Gustin, 2015) and higher elevations (Huang and Gustin, 2015). Diel trends in GOM and PBM have shown higher concentrations in the morning and afternoon (Lindberg and Stratton, 1998; Poissant et al., 2004; Liu et al., 2007; Peterson and Gustin, 2008; Lai et al., 2011; Weiss-Penzias et al., 2011; Gustin et al., 2012; Fang et al., 2013; Fu et al., 2016) and then a decrease at night, potentially due to a reduction in photochemical production of GOM at night (Weiss-Penzias et al., 2011) or due to nighttime inversion (Huang and Gustin, 2015). The seasonal changes in the fluxes depend on geography and site characteristics where, in general, the fluctuations in the GOM and PBM dry deposition seasonal variations have been observed to be more pronounced in arid regions. Higher GOM deposition has been observed in the spring due to GOM events and long-range transport (Liu et al., 2007; Lyman et al., 2009; Weiss-Penzias et al., 2011; Castro et al., 2012; Gustin et al., 2012; Peterson et al., 2012; Sather et al., 2013; Huang and Gustin, 2015).

There are some physical aspects of the various measurement techniques that have been observed to affect measured values. The direction that the filter in the cation-exchange membrane is facing can alter amount of deposited particles, with lower measured dry deposition fluxes of GOM and PBM in downward facing membranes (Lyman et al., 2007; 2009; Weiss-Penzias et al., 2011; Gustin et al., 2012; Huang et al., 2012; Peterson et al., 2012; Sather et al., 2013; 2014; Huang and Gustin, 2015) than upward facing (Huang et al., 2012). In addition to direction of the membrane, higher deposition rates have been observed for rectangular mounts than aerodynamic mounts, possibly due to artificial turbulence created by the rectangular mounts (Lyman et al., 2009). The length of deployment times for surrogate surfaces is under debate, with recommendations that a minimum sampling time of two weeks be implemented at rural sites (Huang et al., 2012; Huang and Gustin, 2015) and for passive samplers (Lyman et al., 2010). Improvements and changes in sampling techniques continue. For example, the ICE 450 passive sampler, used in several GOM studies (e.g. Castro et al., 2012; Huang et al., 2012; Sather et al., 2013; 2014), was replaced in July 2013 with the Mustang S passive sampler (Huang and Gustin, 2015), where fortunately no significant differences in the measurement results from the two samplers have been observed (Huang and Gustin, 2015). Clearly, surrogate surfaces may differ significantly from the real-world natural surfaces, and choices of surrogate surfaces and instrument setup can both affect the measured deposition values.

In Asia, dry deposition fluxes of GOM+PBM are higher than in North America where median and mean values are 10.7 and 22.7 $\mu\text{g m}^{-2} \text{yr}^{-1}$, with a range between 2.4 and 463 $\mu\text{g m}^{-2} \text{yr}^{-1}$. Similar to North America, in Asia, GOM and PBM concentrations and dry deposition fluxes have been observed to be up to five times greater at urban and industrial sites due to the close

proximity to, or downwind from, point source emissions (Sakata and Marumoto, 2005; Liu et al., 2007; Sakata and Asakura, 2008; Fang et al., 2012a) than at rural sites (Sakata and Marumoto, 2005; Sakata and Asakura, 2008). At three sites in Taiwan, PBM concentrations were highest in November and December and in the summer due to industry, such as coal-fired power plants, and GOM events (Fu et al., 2016). Higher PBM has been measured in the winter due to increases in the winter in the use of vehicles, biomass burning, electricity consumption such as home heating, or due to an accumulation in PBM because of lower temperatures and solar radiation (Sakata and Asakura, 2008; Fang et al., 2012a; Fu et al., 2016; Han et al., 2016).

5. Litterfall measurements

Litterfall Hg represents Hg collected by and retained in leaves. It has been observed in recent years to be an important pathway for Hg input from the atmosphere to the floor of forest ecosystems via plant litter. Litterfall is comprised of senesced leaves and needles, twigs, and branches, and other plant tissues. Litterfall Hg can be transmitted to the soil as the litter decomposes. Litterfall Hg is determined by the concentration of Hg in the leaf litter (Rea et al., 1996; Sheehan et al., 2006; Silva-Filho et al., 2006; Wang et al., 2009; Niu et al., 2011). Another method for estimating litterfall Hg has been to multiply the Hg concentration in fresh foliage by 1.5 to account for the assumption that the Hg accumulates in the leaf linearly over an average six month growing season (Fu et al., 2010b; Gong et al., 2014). It is important to note that a litterfall deposition value estimated through multiplying the fresh foliage value by 1.5 does have an associated error factor that comes with it. Litterfall measurement campaigns have ranged from the growing season (Rea et al., 1996; 2002; Fostier et al., 2003; Larssen et al., 2008; Ding et al., 2011; Fisher and Wolfe, 2012) to long-term studies of a year (Schwesig and Matzner, 2000;

2001; St. Louis et al., 2001; de Oliveira et al., 2006; Silva-Filho et al., 2006; Sheehan et al., 2006; da Silva et al., 2009; Wang et al., 2009; Fu et al., 2010a; Teixeira et al., 2012; Zhou et al., 2013; Lang, 2014; Luo et al., 2015) to two years or more (Xiao et al., 1998; Lee et al., 2000; Hall and St. Louis, 2004; Graydon et al., 2008; Selvendiran et al., 2008; Obrist et al., 2011; Risch et al., 2012; Hutnik et al., 2014; Blackwell and Driscoll, 2015b; Richardson and Friedland, 2015). A summary of available measurements of Hg concentrations in leaves and litter (ng g^{-1}) and litterfall Hg ($\mu\text{g m}^{-2} \text{yr}^{-1}$) is provided in Table S4.

Litterfall Hg measurements are highest in Asia, where the median and mean values from 20 sites (34.8 and $42.8 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively) are approximately two to three times higher than the measured litterfall Hg in Europe over 14 sites, (12.8 and $14.2 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively) and more than three times higher than in North America over 92 sites (11.9 and $12.9 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively). Litterfall Hg range from $4.2 - 219.9 \mu\text{g m}^{-2} \text{yr}^{-1}$ in Asia to $0.05 - 42 \mu\text{g m}^{-2} \text{yr}^{-1}$ in Europe. In North America, litterfall Hg ranges between 0.02 and $57.1 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively. Measurements of litterfall Hg in the Amazonian rainforest have been reviewed in detail by Fostier et al. (2015). Atmospheric Hg concentrations affect the levels of Hg concentrations in foliage and leaf litter with higher litterfall Hg in urban forests than remote forests (Fostier et al., 2003; 2015; Wang et al., 2009; Fu et al., 2010a; 2010b; Gong et al., 2014); higher Hg concentrations in close proximity to degassing vents and during eruption activity of volcanoes (Martin et al., 2012); and higher Hg concentrations at intermediate altitudinal mountain zones (Szopka et al., 2011). It is important to note, however, that the atmospheric Hg concentration is not the only factor that affects the levels of Hg concentrations in foliage and leaf litter, as discussed below. In Europe, a decrease in litterfall Hg has been associated with the decrease in

Hg emissions in recent years (Larssen et al., 2008).

Due to coniferous needles remaining on trees for longer periods than deciduous trees (years vs. one season), needle litter tends to have higher Hg concentrations than leaf litter (Grigal et al., 2000; Sheehan et al., 2006; Niu et al., 2011; Blackwell and Driscoll, 2015b; Fisher and Wolfe, 2012; Obrist et al., 2012). This, however, does not translate directly to higher litterfall Hg in coniferous forests due to the contribution from the higher mass of leaf litter than needle litter. As a result, litterfall Hg in deciduous forests has been found to be higher than in coniferous forests (Schwesig and Matzner, 2000; Demers et al., 2007; Kalicin et al., 2008; Fisher and Wolfe, 2012; Obrist et al., 2012; Richardson and Friedland, 2015).

Correlations between Hg concentrations and age of the foliage have been observed with concentrations increasing with age (Barghigiani et al., 1991; Roulet et al., 1999; Ericksen et al., 2003; Larssen et al., 2008; Ding et al., 2011; Martin et al., 2012) and higher in litter than foliage (Iverfeldt, 1991; Bushey et al., 2008; Michelazzo et al., 2010; Ding et al., 2011; Melendez-Perez et al., 2014; Luo et al., 2015).

In three urban forests in Asia, annual litterfall Hg deposition was found to account for up to 75% of the total Hg input fluxes to the forests (Wang et al., 2009). The contribution of Hg from litterfall has been found to be three times the amount from throughfall (Fu et al., 2010a) and anywhere from two times up to 14 times the contribution from wet deposition (Wang et al., 2009; Fu et al., 2010a; Zhou et al., 2013). A detailed review of litterfall, throughfall, and precipitation studies in China can be found in Fu et al., 2015. In Europe, litterfall Hg has been found to represent between 27% and 90% to the total deposited Hg (Schwesig and Matzner, 2000; 2001; Jiskra et al., 2015), from equal to 1.5 times the deposited throughfall Hg and two

times the open-field precipitation (Iverfeldt et al., 1991; Munthe et al., 1995; Lee et al., 2000). In the Amazonian rainforest, litterfall Hg has been observed to be twice the contribution from wet deposition (da Silva et al., 2009) and more than doubled during the dry season (Silva-Filho et al., 2006). In North America, some studies have found contributions to the deposition flux from litterfall and throughfall equal (Iverfeldt et al., 1991; Rea et al., 1996; Grigal et al., 2000; Lee et al., 2000), whereas in other studies, litterfall has contributed up to two times more Hg than throughfall (St. Louis et al., 2001; Johnson, 2002; Fisher and Wolfe, 2012).

Median litterfall deposition of $12.3 \mu\text{g m}^{-2} \text{yr}^{-1}$ accounted for 55% of the Hg deposition to deciduous forest floors at 23 sites in eastern USA (Risch et al., 2012). In this study that spanned 15 states over three years, litterfall Hg ranged between 3.5 and $23.4 \mu\text{g m}^{-2} \text{yr}^{-1}$. Litterfall Hg was found to be more correlated to the amount of litter collected (i.e. the sample catch) than Hg wet deposition, with the highest median litterfall Hg deposition ($14.7 \mu\text{g m}^{-2} \text{yr}^{-1}$) found in deciduous forests. Mixed forests and coniferous forests had the same median litterfall Hg concentration (38.6 ng g^{-1}) but the mixed forest litterfall Hg was higher (9.3 vs. $7.0 \mu\text{g m}^{-2} \text{yr}^{-1}$) due to the higher sample catch. Of the deciduous sites, oak-hickory forests had higher litterfall Hg than maple-beech-birch forests and aspen-birch forests. Litterfall Hg deposition was more dependent on the forest cover type than geographical region. Intraseasonal variations were observed with higher litterfall Hg concentrations in the first autumn sampling than the second. Interannual variations were observed for the sample catch but were not significant for litterfall Hg concentrations.

6. Throughfall measurements

Factors that affect throughfall Hg deposition include: canopy type, with higher

concentrations in coniferous canopies than deciduous canopies due to a combination of higher Hg scavenging efficiency in coniferous forests over deciduous forests and larger surface areas (Kolka et al., 1999; Rea et al., 1996; Grigal et al., 2000; Schwesig and Matzner, 2000; 2001; Johnson, 2002; Demers et al., 2007; Wang et al., 2009; Fu et al., 2010a; 2010b; Fisher and Wolfe, 2012; Åkerblom et al., 2015; Luo et al., 2015); canopy density, with Hg throughfall concentrations increasing with canopy density (Grigal et al., 2000; Witt et al., 2009); meteorology, with higher throughfall concentrations under drier conditions, resulting from increased dry deposition to the foliage (Blackwell and Driscoll, 2015b); and location, with higher concentrations in regions with high atmospheric Hg emissions or close to point sources, such as urban forests (Wang et al., 2009; Luo et al., 2015). A summary of available measurements Hg concentrations in throughfall (ng L^{-1}) and throughfall Hg ($\mu\text{g m}^{-2} \text{yr}^{-1}$) is provided in Table S5.

In Asia, median and mean throughfall Hg for this region are 49.0 and $43.5 \mu\text{g m}^{-2} \text{yr}^{-1}$, respectively, with a range in the means between 10.5 and $71.3 \mu\text{g m}^{-2} \text{yr}^{-1}$. These median and mean are more than two to three times the European values (16.3 and $19.0 \mu\text{g m}^{-2} \text{yr}^{-1}$ for the median and mean, respectively), where the range is between 12.0 and $40.1 \mu\text{g m}^{-2} \text{yr}^{-1}$. For throughfall, the median and mean throughfall Hg in Asia are eight and five times larger, respectively, than those reported in North America, likely due to heavier atmospheric Hg loadings in Asia than North America (Fu et al., 2010b). Fu et al. (2010b) found that while the throughfall in remote regions of China were smaller than in urban regions of China, they were still up to two times higher than in remote regions of North America. Industrial and urban pollution, density of the population, and coal combustion and mining have all resulted in China having the highest Hg emissions in the world (Wang et al., 2009; Fu et al., 2010a; Fu et al.,

2015). A modelling study by Seigneur et al. (2003) found that Hg emissions from Asia were the largest contributor outside of the United States to Hg deposition in New York due to long-range transport. This is similar to detailed discussions in Fu et al. (2015) on the effects of trends in increasing Hg emissions in China on the patterns of long-range transport. In North America, the median and mean values are the lowest at 7.0 and 9.3 $\mu\text{g m}^{-2} \text{yr}^{-1}$, respectively, with a range between 2.1 and 26.6 $\mu\text{g m}^{-2} \text{yr}^{-1}$. The only throughfall Hg measurement in South America is an estimate of 72.0 $\mu\text{g m}^{-2} \text{yr}^{-1}$ in the Brazilian Amazon (Fostier et al., 2000), which has been suggested to be an overestimation due to the temporal sampling distribution where higher Hg concentrations were observed in samples that had a several-day dry period immediately prior to sampling. Without any other Amazonian throughfall studies, it is not known whether this is a typical value for this rainforest or if the high value was related to the monthly collection. In Asia and Europe, Hg contributions from throughfall are higher than from wet deposition, with throughfall ranging from approximately two to four times the wet deposition (Munthe et al., 1995, Lee et al., 2000; Fu et al., 2010a; 2010b; Wang et al., 2009). In North America, the throughfall-precipitation ratio has been found to differ with forest type, where based upon throughfall measurements, dry deposition was found to be less than wet deposition in deciduous forests (Demers et al., 2007; Kalicin et al., 2008; Blackwell and Driscoll, 2015b) and greater than precipitation in coniferous forests (Demers et al., 2007; Kalicin et al., 2008; Witt et al., 2009) and over watersheds (Lindberg et al., 1994; Rea et al., 1996; 2001; Grigal et al., 2000; Graydon et al., 2008).

7. A brief comparison of dry, litterfall, throughfall and wet deposition

The annual contributions to Hg deposition from wet deposition, throughfall, litterfall, and

dry deposition ($\mu\text{g m}^{-2} \text{yr}^{-1}$) are shown in Figure 1 for various sites in (a) Asia, (b) Europe, (c) North American deciduous forests, and (d) North American coniferous forests. Figure 1 includes sites for which measurements for litterfall, throughfall, and dry and wet deposition have all been measured. This figure is provided to give a representation of the range of values for the various measurement methods. In these figures, the wet deposition represents open-field wet deposition measurements, with the exception of the measurements at Whiteface Mountain, in the Adirondack Park, NY (Blackwell et al., 2015b), where the wet deposition at this site is from cloud water. The estimated dry deposition shown in this figure is the sum of the litterfall Hg and throughfall Hg minus the wet deposited Hg.

In China, dry deposition accounted for between 25% up to 90% of the total Hg deposited to subtropical forested catchments. Litterfall Hg was the dominant pathway in these forests contributing between 53 and 90% of the dry deposition to these forests. In forested watersheds in Germany, wet deposition played a stronger role, accounting for 78% of the total Hg deposition in Steinkreuz and almost twice as much at Lehstenbach. At the other European sites, wet deposition only represented between 9 and 45% of the total Hg deposition. At all of the European sites, litterfall Hg was the dominant pathway at 64 to 98% of the dry deposited Hg, with the exception of Buskerud, Norway where wet deposition was not reported, preventing an accurate estimate of dry deposition (Larssen et al., 2008). In North America, the dry deposition of Hg ranged between 13 and 83% of the total deposited Hg over deciduous forests and 15 and 70% over coniferous forests, with the primary factor in these variances being proximity to sources. Exceptions to these values were at Whiteface Mountain, as mentioned above, where wet deposition is reported as high elevation cloudwater Hg rather than open-field precipitation. The contributions from net

throughfall Hg (throughfall Hg – wet deposition) were higher in the coniferous forests than the deciduous forests. In the deciduous forests, litterfall Hg accounted for between 46 and 95% of total deposited Hg.

8. Summary and recommendations

The modelling of GOM and PBM dry deposition on regional and global scales using various chemical transport models shows inconsistencies between the models, up to a factor of ten in some cases, which are caused by differences in model simulated GOM and PBM concentrations and in selected dry deposition algorithms. Recent analysis of large data sets of speciated atmospheric mercury has provided some needed knowledge for improving these models. However, measurement uncertainties in ambient GOM and PBM concentrations, and the significant lack of measurements over the oceans, the Polar Regions, the Southern Hemisphere, Europe, and Asia are inhibiting further evaluation of the models. Progress has been made on the inferential estimation of speciated and total Hg dry deposition at monitoring sites or networks. GEM deposition to forest canopies is more important than previously assumed in chemical transport models. Further development of dry deposition algorithms, especially the bi-directional exchange for GEM, is still needed, but reliable Hg flux data is lacking in validating such models.

Surrogate surfaces are the current preferred method for measuring the dry deposition of GOM and PBM, with a new turf surrogate surface showing promise in extending the length of the sampling times to longer campaigns. Uncertainties in the measured GOM and PBM dry deposition associated with the selected surrogate surfaces and instrument setup are larger than a factor of two. Measurements of PBM dry deposition are very limited around the world due to the

challenges associated with making these measurements. Measured GOM and PBM dry deposition are strongly correlated to emissions from local point sources due to their short lifetime in air. GOM deposition can play a significant role in total Hg deposition, contributing to more than half of Hg dry deposition in some regions, and having been found to be higher than wet deposition in some arid regions and lower in more humid regions due to the high solubility of GOM.

There has been a large increase in the number of field campaigns measuring litterfall and throughfall mercury in Asia, Europe, the Amazon, and North America. Generally, litterfall and throughfall Hg are higher in urban regions of Asia as a result of the high levels of atmospheric Hg in Asia, followed by remote regions in Asia, then Europe, and North America. As can be seen in North American field studies, litterfall Hg is more dominant than throughfall Hg over deciduous canopies due to the larger amounts of litter under these canopies, whereas throughfall is the primary pathway for mercury deposition over coniferous canopies. Net throughfall Hg is the throughfall Hg minus the Hg in precipitation. This deposition, combined with the deposition of litterfall Hg is higher than the precipitation Hg over all canopies and regions. The measurement of litterfall, throughfall and open-field precipitation Hg, and the subsequent comparison with modelled Hg dry deposition, is an effective way of improving our understanding of the mechanisms involved in these transfers.

References

- Åkerblom, S., Meili, M., and Bishop, K.: Organic Matter in Rain: An Overlooked Influence on Mercury Deposition, *Environ. Sci. Technol. Lett.*, 2 (4), 128-132, 2015.
- Ames, M., Gullug, G., and Olmez, I.: Atmospheric mercury in the vapor phase, and in fine and coarse particulate matter at Perch River, New York, *Atmos. Environ.*, 32 (5), 865-872, 1998.

685 Barghigiani, C., Ristori, T., and Cortopassi, M. : Air mercury measurement and interference of
686 atmospheric contaminants with gold traps, *Environ. Technol.*, 12 (10), 935-941, 1991.

687 Baker, K. R. and Bash, J. O.: Regional scale photochemical model evaluation of total mercury
688 wet deposition and speciated ambient mercury, *Atmos. Environ.*, 49, 151-162, 2012.

689 Bash, J. O.: Description and initial simulation of a dynamic bidirectional air-surface exchange
690 model for mercury in Community Multiscale Air Quality (CMAQ) model, *J. Geophys.*
691 *Res.*, 115, D06305, doi:10.1029/2009JD012834, 2010.

692 Bash, J. O., Carlton, A. G., Hutzell, W. T., and Bullock Jr., O. R.: Regional Air Quality Model
693 Application of the Aqueous-Phase Photo Reduction of Atmospheric Oxidized Mercury
694 by Dicarboxylic Acids, *Atmosphere*, 5, 1-15, doi:10.3390/atmos5010001, 2014.

695 Bieser, J., De Simone, F., Gencarelli, C., Geyer, B., Hedgecock, I., Matthias, V., Travníkov, O.,
696 and Weigelt, A.: A diagnostic evaluation of modeled mercury wet depositions in Europe
697 using atmospheric speciated high-resolution observations, *Environ. Sci. Pollut. Res.*, 21,
698 9995-10012, 2014.

699 Blackwell, B. D. and Driscoll, C. T.: Using foliar and forest floor mercury concentrations to
700 assess spatial patterns of mercury deposition, *Environ. Pollut.*, 202, 126-134, 2015a.

701 Blackwell, B. D. and Driscoll, C. T.: Deposition of mercury in forests along a montane elevation
702 gradient, *Environ. Sci. Technol.*, 49, 5363-5370, 2015b.

703 Bullock Jr. and Brehme, K. A.: Atmospheric mercury simulation using the CMAQ model:
704 formulation description and analysis of wet deposition results, *Atmos. Environ.*, 36,
705 2135-2146, 2002.

706 Bullock Jr., O. R., Atkinson, D., Braverman, T., Civerolo, K., Dastoor, A., Davignon, D., Ku, J.-
707 Y., Lohman, K., Myers, T. C., Park, R. J., Seigneur, C., Selin, N. E., Sistla, G., and
708 Vijayaraghavan, L.: The North American Mercury Model Intercomparison Study
709 (NAMMIS): Study description and model-to-model comparisons, *J. Geophys. Res.*,
710 113, D17310, doi:10.1029/2008JD009803, 2008.

711 Bullock Jr., O. R., Atkinson, D., Braverman, T., Civerolo, K., Dastoor, A., Davignon, D., Ku, J.-
712 Y., Lohman, K., Myers, T. C., Park, R. J., Seigneur, C., Selin, N. E., Sistla, G., and
713 Vijayaraghavan, L.: An analysis of simulated wet deposition of mercury from the North
714 American Mercury Model Intercomparison Study, *J. Geophys. Res.*, 114, D08301,
715 doi:10.1029/2008JD011224, 2009.

716 Bushey, J. T., Nallana, A. G., Montesdeoca, M. R., and Driscoll, C. T.: Mercury dynamic of a
717 northern hardwood canopy, *Atmos. Environ.*, 42, 6905-6914, 2008.

718 Caldwell, C. A., Swartzendruber, P., and Prestbo, E.: Concentration and dry deposition of
719 mercury species in arid south central New Mexico (2001-2002), *Environ. Sci. Technol.*,
720 40, 7535-7540, 2006.

721 Castro, M. S., Moore, C., Sherwell, J., and Brooks, S. B.: Dry deposition of gaseous oxidized
722 mercury in Western Maryland, *Sci. Total Environ.*, 417-418, 232-240, 2012.

723 Chen, H. S., Wang, Z. F., Li, J., Tang, X., Ge, B. Z., Wu, X. L., Wild, O., and Carmichael, G. R.:
724 GNAQPMS-Hg v1.0, a global nested atmospheric mercury transport model: model
725 description, evaluation and application to trans-boundary transport of Chinese
726 anthropogenic emissions, *Geosci. Model Dev.*, 8, 2857-2876, 2015.

727 Cheng I., Zhang, L., Blanchard, P.: Regression modeling of gas-particle partitioning of
728 atmospheric oxidized mercury from temperature data. *J. Geophys. Res. –Atmos.*, 119,
729 11,864–11, 876, 2014.

730 Cheng I., Zhang, L., and Mao, H.: Relative contributions of gaseous oxidized mercury and fine
731 and coarse particle-bound mercury to mercury wet deposition at nine monitoring sites in
732 North America, *J. Geophys. Res. –Atmos.*, 120, 8549–8562, 2015.

733 Choi, H.-D., Sharac, T. J., Holsen, T. M.: Mercury deposition in the Adirondacks: A comparison
734 between precipitation and throughfall, *Atmos. Environ.*, 42, 1818-1827, 2008.

735 Christensen, J. H., Brandt, J., Frohn, L. M., and Skov, H.: Modelling of mercury in the arctic
736 with the Danish Eulerian Hemispheric Model, *Atmos. Chem. Phys.*, 4, 2251-2257,
737 2004.

738 Cobbett, F. D. and Van Heyst, B. J.: Measurements of GEM fluxes and atmospheric mercury
739 concentrations (GEM, RGM and Hgp) from an agricultural field amended with
740 biosolids in Southern Ont., Canada (October 2004–November 2004), *Atmos. Environ.*,
741 41, 2270–2282, 2007.

742 Da Silva, G. S., Bisinoti, M. C., Fadini, P. S., Magarelli, G., Jardim, W. F., and Fostier, A. H. :
743 Major Aspects of the Mercury Cycle in the Negro River Basin, Amazon, *J. Braz. Chem.*
744 *Soc.*, 20 (6), 1127-1134, 2009.

745 Dastoor, A., Ryzhkov, A., Durnford, D., Lehnher, I., Steffen, A., and Morrison, H.:
746 Atmospheric mercury in the Canadian Arctic. Part II: Insight from modeling *Sci. Total*
747 *Environ.*, 509-520, 16-27, 2015.

748 De Foy, B., Heo, J., and Schauer, J. J. : Estimation of direct emissions and atmospheric
749 processing of reactive mercury using inverse modeling, *Atmos. Environ.*, 85, 73-82,
750 2014.

751 De Simone, F., Gencarelli, C. N., Hedgecock, I. M., and Pirrone, N. : Global atmospheric cycle
752 of mercury: a model study on the impact of oxidation mechanisms, *Environ. Sci. Pollut.*
753 *R.*, 21, 4110–4123, 2014.

754 Demers, J. D., Driscoll, C. T., Fahey, T. J., and Yavitt, J. B.: Mercury cycling in litter and soil in
755 different forest types in the Adirondack Region, New York, USA, *Ecol. Appl.*, 17 (5),
756 1341-1351, 2007.

757 Ding, Z., Wu, H., Feng, X., Liu, J., Liu, Y., Yuan, Y., Zhang, L., Lin, G., and Jiayong, P.:
758 Distribution of Hg in mangrove trees and its implication for Hg enrichment in the
759 mangrove ecosystem, *Appl. Geochem.*, 26, 205-212, 2011.

760 Eckley, C. S., Gustin, M., Marsik, F., and Miller, M. B.: Measurement of surface mercury fluxes
761 at active industrial gold mines in Nevada (USA), *Sci. Total Environ.*, 409, 514-522,

2011.

Edwards, G. C., Rasmussen, P. E., Schroeder, W. H., Wallace, D. M., Halfpenny-Mitchell, L.,
Dias, G. M., Kemp, R. J., and Ausma, S.: Development and evaluation of a
sampling system to determine gaseous Mercury fluxes using an aerodynamic
micrometeorological gradient method, *J. Geophys. Res.*, D10306,
doi:10.1029/2004JD005187, 2005.

Edwards, G. C. and Howard, D. A.: Air-surface exchange measurements of gaseous elemental
mercury over naturally enriched and background terrestrial landscapes in Australia,
Atmos. Chem. Phys., 13, 5325-5336, 2013.

Engle, M. A., Tate, M. T., Krabbenhoft, D. P., Schauer, J. J., Kolker, A., Shanley, J. B., and
Bothner, M. H.: Comparison of atmospheric mercury speciation and deposition at nine
sites across central and eastern North America, *J. Geophys. Res.*, 115, D18306,
doi:10.1029/2010JD014064, 2010.

Enrico, M., Le Roux, G., Maruszczak, N., Heimbürger, L.-E., Claustres, A., Fu, X., Sun, R., and
Sonke, J. E.: Atmospheric mercury transfer to peat bogs dominated by gaseous
elemental mercury dry deposition, *Environ. Sci. Technol.*, 50, 2405-2412.

Ericksen, J. A., Gustin, M. S., Schorran, D. E., Johnson, D. W., Lindberg, S. E., and Coleman, J.
S.: Accumulation of atmospheric mercury in forest foliage, *Atmos. Environ.*, 37, 1613-
1622, 2003.

Fang, G. C., Tsai, J.-H., Lin, Y.-H., and Chang, C.-Y.: Dry deposition of atmospheric particle-
bound mercury in the middle Taiwan, *Aerosol Air Qual. Res.*, 12, 1298-1308, 2012a.

Fang G. C., Zhang, L., Huang, C. S.: Measurements of size-fractionated concentration and bulk
dry deposition of atmospheric particle bound mercury, *Atmos. Environ.*, 61, 371-377,
2012b.

Fang, G. C., Lin, Y. -H., and Chang, C. -Y.: Use of mercury dry deposition samplers to quantify
dry deposition of particulate-bound mercury and reactive gaseous mercury at a traffic
sampling site, *Environ. Forensics*, 14 (3), 182-186, 2013.

Fang, G. C., Lin, Y.-H., Chang, C.-Y., and Zheng, Y.-C.: Concentrations of particulates in
ambient air, gaseous elementary mercury (GEM), and particulate-bound mercury
(Hg(p)) at a traffic sampling site: a study of dry deposition in daytime and nighttime,
Environ. Geochem. Health, 36, 605-612, 2014.

Fang, G. C., Lin, Y.-H., and Zheng, Y.-C.: Ambient air particulates and particulate-bound
mercury Hg(p) concentrations: dry deposition study over a Traffic, Airport, Park
(T.A.P.) areas during years of 2011–2012, *Environ. Geochem. Health*, 38, 183-194,
2016.

Feddersen, D.M., Taibot, R., Mao, H., and Sive, B.C.: Size distribution of particulate mercury in
marine and coastal atmospheres. *Atmos. Chem. Phys.*, 12, 10899-10909, 2012.

Fisher, L. S., and Wolfe, M. H.: Examination of mercury inputs by throughfall and litterfall in
the Great Smoky Mountains National Park, *Atmos. Environ.*, 47, 554-559, 2012.

801 Fostier, A.-H., Forti, M. C., Guimarães, J. R. D., Merlí, A. J., Boulet, R., Espirito Santo, C. M.,
802 and Krug, F. J.: Mercury fluxes in a natural forested Amazonian catchment (Serra do
803 Navio, Amapá State, Brazil) , *Sci. Total Environ.*, 260, 201-211, 2000.

804 Fostier, A. H., Cecon, K., Forti, M. C.: Urban influence on litterfall trace metals fluxes in the
805 Atlantic forest of São Paulo (Brazil), *J. Physique IV France*, 107, 491-494, 2003.

806 Fostier, A. H., Melendez-Perez, J. J., Richter, L.: Litter mercury deposition in the Amazonian
807 rainforest, *Environ. Pollut.*, 206, 605-610, 2015.

808 Fritsche, J., Obrist, D., Zeeman, M. J., Conen, F., Eugster, W. and Alewell, C.: Elemental
809 mercury fluxes over a sub-alpine grassland determined with two-micrometeorological
810 methods, *Atmos. Environ.*, 42, 2922-2933, 2008.

811 Fu, X. W., Feng, X., Dong, Z. Q., Yin, R. S., Wang, J. X., Yang, Z. R., and Zhang, H.:
812 Atmospheric gaseous elemental mercury (GEM) concentrations and depositions at a
813 high altitude mountain peak in south China, *Atmos. Chem. Phys.*, 10, 2425-2437,
814 2010a.

815 Fu, X.W., Feng, X., Zhu, W., Rothenberg, S., Yao, H., Zhang, H.: Elevated atmospheric
816 deposition and dynamics of mercury in a remote upland forest of southwestern China,
817 *Environ. Pollut.*, 158, 2324-2333, 2010b.

818 Fu, X. W., Zhang, H., Yu, B., Wang, X., Lin, C.-J., and Feng, X. B.: Observations of
819 atmospheric mercury in China: a critical review, *Atmos. Chem. Phys.*, 15, 9455–9476,
820 doi:10.5194/acp-15-9455-2015, 2015.

821 Fu, X. W., Maruszczak, N., Heimbürger, L.-E., Sauvage, B., Gheusi, F., Prestbo, E. M., and
822 Sonke, J. E.: Atmospheric mercury speciation dynamics at the high-altitude Pic du Midi
823 Observatory, southern France, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2015-
824 842, 2016.

825 Fulkerson, M.: Atmospheric mercury deposition in an urban environment, PhD. Thesis,
826 University of Central Florida, Florida, 96 pp., 2006.

827 Gay, D. A., Schmeltz D., Prestbo E., Olson M., Sharac T, and Tordon R.: The Atmospheric
828 Mercury Network: measurement and initial examination of an ongoing atmospheric
829 mercury record across North America, *Atmos. Chem. Phys.*, 13, 11339-11349, 2013.

830 Gong, P., Wang, X.-P., Xue, Y.-G., Xu, B.-Q., Yao, T.-D.: Mercury distribution in the foliage
831 and soil profiles of the Tibetan forest: Processes and implications for regional cycling,
832 *Environ. Pollut.*, 188, 94-101, 2014.

833 Graydon, J. A., St. Louis, V. L., Hintelmann, H., Lindberg, S. E., Sandilands, K. A., Rudd, J. W.
834 M., Kelly, C. A., Hall, B. D., and Mowat, L. D.: Long-Term Wet and Dry Deposition of
835 Total and Methyl Mercury in the Remote Boreal Ecoregion of Canada, *Environ. Sci.*
836 *Technol.*, 42, 8345–8351, 2008.

837 Grigal, D. F., Kolk, R. K., Fleck, J. A., and Nater, E. A.: Mercury budget of an upland-peatland
838 watershed, *Biogeochem.*, 50, 95-109, 2000.

839 Gustin, M. S., Weiss-Penzias, P. S., and Peterson, C.: Investigating sources of gaseous oxidized

mercury in dry deposition at three sites across Florida, USA, *Atmos. Chem. Phys.*, 12, 9201-9219, 2012.

Gustin, M. S., H. M. Amos, J. Huang, M. B. Miller, and K. Heidecorn (2015), Measuring and modeling mercury in the atmosphere: a critical review, *Atmos. Chem. Phys.*, 15, 5697-5713, doi:10.5194/acp-15-5697-2015.

Han, Y.-J., Holsen, T. M., Ever, D. C., and Driscoll, C. T.: Reduced mercury deposition in New Hampshire from 1996 to 2002 due to changes in local sources, *Environ. Pollut.*, 156, 1348–1356, 2008.

Han, J.-S., Seo, Y.-S., Kim, M.-K., Holsen, T. M., and Yi, S.-M.: Total atmospheric mercury deposition in forest areas in Korea, *Atmos. Phys. Chem. Discuss.*, doi:10.5194/acp-2016-7, 2016.

Hall, B. D. and St. Louis, V. L.: Methylmercury and total mercury in plant litter decomposing in upland forests and flooded landscapes, *Environ. Sci. Technol.*, 38, 5010-5021, 2004.

Holloway, T., Voigt, C., Morton, J., Spak, S. N., Rutter, A. P., and Schauer, J. J.: An assessment of atmospheric mercury in the Community Multiscale Air Quality (CMAQ) model at an urban site and a rural site in the Great Lakes Region of North America, *Atmos. Chem. Phys.*, 12, 7117–7133, 2012.

Holmes, H. A., Pardyjak, E. R., Perry, K. D., and Abbott, M. L.: Gaseous dry deposition of atmospheric mercury: A comparison of two surface resistance models for deposition to semiarid vegetation, *J. Geophys. Res.*, 116, S14306, doi:10.1029/2010JD015182, 2011.

Huang, J., Choi, H.-D., Landis, M. S., and Holsen, T. M.: An application of passive samplers to understand atmospheric mercury concentration and dry deposition spatial distributions, *J. Environ. Monit.*, 14, 2012.

Huang, J., and Gustin, M. S.: Use of passive sampling methods and models to understand sources of mercury deposition to high elevation sites in the western United States, *Environ. Sci. Technol.*, 49, 432-441, 2015.

Hultberg, H., Munthe, J., and Iverfeldt, Å.: Cycling of methylmercury and mercury – Responses in the forest roof catchment to three years of decreased atmospheric deposition, *Water Air Soil Poll.*, 80, 415-424, 1995.

Hutnik, R. J., McClenahan, J. R., Long, R. P., and Davis, D. D.: Mercury Accumulation in *Pinus nigra* (Austrian Pine), *Northeast. Nat.*, 21 (4), 529-540, 2014.

Iverfeldt, Å.: Mercury in forest canopy throughfall water and its relation to atmospheric deposition, *Water Air Soil Poll.*, 56, 553-564, 1991.

Jaffe, D. A., S. Lyman, H. M. Amos, M. S. Gustin, J. Huang, N. E. Selin, L. Leonard, A. Ter Schure, R. P. Mason, R. Talbot, A. Rutter, B. Finley, L. Jaeglé, V. Shah, C. McClure, J. Ambrose, L. Gratz, S. Lindberg, P. Weiss-Penzias, G.-R. Sheu, D. Feddersen, M. Horvat, A. Dastoor, A. J. Hynes, H. Mao, J. E. Sonke, F. Slemr, J. A. Fisher, R. Ebinghaus, Y. Zhang, and G. Edwards (2014), Progress on understanding atmospheric mercury hampered by uncertain measurements, *Environ. Sci. Technol.*, 48(13), 7204-

879 7206.

880 Jiskra, M., Wiederhold, J. G., Skjellberg U., Kronberg, R.-M., Hadjas, I., and Kretzschmar, R.:
881 Mercury deposition and re-emission pathways in boreal forest soils investigated with
882 Hg isotope signatures, *Environ. Sci. Technol.*, 49, 7188-7196, 2015.

883 Johnson, K. B.: Fire and its effects on mercury and methylmercury dynamics for two watersheds
884 in Acadia National Park, Maine, MSc. Thesis, the University of Maine, Maine, 73 pp.,
885 2002.

886 Juillerat, J. I., Ross, D. S., and Bank, M. S.: Mercury in litterfall and upper soil horizons in
887 forested ecosystems in Vermont, USA, *Environ. Toxicol. Chem.*, 31 (8), 1720-1729,
888 2012.

889 Kalicin, M. H., Driscoll, C. T., Yavitt, J., Newton, R., and Munson, R.: The Dynamics of
890 Mercury in Upland Forests of the Adirondack Region of New York, in: *Mercury in*
891 *Adirondack wetlands, lakes and terrestrial systems (MAWLTS)*, New York State
892 Energy Research and Development Authority, New York, 8-1-8-15, 2008.

893 Keeler, G., Glinsorn, G., and Pirrone, N.: Particulate mercury in the atmosphere: Its significance,
894 transport, transformation and sources, *Water Air Soil Pollut.*, 80 (1-4), 159-168, 1995.

895

896 Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöcke, P.: Technical Note:
897 An implementation of the dry removal processes DRY DEPosition and SEDImentation
898 in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617–4632,
899 2006.

900 Kim, K. H., Lindberg, S. E. and Meyers, T. P.: Micrometeorological measurements of mercury
901 vapor fluxes over background forest soils in Eastern Tennessee, *Atmos. Environ.*, 29,
902 267–282, 1995.

903 Kim, S.-H., Han, Y.-J., Holsen, T. M., Yi, S.-M.: Characteristics of atmospheric speciated
904 mercury concentrations (TGM, Hg(II) and Hg(p)) in Seoul, Korea, *Atmos. Environ.*, 43
905 (10), 3267-3274, 2009.

906 Kim, P.-R., Han, Y.-J., Olsen, T. M., Yi, S.-M.: Atmospheric particulate mercury:
907 Concentrations and size distributions, *Atmos. Environ.*, 61, 94-102, 2012.

908 Kolka, R. K., Nater, E. A., Grigal, D. F., and Verry, E. S.: Atmospheric inputs of mercury and
909 organic carbon into a forested upland/bogwatershed, *Water Air Soil Poll.*, 113, 273-294,
910 1999.

911 Kos, G., Ryzhkov, A., Dastoor, A., Narayan, J., Steffen, A., Ariya, P. A., and Zhang, L.:
912 Evaluation of discrepancy between measured and modelled oxidized mercury species,
913 *Atmos. Chem. Phys.*, 13, 4839–4863, 2013.

914 Lai, S.-O., Huang, J., Hopke, P. K., and Holsen, T. M.: An evaluation of direct measurement
915 techniques for mercury dry deposition, *Sci. Total Environ.*, 409, 1320-1327, 2011.

916 Landis, M. S., Keeler, G. J., Al-Wali, K. I., and Stevens, R. K.: Divalent inorganic reactive

- 917 gaseous mercury emissions from a mercury cell chlor-alkali plant and its impact on
918 near-field atmospheric dry deposition, *Atmos. Environ.*, 38, 613-622, 2004.
- 919 Lang, X.: Mercury in atmospheric precipitation and litterfall in Mt.Ailao and Mt.Damei, MSc
920 Thesis, Guizhou University, Guiyang, China, 78 pp., 2014.
- 921 Larssen, T., de Wit, H. A., Wiker, M., Halse, K., Lei, H., Wuebbles, D. J., Liang, X.-Z., Tao, Z.,
922 Olsen, S., Artz, R., Ren, X., and Cohen, M.: Mercury budget of a small forested boreal
923 catchment in southeast Norway, *Sci. Total Environ.*, 404, 290-296, 2008.
- 924 Lee, Y. H., Bishop, K. H., Munthe, J., Iverfeldt, Å., Verta, M., Parkman, H., and Hultberg, H.:
925 An Examination of Current Hg Deposition and Export in Fenno-Scandian Catchments,
926 *Biogeochemistry*, 40 (2/3), 125-135, 1998.
- 927 Lee, Y. H., Bishop, K. H., and Munthe, J.: Do concepts about catchment cycling of
928 methylmercury and mercury in boreal catchments stand the test of time? Six years of
929 atmospheric inputs and runoff export at Svartberget, northern Sweden, *Sci. Total*
930 *Environ.*, 260, 11-20, 2000.
- 931 Lee, G.-S., Kim, P.-R., Han, Y.-J., Holsen, T. M., Seo, Y.-S., and Yi, S.-M.: Atmospheric
932 speciated mercury concentrations on an island between China and Korea: sources and
933 transport pathways, *Atmos. Chem. Phys.*, 16, 4119–4133, doi:10.5194/acp-16-4119-
934 2016, 2016.
- 935 Lindberg, S. E., Owens, J. G., and Stratton, W. J.: Application of throughfall methods to estimate
936 dry deposition of mercury, in: *Mercury as a global pollutant*, Huckabee, J. and Watras,
937 C. (Eds.), Lewis Publications, 261-272, 1994.
- 938 Lindberg, S. E., Kim, K.-H., Meyers, T. P., and Owens, J. G.: A micrometeorological gradient
939 approach for quantifying air/surface exchange of mercury vapor: Tests over
940 contaminated soils, *Environ. Sci. Technol.*, 29, 126-135, 1995.
- 941 Lindberg, S. E. and Stratton, W. J.: Atmospheric mercury speciation: Concentrations and
942 behaviour of reactive gaseous mercury in ambient air, *Environ. Sci. Technol.*, 32, 49-57,
943 1998.
- 944 Liu, B., Keeler, G. J., Dvonch, J. T., Barres, J. A., Lynam, M. M., Marsik, F. J., Morgan, J. T.:
945 Temporal variability of mercury speciation in urban air, *Atmos. Environ.*, 41, 1911-
946 1923, 2007.
- 947 Lombard, M. A. S., Brycel, J. G., Mao, H., and Talbot, R.: Mercury deposition in Southern New
948 Hampshire, 2006–2009, *Atmos. Chem. Phys.*, 11, 7657-7668, 2011.
- 949 Luo, Y., Duan, L., Xu, G., and Hao, J.: Inhibition of mercury release from forest soil by high
950 atmospheric deposition of Ca^{2+} and SO_2^- , *Chemosphere*, 134, 113-119, 2015.
- 951 Lyman, S. N., Gustin, M. S., Prestbo, E. M., and Marsik, F. J.: Estimation of dry deposition of
952 atmospheric mercury in Nevada by direct and indirect methods, *Environ. Sci. Technol.*,
953 41, 1970-1976, 2007.
- 954 Lyman, S. N.: Gustin, M. S., Prestbo, E. M., Kilner, P. I., Edgerton, E., and Hartsell, B.: Testing
955 and application of surrogate surfaces for understanding potential gaseous oxidized

956 mercury dry deposition, *Environ. Sci. Technol.*, 43, 6235-6241, 2009.

957 Lyman, S. N., Gustin, M. S., and Presto, E. M.: A passive sampler for ambient gaseous oxidized
 958 mercury concentrations, *Atmos. Environ.*, 44, 246-252, 2010.

959 Lynam, M. M., Dvonch, J. T., Hall, N. L., Morishita, M., and Barres J. A.: Trace elements and
 960 major ions in atmospheric wet and dry deposition across central Illinois, USA, *Air Qual.*
 961 *Atmos. Health*, 8, 135-147, 2015.

962 Ma, M., Wang, D. Y., Du, H. X., Sun, T., Zhao, Z., and Wei, S. Q.: Atmospheric mercury
 963 deposition and its contribution of the regional atmospheric transport to mercury pollution
 964 at a national forest nature reserve, southwest China, *Atmos. Chem. Phys. Discuss.* 22
 965 (24), 20007-20018, 2015.

966 Ma, M., Wang, D., Du, H., Sun, T., Zhao, Z., Wang, Y., and Wei, S.: Mercury dynamics and
 967 mass balance in a subtropical forest, southwestern China, *Atmos. Chem. Phys.*, 16, 4529-
 968 4537, doi: 10.5194/acp-16-4529-2016, 2016.

969 Maestas, M. M.: Characterization of speciated atmospheric mercury concentration measurements
 970 in northern Utah, MSc. Thesis, The University of Utah, Utah, 152 pp., 2011.

971 Marsik, F.J., Keeler, G. J., Lindberg, S. E., and Zhang, H.: The air-surface exchange of gaseous
 972 mercury over a mixed sawgrass-cattail stand within the Florida Everglades, *Environ.*
 973 *Sci. Technol.*, 39, 4739-4746, 2005.

974 Marsik, F. J., Keeler, G. J., and Landis, M. S.: The dry-deposition of speciated mercury to the
 975 Florida Everglades: measurements and modeling, *Atmos. Environ.*, 41, 136-149, 2007.

976 Marsik, F. J.: Mercury Dry Deposition Measurement Intercomparison and Workshop Final
 977 Report,
 978 http://www.glc.org/glad/Projectdocs/Marsik/GLAD_Progress_Report_Oct2009_final.pdf
 979 f, 2009

980 Martin, R. S., Witt, M. L. I., Sawyer, G. M., Thomas, H. E., Watt, S. F. L., Bagnato, E.,
 981 Calabrese, S., Aiuppa, A., Delmelle, P., Pyle, D. M., Mather, T. A.: Bioindication of
 982 volcanic mercury (Hg) deposition around Mt. Etna (Sicily), *Chem. Geolog.*, 310-311,
 983 12-22, 2012.

984 McLaughlin, E., Driscoll, C., Yavitt, J., Newton, R., and Munson, R.: Mercury in upland and
 985 riparian wetland vegetation, in: Mercury in Adirondack wetlands, lakes and terrestrial
 986 systems (MAWLTS), New York State Energy Research and Development Authority,
 987 New York, 9-1-9-6, 2008.

988 Melendez-Perez, J. J., Fostier, A. H., Carvalho Jr., J. A., Windmoller, C. C., Santos, J. C., Carpi,
 989 A.: Soil and biomass mercury emissions during a prescribed fire in the Amazonian rain
 990 forest, *Atmos. Environ.*, 96, 415-422, 2014.

991 Mélières, M.-A., Pourchet, M., Charles-Dominique, P., and Gaucher, P. : mercury in canopy
 992 leaves of French Guiana in remote areas, *Sci. Total Environ.*, 311, 261-267, 2003.

993 Meyers, T. P., Hall, M. E., Lindberg, S. E., and Kim, K.: Use of the modified Bowen-ratio
 994 technique to measure fluxes of trace gases, *Atmos. Environ.*, 30, 3321-3329, 1996.

- 995 Michelazzo, P. A. M., Fostier A. H., Magarelli, G., Santos, J. C., and Andrade de Carvalho Jr.,
 996 A. : Mercury emission from forest burning in southern Amazon, *J. Geophys. Res. Lett.*,
 997 37, L09809, doi:10.1029/2009GL042220, 2010.
- 998 Miller, E. K, Vanarsdale, A., Keeler, G. J., Chalmers, A., Poissant, L., Kanman, N. C., and
 999 Brulotte, R.: Estimation and mapping of wet and dry mercury deposition across
 1000 northeastern North America, *Ecotox.* 14, 53-70, 2005.
- 1001 Munthe, J. Hultberg, H., and Iverfeldt, A.: Mechanisms of deposition of methylmercury and
 1002 mercury to coniferous forests, *Water Air Soil Poll.*, 80, 363-371, 1995.
- 1003 Niu, Z., Zhang, X., Wang, Z., Ci, Z.: Mercury in leaf litter in typical suburban and urban
 1004 broadleaf forests in China, *J. Environ. Sci.*, 23 (12), 2042-2048, 2011.
- 1005 N6voa-Mu6noz, J. C., Pontevedra-Pombal, X., Martinez-Cortizas, E., Garcia-Rodeja Gayoso, E.:
 1006 Mercury accumulation in upland acid forest ecosystems nearby a coal-fired power-plant
 1007 in Southwest Europe (Galicia, NW Spain), *Sci. Total Environ.*, 394, 303-312, 2008.
- 1008 Obrist, D., Johnson, D. W., Lindberg, S. E., Luo, Y., Hararuk, O., Bracho, R., Battles, J. J., Dail,
 1009 D. B., Edmonds, R. L., Monson, R. K., Ollinger, S. V., Pallardy, S. G., Pregitzer, K. S.,
 1010 and Todd, D. E.: Mercury distribution across 14 US Forests. Part I: Spatial patterns of
 1011 concentrations in biomass, litter, and soils, *Environ. Sci. Technol.*, 45, 3974-3981, 2011.
- 1012 Obrist, D.: Mercury distribution across 14 U.S. Forests. Part II: Patterns of methyl mercury
 1013 concentrations and areal mass of total and methyl mercury, *Environ. Sci. Technol.*, 46,
 1014 5921-5930, 2012.
- 1015 de Oliveira, R. R., da Silveira, C. L. P., Magalh6es, A. C., and Firme, R. P.: Turnover of heavy
 1016 metals in the litter of a urban forest at Rio de Janeiro, *Floresta e Ambiente*, 12 (1), 50-
 1017 56, 2005.
- 1018 Peterson, C. and Gustin, M. A.: Mercury in the air, water and biota at the Great Salt Lake (Utah,
 1019 USA), *Sci. Total Environ.*, 405, 255-268, 2008.
- 1020 Peterson, C., Alishahi, M., and Gustin, M. A.: Testing the use of passive sampling systems for
 1021 understanding air mercury concentrations and dry deposition across Florida, USA, *Sci.*
 1022 *Total Environ.*, 424, 297-307, 2012.
- 1023 Poissant, L., Pilote, M., Xu, X., and Zhang, H.: Atmospheric mercury speciation and deposition
 1024 in the Bay St. Francois wetlands, *J. Geophys. Res.*, 109, D11301,
 1025 doi:10.1029/2003JD004364, 2004.
- 1026 Poissant, L. and Casimir, A.: Water-air and soil-air exchange rate of total gaseous mercury
 1027 measured at background sites, *Atmos. Environ.*, 32, 883–893, 1998.
- 1028 Pokharel, A. K., and Obrist, D.: Fate of mercury in tree litter during decomposition,
 1029 *Biogeosciences*, 8, 2507-2521, 2011.
- 1030 Porvari, P. and Verta, M.: Total and methyl mercury concentrations and fluxes from small boreal
 1031 forest catchments in Finland, *Environ. Pollut.*, 123, 181-191, 2003.
- 1032 Prestbo, E. M. and Gay, D. A.: Wet deposition of mercury in the U.S. and Canada, 1996–2005:

1033 Results and analysis of the NADP mercury deposition network (MDN), *Atmos.*
1034 *Environ.*, 43, 4223-4233, 2009.

1035 Rasmussen, P. E., Mierle, G., and Nriagu, J. O. : The analysis of vegetation for total mercury,
1036 *Water Air Soil Poll.*, 56, 379-390, 1991.

1037 Rasmussen, P. E.: Temporal variation of mercury in vegetation, *Water Air Soil Poll.*, 80, 1039-
1038 1042, 1995.

1039 Rea, A. W., Keeler, G. J., and Scherbatskoy, T.: The deposition of mercury in throughfall and
1040 litterfall in the Lake Champlain watershed: a short-term study, *Atmos. Environ.*, 30 (19),
1041 3257-3263, 1996.

1042 Rea, A. W., Lindberg, S. E., and Keeler, G. J.: Assessment of dry deposition and foliar leaching
1043 of mercury and selected trace elements based on washed foliar and surrogate surfaces,
1044 *Environ. Sci. Technol.*, 34, 2418-2425, 2000.

1045 Rea, A. W., Lindberg, S. E., and Keeler, G. J.: Dry deposition and foliar leaching of mercury and
1046 selected trace elements in deciduous forest throughfall, *Atmos. Environ.*, 35, 3453-
1047 3462, 2001.

1048 Rea, A. W., Lindberg, S. E., Scherbatskoy, T., and Keeler, G. J.: Mercury accumulation in
1049 foliage over time in two northern mixed-hardwood forests, *Water Air Soil Poll.*, 133,
1050 49–67, 2002.

1051 Richardson, J. B. and Friedland, A. J.: Mercury in coniferous and deciduous upland forests in
1052 northern New England, USA: implications of climate change, *Biogeosciences*, 12,
1053 6737–6749, 2015.

1054 Risch, M. R., DeWild, J. F., Krabbenoft, D. P., Kolka, R. K., and Zhang, L.: Litterfall mercury
1055 dry deposition in the eastern USA, *Environ. Pollut.*, 161, 284-290, 2012.

1056 Roulet, M., Lucotte, M., Saint-Aubin, A., Tran, S., Rhéault, T. I., Farella, N., Da Silva, E. D. J.,
1057 Dezencourt, J., Sousa Passos, C.-J., Soares, G. S., Guimarães, J. –R., D., Mergler, D.,
1058 and Amorim, M. : The geochemistry of mercury in central Amazonian soils developed
1059 on the Alter-do-Chão formation of the lower Tapajós River Valley, Pará state, Brazil,
1060 *Sci. Total Environ.*, 223, 1-24, 1998.

1061 Roulet, M., Lucotte, M., Farella, N., Serique, G., Coelho, H., Sousa Passos, C. J., de Jesus
1062 Dasilva, E., Scavone de Andrade, P., Mergler, D., Guimarães, J.-R., D., and Amorim,
1063 M. : Effects of recent human colonization on the presence of mercury in Amazonian
1064 ecosystems, *Water Air Soil Poll.*, 112, 297-313, 1999.

1065 Ryaboshapko, A., Bullock Jr., R., Ebinghaus, R., Ilyin, I., Lohman, K., Munthe, J., Petersen, G.,
1066 Seigneur, C., and Wängberg, I. : Comparison of mercury chemistry models, *Atmos.*
1067 *Environ.*, 36, 3881-3898, 2002.

1068 Ryaboshapko, A., Bullock Jr., O. R., Christensen, J., Cohen, M., Dastoor, A., Ilyin, I., Petersen,
1069 G., Syrakov, D., Travnikov, O., Artz R. S., Davignon, D., Draxler, R. R., Munthe, J.,
1070 and Pacyna, J.: Intercomparison study of atmospheric mercury models: 2. Modelling
1071 results vs. long-term observations and comparison of country deposition budgets, *Sci.*

1072 Total Environ., 377, 319-333, 2007.

1073 Sakata, M., and Marumoto, K.: Dry Deposition Fluxes and Deposition Velocities of Trace
1074 Metals in the Tokyo Metropolitan Area Measured with a Water Surface Sampler,
1075 Environ. Sci. Technol., 38, 2190-2197, 2004.

1076 Sakata, M., and Marumoto, K.: Wet and dry deposition fluxes of mercury in Japan, Atmos.
1077 Environ., 39, 3139-3146, 2005.

1078 Sakata, M., and Asakura, K.: Evaluating relative contribution of atmospheric mercury species to
1079 mercury dry deposition in Japan, Water Air Soil Poll., 193, 51-63, 2008.

1080 Sather, M. E., Murkerjee, S., Smith, L., Mathew, J., Jackson, C., Callison, R., Scrapper, L.,
1081 Hathcoat, A., Adam, J., Keese, D., Ketcher, P., Brunette, R., Karlstrom J., and Van der
1082 Jagt, G.: Gaseous oxidized mercury dry deposition measurements in the Four Corners
1083 area and Eastern Oklahoma, U.S.A., Atmos. Poll. Res., 4, 168-180, 2013.

1084 Sather, M. E., Mukerjee, S., Allen, K. L., Smith, L., Mather, J., Jackson, C., Callison, R.,
1085 Scrapper, L., Hathcoat, A., Adam, J., Keese, D., Ketcher, P., Brunette, R., Karlstrom, J.,
1086 and Van der Jagt, G.: Gaseous Oxidized Mercury Dry Deposition Measurements in the
1087 Southwestern USA: A Comparison between Texas, Eastern Oklahoma, and the Four
1088 Corners Area, Sci. World J., 2014, ID 580723, 14 pages, 2014.

1089 Schwesig, D. and Matzner, E.: Pools and fluxes of mercury and methylmercury in two forested
1090 catchments in Germany, Sci. Total Environ., 260, 213-223, 2000.

1091 Schwesig, D. and Matzner, E.: Dynamics of mercury and methylmercury in forest floor and
1092 runoff of a forested watershed in Central Europe, Biogeochemistry, 53, 181-200, 2001.

1093 Seigneur, C., Karamchandani, P., Lohman, K., Vijayaraghavan, K., and Shia, R.-L.: Multiscale
1094 modeling of the atmospheric fate and transport of mercury, J. Geophys. Res., 106
1095 (D21), 27795-27809, 2001.

1096 Seigneur, C., Vijayaraghavan, K., Lohman, K., Karamchandani, P., and Scott, C.: Global source
1097 attribution for mercury deposition in the United States, Environ. Sci. Technol., 38, 555-
1098 569, 2004.

1099 Seigneur, C., Lohman, K., Vijayaraghavan, K., Jansen, J., and Levin, L.: Modeling Atmospheric
1100 Mercury Deposition in the Vicinity of Power Plants, JAPCA J Air Waste Ma., 56 (6),
1101 743-751, 2006.

1102

1103 Selin, N. E., Jacob, D. J., Yantosca, R. M., Strode, S., Jaeglé, L., Sunderland, E. M.: Global 3-D
1104 land-ocean-atmosphere model for mercury: present-day versus preindustrial cycles and
1105 anthropogenic enrichment factors for deposition, Global Biogeochem. Cy., 22, GB2011,
1106 doi:10.1029/2007GB003040, 2008.

1107 Selvendiran, P., Driscoll, C. T., Montesdeoca, M. R., and Bushey, J. T.: Inputs, storage, and
1108 transport of total and methyl mercury in two temperate forest wetlands, J. Geophys.
1109 Res., 113, G00C01, doi:10.1029/2008JG000739, 2008.

- 1110 Sheehan, K. D., Fernandez, I. J., Kahl, J. S., Amirbahman, A.: Litterfall mercury in two forested
1111 watersheds at Acadia National Park, Maine, USA, *Water Air Soil Poll.*, 170, 249-265,
1112 2006.
- 1113 Silva-Filho, E. V., Machado, W., Oliveira, R. R., Sella, S. M., Lacerda, L. D.: Mercury
1114 deposition through litterfall in an Atlantic Forest at Ilha Grande, Southeast Brazil,
1115 *Chemosphere*, 65, 2477-2484, 2006.
- 1116 Slinn, S. A. and Slinn, W. G. N.: Predictions for particle deposition on natural waters, *Atmos.*
1117 *Environ.*, 14 (9), 1013-1016, 1980.
- 1118 Song, S., Selin, N. E., Soerensen, A. L., Angot, H., Artz, R., Brooks, S., Brunke, E.-G., Conley,
1119 G., Dommergue, A., Ebinghaus, R., Holsen, T. M., Jaffe, D. A., Kang, S., Kelley, P.,
1120 Luke, W. T., Magand, O., Marumoto, K., Pfaffhuber, K. A., Ren, X., Sheu, G.-R.,
1121 Slemr, F., Warneke, T., Weigelt, A., Weiss-Penzias, P., Wip, D. C., and Zhang, Q.:
1122 Top-down constraints on atmospheric mercury emissions and implications for global
1123 biogeochemical cycling, *Atmos. Chem. Phys.*, 15, 7103–7125, doi:10.5194/acp-15-
1124 7103-2015, 2015.
- 1125 Sprovieri, F., Pirrone, N., Ebinghaus, R., Kock, H., and Dommergue, A.: A review of worldwide
1126 atmospheric mercury measurements, *Atmos. Chem. Phys.*, 10, 8245-8265,
1127 doi:10.5194/acp-10-8245-2010, 2010.
- 1128 Sprovieri, F., Gratz, L. E., Pirrone, N.: Development of a Ground-Based Atmospheric
1129 Monitoring Network for the Global Mercury Observation System (GMOS). In
1130 proceedings of the 16th International Conference on Heavy Metals in the Environment,
1131 Rome, 23-27 September 2012. E3S Web of Conferences,
1132 doi: 10.1051/e3sconf/20130117007, 2013.
- 1133 St Louis, V. L., Rudd, J. W. M., Kelly, C. A., Hall, B. D., Rolffhus, K. R., Scott, K. J., Lindberg,
1134 S. E., and Dong, W.: Importance of the forest canopy to fluxes of methylmercury and
1135 total mercury to boreal ecosystems, *Environ. Sci. Technol.*, 35, 3089-3098, 2001.
- 1136 Szopka, K., Karczewska, A., and Kabała, C.: Mercury accumulation in the surface layers of
1137 mountain soils: A case study from the Karkonosze Mountains, Poland, *Chemosphere*,
1138 83, 1507-1512, 2011.
- 1139 Teixeira, D. C., Montezuma, R. C., Oliveira, R. R., Silva-Filho, E. W.: Litterfall mercury
1140 deposition in Atlantic forest ecosystem from SE Brazil, *Environ. Pollut.* 164, 11-15,
1141 2012.
- 1142 Travnikov, O. and Ilyin, I.: The EMEP/MSC-E mercury modelling system, in: Pirrone, N. and
1143 Mason, R. (Eds), *Mercury fate and transport in the global atmosphere: emissions,*
1144 *measurements and models*, Springer, New York, 571-587, 2009.
- 1145 Travnikov, O., Lin, C.-J., Dastoor, A., Bullock, O. R., Hedgecock, I. M., Holmes, C., Ilyin, I.,
1146 Jaegle, L., Jung, G. J., Pan, L., Pongprueksa, P., Ryzhkov, A., Seigneur, C., And Skov,
1147 H.: Global and regional modelling, in: *Hemispheric Transport of Air Pollution 2010*
1148 *Part B: Mercury Air Pollution Studies No. 18*, Pirrone, N. and Keating, T., Eds, United
1149 Nations Publications, New York and Geneva, 101-148, 2010.

- 1150 Travnikov, O., Dastoor, A., Friedman, C., Ryzhkov, A., Selin, N., and Song, S: AMAP/UNEP,
1151 2015. Global Mercury Modelling: Update of Modelling Results in the Global Mercury
1152 Assessment 2013. Arctic Monitoring and Assessment Programme, Oslo, Norway/UNEP
1153 Chemicals Branch, Geneva, Switzerland, 32 pp., 2015.
- 1154 Wallschläger, D., Kock, H. H., Schroeder, W. H., Lindberg, S. E., Ebinghaus R., and Wilken, R.-
1155 D.: Estimating gaseous mercury emissions from contaminated floodplain soils to the
1156 atmosphere with simple field measurement techniques, *Water Air Soil Poll.*, 135, 39-54,
1157 2002.
- 1158 Wang, X., Lin, C.-J., and Feng, X.: Sensitivity analysis of an updated bidirectional air-surface
1159 exchange model for elemental mercury vapor, *Atmos. Chem. Phys.*, 14, 6273–6287,
1160 doi:10.5194/acp-14-6273-2014, 2014.
- 1161 Wang, Z., Zhang, X., Xiao, J., Zhijia, C., Yu, P.: Mercury fluxes and pools in three subtropical
1162 forested catchments, southwest China, *Environ. Pollut.*, 157, 801-808, 2009.
- 1163 Weiss-Penzias, P. S., Gustin, M. S., and Lyman, S. N.: Sources of gaseous oxidized mercury and
1164 mercury dry deposition at two southeastern U.S. sites, *Atmos. Environ.*, 45, 4569-4579,
1165 2011.
- 1166 Weiss-Penzias, P. S., Ortiz Jr., C., Acosta, R. P., Heim, W., Ryan, J. P., Fernandez, D., Collett
1167 Jr., J. L., and Flegal, A. R.: Total and monomethyl mercury in fog water from the
1168 central California coast, *Geophys. Res. Lett.*, 39, L03804, doi:10.1029/2011GL05324,
1169 2012.
- 1170 Witt, E. L., Kolka, R. K., Nater, E. A., and Wickman, T. R.: Influence of the Forest Canopy on
1171 Total and Methyl Mercury Deposition in the Boreal Forest, *Water Air Soil Poll.*, 199, 3-
1172 11, 2009.
- 1173 Wright, L. P. and Zhang, L.: An approach estimating bidirectional air-surface exchange for
1174 gaseous elemental mercury at AMNet sites, *J. Adv. Model. Earth Syst.*, 7, 35–49,
1175 doi:10.1002/2014MS000367, 2015.
- 1176 Wright, G., Gustin, M. S., Weiss-Penzias, P., and Miller, M. B.: Investigation of mercury
1177 deposition and potential sources at six sites from the Pacific Coast to the Great Basin,
1178 USA, *Sci. Total Environ.*, 470-471, 1099-1113, 2014.
- 1179 Xiao, Z., Sommar, J., Lindqvist, O., Giouleka, E.: Atmospheric mercury deposition to grass in
1180 southern Sweden, *Sci. Total Environ.*, 213, 85-94, 1998. Xu, L., Chen, J., Yang, L., Niu,
1181 Z., Tong, L., Yin, L., Chen, Y. : Characteristics and sources of atmospheric mercury
1182 speciation in a coastal city, Xiamen, China, *Chemosphere*, 119, 530-539, 2015.
- 1183 Xu, X., Yang, X., Miller, D. R., Helble, J. J., and Carley, R. J.: Formulation of bi-directional
1184 atmosphere-surface exchanges of elemental mercury, *Atmos. Environ.*, 33, 4345-4355,
1185 1999.
- 1186 Zhang, H. Lindberg, S. E., Marsik, F. J., and Keeler, G. J.: Mercury air/surface exchange kinetics
1187 of background soils of the Tahquamenon River watershed in the Michigan Upper
1188 Peninsula, *Water Air Soil Poll.*, 126, 151-169, 2001.

- 1189 Zhang, L., Gong, S., Padro, J., Barrie, L.A.: A size-segregated particle dry deposition scheme for
1190 an atmospheric aerosol module, *Atmos. Environ.*, 35, 549-560, 2001.
- 1191 Zhang, L., Brook, J. R., and Vet, R.: A revised parameterization for gaseous dry deposition in
1192 air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, 2003.
- 1193 Zhang, L., Wright, L., P., Blanchard, P.: A review of current knowledge concerning dry
1194 deposition of atmospheric mercury, *Atmos. Environ.*, 43, 5853-5864, 2009.
- 1195 Zhang, L., Blanchard, P., Johnson, D., Dastoor, A., Ryzhkov, A., Lin, C. J., Vijayaraghavan, K.,
1196 Gay, D., Holsen, T. M., Huang, J., Graydon, J. A., St. Louis, V. L., Castro, M. S.,
1197 Miller, E. K., Marsik, F., Luk, J., Poissant, L., Pilote, M., and Zang, K. M.: Assessment
1198 of modeled mercury dry deposition over the Great Lakes region, *Environ. Pollut.*, 161,
1199 272-283, 2012a.
- 1200 Zhang, L., Blanchard, P., Gay, D. A., Prestbo, E. M., Risch, M. R., Johnson, D., Narayan, J.,
1201 Zsolway, R., Holsen, T. M., Miller, E. K., Castro, M. S., Graydon, J. A., Louis, V. L.
1202 St., and Dalziel, J.: Estimation of speciated and total mercury dry deposition at
1203 monitoring locations in eastern and central North America, *Atmos. Chem. Phys.*, 12,
1204 4327-4340, doi:10.5194/acp-12-4327-2012, 2012b.
- 1205 Zhou, J., Feng, X., Liu, H., Zhang, H., Fu, X., Bao, Z., Wang, X., Zhang, Y.: Examination of
1206 total mercury inputs by precipitation and litterfall in a remote upland forest of
1207 Southwestern China, *Atmos. Environ.*, 81, 364-372, 2013.
- 1208 Zhu, W., Lin, C.-J., Wang, X., Sommar, J., Fu, X. W., and Feng, X.: Global observations and
1209 modeling of atmosphere-surface exchange of elemental mercury – a critical review,
1210 *Atmos. Chem. Phys. Discuss.* doi:10.5194/acp-2015-1064, in review, 2016.

Table 1: Summary of estimated and measured mercury dry deposition, and measured mercury in litterfall and throughfall in Asia, Europe and North America (converted to annual value in the unit of $\mu\text{g m}^{-2} \text{yr}^{-1}$).

	Asia			Europe			North America		
	Median, Mean	Range	<i>n</i>	Median, Mean	Range	<i>n</i>	Median, Mean	Range	<i>n</i>
Modelled GOM+PBM	34.5, 83.8	3.8 – 407	12	3.0, 3.8	0.1 – 17.5	27	2.6, 8.7	0.05 – 318	157
Measured GOM+PBM	10.7, 22.7	2.4 – 463	50				6.1, 30.8	0.26 – 520	105
Litterfall	34.8, 42.8	4.2 – 219.9	20	12.8, 14.2	0.05 – 42.0	14	11.9, 12.9	0.02 – 57.1	92
Throughfall	49.0, 43.5	10.5 – 71.3	11	16.3, 19.0	12 – 40.1	17	7.0, 9.3	2.1 – 26.6	32

