

Supplement of

Air–surface exchange of gaseous mercury over permafrost soil: an investigation at a high-altitude (4700 m a.s.l.) and remote site in the central Qinghai-Tibet Plateau

Zhijia Ci¹, Fei Peng², Xian Xue², and Xiaoshan Zhang¹

¹Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, 100085, China

²Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences, Lanzhou, 730000, China

Correspondence to: Z. J. Ci (zjci@rcees.ac.cn)

There are 5 pages of supplement including **Figure S1–S3** and **Table S1**.




Figure S1. Transmittance of DI water, quartz glass, UVB filter (cut-off wavelength of 320 nm) and UV filter (cut-off wavelength of 405 nm) on the radiation of 200–800 nm. Aluminum foil blocks all radiation (data not shown). UVB filter transmitted 0.7% of incoming UVB, 81.4% of incoming UVA, and 84.4% of the visible light. UV filter transmitted 0.1% of incoming UVB, 3.1% of incoming UVA, and 71.2% of the visible light. The transmittances of materials were determined by UV-VIS-NIR Spectrophotometer (UV-3600, Shimadzu, Japan).




Figure S2. Relationship between the accumulative PAR and accumulative Hg(0) emission flux in the daytime in six days without precipitation during December 2014 and May–June 2015 campaign.




Figure S3. Arrhenius relationships between Hg(0) flux and soil temperature during May–June 2015 campaign.

Table S1. *Ea/R* for Hg(0) emission from various soils with large range of soil Hg concentrations.

Location	Soil type	Soil Hg (ug kg ⁻¹)	<i>Ea/R</i>	References
Qinghai-Tibet Plateau, China*	Background soil	12–13	30.40	This study
Mt. Gongga, South China**	Forest soil	80–880	12.56–23.56	Fu et al. (2008)
Oak Ridge, Tennessee, USA**	Forest soil	61–469	9.08–12.56	Carpi and Lindberg (1998)
Guiyang city, South China**	Urban soil	150–630	9.31–34.75	Feng et al. (2005)
Chongqing city, South China**	Urban soil	136.7–526	15.68	Wang et al. (2006)
Guizhou, South China**	Agricultural soil	249.9	19.62–30.16	Wang et al. (2003)
Idrijca, Slovenia*	Hg-mining soil	4.1–251x10 ³	9.86–13.11	Kocman and Horvat (2010)
Clyde Forks, Canada*	Hg-enriched soils	240 x10 ³	<10	Corbett-Hains et al. (2012)
			7.31	Theoretical value for Hg(0)

*These studies used the Hg(0) flux data in the dark to calculate the *Ea/R*.

**These studies used the bulk Hg(0) flux data in the light to calculate the *Ea/R*.

References

- Boudala, F. S., Folkins, I., Beauchamp, S., Tordon, R., Neima, J., and Johnson, B. Mercury flux measurements over air and water in Kejimikujik National Park, Nova Scotia, *Water Air Soil Pollut.*, 122, 183–202, 2000.
- Carpi, A., and Lindberg, S. E. Application of a Teflon™ dynamic flux chamber for quantifying soil mercury flux: tests and results over background soil, *Atmos. Environ.*, 32, 873–882, 1998.
- Corbett-Hains, H., Walters, N. E., and Van Heyst, B. J. Evaluating the effects of sub-zero temperature cycling on mercury flux from soils, *Atmos. Environ.*, 63, 102–108, 2012.
- Feng, X. B., Wang, S. F., Qiu, G. A., Hou, Y. M., and Tang, S. L. Total gaseous mercury emissions from soil in Guiyang, Guizhou, China, *J. Geophys. Res. Atmos.*, 110, D14306, doi: 10.1029/2004JD005643, 2005.
- Fu, X. W., Feng, X. B., and Wang, S. F. Exchange fluxes of Hg between surfaces and atmosphere in the eastern flank of Mount Gongga, Sichuan province, southwestern China, *J. Geophys. Res. Atmos.*, 113, D20, doi: 10.1029/2008JD009814, 2008.
- Kocman, D., and Horvat, M. A laboratory based experimental study of mercury emission from contaminated soils in the River Idrijca catchment, *Atmos. Chem. Phys.*, 10, 1417–1426, 2010.
- Wang, D. Y., He, L., Shi, X. J., Wei, S. Q., and Feng, X. B. Release flux of mercury from different environmental surfaces in Chongqing, China, *Chemosphere*, 64, 1845–1854, 2006.
- Wang, S., Feng, X., and Qiu, G. The study of mercury exchange rate between air and soil surface in Hongfeng reservoir region, Guizhou, PR China, *J. Phys. IV.*, 107, 1357–1360, 2003.