

Interactive comment on “Development of a high-resolution emission inventory and its evaluation through air quality modeling for Jiangsu Province, China” by Yaduan Zhou et al.

Yaduan Zhou et al.

yuzhao@nju.edu.cn

Received and published: 22 November 2016

Manuscript No.: acp-2016-567

Title: Development of a high-resolution emission inventory and its evaluation and application through air quality modeling for Jiangsu Province, China

Authors: Yaduan Zhou, Yu Zhao, Pan Mao, Qiang Zhang, Jie Zhang, Qiu Liping, Yang Yang

We thank very much for the valuable comments and suggestions from reviewer 2, which help us improve our manuscript. The comments were carefully considered and revisions have been made in response to suggestions. Following is our point-by-point

C1

responses to the comments and corresponding revisions.

Reviewer #2

1. This manuscript presents the development and evaluation of detailed emission inventories for major anthropogenic emission sources in Jiangsu Province, China. Emission is one of the major sources of uncertainty chemical transport modeling in China, where air pollution poses increasing threat to public health. Although it focuses only on one province (a relatively small emitter), which limits its scientific value to the broad ACP readership, this work could help establish procedures to develop and improve similar emission inventories in other provinces in this region. This manuscript could be improved in several key aspects to meet the requirements ACP. First, the presentation, in the current form, needs to be polished. There are several places in the text that the expression is either ambiguous or confusing. Second, the manuscript lacks detailed information on how these inventories were compiled (formula, parameters, etc), and where to find the key information to verify or reproduce the results reported here. Third, some of the materials included here are not relevant to the main theme. For example, the sensitivity of ozone and PM_{2.5} to 50% reduction of NO_x and VOC emissions has nothing to do with development and evaluation of the emission inventories. The authors may consider either broadening the theme (development, evaluation, and application of the EIs) or removing the sensitivity section. Finally, the interpretation of some results need to be toned down or adjusted.

Response and revisions:

We appreciate the reviewer's crucial and important comments. In general, the presentation of the work has been improved, based on specific comments/suggestion from the reviewer. Detailed information on emission inventory development has been added, as indicated in the response to Questions 4, 7, 9, and 11 from the reviewer. In particular, Table S1 has been expanded and a new Table S3 has been added in the supplement to provide more detailed information on data sources of activities and emission factors.

C2

We also take the reviewer's suggestion and broaden the theme by adding the word "application" in the title of the paper. Uncertainties of the analysis have been also discussed to avoid overstatement of the work, as indicated in the response to Questions 14 and 16 from the reviewer.

2. L6: process(es).

Response and revisions:

We thank the reviewer's reminder and it is now corrected.

3. L26-30: Please reword this sentence. Define "unfavorable meteorology". Is the meteorology simulated by WRF inaccurate or the meteorology condition not conducive to pollution formation and accumulation? If the meteorology simulation is problematic, emission may reduce the model bias by offsetting opposite bias caused by meteorology inputs, but that does not warrant the quality of emission data per se.

Response and revisions:

The term "unfavorable meteorology" mentioned in the manuscript referred to the meteorology condition with low wind speed and PBL height, in which horizontal and vertical movement of atmosphere was limited. Under such condition, primary pollutants from large emitters would be easily accumulated over time, and air quality would thus be largely influenced by local sources. The sentence has been revised as "Under the unfavorable meteorology in which horizontal and vertical movement of atmosphere was limited, the simulated SO₂ concentrations at downtown Nanjing (the capital city of Jiangsu) using the regional or national inventories were much higher than observation, implying the overestimated urban emissions when economy or population densities were applied to downscale or allocate the emissions" in lines 26-31 Pages 2-3 in the revised manuscript.

We also agree with the reviewer that uncertainties existed in meteorology field simulation and that the performance of air quality modeling might be offset by the opposite

C3

bias of emission inventory and simulated meteorology. To test the deviation caused by meteorology field simulation with WRF, the model performance of SO₂ was reevaluated excluding the data from 5pm Oct 9th to 5am 10th. The normalized mean bias (NMB) were recalculated at -13%, 66%, and 50%, and the normalized mean errors (NME) were 45%, 88%, and 78% using the provincial, regional, and national inventories, respectively. This result thus implied that the provincial inventory could better support the air quality modeling, even uncertainties existed in meteorology field simulation.

4. Section 2.1: There are at least eight sectors considered here. Is Eq. (1) applied to all sectors or only to point sources? In the case that other formulas are used, please explain in more details (only citations are briefly provided here). It will be very useful to provide a table that includes the following key information: 1) sector; 2) sources included in that sector; 3) method to estimate emission; 4) key parameters and data sources to obtain the information; 5) special adjustment made.

Response and revisions:

We thank the reviewer's important comment. We checked the sectors, and confirmed that there are seven main sectors (residential & commercial is one sector). We have stated in line 133 Page 6 in the revised manuscript that Eq. (1) is applied only to point sources. No equation is provided for mobile sources (i.e., on-road vehicles) as their emissions were estimated using COPERT model. We have also provided the equation and explanation for calculating the emissions from area sources in lines 148-151 Page 7 in the revised manuscript. As suggested by the reviewer, the detailed source categories by sector and the main data sources of activity levels by category were summarized in the revised Table S1 in the supplement. We keep the discussions on data sources of emission factors by category in Section 2.3, with the detailed information provided in Table S2 and a new Table S3 in the supplement, as explained in the response to Q7.

5. L177: which model requirement?

C4

Response and revisions:

We thank the reviewer's reminder and admit the original word was confusing. The "model" here indicates COPERT 4 applied for calculating the emissions from on-road vehicles. Since differences exist in the categories of vehicle types between Chinese and European criteria, population of each Chinese vehicle category should be converted to the number consistent with the COPERT categories, in order to use the model. According to Cai and Xie (2007), for example, the buses, passenger cars (PC), heavy-duty trucks (HDT), light-duty trucks (LDT), and motorcycles (MT) in COPERT 4 indicate the big-size and middle-size passenger cars, small-size and mini passenger cars, heavy-duty and intermediate duty trucks, light-duty and mini trucks, and motorcycles in Chinese statistic system. The original sentence has been modified as "Populations of different vehicle types were derived from statistical yearbooks by city and then converted to the numbers in COPERT 4 categories" in lines 183-184 Page 8 in the revised manuscript.

6. L181-185: Can not understand this sentence. Please rephrase it. It seems a subtraction is involved here, by using data from different sources. What is the implication for uncertainty by subtracting data from different sources?

Response and revisions:

We thank the reviewer's important comment. In this work, as indicated in lines in the revised manuscript, information from the official environmental statistics, Pollution Source Census (PSC), and on-site survey on large emitters were collected to estimate the emissions from point sources. Although most of large industry sources could be investigated through this method, information of certain small plants were not included in those datasets and had to be treated as area sources. Thus the energy consumption and production provided in provincial statistical yearbooks were applied to check the completeness of point source investigation, and the difference between the energy statistics and overall activity levels of point sources was assumed as the activity lev-

C5

els of industrial area sources. The sentence has been revised as "For area sources, the coal consumption of residential activities was directly taken from National Energy Statistic Yearbook (NBSC, 2013c), while that of small industrial plants were calculated by subtracting the coal consumed by industrial point sources from the coal consumption of total industry provided in the provincial energy balance (NBSC, 2013c)" in lines 188-192 Page 8 in the revised manuscript.

7. Section 2.3. Again, a list of emission factors for each sector/source will be very useful. Maybe a table in the supplementary information.

Response and revisions:

We thank the reviewer's important comment. As required by the reviewer, a new Table S3 was provided in the revised supplement, summarizing the detailed emission factors and their sources by sector.

8. L227: "by with"? Please revise the sentence.

Response and revisions:

We thank the reviewer's reminder and admit the typo error. Now the error has been corrected in line 237 Page 10 in the revised manuscript.

9. L233-240. Please provide information of the spatial maps used to distribute emissions for each sector. Currently only the data for open burning have been given.

Response and revisions:

We thank the reviewer's suggestion and agree that more detailed information on spatial distribution of emissions should be provided. For point sources, information of latitude and longitude for each plant was collected from PSC. Locations of the total 6750 point sources shown in Figure S2 in the supplement were verified and modified in Google Earth. The road and hydrographic net in Jiangsu were respectively used to distribute the emissions from on-road transportation and ships city-by city. Densities

C6

of GDP and population in Jiangsu applied to allocate emissions of area sources were from the research by Huang et al. (2014) and Fu et al. (2014). NH₃ emissions from livestock, fertilizer usage and agricultural vehicles were allocated by incorporating the population density and distribution of land types categorized to agricultural activities from the land cover dataset GlobCover2009 (<http://globalchange.nsd.cn>). Such information has been added in lines 243-256 Page 10 in the revised manuscript.

10. L280: "initial (concentration) and boundary conditions"

Response and revisions:

We thank the reviewer's reminder and the text has been corrected in line 293 Page 11 in the revised manuscript.

11. L290-291: Please elaborate how the vertical distribution is determined here. Is this applied to all sectors or just point sources?

Response and revisions:

We thank the reviewer's important suggestion and admit that it was not clearly explained in the original manuscripts. The vertical distributions of emissions were directly taken from L. Wang et al. (2010) except for the power sector, as the height of discharge outlet for each power plant was available for Jiangsu. According to L. Wang et al. (2010), the fractions of emissions of industry sources were 50%, 30% and 20% in layers 1-3, respectively. For the sources near the surface, i.e., transportation, residential & commercial combustion, solvent use, agriculture, and other sources, emissions were overall allocated to the first vertical layer in the model. The emissions of power plants were concentrated in layers 2-5, with the fractions estimated as 14.7%, 45.7%, 34.9% and 4.7%, respectively, based on the height information of the stacks. We have added the information in lines 304-311 Page 12 in the revised manuscript.

12. L447-449: Is the agricultural GDP increase due to change in market price or commodity quantity?

C7

Response and revisions:

We thank the reviewer's crucial comment and agree that application of commodity quantity would be more reasonable and persuasive to illustrate the case. We have checked the growth of livestock and poultry in Jiangsu from the provincial statistics. In lines 465-468 Page 17 in the revised manuscript, the sentence has been modified as "According to the provincial statistics, the total numbers of livestock and poultry increased 6% and 10% from 2010 to 2012 in Jiangsu (JSNBS, 2013). The growth of activity levels was expected to result in enhanced emissions, as very little progress was achieved for NH₃ control for these years."

13. Section 2 may need a sub-section to describe the measurement data (satellite and ground observations) used in this study.

Response and revisions:

We thank the reviewer's comment. The ground observation and retrieval of satellite data were not the main tasks of this study, and the data were directly taken from other publication/groups without any extra modification. Therefore, we do not think it is necessary to describe the data in a specific section, and we have kept the description in corresponding sections in the revised manuscript, i.e., lines 593-599 for ground observations and lines 526-534 Pages 19-20 for satellite observation.

14. Section 4.1 Using satellite NO₂ data for emission evaluation has been well explored (check the literature for more details). There are several issues related to the method used here. Note OMI observes VCD, not emission. Therefore, it may make more sense to compare the VCD from OMI to the equivalent from CMAQ, not the emission distribution directly, unless the nonlinear relationship between emission and VCD is accounted for. In addition, the noise to signal ratio in OMI NO₂ increases with decreasing VCD. The interpretation of the correlation between emission and OMI VCD needs to consider these factors.

C8

Response and revisions:

We thank the reviewer's very important comment and agree that the uncertainty of the method should be discussed in the paper. First, the quality control and reliability of OMI retrieved NO₂ VCDs has been added in lines 529-534 Pages 19-20 in the revised manuscript with literature provided. We then acknowledged that the comparisons between spatial distribution of NO₂ VCDs and emissions should be cautiously interpreted particularly for regions with relatively low values, as the noise to signal ratio in OMI NO₂ increases with decreased VCDs. Although summer NO₂ VCDs were applied in this study to eliminate the effects of long lifetime of NO₂ on pollution plums transport and chemical reaction, non-linear relationship still exists between VCDs and emissions. More detailed comparisons between NO₂ from satellite observation and CTM are thus recommended when improved characterization of NO₂ vertical distribution is available for the region. We have added such discussions in lines 576-582 Page 21 in the revised manuscript.

15. Section 4.3: It is good to see that the new emission inventories can reduce high bias during extreme events when the meteorology is not correctly simulated. Do we have any results showing that this emission data can be used to improve prediction of real pollution events? This manuscript discusses the new emission data for the entire province. Why is only the model prediction over Nanjing discussed in the model-observation comparison?

Response and revisions:

We thank the reviewer's positive remarks and suggestion. As shown in Table 2, better model performances for SO₂, NO_x, O₃ and PM_{2.5} simulation could be achieved with provincial inventory than those with national or regional inventory for the whole October, implying the improvement of provincial emission inventory for Jiangsu. Besides the special case discussed in Section 4.3, the simulated PM_{2.5} concentrations from 8pm on 18th to 5pm on 19th October with different inventories could also indicate the

C9

advantage of the provincial inventory against the regional and national inventories in a pollution event prediction. Taking Caochangmen (CCM) site as the example, the observed PM_{2.5} concentration kept increasing from 8pm on 18th with the highest value reaching 114 $\mu\text{g}/\text{m}^3$ at 2am on 19th. Simulated PM_{2.5} concentrations with provincial, regional and national inventory at that time were 90, 53 and 45 $\mu\text{g}/\text{m}^3$, respectively. The correlation coefficients between observations and simulations with the three inventories were calculated as 0.66, 0.44 and 0.30 in the episode, respectively, indicating the better performance with provincial inventory in real pollution episode simulation. We have added such discussions in lines 619-626 Page 23 in the revised manuscript.

Regarding the second question, the official data on air quality from surface measurements have been routinely published since 2013, thus the complete data for Jiangsu were unavailable for the years before 2013, except for its capital city Nanjing. Therefore only the observations at the nine sites in Nanjing were applied in this work.

16. As mentioned earlier, this section may not be necessary to support the main argument here. The interpretation of the results needs to acknowledge several caveats in the design and scope of these simulations. For instance, the brute force method does not consider nonlinearity in ozone response to precursor change. Large uncertainties exist in the emission dataset, including VOCs that have not been evaluated here, which will affect the chemistry and directionality of the ozone response. In addition, the sensitivity examines ozone response, but NO_x and VOCs changes also result in changes in other pollutants, such as PM_{2.5}. It is hence premature to draw such conclusion as in L770-772.

Response and revisions:

We thank the reviewer's very crucial comment and suggestion. As suggested, we revised the title of the paper, stressing the "application" of the emission inventory. We agree with the reviewer that the brute-force method ignores the nonlinearity of O₃ response to the changes of precursor emissions, and that is a big uncertainty of the anal-

C10

ysis. The results should thus be cautiously interpreted, and the comparisons with other simulation methods that take the nonlinearity mechanisms into account (e.g., OSAT or tagged species method) are further recommended. We have added the discussions in lines 825-831 Page 30 in the revised manuscript.

For VOC, the detailed information on development and evaluation of provincial inventory with source profiles is presented in a separate paper (Zhao et al., in preparation). Regarding the length of current paper, we have briefly stated that in lines 232-234 Page 9, and in lines 681-683 Page 25 in the revised manuscript.

We also agree with the reviewer that the air quality includes many issues besides O3 problem, thus the sentence in lines 770-772 in the original manuscript has been deleted to avoid overstating the case.

References

Cai, H., Xie, S. D.: Estimation of vehicular emission inventories in China from 1980 to 2005, *Atmos. Environ.*, 41, 8963-8979, 2007.

Fu, J. Y., Jiang, D., Huang, Y. H.: 1 Km Grid Population Dataset of China (PopulationGrid_China), Global Change Research Data Publishing & Repository, DOI: 10.3974/geodb.2014.01.06. V1, 2014.

Huang, Y. H., Jiang, D., Fu, J. Y.: 1 Km Grid GDP Data of China (2005, 2010) (GDPGrid_China), Global Change Research Data Publishing & Repository, DOI: 10.3974/geodb.2014.01.07. V1, 2014.

JSNBS (Jiangsu Bureau of Statistics): Statistical Yearbook of Jiangsu, Beijing, China Statistics Press, 2013 (in Chinese).

Wang, L. T., Jang, C., Zhang, Y., Wang, K., Zhang, Q., Streets, D. G., Fu, C. J., Lei, Y., Schreifels, J., He, K. B., Hao, J. M., Lam, Y. F., Lin, J., Meskhidze, N., Voorhees S., Evaris D., Phillips S.: Assessment of air quality benefits from national air pollution control policies in China. Part II: Evaluation of air quality predictions and air quality

C11

benefits assessment. *Atmos. Environ.*, 44, 3449-3457, 2010.

Zhao, Y., Mao, P., Zhou, Y., Yang, Y., Zhang, J., Wang, S., Dong, Y., Xie, F., Yu, Y., and Li, W.: Improved provincial emission inventory and speciation profiles of anthropogenic non-methane volatile organic compounds: a case study for Jiangsu, China, in preparation.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-567, 2016.