

Supplementary Information for publication

Single-particle investigation of summertime and wintertime Antarctic sea spray aerosols using low-Z particle EPMA, Raman microspectrometry, and ATR-FTIR imaging techniques

Hyo-Jin Eom¹, Dhruvajyoti Gupta¹, Hye-Rin Cho¹, HeeJin Hwang², SoonDo Hur², Yeontae Gim³ and Chul-Un Ro^{1,*}

¹*Department of Chemistry, Inha University, Incheon, Republic of Korea*

²*Polar Climate Change Research Division, Korea Polar Research Institute, Incheon, Republic of Korea*

³*Arctic Research Center, Korea Polar Research Institute, Incheon, Republic of Korea*

* *Corresponding author. Tel.: +82 32 860 7676; fax: +82 32 874 9207*
E-mail address: curo@inha.ac.kr (C.-U. Ro)

Table S1. Average elemental concentrations of individual particles in summertime and wintertime PM_{2.5-10} and PM_{1.0-2.5} fractions using low-Z particle EPMA.

sample	Summertime sample S1				Wintertime sample S2			
size fraction	PM _{1.0-2.5} (stage 3)		PM _{2.5-10} (stage 2)		PM _{1.0-2.5} (stage 3)		PM _{2.5-10} (stage 2)	
number of particles	146		148		154		156	
average size (μm)	2.0 (±0.6)		2.9 (±1.5)		1.7 (±0.8)		3.2 (±1.5)	
element	at. conc. (%)	wt. conc. (%)	at. conc. (%)	wt. conc. (%)	at. conc. (%)	wt. conc. (%)	at. conc. (%)	wt. conc. (%)
C	27.1 (±4.8)	14.6	21.9 (±5.9)	11.5	21.6 (±5.1)	11.2	18.8 (±3.8)	9.6
O	17.3 (±3.2)	12.4	19.5 (±4.7)	13.6	17.4 (±3.2)	12.0	18.2 (±4.0)	12.3
Na	25.2 (±3.3)	25.9	26.1 (±3.4)	26.2	27.6 (±4.2)	27.3	28.3 (±4.0)	27.5
Mg	2.3 (±0.4)	2.5	2.7 (±0.7)	2.9	2.9 (±0.5)	3.0	2.7 (±0.6)	2.8
Si	0.8 (±0.5)	1.0	0.3 (±0.4)	0.4	0.0	0.0	0.0	0.0
S	1.6 (±0.3)	2.3	1.8 (±0.5)	2.5	1.6 (±0.2)	2.2	1.6 (±0.3)	2.2
Cl	24.8 (±3.7)	39.4	26.4 (±3.4)	40.9	28.1 (±4.1)	42.8	29.2 (±3.8)	43.8
K	0.5 (±0.2)	0.9	0.5 (±0.2)	0.9	0.4 (±0.3)	0.7	0.5 (±0.2)	0.8
Ca	0.6 (±0.2)	1.1	0.7 (±0.3)	1.2	0.5 (±0.6)	0.9	0.6 (±0.3)	1.0

Figure S1. Three-day (72 h) backward air mass trajectories at 500 m-, 1000 m- and 1500 m-receptor heights in December 9, 2011 and July 23, 2012. HYbrid Lagrangian Single-Particle Integrated Trajectory (HYSPLIT) model available at the NOAA Air Resources Laboratory's web server (<http://www.arl.noaa.gov/ready/hysplit4.html>) was used.





Figure S2. Relative abundances of individual particles in two size fractions of the summer and winter samples, having specific concentration ratios of (a) C, (b) S, (c) Ca, (d) Cl, (e) Mg, and (f) K to Na. The red solid and dotted lines demote the stage 2 (PM_{2.5-10}) and stage 3 (PM_{1.0-2.5}) particles of the summertime sample, respectively, and the blue solid and dotted lines are for stage 2 (PM_{2.5-10}) and stage 3 (PM_{1.0-2.5}) particles of the wintertime sample, respectively. The green lines indicate bulk sea-water ratios corresponding to the elements.




Figure S3. Raman and ATR-FTIR spectra of the aerosols generated by the nebulization of a mixture solution of 0.2 M alanine and 0.1 M MgCl₂ standard chemicals.




Figure S4. Raman and ATR-FTIR spectra of some target chemicals for the organics in SSAs.




Figure S5. Raman and ATR-FTIR spectra of powdery standard Mg palmitate, palmitic acid, Mg stearate, and stearic acid.




Figure S6. ATR-FTIR spectra of Mg palmitate, Mg stearate, a mixture of Mg palmitate and stearate (by 3:1), and MgFAs-containing SSA.




Figure S7. Raman and ATR-FTIR spectra of standard inorganic chemicals (sulfates and nitrates).

