

1 Multi-pollutants emissions from the burning of major 2 agricultural residues in China and the related 3 health-economic effect assessment

4 Chunlin Li¹, Yunjie Hu¹, Fei Zhang¹, Jianmin Chen^{1,2}, Zhen Ma¹, Xingnan Ye¹, Xin
5 Yang^{1,2}, Lin Wang^{1,2}, Xingfu Tang¹, Renhe Zhang², Mu Mu², Guihua Wang²,
6 Haidong Kan³, Xinming Wang⁴, Abdelwahid Mellouki⁵

7 ¹Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention (LAP³), Department
8 of Environmental Science & Engineering, Fudan University, Handan Road 220, Shanghai 200433,
9 China

10 ²Institute of Atmospheric Sciences, Fudan University, Handan Road 220, Shanghai 200433, China

11 ³Public Health School, Fudan University, Dongan Road 120, Shanghai 200032, China

12 ⁴State Key Lab of Organ Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy
13 of Sciences, Kehuajie Road 511, Guangzhou 510640, China

14 ⁵Institut de Combustion, A érothermique, R éactivit é et Environnement, CNRS, 45071 Orl éans
15 cedex 02, France

16 *Correspondence to:* J. M. Chen (jmchen@fudan.edu.cn)

17 **Abstract.** Multi-pollutants in smoke particulate matter (SPM) were identified and
18 quantified for biomass burning of five major agricultural residues such as wheat, rice,
19 corn, cotton, and soybean straws in China by aerosol chamber system combining
20 with various measurement techniques. The primary emission factors (EFs) for PM_{1.0}
21 and PM_{2.5} are 3.04-12.64 and 3.25-15.16 g kg⁻¹. Organic carbon (OC), elemental
22 carbon (EC), char-EC, soot-EC, water-soluble inorganics (WSI), water-soluble
23 organic acids (WSOA), water-soluble amine salts (WSA), trace mineral elements
24 (THM), polycyclic aromatic hydrocarbons (PAHs), and phenols in smoke
25 PM_{1.0}/PM_{2.5} are, 1.34-6.04/1.54-7.42, 0.58-2.08/0.61-2.18, 0.51-1.67/0.56-1.76,
26 0.05-0.41/0.05-0.42, 0.51-3.52/0.52-3.81, 0.13-0.64/0.14-0.77,
27 (4.39-85.72/4.51-104.79)×10⁻³, (11.8-51.1/14.0-131.6)×10⁻³, (1.1-4.0/1.8-8.3)×10⁻³,
28 and (7.7-23.5/9.7-41.5)×10⁻³ g kg⁻¹, respectively. EC and soot-EC mainly exist in

29 PM_{1.0}, heavy metal-bearing particles favor to reside in the range of smoke PM_{1.0-2.5},
30 which are also confirmed by individual particle analysis.

31 With respect to five scenarios of burning activities, the average emissions and
32 overall propagation of uncertainties at 95% confidence interval (CI) of SPM from
33 agricultural open burning in China in 2012 were estimated for PM_{2.5}, PM_{1.0}, OC, EC,
34 char-EC, soot-EC, WSI, WSOA, WSA, THM, PAHs, and phenols to be 1005.7
35 (-24.6% , 33.7%), 901.4 (-24.4%, 33.5%), 432.4 (-24.2%, 33.5%), 134.2 (-24.8%,
36 34.0%), 113.1 (-24.8%, 34.1%), 21.0 (-26.3%, 35.9%), 249.8 (-25.4%, 34.9%), 25.1
37 (-33.3%, 41.4%), 5.8 (-30.1%, 38.5%), 8.7 (-26.6%, 35.6%), 0.5 (-26.0%, 34.9%),
38 and 2.7 (-26.1%, 35.1%) Gg, respectively. The emissions were further
39 temporal-spatially characterized using geographic information system (GIS) at
40 different regions in summer and autumn post-harvest periods. It was found less than
41 25 % of the total emissions were released during summer harvest that was mainly
42 contributed by the North Plain and the Central of China, especially Henan, Shandong,
43 and Anhui, leading the top three provinces of smoke particle emissions.

44 Flux concentrations of primarily emitted smoke PM_{2.5} that were calculated using
45 box-model method based on five versions of emission inventories all exceed the
46 carcinogenic risk permissible exposure limits (PEL). The health impacts and
47 health-related economic losses from the smoke PM_{2.5} short-term exposure were
48 assessed. The results show that China suffered from 7836 (95 % CI: 3232, 12362)
49 premature mortality and 7267237 (95 % CI: 2961487, 1130784) chronic bronchitis
50 in 2012, which led to 8822.4 (95 % CI: 3574.4, 13034.2) million US\$, or 0.1 % of
51 the total GDP losses. We suggest that percentage of open burnt crop straws at
52 post-harvest period should be cut down by over 97 % to ensure risk aversion from
53 carcinogenicity, especially the North Plain and the Northeast, where the emissions
54 should decrease at least by 94% to meet the PEL. Under such emission control, over
55 92 % of the mortality and morbidity attributed to agricultural fire smoke PM_{2.5} can
56 be avoided in China.

57 **Key words:** agricultural straw burning, aerosol chamber, smoke particle, emission
58 factor, emission inventory, health effect, emission control policy

59 **1 Introduction**

60 Biomass burning (BB) is a significant source of particulate- and gaseous- pollutants
61 (Andreae and Merlet, 2001; Clarke et al., 2007; Ram et al., 2011; Saikawa et al.,
62 2009a; Tian et al., 2008). It was estimated that open burning of biomass contributed
63 approximately 40% of the globally averaged annual submicron black carbon (BC)
64 aerosol emissions and 65 % of primary OC emissions (Bond et al., 2013). China is
65 the major contributor that bears over 24 % of global emissions of carbonaceous
66 aerosols, especially from agricultural field burning, about 0.04~0.5 Tg EC and
67 0.4~2.1 Tg OC are released annually (Bond, 2004; Cao et al., 2006; Qin and Xie,
68 2012; Saikawa et al., 2009), resulting in great radiative forcing, air quality
69 deterioration, visibility reduction, premature mortality, and economic loss regionally
70 and globally (Bølling et al., 2009; Bond et al., 2013; Huang et al., 2014; Janssen et
71 al., 2011; Rosenfeld, 2006; Saikawa et al., 2009; Shindell et al., 2012).

72 BB also represents one of the most uncertainties in the emission, climate effect,
73 and public health assessments, which finally relies on the uncertainties in detailed
74 chemical emissions or related properties and burning activities like strength or
75 percentage of biomass fuel burned (Tian et al., 2008; Andreae and Merlet, 2001;
76 Levin et al., 2010). For example, studies have focused on OC and EC emissions due
77 to their specific optical properties (Bond et al., 2013; Cao et al., 2006; Qin and Xie,
78 2012; Ram et al., 2011). OC like sulfate and nitrate can cool the atmosphere by
79 increasing the Earth's reflectivity, however, smoke OC on the other hand has been
80 treated as brown carbon to exhibit pronounced light absorption character (Chen et al.,
81 2015; Ackerman, 2000; Chakrabarty et al., 2010; Christopher et al., 2000). The
82 coated or internal mixed sulfate or nitrate can act as lens to enhance the light
83 absorption activity of BC (Zhang et al., 2008b), probably also the activity of brown
84 carbon (Chen et al., 2015). However, primary emissions for OC, EC, and alkali
85 components are confused and have a wide range (Sen et al., 2014; Cao et al., 2006;
86 Hayashi et al., 2014), and some study still took OC with negative forcing activity
87 (Saikawa et al., 2009; Shindell et al., 2012). Besides, smoke EC is consisting of soot

88 and char, and soot-EC has a higher light-absorption potential compared to char-EC
89 (Arora and Jain, 2015; Reid et al., 2005a). Division and quantification of char- and
90 soot-EC emissions for biomass burning are understudied (Arora and Jain, 2015; Han
91 et al., 2009). Moreover, other components like organic acids, amines, phenols, and
92 mineral elements that enable CCN activity or endow health hazard of smoke aerosol
93 are also deficient, variable, or outdated, which may hinder our overall understanding
94 of biomass burning contributions and also atmospheric process of smoke particles
95 (Li et al., 2015; Akagi et al., 2011; Chan et al., 2005; Dhammapala et al., 2007a; Ge
96 et al., 2011; Reid et al., 2005a, b).

97 Studies using carbon mass-balance (CMB) and pollutant concentration-chamber
98 volume quantification are the two common methods to derive the emission factors
99 for biomass burning aerosols (Akagi et al., 2011; Li et al., 2007; Zhang et al., 2008a).
100 Carbonaceous and inorganics components of smoke particles not only vary with
101 biomass issues (fuel types, water content, or burning strength), but also relate to
102 burning condition and environment (flaming or smoldering, field burning or
103 laboratory simulation), extent of aging, sampling methods, and measurement
104 technologies (Grieshop et al., 2009; Hayashi et al., 2014; Reid et al., 2005b).
105 Comparing to field observations that are closer to the actual burning (Li et al., 2007;
106 Akagi et al., 2011; Rose et al., 2011; Saffari et al., 2013), laboratory studies have a
107 definite advantage over field burning research in emission analysis (Jayarathne et al.,
108 2014; Sun et al., 2016; Zhang et al., 2008a). For example, the environment, amount
109 of fuel, and burning conditions can be precisely controlled, the contamination from
110 ambient atmosphere to the emissions can be excluded, and chemical compositions at
111 different aging extent can be quantified using aerosol chamber system (Li et al.,
112 2015, 2016; Aurell et al., 2015; Dhammapala et al., 2007b).

113 The activity rates of biomass burning (burning rate of biomass fuels) are also
114 response to the great uncertainties in the emission estimates (Sun et al., 2016; Zhang
115 et al., 2008a). Seldom study ever focused on the burning rates, and the limited data
116 were treated as simplex constant or dynamic values in many studies of emission
117 estimation in a certain year or for annual variations with a long time scales, thus,

118 significant difference among the results were founded (Qin and Xie, 2011, 2012;
119 Zhang et al., 2011; Zhao et al., 2012). For instance, Cao et al. (2006, 2011) estimated
120 primary smoke carbonaceous materials emissions for 2000 and 2007 in China with
121 same field burning rates, the results were almost the same for the two year with
122 103-104 Gg yr⁻¹ BC and 425.9-433.3 Gg yr⁻¹ OC emitted. He et al. (2011b) found
123 the declining trends in biomass burning emissions in the Pearl River Delta for the
124 period 2003-2007 based on constant activity data of burning rates. Lu et al. (2011)
125 developed primary carbonaceous aerosol emissions in China for 1996-2010 with
126 time-dependent activity rates extrapolated from 2008 to 2010 based on national
127 fast-track statistic, rapid increase of OC and EC emissions were reported, and OC
128 increased from 1.5 to 2.3 Tg yr⁻¹, BC increased from 418 to 619 Gg yr⁻¹. Qin and Xie
129 (2012) estimated BC emission from crop straw open burning for 1980-2009 with
130 variable burning rates based on peasants' income development, the increasing trend
131 in BC emission was also confirmed, and BC emission increased from 4.3 to 116.6
132 Gg yr⁻¹.

133 As most anthropogenic pollutants are concentrated in submicron particulate
134 matters (PM_{1.0}) (Ripoll et al., 2015), more pronounced relationship of ambient PM_{1.0}
135 to haze formation and adverse health effect has been reported (Huang et al., 2003;
136 Roemer et al., 2001; Shi et al., 2014). Nevertheless, associated chemical
137 characterization of PM_{1.0} is still undefined (Li et al., 2015; Safai et al., 2013; Cheng
138 et al., 2006). The study of source-specific PM_{1.0} chemical compositions and
139 emissions are necessary to replenish database for contribution assessment and model
140 application in atmospheric chemistry, climate changes, and public health evaluation.

141 The emission inventories and forecasting in the emissions of atmospheric
142 pollutants have been widely studied, and the incurred mortality, climatic effect, and
143 economic loss have also been estimated (Ostro and Chestnut, 1998; Saikawa et al.,
144 2009; Shindell et al., 2012), based on which the emission control policies were
145 proposed. Shindell et al. (2012) considered ~400 control measures in tropospheric
146 BC and O₃ emissions for the benefit of global or regional human health and food
147 security, and 14 optimal measures targeting CH₄ and BC emissions were identified.

148 Saikawa et al. (2009) compared different scenarios of OC, EC, and sulfate emissions
149 in China in 2030, concluding that maximum feasible reduction may avoid over
150 480000 premature deaths in China and decrease the radiative force from -97 to -15
151 mW m^{-2} globally. Wang et al. (2008a) reported field burning restriction may save
152 about 5 billion dollars losses from biological resource and air pollution. However,
153 the generalized strategies in emission reduction were inadequate and lack actual
154 practicality (Streets, 2007; Lin et al., 2010).

155 In this study, burning experiments with five major agricultural straws were
156 conducted using a combustion stove in combination with an aerosol chamber system.
157 Accurate compositions and emission factors for SPM in $\text{PM}_{1.0}$ and $\text{PM}_{2.5}$ were
158 characterized and established. Afterwards, up-to-date emissions for agricultural open
159 burning aerosol in 2012 were developed, health and health-related economic impacts
160 from smoke $\text{PM}_{2.5}$ exposure were also assessed. Finally, emission reduction strategy
161 that was implemented in field burning rate control for the carcinogenic risk concern
162 was proposed, which should help establish the policy and provide an idea for the
163 emission control.

164 **2 Methodology**

165 An overview of the research procedures including emission factors acquirement and
166 emission inventory calculation is shown in Fig. 1. Tabulation of emission factors is
167 self-established in our laboratory using a combustion stove to simulate open burning
168 and an aerosol chamber to quantify the emissions. Then, we use a bottom-up
169 approach to calculate the emission inventory of agricultural field burning over China
170 mainland based on crop production data in 2012. Emissions for each species are
171 estimated as:

$$172 E_{k,j} = \sum_i A_{k,i} \times EF_{i,j} \quad (1)$$

173 where E_j is emission, $A_{k,i}$ is effective biofuel consumption, and $EF_{i,j}$ is emission
174 factor. k , i , and j indicates region, agricultural residue type, and particulate chemical
175 species.

176 State-of-the art chemical transport and box models were commonly applied to
177 reproduce or simulate the ambient aerosol concentrations (Ram et al., 2011; Reddy
178 and Venkataraman, 2000; Saikawa et al., 2009). In this study, spatio-temporal
179 dynamic box model is used to calculate the emission flux concentration. Regional
180 crop straws are premised to be combusted proportionally only in the fire occurrence
181 days. Dismissing interaction of emitted pollutants in space and time, pollutants will
182 distribute uniformly in a space covering an area of specific region with mixing
183 height of 0.5 km (atmospheric boundary layer). The flux concentration of
184 agricultural burning smoke can be calculated by Eq. (2):

$$185 \quad C_{k,j} = \frac{E_{k,j}}{S_k \times h \times T_k} \quad (2)$$

186 in Eq. (2), $C_{k,j}$ is flux concentration of smoke aerosol, S_k is regional area, h is
187 boundary layer height, T_k is agricultural field fire duration time.

188 **2.1 Aerosol chamber work and emission factors**

189 **2.1.1 Crop straws**

190 Five kinds of representative crop residues were used for the burning experiments, i.e.,
191 wheat, rice, corn, cotton, and soybean straws. The straws were collected based on
192 regional features of agricultural planting, winter wheat straws were collected from
193 Anhui province, late rice straws from Shanghai, corn straws from Henan province,
194 cotton and soybean residues from Xinjiang. All straws were stored under dark, airy,
195 and cooling condition. Prior to the burning experiment, the dirt and weeds were
196 removed, then straws were dehydrated (at 100 °C for 24 h) to minimize effect of the
197 water content on the burning and pollutant emissions, as study found pollutants
198 emissions and combustion efficiencies (CE) are response to water content, increased
199 moisture content enhances the emissions but also alter the chemical compositions of
200 smoke aerosols (Reid et al., 2005b; Aurell et al., 2015; Hayashi et al., 2014).
201 Although straws in the field are not well dried and moisture contents vary with
202 weather, ventilation, and storing times, for the convenience of practical application

203 and comparison of burnings and emissions, water contents of the straws were
204 controlled within 2 %, which has been applied in many studies (Hayashi et al., 2014;
205 Huo et al., 2016; Li et al., 2015; Oanh et al., 2011; Zhang et al., 2008a, 2011). The
206 dry straws were then cut to a length of approximately 10 cm and weighted 10.0 g per
207 serving.

208 **2.1.2 Burning experiments**

209 The experiments were conducted using an aerosol chamber system (Fig. S1 in
210 supplement information, SI), which was loaded in a temperature-controlled room
211 (18-22 °C, 40%-60% RH). A stainless combustion stove was self-deigned to simulate
212 typical field burning of crop straws by automatic ignition with LPG (Liquid
213 petroleum gas) in particular, albeit on a small scale (ignition time less than 0.1 s).
214 10.0 g conditioned residues were sealed in the 0.227 m³ combustion stove in
215 advance, once ignited, the force-ventilation and HEPA filtrated particle-free air were
216 supplied (300 L min⁻¹). The emissions were immediately injected into a clean,
217 evacuated aerosol chamber. The burning last about 1 min and over 1 m³ particle-free
218 air flushed the stove to ensure all the emissions were transferred into the chamber.

219 The chamber was custom-built to quantify the emissions and characterize the
220 physiochemical properties of smoke aerosols, detailed description of the chamber
221 can be found elsewhere (Zhang et al., 2008a, 2011; Li et al., 2015, 2016). Briefly, the
222 chamber has a volume of 4.5 m³ with 0.3 mm Teflon coating on the inner side, a
223 magnetic fan fixed on the bottom to stir the aerosol uniformly, and a hygrocclip
224 monitor (Rotronic, Model IM-4) equipped inside the chamber to measure the
225 temperature and relative humidity of the aerosol. Before experiment, the chamber
226 was flushed with particle-free air for 6 h, oxidized by high concentration ozone (~3
227 ppm) for 12 h, then flushed and vacuumized, filled with pure dry air to 80 KPa for
228 use. The emissions from straw burning were aspirated into the chamber till room
229 pressure, afterwards, size measurement and sampling of smoke aerosols were
230 conducted from the chamber. For each type of straw, four burning experiments were

231 conducted. The unburned residues were weighted and deducted from 10.0 g after
232 each test.

233 Modified combustion efficiency (MCE) for each burning was monitored with CO
234 and CO₂ measuring to determine the burning phase and ensure the repeatability.
235 MCE is defined as $\Delta\text{CO}_2/(\Delta\text{CO}_2+\Delta\text{CO})$, where ΔCO_2 and ΔCO are the excess molar
236 mixing ratios of CO₂ and CO (Reid et al., 2005b). A gas-chromatograph (GC, model
237 930, Shanghai, Hai Xin Gas Chromatograph Co., LTD) equipped with a flame
238 ionization detector, an Ni-H convertor, and a stainless steel column (2 m long)
239 packed with 15% DNP was used to measure CO and CO₂ concentrations in the
240 chamber. And MCE were 0.89-0.96 for all the experiments, indicating flaming
241 combustion dominated, which were comparable to that in the field burning (Li et al.,
242 2003; Li et al., 2007).

243 **2.1.3 Size and morphology of smoke aerosol**

244 Size distribution (10 nm-10 μm) of smoke particles was measured using a
245 Wide-range Particle Spectrometer (WPS, Model 1000XP, TSI, USA), which has
246 been described by Li et al (2015). Briefly, WPS integrates the function of scan
247 mobility particle sizer (SMPS) and laser particle sizer (LPS), 0.3 L min⁻¹ flow is
248 introduced to SMPS part to classify mobility size from 10 nm to 500 nm in 48 bins,
249 and 0.7 L min⁻¹ flow is introduced to LPS part to measure aerodynamic diameter
250 from 350 nm to 10 μm in 18 bins. Particle density and refractive index were set as
251 1.0 g cm⁻³ and 1.45, and charge correction mode was on for the measurement. A
252 diffusion dryer tube (45 cm in length) filled with descant-silica gel was set prior to
253 the inlet of WPS. Before experiment, WPS was calibrated with certified polystyrene
254 latex spheres (PSL, 40, 80, and 220 nm, Duke Scientific).

255 SPM from the 5 types crop straws burning were sampled onto copper grids coated
256 with carbon film (carbon type-B, 300-mesh copper, Tianld Co., China) using a
257 single-stage cascade impactor with a 0.5 mm diameter jet nozzle at a flow rate of 1.0
258 L min⁻¹. The sampler has a collection efficiency of 100 % at 0.5 μm aerodynamic

259 diameter. More information about the cascade impactor can be found elsewhere (Fu
260 et al., 2012; Hu et al., 2015). Then, a JEOL-2010F field emission high-resolution
261 transmission electron microscope (FE-HRTEM) coupled with an oxford
262 energy-dispersive X-ray spectrum (EDX) was applied to investigate the morphology,
263 composition, and mixing state of individual particles.

264 **2.1.4 Chemical sampling and analysis**

265 PM_{1.0} and PM_{2.5} samples for each burning were collected onto pretreated quartz filter
266 of 90 mm in diameter (Tissuquartz, Pall Corp., USA) from the chamber using a
267 high-volume Particle Sampler (HY-100, Qingdao Hengyuan S.T. Development Co.,
268 Ltd) operating at 100 L min⁻¹. Each filter sampling duration time is 5 min, and total
269 44 samples (including 4 blank samples) were gathered. The quartz microfiber filters
270 were prebaked for 8 h at 450 °C to eliminate contamination. Before and after the
271 sampling, the filters were weighted using a balance (Sartorius BP211D) with an
272 accuracy of 10 µg, and the filters were balanced in an electronic desiccator (40 %
273 RH, 22 °C) for 24 h before usage. After weighting, the loaded filters were stored at
274 -20 °C in a refrigerator for further analysis.

275 Water soluble species including general inorganic ions (ions: F⁻, Cl⁻, NO₂⁻, NO₃⁻,
276 SO₄²⁻, Na⁺, NH₄⁺, K⁺, Ca²⁺, Mg²⁺), organic acids (CH₃COOH, HCOOH, C₂H₂O₄,
277 CH₃SO₃H), and seven protonated amines (MeOH⁺, TeOH⁺, MMAH⁺, DMAH⁺,
278 TMAH⁺, MEAH⁺, and DEAH⁺ for short, corresponding to monoethanolaminium,
279 triethanolaminium, monomethylaminium, dimethylaminium, triethylaminium,
280 monoethylaminium, and diethylaminium) were measured from 1/4 of each filter
281 with ion chromatography (IC, Model 850 Professional IC, Metrohm, USA) consists
282 of a separation column (Metrosep A Supp 7 250/4.0 for anion and organic acids,
283 Metrosep C-4 150/4.0 for cation, and Metrosep C4-250/4.0 for water soluble
284 aminiums). Sampled filters were ultrasonically extracted with 15.0 mL deionized
285 water (Mili-Q water, 18.2 MΩ cm), extracted solutions were filtrated using 0.2 µm
286 filters before injected into IC for measurement. Detection limits (DLs) for the ions

287 and aminiums were within $0.5\sim 3.5\text{ ng mL}^{-1}$, the correlation coefficients for all
288 calibration curves were better than 0.99, and recovery rates for aminiums were in the
289 range of 93%~106% (see in SI, Table S1). Details for the aminium measurements
290 can be found in the work of Tao et al. (2016).

291 1/4 of each filter was acid dissolved to measure the selected elements (As, Pb, Cr,
292 Cd, Ni, V, Zn, Al), of which As, Zn, Pb, Cr, Cd, and Ni are USEPA priority
293 controlled pollutants (Wu et al., 2011). The smashed filters were digested at $170\text{ }^{\circ}\text{C}$
294 for 4 h in high-pressure Teflon digestion vessel with 3.0 mL concentrated HNO_3 , 1.0
295 mL concentrated HClO_4 , and 1.0 mL concentrated HF. Afterwards, the almost dry
296 solution was diluted and characterized using Inductively Coupled Plasma Optical
297 Emission Spectrometer (ICP-OES, Atom Scan 2000, JarroU-Ash, USA). The
298 following wavelength lines of the ICP-OES analysis were used: As 189.042, Pb
299 220.353, Cd 228.802, Cr 205552, Ni 231.604, V 311.071, Zn 206.191, and Al
300 394.401. All reagents used were of highest grades, and recovery tests were
301 conducted with standard additions, recoveries of each element were in the range of
302 93%~102% (see in SI, Table S1).

303 Another 1/4 of each filter was ultrasonically double extracted with 15.0 mL
304 HPLC-grade CH_2Cl_2 . The extracts were then condensed with rotary evaporator and
305 quantified to 1.0 mL. 16 targeted PAHs (2-ring, naphthalene (Nap); 3-ring,
306 acenaphthylene (Ac), acenaphthene (Ace), fluorene (Fl), phenanthrene (Phe),
307 anthracene (Ant); 4-ring, fluoranthene (Flu), pyrene (Pyr), benzo[a]anthracene (BaA),
308 chrysene (Chr); 5-ring, benzo[b]fluoranthene (BbF), benzo[k]fluoranthene (BkF),
309 benzo[a]pyrene (BaP), dibenzo[a,h]anthracene (DBA); and 6-ring: indeno[1,2,3-cd]
310 pyrene (IP), benzo[ghi]perylene (BghiP)) and 5 selected phenols (phenol,
311 2-methoxyphenol, 4-ethylphenol, 4-ethyl-2-methoxyphenol, 2,6-dimethoxyphenol)
312 were measured from the concentrated extracts using an Agilent 6890 Series gas
313 chromatography system coupled with a HP 5973 Mass Selective Detector (GC-MS,
314 Agilent Technologies, Wilmington DE) . A DB-5ms (30 m \times 0.32 mm \times 0.25 mm,
315 Agilent 123-5532) column was installed. The temperature programs were presented
316 as follows: initially at $40\text{ }^{\circ}\text{C}$, hold for 4 min, to $150\text{ }^{\circ}\text{C}$ at $20\text{ }^{\circ}\text{C min}^{-1}$, then to $280\text{ }^{\circ}\text{C}$

317 at 5 °C min⁻¹, hold for 10 min. The interface temperature was kept at 280 °C, the MS
318 was operated in electron impact mode with an ion source temperature of 230 °C, and
319 the high-purity helium (99.999%) carrier gas was maintained at a constant pressure
320 of 16.2 psi with a flow of 2.0 mL min⁻¹. The calibration curves were optimized to be
321 better than 99.9%. Prior to the measurements, PAHs and Phenols recovery studies
322 were undertaken, and recoveries were acceptable with rates of 82%~99% (see in SI,
323 Table S1). In addition, Phenanthrene-d10 (Phe-d10) as internal standard surrogate
324 was added into the PAHs mixture, recovery rate of which was 94%.

325 Organic carbon (OC) and elemental carbon (EC) were measured with the rest
326 quartz filters using a carbon analyzer (Sunset laboratory Inc., Forest Grove, OR)
327 based on the thermal-optical transmittance (TOT) method with a modified
328 NIOSH-5040 (National Institute of Occupational Safety and Health) protocol. Four
329 organic fractions (OC1, OC2, OC3, and OC4 at 150, 250, 450, and 550 °C,
330 respectively), PC fraction (a pyrolyzed carbonaceous component determined when
331 transmitted laser returned to its original intensity after the sample was exposed to
332 oxygen), and three EC fractions (EC1, EC2, and EC3 at 550, 700, and 800 °C,
333 respectively) are produced. And OC is technically defined as OC1 + OC2 + OC3 +
334 OC4 + PC, while EC is defined as EC1 + EC2 + EC3 - PC (Seinfeld et al., 2012).
335 The instrument detection limits for total OC and EC that deposit on the filter are 0.25
336 and 0.12 µg C cm⁻². Moreover, environmental EC in aerosol is a mixture of
337 compounds from slightly charred, biodegradable materials to highly condensed and
338 refractory soot, different EC materials have distinct different thermodynamic
339 properties, study found char-EC decomposes much rapidly than soot when exposed
340 to chemical and thermal oxidation, e.g., EC decomposition temperatures in air
341 increased from ~520 °C for char to ~620 °C for soot, and exceeded 850 °C for
342 graphite, thus, regarding to different oxidation temperatures, Han et al (2007, 2009)
343 furtherly differentiated char and soot-EC from EC measurement as EC2 + EC3
344 equals to soot-EC, and the rest is char-EC.

345 The quality of the data above was guaranteed by standard materials calibration,
346 recovery rate, and operational blank correction. And blank levels were less than 5%

347 of the measured values for all the species.

348 **2.1.5 Calculation of emission factors**

349 The emission quantities derived from the experiment were converted into quantities
350 per unit weight of initial residues as emission factor (EF, unit: g kg⁻¹), which can be
351 calculated from the direct method with effective filter sampling weight, chamber
352 volume, and effective amount of crop straw consumed (Dhammapala et al., 2006,
353 2007a, b; Zhang et al., 2008a), or alternatively from the carbon mass balance method
354 (CMB) via conservation of Carbon in biomass, disregarding the weight of biomass
355 that burnt (Dhammapala et al., 2006; Li et al., 2007). EFs determined from these two
356 methods were found to be in good agreement (Dhammapala et al., 2006),
357 nevertheless, CMB method needs more auxiliary information (e.g., concentrations of
358 CO, CO₂, CH₄, non-methane hydrocarbons, and also particulate carbons), which
359 may result in data redundancy and uncertainty propagation, hence we applied the
360 direct method to calculate EFs in this work. To be more accurate, influence of wall
361 loss and makeup air dilution on smoke particles sampling from the chamber were
362 considered and corrected, details see in SI.

363 In this study, duration for each test (burning, chamber condition, size
364 measurement, and filter sampling) was controlled within 20 min, therefore, the
365 physicochemical processes of pollutants in the chamber can be negligible, and
366 smoke aerosols we measured were primary emissions.

367 **2.2 Emission inventory calculation**

368 **2.2.1 Agricultural field fire survey**

369 Fire sites over China from 2011 to 2013 were statistically analyzed, and the data of
370 mainland agricultural fire sites was derived from the daily report of the Ministry of
371 Environmental Protection of China (MEPC) (website: <http://hjj.mep.gov.cn/jgjs/>).
372 Agricultural fire sites were screened out from MODIS (Moderate Resolution

373 Imaging Spectroradiometer) daily fire products (1 km × 1 km resolution level 3
374 hotspot) using a high resolution real time land use based on geography information
375 system (GIS). Spatial and temporal distributions of fire sites were displayed in Fig.
376 S2 (SI), over 5000 fire sites were allocated into two prominent burning periods
377 corresponding to summer (May to July) and autumn (September to November)
378 harvests, and field burning last 54 days and 60 days on statistical average during the
379 two harvests. In the North of China, open burning occurred primarily in autumn,
380 while temporal-character of field fires was not obvious in the North Plain and the
381 Center of China, where field fires can be observed frequently during the whole
382 investigation time.

383 **2.2.2 Crop straw production**

384 Crop straw production was generally derived from annual or monthly crop production
385 by multiplying crop-specific ratios of residue-to-production (He et al., 2011b; Cao et
386 al., 2011; Zhao et al., 2012). In this study, crop productions were furtherly classified
387 into summer harvest and autumn harvest productions according to field fire sites
388 analysis and traditional seasonal planting and harvesting. The amount of straw
389 produced was calculated by Eq. (3):

$$390 M_{t,k,i} = P_{t,k,i} \times r_i \times H_{t,k,i} \times D_i \quad (3)$$

391 in which M is mass of crop straws produced; P is annual crop-specific amount of
392 crop production; r is the residue-to-production ratio; D is the dry matter
393 content; $H_{t,k,i}$ is production ratio of crop i at region k during summer or autumn
394 harvest period t.

395 Province-level crop production data of wheat, rice, corn, cotton, and soybean were
396 taken directly from the China Yearbook 2013 (National Bureau of Statistics of China,
397 NBSC, 2013). Crop-specific residue-to-production ratios were cited from Chinese
398 Association of Rural Energy Industry (Wang and Zhang, 2008; data available at
399 <http://www.carei.org.cn/index.php>, in Chinese). Dry matter contents of crop straws
400 were referred to He et al. (2011b) and Greenhouse Gas Inventory Reference Manual

401 (IPCC, 2007). The parameters of residue-to-production ratios and dry matter
402 contents were summarized in Table S2 (SI). The regional crop production ratios in
403 summer and autumn harvests were listed in Table S3 (SI).

404 **2.2.3 Field burning rate**

405 Uncertainty of emission estimations mostly relies on intangibility of straw open
406 burning rate (Zhao et al., 2012; He et al., 2011b). However, regional or national
407 percentage of straw open burned was seldom studied, and the limited data were
408 outdated and variable. The available studies indicate national field burning rate of
409 crop straws range from 15.2% to 27.2% in China (Daize, 2000; Wei et al., 2004;
410 Zhang et al., 2008a), and more detailed studies indicate about 31.9% of the crop
411 straw burned in the Pearl River Delta from 2003 to 2007 (He et al., 2011b), while the
412 corresponding figures were almost 100% for the Huabei region in 2003 (Zhao et al.,
413 2012). Two versions of province-level field burning rates were commonly used, one
414 was from Cao et al. (2005, 2006, 2011), who deduced the rates based on regional
415 economic level, and the proposal of the rates to be proportional to peasants' income
416 was confirmed later, the rates were first used to calculate open burning emission in
417 2000. The other version was reported by Wang and Zhang (2008), they obtained
418 provincial percentage of residue open burnt via field survey in 2006. Herein, the two
419 versions were both applied directly into the emission estimation of 2012 in this work
420 and named as business-as-usual scenarios (BAU, BAU-I from Cao et al. and BAU-II
421 from Wang and Zhang in specific).

422 In fact, the burning rates should be dynamic parameters that been influenced by
423 industrial structure, government policy orientation, or public awareness. With crop
424 yields increase and energy consumption structure changes in rural areas, more straws
425 will be discarded and burnt in the field. Nonetheless, rigorous agricultural fire policy
426 may still suppress the condition worsen as it worked during 2008 for Beijing
427 Olympics and 2010 for Shanghai Expo (Huang et al., 2013; Cermak and Knutti,
428 2009; Wang et al., 2010). Qin and Xie (2011; 2012) ever deduced year specific open

429 burning rates in different zone for the period of 1980-2009 according to their
 430 respective peasant income changes in a certain year on the basis of peasant income
 431 and burning rates in 2006. However, the simple linear relationship should be doubted,
 432 as great increase in per capita income after 2006 will surely overestimate the burning
 433 rates. We supposed that the burning rates were inverse proportional to peasants'
 434 agricultural income proportion (AIP), without considering the policy or potential
 435 gain or loss related to agricultural residue treatment. Thus the burning rates
 436 established in 2000 and 2006 from Cao et al. (2005) and Wang and Zhang (2008) can
 437 be converted into that of 2012 based on economic data from equation below:

$$438 \quad R_{k,2012} = \frac{I_{k,2012}}{AI_{k,2012}} \times \frac{AI_{k,y}}{I_{k,y}} \times R_{k,y} \quad (4)$$

439 where R is agricultural straw filed burnt rate, $I_{k,y}$ is peasants' annual income, $AI_{k,y}$
 440 is peasants' annual agricultural income. y indicates reference year (2000 for BAU-I,
 441 and 2006 for BAU-II). $I_{k,y}$ and $AI_{k,y}$ can be found or calculated from China
 442 Yearbook and China Rural Statistic Yearbook (NBSC, 2004-2013).

443 The versions of converted rates based on primary industry level were called
 444 Economic Models I and II (EM -I and EM-II in short) corresponding to BAU-I and
 445 BAU-II. Besides, in 2013, the National Development and Reform Commission of
 446 China published the Chinese agricultural straw treatment report of 2012 (NDRC,
 447 [2014] No.516, data available at <http://www.sdpc.gov.cn/>, in Chinese) for the first
 448 time. The percentages of crop residues discarded in the report were applied in our
 449 estimation, which was called NDRC version.

450 **2.2.4 Emission and flux concentration**

451 From above study, emission of SPM pollutants can be calculated by recount of Eq.
 452 (1) to get Eq. (5) as it was showed below:

$$453 \quad E_{t,k,j} = \sum_i M_{t,k,i} \times R_k \times f_i \times EF_{i,j} \quad (5)$$

454 where $E_{t,k,j}$ is emission amount of chemical species j at region k during harvest
 455 period t; f_i is burning efficiency, the crop specific values were cited as 0.68 for

456 soybean residue and 0.93 for the rest four straws (Zhang et al., 2011; Wang and
457 Zhang, 2008; Zhang et al., 2008a; Koopmans et al., 1997). Thus, flux concentration
458 of corresponded pollutants can be also assessed from box model as mentioned in
459 front.

460 **2.3 Estimate health impacts and health-related economic losses**

461 **2.3.1 Carcinogenic risk of Smoke Particulate Matter (CR_{SPM})**

462 Apart from the enormous climatic effects of smoke particle emissions, new
463 epidemiological and toxicological evidence have also linked carbonaceous aerosol to
464 cardiovascular and respiratory health effects according to the World Health
465 Organization (Bruce et al., 1987; IPCC, 2007). Here, we present the fuel-specific
466 carcinogenic risk of SPM (CR_{SPM}, unit: per $\mu\text{g m}^{-3}$) to assess health hazard from
467 agricultural straw burning particles and help source-specific air quality control. The
468 cancer risk attributed to inhalation exposures of smoke PM_{2.5} from crop straw i
469 burning was calculated as:

$$470 \text{CR}_i = \sum_j f_j \times \text{UnitRisk}_j \quad (6)$$

471 where f_j is mass fraction of individual species j in smoke PM_{2.5}, UnitRisk_j is
472 corresponded unit carcinogenic risk value of species j extracted from database
473 provided by the Integrated Risk Information System (IRIS), California
474 Environmental Protection Agency (CEPA), and related documents (Bruce et al.,
475 1987; Burkart et al., 2013; Tsai et al., 2001; Wu et al., 2009, 2011).

476 CR_i is estimated based on dose addition model of selected hazardous air
477 pollutants (HAPs) including USEPA priority pollutants of PAHs and heavy metals.
478 And UnitRisk values of the selected HAPs presented in Table S4 (SI). Synergistic
479 interactions among pollutants are dismissed, albeit possible. The cancer risk of
480 chromium is adjusted by multiplying a factor of 0.2, assuming that only 20% Cr
481 measured is in the toxic hexavalent form (Bell and Hipfner, 1997). Benzo[a]pyrene
482 (BaP) is used as an indicator compound of carcinogenicity, legally binding threshold

483 of BaP in most countries ranges from 0.7 to 1.3 ng m⁻³, corresponded carcinogenic
 484 risk of BaP is about 1.1×10⁻⁶ per ng m⁻³ (Bruce et al., 1987; Burkart et al., 2013).
 485 Thus, one in million level of carcinogenic potential is frequently used to identify
 486 risks of concern in public health and environmental decision making, and
 487 permissible exposure limits (PEL, unit: µg m⁻³) of crop straw burning particles can
 488 be estimated as:

$$489 \text{ PEL}_i = \frac{10^{-6}}{\text{CR}_i} \quad (7)$$

490 **2.3.2 Human exposure and health impacts**

491 Robust relationship between surface PM_{2.5} and health effects has been revealed and
 492 confirmed by many studies (Pope et al., 2004; Wong et al., 2008). PM_{2.5}-related
 493 health endpoints are composed of a range of elements from sub-clinical effects to the
 494 onset of diseases and the final death (Davidson et al., 2005). In this study, incidence
 495 of commonly studied endpoints like premature mortality, respiratory and
 496 cardiovascular hospital admissions, and chronic bronchitis from primary emitted
 497 smoke PM_{2.5} short-term exposure were assessed using the Poisson regression model,
 498 shown as below (Guttikunda and Kopakka, 2014):

$$499 \Delta E = \Delta \text{Pop} \times \text{IR} \times \left(1 - \frac{1}{e^{\beta \times \Delta C}}\right) \quad (8)$$

500 where ΔE represents the number of estimated cases of mortality and morbidity, ΔC
 501 is the incremental concentration of particulate matter (PM) or flux concentration;
 502 ΔPop is the population exposed to the incremental particulate concentration of ΔC;
 503 IR is short for incidence rate of the mortality and morbidity endpoints, and β is the
 504 coefficient of exposure-response function, defined as the change in number case per
 505 unit change in concentration per capita.

506 Concentration-response function and incidence rate of each health endpoint are
 507 important in health impacts evaluation and they have variation for different
 508 population and regions (Yang et al., 2012; Wong et al., 2008). Here, the variance for
 509 sex and ages were neglected. Region-specific exposure-response coefficients for
 510 individual mortality were summarized from previous studies, as presented in Table

511 S5 (SI). The coefficients for individual respiratory and cardiovascular hospital
512 admission, and chronic bronchitis were cited as 1.2 %, 0.7 %, and 4.4 % (per 10 μg
513 m^{-3} , 95% CI) from Aunan and Pan's work (Aunan and Pan, 2004). This is the case
514 because seldom studies ever confirmed these topics in China. Region-specific
515 mortality and hospitalization IRs were taken from statistical reports authorized by
516 National Health and Family Planning Commission of the People's Republic of China
517 (NHFPC, 2013), and morbidity of chronic bronchitis were defined as 13.8 % based
518 on the forth national health survey, which was released by the Chinese Ministry of
519 Health in 2008 (CMH, 2009).

520 2.3.3 Economic valuation of the health impacts

521 The economic losses of the health impacts associated with smoke $\text{PM}_{2.5}$ exposure in
522 2012 were further evaluated. The amended human capital (AHC) approach was
523 employed to calculate the unit economic cost of premature mortality. The commonly
524 applied AHC method uses per capita GDP to measure the value of a statistical year
525 of life (IBRD and SEPA, 2007) based on Eq. (9). It can be used as a social statement
526 of the value of avoiding premature mortality and estimates human capital (HC) from
527 the perspective of entire society, neglecting individual differences (Hou et al., 2012).

$$528 \text{HC}_k = \frac{\text{GDP}_k}{\text{POP}_k} \times \sum_{i=1}^{\tau} \frac{(1+\alpha)^i}{(1+\gamma)^i} \quad (9)$$

529 GDP_k and POP_k are gross domestic production and population of target region k
530 that were reported in the statistical yearbook in 2012; α and γ are economic
531 parameters referring to national GDP growth rate and social discount rate, which
532 were 7.7 % and 8.0 % in 2012 from National Bureau of Statistics of China (NBSC,
533 2013, data available at <http://www.stats.gov.cn/tjsj/ndsj/>, in Chinese). τ is the
534 life-expectancy lost due to aerosol pollution, and 18 year of life was widely applied
535 (Hou et al., 2012). The annual exchange rate of US dollar to RMB was 6.31 in 2012.
536 One can deduce the HC values of the provinces, municipalities, and autonomous
537 regions in the country, and the calculated regional HC values were listed in Table S6
538 (SI). In this paper, the cost of respiratory, cardiovascular hospital admissions, and

539 chronic bronchitis were 632.2, 1223.4, and 948.6 US\$ per case in 2012, which were
540 derived from the national health statistical reports (NHFPC, 2013).

541 The regional and national health-related economic loss from smoke PM_{2.5}
542 exposure can be calculated based on the excess mortality and morbidity multiplied
543 by the corresponding unit economic values.

544 **3 Result**

545 **3.1 Particulate chemical compositions and emission factors**

546 **3.1.1 Organic carbon and elemental carbon**

547 An overview of particulate chemical compositions for smoke PM_{2.5} and PM_{1.0} is
548 pie-graphically profiled in Fig. 2, and the corresponded emission factors are given in
549 Table 1-2 (detailed EFs for elements, PAHs, and Phenols in Table S7 and S8, SI).
550 From multivariate statistical analysis ($P < 0.05$ at 95% CI), significant differences of
551 chemical compositions and emissions in size range and fuel types can be observed,
552 implying the nonuniform mixing and distribution of particulate pollutants from
553 biomass burning, which is consistent with the conclusion from Lee et al. (2015) and
554 Giordano et al. (2015). EFs of particulate species from this study were compared
555 with that from literature as summarized in Table 3, since EFs in smoke PM_{1.0} were
556 seldom reported, only smoke PM_{2.5} or total particulate matter emissions were
557 collected, which were comparable with the results in this work. EFs of smoke PM_{2.5}
558 and PM_{1.0} were in range of 3.25~15.16 and 3.04~13.20 g kg⁻¹ for the five kinds of
559 crop straws, a high ratio of PM_{1.0}/PM_{2.5} was observed to be over 90 wt.%, which was
560 in line with size distribution analysis of smoke particles given in Fig. S3 (SI). Li et al.
561 (2007) measured the emissions from field burning of crop straws via CMB method,
562 PM_{2.5} EFs for wheat and corn straw were estimated to be 7.6 ± 4.1 and 11.7 ± 1.0 g
563 kg⁻¹ (dry basis, MCE > 0.9), which were higher and presented more uncertainties
564 than our result. As study ever found a positive relationship between particulate EFs
565 and moisture content of agricultural residue (Hayashi et al., 2014), it was reasonable

566 that combustion of the dehydrated crop straw produced less smoke aerosol in this
567 work. Hayashi et al. (2014) measured particulate EFs to be 2.2 and 15.0 g kg⁻¹ for
568 rice and wheat straw of ~10 wt.% moisture content, while corresponded EFs
569 increased to 9.1 and 19.5 g kg⁻¹ when water content of straw was ~20 wt.%, and the
570 linear equations between smoke EFs and straw moisture content were furtherly
571 proposed. However, the simple linearity and its application scope should be doubted,
572 as Hayashi et al. only considered two water content levels (10 wt.% vs 20 wt.%) and
573 disregarded influence of combustion efficiency for the fires. PM_{2.5} EFs given by
574 Dhammapala et al (2006, 2007a, b) were 4.7 ± 0.4 g kg⁻¹ for wheat straw and 12.1 ±
575 1.4 g kg⁻¹ for herbaceous fuel that were burnt using a chamber under flaming phase,
576 and negative response for particulate EFs to combustion efficiency was observed.
577 After all, smoke EFs vary with fires depend on fuel type and moisture, combustion
578 phase, environmental conditions, and some other variables (Reid et al., 2005b).

579 The carbonaceous materials (Organic matter and EC) are dominated in SPM,
580 accounting for about 73.4 wt.% for PM_{2.5} and 71.3 wt.% for PM_{1.0} on average.
581 Organic matter (OM) was converted from OC by multiplying a factor of 1.3 to
582 account for noncarbon materials like oxygen, hydrogen, and other minor species (Li
583 et al., 2007; Li et al., 2015), and Li et al. (2016) ever measured OM/OC ratio as ~1.3
584 for fresh smoke particles via volatility analysis. EFs of EC and OC from this work
585 were consistent with most studies, average OC EFs were 4.21 and 3.58 g kg⁻¹ in
586 smoke PM_{2.5} and PM_{1.0}, and the corresponded EC EFs were 1.09 and 1.01 g kg⁻¹,
587 respectively. These values fell within the ranges (0.9~9.3 g kg⁻¹ for OC and 0.2~1.7
588 g kg⁻¹ for EC) found in other similar sources (Dhammapala et al., 2007; Hayashi et
589 al., 2014; Li et al., 2007; May et al., 2014). Due to the technical limitation and
590 ambiguous artificial boundary, carbon contents of biomass burning particles have
591 vast variability and uncertainty (Lavanchy et al., 1999; Levin et al., 2010). It was
592 ever reported chamber burn study may overestimate EC EFs due to a misassigned
593 OC-EC split for the heavily mass loaded filter samples (Dhammapala et al., 2007b).
594 Moreover, carbon measurement based on TOT method with NIOSH protocol may
595 overestimate OC fraction by sacrificing EC part compared with that of TOR

596 (Thermal-Optical Reflectance) method with IMPROVE program (Han et al., 2016).
597 Mass ratio of OC/EC is a practical parameter to indicate the primary organic aerosol
598 (OA) emission and secondary organic aerosol (SOA) production. The ratio is
599 influenced by burning conditions, source, aging extent, and particle size (Engelhart
600 et al., 2012; Grieshop et al., 2009). Smoke emitted from smoldering fires is
601 OC-dominated while flaming combustion produces more EC, and the discrepancy of
602 OC/EC ratio can be an order of magnitude regarding to different combustion phase
603 (Grieshop et al., 2009). SOA production upon photo-oxidation will enlarge OC/EC
604 ratio, and positive relation between oxidation level of OA loading and OC/EC ratio
605 was reported (Grieshop et al., 2009). Here, OC/EC ratio in primary emissions varied
606 from 2.4 to 6.2 under flaming phase, similar to previous studies (Arora and Jain,
607 2015; Dhammapala et al., 2007a, b; Hayashi et al., 2014; Lewis et al., 2009). The
608 OC/EC ratios were larger in PM_{2.5} with average value of 3.8, while it was 3.6 in
609 PM_{1.0}, indicating more EC resides in PM_{1.0}.

610 EC in smoke particle can be furtherly classified as char-EC and soot-EC based on
611 the distinct different physiochemical properties and formation mechanisms of soot
612 and char (Arora and Jain, 2015; Lin et al., 2011; Reid et al., 2005a; Richter et al.,
613 2000). Both char- and soot-EC represent the major light-absorbing fraction of PM,
614 however, soot-EC exhibits much stronger light-absorbing character and longer
615 lifetime in the atmosphere than char-EC, and merits greater focus in climate research
616 (Arora and Jain, 2015; Han et al., 2010). Char-EC can be distinguished as brown
617 carbon, as carbonaceous materials that are optically between the strongly absorbing
618 soot and non-absorbing organics are operationally defined as brown carbon (Yang et
619 al., 2009; Andreae and Gelencsér, 2006; Gustafsson et al., 2009). Char-EC is formed
620 from solid residues during relative low-temperature combustion, while generation of
621 soot-EC takes place under high-temperature conditions from recondensation and
622 dihydrogen-carbonization of gaseous materials like PAHs (Han et al., 2007, 2009).
623 To our knowledge, seldom study ever reported source specific EFs of char- and
624 soot-EC for crop straw burnings. Here, particulate char- and soot-EC EFs in fine
625 mode were estimated to be 0.56 ~ 1.76 and 0.05 ~ 0.42 g kg⁻¹, while char- and

626 soot-EC EFs in smoke $PM_{1.0}$ were 0.51 ~ 1.67 and 0.06 ~ 0.41 $g\ kg^{-1}$, respectively.
627 Mass ratio of char-EC/soot-EC is a more effective indicator for source identification
628 and apportionment than OC/EC, as atmospheric process can hardly change the mass
629 loading and dissociation of inert EC fractions (Han et al., 2007, 2009, 2010). The
630 ratios of char-EC/soot-EC are also fuel types and PM size dependent. Generally,
631 char-EC/soot-EC is also controlled by combustion mode or even moisture content of
632 biomass fuel, and biomass burning by smoldering at low temperatures results in high
633 char-EC/soot-EC. Chuang et al. (2013) reported char-EC/soot-EC in smoke $PM_{2.5}$
634 was 9.4 ± 3.8 for biomass burning (BB), and Cao et al. (2005) proposed the ratio to
635 be 11.6 for BB sources. These values were larger than the present study, as we
636 estimated char-EC/soot-EC in $PM_{2.5}$ to be 7.28 ± 1.98 on average. It can be
637 explained by different techniques for EC measurement, char-EC and soot-EC were
638 mostly measured using TOR-IMPROVE method, while TOT-NIOSH method used in
639 this study will overestimate PC fraction in OC-EC split, resulting in less char-EC
640 fraction (EC1-PC) and lower char-EC/soot-EC ratio. Nonetheless, the results were
641 still comparable for the two methods (Han et al., 2016).

642 The char-EC/soot-EC ratio was 6.29 in $PM_{1.0}$, which was smaller than that in
643 smoke $PM_{2.5}$, the result indicates that SPM comprises a considerable amount of
644 char-EC and char particle has a larger size than soot, in consistent with the
645 conclusion that soot particles are mainly tens of nanometers in size and cluster
646 together into loose aggregates of hundred nanometers, while char particles were
647 reported to be larger with diameter in the range of 1~100 μm (Arora and Jain, 2015;
648 China et al., 2014; Lin et al., 2011; Wornat et al., 2007). Besides, correlation among
649 the multi-pollutants was analyzed by relevance matrix as shown in Table S9 (SI), the
650 strong positive linear relationship ($R^2 > 0.99$, $p < 0.05$ at 95% CI) between EC and
651 char-EC also confirms the reliable source of biomass burning to produce char-EC
652 (Lin et al., 2011; Arora and Jain, 2015).

653 3.1.2 Water soluble organic acids

654 Smoke particles comprise a considerable amount of water soluble organic acids
655 (WSOA), it was 3.35 wt.% in PM_{2.5} and 3.17 wt.% in PM_{1.0} on average, which was
656 in line with previous work that organic acids measured represent less than 5 wt.% of
657 the total smoke aerosol mass load and favor to partition in larger size (Falkovich et
658 al., 2005; Gao et al., 2003). Acetic acid followed by methanesulfonic acid contributed
659 the most of the measured low molecule weight acids. Oxalic acid is the dominated
660 dicarboxylic acids measured in the ambient environment and biomass burning
661 aerosol (Falkovich et al., 2005; Kundu et al., 2010), and oxalic acid EF was
662 measured to be 2.2 ~ 4.8 and 1.6 ~ 3.6 mg kg⁻¹ for smoke PM_{2.5} and PM_{1.0} in present
663 work. The sums of WSOA EFs ranged from 46.7 to 770.0 mg kg⁻¹, and they were
664 highly correlated with emissions of OC and PM from Table S9 (SI). Study has
665 confirmed organic acids contribute a significant fraction of both oxygenated volatile
666 organic compounds (OVOCs) in gaseous phase and SOA in particulate phase, the
667 direct emission of particulate organic acids from biomass burning also represents a
668 significant source of precursors for SOA formation, as the low molecular organic
669 acids will evaporate into gas phase or involve in the heterogeneous reaction directly
670 (Takegawa et al., 2007; Veres et al., 2010; Yokelson et al., 2007; Carlton et al., 2006).
671 Moreover, as the significant fraction of water soluble organic carbon, organic acids
672 play major response to CCN activity of smoke particles, and organic acids coating or
673 mixing can amplify hygroscopic growth of inorganic salts by decreasing the
674 deliquescence RH, enable the particle to be CCN at relative low degree of
675 supersaturation (Falkovich et al., 2005; Ghorai et al., 2014). In the ambient
676 environment, organic acids can enhance atmospheric new particle formation by
677 impairing nucleation barrier (Zhang et al., 2004), besides, particulate organic acids
678 can also mobilize the solubility of mineral species, like iron, altering the chemical
679 process of particles (Cwiertny et al., 2008). And prominent optical properties of
680 organic acids like humic/fulvic substance make them as potential contributors to the
681 global warming (Yang et al., 2009; Andreae and Gelencsér, 2006).

682 3.1.3 Water soluble aminiums

683 Interest has been focused on the vital role of amines in particle nucleation-growth
684 process and acidity regulating due to their strong base (Tao et al., 2016; Bzdek et al.,
685 2010, 2011). Though ultratrace gaseous amines and particulate aminiums were on
686 the order of pptv or ng m^{-3} , aminium salts exhibit potential climatic and health effect
687 due to their significant different properties in hygroscopicity, optics, and also
688 toxicology (Qiu and Zhang, 2012; Qiu et al., 2011; Samy and Hays, 2013; Zheng et
689 al., 2015; Ho et al., 2015; Tao et al., 2016). It ever proposed that biomass burning is
690 an important source for gaseous amines, especially from smoldering burning, and
691 alkyl amides can be served as biomarkers in particular (Ge et al., 2011; Ho et al.,
692 2015; Lee and Wexler, 2013; Lobert et al., 1990; Simoneit et al., 2003). However,
693 seldom study ever quantitatively explored the particulate water soluble amine salts
694 (WSA) in primary smoke emissions (Schade and Crutzen, 1995; Ge et al., 2011).
695 From this study, WSA contributed about 4.81 wt.% of smoke $\text{PM}_{2.5}$ and 4.69 wt.%
696 of $\text{PM}_{1.0}$, implicating aminium favored to be abundant in fine-mode of smoke
697 particles, especially in $\text{PM}_{2.5-1.0}$. DEAH^+ , TMAH^+ , TEOH^+ and DMAH^+ made up
698 over 80 wt. % of the measured WSA. Fuel-dependence of WSA distribution and
699 emissions were evident. EFs of WSA ranged from 4.5 to 104.8 mg kg^{-1} in smoke
700 $\text{PM}_{2.5}$, the least was from burning of soybean straw and the largest from cotton and
701 rice straws. We used mass ratio of WSA to NH_4^+ to denote the enrichment of
702 aminium in particulate phase. Statistical analysis showed WSA/NH_4^+ was $0.16 \pm$
703 0.03 and 0.18 ± 0.06 in smoke $\text{PM}_{1.0}$ and $\text{PM}_{2.5}$, respectively, which were almost one
704 order of magnitude larger than that in the ambient aerosol (Liu and Bei, 2016; Tao et
705 al., 2016). Tao et al. (2016) ever measured the ratio as a function of particle size
706 during NPF days in Shanghai, and a noticeable enrichment of aminiums for ultrafine
707 particles ($<56 \text{ nm}$) was observed with WSA/NH_4^+ over 0.2, highlighting the
708 competitive role for amines to ammonia in particle nucleation and initial growth of
709 the nuclei, the ratio was then decreased with the increasing particle size, and the final
710 increasing trend was found after $\sim 1.0 \mu\text{m}$, and average WSA/NH_4^+ for ambient bulk

711 PM_{1.0} and PM_{2.5} were 3.2% and 3.5% , respectively.

712 **3.1.4 PAHs and Phenols**

713 Atmospheric PAHs are primarily the byproduct of incomplete combustion of
714 biomass and fossil fuels (Simcik et al., 1999; Galarneau, 2008). Due to their high
715 degree of bioaccumulation and carcinogenic or mutagenic effect, the sources and
716 environmental fate of the ubiquitous PAHs have been the subjects of extensive
717 studies (Santodonato, 1997; Kim et al., 2013). PAHs can involve in photochemical
718 reaction to form SOA, the process is influenced by gas-to-particle partition and
719 meteorological conditions. Moreover, oxidation may increase the toxicity of PAHs
720 (Arey and Atkinson, 2003; Wang et al., 2011). Biomass burning is one of the main
721 sources of gaseous and particulate PAHs, which even contributes to about half of
722 anthropogenic PAHs emissions in China (Xu et al., 2006; Zhang et al., 2011).
723 Burning conditions can significantly influence the emission of PAHs, under the
724 flaming phase in this study, PAHs contributed 0.46 wt.% of smoke PM_{2.5} and 0.28
725 wt.% of PM_{1.0}, over 60% of the total PAHs were associated to respiratory submicron
726 particles. The sum of EFs of 16 PAHs in smoke PM_{2.5} ranged from 1.81 to 8.30 mg
727 kg⁻¹, which were consistent with the values from literature (Dhammapala et al.,
728 2007a, b; Lee et al., 2005; Zhang et al., 2011). Hays et al. (2005) estimated total EFs
729 of 16 PAHs to be 3.3 mg Kg⁻¹ in wheat straw burning PM_{2.5}. Korenaga et al. (2001)
730 measured PAHs EFs from rice straw burning to be 1.9 mg Kg⁻¹ in particulate phase,
731 while the value from Jenkins et al. (1996) was 16 mg Kg⁻¹. Dhammapala et al.
732 (2007b) found negative linear response for biomass burning source PAHs emissions
733 to burning efficiency, and under flaming combustion, particulate total 16 PAHs EFs
734 were 2 ~ 4 mg Kg⁻¹. Zhang et al. (2011) simulated burning of rice, corn, and wheat
735 straws, the corresponded PAHs EFs were measured as 1.6, 0.9, and 0.7 mg Kg⁻¹ in
736 fine smoke particles, respectively. Great uncertainties for PAHs EFs were evident
737 that relied on burning phase, fuel types, moisture content, and also measurement
738 techniques. Dhammapala et al. (2007a) also found laboratory simulation might

739 overestimate the emission factors of PAHs compared with field burnings. EFs for
 740 individual PAHs were included in Table S7 and S8 (SI). The distribution of
 741 particulate PAHs emission factors was presented in Fig. 3a. Of the particle bound
 742 PAHs, 3~4-rings components were the primary ones, including Pyr, Ant, Ace, Flu,
 743 Phe, and Chr. Concentration ratios of selected PAHs, namely diagnostic ratios, were
 744 usually used to trace the source and make apportionment of specific pollutions
 745 (Yunker et al., 2002; Simcik et al., 1999). In this work, average Ant/(Ant+Phe),
 746 Flu/(Flu+Pyr), BaA/(BaA+Chr), and IP/(IP+BghiP) ratios of 5 types agricultural
 747 residue burning smokes were 0.72, 0.36, 0.47, and 0.58, respectively. There was no
 748 significant difference ($P < 0.05$ at 95% CI) of the ratios in $PM_{1.0}$ and $PM_{2.5}$. According
 749 to previous work, Ant/(Ant+Phe) above 0.1 and BaA/(BaA+Chr) above 0.35 indicate
 750 the dominance of combustion and pyrolytic sources, Flu/(Flu+Pyr) and IP/(IP+BghiP)
 751 ratios greater than 0.50 suggest coal or biomass burnings dominate (Simcik et al.,
 752 1999; Yunker et al., 2002). However, validation of source apportionment using
 753 specific diagnostic ratios should have its constraints, because of variations in source
 754 strengths and atmospheric processing of PAHs (Arey and Atkinson, 2003; Galarneau,
 755 2008).

756 From Table S9 (SI), PAHs in smoke particles were highly correlated with EC and
 757 OC contents. PAHs primarily originate from pyrolysis of organic materials during
 758 combustion, and formation mechanisms of PAHs and soot are closely intertwined in
 759 flames. High-molecular-weight PAHs (>500 atomic mass unit) act as precursors of
 760 soot particles (Lima et al., 2005; Richter et al., 2000). Thus, PAHs with 3, 4, and 5
 761 rings accumulate and dominate in the emissions of biomass burning, as larger
 762 molecular weight PAHs tend to incorporate into soot particles. PAHs
 763 expulsion-accumulation in OC and EC fractions were analyzed by linear fitting of
 764 PAHs mass fractions and EC mass fractions in carbonaceous materials (EC+OC) in
 765 Fig. 3b. The partitions can be parameterized as Eq. (10):

$$766 \quad f_{\text{PAHs}} = \frac{m_{\text{PAHs}}}{m_{\text{OC}} + m_{\text{EC}}} = \beta_{\text{EC}} \times \frac{m_{\text{EC}}}{m_{\text{OC}} + m_{\text{EC}}} + \beta_{\text{OC}} \times \frac{m_{\text{OC}}}{m_{\text{OC}} + m_{\text{EC}}} = \beta_{\text{EC}} \times f_{\text{EC}} + \beta_{\text{OC}} \times f_{\text{OC}} \quad (10)$$

767 where f_{EC} and f_{OC} are the mass fraction of OC and EC in carbonaceous materials

768 (EC+OC). β_{EC} and β_{OC} are expulsion-accumulation coefficients of PAHs in OC and
769 BC. The coefficient of β_{EC} is 1.1×10^{-3} in smoke $PM_{1.0}$ and 1.9×10^{-3} in $PM_{2.5}$, the
770 corresponded β_{OC} is 0.3×10^{-3} and 0.5×10^{-3} .

771 Phenols are the most common SOA precursor/product and organic pollutants in
772 the atmosphere (Berndt and Böge, 2006; Schauer et al., 2001). Hydroxyl functional
773 group and aromatic benzene ring make phenols a paradigm in heterogeneous
774 reaction upon photo oxidation research and aqueous phase reaction research. Phenols
775 are also ROS (reactive oxidized species) precursors that present health hazard (Bruce
776 et al., 1987). Phenol and substituted phenols are thermal products of lignin pyrolysis
777 during biomass burning (Dhammapala et al., 2007a), and the most abundant
778 methoxyphenols can be markers of biomass burning sources (Urban et al., 2016).
779 The five measured phenols contributed 3.0 wt.% and 2.5 wt.% of $PM_{2.5}$ and $PM_{1.0}$.
780 2, 6-dimethoxyphenol was the major one of the measured phenols. Mass fraction of
781 phenols was about 7~9 times of PAHs in smoke aerosols. EFs for the sum phenols
782 were 9.7 ~ 41.5 and 7.7 and 23.5 $mg\ Kg^{-1}$ for smoke $PM_{2.5}$ and $PM_{1.0}$, respectively.
783 Dhammapala et al. (2007a) estimated particulate methoxyphenols emissions to be 35
784 $\pm 24\ mg\ Kg^{-1}$ for wheat straw burning, while Hays et al. (2005) measured the same
785 compounds to be 6.8 $mg\ Kg^{-1}$. Carbonaceous materials like PAHs and Phenols or
786 aromatic and phenolic deviates are the main chromophores in the atmosphere, and
787 the considerable fractions of PAHs and Phenols justify biomass burning as a
788 significant source of brown carbon (Laskin et al., 2015), study has proved ~ 50% of
789 the light absorption in the solvent-extractable fraction of smoke aerosol can be
790 attributed to these strong BrC chromophores (Lin et al., 2016).

791 **3.1.5 Inorganic components**

792 From Fig. 2, smoke particles consisted of approximately 24 wt.% water soluble
793 inorganics (WSI), and the inorganic salts resided more in submicron particles. Great
794 amount of inorganics enable smoke particles to be efficient CCN, and the distinct
795 optical scattering characters of the inorganic fractions may neutralize the warming

796 effect of brown carbon for smoke aerosol, otherwise, inorganics coating or mixing
797 will enhance light absorbing of BC. K^+ , NH_4^+ , Cl^- , and SO_4^{2-} were the principle
798 inorganic ions. Particulate enriched K^+ together with levoglucosan are treated as tracer
799 of pyrogenic source (Andreae et al., 1998). And specific mass ratio of K^+/OC or
800 K^+/EC will help make source apportionment of particulate pollutants with PMF
801 (Positive Matrix Factorization) and PFA (Principle Balance Analysis) models (Lee et
802 al., 2015). K^+/OC in smoke particles ranged from 0.11 to 0.25 with average value of
803 0.17 in $PM_{1.0}$ and 0.14 in $PM_{2.5}$, which were similar to those reported for the
804 Savannah burning and agricultural waste burning emissions in India and China
805 (Echalar et al., 1995; Ram and Sarin, 2011; Li et al., 2015). However, OC represents
806 large uncertainty arise from degree of oxidization and burning condition, K^+/EC is
807 more practical parameter to distinguish the pyrogenic pollutants in ambient study. To
808 smoke particle emitted from flaming fires, K^+/EC was 0.58 ± 0.24 in $PM_{1.0}$ and 0.53
809 ± 0.18 in $PM_{2.5}$. Cl^- was the main anion to balance the charge of WSI in smoke
810 particles. Mean charge ratio of $Cl^- : K^+$ was 1.46 and 1.49 in $PM_{1.0}$ and $PM_{2.5}$,
811 implicating surplus chloride will associate with other cations. With atmospheric
812 aging, the Cl/K ratio will decrease as chloride being replaced by secondary sulfate
813 and nitrate (Li et al., 2015; Li et al., 2003). Equivalent charge ratio of primary
814 cations ($NH_4^+ + K^+$) to primary anions ($SO_4^{2-} + Cl^-$) was 1.05 in $PM_{1.0}$ and 1.01 in
815 $PM_{2.5}$ on average, and charge ratios of total cations to anions ($R_{C/A}$) was 1.09 and
816 1.07 in $PM_{1.0}$ and $PM_{2.5}$. $R_{C/A}$ was used to indicate the neutralizing level of
817 particulate matters in many studies. $R_{C/A} \geq 1$ indicates most of the acids can be
818 neutralized, while $R_{C/A} < 1$ means atmospheric ammonia is deficient and the aerosol is
819 acidic (Adams et al., 1999; He et al., 2011a; Kong et al., 2014). In ambient
820 environment, acidic aerosol was prevailing urban pollutants in many cities from field
821 investigation (He et al., 2011a; Kong et al., 2014). Acidic aerosols can increase the
822 risks to human health and affect the atmospheric chemistry by activating hazardous
823 materials and promoting the solubility of particulate iron and phosphorus (Amdur
824 and Chen, 1989; Meskhidze, 2005). The emission and transport of biomass burning
825 particles may neutralize the acidity of ambient particles. However, only limited WSI

826 were brought into in the analytical system, it is not really to tell the acidity or base of
827 smoke particles, considering the existence of massive organic acids and ammoniums.

828 Trace mineral elements attracted great attention for the role as catalyst in
829 atmospheric heterogeneous reaction and health cares (Davidson et al., 2005;
830 Dentener et al., 1996). Wet/dry deposition of particles during long range transport
831 will affect the ecological balance by releasing mineral elements (Jickells et al., 2005).
832 Dust storm, weathering, and industrial process are the main sources of particulate
833 metals, and incineration can also produce a lot of mineral elements (Moreno et al.,
834 2013). However, the emissions of trace metals from biomass burning are highly
835 uncertain (Li et al., 2007; Zhang et al., 2012), the great influence from local soil
836 environment and soil heavy metal pollution will certainly affect the metal content in
837 biomass fuel and smoke particle. In this study, THM resided more in PM_{2.5} than in
838 PM_{1.0}. Smoke PM_{2.5} consisted of 6.7 wt.% THM on average, PM_{1.0} comprised 4.1
839 wt.% THM. Average EFs of THM in PM_{2.5} and PM_{1.0} were 0.056 g kg⁻¹ and 0.028 g
840 kg⁻¹ in this work, of which Al contributed over 90 wt.%, in line with result from
841 domestic burning of wood and field investigation of crop straw burning (Li et al.,
842 2007; Zhang et al., 2012). Smoke particles from wheat, rice, and corn straws
843 contained more mineral elements than that from cotton and soybean residues
844 combustion. Regardless the difference in biomass fuels, the result may imply that
845 soil heavy metal pollution is heavier in the East China than that in Xinjiang in the
846 West North of China (Wei and Yang, 2010).

847 **3.2 Size, morphology, and mixing state of smoke particles**

848 Fresh smoke particles exhibited unimodal size distribution within 500 nm (Fig. S3,
849 SI), and previous chamber study has also confirmed that agricultural fire produces
850 large amount of ultrafine particles, implying the great potential role to act as CCN
851 and more profound threat to human health (Araujo et al., 2008; Delfino et al., 2005;
852 Zhang et al., 2011). However, the role of particles in the atmospheric process and
853 health hazard depends not only size, but also morphology and chemical mixing

854 states (Dusek et al., 2006; Kennedy, 2007; Mikhailov et al., 2006; Schlesinger, 1985).
855 From TEM images in Fig. 4, agricultural straw burning aerosols comprised a broad
856 class of morphological and chemically heterogeneous particles. Non-uniformly
857 internal mixing of the agglomerates was noticeable, including the major
858 carbonaceous particles and a considerable amount of inorganic salt particles, which
859 was consistent with previous particulate chemical analysis. KCl particles containing
860 minor sulfate or nitrate were the primary inorganic particles, which presented crystal
861 or amorphous state from X-ray diffraction analysis (Fig. 4 a, b, c). And
862 potassium-bearing particles have been used as a tracer of ambient biomass burning
863 pollutants. Fly ash particles were arresting due to visible morphology difference and
864 mineral chemical composition (Fig. 4 d, e, f). Fly ash particles were more compact
865 and rich in mineral elements like Ca, Si, Fe, Al, Mn, and Cr. Besides, these particles
866 had larger size, statistical average diameter of fly ash particles obtained from bulk
867 analysis was $2.2 \pm 1.6 \mu\text{m}$. The result also proved heavy metals resided more in
868 $\text{PM}_{2.5}$ than $\text{PM}_{1.0}$. Fly ashes are by products of incineration process (Buha et al.,
869 2014), including coagulation of fuel issue debris, condensation of evaporated
870 mineral metal from biomass fuels or adhered dirt at different burning phase. These
871 fly ashes coated by or agglomerated with carbonaceous materials were like mash of
872 mineral without clear lattice. Tar ball as a specific form of brown carbon and soot
873 were representative particles of biomass burning aerosol (Wilson et al., 2013;
874 Chakrabarty et al., 2010; Tóth et al., 2013). From Fig. 4 g, chain-like soot particles
875 were coagulated with tar ball. Soot particles were agglomerates of small roughly
876 spherical elementary carbonaceous particles, these chemical consistent particles were
877 within 20~30 nm, and high-resolution TEM showed the soot spheres consisted of
878 concentrically wrapped graphitic layers, while monomeric tar balls possessed
879 disordered microstructure. Tar balls and soot corresponded to different stages in the
880 aging of organic particles; tar balls abundant in fresh or slightly aged biomass smoke
881 are formed by gas-to-particle conversion of high-molecular weight organic species
882 or from aged primary tar droplets upon biomass burning. Soot represents further
883 aged carbon-bearing particles, formed from the pyrolysis of lignin, cellulose, or tar

884 balls (Pósfai, 2004; Tóth et al., 2013). The botryoid aggregates in Fig. 4 g can be
885 viewed as transformation of tar ball to soot. Tar ball and soot were also internal
886 mixed with inorganic salt including sulfate and nitrate (Fig. 4 g, h, i), which made
887 the physiochemical properties of BC even complicated, as study has confirmed
888 inorganic sulfate mixing will enhance light absorption and hygroscopicity of BC
889 (Zhang et al., 2008b). Dark-ring like shell of tar ball (Fig. 4 g, h) and spot-like
890 particles adhered to the surface of tar ball (Fig. 4 i) were K-rich materials. And size
891 of soot particles was mainly within 200 nm, while tar ball and other carbonaceous
892 particles can be over one micrometer.

893 **3.3 Open burning emissions**

894 **3.3.1 Crop straw production**

895 The agricultural straw productions were calculated and geographically displayed in
896 Fig. 5 a-c. Totally 647.3 Tg agricultural straws were produced in 2012 and dispersed
897 mainly in the North and Northeast of China. The distributions of the straws clearly
898 correspond to the distinct planting regions that are divided by Qinling
899 Mountain-Huaihe River line and the Yangtze River. Rice is primarily planted in the
900 south of Qinling Mountain-Huaihe River line, only 10 % rice (single cropping rice
901 dominate) is planted in Heilongjiang, Jilin, and Liaoning province, while wheat and
902 corn are grew mostly in the north of the Yangtze River. Over 90 % of the wheat
903 planted in China is winter wheat that gets ripe in summer, and more than 80 % rice
904 including middle and late rice grows mature in autumn. Summer harvest contributed
905 about 25 % of the agricultural straw production, which solely consists of rice and
906 wheat straws in this period and distributes uniformly in the central and east of China.
907 493.9 Tg crop straws were produced mainly from corn and rice harvesting in autumn.
908 Soybean and cotton straws account for about 8.6 % of autumn straw production that
909 were primarily produced in Heilongjiang and Xinjiang province.

910 3.3.2 Open burning rate

911 The five scenarios of field burning rates and regional AIP ($\frac{I_{k,y}}{AI_{k,y}}$) in the year of
912 2000, 2006, and 2012 were listed in Table 4 and statistically analyzed in Fig. 6. A
913 significant difference ($P < 0.05$ at 95% CI) of regional burning rates among the
914 versions was observed, and the rates from NDRC report were generally higher. For
915 convenience, six zones were classified by geographic divisions and economic areas
916 in China, including the North Plain of China (NPC: Anhui, Shandong, Hebei, Shanxi,
917 Tianjin, Beijing), the Central of China (CC: Hunan, Henan, Hubei), the Yangtze
918 River Delta (YRD: Zhejiang, Jiangsu, Shanghai), the Northeast of China (NC:
919 Heilongjiang, Liaoning, Jilin), the Pan-Pearl River Delta (PRD: Hainan, Guangdong,
920 Fujian, Guangxi, Guizhou, Sichuan, Yunnan, Jiangxi), the West of China (WC:
921 Shanxi, Chongqing, Xinjiang, Qinghai, Ningxia, Tibet, Inner Mongolia, Gansu).
922 And the bulk-weighted burning rates that averaged from BAU, EM, and NDRC
923 versions for the six zones were $22.3 \% \pm 3.1 \%$, $21.1 \% \pm 3.3 \%$, $28.4 \% \pm 6.2 \%$,
924 $23.3 \% \pm 9.2 \%$, $21.4 \% \pm 6.5 \%$, and $14.2 \% \pm 8.0 \%$, respectively. It was obvious
925 that agricultural field burning was most serious in the Yangtze River Delta,
926 especially in the Zhejiang province. The nationwide field burning rate was 21.4 %,
927 16.3 %, 26.0 %, 14.9 %, and 26.8 % for BAU-I, BAU-II, EM-I, EM-II, and NDRC,
928 respectively, which were comparable with the document values (Daize, 2000; Wei et
929 al., 2004; Zhang et al., 2008a).

930 3.3.3 Agricultural open burning emissions

931 $PM_{2.5}$ emissions from agricultural field burnings based on BAU, EM, and NDRC
932 versions were calculated and geographically presented in Fig. 7 (emissions of
933 detailed individual species in SI). A similar spatial character of regional emission
934 distribution was observed for BAU, EM, and NDRC versions, most emissions were
935 allocated in the North Plain and the Central of China, where the primary agricultural
936 regions locate, echoing the agricultural fire sites in Fig. S2 (SI). Although filed

937 burning rates were higher in the Yangtze River Delta, the crop residue productions in
938 this zone were much less, which only contributed 4.3 % of the national straw
939 productions. Take NDRC as the basis, BAU and EM scenarios all underestimated the
940 emissions in the Northeast of China, especially in Heilongjiang.

941 The temporal distributions of field burning emissions also echoed the crop residue
942 productions and the agricultural fire sites in summer and autumn harvest. Apart from
943 Henan and Tibet where the main crop straws were produced in summertime, more
944 pollutants were emitted in autumn harvest period to the rest place, which has been
945 confirmed by many studies (He et al., 2011; Wang and Zhang, 2008). And the large
946 scale field burning emissions in autumn exhibited great influence on the haze
947 formation and visibility degradation in the North and East of China (Leng et al.,
948 2014; Shi et al., 2014), Huang et al. (2012a) has identified biomass burning together
949 with secondary inorganic aerosol (SIA) and dust pollution as three typical haze types
950 in Shanghai. In summertime, field burning emissions concentrated in the North Plain,
951 the Central, and the South regions. While in autumn, the emissions became more
952 ubiquitous and serious in the Northeast of China.

953 Nationwide emission inventories and flux concentrations were graphically
954 displayed in Fig. 8 and tabular presented in Table 5. The total PM_{2.5} emission from
955 agricultural field burnings was 738.36-1241.69 Gg in 2012, and rice, corn, and
956 wheat straw burnings made up 93.5 % ~ 95.6 % of the total emissions. The largest
957 quantities of PM_{2.5} emissions were emitted from Heilongjiang, Shandong, Henan,
958 Jilin, Jiangsu, Anhui and Hebei, distinct difference in the emissions from various
959 scenarios were observed, especially for Heilongjiang province which contributed
960 5.5 % (55.4 Gg) of PM_{2.5} emissions under BAU-II scenarios, while the figure was
961 22.9 % (231.0 Gg) under EM-I scenarios. Annual emissions of PM_{1.0}, OC, and EC
962 was 661.81-1111.90, 318.84-533.19, and 98.06-164.97 Gg, respectively, which were
963 comparable with the precious studies (Cao et al., 2006, 2011; Wang et al., 2012). Qin
964 and Xie (2011, 2012) developed national carbonaceous aerosol emission inventories
965 from biomass open burning for multi-years with dynamic burning activity, they
966 believed BC and OC emissions followed an exponential growth from 14.03 and

967 57.37 Gg in 1990 to 116.58 and 476.77 Gg in 2009. Cao et al. (2006, 2011)
968 calculated smoke aerosol emissions from biomass burning in China for 2000 and
969 2007 using the same activity data from BAU-I scenarios, national OC and EC
970 emissions were reported to be 425.9 and 103.0 Gg in 2000, however, no evident
971 changes were found for the emissions in 2007, which were assessed to be 433.0 and
972 104.0 Gg. Huang et al. (2012b) estimated crop burning in the fields with unified EFs
973 and burning rate (~6.6 %) for all kinds of crops across China in 2006, the estimated
974 annual agricultural fire emissions were about 270, 100, and 30 Gg for PM_{2.5}, OC,
975 and BC, respectively. In present work, agricultural fire PM_{2.5} emissions in 2012 were
976 allocated into six zones, average contribution in percentage for each zone was
977 compared: NPC (23.1 %) ≥ NC (21.6 %) > PRD (18.4 %) ≥ CC (18.2 %) > WC
978 (9.8 %) > YRD (8.8 %). Furtherly, contribution for summertime emissions was: NPC
979 (35.5 %) > CC (28.8 %) ≥ PRD (21.1 %) > YRD (9.1 %) > WC (5.4 %) > NC
980 (0.1 %), and for autumn harvest emissions: NC (27.8 %) > NPC (19.6 %) > PRD
981 (17.6 %) > CC (15.1 %) > WC (11.1 %) > YRD (8.8 %). It was obviously that the
982 North Plain experienced extensive crop fire emissions during the whole harvest
983 periods, where PM_{2.5}, PM_{1.0}, OC, and BC emissions in 2012 were 233.6, 209.8,
984 102.3, and 29.4 Gg on average. Liu et al. (2015) developed emission inventories
985 from agricultural fires in the North Plain based on MODIS fire radiative power,
986 emission for PM_{2.5}, OC, and BC in 2012 was reported to be 102.3, 37.4, and 13.0 Gg,
987 respectively. However, EFs were also treated as unified values (e.g., Crop burning
988 EFs for PM_{2.5}, OC, and BC was 6.3, 2.3, and 0.8 g Kg⁻¹) in the work of Liu et al.
989 (2015) that was cited directly from Akagi et al. (2011) without considering fuel type
990 dependence of EFs. Zhao et al. (2012) established comprehensive anthropogenic
991 emission inventories for Huabei Region including the North Plain, Inner Mongolia,
992 and Liaoning province, all crop straws were assumed to be burnt in the field,
993 resulting in much more emissions of 446 Gg OC and 160 Gg BC in 2003. A specific
994 temporal pattern for agricultural fire emissions was observed in the Northeast of
995 China (Heilongjiang, Liaoning, and Jilin), where the open burning were mainly
996 occurred in autumn harvest to produce great amount of pollutants (217.5 Gg PM_{2.5},

997 89.4 Gg OC, and 29.7 Gg EC), while emissions in the summertime can be neglected.

998 In 2012, 20-25 % of national emissions were released from summertime field
999 burnings, that was 226.0 Gg PM_{2.5}, 205.2 Gg PM_{1.0}, 105.9 Gg OC, 28.4 Gg EC, 24.6
1000 Gg char-EC, 3.8 Gg soot-EC, 6.8 Gg WSOA, 1.0 Gg WSA, 0.1 Gg PAHs, 0.9 Gg
1001 phenols, and 2.1 Gg THM on average. The corresponded values for autumn harvest
1002 were 781.6, 697.9, 327.3, 106.0, 88.8, 17.2, 18.4, 4.8, 0.4, 1.9, and 6.6 Gg,
1003 respectively. Integrated smoke OC/EC was 3.7 from national summertime emission
1004 and 3.1 from autumn harvest emission, the figure for smoke char-EC/soot-EC was
1005 6.5 and 5.2, regarding to different locations, integrated OC/EC in the North Plain
1006 was 4.1 in summertime emission and 3.2 in autumn harvest, while OC/EC in the
1007 Central of China was 3.1 for both summer and autumn harvest emissions, implying
1008 temporal-spatial characters of agricultural field fires exhibit potential influence on
1009 composition of smoke emissions and its related physiochemical properties. Zhang et
1010 al. (2011) estimated particulate PAHs emissions form three types of crop residues to
1011 be 0.46 Gg in 2003. Xu et al. (2006) counted PAHs from all straws with the
1012 assumption that burning rates to be unit, and they calculated 5-10 Gg PAHs
1013 emissions in 2003, which was ten times of our result.

1014 The nationwide flux concentration of smoke PM_{2.5} was 0.7-1.0 $\mu\text{g m}^{-3} \text{d}^{-1}$ in
1015 summer harvest and 1.4-3.5 $\mu\text{g m}^{-3} \text{d}^{-1}$ in autumn harvest, while average annual flux
1016 concentrations for OC and EC were 0.80 and 0.25 $\mu\text{g m}^{-3} \text{d}^{-1}$. Saikawa et al. (2009)
1017 assessed the annual concentrations of OC and BC from biomass burning primary
1018 emission in China using global models of chemical transport (MOZART-2) to be 1.8
1019 and 0.35 $\mu\text{g m}^{-3}$. The most polluted areas were Anhui, Henan, Shandong, Jiangsu,
1020 Liaoning, and Hunan.

1021 **3.3.4 Uncertainties of the emissions**

1022 The fuzziness and uncertainties of major pollutants emissions from fuel combustion
1023 in China came from the uncertainties in specific-source emission factors and
1024 effective consumption of bio- or fossil fuel. Frey et al. analyzed uncertainties in

1025 emission factors and emissions of air toxic pollutants and technology dependent
1026 coal-fire power plants via bootstrap simulation method (Frey and Zhao, 2004; Frey
1027 and Zheng, 2002). Zhao et al. estimated uncertainties in national anthropogenic
1028 pollutants emissions based on Monte Carlo simulation, and they believed activity
1029 rates (e.g. fuel consumption) are not the main source of emissions uncertainties at the
1030 national level (Zhao et al., 2011; Zhao et al., 2012). The uncertainties in emission
1031 inventory can also be estimated by comparing different emission inventories for the
1032 same region and period (Ma and Van Aardenne, 2004).

1033 In this study, we investigated the uncertainties of multi-pollutants emissions for
1034 agricultural residue open burning using Monte Carlo Simulation. Detailed
1035 methodology was referred to Qin and Xie (2011). We followed the assumption: a
1036 normal distribution with coefficient of variation (CV) of 30% for the official
1037 statistics (e.g., crop production and GDP economic data obtained from Statistic
1038 Yearbooks, field burning rates for agricultural straw derived from NDRC report,
1039 etc.), a normal distribution with 50% CV for open burning rates from literature
1040 (BAU-I and BAU-II), and a uniform distribution with $\pm 30\%$ deviation for the rest
1041 activity data (crop-to-residue ratio, dry matter fraction, and burning efficiency).
1042 Regarding the emission factors, Bond et al. (2004) assumed that most particulate
1043 EFs followed lognormal distributions with CV of $\pm 50\%$ for domestic EFs, and of \pm
1044 150% for EFs obtained from foreign studies. Here, we applied the CV of smoke
1045 EFs as we measured ones, which were chemical species and fuel type dependent.
1046 With randomly selected values within the respective probability density functions
1047 (PDFs) of EFs and activity data for each biomass type, Monte Carlo simulation was
1048 implemented for 10,000 times, and the uncertainties in national yearly
1049 multi-pollutants emissions at 95% CI were obtained for all the 5 versions.
1050 Afterwards, uncertainties for the average emission inventories were assessed using
1051 the propagation of uncertainty calculation that suggested by IPCC (1997) (method
1052 in SI), and all the emission uncertainties were presented in percentage in Table 6.
1053 Emissions for water soluble aminiums and organic acids had the vast uncertainties,
1054 due to their large deviation in EFs compared with other smoke species. Besides,

1055 emissions of BAU versions were more accurate than EM versions, because of more
1056 uncertainty addition in the burning rates conversion using economic data for EM
1057 versions. Otherwise, burning rates derived from NDRC report were assumed to
1058 have less uncertainty, resulting in the least uncertainties in smoke emission
1059 assessments. On average of all the 5 versions, mean, 2.5th percentile, and 97.5th
1060 percentile values for smoke PM_{2.5} emissions in 2012 were 1005.7, 758.3, and
1061 1344.6 Gg, respectively. As to OC emissions, mean, 2.5th percentile, and 97.5th
1062 percentile values were 432.4, 327.8, and 576.4 Gg, the figure for EC was 134.2,
1063 100.9, and 187.9 Gg. Therefore, the overall propagation of uncertainties for smoke
1064 PM_{2.5}, OC, and EC at 95% CI was (-24.6%, 33.7%), (-24.4%, 33.5%), and (-24.2%,
1065 33.3%), respectively. The uncertainties for OC and EC emissions were much less
1066 than the work of Qin and Xie (2011), in which emission and uncertainties were
1067 266.7 Gg (-55.9%, 96.1%) for OC and 66.9 Gg (-53.9%, 92.6%) for EC in 2005.

1068 **3.4 Health and health-related economic impacts**

1069 **3.4.1 Carcinogenic risk**

1070 Calculated CR_{SPM} for smoke PM_{2.5} from wheat, corn, rice, cotton, and soybean
1071 straw burning were 5.3×10^{-6} , 3.8×10^{-6} , 2.6×10^{-6} , 0.7×10^{-6} , and 1.3×10^{-6} per $\mu\text{g m}^{-3}$,
1072 respectively. And the corresponded one in million PEL was 0.2, 0.3, 0.4, 1.4, and 0.8
1073 $\mu\text{g m}^{-3}$. Wu et al. (2009) ever assessed unit risk of wood and fuel burning particles
1074 using metals merely, the results were 3.2×10^{-6} and 1.5×10^{-6} per $\mu\text{g m}^{-3}$, which were
1075 close to that in our study. In actual application, PEL of smoke particles should be
1076 bulk mass concentration of mixed aerosols.

1077 It was noticeable that apart from Tibet and Qinghai, the flux concentration of
1078 smoke PM_{2.5} among all the five emission versions in other regions far surpassed the
1079 PEL, especially the North Plain and the Central of China, exhibiting great potential
1080 inhalable cancer risk. For the health care, emission flux concentration should be
1081 constrained within the PEL of crop straw burning aerosol. Thus the critical filed
1082 burning rates can be derived to ensure risk aversion following Eq. (11):

$$R_k \leq \frac{10^{-6} \times S_k \times h \times T_k}{\sum_j \sum_i P_{t,k,i} \times r_i \times H_{t,k,i} \times D_i \times f_i \times EF_{i,j} \times CRF_i} \quad (11)$$

1084 The conservative values of regional field burning rates from Eq. (11) were named
 1085 as Carcinogenic Risk Control scenarios (CRC) and listed in Table S10 (SI), which
 1086 would be instructive in emission control. Under CRC, national crop straw field
 1087 burning rate was less than 3%, emissions of PM_{2.5} were geographically presented in
 1088 Fig. S4 (SI), and 146.3 Gg yr⁻¹ smoke PM_{2.5} should be released at largest in China,
 1089 the corresponded annual flux concentration of PM_{2.5} was within 0.3 μg m⁻³ d⁻¹
 1090 (detailed emission inventories under CRC version see in SI).

1091 3.4.2 Health impacts

1092 Regional health impacts from acute exposure of agricultural residue burning aerosol
 1093 were assessed using average daily flux concentrations of smoke PM_{2.5}, the result was
 1094 tabulated in Table S11 (SI). The impacts from smoke PM_{2.5} exposure were severest
 1095 in Jiangsu, Shandong, and Henan province, where annual premature mortality was
 1096 over one thousand. Overall, China suffered from 7836 (95% CI: 3232, 12362)
 1097 premature death, 31181 (95% CI: 21145, 40881) respiratory hospital admissions,
 1098 29520 (95% CI: 12873, 45602) cardiovascular hospital admissions, and 7267237 (95%
 1099 CI: 2961487, 1130784) chronic bronchitis related to agricultural fire smoke in 2012
 1100 from Table 7. According to national health statistical reports (NHFPC, 2013), the
 1101 hospital admission due to respiratory and cardiovascular disease was 5071523 in
 1102 China in 2012, and smoke PM_{2.5} exposure might contribute ~1.2% of the hospital
 1103 admissions from this study. Saikawa et al. (2009) ever reported 70000 premature
 1104 deaths in China and an additional 30000 deaths globally due to OC, EC, and sulfate
 1105 exposure that were primarily emitted from biofuel combustion in China in 2000,
 1106 however, the results should be overestimated not only in the exaggerated pollutant
 1107 emissions but also in the iterative operations of respective species induced mortality,
 1108 besides, the exposure-response coefficient β and incidence rate he applied from Pope
 1109 et al. (2002) and WHO (2000) were higher than the practical values from local
 1110 research (Cao et al., 2012; Chen et al., 2011; Hou et al., 2012). From Table 7, under

1111 CRC version, over 92 % mortality and morbidity can be avoided.

1112 **3.4.3 Health-related economic losses**

1113 Health-related total economic losses from straw open burning smoke PM_{2.5} exposure
1114 were assessed to be 8822.4 (95% CI: 3574.4, 13034.2) million US\$ on average from
1115 Table 8, accounting for 0.1 % of the total GDP in 2012, and detailed regional
1116 economic losses were listed in Table S12. Economic losses from premature death
1117 contributed about 17% of total losses, and loss from chronic bronchitis dominated.
1118 Hou et al. (2012) ever estimated 106.5 billion US\$ lost due to ambient PM₁₀
1119 exposure in China in 2009; even a severe haze episode (PM_{2.5} be focused on) in
1120 January 2013 may cause 690 premature death and 253.8 million US\$ loss in Beijing,
1121 and source-specification analysis stressed the emission from biomass burning (Yang
1122 et al., 2015; Gao et al., 2015). It was obvious that smoke PM_{2.5} contributed a
1123 noticeable damage to public health and social welfare. According to CRC version
1124 estimation, the carcinogenic risk control policy can save over 92 % of the economic
1125 loss.

1126 **4 Conclusion**

1127 Detailed chemical compositions of smoke aerosol from five major agricultural
1128 straws burning were characterized using an aerosol chamber system. And
1129 corresponded emission factors for particulate OC-EC, char-/soot-EC, WSI, WSOA,
1130 WSA, PAHs, Phenols, and THM in smoke PM_{2.5} and PM_{1.0} were established.

1131 Permissible exposure limits (PEL) of the smoke particles were assessed for
1132 carcinogenic risk concern based on selected hazard pollutants including PAHs and
1133 THM in smoke PM_{2.5}. Daily exposure concentration should be constrained within
1134 0.2, 0.3, 0.4, 1.4, and 0.8 $\mu\text{g m}^{-3}$ for wheat, corn, rice, cotton, and soybean straw,
1135 respectively.

1136 Emission inventories of primary particulate pollutants from agricultural field
1137 burning in 2012 were estimated based on BAU-I, BAU-II, EM-I, EM-II, and NDRC

1138 scenarios, which were further allocated into different regions at summer and autumn
1139 open burning periods. The estimated total emissions were 1005.7 Gg PM_{2.5} (95%CI:
1140 -24.6% , 33.7%), 901.4 Gg PM_{1.0} (95%CI: -24.4%, 33.5%), 432.4 Gg OC (95%CI:
1141 -24.2%, 33.5%), 134.2 Gg EC (95%CI: -24.8%, 34.0%), 113.1 Gg char-EC (95%CI:
1142 -24.8%, 34.1%), 21.0 Gg soot-EC (95%CI: -26.3%, 35.9%), 249.8 Gg WSI (95%CI:
1143 -25.4%, 34.9%), 25.1 Gg WSOA (95%CI: -33.3%, 41.4%), 5.8 Gg WSA (95%CI:
1144 -30.1%, 38.5%), 8.7 Gg THM (95%CI: -26.6%, 35.6%), 0.5 Gg PAHs (95%CI:
1145 -26.0%, 34.9%), and 2.7 Gg Phenols (95%CI: -26.1%, 35.1%), respectively. The
1146 spatial and temporal distributions of the five versions have similar characters that
1147 echo to the agricultural fires sites from satellite remote sensing. Less than 25 % of
1148 the emissions were released from summer field burnings that were mainly
1149 contributed by the North Plain and the Central of China. Flux concentrations of
1150 annual smoke PM_{2.5} that were calculated using box-model method based on five
1151 versions all exceed the PEL. From assessment of health impacts and health-related
1152 economic losses due to smoke PM_{2.5} short-term exposure, China suffered from 7836
1153 (95%CI: 3232, 12362) premature mortality and 7267237 (95% CI: 2961487,
1154 1130784) chronic bronchitis in 2012, which led to 8822.4 (95%CI: 3574.4, 13034.2)
1155 million US\$, or 0.1 % of the total GDP losses.

1156 Percentage of open burned crop straws at post-harvest period should cut down to
1157 less than 3% to ensure risk aversion from carcinogenicity, especially the North Plain
1158 and the Northeast, where the emissions should decrease at least by 94% to meet the
1159 PEL. And by applying such emission control policy, over 92% of the mortality and
1160 morbidity attributed to agricultural fire smoke PM_{2.5} can be avoided in China.

1161 **Supplementary material related to this article is available online at:**

1162 *Acknowledgment.* This work is supported by National Natural Science Foundation of
1163 China (No. 21190053, 21177025), Cyrus Tang Foundation (No. CTF-FD2014001),
1164 Shanghai Science and Technology Commission of Shanghai Municipality (No.
1165 13XD1400700, 12DJ1400100), Priority fields for Ph.D. Programs Foundation of
1166 Ministry of Education of China (No. 20110071130003) and Strategic Priority
1167 Research Program of the Chinese Academy of Sciences (Grant No. XDB05010200).

1168

References:

- 1169 Ackerman, A. S.: Reduction of Tropical Cloudiness by Soot, *Science*, 5468, 1042-1047, 2000.
- 1170 Adams, P. J., Seinfeld, J. H. and Koch, D. M.: Global concentrations of tropospheric sulfate, nitrate,
1171 and ammonium aerosol simulated in a general circulation model, *J. Geophys Res*, D11:13791-13823,
1172 1999.
- 1173 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D. and
1174 Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric
1175 models, *Atmos Chem. Phys.*, 9, 4039-4072, 2011.
- 1176 Amdur, M. O. and Chen, L. C.: Furnace-Generated Acid Aerosols: Speciation and Pulmonary Effects,
1177 *Environ. Health Persp.*, 79, 147-150, 1989.
- 1178 Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing
1179 carbonaceous aerosols, *Atmos Chem. Phys.*, 10, 3131-3148, 2006.
- 1180 Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global*
1181 *Biogeochem Cy.*, 4, 955-966, 2001.
- 1182 Andreae, M. O., Andreae, T. W., Annegarn, H., Beer, J., Cachier, H., le Canut, P., Elbert, W.,
1183 Maenhaut, W., Salma, I., Wienhold, F. G., and Zenke, T. :, Airborne studies of aerosol emissions from
1184 savanna fires in southern Africa: 2. Aerosol chemical composition, *J. Geophys Res.*, D24, 32119-32128,
1185 1998.
- 1186 Araujo, J. A., Barajas, B., Kleinman, M., Wang, X., Bennett, B. J., Gong, K. W. Navab, M., Harkema,
1187 J., Sioutas, C., Lulis, A. J., and Nel, A. E.: Ambient particulate pollutants in the ultrafine range promote
1188 early atherosclerosis and systemic oxidative stress, *Circ Res.*, 5, 589-596, 2008.
- 1189 Arey, J. and Atkinson, R.: Photochemical reactions of PAHs in the atmosphere, PAHs: An
1190 Ecotoxicological Persp., 47- 63, doi: 10.1002/0470867132.ch4, 2003.
- 1191 Arora, P. and Jain, S.: Estimation of Organic and Elemental Carbon Emitted from Wood Burning in
1192 Traditional and Improved Cookstoves Using Controlled Cooking Test, *Environ. Sci. Technol.*, 6,
1193 3958-3965, 2015.
- 1194 Aunan, K. and Pan, X.: Exposure-response functions for health effects of ambient air pollution
1195 applicable for China-a meta-analysis, *Sci. Total Environ.*, 329, 3-16, 2004.
- 1196 Aurell, J., Gullett, B. K. and Tabor, D.: Emissions from southeastern U.S. Grasslands and pine
1197 savannas: Comparison of aerial and ground field measurements with laboratory burns, *Atmos. Environ.*,
1198 111, 170-178, 2015.
- 1199 Bell, R. W. and Hipfner, J. C.: Airborne Hexavalent Chromium in Southwestern Ontario, *J. Air Waste*
1200 *Manage*, 8, 905-910, 1997.
- 1201 Berndt, T. and Böge, O.: Formation of phenol and carbonyls from the atmospheric reaction of OH
1202 radicals with benzene, *Phys. Chem. Chem. Phys.*, 10, 1205-1214, doi:10.1039/B514148F, 2006.
- 1203 Bølling, A. K., Pagels, J., Yttri, K. E., Barregard, L., Sallsten, G., Schwarze, P. E. and Boman, C.:
1204 Health effects of residential wood smoke particles: the importance of combustion conditions and
1205 physicochemical particle properties, *Part. Fibre Toxicol.*, 29, doi:10.1186/1743-8977-6-29, 2009.
- 1206 Bond, T. C.: A technology-based global inventory of black and organic carbon emissions from
1207 combustion, *J. Geophys Res.*, 109, D14203, doi:10.1029/2003JD003697, 2004.
- 1208 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J. Flanner, M. G.,
1209 Ghan, S., Köcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G.,
1210 Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K.,

1211 Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J., PShindell, D., Storelvmo, T.,
1212 Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A
1213 scientific assessment, *J. Geophys Res: Atmos*, 11, 5380-5552, 2013.

1214 Bruce, R. M., Santodonato, J. and Neal, M. W.: Summary Review of the Health Effects Associated
1215 With Phenol, *Toxicol Ind. Health*, 4, 535-568, 1987.

1216 Buha, J., Mueller, N., Nowack, B., Ulrich, A., Losert, S. and Wang, J.: Physical and Chemical
1217 Characterization of Fly Ashes from Swiss Waste Incineration Plants and Determination of the Ash
1218 Fraction in the Nanometer Range, *Environ. Sci. Technol.*, 9, 4765-4773, 2014.

1219 Burkart, K., Nehls, I., Win, T. and Endlicher, W.: The carcinogenic risk and variability of
1220 particulate-bound polycyclic aromatic hydrocarbons with consideration of meteorological conditions,
1221 *Air Quality, Atmos. Health*, 1, 27-38, 2013.

1222 Bzdek, B. R., Ridge, D. P. and Johnston, M. V.: Amine reactivity with charged sulfuric acid clusters,
1223 *Atmos. Chem. Phys.*, 16, 8735-8743, 2011.

1224 Bzdek, B. R., Ridge, D. P. and Johnston, M. V., Amine exchange into ammonium bisulfate and
1225 ammonium nitrate nuclei, *Atmos Chem Phys*, 8:3495-3503, 2010.

1226 Cao, J. J., Wu, F., Chow, J. C., Lee, S. C., Li, Y., Chen, S. W., An, Z. S., Fung, K. K., Watson, J. G.
1227 and Zhu, C. S.: Characterization and source apportionment of atmospheric organic and elemental
1228 carbon during fall and winter of 2003 in Xi'an, China, *Atmos. Chem. Phys.*, 11, 3127-3137, 2005.

1229 Cao, G., Zhang, X. and Zheng, F.: Inventory of black carbon and organic carbon emissions from China,
1230 *Atmos. Environ.*, 34, 6516-6527, 2006.

1231 Cao, G., Zhang, X., Gong, S., An, X. and Wang, Y.: Emission inventories of primary particles and
1232 pollutant gases for China, *Chinese Sci. Bull*, 8, 781-788, 2011.

1233 Cao, G., Zhang, X., Wang, D. and Zheng, F.: Inventory of Emissions of Pollutants from Open Burning
1234 Crop Residues, *J. Agro-Environ. Sci.*, 4, 800-804, 2005.

1235 Cao, J., Xu, H., Xu, Q., Chen, B. and Kan, H.: Fine particulate matter constituents and
1236 cardiopulmonary mortality in a heavily polluted Chinese city, *Environ. Health Persp.*, 3, 373, 2012.

1237 Carlton, A. G., Turpin, B. J., Lim, H., Altieri, K. E. and Seitzinger, S.: Link between isoprene and
1238 secondary organic aerosol (SOA): Pyruvic acid oxidation yields low volatility organic acids in clouds,
1239 *Geophys. Res. Lett.*, L06822, doi:10.1029/2005GL025374, 2006.

1240 Cermak, J. and Knutti, R.: Beijing Olympics as an aerosol field experiment, *Geophys. Res. Lett.*, 36,
1241 L10806, doi:10.1029/2009GL038572, 2009.

1242 Chakrabarty, R. K., Moosmüller, H., Chen, L. W. A., Lewis, K., Arnott, W. P., Mazzoleni, C., Dubey,
1243 M. K., Wold, C. E., Hao, W. M., and Kreidenweis, S. M.: Brown carbon in tar balls from smoldering
1244 biomass combustion, *Atmos. Chem. Phys.*, 13, 6363-6370, 2010.

1245 Chan, M. N., Choi, M. Y., Ng, N. L. and Chan, C. K.: Hygroscopicity of Water-Soluble Organic
1246 Compounds in Atmospheric Aerosols: Amino Acids and Biomass Burning Derived Organic Species,
1247 *Environ. Sci. Technol.*, 6, 1555-1562, 2005.

1248 Chen, H., Hu, D., Wang, L., Mellouki, A. and Chen, J.: Modification in light absorption cross section
1249 of laboratory-generated black carbon-brown carbon particles upon surface reaction and hydration,
1250 *Atmos. Environ.*, 116, 253-261, 2015.

1251 Chen, R., Li, Y., Ma, Y., Pan, G., Zeng, G., Xu, X., Chen, B. and Kan, H.: Coarse particles and
1252 mortality in three Chinese cities: The China Air Pollution and Health Effects Study (CAPES), *Sci.*
1253 *Total Environ.*, 23, 4934-4938, 2011.

1254 Cheng, Y., Ho, K. F., Lee, S. C. and Law, S. W.: Seasonal and diurnal variations of PM_{1.0}, PM_{2.5} and

1255 PM₁₀ in the roadside environment of Hong Kong, *China Particuology*, 06:312-315, 2006.

1256 China, S., Salvadori, N. and Mazzoleni, C.: Effect of Traffic and Driving Characteristics on
1257 Morphology of Atmospheric Soot Particles at Freeway On-Ramps, *Environ. Sci. Technol.*, 6,
1258 3128-3135, 2014.

1259 China Ministry of Health (CMH): China statistical yearbook of public health, Peking Union Medical
1260 College Press, 172–189, 2009. (*In Chinese*)

1261 Christopher, S. A., Chou, J., Zhang, J., Li, X., Berendes, T. and Welch, R. M.: Shortwave direct
1262 radiative forcing of biomass burning aerosols estimated using VIRS and CERES data, *Geophys. Res.*
1263 *Let.*, 15, 2197-2200, 2000.

1264 Chuang, M., Chou, C. C., Sopajaree, K., Lin, N., Wang, J., Sheu, G., Chang, Y. and Lee, C.:
1265 Characterization of aerosol chemical properties from near-source biomass burning in the northern
1266 Indochina during 7-SEAS/Dongsha experiment, *Atmos. Environ.*, 78, 72-81, 2013.

1267 Clarke, A., McNaughton, C., Kapustin, V., Shinozuka, Y., Howell, S., Dibb, J., Zhou, J., Anderson, B.,
1268 Brekhovskikh, V., Turner, H. and Pinkerton, M.: Biomass burning and pollution aerosol over North
1269 America: Organic components and their influence on spectral optical properties and humidification
1270 response, *J. Geophys. Res.*, D12, doi:10.1029/2006JD007777, 2007.

1271 Cwiertny, D. M., Baltrusaitis, J., Hunter, G. J., Laskin, A., Scherer, M. M. and Grassian, V. H.:
1272 Characterization and acid-mobilization study of iron-containing mineral dust source materials, *J.*
1273 *Geophys. Res.: Atmos.*, D5, doi:10.1029/2007JD009332, 2008.

1274 Daize, H.: The Utilizing Status and Prospects of the Crop Straw Resources in China, *Resource*
1275 *Development & Market*:12, 2000.

1276 Davidson, C. I., Phalen, R. F. and Solomon, P. A.: Airborne Particulate Matter and Human Health: A
1277 Review, *Aerosol Sci. Tech.*, 8, 737-749, 2005.

1278 Delfino, R. J., Sioutas, C. and Malik, S.: Potential role of ultrafine particles in associations between
1279 airborne particle mass and cardiovascular health, *Environ. Health Perspect.*, 8, 934-946, 2005.

1280 Dentener, F. J., Carmichael, G. R., Zhang, Y., Lelieveld, J. and Crutzen, P. J.: Role of mineral aerosol
1281 as a reactive surface in the global troposphere, *J. Geophys. Res.*, D17, 22869-22889, doi:
1282 10.1029/96JD01818, 1996.

1283 Dhammapala, R., Claiborn, C., Corkill, J. and Gullett, B.: Particulate emissions from wheat and
1284 Kentucky bluegrass stubble burning in eastern Washington and northern Idaho, *Atmos. Environ.*, 6,
1285 1007-1015, 2006.

1286 Dhammapala, R., Claiborn, C., Jimenez, J., Corkill, J., Gullett, B., Simpson, C. and Paulsen, M.:
1287 Emission factors of PAHs, methoxyphenols, levoglucosan, elemental carbon and organic carbon from
1288 simulated wheat and Kentucky bluegrass stubble burns, *Atmos. Environ.*, 12, 2660-2669, 2007a.

1289 Dhammapala, R., Claiborn, C., Simpson, C. and Jimenez, J.: Emission factors from wheat and
1290 Kentucky bluegrass stubble burning: Comparison of field and simulated burn experiments, *Atmos.*
1291 *Environ.*, 7, 1512-1520, 2007b.

1292 Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F.,
1293 Hings, S., Jung, D.: Size matters more than chemistry for cloud-nucleating ability of aerosol particles,
1294 *Science*, 5778, 1375-1378, 2006.

1295 Echalar, F., Gaudichet, A., Cachier, H. and Artaxo, P.: Aerosol emissions by tropical forest and
1296 savanna biomass burning: Characteristic trace elements and fluxes, *Geophys. Res. Lett.*, 22, 3039-3042,
1297 doi:10.1029/95GL03170, 1995.

1298 Engelhart, G. J., Hennigan, C. J., Miracolo, M. A., Robinson, A. L. and Pandis, S. N.: Cloud

1299 condensation nuclei activity of fresh primary and aged biomass burning aerosol, *Atmos. Chem. Phys.*,
1300 15, 7285-7293, doi:10.5194/acp-12-7285-2012, 2012.

1301 Falkovich, A. H., E., R. G., G., S., Y., R., Maenhaut, W. and Artaxo, P.: Low molecular weight organic
1302 acids in aerosol particles from Rondônia, Brazil, during the biomass-burning, transition and wet
1303 periods, *Atmos. Chem. Phys.*, 5, 781-797, doi:10.5194/acp-5-781-2005, 2005.

1304 Frey, H. C. and Zhao, Y.: Quantification of Variability and Uncertainty for Air Toxic Emission
1305 Inventories with Censored Emission Factor Data, *Environ. Sci. Technol.*, 22, 6094-6100, 2004.

1306 Frey, H. and Zheng, J.: Quantification of variability and uncertainty in air pollutant emission
1307 inventories: method and case study for utility NO_x emissions, *J. Air Waste Manag. Assoc.*, 9,
1308 1083-1095, 2002.

1309 Fu, H., Zhang, M., Li, W., Chen, J., Wang, L., Quan, X. and Wang, W.: Morphology, composition and
1310 mixing state of individual carbonaceous aerosol in urban Shanghai, *Atmos. Chem. Phys.*, 2, 693-707,
1311 2012.

1312 Galarneau, E.: Source specificity and atmospheric processing of airborne PAHs: Implications for
1313 source apportionment, *Atmos. Environ.*, 35, 8139-8149, 2008.

1314 Gao, M., Guttikunda, S. K., Carmichael, G. R., Wang, Y., Liu, Z., Stanier, C. O., Saide, P. E. and Yu,
1315 M.: Health impacts and economic losses assessment of the 2013 severe haze event in Beijing area, *Sci.*
1316 *Total Environ.*, 511, 553-561, 2015.

1317 Gao, S., Hegg, D. A., Hobbs, P. V., Kirchstetter, T. W., Magi, B. I. and Sadilek, M.: Water-soluble
1318 organic components in aerosols associated with savanna fires in southern Africa: Identification,
1319 evolution, and distribution, *J. Geophys. Res.: Atmos.*, D13, doi:10.1029/2002JD002324, 2003.

1320 Ge, X., Wexler, A. S. and Clegg, S. L.: Atmospheric amines-Part I. A review, *Atmos. Environ.*, 3,
1321 524-546, 2011.

1322 Ghorai, S., Wang, B., Tivanski, A. and Laskin, A.: Hygroscopic Properties of Internally Mixed
1323 Particles Composed of NaCl and Water-Soluble Organic Acids, *Environ. Sci. Technol.*, doi:
1324 10.1021/es404727u, 2014.

1325 Giordano, M., Espinoza, C. and Asa-Awuku, A.: Experimentally measured morphology of biomass
1326 burning aerosol and its impacts on CCN ability, *Atmos. Chem. Phys.*, 4, 1807-1821, 2015.

1327 Grieshop, A. P., Logue, J. M., Donahue, N. M. and Robinson, A. L.: Laboratory investigation of
1328 photochemical oxidation of organic aerosol from wood fires 1: measurement and simulation of organic
1329 aerosol evolution, *Atmos. Chem. Phys.*, 4, 1263-1277, 2009.

1330 Gustafsson, Ö., Kruså M., Zencak, Z., Sheesley, R. J., Granat, L., Engström, E., Praveen, P. S., Rao, P.,
1331 Leck, C. and Rodhe, H.: Brown clouds over South Asia: biomass or fossil fuel combustion? *Science*,
1332 5913, 495-498, 2009.

1333 Guttikunda, S. K. and Kopakka, R. V.: Source emissions and health impacts of urban air pollution in
1334 Hyderabad, India, *Air Quality, Atmos. & Health*, 2, 195-207, 2014.

1335 Han, Y. M., Cao, J. J., Ho, S. C. L. F. and An, Z. S.: Different characteristics of char and soot in the
1336 atmosphere and their ratio as an indicator for source identification in Xi 'an, China, *Atmos. Chem.*
1337 *Phys.*, 10, 595-607, 2010.

1338 Han, Y. M., Chen, L., Huang, R., Chow, J. C., Watson, J. G., Ni, H. Y., Liu, S. X., Fung, K. K., Shen,
1339 Z. X. and Wei, C.: Carbonaceous aerosols in megacity Xi'an, China: Implications of thermal/optical
1340 protocols comparison, *Atmos. Environ.*, 132, 58-68, 2016.

1341 Han, Y. M., Lee, S. C., Cao, J. J., Ho, K. F. and An, Z. S.: Spatial distribution and seasonal variation of
1342 char-EC and soot-EC in the atmosphere over China, *Atmos. Environ.*, 38, 6066-6073, 2009.

1343 Han, Y., Cao, J., Chow, J. C., Watson, J. G., An, Z., Jin, Z., Fung, K. and Liu, S.: Evaluation of the
1344 thermal/optical reflectance method for discrimination between char- and soot-EC, *Chemosphere*, 4,
1345 569-574, 2007.

1346 Hayashi, K., Ono, K., Kajiura, M., Sudo, S., Yonemura, S., Fushimi, A., Saitoh, K., Fujitani, Y. and
1347 Tanabe, K.: Trace gas and particle emissions from open burning of three cereal crop residues: Increase
1348 in residue moistness enhances emissions of carbon monoxide, methane, and particulate organic carbon,
1349 *Atmos. Environ.*, 95, 36-44, 2014.

1350 Hays, M. D., Fine, P. M., Geron, C. D., Kleeman, M. J. and Gullett, B. K.: Open burning of agricultural
1351 biomass: physical and chemical properties of particle-phase emissions, *Atmos. Environ.*, 36,
1352 6747-6764, 2005.

1353 He, K., Zhao, Q., Ma, Y., Duan, F. and Yang, F.: Spatial and seasonal variability of PM_{2.5} acidity at
1354 two Chinese megacities: insights into the formation of secondary inorganic aerosols, *Atmos. Chem.*
1355 *Phys. Dis.*, 25557-25603, doi:10.5194/acpd- 11- 25557- 2011, 2011a.

1356 He, M., Zheng, J., Yin, S. and Zhang, Y.: Trends, temporal and spatial characteristics, and uncertainties
1357 in biomass burning emissions in the Pearl River Delta, China, *Atmos. Environ.*, 24, 4051-4059, 2011b.

1358 Ho, K. F., Ho, S. S. H., Huang, R., Liu, S. X., Cao, J., Zhang, T., Chuang, H., Chan, C. S., Hu, D. and
1359 Tian, L.: Characteristics of water-soluble organic nitrogen in fine particulate matter in the continental
1360 area of China, *Atmos. Environ.*, 106, 252-261, 2015.

1361 Hou, Q., An, X., Wang, Y., Tao, Y. and Sun, Z.: An assessment of China's PM₁₀-related health
1362 economic losses in 2009, *Sci. Total. Environ.*, 61-65, 2012.

1363 Hu, Y., Lin, J., Zhang, S., Kong, L., Fu, H. and Chen, J.: Identification of the typical metal particles
1364 among haze, fog, and clear episodes in the Beijing atmosphere, *Sci. Total Environ.*, 369-380, 2015.

1365 Huang, K., Zhuang, G., Lin, Y., Fu, J. S., Wang, Q., Liu, T., Zhang, R., Jiang, Y., Deng, C. and Fu, Q.:
1366 Typical types and formation mechanisms of haze in an Eastern Asia megacity, Shanghai, *Atmos. Chem.*
1367 *Phys.*, 2012a.

1368 Huang, K., Zhuang, G., Lin, Y., Wang, Q., Fu, J. S., Fu, Q., Liu, T. and Deng, C.: How to improve the
1369 air quality over megacities in China: pollution characterization and source analysis in Shanghai before,
1370 during, and after the 2010 World Expo, *Atmos. Chem. Phys.*, 12, 5927-5942, 2013.

1371 Huang, R., Zhang, Y., Bozzetti, C., Ho, K., Cao, J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S.
1372 M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S., Brun, E., Crippa, M., Ciarelli, G., Piazzalunga, A.,
1373 Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. and Prevot, A.:
1374 High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 2014.

1375 Huang, S., Hsu, M. and Chan, C.: Effects of submicrometer particle compositions on cytokine
1376 production and lipid peroxidation of human bronchial epithelial cells., *Environ. Health Persp.*, 4, 478,
1377 2003.

1378 Huang, X., Li, M., Li, J. and Song, Y.: A high-resolution emission inventory of crop burning in fields
1379 in China based on MODIS Thermal Anomalies/Fire products, *Atmos. Environ.*, 9-15, 2012b.

1380 Huo, J., Lu, X., Wang, X., Chen, H., Ye, X., Gao, S., Gross, D. S., Chen, J. and Yang, X.: Online
1381 single particle analysis of chemical composition and mixing state of crop straw burning particles: from
1382 laboratory study to field measurement, *Front Env. Sci. Eng.*, 2, 244-252, 2016.

1383 IBRD and SEPA, Cost of pollution in China: economic estimates of physical damages, 2007, pp.
1384 1-128.

1385 IPCC, Greenhouse Gas Inventory Reference Manual: Revised 2006 IPCC Guidelines for National
1386 Greenhouse Gas Inventories. IPCC/OECD/IES, UK. Meteorological Office, Bracknell, UK., 2007.

1387 IPCC, Quantifying Uncertainties in Practice, Chapter 6: Good Practice Guidance and Uncertainty
1388 Management in National Greenhouse Gas Inventories: In: IES, IPCC, OECD, et al. Bracknell, UK,
1389 1997.

1390 Janssen, N. A. H., Hoek, G., Simic-Lawson, M., Fischer, P., van Bree, L., Ten Brink, H., Keuken, M.,
1391 Atkinson, R., Anderson, H., Brunekreef, B. and Cassee, F.: Black Carbon as an Additional Indicator of
1392 the Adverse Health Effects of Airborne Particles Compared with PM₁₀ and PM_{2.5}, Environ. Health
1393 Persp., 12, 1691-1699, 2011.

1394 Jayarathne, T., Stockwell, C. E., Yokelson, R. J., Nakao, S. and Stone, E. A.: Emissions of fine particle
1395 fluoride from biomass burning, Environ. Sci. Technol., 21, 12636-12644, 2014.

1396 Jenkins, B. M., Jones, A. D., Turn, S. Q. and Williams, R. B.: Emission factors for polycyclic aromatic
1397 hydrocarbons from biomass burning, Environ. Sci. Technol., 8, 2462-2469, 1996.

1398 Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., J, N. B. J., Duce, R. A.,
1399 Hunter, H., Mahowald, N. and Prospero, A.: Global Iron Connections Between Desert Dust, Ocean
1400 Biogeochemistry, and Climate, Science, 308, 67-71, 2005.

1401 Kennedy, I. M.: The health effects of combustion-generated aerosols, P. Combust Inst, 2, 2757-2770,
1402 2007.

1403 Kim, K., Jahan, S. A., Kabir, E. and Brown, R. J.: A review of airborne polycyclic aromatic
1404 hydrocarbons (PAHs) and their human health effects, Environ Int., 71-80, 2013.

1405 Kong, L., Yang, Y., Zhang, S., Zhao, X., Du, H., Fu, H., Zhang, S., Cheng, T., Yang, X. and Chen, J.:
1406 Observations of linear dependence between sulfate and nitrate in atmospheric particles, J. Geophys.
1407 Res.: Atmos., 1, 341-361, doi:10.1002/2013JD020222, 2014.

1408 Koopmans, A. and Koppejan, J.: Agricultural and forest residue-generation, utilization and availability,
1409 Modern Applications of Biomass Energy, 1997.

1410 Korenaga, T., Liu, X. and Huang, Z.: The influence of moisture content on polycyclic aromatic
1411 hydrocarbons emission during rice straw burning, Chemosphere-Global Change Science, 1, 117-122,
1412 2001

1413 Kundu, S., Kawamura, K., Andreae, T. W., Hoffer, A. and Andreae, M. O.: Molecular distributions of
1414 dicarboxylic acids, ketocarboxylic acids and α -dicarbonyls in biomass burning aerosols: implications
1415 for photochemical production and degradation in smoke layers, Atmos. Chem. Phys., 5, 2209-2225,
1416 2010.

1417 Laskin, A., Laskin, J. and Nizkorodov, S. A.: Chemistry of atmospheric brown carbon, Chem. Rev., 10,
1418 4335-4382, 2015.

1419 Lavanchy, V. M. H., G Ggeler, H. W., Nyeki, S. and Baltensperger, U.: Elemental carbon (EC) and
1420 black carbon (BC) measurements with a thermal method and an aethalometer at the high-alpine
1421 research station Jungfraujoch, Atmos. Environ., 17, 2759-2769, 1999.

1422 Lee, A. K. Y., Willis, M. D., Healy, R. M., Wang, J. M., Jeong, C. H., Wenger, J. C., Evans, G. J. and
1423 Abbatt, J.: Single particle characterization of biomass burning organic aerosol (BBOA): evidence for
1424 non-uniform mixing of high molecular weight organics and potassium, Atmos. Chem. Phys. Dis., 22,
1425 32157-32183, 2015.

1426 Lee, D. and Wexler, A. S.: Atmospheric amines-Part III: Photochemistry and toxicity, Atmos. Environ.,
1427 95-103, 2013.

1428 Lee, R. G. M., Coleman, P., Jones, J. L., Jones, K. C. and Lohmann, R.: Emission Factors and
1429 Importance of PCDD/Fs, PCBs, PCNs, PAHs and PM₁₀ from the Domestic Burning of Coal and Wood
1430 in the U.K., Environ. Sci. Technol., 6, 1436-1447, 2005.

1431 Leng, C., Zhang, Q., Zhang, D., Xu, C., Cheng, T., Zhang, R., Tao, J., Chen, J., Zha, S. and Zhang, Y.:
1432 Variations of cloud condensation nuclei (CCN) and aerosol activity during fog-haze episode: a case
1433 study from Shanghai, *Atmos. Chem. Phys.*, 22, 12499-12512, doi:10.5194/acp-14-12499-2014, 2014.

1434 Lewis, K. A., Arnott, W. P., Moosmuller, H., Chakrabarty, R. K., Carrico, C. M., Kreidenweis, S. M.,
1435 Day, D. E., Malm, W., Laskin, A., Jimenez, J., Ulbrich, I., Huffman, J., Onasch, T., Trimborn, A., Liu,
1436 L. and Mishchenko, M.: Reduction in biomass burning aerosol light absorption upon humidification:
1437 roles of inorganically-induced hygroscopicity, particle collapse, and photoacoustic heat and mass
1438 transfer, *Atmos. Chem. Phys.*, 9, 8949-8966, 2009.

1439 Levin, E. J. T., McMeeking, G. R., Carrico, C. M., Mack, L. E., Kreidenweis, S. M., Wold, C. E.,
1440 Moosmüller, H., Arnott, W., Hao, W., Collett, J. and Malm, W.: Biomass burning smoke aerosol
1441 properties measured during Fire Laboratory at Missoula Experiments (FLAME), *J. Geophys Res.*,
1442 D18210, doi:10.1029/2009JD013601, 2010.

1443 Li, C., Ma, Z., Chen, J., Wang, X., Ye, X., Wang, L., Yang, X., Kan, H., Donaldson, D. and Mellouki,
1444 A.: Evolution of biomass burning smoke particles in the dark, *Atmos. Environ.*, 120, 244-252, 2015.

1445 Li, C., Hu, Y., Chen, J., Ma, Z., Ye, X., Yang, X., Wang, L., Wang, X. and Mellouki, A.:
1446 Physicochemical properties of carbonaceous aerosol from agricultural residue burning: Density,
1447 volatility, and hygroscopicity, *Atmos. Environ.*, 2016.

1448 Li, J., Pósfai, M., Hobbs, P. V. and Buseck, P. R.: Individual aerosol particles from biomass burning in
1449 southern Africa: 2, Compositions and aging of inorganic particles, *J. Geophys. Res.: Atmos.*,
1450 (1984-2012), D13, doi:10.1029/2002JD002310, 2003.

1451 Li, X., Wang, S., Duan, L., Hao, J., Li, C., Chen, Y. and Yang, L.: Particulate and trace gas emissions
1452 from open burning of wheat straw and corn stover in China, *Environ. Sci. Technol.*, 17, 6052-6058,
1453 2007.

1454 Lima, A. L. C., Farrington, J. W. and Reddy, C. M.: Combustion-Derived Polycyclic Aromatic
1455 Hydrocarbons in the Environment-A Review, *Environ. Forensics.*, 2, 109-131, 2005.

1456 Lin, J., Nielsen, C. P., Zhao, Y., Lei, Y., Liu, Y. and McElroy, M. B.: Recent changes in particulate air
1457 pollution over China observed from space and the ground: effectiveness of emission control, *Environ.*
1458 *Sci. Technol.*, 20, 7771-7776, 2010.

1459 Lin, L., Gustafsson, E. and Strand, M.: Aerosol-based method for investigating biomass char reactivity
1460 at high temperatures, *Combust Flame*, 7, 1426-1437, 2011.

1461 Lin, P., Aiona, P. K., Li, Y., Shiraiwa, M., Laskin, J., Nizkorodov, S. A. and Laskin, A.: Molecular
1462 Characterization of Brown Carbon in Biomass Burning Aerosol Particles, *Environ. Sci. Technol.*, 21,
1463 11815-11824, 2016.

1464 Liu, M., Song, Y., Yao, H., Kang, Y., Li, M., Huang, X. and Hu, M.: Estimating emissions from
1465 agricultural fires in the North China Plain based on MODIS fire radiative power, *Atmos. Environ.*,
1466 326-334, 2015.

1467 Liu, Q. and Bei, Y.: Impacts of crystal metal on secondary aliphatic amine aerosol formation during
1468 dust storm episodes in Beijing, *Atmos. Environ.*, 227-234, 2016.

1469 Lobert, J. M., Scharffe, D. H., Hao, W. M. and Crutzen, P. J.: Importance of biomass burning in the
1470 atmospheric budgets of nitrogen-containing gases, 6284, 552-554, 1990.

1471 Lu, Z., Zhang, Q. and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in
1472 China and India, 1996-2010, *Atmos. Chem. Phys.*, 11, 9839-9864, doi:10.5194/acp-11-9839-2011,
1473 2011.

1474 Ma, J. and Van Aardenne, J. A.: Impact of different emission inventories on simulated tropospheric

1475 ozone over China: a regional chemical transport model evaluation, *Atmos. Chem. Phys.*, 4, 877-887,
1476 2004.

1477 May, A. A., McMeeking, G. R., Lee, T., Taylor, J. W., Craven, J. S., Burling, I., Sullivan, A. P., Akagi,
1478 S., Collett, J., Flynn, M., Coe, H., Urbanski, S., Seinfeld, J., Yokelson, R. and Kreidenweis, S.: Aerosol
1479 emissions from prescribed fires in the United States: A synthesis of laboratory and aircraft
1480 measurements, *J. Geophys. Res.: Atmos.*, 20, 11826-11849, doi:10.1002/2014JD021848, 2014.

1481 Meskhidze, N.: Dust and pollution: A recipe for enhanced ocean fertilization? *J. Geophys. Res.*, D3,
1482 doi:10.1029/2004JD005082, 2005.

1483 Mikhailov, E. F., Vlasenko, S. S., Podgorny, I. A., Ramanathan, V. and Corrigan, C. E.: Optical
1484 properties of soot-water drop agglomerates: An experimental study, *J. Geophys. Res.*, 111, D07209,
1485 doi:10.1029/2005JD006389, 2006.

1486 Moreno, T., Karanasiou, A., Amato, F., Lucarelli, F., Nava, S., Calzolari, G., Chiari, M., Coz, E.,
1487 Artinano, B., Lumberras, J., Borge, R., Boldo, R., Linares, C., Alastursy, A., Querol, X. and Gibbons,
1488 X.: Daily and hourly sourcing of metallic and mineral dust in urban air contaminated by traffic and
1489 coal-burning emissions, *Atmos. Environ.*, 68, 33-44, 2013.

1490 National Bureau of Statistics of China (NBSC): National Public Health Yearbook, China Statistics
1491 Press, Beijing (in Chinese), 2009.

1492 National Bureau of Statistics of China (NBSC): China Statistical Yearbook 2001~2013, China
1493 Statistics Press, Beijing (in Chinese), 2001~2013.

1494 National Bureau of Statistics of China (NBSC): China Rural Statistical Yearbook 2004~2013, China
1495 Statistics Press, Beijing (in Chinese), 2004~2013.

1496 National Health and Family Planning Commission (NHFPC): National Health and Family Planning
1497 Yearbook, Peking Union Medical College Press, 2013.

1498 Oanh, N. T. K., Ly, B. T., Tipayarom, D., Manandhar, B. R., Prapat, P., Simpson, C. D. and Liu, L. S.:
1499 Characterization of particulate matter emission from open burning of rice straw, *Atmos. Environ.*, 2,
1500 493-502, 2011.

1501 Ostro, B. and Chestnut, L.: Assessing the health benefits of reducing particulate matter air pollution in
1502 the United States, *Environ. Res.*, 2, 94-106, 1998.

1503 Pope III, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K. and Thurston, G. D.: Lung
1504 cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *Jama*, 9,
1505 1132-1141, 2002.

1506 Pope, C. A., Burnett, R. T., Thurston, G. D., Thun, M. J., Calle, E. E., Krewski, D. and Godleski, J. J.:
1507 Cardiovascular mortality and long-term exposure to particulate air pollution epidemiological evidence
1508 of general pathophysiological pathways of disease, *Circulation*, 1, 71-77, 2004.

1509 Pósfai, M.: Atmospheric tar balls: Particles from biomass and biofuel burning, *J. Geophys. Res.*, 109,
1510 D06213, doi:10.1029/2003JD004169, 2004.

1511 Qin, Y. and Xie, S. D.: Spatial and temporal variation of anthropogenic black carbon emissions in
1512 China for the period 1980-2009, *Atmos. Chem. Phys.*, 11, 4825-4841, 2012.

1513 Qin, Y. and Xie, S. D.: Historical estimation of carbonaceous aerosol emissions from biomass open
1514 burning in China for the period 1990-2005, *Environ Pollut.*, 12, 3316-3323, 2011.

1515 Qiu, C. and Zhang, R.: Physicochemical Properties of Alkylammonium Sulfates: Hygroscopicity,
1516 Thermostability, and Density, *Environ. Sci. Technol.*, 8, 4474-4480, 2012.

1517 Qiu, C., Wang, L., Lal, V., Khalizov, A. F. and Zhang, R.: Heterogeneous Reactions of Alkylamines
1518 with Ammonium Sulfate and Ammonium Bisulfate, *Environ. Sci. Technol.*, 11, 4748-4755, 2011.

1519 Ram, K., Sarin, M. M. and Tripathi, S. N.: Temporal Trends in Atmospheric PM_{2.5}, PM₁₀, Elemental
1520 Carbon, Organic Carbon, Water-Soluble Organic Carbon, and Optical Properties: Impact of Biomass
1521 Burning Emissions in The Indo-Gangetic Plain, *Environ. Sci. Technol.*, 2, 686-695, 2011.

1522 Ram, K. and Sarin, M. M.: Day-night variability of EC, OC, WSOC and inorganic ions in urban
1523 environment of Indo-Gangetic Plain: implications to secondary aerosol formation, *Atmos. Environ.*, 2,
1524 460-468, 2011.

1525 Reddy, M. S. and Venkataraman, C.: Atmospheric optical and radiative effects of anthropogenic
1526 aerosol constituents from India, *Atmos. Environ.*, 34, 4511-4523, 2000.

1527 Reid, J. S., Eck, T. F., Christopher, S. A., Koppmann, R., Dubovik, O., Eleuterio, D. P., Holben, B. N.,
1528 Reid, E. A. and Zhang, J.: A review of biomass burning emissions part III: intensive optical properties
1529 of biomass burning particles, *Atmos. Chem. Phys.*, 5, 827-849, 2005a.

1530 Reid, J. S., Koppmann, R., Eck, T. F. and Eleuterio, D. P.: A review of biomass burning emissions part
1531 II: intensive physical properties of biomass burning particles, *Atmos. Chem. Phys.*, 3, 799-825, 2005b.

1532 Richter, H. and J, H.: Formation of polycyclic aromatic hydrocarbons and their growth to soot-a review
1533 of chemical reaction pathways, *Prog Energ Combust*, 4, 565-608, 2000.

1534 Ripoll, A., Minguillón, M. C., Pey, J., Pérez, N., Querol, X. and Alastuey, A.: Joint analysis of
1535 continental and regional background environments in the western Mediterranean: PM₁ and PM₁₀
1536 concentrations and composition, *Atmos. Chem. Phys.*, 2, 1129-1145, doi:10.5194/acp-15-1129-2015,
1537 2015.

1538 Roemer, W. H. and van Wijnen, J. H.: Differences among Black Smoke, PM₁₀, and PM_{1.0} Levels at
1539 Urban Measurement Sites, *Environ. Health Persp.*, 2, 151-153, 2001.

1540 Rose, D., Gunthe, S. S., Su, H., Garland, R. M., Yang, H., Berghof, M., Cheng, Y. F., Wehner, B.,
1541 Achtert, P., Nowak, A., Wiedensohler, A., Takegawa, N., Kondo, Y., Hu, M., Zhang, Y., Andreae, M.
1542 and Poschl, U.: Cloud condensation nuclei in polluted air and biomass burning smoke near the
1543 mega-city Guangzhou, China-Part 2: Size-resolved aerosol chemical composition, diurnal cycles, and
1544 externally mixed weakly CCN-active soot particles, *Atmos. Chem. Phys.*, 6, 2817-2836,
1545 doi:10.5194/acp-11-2817-2011, 2011.

1546 Rosenfeld, D.: Atmosphere: Aerosols, Clouds, and Climate, *Science*, 5778, 1323-1324, 2006.

1547 Safai, P. D., Raju, M. P., Budhavant, K. B., Rao, P. and Devara, P.: Long term studies on
1548 characteristics of black carbon aerosols over a tropical urban station Pune, India, *Atmos. Res.*, 173-184,
1549 2013.

1550 Saffari, A., Daher, N., Samara, C., Voutsas, D., Kouras, A., Manoli, E., Karagkiozidou, O.,
1551 Vlachokostas, C., Moussiopoulos, N., Shafer, M., Schauer, J. and Sioutas, C.: Increased Biomass
1552 Burning Due to the Economic Crisis in Greece and Its Adverse Impact on Wintertime Air Quality in
1553 Thessaloniki, *Environ. Sci. Technol.*, 23, 13313-13320, 2013.

1554 Saikawa, E., Naik, V., Horowitz, L. W., Liu, J. and Mauzerall, D. L.: Present and potential future
1555 contributions of sulfate, black and organic carbon aerosols from China to global air quality, premature
1556 mortality and radiative forcing, *Atmos. Environ.*, 17, 2814-2822, 2009.

1557 Samy, S. and Hays, M. D.: Quantitative LC-MS for water-soluble heterocyclic amines in fine aerosols
1558 (PM_{2.5}) at Duke Forest, USA, *Atmos. Environ.*, 77-80, 2013.

1559 Santodonato, J.: Review of the estrogenic and antiestrogenic activity of polycyclic aromatic
1560 hydrocarbons: relationship to carcinogenicity, *Chemosphere*, 4, 835-848, 1997.

1561 Schade, G. W. and Crutzen, P. J.: Emission of aliphatic amines from animal husbandry and their
1562 reactions: Potential source of N₂O and HCN, *J. Atmos. Chem.*, 3, 319-346, 1995.

1563 Schauer, J. J., Kleeman, M. J., Cass, G. R. and Simoneit, B. R. T.: Measurement of Emissions from Air
1564 Pollution Sources. 3. C1–C29 Organic Compounds from Fireplace Combustion of Wood, *Environ. Sci.*
1565 *Technol.*, 9, 1716-1728, 2001.

1566 Schlesinger, R. B.: Comparative deposition of inhaled aerosols in experimental animals and humans: a
1567 review, *Journal of Toxicology and Environmental Health, Part A Current Issues*, 2, 197-214, 1985.

1568 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate*
1569 *change: John Wiley & Sons*, 2012.

1570 Sen, A., Mandal, T. K., Sharma, S. K., Saxena, M., Gupta, N. C., Gautam, R., Gupta, A., Gill, T., Rani,
1571 S., Saud, T., Singh, D. and Gadi, R.: Chemical properties of emission from biomass fuels used in the
1572 rural sector of the western region of India, *Atmos. Environ.*, 411-424, 2014.

1573 Shi, Y., Chen, J., Hu, D., Wang, L., Yang, X. and Wang, X.: Airborne submicron particulate (PM₁)
1574 pollution in Shanghai, China: Chemical variability, formation/dissociation of associated semi-volatile
1575 components and the impacts on visibility, *Sci. Total. Environ.*, 199-206, 2014.

1576 Shindell, D., Kuylensstierna, J. C. I., Vignati, E., van Dingenen, R., Amann, M., Klimont, Z., Anenberg,
1577 S. C., Muller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L., Kupiainen,
1578 K., Hoglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh, N., Milly, G.,
1579 Williams, M., Demkin, V. and Fowler, D.: Simultaneously Mitigating Near-Term Climate Change and
1580 Improving Human Health and Food Security, *Science*, 6065, 183-189, 2012.

1581 Simcik, M. F., Eisenreich, S. J. and Lioy, P. J.: Source apportionment and source/sink relationships of
1582 PAHs in the coastal atmosphere of Chicago and Lake Michigan, *Atmos. Environ.*, 30, 5071-5079,
1583 1999.

1584 Simoneit, B. R. T., Rushdi, A. I., Bin Abas, M. R. and Didyk, B. M.: Alkyl Amides and Nitriles as
1585 Novel Tracers for Biomass Burning, *Environ. Sci. Technol.*, 1, 16-21, 2003.

1586 Streets, D. G.: Dissecting future aerosol emissions: Warming tendencies and mitigation opportunities,
1587 *Climatic Change*, 3-4, 313-330, 2007.

1588 Sun, J., Peng, H., Chen, J., Wang, X., Wei, M., Li, W., Yang, L., Zhang, L., Wang, W. and Mellouki,
1589 A.: An estimation of CO₂ emission via agricultural crop residue open field burning in China from 1996
1590 to 2013, *J. Clean Prod.*, 2625-2631, 2016.

1591 Takegawa, N., Miyakawa, T., Kawamura, K. and Kondo, Y.: Contribution of Selected Dicarboxylic
1592 and ω-Oxocarboxylic Acids in Ambient Aerosol to the m/z 44 Signal of an Aerodyne Aerosol Mass
1593 Spectrometer, *Aerosol Sci. Tech.*, 41, 418-437, 2007.

1594 Tao, Y., Ye, X., Jiang, S., Yang, X., Chen, J., Xie, Y. and Wang, R.: Effects of amines on particle
1595 growth observed in new particle formation events, *J. Geophys. Res.: Atmos.*, 121, 324-335,
1596 doi:10.1002/2015JD024245, 2016.

1597 The World Bank (IBRD), State Environmental Protection Administration (SEPA): *Cost of pollution*
1598 *in China: economic estimates of physical damages*, 1-128, 2007.

1599 Tian, D., Hu, Y., Wang, Y., Boylan, J. W., Zheng, M. and Russell, A. G.: Assessment of Biomass
1600 Burning Emissions and Their Impacts on Urban and Regional PM_{2.5}: A Georgia Case Study, *Environ*
1601 *Sci. Technol.*, 2, 299-305, 2008.

1602 Tóth, A., Hoffer, A., Nyirő-Kósa, I., Pósfai, M. and Gelencsér, A.: Atmospheric tar balls: aged primary
1603 droplets from biomass burning? *Atmos. Chem. Phys. Dis.*, 12, 33089-33104, doi:10.5194/acp-14-
1604 6669-2014, 2013.

1605 Tsai, P. J., Shieh, H. Y., Lee, W. J. and Lai, S. O.: Health-risk assessment for workers exposed to
1606 polycyclic aromatic hydrocarbons (PAHs) in a carbon black manufacturing industry, *Sci. Total.*

1607 Environ., 1-3, 137-150, 2001.

1608 Urban, R. C., Alves, C. A., Allen, A. G., Cardoso, A. A. and Campos, M.: Organic aerosols in a
1609 Brazilian agro-industrial area: Speciation and impact of biomass burning, *Atmos. Res.*, 271-279, 2016.

1610 Veres, P., Roberts, J. M., Burling, I. R., Warneke, C., de Gouw, J. and Yokelson, R. J.: Measurements
1611 of gas-phase inorganic and organic acids from biomass fires by negative-ion proton-transfer
1612 chemical-ionization mass spectrometry, *J. Geophys Res*, 115, D23302, doi:10.1029/2010JD014033,
1613 2010.

1614 Wang, L., Li, X. and Xu, Y.: The Economic Losses Caused By Crop Residues Burnt in Open Field in
1615 China, *J. Arid Land Resources and Environment*, 2, 170-175, 2008a.

1616 Wang, R., Tao, S., Wang, W., Liu, J., Shen, H., Shen, G., Wang, B., Liu, X., Li, W., Huang, Y., Zhang,
1617 Y., Lu, Y., Chen, H., Chen, Y., Wang, C., Zhu, D., Wang, X., Li, B., Liu, W., Ma, J. and Prospero, A.:
1618 Black carbon emissions in China from 1949 to 2050, *Environ. Sci. Technol.*, 14, 7595-7603, 2012.

1619 Wang, S. and Zhang, C.: Spatial and temporal distribution of air pollutant emissions from open burning
1620 of crop residues in China, *Sciencepaper Online*, 5:329-333, 2008b.(*in chinese*)

1621 Wang, S., Zhao, M., Xing, J., Wu, Y., Zhou, Y., Lei, Y., He, K., Fu, L. and Hao, J.: Quantifying the air
1622 pollutants emission reduction during the 2008 Olympic Games in Beijing, *Environ. Sci. Technol.*, 7,
1623 2490-2496, 2010.

1624 Wang, W., Jariyasopit, N., Schrlau, J., Jia, Y., Tao, S., Yu, T., Dashwood, R. H., Zhang, W., Wang, X.
1625 and Simonich, S.: Concentration and photochemistry of PAHs, NPAHs, and OPAHs and toxicity of
1626 PM_{2.5} during the Beijing Olympic Games, *Environ. Sci. Technol.*, 16, 6887-6895, 2011.

1627 Wei, B. and Yang, L.: A review of heavy metal contaminations in urban soils, urban road dusts and
1628 agricultural soils from China, *Microchem J.*, 2, 99-107, 2010.

1629 Wei, W., Jitao, Y., Qingling, Z. and Bailiang, Z.: Current Situation and Developing Direction of Straw
1630 Utilization Technology in China, *China Resources Comprehensive Utilization*, 11, 2004.

1631 WHO, Life Database in 2000, World Health Organization, 2000.

1632 Wilson, J. M., Baeza-Romero, M. T., Jones, J. M., Pourkashanian, M., Williams, A., Lea-Langton, A.
1633 R., Ross, A. B. and Bartle, K.: Soot Formation from the Combustion of Biomass Pyrolysis Products
1634 and a Hydrocarbon Fuel, n-Decane: An Aerosol Time Of Flight Mass Spectrometer (ATOFMS) Study,
1635 *Energ Fuel*, 3, 1668-1678, 2013.

1636 Wong, C., Vichit-Vadakan, N., Kan, H., Qian, Z. and Teams, T. P. P.: Public Health and Air Pollution
1637 in Asia (PAPA): A Multicity Study of Short-Term Effects of Air Pollution on Mortality, *Environ.*
1638 *Health Persp.*, 9, 1195-1202, 2008.

1639 Wornat, M. J. and Sarofim, A. F.: Char- and Aerosol-associated Polycyclic Aromatic Compounds from
1640 Coal Pyrolysis, *Aerosol Sci. Tech.*, 12, 832-841, 2007.

1641 Wu, C., Liu, L. J. S., Cullen, A., Westberg, H. and Williamson, J., Spatial-temporal and cancer risk
1642 assessment of selected hazardous air pollutants in Seattle, *Environ Int.*, 1, 11-17, 2011.

1643 Wu, C., Wu, S., Wu, Y., Cullen, A. C., Larson, T. V., Williamson, J. and Liu, L. J. S.: Cancer risk
1644 assessment of selected hazardous air pollutants in Seattle, *Environ Int.*, 3, 516-522, 2009.

1645 Xu, S., Liu, W. and Tao, S.: Emission of Polycyclic Aromatic Hydrocarbons in China, *Environ Sci.*
1646 *Technol.*, 3, 702-708, 2006.

1647 Yang, C., Peng, X., Huang, W., Chen, R., Xu, Z., Chen, B. and Kan, H. . A time-stratified
1648 case-crossover study of fine particulate matter air pollution and mortality in Guangzhou, China, *Int*
1649 *Arch. Occ. Env. Hea.*, 5, 579-585, 2012.

1650 Yang, M., Howell, S. G., Zhuang, J. and Huebert, B. J.: Attribution of aerosol light absorption to black

1651 carbon, brown carbon, and dust in China-interpretations of atmospheric measurements during
1652 EAST-AIRE, *Atmos. Chem. Phys.*, 6, 2035-2050, doi:10.5194/acp-9-2035-2009, 2009.

1653 Yang, Y., Liu, X., Qu, Y., Wang, J., An, J., Zhang, Y. and Zhang, F.: Formation mechanism of
1654 continuous extreme haze episodes in the megacity Beijing, China, in January 2013, *Atmos. Res.*,
1655 192-203, 2015.

1656 Yokelson, R. J., Karl, T., Artaxo, P. and Blake, D. R.: The Tropical Forest and Fire Emissions
1657 Experiment: overview and airborne fire emission factor measurements, *Atmos. Chem. Phys.*, 7,
1658 5175-5196, doi:10.5194/acp-7-5175-2007, 2007.

1659 Yunker, M. B., Macdonald, R. W., Vingarzan, R., Mitchell, R. H., Goyette, D. and Sylvestre, S.: PAHs
1660 in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and
1661 composition, *Org Geochem*, 4, 489-515, 2002.

1662 Zhang, H., Hu, D., Chen, J., Ye, X., Wang, S. X., Hao, J. M., Wang, L., Zhang, R. and An, Z.: Particle
1663 Size Distribution and Polycyclic Aromatic Hydrocarbons Emissions from Agricultural Crop Residue
1664 Burning, *Environ. Sci. Technol.*, 13, 5477-5482, 2011.

1665 Zhang, H., Wang, S., Hao, J., Wan, L., Jiang, J., Zhang, M., Mestl, H. E., Mestl, H., Alnes, L., Aunan,
1666 K. and Mellouki, A.: Chemical and size characterization of particles emitted from the burning of coal
1667 and wood in rural households in Guizhou, China, *Atmos. Environ.*, 94-99, 2012.

1668 Zhang, H., Ye, X., Cheng, T., Chen, J., Yang, X., Wang, L. and Zhang, R.: A laboratory study of
1669 agricultural crop residue combustion in China: Emission factors and emission inventory, *Atmos.*
1670 *Environ.*, 36, 8432-8441, 2008a.

1671 Zhang, R., Khalizov, A. F., Pagels, J., Zhang, D., Xue, H. and McMurry, P. H.: Variability in
1672 morphology, hygroscopicity, and optical properties of soot aerosols during atmospheric processing,
1673 *Proc Natl Acad Sci USA*, 30, 10291-10296, 2008b.

1674 Zhang, R., Suh, I., Zhao, J., Zhang, D., Fortner, E. C., Tie, X., Molina, L. T. and Molina, M. J.:
1675 Atmospheric new particle formation enhanced by organic acids, *Science*, 5676, 1487-1490, 2004.

1676 Zhao, B., Wang, P., Ma, J. Z., Zhu, S., Pozzer, A. and Li, W.: A high-resolution emission inventory of
1677 primary pollutants for the Huabei region, China, *Atmos. Chem. Phys.*, 1, 481-501, 2012.

1678 Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B. and Hao, J.: Quantifying the Uncertainties of a
1679 Bottom-Up Emission Inventory of Anthropogenic Atmospheric Pollutants in China, *Atmos. Chem.*
1680 *Phys.*, 11, 2295-2308, 2011.

1681 Zheng, J., Ma, Y., Chen, M., Zhang, Q., Wang, L., Khalizov, A. F., Yao, L., Wang, Z., Wang, X. and
1682 Chen, L.: Measurement of atmospheric amines and ammonia using the high resolution time-of-flight
1683 chemical ionization mass spectrometry, 249-259, 2015.

1684

1685

1686 **Tables and figure captions**

1687 **Table 1.** Emission factors of particulate chemical species in smoke PM_{2.5} from
1688 agricultural residue burning (mean value ± standard deviation).

1689 **Table 2.** Emission factors of particulate chemical species in smoke PM_{1.0} from
1690 agricultural residue burning (mean value ± standard deviation).

1691 **Table 3.** Comparison of emission factors with literature (specific chemical materials
1692 in form of PM_{2.5})

1693 **Table 4.** Summary of field burning rates and economic data in China

1694 **Table 5.** National agricultural field burning emissions of BAU, EM, and NDRC
1695 scenarios in China, 2012.

1696 **Table 6.** Uncertainties for national smoke aerosol emissions in 2012.

1697 **Table 7.** Estimated number of cases (95% CI) attributable to agricultural fire smoke
1698 PM_{2.5} exposure in China, 2012

1699 **Table 8.** Health-related economic loss (95% CI) from agricultural fire smoke PM_{2.5}
1700 exposure in China, 2012

1701 **Figure 1.** Schematic methodology for developing emission estimations

1702 **Figure 2.** Chemical profiles of smoke PM_{2.5} and PM_{1.0} from 5 types agricultural
1703 residue burnings. OM (organic matter = 1.3×OC). OWSI, other water soluble ions
1704 including F⁻, NO₂⁻, Na⁺, Ca²⁺, Mg²⁺.

1705 **Figure 3.** a) Emission factors of 16 USEPA priority PAHs in smoke PM_{2.5} and PM_{1.0};
1706 b) expulsion-accumulation of PAHs in OC-EC of smoke PM_{2.5} and PM_{1.0}

1707 **Figure 4.** Transmission electron microscope (TEM) images and EDX analysis of
1708 fresh agricultural residue burning particles. (a)-(c) Crystal and amorphous KCl
1709 particles internally mixed with sulfate, nitrate, and carbonaceous materials. (d)-(f)
1710 Heavy metal-bearing fractal-like fly ash particles. (e)-(g) Chain-like soot particles and
1711 tar ball.

1712 **Figure 5.** Annual agricultural residue production of five major crops and allocated
1713 into two harvest (summer and autumn harvest) based on agricultural yield in China,
1714 2012.

1715 **Figure 6.** Statistical analysis of field burning rates from BAU, EM, and NDRC
1716 versions

1717 **Figure 7.** Spatial and temporal distribution of smoke PM_{2.5} emissions and flux
1718 concentrations from agricultural field burning over China, 2012

1719 **Figure 8.** Nationwide PM_{2.5} emissions and flux concentrations based on different
1720 burning versions. The inset pie-graphs are chemical compositions of integrated PM_{2.5}
1721 from five major agricultural residue burning.
1722

1723
1724

Table 1. Emission factors of particulate chemical species in smoke PM_{2.5} from agricultural residue burning (mean value ± standard deviation).

Chemical Species (g kg ⁻¹)	wheat straw	corn straw	rice straw	cotton residue	soybean residue
PM _{2.5}	5.803 ± 0.363	5.988 ± 0.723	14.732 ± 2.417	15.162 ± 2.053	3.249 ± 0.350
OC	2.813 ± 0.147	2.393 ± 0.351	6.882 ± 0.689	7.415 ± 0.547	1.539 ± 0.253
EC	0.676 ± 0.027	0.778 ± 0.152	2.182 ± 0.278	1.192 ± 0.171	0.614 ± 0.190
Char-EC	0.606 ± 0.024	0.667 ± 0.132	1.761 ± 0.166	1.072 ± 0.154	0.564 ± 0.177
Soot-EC	0.069 ± 0.007	0.110 ± 0.043	0.421 ± 0.061	0.120 ± 0.034	0.051 ± 0.031
Inorganic ions (g kg⁻¹)	1.273 ± 0.072	1.810 ± 0.030	3.086 ± 0.266	3.810 ± 0.246	0.523 ± 0.149
SO ₄ ²⁻	0.084 ± 0.028	0.217 ± 0.041	0.409 ± 0.127	0.701 ± 0.081	0.073 ± 0.014
Cl ⁻	0.576 ± 0.038	0.709 ± 0.034	1.158 ± 0.232	1.351 ± 0.114	0.178 ± 0.030
F ⁻	0.023 ± 0.061	0.061 ± 0.005	0.073 ± 0.024	0.265 ± 0.012	0.009 ± 0.004
NO ₃ ⁻	0.023 ± 0.000	0.032 ± 0.002	0.051 ± 0.025	0.072 ± 0.004	0.009 ± 0.004
NO ₂ ⁻	0.006 ± 0.001	0.016 ± 0.002	0.018 ± 0.002	0.036 ± 0.001	0.004 ± 0.003
Ca ²⁺	0.030 ± 0.011	0.036 ± 0.003	0.046 ± 0.007	0.060 ± 0.003	0.010 ± 0.002
Na ⁺	0.005 ± 0.001	0.012 ± 0.001	0.028 ± 0.004	0.050 ± 0.004	0.005 ± 0.001
NH ₄ ⁺	0.152 ± 0.005	0.197 ± 0.010	0.542 ± 0.107	0.347 ± 0.008	0.029 ± 0.004
Mg ²⁺	0.005 ± 0.000	0.017 ± 0.002	0.023 ± 0.004	0.032 ± 0.002	0.005 ± 0.001
K ⁺	0.368 ± 0.041	0.514 ± 0.009	0.739 ± 0.049	0.947 ± 0.070	0.200 ± 0.023
Organic Acids (mg kg⁻¹)	156.680 ± 81.830	46.670 ± 9.000	557.130 ± 269.380	769.990 ± 317.550	143.310 ± 39.770
CH ₃ COOH	148.900 ± 79.290	36.640 ± 8.210	417.930 ± 186.140	743.320 ± 159.600	135.500 ± 62.320
MSA	7.170 ± 2.110	10.030 ± 30.000	136.990 ± 81.700	12.980 ± 1.530	3.200 ± 1.530
H ₂ C ₂ O ₄	2.610 ± 0.430	ND	2.210 ± 1.560	4.760 ± 2.640	2.170 ± 2.380
HCOOH	ND	ND	ND	8.930 ± 2.630	2.440 ± 1.450
Amine salts (mg kg⁻¹)	19.246 ± 9.368	32.877 ± 19.141	104.787 ± 15.635	102.409 ± 13.379	4.514 ± 1.776
MeOH ⁺ + MMAH ⁺	1.322 ± 0.086	5.735 ± 0.102	17.226 ± 1.454	19.888 ± 0.351	0.456 ± 0.196
MEAH ⁺	0.201 ± 0.055	0.675 ± 0.135	4.175 ± 0.920	3.690 ± 1.959	ND
TEOH ⁺	2.562 ± 0.962	4.118 ± 0.741	25.129 ± 0.343	14.376 ± 8.688	0.672 ± 0.558
DEAH ⁺ + TMAH ⁺	13.728 ± 7.512	18.973 ± 0.466	46.148 ± 12.185	28.568 ± 5.321	2.012 ± 0.878
DMAH ⁺	1.434 ± 0.925	3.376 ± 0.674	12.110 ± 6.166	35.887 ± 2.940	1.374 ± 0.144
Elemental Species (mg kg⁻¹)	53.813 ± 18.860	53.546 ± 9.070	131.612 ± 5.920	27.577 ± 3.700	14.003 ± 8.710
Phenols (mg kg⁻¹)	26.785 ± 8.582	16.390 ± 2.652	27.238 ± 4.861	41.481 ± 5.517	9.673 ± 2.272
PAHs (mg kg⁻¹)	1.814 ± 0.348	2.706 ± 0.798	7.267 ± 1.722	8.302 ± 2.856	1.832 ± 0.353

1725

ND means not detected

1726
1727

Table 2. Emission factors of particulate chemical species in smoke PM_{1.0} from agricultural residue burning (mean value ± standard deviation).

Chemical Species (g kg ⁻¹)	wheat straw	corn straw	rice straw	cotton residue	soybean residue
PM _{1.0}	5.298 ± 0.295	5.360 ± 0.551	13.200 ± 1.440	12.635 ± 1.243	3.036 ± 0.257
OC	2.419 ± 0.126	2.063 ± 0.340	6.024 ± 0.602	6.036 ± 0.360	1.338 ± 0.128
EC	0.650 ± 0.037	0.728 ± 0.122	2.083 ± 0.413	1.023 ± 0.205	0.575 ± 0.260
Char-EC	0.567 ± 0.033	0.580 ± 0.098	1.671 ± 0.331	0.916 ± 0.184	0.511 ± 0.233
Soot-EC	0.083 ± 0.014	0.148 ± 0.057	0.411 ± 0.073	0.107 ± 0.048	0.063 ± 0.057
Inorganic ions (g kg⁻¹)	1.215 ± 0.040	1.768 ± 0.010	2.940 ± 0.249	3.516 ± 0.145	0.510 ± 0.156
SO ₄ ²⁻	0.078 ± 0.011	0.199 ± 0.032	0.333 ± 0.107	0.581 ± 0.054	0.073 ± 0.056
Cl ⁻	0.544 ± 0.033	0.712 ± 0.027	1.145 ± 0.118	1.243 ± 0.067	0.175 ± 0.031
F ⁻	0.022 ± 0.007	0.041 ± 0.004	0.078 ± 0.030	0.151 ± 0.011	0.001 ± 0.001
NO ₃ ⁻	0.021 ± 0.005	0.027 ± 0.002	0.043 ± 0.016	0.061 ± 0.003	0.009 ± 0.002
NO ₂ ⁻	0.006 ± 0.001	0.010 ± 0.003	0.013 ± 0.004	0.019 ± 0.002	0.004 ± 0.003
Ca ²⁺	0.027 ± 0.013	0.028 ± 0.002	0.045 ± 0.008	0.067 ± 0.005	0.010 ± 0.002
Na ⁺	0.004 ± 0.000	0.012 ± 0.000	0.027 ± 0.003	0.056 ± 0.006	0.005 ± 0.002
NH ₄ ⁺	0.147 ± 0.005	0.191 ± 0.009	0.511 ± 0.067	0.401 ± 0.004	0.031 ± 0.005
Mg ²⁺	0.005 ± 0.001	0.035 ± 0.001	0.024 ± 0.006	0.033 ± 0.002	0.005 ± 0.001
K ⁺	0.359 ± 0.040	0.513 ± 0.015	0.721 ± 0.073	0.994 ± 0.067	0.197 ± 0.035
Organic Acids (mg kg⁻¹)	124.310 ± 25.170	47.830 ± 10.610	427.400 ± 221.270	639.820 ± 244.960	130.760 ± 59.310
CH ₃ COOH	115.790 ± 21.940	38.960 ± 9.610	383.360 ± 179.050	615.790 ± 232.860	124.310 ± 69.000
MSA	6.830 ± 2.030	8.870 ± 2.730	41.380 ± 38.480	11.380 ± 2.360	3.200 ± 1.730
H ₂ C ₂ O ₄	1.690 ± 1.200	ND	2.660 ± 1.760	3.620 ± 1.250	1.560 ± 1.670
HCOOH	ND	ND	ND	9.030 ± 7.710	1.690 ± 1.390
Amine salts (mg kg⁻¹)	18.191 ± 5.351	29.891 ± 13.480	81.726 ± 11.455	85.720 ± 21.337	4.385 ± 1.445
MeOH ⁺ + MMAH ⁺	1.300 ± 0.282	5.647 ± 0.342	16.627 ± 0.104	18.834 ± 1.991	0.464 ± 0.265
MEAH ⁺	0.157 ± 0.037	0.787 ± 0.211	3.581 ± 0.602	2.771 ± 1.304	ND
TEOH ⁺	1.719 ± 0.283	5.115 ± 0.732	17.575 ± 0.844	11.441 ± 3.229	0.529 ± 0.304
DEAH ⁺ + TMAH ⁺	13.716 ± 9.047	15.921 ± 1.620	33.565 ± 6.795	29.057 ± 3.793	2.278 ± 0.533
DMAH ⁺	1.300 ± 0.702	2.420 ± 0.575	10.377 ± 4.521	23.617 ± 20.086	1.115 ± 0.343
Elemental Species (mg kg⁻¹)	31.586 ± 10.630	29.265 ± 4.240	51.062 ± 5.920	16.738 ± 3.480	11.817 ± 6.650
Phenols (mg kg⁻¹)	20.774 ± 4.972	13.193 ± 2.181	20.480 ± 1.403	23.521 ± 8.521	7.689 ± 1.356
PAHs (mg kg⁻¹)	1.257 ± 0.398	1.420 ± 0.232	3.967 ± 0.970	4.359 ± 1.373	1.123 ± 0.205

1728

ND means not detected

Table 3. Comparison of emission factors with literature (specific chemical materials in form of PM_{2.5}).

Species	Emission factors (g kg ⁻¹)		Reference
	This work	Reference value	
PM _{2.5}	8.99 ± 5.55	7.6~11.7(AR), 6.26~15.3 (TL), ~3.0 (AR), 2.2~15.0 (AR)	Li et al., 2007; Akagi et al., 2011; Dhammapala et al., 2007; Hayashi et al., 2014
PM _{1.0}	7.91 ± 4.67	4.4.3~12.1 (TL)	May et al., 2014
OC	4.21 ± 2.73	2.7~3.9 (AR), 2.3~9.7(TL), ~1.9(AR) , 1.0~9.3 (AR), 0.8~5.9 (TL)	Li et al., 2007; Akagi et al., 2011; Dhammapala et al., 2007; Hayashi et al., 2014; May et al., 2014
EC	1.09 ± 0.65	0.35~0.49 (AR), 0.37~0.91(TL), ~0.4(AR), 0.21~0.81(AR), 1.13~1.73 (TL)	Li et al., 2007; Akagi et al., 2011; Dhammapala et al., 2007; Hayashi et al., 2014; May et al., 2014
WSOA	0.33 ± 0.31	0.039~0.109 (TL)	Akagi et al., 2011
WSA	0.05 ± 0.05	0.08~0.13 (TL), ~0.55 (TL)	Akagi et al., 2011; Andreae et al., 2001
WSI	2.10 ± 1.34	1.84~4.9 (AR),0.8~1.31(TL), 0.43~1.63 (AR)	Li et al., 2007; Akagi et al., 2011; Hayashi et al., 2014
THM	0.06 ± 0.05	0.06~0.09 (AR)	Li et al., 2007
PAHs (×10 ³)	4.38 ± 3.15	~17(AR), 0.72~1.64(AR), ~9.0 (W)	Dhammapala et al., 2007; Zhang et al., 2011; Lee et al.2005
Phenols (×10 ³)	24.31 ± 12.11	~35(AR), ~5 (AR), ~13 (TL)	Dhammapala et al., 2007; Hays et al., 2005; Andreae et al., 2001

AR: agricultural residue; TL: total, including forest fires and straw burning; W: wood

Table 4. Summary of field burning rates and economic data in China.

Province	Burning rate from literature		Agricultural income ratio ^c			Estimated burning rate		NDRC report ^d	Average rate
	BAU-I ^a	BAU-II ^b	2000	2006	2012	EM-I	EM-II	NDRC	
Beijing	0.00	0.17	0.08	0.06	0.06	0.00	0.19	0.13	0.10 ±0.08
Tianjin	0.00	0.17	0.10	0.14	0.12	0.00	0.20	0.30	0.13 ±0.12
Hebei	0.20	0.17	0.27	0.22	0.24	0.22	0.16	0.19	0.19 ±0.02
Shanxi	0.20	0.17	0.20	0.21	0.25	0.16	0.14	0.22	0.18 ±0.03
Inner Mongolia	0.00	0.12	0.44	0.49	0.66	0.00	0.09	0.27	0.10 ±0.10
Liaoning	0.20	0.12	0.30	0.29	0.39	0.16	0.09	0.34	0.18 ±0.09
Jilin	0.30	0.12	0.73	0.73	0.77	0.28	0.11	0.25	0.21 ±0.08
Heilongjiang	0.30	0.12	0.99	0.83	0.59	0.50	0.17	0.25	0.27 ±0.13
Shanghai	0.00	0.32	0.10	0.08	0.09	0.00	0.29	0.12	0.15 ±0.14
Jiangsu	0.30	0.32	0.32	0.22	0.30	0.32	0.23	0.19	0.27 ±0.05
Zhejiang	0.30	0.32	0.19	0.08	0.09	0.64	0.28	0.22	0.35 ±0.15
Anhui	0.20	0.32	0.44	0.39	0.43	0.21	0.29	0.43	0.29 ±0.08
Fujian	0.30	0.32	0.18	0.10	0.14	0.39	0.22	0.17	0.28 ±0.08
Jiangxi	0.20	0.11	0.45	0.31	0.44	0.20	0.08	0.25	0.17 ±0.06
Shandong	0.30	0.17	0.33	0.25	0.24	0.40	0.17	0.21	0.25 ±0.09
Henan	0.20	0.17	0.39	0.35	0.33	0.23	0.18	0.22	0.20 ±0.02
Hubei	0.20	0.11	0.42	0.30	0.41	0.21	0.08	0.30	0.18 ±0.08
Hunan	0.20	0.33	0.47	0.31	0.43	0.22	0.24	0.35	0.27 ±0.06
Guangdong	0.30	0.33	0.19	0.10	0.13	0.44	0.25	0.18	0.30 ±0.09
Guangxi	0.20	0.33	0.40	0.25	0.33	0.25	0.25	0.35	0.28 ±0.06
Hainan	0.30	0.33	0.35	0.16	0.21	0.51	0.25	0.56	0.39 ±0.12
Chongqing	0.20	0.11	0.35	0.23	0.30	0.24	0.08	0.45	0.22 ±0.13
Sichuan	0.20	0.11	0.37	0.22	0.28	0.26	0.09	0.30	0.19 ±0.08
Guizhou	0.20	0.11	0.38	0.23	0.25	0.31	0.10	0.43	0.23 ±0.13
Yunnan	0.20	0.11	0.36	0.26	0.31	0.24	0.09	0.28	0.18 ±0.07
Tibet	0.00	0.16	0.15	0.09	0.05	0.00	0.30	0.16	0.12 ±0.11
Shannxi	0.20	0.17	0.33	0.27	0.26	0.25	0.18	0.28	0.22 ±0.04
Gansu	0.10	0.16	0.25	0.20	0.28	0.09	0.11	0.33	0.16 ±0.09
Qinghai	0.00	0.16	0.23	0.10	0.08	0.00	0.20	0.28	0.13 ±0.11
Ningxia	0.10	0.16	0.42	0.38	0.45	0.09	0.13	0.16	0.13 ±0.03
Xinjiang	0.10	0.16	0.43	0.61	0.73	0.06	0.13	0.30	0.15 ±0.08
Nationwide	0.21	0.16	0.34	0.27	0.31	0.26	0.15	0.27	0.21 ±0.05

a. Zhao et al., 2012; Cao et al., 2006; Cao et al., 2011

b. Wang and Zhang., 2008

c. Calculated based on data from China Yearbook 2001~2013 (NBSC, 2001-2013), China Rural Statistic Yearbook 2001~2013, data available at <http://www.grain.gov.cn/Grain/>

d. Data from the National Development and Reform Commission report ([2014]No.516) : <http://www.sdpc.gov.cn/>

Table 5. National agricultural field burning emissions of BAU, EM, and NDRC scenarios in China in 2012.

Unit: Gg	BAU-I			BAU-II			EM-1			EM-2			NDRC			Average		
	Total	Summer	Autumn	Total	Summer	Autumn	Total	Summer	Autumn	Total	Summer	Autumn	Total	Summer	Autumn	Total	Summer	Autumn
PM _{2.5}	1001.05	218.99	782.06	835.42	209.29	626.13	1211.92	258.58	953.34	738.36	182.34	556.02	1241.69	258.24	983.46	1007.650	226.007	781.646
PM _{1.0}	897.52	198.93	698.59	748.57	189.92	558.65	1087.05	234.85	852.20	661.81	165.61	496.20	1111.90	234.44	877.46	903.125	205.217	697.911
OC	429.51	102.87	326.64	360.99	97.67	263.32	519.26	121.33	397.94	318.84	85.55	233.29	533.19	120.86	412.33	433.184	105.885	327.300
EC	133.61	27.37	106.24	111.40	26.52	84.88	162.71	32.39	130.32	98.06	22.85	75.21	164.97	32.53	132.45	134.414	28.404	106.010
char-EC	112.75	23.76	88.99	93.82	22.88	70.94	137.15	28.09	109.06	82.79	19.81	62.98	139.21	28.14	111.07	113.366	24.596	88.770
soot-EC	20.80	3.59	17.21	17.54	3.62	13.91	25.50	4.28	21.22	15.23	3.02	12.21	25.70	4.36	21.33	20.992	3.787	17.205
SO ₄ ²⁻	30.22	3.96	26.26	24.97	3.94	21.04	36.39	4.71	31.68	22.09	3.32	18.76	38.21	4.78	33.44	30.440	4.155	26.285
NO ₃ ⁻	4.35	0.84	3.51	3.55	0.80	2.75	5.24	0.99	4.25	3.17	0.70	2.47	5.40	0.99	4.41	4.350	0.864	3.486
NH ₄ ⁺	32.08	6.37	25.71	26.65	6.21	20.44	39.09	7.54	31.55	23.43	5.32	18.11	39.46	7.59	31.87	32.202	6.623	25.580
K ⁺	67.49	13.12	54.38	54.75	12.38	42.37	81.40	15.45	65.95	49.10	10.90	38.20	83.62	15.36	68.26	67.412	13.469	53.943
WSOA	24.44	6.55	17.89	21.94	6.39	15.55	29.69	7.76	21.93	18.77	5.48	13.30	30.82	7.81	23.01	25.174	6.815	18.360
WSA	5.75	0.95	4.80	4.85	0.95	3.90	6.99	1.13	5.86	4.23	0.80	3.43	7.19	1.15	6.04	5.815	1.000	4.815
PAHs	0.48	0.11	0.37	0.40	0.10	0.30	0.58	0.12	0.45	0.35	0.09	0.26	0.59	0.13	0.47	0.480	0.109	0.371
Phenols	2.71	0.85	1.87	2.25	0.78	1.47	3.25	0.99	2.26	2.02	0.70	1.323	3.40	0.98	2.36	2.721	0.861	1.861
THM	8.68	2.01	6.67	7.19	1.92	5.27	10.56	2.37	8.19	6.36	1.67	4.69	10.64	2.37	8.27	8.702	2.073	6.628
WSI	249.96	47.46	202.50	204.46	45.24	159.22	301.75	56.01	245.74	182.31	39.50	142.82	310.31	55.88	254.43	250.269	48.927	201.342

Table 6. Uncertainties for the national smoke aerosol emissions in 2012 (pollutant emission in unit of Gg/yr, 95% CI in percentage)


Species	BAU-I		BAU-II		EM-I		EM-II		NDRC		Average	
PM _{2.5}	1001.1	(-52.3% , 73.5%)	835.4	(-48.7% , 68.8%)	1211.9	(-63.6% , 84.3%)	738.4	(-55.9% , 74.3%)	1241.7	(-46.2% , 65.1%)	1005.7	(-24.6% , 33.7%)
PM _{1.0}	897.5	(-51.6% , 73.0%)	748.6	(-48.4% , 68.6%)	1087.1	(-62.9% , 83.8%)	661.8	(-55.5% , 74.1%)	1111.9	(-45.7% , 64.7%)	901.4	(-24.4% , 33.5%)
OC	429.5	(-50.5% , 71.5%)	361.0	(-48.9% , 69.2%)	519.3	(-61.4% , 81.8%)	318.8	(-55.6% , 74.1%)	533.2	(-47.1% , 66.7%)	432.4	(-24.2% , 33.3%)
EC	133.6	(-52.1% , 73.6%)	111.4	(-50.1% , 71.0%)	162.7	(-63.3% , 84.3%)	98.1	(-56.8% , 75.7%)	165.0	(-46.7% , 66.0%)	134.2	(-24.8% , 34.0%)
char-EC	112.8	(-51.1% , 73.3%)	93.8	(-49.4% , 69.9%)	137.2	(-63.1% , 84.0%)	82.8	(-60.8% , 80.7%)	139.2	(-46.2% , 65.4%)	113.1	(-24.8% , 34.1%)
soot-EC	20.8	(-53.7% , 74.7%)	17.5	(-55.3% , 77.6%)	25.5	(-65.9% , 87.4%)	15.2	(-61.8% , 81.9%)	25.7	(-50.6% , 71.1%)	21.0	(-26.3% , 35.9%)
WSOA	24.4	(-68.5% , 86.2%)	21.9	(-75.7% , 95.2%)	29.7	(-78.7% , 96.2%)	18.8	(-77.8% , 95.4%)	30.8	(-67.5% , 85.1%)	25.1	(-33.3% , 41.4%)
WSA	5.8	(-62.8% , 82.1%)	4.9	(-65.9% , 84.1%)	7.0	(-73.9% , 93.2%)	4.2	(-69.3% , 86.3%)	7.2	(-58.7% , 75.9%)	5.8	(-30.1% , 38.5%)
WSI	250.0	(-54.4% , 77.2%)	204.5	(-47.5% , 67.4%)	301.8	(-66.9% , 89.3%)	182.3	(-56.1% , 74.8%)	310.3	(-46.9% , 66.4%)	249.8	(-25.4% , 34.9%)
THM	8.7	(-56.2% , 77.5%)	7.2	(-52.8% , 71.4%)	10.6	(-67.5% , 88.3%)	6.4	(-61.2% , 79.5%)	10.6	(-50.8% , 69.4%)	8.7	(-26.6% , 35.6%)
PAHs	0.5	(-55.2% , 75.7%)	0.4	(-52.4% , 72.2%)	0.6	(-66.5% , 86.8%)	0.4	(-58.8% , 76.9%)	0.6	(-49.3% , 67.8%)	0.5	(-26.0% , 34.9%)
Phenols	2.7	(-56.1% , 77.6%)	2.3	(-51.4% , 70.6%)	3.3	(-67.3% , 88.3%)	2.0	(-59.9% , 78.4%)	3.4	(-48.7% , 67.1%)	2.7	(-26.1% , 35.1%)

Table 7. Estimated number of cases (95% CI) attributable to agricultural fire smoke PM_{2.5} exposure in China, 2012.

Emission version	Mortality	Respiratory hospital admission	Cardiovascular hospital admission	Chronic bronchitis
BAU-I	7864 (3154, 12489)	31123 (21114, 40788)	29454 (12849, 45481)	7577067 (2952006, 11024705)
BAU-II	7187 (3056, 11260)	28711 (19443, 37693)	27156 (11825, 42007)	7132581 (2735111, 10523803)
EM-I	9435 (3817, 14933)	36950 (25151, 48269)	35116 (15373, 54042)	8712880 (3484325, 12430411)
EM-II	6175 (2554, 9751)	25166 (17004, 33112)	23745 (10316, 36816)	6383442 (2407643, 9526727)
NDRC	8523 (3581, 13377)	33957 (23015, 44542)	32131 (14003, 49664)	8332216 (3228351, 12148274)
Average	7836 (3232, 12362)	31181 (21145, 40881)	29520 (12873, 45602)	7267237 (2961487, 1130784)
CRC	538 (227, 850)	2191 (1462, 2920)	2038 (874, 3199)	636650 (214617, 1052153)


Table 8. Health-related economic loss (95% CI) from agricultural fire smoke PM_{2.5} exposure in China, 2012.

Emission version	Economic cost (million US\$)				Total cost (million US\$)	GDP ratio (‰)
	Mortality	Respiratory hospital admission	Cardiovascular hospital admission	Chronic bronchitis		
BAU-1	1544.5 (730.7, 2430.0)	19.6 (13.3, 25.7)	36.0 (15.7, 55.6)	7187.6 (2800.3, 10458.3)	8787.8 (3560.0, 12969.4)	1.0 (0.4 , 1.5)
BAU-2	1453.9 (719, 2252.2)	18.1 (12.2, 23.8)	33.2 (14.4, 51.3)	6766.0 (2594.5, 9982.9)	8271.2 (3340.3, 12310.3)	1.0 (0.4 , 1.4)
EM-1	1855.2 (870.3, 2913.7)	23.3 (15.9, 30.5)	42.9 (18.8, 66.1)	8265.0 (3305.2, 11791.5)	10186.5 (4210.2, 14801.8)	1.2 (0.5 , 1.7)
EM-2	1228.1 (600.6, 1917.6)	15.9 (10.7, 20.9)	29.0 (12.6, 45.0)	6055.3 (2283.9, 9037.1)	7328.4 (2907.9, 11020.7)	0.9 (0.3 , 1.3)
NDRC	1573.4 (759.3, 2456.2)	21.4 (14.5, 28.1)	39.3 (17.1, 60.7)	7903.9 (3062.4, 11523.9)	9538.2 (3853.4, 14069.0)	1.1 (0.4 , 1.6)
Average	1531.0 (736.0, 2393.9)	19.7 (13.3, 25.8)	36.1 (15.7, 55.7)	7235.6 (2809.3, 10558.7)	8822.4 (3574.4, 13034.2)	1.0 (0.4 , 1.5)
CRC	100.0 (48.0, 157.1)	1.3 (0.9, 1.8)	2.4 (1.0, 3.9)	603.9 (203.6, 998.1)	707.8 (253.6, 1160.9)	0.1 (0.0 , 0.1)




1
2
3

Figure 1. Schematic methodology for developing emission estimations.




5

6 **Figure 2.** Chemical profiles of smoke PM_{2.5} and PM_{1.0} from 5 types agricultural
 7 residue burnings. OM (organic matter = 1.3×OC). OWSI, other water soluble ions
 8 including F⁻, NO₂⁻, Na⁺, Ca²⁺, and Mg²⁺.




9
10
11
12

Figure 3. (a) Emission factors of 16 USEPA priority PAHs in smoke PM_{2.5} and PM_{1.0};
(b) expulsion-accumulation of PAHs in OC-EC of smoke PM_{2.5} and PM_{1.0}.



13
 14 **Figure 4.** Transmission electron microscope (TEM) images and EDX analysis of
 15 fresh agricultural residue burning particles. (a)-(c) Crystal and amorphous KCl
 16 particles internally mixed with sulfate, nitrate, and carbonaceous materials. (d)-(f)
 17 Heavy metal-bearing fractal-like fly ash particles. (e)-(g) Chain-like soot particles and
 18 tar ball.


19



20

21 **Figure 5.** Annual agricultural residue production of five major crops and allocated
 22 into two harvest (summer and autumn harvest) based on agricultural yield in China,
 23 2012. (Abbreviation, BJ: Beijing; TJ: Tianjin; HB: Hebei; SX: Shanxi; IM: Inner Mongolia; LN: Liaoning; JL:
 24 Jilin; HLJ: Heilongjiang; SH: Shanghai; JS: Jiangsu; ZJ: Zhejiang; AH: Anhui; FJ: Fujian; JX: Jiangxi; SD:
 25 Shandong; HN: Henan; HUB: Hubei; HUN: Hunan; GD: Guangdong; GX: Guangxi; HAN: Hainan; CQ:
 26 Chongqing; SC: Sichuan; GZ: Guizhou; YN: Yunnan; SHX: Shannxi; GS: Gansu; QH: Qinghai; NX: Ningxia; XJ:
 27 Xinjiang)


28



29


30 **Figure 6.** Statistical analysis of field burning rates from BAU, EM, and NDRC
 31 versions. The North Plain (Anhui, Shandong, Hebei, Shanxi, Tianjin, Beijing), the
 32 Central of China (Hunan, Henan, Hubei), the Yangtze River Delta (Zhejiang, Jiangsu,
 33 Shanghai), the Northeast of China (Heilongjiang, Liaoning, Jilin), the Pan-Pearl River
 34 Delta (Hainan, Guangdong, Fujian, Guangxi, Guizhou, Sichuan, Yunnan, Jiangxi), the
 35 West of China (Shannxi, Chongqing, Xinjiang, Qinghai, Ningxia, Tibet, Inner
 36 Mongolia, Gansu)

37



38

39 **Figure 7.** Spatial and temporal distribution of smoke $PM_{2.5}$ emissions and flux
 40 concentrations from agricultural field burning over China, 2012.



41

42 **Figure 8.** Nationwide $PM_{2.5}$ emissions and flux concentrations based on different
 43 burning versions. The inset pie-graphs are chemical compositions of integrated $PM_{2.5}$
 44 from five major agricultural residue burning.