

Interactive comment on “Sensitivity model study of regional mercury dispersion in the atmosphere” by Christian N. Gencarelli et al.

T. S. Dibble

tsdibble@esf.edu

Received and published: 8 August 2016

The mechanism used here for Br-initiated oxidation of mercury (from Gencarelli et al (2015)) cites Goodsite et al (2004) for the rate constants for HgBr reacting with Br and OH. These two rate constants are given as equal in Gencarelli et al. However, Goodsite et al did not report a rate constant for HgBr reaction with OH, so Gencarelli et al (2015) assumed that these two rate constants are equal.

If a model is going to treat these two rate constants as equal, it seems reasonable to include the reactions of HgBr with NO₂, HOO, ClO, and BrO with the same rate constant, as was suggested by Dibble et al (2012).

References

C1

Dibble, T. S., Zelic, M. J., and Mao, H. Thermodynamics of reactions of HgCl and HgBr radicals with atmospherically abundant free radicals. *Atmos. Chem. Phys.* 12, 10271-10279, 2012.

Gencarelli, C. N., De Simone, F., Hedgecock, I. M., Sprovieri, F., Yang, X., and Pirrone, N. European and Mediterranean mercury modelling: Local and long-range contributions to the deposition flux. *Atmos. Env.* 117, 162-168, 2015.

Goodsite, M. E., Plane, J. M. C., and Skov, H. A Theoretical Study of the Oxidation of Hg⁰ to HgBr₂ in the Troposphere. *Environ. Sci. Technol.*, 38, 1772-1776, 2004.

Interactive comment on Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2016-663, 2016.

C2