

We thank Referee N°1 for his very helpful comments and suggestions. We feel they have greatly improved our manuscript. We have addressed the comments below.

Comment 1: *Introduction: The introduction is well written; however, I think that some major studies have been forgotten or omitted. Recently, studies conducted at the Jungfraujoch appeared in Science and JGR-A. These studies should be mentioned as a comparison, especially in terms of CCN production, would be extremely valuable (Herrmann et al. 2015, Bianchi et al. 2016, Tröstl et al. 2016)*

Reply 1: Some of these studies are very recent and were not published before the initial submission of the present study to ACPD. They are now mentioned in the introduction, as well as in the discussion:

Introduction: “However, observations to validate these predictions are scarce, especially for the FT, where measurements are often technically challenging. Recent studies conducted at the Jungfraujoch station (Switzerland, 3580 m a.s.l.) reported significant enhancement of the particle concentration below 50 nm by NPF in the FT, while only a minor fraction of these particles grow beyond 90 nm, even on a time scale of several days (Herrmann et al., 2015; Tröstl et al., 2016). The contribution of NPF to the production of CCN is thus likely to be very limited in this part of the FT, while boundary layer originating particles were observed to dominate the CCN concentrations measured at Jungfraujoch. The occurrence of the NPF process itself in the FT was reported to be tightly connected with the strength of boundary layer influence at the site, together with global radiation (Bianchi et al., 2016; Tröstl et al., 2016).”

Comment 2: *Page 2 Line 29: The reference Yli-Juuti is only about one site (Hyytiälä). I would recommend to consider the Manninen et al. (2010) EUCAARI paper which reports findings from 12 European sites.*

Reply 2: It is true that the study by Yli-Juuti et al. investigate measurements from Hyytiälä. However, it provides a very useful comparison between the GRs obtained at various locations, including all observations by Manninen et al. 2010 along with other additional studies.

Comment 3: *Page 2 Line 30: Reference is needed.*

Reply 3: the synthesis by Kerminen et al. (2012) is now mentioned.

Comment 4: *Page 3 Line 17: “...However, observations to validate these predictions are scarce, especially for the FT...” It is true that little information is available in this specific field of research. However, new study by Tröstl et al., (2016) has just been published in the Journal of Geophysical Research (Atmospheres) which (among other things) investigates the contribution of new particle formation to the CCN concentration in the Alps in some detail. I think a comparison to this work (Alps vs Andes) would be quite interesting, especially considering the general scarcity of similar research.*

Reply 4: We do agree, however the paper by Tröstl et al. (2016) was not published (available online on September 8th) when the present study was submitted to ACPD (August 15th). The results from Jungfraujoch are now mentioned (See Reply 1).

Comment 5: *Page 4 Line 1,2: The authors underline the vicinity of the site to a city like La Paz that has a large population and is assumingly rather polluted. This fact seems weirdly underused in the study. Why not determine La Paz air masses to find out what (if any) effect polluted air masses have on NPF and CCN production? A backtrajectory analysis might actually be quite interesting.*

Reply 5: The effect of wind direction and air mass back trajectories on NPF occurrence and characteristics was already investigated by Rose et al. (2015b). NPF events are more frequent and also more intense during the dry season, in dominant north western winds corresponding to oceanic air masses (Fig. 11 of the mentioned study). The fastest particle growth, which are observed during the

west season, are in contrast observed in air masses that passed over the Amazon basin (Fig. 14 of the mentioned study). No distinctive feature connected to air masses from La Paz came out of this analysis. This is now shown on Fig S4 in the supplementary, and explicitly mentioned in Section 3.1.2:

“It is worth noticing that winds originating from the more polluted sector of La Paz – El Alto (south) do not seem to be over-represented neither on event nor on non-event days. However, because of the close proximity of this area, it is complex to further assess how it contributes to CCN concentration from wind direction alone, and we cannot exclude a bias related to the variability of this specific source between event and non-event days.”

Comment 6: *Section 2.2 Indirect method for the estimation of the NPF contribution to the CCN production I'm not sure that this method fulfilled what the authors claimed. I agree with the fact that the CCN increased during NPF is not just a pure coincidence since the NPF precursors certainly also facilitate growth. However, it's not possible to distinguish the CCN formed by the NPF events with the growth of pre-existing particles during the same time. An easy way to fix that somehow would be to assume that ALL pre-existing particles to become CCN before any new particles. I.e. the number of particles below 100 nm before the event must be subtracted from the CCN100 you are now using and so far for the other sizes. This still would not account for, say, 90 nm particles that are transported to the site during NPF and grow above the threshold and contribute to CCNmax but it would be better than the current approach.*

Comment 8: *Page 5 Line 1: "...The CCN production during an event was obtained from the comparison of the CCN concentration N_{init} prior to and the maximum CCN concentration N_{max} during the event..." I agree with the authors that this is the CCN production during an event. However, the authors treat this as the CCN increase due to NPF. As already mentioned, CCN increase DURING and BECAUSE OF NPF are not the same thing. This distinction demands for clarity of concept and language whenever the topic is discussed. The manuscript in its current form, however, conflates those things. Besides the need for exact language, I actually think the issue can be addressed (to an extent) as lined out above.*

Comment 9: *Page 5 Line4 "...tinit, when nucleated particles reach the threshold size..." I don't think is possible to know that nucleated particles reached CCN size instead of larger particles that simply have grown above the threshold. The respective figure1 actually shows that t_{init} isn't found as the text claims: if the text was true, then t_{init_100} should be well after t_{init_50} because growth takes time. In the figure, however, all t_{init} are the same. That means t_{init} is really just the time when CCN numbers start to increase. But that increase doesn't likely come from NPF. Figure 1 illustrates a further problem with this claim: t_{init_100} is roughly 1.5...2 h after nucleation onset. If those were really newly nucleated particles, we would need growth rates of 50 nm/h. I find that hard to believe as such numbers have never (to my knowledge) been reported in the literature for atmospheric nucleation.*

Comments 6, 8 and 9 were addressed together since they all refer to the same topic.

Reply 6 – 8 - 9: Our approach to deal with the growth to CCN size of particles that are not issued from NPF is to subtract the CCN increase observed during non NPF days from the CCN increase observed during NPF days (section 3.1.2.). However, we first used in section 3.1.1 the classical approach found in most papers dealing with the evaluation of CCN production from NPF, in order to be able to compare with existing values reported in the literature from other sites. We do agree that using this first analysis could be confusing if not well explained. We now use a different terminology for section 3.1.1, using the term « CCN production DURING NPF » instead of CCN production FROM NPF. Also, in order to make our approach clearer the methodologies and discussion on the uncertainties on these methodologies are included in the results sections.

If we understand correctly the second part of Comment 6, the reviewer suggests, for each threshold size D , to subtract the concentration of particles below D measured before the event to what we actually calculate as CCN_D . We believe that this methodology does not provide significant progress compared to what we did for two main reasons:

- This would only remove the particles that arrived at the station before the events, assuming that their concentration remains the same during the event. In any case, the contribution of the particles which reach the station during the event and further grow would still not be filtered out.
- This suggestion relies on the strong assumption that all particles below threshold D , regardless their origin, will reach D , which is unrealistic due to coagulation process.

In brief, we believe that applying such a methodology would consist to assume different hypothesis, which might not be more suitable than ours to describe what is actually occurring.

Comment 7: Page 4 Line 28: I found CCN_{high} and CCN_{low} quite confusing. I would rather prefer to use the size of the particles, therefore, I would call them CCN_{50} and CCN_{100} .

Reply 7: Notations were changed accordingly.

Comment 10: Page 5 Line 14: The authors acknowledge only partially the previous point. They say that the particles can be transported during that period but they still don't mention that small particles transported there can then grow to the threshold and being considered as formed by NPF. They also correct the transported particles by comparing NPF days with non-NPF days. This assumption is valid only if the physic dynamic is the same. If the NPF is triggered by the wind convection might be that during nucleation (more wind) the particles transported up there are more. This point needs further investigation or at least being commented.

Reply 10: The possibility for particles formed off-site and transported to the station to contribute to CCN concentrations is now explicitly mentioned (see Reply 6 – 8 - 9).

The revised version of the manuscript now includes additional discussion regarding environmental conditions on event and non-event days, and the effect it can have when estimating transport contribution to CCN concentrations (Section 3.1.2):

“These calculations rely on the hypothesis that the specific environmental conditions on which NPF occurs are not influencing the transport from lower atmospheric layers. In order to further evaluate the reliability of this assumption, wind direction and speed as well as global radiation were investigated on event and non-event days (Figures S3 and S4 in the supplementary material). As previously reported by Rose et al. (2015a), NPF events are favoured during clear sky conditions, when radiation is higher (Fig. S3). Thus, there is likely a bias towards an underestimation of radiative driven transport from lower atmospheric layers due to the fact that cloudy days are over-represented for non-event days. Regarding wind, contrasting directions are also observed between event and non-event days (Fig. S4), with patterns closely related to those observed for the dry and wet seasons, respectively (Rose et al., 2015a). It is worth noticing that winds originating from the more polluted sector of La Paz – El Alto (south) do not seem to be over-represented neither on event nor on non-event days. However, because of the close proximity of this area, it is complex to further assess how it contributes to CCN concentration from wind direction alone, and we cannot exclude a bias related to the variability of this specific source between event and non-event days. Nonetheless, the particle number concentrations observed at the time preceding the usual occurrence of the NPF events are similar for event and non-event days (Fig 3, S1, S2). Moreover, higher wind speeds are on average recorded on non-event days,

that likely lead to an enhanced transport of particles to the site compared to event days, and hence lead to an underestimation of the contribution of NPF to the increase of CCN. In any case, taking into account the contribution of transport when calculating the increase of CCN concentrations after NPF events was never done in the past, and certainly helps approaching a more realistic view of the real contribution of NPF to CCN number concentrations.

Two additional figures are provided in the Supplementary material to show wind speed and direction (Fig. S4) and radiation (Fig. S3).

Comment 11: *Section 2.3 Method to assess the influence of the boundary layer in Chacaltaya. To my understanding, this method only takes into account the local PBL influence at the time of nucleation. It does little to actually describe the air mass in which nucleation occurs. Bianchi et al. (2016) have shown that strong PBL contact 1-2 days before NPF is crucial in the case of the Alps (Jungfraujoch). While conditions are certainly different in the Andes, there is no reason to believe that local wind conditions could accurately describe an air mass and its history which is what one must do to get a handle on PBL influence. There is a good body of literature dealing with the assessment of PBL influence. Much of it has been summarized in recent papers by Bukowiecki et al. (2016) and Herrmann et al. (2015).*

Reply 11: Indeed the methodology that we propose only takes into account the local BL influence at the time of the event. But this is obviously an interesting parameter to take into account since we do observe different event types as a function of the different evolution patterns of the BL (Fig. 6). The fact that only local influence is considered is now clearly stated in the title of Section 2.3 “Method to assess the local influence of the boundary layer in Chacaltaya”, as well as in the introducing lines of the same section “In order to assess whether the site is under the influence of the planetary boundary layer or the low free troposphere at a local scale, regardless the history of the air mass, we employed the hourly-averaged value of the standard deviation of the horizontal wind direction (σ_θ)”.

Also, the valuable additional information which could be provided by an analysis of the air mass history (not performed here due to lack of proxy measurement and modelling tools such as those used at Jungfraujoch) is discussed at the end of Section 3.2.3, in the light of the work by Tröstl et al. (2016) and Bianchi et al. (2016):

“Additional analysis regarding the history of the air mass and BL influence along its trajectory would provide valuable information to even more assess the role of the exchanges between the BL and the FT on the occurrence of NPF and its contribution to the formation of new CCN. Indeed, observations conducted at the Jungfraujoch showed that stronger NPF events (type I) occurred in air masses one or two days after contact with the BL (Bianchi et al., 2016; Tröstl et al., 2016). These results are however based on proxies (CO, NO_y) and modelling tools which were unfortunately not available for Chacaltaya. Nevertheless, our results goes to some extent into the same direction as the work by Tröstl et al. (2016) and Bianchi et al. (2016), at least supporting the major role of BL intrusion (regardless of its kind, before or during the event) to sustain particle growth. Similar FT feeding process from the BL was also shown by Rose et al. (2015b) at the puy de Dôme (France, 1465 m a.s.l.)”

Comment 12: *Page 8 Line 14: “...when particles reached the lowest activation diameter, i.e. 50 nm, they systematically grew up to at least 100 nm...” This statement is stronger than what the data seem to support. We don’t know for certain that those are not pre-existing particles that simply did a bit of growing above the considered threshold, or do we?*

Reply 12: As previously mentioned in reply 6 – 8 - 9, the contribution of pre-existing particles cannot be excluded. Rephrasing and additional explanation should however help in this particular case:

“Over the whole year, 61% of the studied NPF events were apparently growing to CCN-relevant sizes, and when observed, the contribution of growing particles to CCN concentrations was systematically seen up to at least 100 nm. During the wet season, the frequency of aerosol particles reaching CCN sizes during a NPF event was higher compared to the dry season (79 % and 56%, respectively). This last observation can be ascribed to the larger growth rates which were detected during the wet season, being on average enhanced by a factor 1.7 compared to the dry season (Rose et al., 2015a). It is however worth noticing that at this stage, the contribution of pre-existing particles transported to the site at already grown sizes cannot be excluded.”

Comment 13: Page 8 Line 16: “...aerosol particles originating from NPF event and reaching CCN sizes...” Yet again the same problem that I think it should be fixed. I haven’t seen any evidence that all those new CCN come directly from NPF, and, indeed, I find it highly unlikely: as long as there are pre-existing particles their chances to add to the CCN concentration are MUCH higher than the chances of newly formed particles.

Reply 13: Yes, we rephrased: “During the wet season, the frequency of aerosol particles reaching CCN sizes during a NPF event was higher compared to the dry season”.

Comment 14: Page 9 from Line 15 to Line 32: The paragraph lacking a message. First the authors give us comparisons to sites that are hardly comparable to a 5000 m peak, and then they tell us in the last few lines that those comparisons are more or less pointless and I actually i would agree because these sites are just different and the comparison does not provide useful information. A comparison to Tröstl et al., (2016) might be more interesting, especially since those results are quite different.

Reply 14: The studies that are mentioned in the initial version of the manuscript were, at that time, the only ones that focussed on CCN production from NPF using a method that is similar to ours. A comparison with the recent results by Tröstl et al. (2016) is now included:

“The potential of NPF to contribute to CCN production at high altitude was more particularly investigated by Tröstl et al. (2016) at the Jungfraujoch station. Tröstl and co-workers found that newly formed particles did not directly grow to CCN sizes (90 nm at Jungfraujoch) within observable time scale (up to two days) but rather experienced a multi-step growth process over several days. As a consequence, the contribution of NPF to the CCN budget was complex to distinguish from that of other sources such as BL entrainment of larger particles, which was likely the main source of measured CCN.”

Comment 15: Page 10 Line 8: Sunrise is typically a well-defined point in time and not a process that has an onset.

Reply 15: Changed to “that takes place after sunrise”.

Comment 16: Section 3.1.2 Correction for the contribution of particles transported to the site I’m a bit concerned regarding this method to correct the contribution of particles transported. As mentioned earlier, this method is valid only in case every day we have the same physics and nucleation only depends on the vapors present. However, if nucleation is triggered by the wind coming up the valley than during nucleation we would have more transportation of big particles and therefore the correction method is not ideal. Would be nice to know what are the differences (Wind direction, wind speed etc..) during nucleation and during no nucleation where these background values is taken in account.

Reply 16: This point is now discussed in Section 3.1.2, in the light of supplementary Figure S4.:

“Regarding wind, contrasting directions are also observed between event and non-event days (Fig. S4), with patterns closely related to those observed for the dry and wet seasons, respectively (Rose et al.,

2015a). It is worth noticing that winds originating from the more polluted sector of La Paz – El Alto (south) do not seem to be over-represented neither on event nor on non-event days. However, because of the close proximity of this area, it is complex to further assess how it contributes to CCN concentration from wind direction alone, and we cannot exclude a bias related to the variability of this specific source between event and non-event days. Nonetheless, the particle number concentrations observed at the time preceding the usual occurrence of the NPF events are similar for event and non-event days (Fig 3, S1, S2). Moreover, higher wind speeds are on average recorded on non-event days, that likely lead to an enhanced transport of particles to the site compared to event days, and hence lead to an underestimation of the contribution of NPF to the increase of CCN.”

Comment 17: Section 3.2 How layering influences growth to CCN-sizes I do understand the need of knowing where the nucleation events take place and especially if this lead to a big production of particles in the free troposphere. However, I believe that dividing in 10 scenarios is a bit over exaggerated and probably not quite realistic. I think it would be better if the authors can simplify this section. I don't think that selecting more than 3 scenarios is feasible. In addition to that the split into different scenarios seems illconceived since most scenarios are quite irrelevant with very little occurrences. This might all be a nice exercise in data analysis but the text fails to tell us what the actual results are. What do we learn in this section apart from some minuscule details? This section has the feel of filler material and needs to be improved with a fair amount of actual substance.

Reply 17: Sections 3.2.1 and 3.2.2 were simplified, with a reduced number of scenarios (3) focussed on BL and FT tropospheric conditions, which are the most frequent.

Comment 18: Page 12 Line 4: “...regarding the location of the station in the tropospheric layers...” I wonder if this is actually relevant at all? The NPF events are mainly driven by the air mass history and not so much by the atmospheric layer when the event begins.

Reply 18: If air mass history has an influence on the nucleation process, based on Fig. 6 it seems that local atmospheric layering at the station during NPF does also have an effect, since different event types are observed as a function of the different evolution pattern of the BL. For example, we clearly observe that for events triggered in the FT, the probability for type I events to occur is increased when initial free tropospheric conditions are changing in the course of the events. Additional discussion on air mass history is now provided at the end of Section 3.2.3.

Comment 19: Page 12 Line 8: “389 NPF events” Is that a different data set?

Reply 19: The events included in the CCN production investigation are only those belonging to class I, as stated P8, L10: “147 days showing type I NPF events”. In contrast, the analysis regarding the influence of atmospheric layering on the occurrence of NPF (Sections 3.2.1 and 3.2.2) includes all the events previously discussed by (Rose et al., 2015b), not only type I events. It is now clearly stated: “389 NPF events (including all event types, i.e. I, II or bump, Hirsikko et al., 2007) previously discussed by (Rose et al., 2015a)...”

Minor edits:

Figures: In general no need to state Chacaltaya at the end of every captions

Ok, removed.

Figure 1: Why Tinit is not before the nucleation but already a after the start of the event? Please also describe the figure, Particle size distribution measured by..... and so on.

P5, L9: t_{init} is defined as the time “ when nucleated particles reach the threshold size”, or let’s say growing particles, since their origin might be uncertain. In other words, t_{init} is the time when the “banana” reach the threshold size, further leading to the CCN concentration increase, as shown on Fig. 1.

Additional information is provided: “Determination of the CCN concentration increase for the 3 threshold diameters (50, 80 and 100 nm) from the particle size distribution measured by SMPS” .

We thank referee N°2 for his comments and remarks which contributed to improve and clarify the present paper. Our answers to the suggestions are listed below.

SPECIFIC COMMENTS

Comment 1: How do the authors separate CCN formation from freshly nucleated particles and CCN formation from the growth of pre-existing particles during nucleation events? Here, I refer to the very detailed comments of Anonymous Referee #1. There is not much to add.

Reply 1: We agree with the fact that the manuscript needed for more clarity regarding this aspect. We now use a different terminology for section 3.1.1, using the term « CCN production DURING NPF » instead of CCN production FROM NPF. Also, in order to make our approach clearer the methodologies and discussion on the uncertainties on these methodologies are included in the results sections. In addition, several sentences were rephrased throughout the manuscript in order to further avoid any misunderstanding.

Comment 2: How robust is the treatment of advection of different air masses by the selected approach? Are there other observables (e.g., trace gases) available which allow a more robust treatment of particle transport than the simple method deployed in the study? The authors use the hypothesis that similar particle number concentrations are transported to the site during days with and without particle formation events. They state in Section 3.1.2 that at hours outside of NPF events, particle number concentrations were on average similar for event and non-event days. However, what is the variability of the particle background and does it depend on the wind direction where the air masses came from, etc.? In particular the variability of the particle background needs to be presented more quantitatively since this parameter determines the level of uncertainty of the reported CCN increases by NPF. Concerning the structure, the presentation of results in Section 3.1 is confusing. The authors start with a detailed description of CCN production and list all obtained numbers in detail and show the min Fig. 2. Then in Section 3.1.2 they introduce a correction of the presented CCN number concentrations. It is confusing that the CCN production neglecting the influence of advection is shown in Fig. 2 while the more important CCN production from NPF only is not shown but only listed in Table 2. If I understood right, the authors focus on CCN from NPF. If this is true then the way of presenting the data in Section 3.1 should be revised. One possibility is to start with a quantitative analysis of the “particle background” during non-event days, including its variability, introduce then the method for determining CCN production and present finally the CCN production values corrected for particle transport.

Reply 2: We know that the correction that we applied for particle transport is not completely accurate, since it relies on a strong hypothesis which is explicitly mentioned in the text (“similar number concentrations of particles are transported to the site on event and non-event days”), and which might actually not be verified because of some reasons which are now mentioned with more details in the revised version of the manuscript (Section 3.1.2):

“These calculations rely on the hypothesis that the specific environmental conditions on which NPF occurs are not influencing the transport from lower atmospheric layers. In order to further evaluate the reliability of this assumption, wind direction and speed as well as global radiation were investigated on event and non-event days (Figures S3 and S4 in the supplementary material). As previously reported by Rose et al. (2015a), NPF events are favoured during clear sky conditions, when radiation is higher (Fig. S3). Thus, there is likely a bias towards an underestimation of radiative driven transport from lower atmospheric layers due to the fact that cloudy days are over-represented for non-event days. Regarding wind, contrasting directions are also observed between event and non-event days (Fig. S4), with patterns closely related to those observed for the dry and wet seasons, respectively (Rose et al., 2015a). It is worth noticing that winds originating from the more polluted sector of La Paz – El Alto

(south) do not seem to be over-represented neither on event nor on non-event days. However, because of the close proximity of this area, it is complex to further assess how it contributes to CCN concentration from wind direction alone, and we cannot exclude a bias related to the variability of this specific source between event and non-event days. Nonetheless, the particle number concentrations observed at the time preceding the usual occurrence of the NPF events are similar for event and non-event days (Fig 3, S1, S2). Moreover, higher wind speeds are on average recorded on non-event days, that likely lead to an enhanced transport of particles to the site compared to event days, and hence lead to an underestimation of the contribution of NPF to the increase of CCN. In any case, taking into account the contribution of transport when calculating the increase of CCN concentrations after NPF events was never done in the past, and certainly helps approaching a more realistic view of the real contribution of NPF to CCN number concentrations.”

Additional figures are also provided in the supplementary, showing wind speed and direction (Fig. S4) and radiation (Fig. S3)

However, the aim of this correction is not to provide an exact estimation of transport contribution, which, we believe, cannot be retrieved from these measurements, even if including additional parameters, such as trace gases concentrations. Our objective, less ambitious, is rather to go one step further compared with previous analysis published in the literature in order to estimate NPF contributions to CCN number concentrations which are closer to actual values, as mentioned in the text.

We agree that the terminology to describe sections 3.1.1 and 3.1.2. was confusing. We now use the term « CCN formation during NPF » for section 3.1.1 and CCN formation from NPF for section 3.1.2.

We agree that the most important results should be better emphasised for section 3.1.2. For that purpose, an additional Figure (4) is now provided and the text has been slightly changed:

“The contributions of NPF particles to the increase of CCN, all shown on Fig. 4.a. and reported in Table 2 for the different seasons and sizes, hence represent a significant fraction of the CCN increase shown on Fig. 2.a. and reported in Table 1. The contribution of NPF to CCN concentrations are comparable or even higher than those previously mentioned for other stations in the literature, which probably also include CCN sources other than NPF. The relative impact of NPF are estimated to increase the CCN₅₀ number concentrations by more than 250 % during both seasons, and the CCN₁₀₀ number concentrations by more than 100% and 200% during the dry season and wet season, respectively.”

***Comment 3:** How robust is the separation between air masses from the boundary layer and from the free troposphere, and what is the expected impact of air mass history on the occurrence of new particle formation events? This question refers to Section 3.2 which in its current form is difficult to understand. The attempt of the authors is quite understandably to study if NPF events occur preferably in air masses originating from the BL or from the FT. However, doing this requires a clear presentation of event types and characteristics before going into details. Here the authors should restructure Section 3.2, start with a clear presentation of event types and scenarios. One table including all considered cases (with more detail than stated in Table 3) etc. might help. Looking at Fig. 5, there is no big difference between the scenarios, except for S1, S6 and likely S7. The authors may rethink the choice of scenarios in order to get a more precise conclusion from this part of the study. In addition, the expected impact of air mass history should be investigated / discussed.*

Reply 3:

The method used to distinguish between boundary layer and free troposphere air masses performed in the present study is the only one we could apply given the measurements conducted in Chacaltaya.

The impact of air mass history, including occurrence and length of BL contact, has not been investigated due to lack of proxy measurement and modelling tools (such as those used by Tröstl et al. (2016)). However, the fact that air mass history may, in addition to local conditions, influence the occurrence of NPF is now mentioned at the end of Section 3.2.3 in the light of the work by Tröstl et al. (2016) and Bianchi et al. (2016):

“Additional analysis regarding the history of the air mass and BL influence along its trajectory would provide valuable information to even more assess the role of the exchanges between the BL and the FT on the occurrence of NPF and its contribution to the formation of new CCN. Indeed, observations conducted at the Jungfraujoch showed that stronger NPF events (type I) occurred in air masses one or two days after contact with the BL (Bianchi et al., 2016; Tröstl et al., 2016). These results are however based on proxies (CO, NO_y) and modelling tools which were unfortunately not available for Chacaltaya. Nevertheless, our results goes to some extent into the same direction as the work by Tröstl et al. (2016) and Bianchi et al. (2016), at least supporting the major role of BL intrusion (regardless of its kind, before or during the event) to sustain particle growth. Similar FT feeding process from the BL was also shown by Rose et al. (2015b) at the puy de Dôme (France, 1465 m a.s.l.).”

We agree with the fact that the number of scenarios used to assess the impact of BL on the occurrence of NPF was maybe too high. It was thus reduced in order to clarify the message:

” The most frequent scenarios, which include more than 88% of the documented events, are listed, together with their frequency of occurrence, in Table 3. Scenario S1 refers to those days when the first steps of the NPF process were observed to occur in the BL, while scenario S2 refer to the events started in the FT. Scenario S2 is further divided into two sub-classes to distinguish between the events which first steps occur exclusively in the FT (S2.1) from those during which BL dynamics lead to changing conditions in the course of the event (S2.2). Events triggered in the IL are not frequently observed compared to those initiated in the BL or in the FT, and are thus not highlighted in this classification.” (Section 3.2.1)

The description of the scenarios in Table 3 was also detailed.

MINOR COMMENTS

Comment 1: Since the classification of NPF events is crucial for understanding the manuscript, a brief description of types should be given at the end of section 2.2, instead of referring to the references Hirsikko et al. (2007) and Rose et al. (2015).

Reply 1: As suggested, brief description of the event types is now provided in Section 3.1.1:

“First, only those NPF events referred as type I, i.e. with clear particle growth from smallest sizes, were considered; they contrast with type II events, during which the growth is more irregular and may be interrupted in certain size ranges, and bump type events, which completely miss the growth of the newly formed clusters (Hirsikko et al., 2007; Yli-Juuti et al., 2009).”

Comment 2: Please add brief descriptions of quantities J and GR to x axis of Figure 6.

Reply 2: Figure caption was slightly modified:

“Median formation rate of 2 nm particles (J_2) and growth rate in the range 1-3 nm (GR_{1-3}) reported separately for type I events initiated in the BL (scenario S1) and in the FT (scenario S2). Lower and upper limits of the error bars stand for the 1st and 3rd quartile, respectively.”

Additional information is also provided in the text, before the description of Fig. 7:

“In order to further characterize the NPF events in the different atmospheric layers, statistics regarding the formation rate of 2 nm particle and the growth rate (GR) in the size range 1-3 nm as a function of the scenarios were performed for type I events. Growth rates were derived from the particle number size distribution using the “maximum” method from Hirsikko et al. (2005), while formation rates were calculated according to Kulmala et al. (2007).”

1 **New Particle Formation and impact on CCN concentrations**
2 **in the boundary layer and free troposphere at the high**
3 **altitude station of Chacaltaya (5240 m a.s.l.), Bolivia**

4 C. Rose¹, K. Sellegri¹, I. Moreno², F. Velarde², M. Ramonet³, K. Weinhold⁴, R. Krejci⁵, M.
5 Andrade², A. Wiedensohler⁴, P. Ginot^{6,7} and P. Laaj⁶

6 ¹ Laboratoire de Météorologie Physique CNRS UMR 6016, Observatoire de Physique du
7 Globe de Clermont-Ferrand, Université Blaise Pascal, 24 avenue des Landais, 63171 Aubière,
8 France

9 ² Universidad Mayor de San Andres, LFA-IIF-UMSA, Laboratory for Atmospheric Physics,
10 Campus Universitario Cota Cota calle 27, Edificio FCPN piso 3, Casilla 4680, La Paz,
11 Bolivia

12 ³ Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-
13 UVSQ, Université Paris-Saclay, F-91191, Gif-sur-Yvette, France

14 ⁴ Leibniz Institute for Tropospheric Research, Permoserstr. 15, 04318 Leipzig, Germany

15 ⁵ Department Environmental Science and Analytical Chemistry (ACES), Atmospheric
16 Science Unit, Stockholm University, S 10691 Stockholm, Sweden

17 ⁶ Laboratoire de Glaciologie et Géophysique de l'Environnement, UMR 5183, UGA/CNRS,
18 Grenoble, France

19 ⁷ Observatoire des sciences de l'Univers de Grenoble, UMS 222, IRD/UGA/CNRS, Grenoble,
20 France

21

22 **Abstract**

23 Global models predict that new particle formation (NPF) is, in some environments,
24 responsible for a substantial fraction of the total atmospheric particle number concentration
25 and subsequently contribute significantly to cloud condensation nuclei (CCN) concentrations.
26 NPF events were frequently observed at the highest atmospheric observatory in the world,
27 Chacaltaya (5240 m a.s.l.), Bolivia. The present study focuses on the impact of NPF on CCN
28 population. Neutral cluster and Air Ion Spectrometer and mobility particle size spectrometer
29 measurements were simultaneously used to follow the growth of particles from cluster sizes
30 down to ~2 nm up to CCN threshold sizes set to 50, 80 and 100 nm. Using measurements
31 performed between January 1 and December 31 2012, we found that 61% of the 94 analysed
32 events showed a clear particle growth and significant enhancement of the CCN-relevant
33 particle number concentration. We evaluated the contribution of NPF relative to the transport
34 and growth of pre-existing particles to the CCN-size. The averaged production of 50 nm

1 particles during those events was 5072 cm^{-3} , and 1481 cm^{-3} for 100 nm particles, with a larger
2 contribution of NPF compared to transport, especially during the wet season. The data set was
3 further segregated into boundary layer (BL) and free troposphere (FT) conditions at the site.
4 The NPF frequency of occurrence was higher in the BL (48%) compared to the FT (39%).
5 Particle condensational growth was more frequently observed for events initiated in the FT,
6 but on average faster for those initiated in the BL, when the amount of condensable species
7 was most probably larger. As a result, the potential to form new CCN was higher for events
8 initiated in the BL (67% against 53% in the FT). In contrast, higher CCN number
9 concentration increases were found when the NPF process initially occurred in the FT, under
10 less polluted conditions. This work highlights the competition between particle growth and
11 the removal of freshly nucleated particles by coagulation processes. The results support model
12 predictions which suggest that NPF is an effective source of CCN in some environments, and
13 thus may influence regional climate through cloud related radiative processes.

14 **1 Introduction**

15 Atmospheric aerosol particles are known to affect air quality, health (Seaton et al., 1995) and
16 climate. Beside their direct interaction with the solar and telluric radiations, aerosol particles
17 also act as condensation nuclei for cloud droplets. Cloud effects such as cloud albedo
18 (Twomey, 1977) and lifetime (Albrecht, 1989) constitute the largest uncertainty in the
19 estimation of the radiative forcing of the Earth's atmosphere (IPCC, 2013).

20 The interaction between aerosol particles and the formation of warm clouds relies on the
21 ability of the particles to serve as cloud condensation nuclei (CCN), which depends on the
22 water vapour supersaturation, particle size distribution and also the chemical composition
23 (e.g.: Roberts et al., 2010; Wex et al., 2010; Asmi et al., 2012). Besides the processing of
24 primary particles, other CCN sources were identified, such as regional new particle formation
25 (NPF) events (Kerminen et al., 2012).

26 NPF is a frequent atmospheric phenomenon including the formation of nanometer-sized
27 clusters from gaseous precursors and their subsequent growth to larger sizes (eg. Kulmala and
28 Kerminen, 2008). Typical growth rates between 1.8 and 10.7 nm h^{-1} were found for particles
29 in the range $1.5 - 20 \text{ nm}$ (Yli-Juuti et al., 2011), meaning that a few hours to a few days are
30 needed for nucleated particles to grow to CCN sizes, around $50-150 \text{ nm}$ (Kerminen et al.,
31 2012). The chance for these clusters to grow to CCN sizes strongly depends on the

1 competition between condensational growth and their removal by coagulation onto pre-
2 existing particles.

3 During the last few years, several global model investigations were dedicated to the study of
4 the CCN-size aerosol production attributed to atmospheric NPF (Makkonen et al., 2012;
5 Merikanto et al., 2009; Reddington et al., 2011; Spracklen et al., 2008). While the outcomes
6 of these different models may vary according to the way they treat NPF and aerosol particle
7 processes (Lee et al., 2013), most of them show an enhancement of the CCN number
8 concentration due to NPF, both in the boundary layer (BL) and in the free troposphere (FT).
9 Based on the study by Makkonen et al. (2012), predictions of the present day annual global
10 average CCN concentration in the BL show almost a fivefold increase when taking into
11 account NPF. According to Merikanto et al. (2009), 45% of global low-level cloud CCN at
12 0.2% supersaturation originate from nucleation, and 35% have been formed in the free and
13 upper troposphere. Slightly contrasting results are provided by Reddington et al. (2011) using
14 the global model GLOMAP against measurements conducted at 15 European ground based
15 stations in the frame of the EUCAARI project. Reddington and co-workers found that CCN-
16 sized particle concentrations in the BL were mainly driven by processes other than NPF,
17 which contributed significantly to the CCN budget at little less than a quarter of observational
18 sites included in the study.

19 However, observations to validate these predictions are scarce, especially for the FT, where
20 measurements are often technically challenging. Recent studies conducted at the Jungfraujoch
21 station (Switzerland, 3580 m a.s.l.) reported significant enhancement of the particle
22 concentration below 50 nm by NPF in the FT, while only a minor fraction of this particles
23 grow beyond 90 nm, even on a time scale of several days (Herrmann et al., 2015; Tröstl et al.,
24 2016). The contribution of NPF to the production of CCN is thus likely to be very limited in
25 this part of the FT, while boundary layer originating particles were observed to dominate the
26 CCN concentrations measured at Jungfraujoch. The occurrence of the NPF process itself in
27 the FT was reported to be tightly connected with the strength of boundary layer influence at
28 the site, together with global radiation (Bianchi et al., 2016; Tröstl et al., 2016).

29 In this context, the purpose of the present study is to estimate the contribution of NPF to CCN
30 formation at the station of Chacaltaya (5240 m a.s.l., Bolivia) with a special attention in
31 differentiating the CCN number concentrations attributed to NPF and particle growth
32 occurring at the station from those attributed to the transport of pre-existing CCN-size

1 particles to the site. This analysis was performed using an indirect method based on the NPF
2 event classification previously reported by Rose et al., (2015a) and particle number size
3 distribution measurements in the range 10-500 nm. In addition to global CCN number
4 concentrations, a more detailed analysis of NPF and subsequent CCN production in the BL or
5 in the FT is also reported.

6 **2 Measurements and methods**

7 **2.1 Observation site and instruments**

8 Aerosol particle number size distributions, together with routine meteorological parameters,
9 were measured at the Chacaltaya GAW station, located in a range of the Bolivian Andes at the
10 summit of Mount Chacaltaya (16°21.014' S, 68°07.886' W), 15 km North of La Paz – El Alto
11 metropolitan area (2 million inhabitants).

12 The mobility distribution of charged particles and ions ($3.2 - 0.0013 \text{ cm}^2\text{V}^{-1}\text{s}^{-1}$) and the size
13 distribution of total particles (2 – 42 nm) were measured by a Neutral cluster and Air Ion
14 Spectrometer (NAIS, Airel Ltd., Mirme and Mirme, 2013). The NAIS sampled the ambient
15 aerosol through an individual non-heated short inlet (~ 50 cm) with a 5 minute time
16 resolution. Since the NAIS was likely to overestimate particle number concentrations above
17 20 nm (Manninen et al., 2016), particles in the range from 20 nm to CCN relevant sizes were
18 preferentially measured using a mobility particle size spectrometer type TROPOS-SMPS
19 (Wiedensohler et al., 2012). The SMPS operated behind a Whole Air Inlet equipped with an
20 automatic dryer.

21 More details on the measurement site as well as the instrumental setup and the data quality
22 assurance can be found in Rose et al. (2015a) and Andrade et al. (2015).

23 **Section 2.2 moved to Sect. 3.1.1 and 3.1.2**

24 **2.2 Method to assess the local influence of the boundary layer in Chacaltaya**

25 In order to assess whether the site is under the influence of the planetary boundary layer or the
26 low free troposphere at a local scale, regardless the history of the air mass, we employed the
27 hourly-averaged value of the standard deviation of the horizontal wind direction (σ_θ).

28 The value of σ_θ has been extensively used in air pollution monitoring (EPA, 2008; Mitchell,
29 1982; Mitchell and Timbre, 1979; Weber, 1997) and dispersion models as an indicator of the

1 stability of the lower atmosphere. Instable atmospheric conditions produce turbulence and
 2 therefore high wind variability. Conversely, low wind variability due to stable conditions
 3 produces low σ_θ values. In Chacaltaya, σ_θ was used from a mountain perspective, .i.e.
 4 assuming that turbulent conditions ($\sigma_\theta \geq 12.5$) reflect the influence of the BL at the observatory
 5 and, contrarily, that non-turbulent (or stable) conditions are equivalent of being in the FT (σ_θ
 6 < 12.5).

7 In Chacaltaya, σ_θ is obtained at the summit (5380 m a.s.l., 10 m above the surface) by means
 8 of a wind vane and propeller (Young 05103) and processed directly on a CS-CR1000
 9 datalogger. Sigma theta is defined as the standard deviation of the horizontal wind direction
 10 itself according to Eq. (1), but its value is approximated by the Yamartino (1984) single-pass
 11 method (set of Eq. (2)) directly in the datalogger.

$$12 \quad \sigma_\theta = \left[\frac{\sum_{i=1}^N (\theta_i - \theta_A)^2}{N-1} \right]^{\frac{1}{2}} \quad (1)$$

13 where θ_i is the instantaneous wind direction and θ_A the average wind direction.

$$14 \quad \sigma_\theta = \arcsin(\varepsilon) \left[1 + \left(\frac{2}{\sqrt{3}} - 1 \right) \varepsilon^3 \right]$$

$$\varepsilon \equiv \sqrt{1 - (S^2 + C^2)}$$

$$S = \frac{1}{N} \sum_{i=1}^N \sin \theta_i$$

$$C = \frac{1}{N} \sum_{i=1}^N \cos \theta_i \quad (2)$$

15 The synoptically driven change of wind direction may affect the calculation of σ_θ for short
 16 time periods. This low-frequency horizontal wind oscillation is called “meandering” and may
 17 produce overestimation of σ_θ during situations of low wind speed ($\leq 2 \text{m.s}^{-1}$), which usually
 18 take place during daytime in Chacaltaya. Therefore, 15-min averaged values are calculated
 19 offline according to Eq. (3) to avoid wind meandering effects.

$$20 \quad \sigma_{\theta(1-hr)}^2 = \frac{\sigma_{\theta(15)}^2 + \sigma_{\theta(30)}^2 + \sigma_{\theta(45)}^2 + \sigma_{\theta(60)}^2}{4} \quad (3)$$

21 where every $\sigma_{\theta(15x)}$ equation is a 15-minute deviation of the wind direction.

1 The threshold set for stable FT conditions is $\sigma_{\theta} \geq 12.5$, following Mitchell's recommendations
2 (1982). In Chacaltaya, FT conditions take place usually during night-time and before sunrise,
3 as it would be expected for mountain sites. Nevertheless, in many cases σ_{θ} values lower than
4 18 are observed in a persistent pattern (more than 4 hours of this condition). This may
5 indicate the existence of a residual or interface layer (IL). This intermediate layer would not
6 correspond neither to the FT nor the proper BL. Moreover, during the wet season, convective
7 and unstable conditions produce more turbulence at the site, shifting the σ_{θ} towards higher
8 values, typically below 18. Therefore other secondary site specific thresholds are applied,
9 namely 18 and 22.5.

10 Obtained hourly dataset is then checked for consistency, in particular with black carbon
11 measurements, and the following smoothing is applied. We establish a 4-hour window (2h
12 before and 2h after the data point of interest) into which the following criteria are applied:

- 13 • If the σ_{θ} value is lower than 12.5 (classified as FT), but if it is the only data point in
14 the 4-hour window, it is not considered as FT and it is reclassified as an IL point
15 instead.
- 16 • If the σ_{θ} value is lower than 18 and 75% of the points in the 4-hour window are lower
17 than 12.5, the point is classified as a FT point (stable).
- 18 • If the σ_{θ} value is lower than 22.5 and 75% of the points in the 4-hour window are
19 lower than 18, the point is classified as an IL point (this takes place mostly during the
20 wet season).

21

22 **3 Results**

23 **3.1 CCN formation during and from NPF**

24 **3.1.1 CCN formation during NPF events (part of Sect. 2.2 moved here)**

25 In absence of direct CCN measurements at Chacaltaya, the contribution of NPF to CCN
26 production was estimated from the continuous monitoring of the particle number size
27 distribution. This indirect method was first introduced by Lihavainen et al. (2003) and has
28 already been used in several other studies (Asmi et al., 2011; Kerminen et al., 2012; Laakso et
29 al., 2013; Laaksonen et al., 2005).

30 The basic hypothesis is that the lower cloud droplet activation diameter of aerosol particles is
31 in the range 50-150 nm for the usual supersaturations encountered in natural clouds (Asmi et
32 al., 2011, 2012; Komppula et al., 2005) including those forming at altitudes up to 3580 m

1 a.s.l., as observed at the Jungfraujoch station (Switzerland) (Hammer et al., 2014; Jurányi et
2 al., 2011). Although these conditions might be slightly different from those found in clouds
3 forming above 5000 m, we assume that on a first approach the CCN sizes previously
4 mentioned apply the same way at such altitudes. Thus, CCN number concentrations are
5 assimilated to a range of three different CN concentrations: hereafter, CCN_{50} and CCN_{100}
6 refer to the higher and lower limits of the CCN concentration estimated from the number
7 concentrations of particles larger than 50 nm and 100 nm, respectively; as additional
8 information, an intermediate CCN concentration (CCN_{80}) was deduced from the number
9 concentration of particles larger than 80 nm. The CCN production during an event was
10 obtained from the comparison of the CCN concentration N_{init} prior to and the maximum CCN
11 concentration N_{max} during the event. For each particle diameter range, N_{init} is defined as the
12 30 minute average concentration obtained at t_{init} , when growing particles reach the threshold
13 size, whereas N_{max} is the 30 minute average concentration calculated when the CCN
14 concentration reaches a maximum during an event, at t_{max} . The determination of N_{init} and N_{max}
15 is depicted on Fig. 1. It is worth noticing that this indirect method based on particle size only
16 provides estimations of potential CCN concentrations instead of real concentrations as
17 measured by CCN chambers (Roberts and Nenes, 2005). However, for simplicity, we refer to
18 these potential CCN as CCN hereafter.

19 The selection of the NPF events to be analyzed was performed based on the following criteria
20 First, only those NPF events referred as type I, i.e. with clear particle growth from smallest
21 sizes, were considered; they contrast with type II events, during which the growth is more
22 irregular and may be interrupted in certain size ranges, and bump type events, which
23 completely miss the growth of the newly formed clusters (Hirsikko et al., 2007; Yli-Juuti et
24 al., 2009). Second, the days showing an eventual contribution from NPF events triggered the
25 day before were rejected. Especially, those days when the NPF contribution superimpose on
26 that from a strong growing pre-existing Aitken mode band (of similar or even larger intensity
27 in terms of particle concentration), as previously described by Tröstl et al. (2016), were
28 removed from the analysis. Regarding this aspect, our analysis is thus a lower limit of the
29 contribution of NPF to CCN-size relevant aerosol concentrations.

30 During the measurement period January 1 to December 31 2012, 147 days showing type I
31 NPF events were detected: 112 during the dry season, from May to October, and 35 during
32 the wet season, from November to April (Rose et al., 2015a). However, because of missing

1 data of particle number size distribution measurements, only 94 of them were further analysed
2 (75 from the dry season and 19 from the wet season).

3 Over the whole year, 61% of the studied NPF events were apparently growing to CCN-
4 relevant sizes, and when observed, the contribution of growing particles to CCN
5 concentrations was systematically seen up to at least 100 nm. During the wet season, the
6 frequency of aerosol particles reaching CCN sizes during a NPF event was higher compared
7 to the dry season (79 % and 56%, respectively). This last observation can be ascribed to the
8 larger growth rates which were detected during the wet season, being on average enhanced by
9 a factor 1.7 compared to the dry season (Rose et al., 2015a). It is however worth noticing that
10 at this stage, the contribution of pre-existing particles transported to the site at already grown
11 sizes cannot be excluded.

12 Our results of CCN concentration increase during NPF events can be compared to literature
13 values obtained using similar methodologies for other sites. The results reported by Asmi et
14 al. (2011) for Pallas (560 m a.s.l., Finland) slightly contrast with these observations. Indeed,
15 the CCN number concentration increase during NPF events showed a seasonal variation but
16 also decreased with increasing activation diameter. This might be explained by a decreasing
17 availability of condensing vapours over the course of the particle growth time period. At
18 Chacaltaya, the availability of condensing gases appears to increase over a large time period,
19 sometimes reaching concentrations that trigger a second (and third) nucleation event during
20 the same day, in spite of the raising condensable sink due to the first nucleation event (Rose et
21 al., 2015a). Coagulation processes however lead to a decrease of CCN_{100} compared to CCN_{50} .
22 This is illustrated on Figure 2.a, which shows, for the three threshold sizes and for each
23 season, the median CCN concentration increase observed during NPF events and calculated
24 as the difference between N_{max} and N_{init} . Considering all type I event days over the whole
25 year, the median number concentration of new CCN produced during a NPF event was 5072
26 cm^{-3} for CCN_{50} , 2254 and 1481 cm^{-3} for CCN_{80} and CCN_{100} , respectively. The number
27 concentration of new CCN was on average higher during the dry season, especially for
28 CCN_{50} .

29 Corresponding relative increases in CCN number concentration were calculated as the ratio of
30 the absolute increases previously reported over N_{init} , i.e. the 30 min average CCN number
31 concentration measured when growing particles initially reach the threshold sizes (Fig. 2.b).

1 CCN concentrations were found to increase by 168 to 996% at Chacaltaya during NPF events,
2 with no clear differences between seasons or threshold sizes.

3 One should note that when several consecutive type I events were detected on a same day
4 (this occurred on 7 occasions), it was complex to extract the contribution of each individual
5 event, so the calculated CCN production was the result of the contribution of all events as a
6 whole. During multiple events days, the median number concentration of CCN produced was
7 on average 1.7 times higher compared to single type I event days.

8 As previously mentioned, similar methodologies were used in previous studies to evaluate the
9 increase of CCN concentrations during NPF events. The average absolute CCN production
10 observed during NPF events at Chacaltaya is lower compared to that reported by Laaksonen
11 et al., (2005) at the station of San Pietro Capofiume located in the polluted region of the Pô
12 valley (11 m a.s.l., Italy): on the basis of 304 NPF events, the average number of new CCN
13 produced during an event are $7.3 \times 10^3 \text{ cm}^{-3}$ and $2.4 \times 10^3 \text{ cm}^{-3}$, for CCN₅₀ and CCN₁₀₀,
14 respectively. In contrast, the values from both Chacaltaya and San Pietro Capofiume are
15 significantly higher than those reported by Kerminen et al. (2012) for the stations of
16 Botsalano (1420 m a.s.l., South Africa), Vavihill (172 m a.s.l., Sweden), Pallas and Hyytiälä
17 (182 m a.s.l., Finland). Among these four sites, the highest CCN concentration increases are
18 on average observed at Botsalano (2500 cm^{-3} , 1400 cm^{-3} and 800 cm^{-3} for CCN₅₀, CCN₈₀ and
19 CCN₁₀₀, respectively), whereas Pallas displays the lowest CCN production (1000 cm^{-3} , 250
20 cm^{-3} and 150 cm^{-3} for CCN₅₀, CCN₈₀ and CCN₁₀₀, respectively). Corresponding relative
21 increases in CCN concentrations found in the literature are always larger than 100% but never
22 exceed 400%, being thus on average significantly lower than those observed at Chacaltaya.
23 However, it is worth noticing that these contrasting results may arise from the various
24 conditions that are found at the different stations, especially regarding altitude and pollution
25 levels, thus influencing NPF both in terms of strength, spatial extend and temporal evolution.

26 The potential of NPF to contribute to CCN production at high altitude was more particularly
27 investigated by Tröstl et al. (2016) at the Jungfrauoch station. Tröstl and co-workers found
28 that newly formed particles did not directly grow to CCN sizes (90 nm at Jungfrauoch)
29 within observable time scale (up to two days) but rather experienced a multi-step growth
30 process over several days. As a consequence, the contribution of NPF to the CCN budget was
31 complex to distinguish from that of other sources such as BL entrainment of larger particles,
32 which was likely the main source of measured CCN. At Mount Whistler (2182 m a.s.l.,

1 Canada), Pierce et al. (2012) followed a different approach including calculations of the
2 probability for freshly nucleated particles to reach CCN relevant sizes. Based on a five event
3 day period, they found that in absence of high coagulation/condensation sinks, up to 24% of
4 the newly formed clusters could grow to at least 100 nm, thus forming potential CCN.

5 As previously mentioned, the vertical transport of aerosol particles from lower atmospheric
6 levels that takes place after sunrise concurrently to NPF may represent a significant
7 contribution to the increase of CCN-relevant size particle number concentrations at these
8 mountain sites. This aspect will be addressed in the next section, in which the contribution of
9 NPF is further compared with the CCN number concentration increase resulting from the
10 transport of particles to the site.

11 The seasonal and annual CCN productions related to NPF events were estimated from 1) the
12 average fraction of type I NPF events contributing to the formation of new CCN reported
13 above, 2) the frequency of occurrence of type I NPF events at the site and 3) the average CCN
14 number concentration increase measured for those type I events during which growing
15 particles reached the potential CCN activation diameter. As an example, the CCN_{50}
16 production during the wet season was calculated as follows:

$$17 \quad CCN_{50-wet} = frac_{wet} \times tot_nb_{wet} \times avg_conc_{wet} = 79\% \times 35 \times 3070 = 8.48 \times 10^4 \text{ cm}^{-3} \quad (4)$$

18 where, for each season, *frac* is the fraction of NPF events leading to CCN concentration
19 increase, *tot_nb* is the total number of days showing type I events and *avg_conc* is the
20 median number of new CCN formed during an event. Similar calculations were done for each
21 season and CCN class, leading to the values reported in Table 1. The annual CCN production
22 was calculated as the sum of the seasonal productions.

23 Based on Table 1, the CCN production at Chacaltaya was higher during the dry season
24 compared to the wet season for all CCN classes, but especially for CCN_{50} , which was more
25 than 4 times higher compared to the wet season. The annual CCN production calculated at
26 San Pietro Capofiume is $3.4 \times 10^5 \text{ cm}^{-3}$ and $1.1 \times 10^5 \text{ cm}^{-3}$, for CCN_{50} and CCN_{100} , respectively
27 (Laaksonen et al., 2005). These values are slightly lower than those obtained at Chacaltaya,
28 despite the fact that the median number of potential new CCN formed during an event is on
29 average higher in San Pietro Capofiume. This last observation can be ascribed to the high
30 NPF frequency at Chacaltaya, together with the significant fraction of type I events and high
31 growth rates (Rose et al., 2015a).

1 3.1.2 CCN formation from NPF events (part of Sect. 2.2 moved here)

2 The aim of this section is to evaluate the contribution of particles transported to the site to the
3 total CCN concentration and give an estimation of the CCN production from NPF alone.

4 In fact, in addition to the previous analysis classically used in the literature, further
5 calculations are needed to take into consideration the geographical specificity of the site.
6 Indeed, if NPF contributes to the formation of potential new CCN, pre-existing particles in the
7 CCN size range transported to the site by diurnal forced or heat convection might also, in
8 parallel, lead to an apparent increase of the CCN number concentration. Thus, the CCN
9 number concentrations estimated using the methodology previously described, and attributed
10 to NPF in a first approach, might in fact result from both NPF and transport. The transport of
11 particles to the site is taken into account based on the hypothesis that similar number
12 concentrations of particles are transported to the site on event and non-event days. The
13 contribution of NPF to the production of new CCN was thus estimated from the difference
14 between the median CCN increases obtained on event (contributions from NPF and transport)
15 and non-event days (transport only).

16 Among the 362 days included in this analysis, 108 (23 and 85 during the dry and wet season
17 respectively) were identified as non-event days, but only 78 of them (22 from the dry season
18 and 56 from the wet season) were further analysed because of instrumental failures. The
19 median diurnal variation of CCN_{50} obtained on these non-event days and attributed to
20 transport is shown on Fig. 3, together with the median number concentrations obtained on
21 event days and ascribed to both NPF and transport (upper panel). Similar figures are reported
22 in the supplementary material for CCN_{80} and CCN_{100} (Figures S1 and S2, respectively). As
23 previously mentioned, the contribution of NPF to the production of new CCN was estimated
24 from the difference between the median CCN_{50} increases obtained on event and non-event
25 days and is shown on Fig. 3 (lower panel). This absolute CCN production from NPF alone is
26 also reported, together with the corresponding relative concentration increase, on Fig. 4 for
27 further comparison with Fig. 2 (showing both transport and NPF contributions as a whole).

28 During the dry season, transport contributes to CCN_{80} and CCN_{100} to the median level of 1139
29 and 863 cm^{-3} , which is similar to the contribution of NPF (1229 and 784 cm^{-3} for CCN_{80} and
30 CCN_{100} , respectively, Fig. S1, S2). In contrast, CCN_{50} attributed to NPF (3197 cm^{-3})
31 significantly exceeds the median number of particles transported to the site (1610 cm^{-3}) (Fig.
32 3). During the wet season, NPF is likely to be the dominant CCN source, with productions of

1 1950, 771 and 535 cm^{-3} for CCN_{50} , CCN_{80} and CCN_{100} , respectively, compared to median
2 concentrations attributed to transport which do not exceed 690, 404 and 321 cm^{-3} . The
3 contributions of NPF particles to the increase of CCN, all shown on Fig. 4.a. and reported in
4 Table 2 for the different seasons and sizes, hence represent a significant fraction of the CCN
5 increase shown on Fig. 2.a. and reported in Table 1. The contribution of NPF to CCN
6 concentrations are comparable or even higher than those previously mentioned for other
7 stations in the literature, which probably also include CCN sources other than NPF. The
8 relative impact of NPF are estimated to increase the CCN_{50} number concentrations by more
9 than 250 % during both seasons, and the CCN_{100} number concentrations by more than 100%
10 and 200% during the dry season and wet season, respectively.

11 These calculations rely on the hypothesis that the specific environmental conditions on which
12 NPF occurs are not influencing the transport from lower atmospheric layers. In order to
13 further evaluate the reliability of this assumption, wind direction and speed as well as global
14 radiation were investigated on event and non-event days (Figures S3 and S4 in the
15 supplementary material). As previously reported by Rose et al. (2015a), NPF events are
16 favoured during clear sky conditions, when radiation is higher (Fig. S3). Thus, there is likely
17 a bias towards an underestimation of radiative driven transport from lower atmospheric layers
18 due to the fact that cloudy days are over-represented for non-event days. Regarding wind,
19 contrasting directions are also observed between event and non-event days (Fig. S4), with
20 patterns closely related to those observed for the dry and wet seasons, respectively (Rose et
21 al., 2015a). It is worth noticing that winds originating from the more polluted sector of La Paz
22 – El Alto (south) do not seem to be over-represented neither on event nor on non-event days.
23 However, because of the close proximity of this area, it is complex to further assess how it
24 contributes to CCN concentration from wind direction alone, and we cannot exclude a bias
25 related to the variability of this specific source between event and non-event days.
26 Nonetheless, the particle number concentrations observed at the time preceding the usual
27 occurrence of the NPF events are similar for event and non-event days (Fig 3, S1, S2).
28 Moreover, higher wind speeds are on average recorded on non-event days, that likely lead to
29 an enhanced transport of particles to the site compared to event days, and hence lead to an
30 underestimation of the contribution of NPF to the increase of CCN. In any case, taking into
31 account the contribution of transport when calculating the increase of CCN concentrations
32 after NPF events was never done in the past, and certainly helps approaching a more realistic
33 view of the real contribution of NPF to CCN number concentrations.

1

2 **3.2 How layering influences growth to CCN-sizes**

3 3.2.1 Occurrence of NPF in the different tropospheric layers

4 The purpose of this section is to further investigate NPF in terms of occurrence, event type
5 and characteristics (particle formation and growth rate) regarding the location of the station in
6 the tropospheric layers (i.e. BL, FT or IL) at the onset of the NPF process. The classification
7 of air mass types into BL, IL and FT was obtained using the standard deviation of wind
8 direction (Section 2.2).

9 389 NPF events (including all event types, i.e. I, II or bump) previously discussed by Rose et
10 al. (2015a) were included in this analysis. For each event, the air mass type (BL, IL or FT)
11 prevailing at the station was investigated on an hourly basis during the first steps of the NPF
12 process, i.e. from the appearance of the newly formed clusters (< 3nm) to the time at which
13 the concentration of 3-7 nm particles was maximum. There was no information available
14 regarding the classification into BL, IL and FT for 56 events.

15 Various scenarios were observed during this part of the NPF process, which on average
16 lasted for 2.7 ± 1.3 hours. The most frequent scenarios, which include more than 88% of the
17 documented events, are listed, together with their frequency of occurrence, in Table 3.
18 Scenario S1 refers to those days when the first steps of the NPF process were observed to
19 occur in the BL, while scenario S2 refer to the events started in the FT. Scenario S2 is further
20 divided into two sub-classes to distinguish between the events which first steps occur
21 exclusively in the FT (S2.1) from those during which BL dynamics lead to changing
22 conditions in the course of the event (S2.2). Events triggered in the IL are not frequently
23 observed compared to those initiated in the BL or in the FT, and are thus not highlighted in
24 this classification. Since multiple events were frequently detected at Chacaltaya, additional
25 information regarding the occurrence of the scenarios as a function of the event position (first
26 event, second event, third and following events) is also provided. For that purpose, single
27 events and events occurring first on multiple event days were considered all together, while
28 second and following events were considered in a second category.

29 Based on Table 3, constant conditions, i.e. scenarios S1 (BL conditions only) and S2.1 (FT
30 conditions only), were found in 64% of the selected single and first position events and 97%

1 of the second and following events. In each case, scenario S1, corresponding to BL
2 conditions, was the most frequent, representing 93% and 96% of the events initiated in
3 constant conditions, respectively for single and first position events and for second and
4 following ones. The fact that scenario S2.2 related to changing conditions was more
5 frequently observed for single and first position events (36% compared to 3% for following
6 events), i.e. occurring earlier in the morning compared to following events, is mainly
7 explained by the development of the BL during the first part of the day, as shown on Fig. 5.

8 NPF frequencies in the FT and in the BL were also deduced from the previous classification.
9 For that purpose, the analysis was focused on the time period 08:00 - 12:00 (Local), which
10 includes the most probable nucleation hours (Rose et al., 2015a). 72 days (including both
11 event, non-event and undefined days) were rejected from the analysis because of missing
12 information regarding the location of the station in the tropospheric layers. Free tropospheric
13 conditions were detected during at least one hour on 122 days, and among these days, 48
14 showed NPF events initiated in the FT, leading to a NPF frequency of 39%. In contrast, the
15 station laid in the BL during at least one hour on 248 days, and among these days, 119
16 showed events starting in the BL, leading to a NPF frequency of 48%.

17 3.2.2 Event type and characteristics

18 An additional analysis concerning the event type (i.e. I, II or bump) as a function of the
19 scenario was performed using the event classification from Rose et al. (2015a). The results of
20 this analysis are shown on Fig. 6. Almost half of the 77 events triggered in the FT (scenarios
21 S2, Table 3) were identified as type I events (38 events), while types II and bump events were
22 observed on 18 and 21 occasions, respectively, which represent 23 and 27% of scenario S2.
23 When considering the scenarios S2.1 and S2.2 independently from one another, we found that
24 type I events were predominant when changing conditions were detected (S2.2), whereas they
25 displayed similar probabilities of occurrence as other event types in constant free tropospheric
26 conditions (S2.1). This observation suggests that the probability for type I events to occur is
27 increased when initial free tropospheric conditions are changing in the course of the events.
28 This could be explained by favorable conditions for the onset of nucleation events, followed
29 by increased input of condensable species from the BL promoting particle growth. However,
30 this hypothesis must be considered with caution regarding the limited number of events
31 occurring under scenario S2.1. Regarding scenario S1, in the BL, comparable number of

1 events belonging to class I and II were reported (87 and 92 events, thus representing 40 and
2 42% of scenario S1, respectively).

3 In order to further characterize the NPF events in the different atmospheric layers, statistics
4 regarding the formation rate of 2 nm particle and the growth rate (GR) in the size range 1-3
5 nm as a function of the scenarios were performed for type I events. Growth rates were derived
6 from the particle number size distribution using the “maximum” method from Hirsikko et al.
7 (2005), while formation rates were calculated according to Kulmala et al. (2007). Given the
8 limited number of type I events observed in scenario S2.1 (4 events), scenarios S2.1 and S2.2
9 were not distinguished from each other in the statistics. As reported on Fig. 7, increased
10 values are on average reported in the BL, with higher variability, especially for the GR.
11 Additional analysis was performed to investigate the correlation between the GR in the size
12 range 3-7 nm and the location of the station at the end of the scenarios. However, because of
13 an insufficient number of values for events occurring under scenarios ending in the FT
14 (scenario S2.1, 4 values), these results will not be further discussed.

15 We have shown so far that while higher NPF frequencies were found in the BL compared to
16 the FT, higher probabilities for type I events to occur were associated to scenarios starting in
17 the FT and ending in the BL or IL. However, when events belonging to class I are initiated in
18 the BL, they show on average higher particle formation and growth rates compared to those
19 started in the FT. Thus, it is likely that on the one hand higher amounts of gaseous precursors
20 usually associated with the BL could favor nucleation events of higher intensity and explain
21 both higher NPF frequencies and enhanced particle formation and growth rates. On the other
22 hand, cleaner conditions found in the FT at the very beginning of the NPF process may reduce
23 the sink for the newly formed clusters and favor their growth to larger sizes. This observation
24 suggests that the amount of condensable species could directly influence the occurrence of the
25 NPF process and determine the particle growth rate while the occurrence of the growth
26 process itself could rather depend on the strength of the particle sink. Overall, the difference
27 of occurrence frequency, nucleation rates and GR between FT and BL are not very large, and
28 we show that nucleation is initiated in the FT with a rather high frequency.

29 The purpose of the next section is now to investigate the impact of these NPF events on the
30 CCN number concentration in each of the atmospheric layers.

1 3.2.3 CCN production during NPF events in the different tropospheric layers

2 Based on the results discussed in section 3.1.1, 57 NPF event days showing particle growth up
3 to CCN activation diameter were detected at Chacaltaya. 13 of them were not further analyzed
4 due to missing information regarding the location of the station in the tropospheric layers.
5 Among the remaining 44 days, 31 showed events initiated in the BL, 10 in the FT, 2 at the
6 interface between the BL and the FT and 1 in random conditions. Given their limited number,
7 events started in the IL will not be further discussed. The frequency of NPF contribution to
8 the production of new CCN in the BL and in the FT was calculated as the ratio of NPF events
9 growing to the CCN sizes to the total number of type I events occurring in each atmospheric
10 layer, i.e. 46 in the BL and 19 in the FT. The resulting frequency of CCN production from
11 NPF was 67% in the BL, being slightly higher compared to the FT (53%).

12 The number concentration of CCN formed during an event was also analyzed as a function of
13 the air mass type (BL or FT) prevailing at the station (Table 4). Using the three threshold
14 sizes, median CCN productions were comparable for events initiated in the BL and in the FT.
15 In contrast, the third quartiles of CCN_{80} and CCN_{100} were higher for the events initiated in the
16 FT.

17 The fact that the contribution of NPF to the formation of new CCN was more frequently
18 observed for events initiated in the BL might be explained by faster particle growths sustained
19 by higher amounts of condensable material, thus increasing the chances for particles to reach
20 CCN sizes. The tendency for CCN_{80} and CCN_{100} to reach higher values when the NPF
21 process was started in the FT can be due to smaller initial concentrations prior to the NPF
22 event, and thus weaker coagulation associated to less polluted conditions in the FT.

23 Additional analysis regarding the history of the air mass and BL influence along its trajectory
24 would provide valuable information to even more assess the role of the exchanges between
25 the BL and the FT on the occurrence of NPF and its contribution to the formation of new
26 CCN. Indeed, observations conducted at the Jungfraujoch showed that stronger NPF events
27 (type I) occurred in air masses one or two days after contact with the BL (Bianchi et al., 2016;
28 Tröstl et al., 2016). These results are however based on proxies (CO, NO_y) and modelling
29 tools which were unfortunately not available for Chacaltaya. Nevertheless, our results goes to
30 some extent into the same direction as the work by Tröstl et al. (2016) and Bianchi et al.
31 (2016), at least supporting the major role of BL intrusion (regardless of its kind, before or

1 during the event) to sustain particle growth. Similar FT feeding process from the BL was also
2 shown by Rose et al. (2015b) at the puy de Dôme (France, 1465 m a.s.l.).

3 **4 Conclusion**

4 In this paper, the contribution of NPF to the production of potential new CCN was
5 investigated at the highest station in the world, Chacaltaya (5240 m a.s.l., Bolivia), between
6 January 1 and December 31 2012.

7 Using potential CCN activation diameters 50, 80 and 100 nm, we found that 61% of the type I
8 NPF events included in the analysis lead to CCN number concentration increase, with higher
9 probabilities during the wet season (79%) explained by faster particle growth. Because of
10 coagulation on pre-existing particles, the number concentration of CCN formed was observed
11 to decrease with increasing activation diameter, but the frequency of particles reaching the
12 highest potential CCN activation diameter (100nm) was not reduced compared to the lowest
13 CCN size (50 nm). When comparing the CCN production from NPF with the number
14 concentration of pre-existing CCN transported to the site, we found that NPF was on average
15 responsible for the largest contribution to the CCN concentration, especially during the wet
16 season.

17 When segregating into BL and FT air masses sampled at the site, we found slightly higher
18 NPF frequency in the BL (48%) but still an important frequency of occurrence in the FT
19 (nucleation frequency of 39%). This observation is, to our knowledge, the first of its kind.
20 Particle growth was more frequently observed for events initiated in the FT but was on
21 average faster for events started in the BL, most probably because of increased amounts of
22 condensable vapours. As a result, the chance for particles to grow up to potential CCN
23 activation diameters was higher when the NPF process occurred in the BL. In contrast, the
24 impact of NPF initiated in the FT on CCN number concentrations was higher than for NPF
25 initiated in the BL, most likely because of the decreased pollution levels and weaker
26 coagulation sink. The previous observations clearly highlight the competition that exists
27 between particle growth and their removal by coagulation processes on pre-existing particles,
28 and thus the complex balance between sources and sinks that is required to observe the
29 formation of new particles and their subsequent growth to climate relevant sizes. Such
30 conditions are often fulfilled at Chacaltaya, where NPF seems to often play a dominant role in
31 the formation of new CCN.

32

1 *Acknowledgments*

2 *This work was performed within the framework of ACTRIS2 (Aerosols, Clouds and Trace*
3 *gases Research Infra Structure Network) and was also supported by SNO CLAP, IRD, LEFE-*
4 *CHAT, OSUG@2020 as well as STINT and FORMAS funding agencies.*

5

6 **References**

- 7 Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245(4923),
8 1227–1230, 1989.
- 9 Andrade, M., Zaratti, F., Forno, R., Gutiérrez, R., Moreno, I., Velarde, F., Ávila, F., Roca, M.,
10 Sánchez, M. F., Laj, P., Jaffrezo, J. L., Ginot, P., Sellegri, K., Ramonet, M., Laurent, O.,
11 Weinhold, K., Wiedensohler, A., Krejci, R., Bonasoni, P., Cristofanelli, P., Whiteman, D.,
12 Vimeux, F., Dommergue, A. and Magand, O.: Puesta en marcha de una nueva estación de
13 monitoreo climático en los andes centrales de Bolivia: la estación Gaw/Chacaltaya, *Revista*
14 *Boliviana de Física*, 26(26), 06-15, 2015.
- 15 Asmi, E., Kivekäs, N., Kerminen, V.-M., Komppula, M., Hyvärinen, A.-P., Hatakka, J.,
16 Viisanen, Y. and Lihavainen, H.: Secondary new particle formation in Northern Finland
17 Pallas site between the years 2000 and 2010, *Atmos. Chem. Phys.*, 11(24), 12959–12972,
18 doi:10.5194/acp-11-12959-2011, 2011.
- 19 Asmi, E., Freney, E., Hervo, M., Picard, D., Rose, C., Colomb, A. and Sellegri, K.: Aerosol
20 cloud activation in summer and winter at puy-de-Dôme high altitude site in France, *Atmos.*
21 *Chem. Phys.*, 12(23), 11589–11607, doi:10.5194/acp-12-11589-2012, 2012.
- 22 Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U.,
23 Herrmann, E., Adamov, A., Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M.,
24 Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H. E., Münch, S., Peräkylä, O.,
25 Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala,
26 M., Dommen, J. and Baltensperger, U.: New particle formation in the free troposphere: A
27 question of chemistry and timing, *Science*, 352(6289), 1109–1112,
28 doi:10.1126/science.aad5456, 2016.
- 29 EPA: Quality Assurance Handbook for Air Pollution Measurement Systems, Volume IV:
30 Meteorological measurements, 2008.

1 Hammer, E., Bukowiecki, N., Gysel, M., Jurányi, Z., Hoyle, C. R., Vogt, R., Baltensperger,
2 U. and Weingartner, E.: Investigation of the effective peak supersaturation for liquid-phase
3 clouds at the high-alpine site Jungfraujoch, Switzerland (3580 m a.s.l.), *Atmos. Chem. Phys.*,
4 14(2), 1123–1139, doi:10.5194/acp-14-1123-2014, 2014.

5 Herrmann, E., Weingartner, E., Henne, S., Vuilleumier, L., Bukowiecki, N., Steinbacher, M.,
6 Conen, F., Collaud Coen, M., Hammer, E., Jurányi, Z., Baltensperger, U. and Gysel, M.:
7 Analysis of long-term aerosol size distribution data from Jungfraujoch with emphasis on free
8 tropospheric conditions, cloud influence, and air mass transport, *J. Geophys. Res. Atmos.*,
9 120(18), 2015JD023660, doi:10.1002/2015JD023660, 2015.

10 Hirsikko, A., Laakso, L., Horrak, U., Aalto, P. P., Kerminen, V. and Kulmala, M.: Annual and
11 size dependent variation of growth rates and ion concentrations in boreal forest, *Boreal
12 environment research*, 10(5), 357, 2005.

13 Hirsikko, A., Bergman, T., Laakso, L., Dal Maso, M., Riipinen, I., Hörrak, U. and Kulmala,
14 M.: Identification and classification of the formation of intermediate ions measured in boreal
15 forest, *Atmos. Chem. Phys.*, 7(1), 201–210, doi:10.5194/acp-7-201-2007, 2007.

16 IPCC: IPCC (AR5):Climate change 2013: The Physical Science Basis, Summary for
17 policymakers. Contribution of Working Group I to the Fifth Assessment report, 2013.

18 Jurányi, Z., Gysel, M., Weingartner, E., Bukowiecki, N., Kammermann, L. and Baltensperger,
19 U.: A 17 month climatology of the cloud condensation nuclei number concentration at the
20 high alpine site Jungfraujoch, *J. Geophys. Res.*, 116(D10), D10204,
21 doi:10.1029/2010JD015199, 2011.

22 Kerminen, V.-M., Paramonov, M., Anttila, T., Riipinen, I., Fountoukis, C., Korhonen, H.,
23 Asmi, E., Laakso, L., Lihavainen, H., Swietlicki, E., Svenningsson, B., Asmi, A., Pandis, S.
24 N., Kulmala, M. and Petäjä, T.: Cloud condensation nuclei production associated with
25 atmospheric nucleation: a synthesis based on existing literature and new results, *Atmos.
26 Chem. Phys.*, 12(24), 12037–12059, doi:10.5194/acp-12-12037-2012, 2012.

27 Komppula, M., Lihavainen, H., Kerminen, V.-M., Kulmala, M. and Viisanen, Y.:
28 Measurements of cloud droplet activation of aerosol particles at a clean subarctic background
29 site, *Journal of Geophysical Research: Atmospheres* (1984–2012), 110(D6) [online] Available
30 from: <http://www.agu.org/journals/jd/jd0506/2004JD005200/2004jd005200-t03.txt> (Accessed
31 21 February 2014), 2005.

1 Kulmala, M. and Kerminen, V.-M.: On the formation and growth of atmospheric
2 nanoparticles, *Atmospheric Research*, 90(2–4), 132–150, doi:10.1016/j.atmosres.2008.01.005,
3 2008.

4 Kulmala, M., Riipinen, I., Sipilä, M., Manninen, H. E., Petäjä, T., Junninen, H., Maso, M. D.,
5 Mordas, G., Mirme, A., Vana, M., Hirsikko, A., Laakso, L., Harrison, R. M., Hanson, I.,
6 Leung, C., Lehtinen, K. E. J. and Kerminen, V.-M.: Toward Direct Measurement of
7 Atmospheric Nucleation, *Science*, 318(5847), 89–92, doi:10.1126/science.1144124, 2007.

8 Laakso, L., Merikanto, J., Vakkari, V., Laakso, H., Kulmala, M., Molefe, M., Kgabi, N.,
9 Mabaso, D., Carslaw, K. S., Spracklen, D. V., Lee, L. A., Reddington, C. L. and Kerminen,
10 V.-M.: Boundary layer nucleation as a source of new CCN in savannah environment, *Atmos.*
11 *Chem. Phys.*, 13(4), 1957–1972, doi:10.5194/acp-13-1957-2013, 2013.

12 Laaksonen, A., Hamed, A., Joutsensaari, J., Hiltunen, L., Cavalli, F., Junkermann, W., Asmi,
13 A., Fuzzi, S. and Facchini, M. C.: Cloud condensation nucleus production from nucleation
14 events at a highly polluted region, *Geophysical research letters*, 32(6) [online] Available
15 from: <http://onlinelibrary.wiley.com/doi/10.1029/2004GL022092/full> (Accessed 21 February
16 2014), 2005.

17 Lee, L. A., Pringle, K. J., Reddington, C. L., Mann, G. W., Stier, P., Spracklen, D. V., Pierce,
18 J. R. and Carslaw, K. S.: The magnitude and causes of uncertainty in global model
19 simulations of cloud condensation nuclei, *Atmos. Chem. Phys.*, 13(17), 8879–8914,
20 doi:10.5194/acp-13-8879-2013, 2013.

21 Lihavainen, H., Kerminen, V.-M., Komppula, M., Hatakka, J., Aaltonen, V., Kulmala, M. and
22 Viisanen, Y.: Production of “potential” cloud condensation nuclei associated with
23 atmospheric new-particle formation in northern Finland, *Journal of Geophysical Research:*
24 *Atmospheres* (1984–2012), 108(D24) [online] Available from:
25 <http://onlinelibrary.wiley.com/doi/10.1029/2003JD003887/full> (Accessed 21 February 2014),
26 2003.

27 Makkonen, R., Asmi, A., Kerminen, V.-M., Boy, M., Arneth, A., Hari, P. and Kulmala, M.:
28 Air pollution control and decreasing new particle formation lead to strong climate warming,
29 *Atmos. Chem. Phys.*, 12(3), 1515–1524, doi:10.5194/acp-12-1515-2012, 2012.

30 Manninen, H. E., Mirme, S., Mirme, A., Petäjä, T. and Kulmala, M.: How to reliably detect
31 molecular clusters and nucleation mode particles with Neutral cluster and Air Ion

1 Spectrometer (NAIS), Atmospheric Measurement Techniques Discussions, 1–57,
2 doi:10.5194/amt-2016-25, 2016.

3 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J. and Carslaw, K. S.: Impact of
4 nucleation on global CCN, *Atmos. Chem. Phys.*, 9(21), 8601–8616, doi:10.5194/acp-9-8601-
5 2009, 2009.

6 Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS – a
7 spectrometer for the measurement of cluster ion and nanometer aerosol size distributions,
8 *Atmospheric Measurement Techniques*, 6(4), 1061–1071, doi:10.5194/amt-6-1061-2013,
9 2013.

10 Mitchell, A. E.: A comparison of short-term dispersion estimates resulting from various
11 atmospheric stability classification methods, *Atmospheric Environment* (1967), 16(4), 765–
12 773, doi:10.1016/0004-6981(82)90394-8, 1982.

13 Mitchell, A. E. and Timbre, K. O.: Atmospheric stability class from horizontal wind
14 fluctuation, *Proceedings of the LXXII Annual Meeting of the Air Pollution Control*
15 *Association*, 79–29.2, 1979.

16 Pierce, J. R., Leaitch, W. R., Liggio, J., Westervelt, D. M., Wainwright, C. D., Abbatt, J. P.
17 D., Ahlm, L., Al-Basheer, W., Cziczo, D. J., Hayden, K. L. and others: Nucleation and
18 condensational growth to CCN sizes during a sustained pristine biogenic SOA event in a
19 forested mountain valley, *Atmospheric Chemistry and Physics*, 12(7), 3147–3163, 2012.

20 Reddington, C. L., Carslaw, K. S., Spracklen, D. V., Frontoso, M. G., Collins, L., Merikanto,
21 J., Minikin, A., Hamburger, T., Coe, H., Kulmala, M., Aalto, P., Flentje, H., Plass-Dülmer,
22 C., Birmili, W., Wiedensohler, A., Wehner, B., Tuch, T., Sonntag, A., O’Dowd, C. D.,
23 Jennings, S. G., Dupuy, R., Baltensperger, U., Weingartner, E., Hansson, H.-C., Tunved, P.,
24 Laj, P., Sellegri, K., Boulon, J., Putaud, J.-P., Gruening, C., Swietlicki, E., Roldin, P.,
25 Henzing, J. S., Moerman, M., Mihalopoulos, N., Kouvarakis, G., Ždímal, V., Zíková, N.,
26 Marinoni, A., Bonasoni, P. and Duchi, R.: Primary versus secondary contributions to particle
27 number concentrations in the European boundary layer, *Atmos. Chem. Phys.*, 11(23), 12007–
28 12036, doi:10.5194/acp-11-12007-2011, 2011.

29 Roberts, G. C. and Nenes, A.: A continuous-flow streamwise thermal-gradient CCN chamber
30 for atmospheric measurements, *Aerosol Science and Technology*, 39(3), 206–221, 2005.

1 Roberts, G. C., Day, D. A., Russell, L. M., Dunlea, E. J., Jimenez, J. L., Tomlinson, J. M.,
2 Collins, D. R., Shinozuka, Y. and Clarke, A. D.: Characterization of particle cloud droplet
3 activity and composition in the free troposphere and the boundary layer during INTEX-B,
4 *Atmos. Chem. Phys.*, 10(14), 6627–6644, doi:10.5194/acp-10-6627-2010, 2010.

5 Rose, C., Sellegri, K., Velarde, F., Moreno, I., Ramonet, M., Weinhold, K., Krejci, R.,
6 Andrade, M., Wiedensohler, A. and Laj, P.: Frequent nucleation events at the high altitude
7 station of Chacaltaya (5240 m a.s.l.), Bolivia, *Atmospheric Environment*, 102, 18–29,
8 doi:10.1016/j.atmosenv.2014.11.015, 2015a.

9 Rose, C., Sellegri, K., Asmi, E., Hervo, M., Freney, E., Colomb, A., Junninen, H., Duplissy,
10 J., Sipilä, M., Kontkanen, J., Lehtipalo, K. and Kulmala, M.: Major contribution of neutral
11 clusters to new particle formation at the interface between the boundary layer and the free
12 troposphere, *Atmos. Chem. Phys.*, 15(6), 3413–3428, doi:10.5194/acp-15-3413-2015, 2015b.

13 Seaton, A., Godden, D., MacNee, W. and Donaldson, K.: Particulate air pollution and acute
14 health effects, *The Lancet*, 345(8943), 176–178, doi:10.1016/S0140-6736(95)90173-6, 1995.

15 Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I.,
16 Merikanto, J., Mann, G. W., Chipperfield, M. P. and Wiedensohler, A.: Contribution of
17 particle formation to global cloud condensation nuclei concentrations, *Geophysical Research*
18 *Letters*, 35(6) [online] Available from:
19 <http://onlinelibrary.wiley.com/doi/10.1029/2007GL033038/full> (Accessed 17 October 2013),
20 2008.

21 Tröstl, J., Herrmann, E., Frege, C., Bianchi, F., Molteni, U., Bukowiecki, N., Hoyle, C. R.,
22 Steinbacher, M., Weingartner, E., Dommen, J., Gysel, M. and Baltensperger, U.: Contribution
23 of new particle formation to the total aerosol concentration at the high altitude site
24 Jungfraujoch (3'580 m a.s.l., Switzerland), *J. Geophys. Res. Atmos.*, 2015JD024637,
25 doi:10.1002/2015JD024637, 2016.

26 Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *Journal of the*
27 *atmospheric sciences*, 34(7), 1149–1152, 1977.

28 Weber, R. O.: Estimators for the Standard Deviation of Horizontal Wind Direction, *J. Appl.*
29 *Meteor.*, 36(10), 1403–1415, doi:10.1175/1520-0450(1997)036<1403:EFTSDO>2.0.CO;2,
30 1997.

1 Wex, H., McFiggans, G., Henning, S. and Stratmann, F.: Influence of the external mixing
2 state of atmospheric aerosol on derived CCN number concentrations, *Geophysical Research*
3 *Letters*, 37(10) [online] Available from:
4 <http://onlinelibrary.wiley.com/doi/10.1029/2010GL043337/full> (Accessed 21 February 2014),
5 2010.

6 Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner,
7 B., Tuch, T., Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac,
8 H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P.,
9 Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F.,
10 Santos, S., Gröning, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G.,
11 O'Dowd, C. D., Marinoni, A., Horn, H.-G., Keck, L., Jiang, J., Scheckman, J., McMurry, P.
12 H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Löschau, G. and
13 Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and
14 data structure to facilitate high quality long-term observations of atmospheric particle number
15 size distributions, *Atmospheric Measurement Techniques*, 5(3), 657–685, doi:10.5194/amt-5-
16 657-2012, 2012.

17 Yamartino, R. J.: A Comparison of Several “Single-Pass” Estimators of the Standard
18 Deviation of Wind Direction, *Journal of Climate and Applied Meteorology*, 23(9), 1362–
19 1366, doi:10.1175/1520-0450(1984)023<1362:ACOSPE>2.0.CO;2, 1984.

20 Yli-Juuti, T., Riipinen, I., Aalto, P. P., Nieminen, T., Maenhaut, W., Janssens, I. A., Claeys,
21 M., Salma, I., Ocskay, R. and Hoffer, A.: Characteristics of new particle formation events and
22 cluster ions at K-pusztá, Hungary, *Boreal Environment Research*, 14(4), 683–698, 2009.

23 Yli-Juuti, T., Nieminen, T., Hirsikko, A., Aalto, P. P., Asmi, E., Hörrak, U., Manninen, H. E.,
24 Patokoski, J., Dal Maso, M., Petäjä, T., Rinne, J., Kulmala, M. and Riipinen, I.: Growth rates
25 of nucleation mode particles in Hyytiälä during 2003–2009: variation with particle size,
26 season, data analysis method and ambient conditions, *Atmos. Chem. Phys.*, 11(24), 12865–
27 12886, doi:10.5194/acp-11-12865-2011, 2011.

28
29
30
31

1
2
3
4
5
6
7
8
9
10
11
12
13

14 Table 1 Estimation of the median seasonal and annual CCN productions during NPF events.

	CCN ₅₀ (cm ⁻³)	CCN ₈₀ (cm ⁻³)	CCN ₁₀₀ (cm ⁻³)
Dry season	3.96×10 ⁵	1.60×10 ⁵	9.40×10 ⁴
Wet season	8.48×10 ⁴	4.98×10 ⁴	3.90×10 ⁴
Whole year	4.81×10 ⁵	2.10×10 ⁵	1.33×10 ⁵

15
16

17 Table 2 Estimation of the median seasonal and annual CCN increases from NPF, i.e.
18 corrected for the contribution of particles transported to the site.

	CCN ₅₀ (cm ⁻³)	CCN ₈₀ (cm ⁻³)	CCN ₁₀₀ (cm ⁻³)
Dry season	2.00×10 ⁵	7.71×10 ⁴	4.92×10 ⁴
Wet season	5.39×10 ⁴	2.13×10 ⁴	1.48×10 ⁴
Whole year	2.54×10 ⁵	9.84×10 ⁴	6.40×10 ⁴

1
2
3
4
5
6
7
8
9
10

11 Table 3 Description of the scenarios concerning the location of the station in the troposphere
12 (boundary layer (BL), interface layer (IL) and free troposphere (FT)) during the first steps of
13 the NPF process, i.e. from the appearance of the newly formed clusters (< 3nm) to the time at
14 which the concentration of 3-7 nm particles was maximum. The total number of occurrence is
15 provided for each scenario in the second column. Since multiple events are frequently
16 observed at Chacaltaya, a more detailed classification including the event position is specified
17 in the last two columns.

Scenario	Description	Total number of occurrence	Single and first position events	Second and following events
S1	First steps of NPF occur in BL	217	100	117
S2		77	68	9
<i>S2.1</i>	First steps of NPF occur in FT	12	7	5
<i>S2.2</i>	Nucleation occurs in FT and initial particle growth is observed in changing conditions, from FT to IL/BL	65	61	4

18

1 Table 4. CCN production as a function of the location of the station (BL or FT) at the onset of
 2 the NPF process.

Threshold CCN size	CCN increase for events started in the BL (cm^{-3})			CCN increase for events started in the FT (cm^{-3})		
	25 th perc.	Median	75 th perc.	25 th perc.	Median	75 th perc.
50 nm	2556	5072	10110	3070	5137	9378
80 nm	1155	2416	3919	1483	2138	5173
100 nm	820	1518	2338	960	1447	3568

3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20





Fig. 1. Determination of the CCN concentration increase for the 3 threshold diameters (50, 80 and 100 nm) from the particle size distributions measured by SMPS. t_{init} and t_{max} denote, for each diameter, the times from which concentration increases are calculated. July 24th 2012.


1
2
3
4
5
6
7
8
9
10



20
21
22
23
24
25
26


Fig. 2. Median a. absolute and b. relative CCN productions observed during type I events for the different activation diameters and seasons (wet and dry). Lower and upper limits of the error bars stand for the 1st and 3rd quartile, respectively.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22



23 Fig. 3. Median diurnal variation of CCN₅₀ on event (upper panel, “Transport and NPF”) and
24 non-event days (upper panel, “Transport”). CCN₅₀ attributed to NPF (lower panel) is
25 calculated as the difference of the concentrations recorded on event and non-event days.
26 Lower and upper limits of the error bars stand for the 1st and 3rd quartile, respectively.


1
2
3
4
5
6
7
8
9
10
11
12



21
22
23
24
25
26

Fig. 4. Median a. absolute and b. relative CCN productions from NPF, i.e. corrected for the transport of CCN-size particles to the site, for the different activation diameters and seasons (wet and dry).

1
2
3
4
5
6
7
8
9
10
11



21
22
23
24
25
26

Fig. 5. Statistics on the location of the station in the tropospheric layers (boundary layer (BL), interface layer (IL) and free troposphere (FT)) between 8:00 and 12:00 (Local), separately for the dry and wet seasons.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26




Fig. 6. Statistics on the event type (I, II or bump) as a function of the scenario describing the location of the station in the tropospheric layers (see Table 3).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24




Fig. 7. Median formation rate of 2 nm particles (J_2) and growth rate in the range 1-3 nm (GR_{1-3}) reported separately for type I events initiated in the BL (scenario S1) and in the FT (scenario S2). Lower and upper limits of the error bars stand for the 1st and 3rd quartile, respectively.