

Interactive comment on “Ten years of atmospheric methane from ground-based NDACC FTIR observations” by Whitney Bader et al.

Anonymous Referee #2

Received and published: 20 September 2016

This manuscript presents the analysis of atmospheric methane trends derived from FTIR measurements at ten NDACC sites that cover both, Northern and Southern Hemisphere over the period of 2005-2014. Using GEOS-Chem simulations the authors found that - anthropogenic sources of atmospheric CH₄ are responsible for the renewed growth of CH₄ that has been registered by different observational systems since 2005; - main contributors into the interannual variations of CH₄ total columns are the natural sources (wetlands and biomass burning).

General comments:

In Page 5 line 15. The authors noted that CH₄ total columns for the Toronto site have a systematic error due to unknown instrument artifact and then made some manipulations with the data which seem to be doubtful. The main issue is how to separate

C1

(in data) the signals, which come from real atmospheric processes and from the instrument that doesn't work in a proper way. Could authors suggest more reasonable criteria/way for the correction of CH₄ time series for Toronto? Or, maybe, it would be better to omit Toronto site's data for the period of 2008-2009 from the analysis?

In Page 9 line 34. The explanation of the the lower value of CH₄ trend (~0.22%/yr for 2005-2012) for the Jungfraujoch site given in the paper is in contradiction to the following: - according to a reference (Collaud Coen,2011), the coming of polluted air to the Jungfraujoch site was usually detected using the monitoring of CO, NO_x and SO₂ concentrations in the ambient air by local sensors. Authors need to bring compelling arguments proving that portions of polluted air, which can reach high altitude site, will significantly influence not only the concentrations of some gases but also the CH₄ total column. - for Zugspitze (also a high altitude site), which is located not so far from Jungfraujoch, CH₄ trend has the value of 0.31%/yr (2005-2012). Therefore it is worth to explain such noticeable difference between trends for Jungfraujoch and Zugspitze.

In Page 10 line 13. This is not quite clear why “small annual changes of methane and smaller uncertainty ... complicates the agreement between the FTIR and GEOS-Chem ...”.

In Page30 Table 3. Methane trends derived from FTIR measurements over 2005-2012 are higher for the stations that are located in the Northern Hemisphere than for sites in the Southern Hemisphere. In comparison to FTIR trends, GEOS-Chem simulations give us an opposite tendency: CH₄ trends have lower values for the Northern Hemisphere. Could authors suggest any reasons of such inconsistency between observational and modeling estimations of CH₄ trend?

Technical corrections: Table 3. Column “GEOS-Chem trend [2005-2012]”, row “Unit”: please, check units.

Taking to account the above described considerations the manuscript can be published in ACP.

C2

